Autophagy is required for the rejection of self-incompatible pollen in two accessions of

transgenic *Arabidopsis thaliana*

Stuart R. Macgregor,[a] Hyun Kyung Lee,[a,1] Hayley Nelles,[a] Daniel C. Johnson,[a] Tong Zhang,[b]

Chaozhi Ma,[b] and Daphne R. Goring[a,c,2,3]

[a]Department of Cell & Systems Biology, University of Toronto, Toronto, Canada M5S 3B2.

[b]National Key Laboratory of Crop Genetic Improvement, National Center of Rapeseed

Improvement in Wuhan, Huazhong Agricultural University, Wuhan 430070, China.

[c]Centre for the Analysis of Genome Evolution & Function, University of Toronto, Toronto,

Canada M5S 3B2.

[1]Current address: Dept. Plant Molecular Biology, University of Lausanne, Switzerland.

[2]Author for Contact, [3]Senior author: Daphne Goring d.goring@utoronto.ca

Short Title: Autophagy in *Arabidopsis* self-incompatibility

One Sentence Summary: In self-incompatible transgenic *Arabidopsis thaliana* lines, autophagy

is an integral part of the cellular responses in the stigma to efficiently block fertilization by self-

incompatible pollen.

Author Contributions: DRG conceived the project; DRG, SRM, HKL, HN, DCJ and TZ

conceptualized and designed experiments; SRM performed crosses, generated and analyzed data

for the SI-C24 *atg7-9* plants, SI-Col-0 *atg5-1* plants, and the SI-Col-0 *atg7-2* GFP-ATG8a

imaging; HKL and SRM performed crosses, generated and analyzed data for the SI-Col-0 *atg7-2*

plants; HN and DCJ performed preliminary work on the SI lines and imaging; TZ generated the

SI-Col-0 line; DRG and CM supervised the experiments; SRM, HKL, and DRG wrote the

manuscript; All authors commented on the manuscript before submission; DRG agrees to serve

as the author responsible for contact and ensures communication.

Funding information: This research was supported by a grant from Natural Sciences and

Engineering Research Council of Canada to DRG, SRM was supported by the Rustom H. Dastur

Graduate Scholarship, and HKL was supported by an Ontario Graduate Scholarship.

Abstract

Successful reproduction in the Brassicaceae is mediated by a complex series of interactions

between the pollen and the pistil, and some species have an additional layer of regulation with

the self-incompatibility trait. While the initial activation of the self-incompatibility pathway by

the pollen S-locus protein11/*S*-locus cysteine-rich peptide and the stigma *S* Receptor Kinase is

well characterized, the downstream mechanisms causing self-pollen rejection are still not fully

understood. In previous studies, we had detected the presence of autophagic bodies with self-

incompatible pollinations in *Arabidopsis lyrata* and transgenic *A. thaliana* lines, but it was not

known if autophagy was essential for self-pollen rejection. Here, we investigated the requirement

of autophagy in this response by crossing mutations in the essential *AUTOPHAGY7* (*ATG7*) or

AUTOPHAGY5 (*ATG5*) genes into two different transgenic self-incompatible *A. thaliana* lines in

the Col-0 and C24 accessions. By using these previously characterized transgenic lines that

express *A. lyrata* and *A. halleri* self-incompatibility genes, we demonstrated that disrupting

autophagy can weaken their self-incompatible responses in the stigma. When the *atg7* or *atg5*

mutations were present, an increased number of self-incompatible pollen were found to hydrate

and form pollen tubes that successfully fertilized the self-incompatible pistils. Additionally, we

confirmed the presence of GFP-ATG8a labelled autophagosomes in the stigmatic papillae

following self-incompatible pollinations. Together, these findings support the requirement of

autophagy in the self-incompatibility response and add to the growing understanding of the

cellular events that take place in the stigma to reject self-pollen.

Key words

Autophagy, self-incompatibility, pollen-pistil interactions, reproduction, *SP11/SCR*, *SRK*, *ARC1*,

ATG7, ATG5, GFP-ATG8a

Introduction

Reproduction in the Brassicaceae is a tightly controlled process, with many components contributing to successful pollen-pistil interactions (reviewed in Doucet et al., 2016; Mizuta and Higashiyama, 2018; Johnson et al., 2019; Adhikari et al., 2020). The Brassicaceae dry stigma lacks surface secretions and instead regulates the delivery of resources to the pollen shortly after pollination. This leads to the selective hydration of compatible pollen grains, followed by the penetration of the pollen tube into the stigmatic barrier to grow down the reproductive tract (reviewed in Dickinson, 1995; Palanivelu and Tsukamoto, 2012; Doucet et al., 2016; Adhikari et al., 2020). It is also at this initial cell-cell communication stage that self-incompatible pollen is recognized and blocked by the stigma to prevent inbreeding (reviewed in Jany et al., 2019; Nasrallah, 2019; Durand et al., 2020). A recent live-cell imaging study in *Arabidopsis thaliana* compared pollen hydration rates of compatible and self-incompatible pollen grains and found that while compatible pollen grains rapidly hydrated, self-incompatible pollen failed to fully hydrate which then led to a failure in pollen germination or a failure of pollen tubes to penetrate the stigmatic papilla cell wall (Rozier et al., 2020). Furthermore, actin foci were present in the stigmatic papillae at the site of emerging pollen tubes for compatible pollen only. The actin focalization is likely associated with the stigmatic papilla cellular responses to enable compatible pollen hydration and pollen tube penetration (Rozier et al., 2020).

The Brassicaceae self-incompatible response is initiated by the binding of the *S*-haplotype-specific pollen ligand, *S*-locus protein 11/*S*-locus cysteine-rich peptide (SP11/SCR) to the corresponding stigma-specific *S* Receptor Kinase (SRK). This leads to the phosphorylation of the SRK kinase domain and activation of the downstream signaling responses, ultimately leading to the rejection of the self-incompatible pollen grain reviewed in Jany et al., 2019; Nasrallah, 2019). Not all Brassicaceae species possess this trait due to mutations in the key self-incompatibility genes, *SP11/SCR* and *SRK* (reviewed in Durand et al., 2020). For example, the transition of *A. thaliana* to a fully selfing species is associated with mutations in the *SCR* and *SRK* genes (Kusaba et al., 2001; Shimizu et al., 2008; Boggs et al., 2009; Tsuchimatsu et al., 2010; Guo et al., 2011; Tsuchimatsu et al., 2017). In contrast, other family members like *A. lyrata* and *A. halleri* have intact *SCR* and *SRK* genes, which result in a robust self-incompatibility response upon self-pollination (Kusaba et al., 2001; Schierup et al., 2001; Goubet et al., 2012).

A number of studies have reintroduced the self-incompatibility trait back into *A. thaliana* by

introducing *A. lyrata* or *A. halleri SP11/SCR* and *SRK* transgenes (Nasrallah et al., 2004; Boggs

et al., 2009; Indriolo et al., 2014; Iwano et al., 2015; Zhang et al., 2019; Rozier et al., 2020).

However, the success of this approach was accession-dependent; for example, transgenic

experiments in *A. thaliana* Col-0 failed to produce self-incompatible plants, whereas experiments

using *A. thaliana* C24 were very successful (Nasrallah et al., 2002; Boggs et al., 2009; Iwano et

al., 2015; Rozier et al., 2020). The cause of this was linked to inverted repeats within the *A.*

thaliana SRKA pseudogene which triggers RNA silencing of the introduced *SRK* transgenes

(Fujii et al., 2020). For the Col-0 accession, the inverted repeat was found in a conserved region

of the predicted *pseudoSRKA* kinase domain resulting in the silencing of closely related *SRK*

transgenes. For the C24 accession, the inverted repeat was found in a more polymorphic region

of the predicted *pseudoSRKA* ectodomain and appeared to be less effective at inducing RNA

silencing of related *SRK* transgenes (Fujii et al., 2020).

We have found that a strong self-incompatibility response can also be recapitulated in *A.*

thaliana Col-0 when the *A. lyrata* or *A. halleri ARC1* transgene was added along with the *A.*

lyrata or *A. halleri SP11/SCR* and *SRK* transgenes (Indriolo et al., 2014; Zhang et al., 2019). The

ARC1 E3 ubiquitin ligase is proposed to act directly downstream of SRK in the self-

incompatibility signaling pathway (Gu et al., 1998; Stone et al., 1999; Stone et al., 2003). A large

part of the *ARC1* gene has been deleted in the *A. thaliana* genome, and knockdowns of *ARC1* in

self-incompatible *Brassica napus* and *A. lyrata* resulted in a partial breakdown of self-

incompatibility (Stone et al., 1999; Indriolo et al., 2012). ARC1 is proposed to target cellular

components that are required in the stigma to accept compatible pollen, such as the EXO70A1

exocyst subunit (Samuel et al., 2009) and *Brassica* GLO1 (Sankaranarayanan et al., 2015). Self-

incompatible pollinations also induced a strong and rapid cytosolic calcium influx in the

stigmatic papillae (Iwano et al., 2015). These combined self-incompatibility responses are

proposed to disrupt the secretory activity in the stigmatic papillae which would be required for

compatible pollen acceptance (Dickinson, 1995; Elleman and Dickinson, 1996; Samuel et al.,

2009; Safavian and Goring, 2013; Safavian et al., 2015; Goring, 2017).

In previous work, we uncovered signs of autophagy activation in *A. lyrata* and *A. thaliana*

stigmatic papillae within 10 minutes of self-incompatible pollinations, but not with compatible

pollinations (Safavian and Goring, 2013; Indriolo et al., 2014). TEM images revealed structures

resembling autophagic bodies in the vacuoles of stigmatic papilla following self-incompatible

pollinations (Safavian and Goring, 2013; Indriolo et al., 2014). In contrast, TEM images of

compatible pollinations revealed vesicle-like structures in the stigmatic papillae under the

compatible pollen contact site suggestive of secretory activity (Safavian and Goring, 2013;

Indriolo et al., 2014). The presence of autophagosomes in *A. lyrata* self-incompatible

pollinations was further confirmed using the GFP-ATG8a autophagosome marker (Safavian and

Goring, 2013). Although these studies indicated that autophagy was activated following self-

incompatible pollinations, it was unclear whether autophagy is essential for the rejection of self-

incompatible pollen. Here, we have used two different self-incompatible (SI) transgenic *A.*

thaliana lines (in the Col-0 and C24 accessions; Iwano et al., 2014; Zhang et al., 2019) to test the

requirement of the autophagy system for self-pollen rejection. When loss-of-function mutants in

the *AUTOPHAGY7* (*ATG7*) and *AUTOPHAGY5(ATG5)* genes (Chung et al., 2010; Ding et al.,

2018) were combined with the SI-Col-0 and SI-C24 lines, we observed a partial breakdown of

the self-incompatibility response in both accessions. This indicates that autophagic activity is

required for the effective rejection of self-incompatible pollen, and thus, elucidates an additional

downstream cellular event in the self-incompatibility pathway.

Results

Generation of self-incompatible *A. thaliana* transgenic lines carrying the *atg7* and *atg5*

mutations to disrupt autophagy

To investigate the involvement of autophagy in self-incompatibility, two different transgenic *A.*

thaliana SI lines were used to cross with *atg7* and *atg5* autophagy mutants. ATG7 and ATG5 are

essential components in the formation of autophagosomes (reviewed in Ding et al., 2018), and

both the *atg7* and *atg5* mutations lead to a loss of autophagy (Thompson et al., 2005; Chung et

al., 2010; Young et al., 2019). The first self-incompatible line is in the Col-0 accession (SI-Col-

0) and carries the *A. halleri SCR13*, *SRK13* and *ARC1* transgenes in a single T-DNA (*Ah-SCR13*

Ah-SRK13 Ah-ARC1 #2 line; Zhang et al., 2019). This SI-Col-0 line was crossed to the

previously characterized *atg7-2* mutant in the Col-0 accession (Chung et al., 2010). The F_1 plants

from this cross were selfed, and all pollination assays were carried out in the F_2 generation. To

confirm that any observed effects were not specific to the loss of *ATG7*, the previously

characterized *atg5-1* mutation in the Col-0 accession (Thompson et al., 2005) was also crossed

into the SI-Col-0 line, and taken to the F_2 generation for pollination assays. The second self-
incompatible line is in the C24 accession (SI-C24) and carries the *A. lyrata SCRb* and *SRKb*
transgenes (SI-C24 #15-1 line; Iwano et al., 2015). As there were no *atg7* mutants available in
the C24 accession, CRISPR/Cas9 technology was used to generate additional *atg7* mutants.
Deletion of the *ATG7* gene was confirmed through PCR and sequencing of genomic regions
flanking gRNA target sites (Supplemental Fig. S1). The *atg7-9* C24 mutant displayed the same
overall wild-type plant and flower morphology (Supplemental Fig. S2) as seen for the *atg7-2*
Col-0 mutant. The C24 *atg7-9* mutant was crossed with the SI-C24 line, and following selfing of
the F_1 plants, all pollination assays were carried out in F_2 generation (T_4 generation for the *atg7-*
9 mutation). For all three sets of F_2 progeny, genotyping was first carried out to identify the
transgenic plants carrying the SI transgenes, followed by genotyping for the *atg7* or atg5
mutations to identify plants that were wild-type or homozygous for the *atg* mutations.
166
Disrupting autophagy breaks down the initial self-incompatible response of blocking pollen
hydration
To assess the early stages of pollen rejection, the pollen hydration checkpoint was first
investigated. Self-incompatible pollen from the parental SI-Col-0 or SI-C24 transgenic lines
were used to pollinate pistils from the respective F_2 progeny segregating for the *atg7* or *atg5*
mutations, and pollen hydration assays were carried out by measuring the increase in pollen
diameter at 10 min post-pollination (Lee et al., 2020). Positive controls of the SI-Col-0 and SI-
C24 pollen placed on compatible wildtype Col-0 and C24 pistils, respectively, resulted in a large
increase in diameter as a result of pollen hydration (Fig. 1A, B). In contrast, self-pollen placed on
the SI-Col-0 and SI-C24 pistils resulted in very little pollen hydration at 10-minutes post-
pollination (Fig. 1A, B). This lack of hydration was consistent between the SI-Col-0 and SI-C24
lines and indicative of the early recognition and rejection of self-incompatible pollen.
Importantly, stigmas from the *atg7-2, atg5-1 and atg7-9* mutants showed normal SI-Col-0 and
SI-C24 pollen hydration, respectively, indicating that disruption in the autophagic process did
not perturb the compatible post-pollination responses (Fig. 1A, B).
For the F_2 progeny from the SI-Col-0 x *atg7-2* cross, the SI-Col-0 plants homozygous for
wildtype *ATG7* showed the same small increase in pollen hydration as the control self-
incompatible pollinations at 10 min post-pollination. In contrast, when SI-Col-0 pollen was

applied to the SI-Col-0 stigmas that were homozygous for the *atg7-2* mutation, an increase in

pollen hydration was observed, indicating an incomplete rejection of the self-incompatible pollen

(Fig. 1A). Similarly, pollen hydration assays on the F_2 progeny from the SI-Col-0 x *atg5-1* cross

showed that hydration of SI-Col-0 pollen on SI-Col-0 stigmas wild-type for *ATG5* displayed a

small increase while SI-Col-0 pollen hydration on SI-Col-0 stigmas homozygous for the *atg5-1*

mutation was significantly increased (Fig. 1B). A comparable set of pollen hydration

experiments on the F_2 progeny from the SI-C24 x *atg7-9* cross also showed a similar trend (Fig.

1C). SI-C24 stigmas that were wild-type for *ATG7* supported similar small increases in SI-C24

pollen hydration as the control self-incompatible pollinations, and the introduction of the *atg7-9*

mutation into this background resulted in increased hydration of the self-incompatible pollen.

Together, these results indicate that autophagy is required to fully prevent the hydration of self-

incompatible pollen.

Disruption of autophagy increases self-incompatible pollen tube penetration

With the observed increase in self-incompatible pollen hydration, the next stage of preventing

self-incompatible pollen germination and pollen tube penetration was investigated. To determine

if this block was affected by the *atg7* and *atg5* mutations, pollinated pistils were collected at 2-

and 24-hours post-pollination and stained with aniline blue to visualize pollen tubes. At 2-hours

post-pollination, SI-Col-0 and SI-C24 pollen was completely rejected by the respective self-

incompatible pistils, and no pollen tubes were visible (Supplemental Fig. S2 and S3). Although

we did not see any obvious difference at 2-hours post-pollination, we did observe an occasional

pollen tube that penetrated the stigma barrier of the SI-Col-0 and SI-C24 *atg* homozygous

mutants. To address whether we could observe a greater number of SI-Col-0 and SI-C24 pollen

tubes after a longer period, pollinated pistil samples were then harvested at 24-hours post-

pollination. A later time point following pollination is more reflective of the natural pollination

process and provides more time for the self-incompatible pollen to germinate and grow into the

pistil. Self-pollination of the SI-Col-0 and SI-C24 parental lines resulted in an absence of pollen

tubes (Fig. 2A, 3A and 4A) while pollinations with compatible wild-type pollen led to an

abundance of pollen tubes growing in the pistils (Fig. 2B, 3B and 4B). The results indicated that

the SI-Col-0 and SI-C24 were able to fully reject self-incompatible pollen and accept wild-type

compatible pollen at 24-hours post-pollination. The SI-Col-0 and SI-C24 pollen also produced

numerous pollen tubes when placed on the *atg* mutant pistils, indicating that the absence of

autophagy did not affect these compatible pollinations (Fig. 2C and 3C). The impact of the *atg*

mutations on self-incompatibility at 24-hours post-pollination was examined by applying the SI-

Col-0 and SI-C24 pollen on their respective F_2 progeny homozygous for the *atg* mutations.

When SI-Col-0 *atg7-2* stigmas or SI-Col-0 *atg5-1* stigmas were pollinated with SI-Col-0 pollen,

there was a general increase in the number of pollen grains adhering to the stigma and forming

pollen tubes when compared to the SI-Col-0 pistils (Fig. 2D-F and 3D-F). The pollinations were

repeated for the SI-C24 F_2 plants, and again the SI-C24 *atg7-9* pistils did show some SI-C24

pollen adhering to the stigmatic surface and successfully growing pollen tubes through the pistil

while the SI-C24 *ATG7* pistils rejected the SI-C24 pollen (Fig. 4D-F). The number of pollen

tubes were counted in the SI-Col-0 and SI-C24 pistils that were wild-type or homozygous for the

atg mutations, and a significant increase in the number of pollen tubes were counted in the

homozygous *atg7* and *atg5* mutants (Fig. 2F, 3F and 4F). Thus, the 24-hours post-pollination

samples demonstrated that autophagy is required for fully blocking self-incompatible pollen

germination and pollen tube growth through the self-incompatible pistil.

Loss of autophagy increases seed set in the self-incompatible lines

With the observation that self-incompatibility is partially lost in the SI-Col-0 *atg7-2*, SI-Col-0

atg5-1 and SI-C24 *atg7-9* mutant pistils, resulting in more pollen tubes growing through the

pistils, the impact of this on seed set was assessed. Manual pollinations were carried out using

SI-Col-0 and SI-C24 pollen on pistils from the respective parental lines and F_2 progeny

homozygous for the *atg7* or *atg5* mutations, and 2-week old siliques were harvested for seed

counting (Fig. 5; Supplemental Fig. S5). As expected for compatible pollinations in Col-0, C24,

and the *atg* mutants, abundant seeds/silique were produced (Fig. 5A-C). As well, self-

pollinations of SI-Col-0 and SI-C24 pistils resulted in a very limited seed yield consistent with

the rejection of self-incompatible pollen (Fig. 5A-C). When the SI-Col-0 *atg7-2* pistils were

pollinated with SI-Col-0 pollen, there was a significant increase in yield to 27 seeds/silique

compared to 12.8 seeds/silique for SI-Col-0 indicating an incomplete rejection of the SI-Col-0

pollen by the SI-Col-0 *atg7-2* pistils (Fig. 5A). Similarly, when SI-Col-0 *atg5-1* pistils were

pollinated with SI-Col-0 pollen, there was an increased average yield of 31.9 seeds/silique

compared to 10.8 seeds/silique for SI-Col-0 (Fig. 4B). Finally, similar trends in seed yields were

observed in the C24 accession. Self-incompatible pollinations with SI-C24 pistils resulted in a

low average of 6.8 seeds/silique as expected yet when the SI-C24 *atg7-9* pistils were pollinated

with SI-C24 pollen, there was a significant increase in seed yields to17.2 seeds/silique (Fig. 5C).

Altogether, these results indicate that loss of autophagy was sufficient to disrupt the self-

incompatibility pathway allowing for the growth of more SI pollen tubes through the SI-Col-0

and SI-C24 pistils carrying *atg7* or *atg5* mutations to initiate fertilization and seed production,

and thus, confirming that autophagy is required for full rejection of self-pollen.

Autophagosomes are produced following self-pollination in the SI-C24 line

Previously, autophagosomes were detected by TEM following self-incompatible pollinations in

A. lyrata and in the transgenic self-incompatible *A. thaliana* in the Col-0 accession transformed

with *A. lyrata* S-locus genes (Safavian and Goring, 2013; Indriolo et al., 2014). To confirm that

autophagosomes are also forming in the SI-Col-0 *Ah-SCR13 Ah-SRK13 Ah-ARC1* #2 line used in

this study, *GFP-ATG8a* was used as an autophagosome marker (Thompson et al., 2005). Col-0

plants carrying the *ProUBQ10:GFP-ATG8a* construct (Kim et al., 2013) were crossed with the

SI-Col-0 line, and the subsequent F_1 generation was used for imaging. The SI-Col-0 *GFP-*

ATG8a stigmas were imaged as unpollinated or manually pollinated with SI-Col-0 pollen. At 15-

minutes post-pollination, many GFP punctae were observed in the SI-Col-0 *GFP-ATG8a*

stigmatic papillae contacting SI-Col-0 pollen (Fig. 6A). Conversely, few puncta were observed

in unpollinated SI-Col-0 GFP-ATG8a stigmatic papillae, consistent with a low background level

of autophagy (Fig. 6C). The number of puncta were counted for both SI-Col-0 pollinated and

unpollinated SI-Col-0 GFP-ATG8a stigmatic papillae (Fig. 6E), and there was a significant

increase in the number of GFP-ATG8a-labelled punctae in the SI-pollinated SI-Col-0 *GFP-*

ATG8a stigmatic papillae indicated a strong induction of autophagy. These observations confirm

our previous findings that autophagy is activated in the stigmatic papillae with self-incompatible

pollinations when compared to an unpollinated stigma (Safavian and Goring, 2013; Indriolo and

Goring, 2014).

Discussion

Self-incompatibility plays an important regulatory role in many outcrossing Brassicaceae species

where producing inbred progeny would be disadvantageous (Shimizu and Tsuchimatsu, 2015;

Durand et al., 2020). Previous studies have uncovered many important steps in the self-

incompatibility pathway, but there are still outstanding questions on the cellular mechanisms by

which self-pollen is rejected (reviewed in Doucet et al., 2016; Goring, 2017; Jany et al., 2019). In

previous studies, we observed the presence of autophagosomes in the cytoplasm and autophagic

bodies in the vacuole with self-incompatible pollinations in self-incompatible *A. lyrata* and

transgenic self-incompatible *A. thaliana* Col-0 (Safavian and Goring, 2013; Indriolo and Goring,

2014). What was not known, however, was whether autophagy was necessary for self-

incompatible pollen rejection. In this study, we addressed this question by crossing two different

autophagy-deficient mutants (*atg7* and *atg5*) into two different transgenic self-incompatible *A.*

thaliana lines, and demonstrated that a functioning autophagy system is required in the stigma

for the full rejection of self-incompatible pollen. It is important to note that disrupting autophagy

did not cause a full loss of self-incompatibility indicating that other cellular responses are

required in conjunction with autophagy for self-incompatible pollen rejection. Importantly, this

partial breakdown in the stigma was seen for all stages from increased pollen hydration to more

pollen tubes growing through pistil to increase seed yields. Finally, the requirement of autophagy

in the rejection of self-incompatible pollen was consistent between the three different autophagy

deficient mutants *(atg7-2, atg7-9* and *atg5)* and the two different *Arabidopsis thaliana*

accessions (Col-0 and C24) used in the SI lines indicating that there is a shared mechanism for

self-incompatible pollen rejection.

The rationale for using both the SI-Col-0 and SI-C24 lines was to determine if there were any

accession- or transgene-specific associations with autophagy. The SI-Col-0 carries the *A. halleri*

SCR13, *SRK13* and *ARC1* transgenes (Zhang et al., 2019) while the SI-C24 line carries the *A.*

lyrata SCRb and *SRKb* transgenes (Iwano et al., 2015). In addition to two different S-haplotypes

from two different *Arabidopsis* species, the SI-Col-0 line also carries the *A. halleri ARC1*

transgene. The fact that both accessions require an intact autophagy system in the stigma for full

self-incompatible pollen rejection indicates that autophagy activation does not require ARC1,

though the potential involvement of another related PUB E3 ligase cannot be ruled out (Indriolo

et al., 2012; Indriolo and Goring, 2016). Our results indicate that autophagy is a clear

commonality in the self-incompatibility response between these two accessions, although

variation may exist in the execution of this cellular function. The question then arises to the

purpose of autophagy activation in self-incompatibility.

Self-incompatible pollen carries the signals to be recognized as both self-incompatible pollen and

compatible pollen, and it has been hypothesized that self-incompatibility blocks compatible

pollen responses in the stigma (Dickinson, 1995). With the working model that secretory activity

in the stigma is required for compatible pollen acceptance (reviewed in Dickinson, 1995; Doucet

et al., 2016; Goring, 2017), several of the self-incompatibility cellular responses such as

EX070A1 inhibition (Samuel et al., 2009; Sankaranarayanan et al., 2015), a strong and rapid

calcium influx (Iwano et al., 2015), and the absence of actin focalization (Rozier et al., 2020) are

consistent with blocking secretion (reviewed in Goring, 2017). Autophagy could also be an

additional step to clear any secretory activity that was initially activated. Another possibility is

that autophagy is resetting the cellular program to reject self-pollen. (Rodriguez et al., 2020)

recently demonstrated a role for autophagy in switching *A. thaliana* cell fate determination

following treatments with different signals. Autophagy was shown to be required for this switch

and was proposed to "clean-up" the cell to allow for the new cellular program to be established.

In the context of self-incompatibility, autophagy activation in the stigma could be linked to

reprogramming the cells from the default compatible pollen responses to the self-pollen rejection

pathway.

In conclusion, this study now firmly places autophagy as part of the cellular mechanisms leading

to self-incompatible pollen rejection. Future studies will need to focus on how autophagy is

activated in the self-incompatibility pathway and whether there are specific cytoplasmic

components targeted for autophagic destruction. Understanding the mechanisms underlying self-

incompatibility is important as seed yield of many important crop species is directly tied to their

reproductive success, and the ability to modulate selfing would be a powerful tool in increasing

yield.

Materials and methods

Plant materials and growth conditions

All *Arabidopsis thaliana* seeds (Col-0, C24, transgenic lines, T-DNA insertion mutants and

CRISPR/Cas9 deletion mutants) were sterilized and cold stratified at 4°C for 3 days. Seedlings

were germinated on ½ Murashige and Skoog (MS) medium plates supplemented with 0.4 (w/v)

phytoagar (pH 5.8) and 1% (w/v) sucrose. Seedlings were grown at 22°C under 16h light for 7

days, and then transferred to Sunshine #1 soil supplemented with Plant Prod All Purpose

Fertilizer 20-20-20 (1g/L water). All plants were grown in chambers under a 16-hour light/8-

hour dark growth cycle at 22°C, with humidity kept under 50%.

For the Col-0 accession experiments, the *Ah-SCR13 Ah-SRK13 Ah-ARC1* #2 Col-0 line (Zhang et

al., 2019) was crossed to the *atg7-2* mutant (Col-0 accession; Chung et al., 2010) and *atg5-1*

mutant (Thompson et al., 2005). For the C24 accession experiments, the *SI-C24 #15-1* line (with

Al-SCRb AlSRKb; Iwano et al., 2015) was crossed to the *atg7-9* mutant (C24 accession; see

below). All transgenic lines were genotyped using primers found in Supplemental Table 1. For

the C24 accession, plants were genotyped for *AlSCR$_b$*, *AlSRK$_b$* and *atg7-9*. Progeny were

analyzed in the F$_2$ generation for all three crosses. For SI-Col-0 *GFP-ATG8a*, the *Ah-SCR13 Ah-*

SRK13 Ah-ARC1 #2 Col-0 line (Zhang et al., 2019), was crossed into the *ProUBQ10:GFP-*

ATG8a line (Col-0 accession; Kim et al., 2013), and the resulting F$_1$ progeny was used for all

imaging.

Plasmid construction and plant transformation

The *atg7-9* CRISPR/Cas9 deletion mutation was generated in C24 using previously described

methods in the Col-0 accession (Doucet et al., 2019; Doucet et al., 2019). The final pBEE401E

vector contained two guide RNAs targeting exons 3 and 8 of *ATG7* (Supplemental Fig. S1).

Arabidopsis thaliana C24 plants were transformed using floral dip (Clough and Bent, 1998), and

plants were dipped twice, 3 days apart, to maximize transformation efficiency. T1 seeds were

collected and germinated on soil, and one-week old seedlings were sprayed with BASTA

herbicide to select for transformants. PCR was used to confirm transformants (BASTA marker)

and the presence of *atg7* mutations (deletions of the exon 3-8 region; Supplemental Figure S1).

These *atg7* deletion mutants were then verified through sequencing of regions flanking the

deletions. The *atg7* mutants showed a wild-type plant and flower morphology, along with the

shorter life span typically seen in *atg* mutants (Thompson et al., 2005; Avila-Ospina et al., 2014).

One homozygous mutant, *atg7-9*, from the T2 generation was crossed to the *SI-C24 #15-1* line,

and all data was collected in the F$_2$ generation of this cross (F$_4$ generation for *atg7-9*).

Confocal Microscopy

A. thaliana SI-Col-0 plants carrying the *ProUBQ10:GFP-ATG8a* construct were manually

pollinated using pollen from the self-incompatible SI-Col-0 line. Pistils were removed at 10-

minutes post-pollination and images of autophagosomes were taken at 15-minutes post-

pollination. All images were taken using a Leica SP8 confocal microscope, with standard GFP

excitation-emission wavelengths (488nm excitation, 495-545nm emission) with a pinhole

diameter of 2.0AU. Leica LAS AF lite in combination with ImageJ was used for image

processing and puncta counting. GFP punctae were counted in stigmatic papillae cell areas

directly adjacent to SI-Col-0 pollen, and an equivalent cell area in unpollinated samples.

Assays for Pollen hydration, Aniline blue staining, and Seed set

All post-pollination assays used manual pollinations and were conducted as described in Lee et

al., 2020). For the SI-C24 plants, late stage 12 pistils were first emasculated and wrapped in

plastic wrap overnight prior to pollination. For the SI-Col-0 plants, the pistils did not require

emasculation due to the approach herkogamy phenotype (Zhang et al. 2019*)*, and freshly opened

flowers with unpollinated pistils were selected for these assays. All pollinations were completed

at humidity levels under 50% for consistency. Pollen hydration assays were conducted by

removing and mounting pistils on ½ MS, and lightly pollinating the stigmatic surface using a

single anther. Images were taken at 0- and 10-min post-pollination using a Nikon sMz800

microscope, and pollen grains on the stigmatic surface were randomly selected for measurements

using the NIS-elements imaging software. For the aniline blue staining, one anther was used to

lightly pollinate each pistil, and the pistils were then collected at 2- and 24-hours post-pollination

for staining as described in Lee et al., 2020). The aniline blue-stained pistils were mounted on a

slide with sterile water, flattened, and imaged using a Zeiss Axioscope2 Plus fluorescent

microscope. For seeds counts, pistils were manually pollinated with a single anther and were left

for 10-14 days. The maturing siliques were then removed and cleared in 70% (v/v) ethanol for 3-

5 days, or until seeds were visible (Beuder et al., 2020). Siliques were then mounted on a dry

slide with the septum facing upwards and imaged using a Nikon sMz800 microscope, and seeds

were counted using the NIS-elements imaging software.

Supplemental Material

Supplemental Figure S1. Generation of the *ATG7* CRISPR/Cas9 deletion mutation in the C24 accession.

Supplemental Figure S2. Representative images of flowering plants for the SI-C24 and *atg7-9* mutant plants in the C24 accession.

Supplemental Figure S3. Representative images of aniline blue-stained pistils at 2 hrs following pollination with SI-Col-0 or Col-0 pollen.

Supplemental Figure S4. Representative images of aniline blue-stained pistils at 2 hrs following pollination with SI-C24 or C24 pollen.

Supplemental Figure S5. Representative images of cleared siliques used in seed counting for control and SI plants with the *atg7* mutation.

Supplemental Table 1. Primer Sequences.

Acknowledgements

We thank Laura Canales and Arjun Sharma for technical assistance, and members of the Goring lab for critically reading this article. We are very grateful to Seiji Takayama (Nara Institute of Science and Technology; University of Tokyo) for providing the *SI-C24 #15-1* seeds (*A. thaliana* C24), Richard Vierstra (Washington University in St Louis) and Peter Bozhkov (Swedish University of Agricultural Sciences) for providing *atg7-2* and *atg5-1* mutant seeds (*A. thaliana* Col-0), and Marisa Otegui (University of Wisconsin-Madison) for providing the *ProUBQ10:GFP-ATG8a* seeds (*A. thaliana* Col-0).

Figure 1. Pollen hydration assays on control and SI F$_2$ progeny segregating for the *atg7* and

***atg5* mutations.**

Pollen hydration assays for **A,** SI-Col-0 pollen grains placed on stigmas from Col-0, SI-Col-0,

atg7-2, and F$_2$ progeny from the SI-Col-0 x *atg7-2* cross; **B,** SI-Col-0 pollen grains placed on

stigmas from Col-0, SI-Col-0, *atg5-1*, and F$_2$ progeny from the SI-Col-0 x *atg5-1* cross; and **C,**

SI-C24 pollen grains placed on stigmas from C24, SI-C24, *atg7-9*, and F$_2$ progeny stigmas from

the SI-C24 x *atg7-9* cross. Pollen hydration was measured by taking pollen grain diameters at 0-

and 10-minutes post-pollination. Data is shown as Mean +/- SE with all data points displayed.

n=30 pollen grains per line, P<0.05 (One-way ANOVA with Tukey-HSD post-hoc test).

Figure 2. Representative images of aniline blue-stained SI-Col-0 or SI-Col-0 *atg7-2* pistils at

24 hrs following pollination.

A-E, All stigmas were hand-pollinated with SI-Col-0 pollen or wild-type Col-0 pollen as

indicated and collected at 24 hours post-pollination for aniline blue staining. The genotypes of

the pistils are indicated for each panel. Pollen tubes penetrating through the stigma and style

indicate a breakdown of self-incompatibility as seen in **E**. For each panel, top image =

brightfield, bottom image = aniline blue. Scale bar = 100μm

F, Bar graph showing the mean number of pollen tubes at 24 hrs post-pollination. Data is shown

as Mean +/- SE with all data points displayed. n = 11-12 pistils per line. At 24 hours, pistils

from SI-Col-0 *atg7-2* mutants allowed a significantly higher number of SI-Col-0 pollen tubes

compared to the SI-Col-0 pistils (P<0.05).

Figure 3. Representative images of aniline blue-stained SI-Col-0 or SI-Col-0 *atg5-1* pistils at

24 hrs following pollination.

A-E, All stigmas were hand-pollinated with SI-Col-0 pollen or wild-type Col-0 pollen as

indicated and collected at 24 hours post-pollination for aniline blue staining. The genotypes of

the pistils are indicated for each panel. Pollen tubes penetrating through the stigma and style

indicate a breakdown of self-incompatibility as seen in **E**. For each panel, top image =

brightfield, bottom image = aniline blue. Scale bar = 100μm

F, Bar graph showing the mean number of pollen tubes at 24 hrs post-pollination. Data is shown

as Mean +/- SE with all data points displayed. n = 10 pistils per line. At 24 hours, pistils from

SI-Col-0 *atg5-1* mutants allowed a significantly higher number of SI-Col-0 pollen tubes

compared to the SI-Col-0 pistils (P<0.05).

Figure 4. Representative images of aniline blue-stained SI-C24 or SI-C24 *atg7-9* pistils at 24

hrs following pollination.

A-E, All stigmas were hand-pollinated with SI-C24 pollen or wild-type C24 pollen as indicated

and collected at 24 hours post-pollination for aniline blue staining. The genotypes of the pistils

are indicated for each panel. Pollen tubes penetrating through the stigma and style indicate a

breakdown of self-incompatibility as seen in **E**. For each panel, top image = brightfield, bottom

image = aniline blue. Scale bar = 100µm

F, Bar graph showing the mean number of pollen tubes at 24 hrs post-pollination. Data is shown

as Mean +/- SE with all data points displayed. n = 8-10 pistils per line. At 24 hours, pistils from

SI-C24 *atg7-9* mutants allowed a significantly higher number of SI-C24 pollen tubes compared

to the SI-C24 pistils (P<0.05).

Figure 5. Seeds per silique counts for control and SI F$_2$ progeny with the *atg7* or *atg5*

mutations.

Seeds per silique counts following manual pollination with **A, B,** SI-Col-0 pollen or **C,** SI-C24

pollen on pistils with the indicated genotypes. Siliques were harvested at 2-weeks post-

pollination and cleared for seeds counts. Data is shown as Mean +/- SE with all data points

displayed. n= 8-10 siliques per line. Compared to the control SI pollinations, significant

increases in the number of seeds per silique were observed for the SI plants that were

homozygous for the *atg7* or *atg5* mutations, P<0.05 (One-way ANOVA with Tukey-HSD post-

hoc test).

Figure 6. Autophagosomes are more abundant in SI-Col-0 *GFP-ATG8a* papillae following

pollination with SI-Col-0 pollen.

More autophagosomes marked by GFP-ATG8a are visible at 15-minutes post-pollination in the

SI-Col-0 *GFP-ATG8a* papillae (**A-B**) compared to unpollinated SI-Col-0 *GFP-ATG8a* papillae

(**C-D**). Scale bar = 10µm.

E, Bar graph showing the mean number of autophagosomes present in each sample. Data is

shown as Mean +/- SE with all data points displayed. n=14 papillae. At 15 min post-pollination

with SI-Col-0 pollen, there is a significant increase in the number of GFP-labelled punctae in the

SI-Col-0 *GFP-ATG8a* papillae ($P<0.05$). The large punctae are chloroplasts with

autofluorescence and were not counted.

Literature Cited

Adhikari PB, Liu X, Wu X, Zhu S, Kasahara RD (2020) Fertilization in flowering plants: an odyssey of sperm cell delivery. Plant Mol Biol 103: 9-32

Avila-Ospina L, Moison M, Yoshimoto K, Masclaux-Daubresse C (2014) Autophagy, plant senescence, and nutrient recycling. J Exp Bot 65: 3799-3811

Beuder S, Dorchak A, Bhide A, Moeller SR, Petersen BL, MacAlister CA (2020) Exocyst mutants suppress pollen tube growth and cell wall structural defects of hydroxyproline O-arabinosyltransferase mutants. Plant J

Boggs NA, Nasrallah JB, Nasrallah ME (2009) Independent S-Locus Mutations Caused Self-Fertility in Arabidopsis thaliana. Plos Genetics 5

Chung T, Phillips AR, Vierstra RD (2010) ATG8 lipidation and ATG8-mediated autophagy in Arabidopsis require ATG12 expressed from the differentially controlled ATG12A AND ATG12B loci. Plant J 62: 483-493

Clough SJ, Bent AF (1998) Floral dip: A simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. Plant Journal 16: 735-743

Dickinson H (1995) Dry stigmas, water and self-incompatibility in Brassica. Sexual Plant Reproduction 8: 1-10

Ding X, Zhang X, Otegui MS (2018) Plant autophagy: new flavors on the menu. Curr Opin Plant Biol 46: 113-121

Doucet J, Lee HK, Goring DR (2016) Pollen acceptance or rejection: a tale of two pathways. Trends in plant science 21: 1058-1067

Doucet J, Lee HK, Udugama N, Xu J, Qi B, Goring DR (2019) Investigations into a putative role for the novel BRASSIKIN pseudokinases in compatible pollen-stigma interactions in Arabidopsis thaliana. BMC Plant Biol 19: 549

Doucet J, Truong C, Frank-Webb E, Lee HK, Daneva A, Gao Z, Nowack MK, Goring DR (2019) Identification of a role for an E6-like 1 gene in early pollen-stigma interactions in Arabidopsis thaliana. Plant Reprod 32: 307-322

Durand E, Chantreau M, Le Veve A, Stetsenko R, Dubin M, Genete M, Llaurens V, Poux C, Roux C, Billiard S, Vekemans X, Castric V (2020) Evolution of self-incompatibility in the Brassicaceae: Lessons from a textbook example of natural selection. Evol Appl 13: 1279-1297

Elleman C, Dickinson H (1996) Identification of pollen of components regulating the responses
in stigmatic Brassica oleracea. New Phytologist 133: 197-205

Fujii S, Shimosato-Asano H, Kakita M, Kitanishi T, Iwano M, Takayama S (2020) Parallel
evolution of dominant pistil-side self-incompatibility suppressors in Arabidopsis. Nat
Commun 11: 1404

Goring DR (2017) Exocyst, exosomes, and autophagy in the regulation of Brassicaceae pollen-
stigma interactions. J Exp Bot 69: 69-78

Goubet PM, Berges H, Bellec A, Prat E, Helmstetter N, Mangenot S, Gallina S, Holl AC, Fobis-
Loisy I, Vekemans X, Castric V (2012) Contrasted patterns of molecular evolution in
dominant and recessive self-incompatibility haplotypes in Arabidopsis. PLoS Genet 8:
e1002495

Gu T, Mazzurco M, Sulaman W, Matias DD, Goring DR (1998) Binding of an arm repeat
protein to the kinase domain of the *S*-locus receptor kinase. Proceedings of
the National Academy of Sciences 95: 382-387

Guo YL, Zhao X, Lanz C, Weigel D (2011) Evolution of the S-locus region in Arabidopsis
relatives. Plant Physiol 157: 937-946

Indriolo E, Goring DR (2014) A conserved role for the ARC1 E3 ligase in Brassicaceae self-
incompatibility. Front Plant Sci 5: 181

Indriolo E, Goring DR (2016) Yeast two-hybrid interactions between Arabidopsis lyrata S
Receptor Kinase and the ARC1 E3 ligase. Plant Signal Behav 11: e1188233

Indriolo E, Safavian D, Goring DR (2014) The ARC1 E3 ligase promotes two different self-
pollen avoidance traits in Arabidopsis. The Plant Cell 26: 1525-1543

Indriolo E, Tharmapalan P, Wright SI, Goring DR (2012) The ARC1 E3 ligase gene is frequently
deleted in self-compatible Brassicaceae species and has a conserved role in Arabidopsis
lyrata self-pollen rejection. Plant Cell 24: 4607-4620

Iwano M, Igarashi M, Tarutani Y, Kaothien-Nakayama P, Nakayama H, Moriyama H, Yakabe
R, Entani T, Shimosato-Asano H, Ueki M, Tamiya G, Takayama S (2014) A Pollen
Coat–Inducible Autoinhibited Ca^{2+}-ATPase Expressed in Stigmatic Papilla
Cells Is Required for Compatible Pollination in the Brassicaceae. The Plant Cell 26: 636-
649

Iwano M, Ito K, Fujii S, Kakita M, Asano-Shimosato H, Igarashi M, Kaothien-Nakayama P,
Entani T, Kanatani A, Takehisa M, Tanaka M, Komatsu K, Shiba H, Nagai T, Miyawaki
A, Isogai A, Takayama S (2015) Calcium signalling mediates self-incompatibility
response in the Brassicaceae. Nat Plants 1: 15128

Jany E, Nelles H, Goring DR (2019) The Molecular and Cellular Regulation of Brassicaceae
Self-Incompatibility and Self-Pollen Rejection, Vol 343. Elsevier Ltd

Johnson MA, Harper JF, Palanivelu R (2019) A Fruitful Journey: Pollen Tube Navigation from
Germination to Fertilization. Annual review of plant biology 70: 809-837

Kim J, Lee H, Lee HN, Kim SH, Shin KD, Chung T (2013) Autophagy-related proteins are
required for degradation of peroxisomes in Arabidopsis hypocotyls during seedling
growth. Plant Cell 25: 4956-4966

Kusaba M, Dwyer K, Hendershot J, Vrebalov J, Nasrallah JB, Nasrallah ME (2001) Self-
Incompatibility in the Genus Arabidopsis: Characterization of the S Locus in the
Outcrossing A. Lyrata and Its Autogamous Relative A. Thaliana. The Plant Cell 13: 627-
644

Lee HK, Macgregor S, Goring DR (2020) A Toolkit for Teasing Apart the Early Stages of
Pollen-Stigma Interactions in Arabidopsis thaliana. Methods Mol Biol 2160: 13-28

Mizuta Y, Higashiyama T (2018) Chemical signaling for pollen tube guidance at a glance.
Journal of Cell Science 131

Nasrallah JB (2019) Self-incompatibility in the Brassicaceae: Regulation and mechanism of self-
recognition. Curr Top Dev Biol 131: 435-452

Nasrallah ME, Liu P, Nasrallah JB (2002) Generation of self-incompatible Arabidopsis thaliana
by transfer of two S locus genes from A. lyrata. Science 297: 247-249

Nasrallah ME, Liu P, Sherman-Broyles S, Boggs NA, Nasrallah JB (2004) Natural variation in
expression of self-incompatibility in Arabidopsis thaliana: Implications for the evolution
of selfing. Proceedings of the National Academy of Sciences of the United States of
America 101: 16070-16074

Palanivelu R, Tsukamoto T (2012) Pathfinding in angiosperm reproduction: pollen tube guidance
by pistils ensures successful double fertilization. Wiley Interdisciplinary Reviews:
Developmental Biology 1: 96-113

Rodriguez E, Chevalier J, Olsen J, Ansbol J, Kapousidou V, Zuo Z, Svenning S, Loefke C,
Koemeda S, Drozdowskyj PS, Jez J, Durnberger G, Kuenzl F, Schutzbier M, Mechtler K,
Ebstrup EN, Lolle S, Dagdas Y, Petersen M (2020) Autophagy mediates temporary
reprogramming and dedifferentiation in plant somatic cells. EMBO J 39: e103315
Rozier F, Riglet L, Kodera C, Bayle V, Durand E, Schnabel J, Gaude T, Fobis-Loisy I (2020)
Live-cell imaging of early events following pollen perception in self-incompatible
Arabidopsis sthaliana. J Exp Bot
Safavian D, Goring DR (2013) Secretory activity is rapidly induced in stigmatic papillae by
compatible pollen, but inhibited for self-incompatible pollen in the brassicaceae. PLoS
ONE 8
Safavian D, Zayed Y, Indriolo E, Chapman L, Ahmed A, Goring DR (2015) RNA silencing of
exocyst genes in the stigma impairs the acceptance of compatible pollen in Arabidopsis.
Plant Physiology 169: 2526-2538
Samuel MA, Chong YT, Haasen KE, Aldea-Brydges MG, Stone SL, Goring DR (2009) Cellular
pathways regulating responses to compatible and self-incompatible pollen in Brassica and
Arabidopsis stigmas intersect at Exo70A1, a putative component of the exocyst complex.
Plant Cell 21: 2655-2671
Sankaranarayanan S, Jamshed M, Samuel MA (2015) Degradation of glyoxalase I in Brassica
napus stigma leads to self-incompatibility response. Nat Plants 1: 15185
Schierup MH, Mable BK, Awadalla P, Charlesworth D (2001) Identification and characterization
of a polymorphic receptor kinase gene linked to the self-incompatibility locus of
Arabidopsis lyrata. Genetics 158: 387-399
Shimizu KK, Shimizu-Inatsugi R, Tsuchimatsu T, Purugganan MD (2008) Independent origins
of self-compatibility in Arabidopsis thaliana. Mol Ecol 17: 704-714
Shimizu KK, Tsuchimatsu T (2015) Evolution of Selfing: Recurrent Patterns in Molecular
Adaptation. Annual Review of Ecology, Evolution, and Systematics, Vol 46 46: 593-622
Stone SL, Anderson EM, Mullen RT, Goring DR (2003) ARC1 is an E3 ubiquitin ligase and
promotes the ubiquitination of proteins during the rejection of self-incompatible Brassica
pollen. The Plant Cell 15: 885-898
Stone SL, Arnoldo M, Goring DR (1999) A breakdown of Brassica self-incompatibility in ARC1
antisense transgenic plants. Science 286: 1729-1731

Thompson AR, Doelling JH, Suttangkakul A, Vierstra RD (2005) Autophagic nutrient recycling

in Arabidopsis directed by the ATG8 and ATG12 conjugation pathways. Plant Physiol

138: 2097-2110

Tsuchimatsu T, Goubet PM, Gallina S, Holl AC, Fobis-Loisy I, Berges H, Marande W, Prat E,

Meng D, Long Q, Platzer A, Nordborg M, Vekemans X, Castric V (2017) Patterns of

Polymorphism at the Self-Incompatibility Locus in 1,083 Arabidopsis thaliana Genomes.

Mol Biol Evol 34: 1878-1889

Tsuchimatsu T, Suwabe K, Shimizu-Inatsugi R, Isokawa S, Pavlidis P, Stadler T, Suzuki G,

Takayama S, Watanabe M, Shimizu KK (2010) Evolution of self-compatibility in

Arabidopsis by a mutation in the male specificity gene. Nature 464: 1342-1346

Young PG, Passalacqua MJ, Chappell K, Llinas RJ, Bartel B (2019) A facile forward-genetic

screen for Arabidopsis autophagy mutants reveals twenty-one loss-of-function mutations

disrupting six ATG genes. Autophagy 15: 941-959

Zhang T, Zhou G, Goring DR, Liang X, Macgregor S, Dai C, Wen J, Yi B, Shen J, Tu J, Fu T,

Ma C (2019) Generation of Transgenic Self-Incompatible Arabidopsis thaliana Shows a

Genus-Specific Preference for Self-Incompatibility Genes. Plants (Basel, Switzerland) 8:

570

Figure 1. Pollen hydration assays on control and SI F2 progeny segregating for the *atg7* and *atg5* mutations.
Pollen hydration assays for **A,** SI-Col-0 pollen grains placed on stigmas from Col-0, SI-Col-0, *atg7-2*, and F2 progeny from the SI-Col-0 x *atg7-2* cross; **B,** SI-Col-0 pollen grains placed on stigmas from Col-0, SI-Col-0, *atg5-1*, and F2 progeny from the SI-Col-0 x *atg5-1* cross; and **C,** SI-C24 pollen grains placed on stigmas from C24, SI-C24, *atg7-9*, and F2 progeny stigmas from the SI-C24 x *atg7-9* cross.
Pollen hydration was measured by taking pollen grain diameters at 0- and 10-minutes post-pollination. Data is shown as Mean +/- SE with all data points displayed. n=30 pollen grains per line, P<0.05 (One-way ANOVA with Tukey-HSD post-hoc test).

Figure 2. Representative images of aniline blue-stained SI-Col-0 or SI-Col-0 *atg7-2* pistils at 24 hrs following pollination.

A-E, All stigmas were hand-pollinated with SI-Col-0 pollen or wild-type Col-0 pollen as indicated and collected at 24 hours post-pollination for aniline blue staining. The genotypes of the pistils are indicated for each panel. Pollen tubes penetrating through the stigma and style indicate a breakdown of self-incompatibility as seen in E. For each panel, top image = brightfield, bottom image = aniline blue. Scale bar = 100μm

F, Bar graph showing the mean number of pollen tubes at 24 hrs post-pollination. Data is shown as Mean +/- SE with all data points displayed. n = 11-12 pistils per line. At 24 hours, pistils from SI-Col-0 *atg7-2* mutants allowed a significantly higher number of SI-Col-0 pollen tubes compared to the SI-Col-0 pistils (P<0.05).

Figure 3. Representative images of aniline blue-stained SI-Col-0 or SI-Col-0 *atg5-1* pistils at 24 hrs following pollination.

A-E, All stigmas were hand-pollinated with SI-Col-0 pollen or wild-type Col-0 pollen as indicated and collected at 24 hours post-pollination for aniline blue staining. The genotypes of the pistils are indicated for each panel. Pollen tubes penetrating through the stigma and style indicate a breakdown of self-incompatibility as seen in E. For each panel, top image = brightfield, bottom image = aniline blue. Scale bar = 100μm

F, Bar graph showing the mean number of pollen tubes at 24 hrs post-pollination. Data is shown as Mean +/- SE with all data points displayed. n = 10 pistils per line. At 24 hours, pistils from SI-Col-0 *atg5-1* mutants allowed a significantly higher number of SI-Col-0 pollen tubes compared to the SI-Col-0 pistils (P<0.05).

Figure 4. Representative images of aniline blue-stained SI-C24 or SI-C24 *atg7-9* pistils at 24 hrs following pollination.

A-E, All stigmas were hand-pollinated with SI-C24 pollen or wild-type C24 pollen as indicated and collected at 24 hours post-pollination for aniline blue staining. The genotypes of the pistils are indicated for each panel. Pollen tubes penetrating through the stigma and style indicate a breakdown of self-incompatibility as seen in E. For each panel, top image = brightfield, bottom image = aniline blue. Scale bar = 100μm

F, Bar graph showing the mean number of pollen tubes at 24 hrs post-pollination. Data is shown as Mean +/- SE with all data points displayed. n = 8-10 pistils per line. At 24 hours, pistils from SI-C24 *atg7-9* mutants allowed a significantly higher number of SI-C24 pollen tubes compared to the SI-C24 pistils (P<0.05).

Figure 5. Seeds per silique counts for control and SI plants with the *atg7* or *atg5* mutations.
Seeds per silique counts following manual pollination with **A, B,** SI-Col-0 pollen or **C,** SI-C24 pollen on pistils with the indicated genotypes. Siliques were harvested at 2-weeks post-pollination and cleared for seeds counts. Data is shown as Mean +/- SE with all data points displayed. n= 8-10 siliques per line. Compared to the control SI pollinations, significant increases in the number of seeds per silique were observed for the SI plants that were homozyogous for the *atg7* or *atg5* mutations, P<0.05 (One-way ANOVA with Tukey-HSD post-hoc test).

Figure 6. Autophagosomes are more abundant in SI-Col-0 *GFP-ATG8a* papillae following pollination with SI-Col-0 pollen

More autophagosomes marked by GFP-ATG8a are visible at 15-minutes post-pollination in the SI-Col-0 *GFP-ATG8a* papillae (**A-B**) compared to unpollinated SI-Col-0 *GFP-ATG8a* papillae (**C-D**). Scale bar = 10µm.

E, Bar graph showing the mean number of autophagosomes present in each sample. Data is shown as Mean +/- SE with all data points displayed. n=14 papillae. At 15 min post-pollination with SI-Col-0 pollen, there is a significant increase in the number of GFP-labelled punctae in the SI-Col-0 GFP-ATG8a papillae (P<0.05). The large punctae are chloroplasts with autofluorescence and were not counted.