Original Article

Estimating fecundity and density dependence from mark-recapture data for

making population projections

Bilgecan Şen[1*](Orcid ID: 0000-0002-7393-1805) and H. Reşit Akçakaya[1] (Orcid ID: 0000-0002-

8679-5929)

[1]Department of Ecology and Evolution, Stony Brook University, Stony Brook, NY, 11794,

U.S.A.

[*]Corresponding Author: bilgecan.sen@gmail.com

Abstract

Forecasting changes in size and distributions of populations is at the forefront of

ecological sciences in the 21st century. Such forecasts require robust estimators of fecundity,

survival and density-dependence. While survival estimation is the main focus of mark-recapture

modelling, fecundity and density dependence are rarely the subject of these models. Here, we

demonstrate that these parameters can be simultaneously estimated in a Bayesian framework

using only robust design mark-recapture data. Using simulated capture histories, we show that

this framework (which we named CJS-pop) can estimate vital rates and their density dependence

with little bias. When CJS-pop is applied to capture history data from Brown Creeper (*Certhia*

americana), it provides estimates of fecundity that is expected from the breeding biology of this

species. Finally, we illustrate that density dependence, even when estimated with uncertainty in

the CJS-pop framework, regularizes population dynamics and reduces the frequent population

extinctions and explosions observed under density-independent models. While CJS-pop as a

whole is a useful addition to the current mark-recapture modelling toolbox, we argue that the

independent components of this framework in estimating fecundity and density dependence can

be integrated to other CJS frameworks, potentially creating models capable of population

projections.

Keywords: CJS-pop, Fecundity, Mark-recapture, Population projections, Density Dependence

Introduction

Mark-recapture data analysis is a staple in population ecology for estimating survival, abundance, and recruitment rates (Lebreton et al. 1992; Williams et al. 2002; Cooch and White 2016). More recently, mark-recapture methods have been extended to work in parallel with different types of data in frameworks such as integrated population models (IPMs; Schaub and Abadi 2011) and to estimate dispersal and animal movement in spatial capture-recapture analysis (Ergon and Gardner 2014; Schaub and Royle 2014). When considering the wide applicability of mark-recapture methods for estimating parameters related to population dynamics, rarity of two types of parameters in the mark-recapture literature stands out: 1) fecundity as defined in a population modelling setting; and 2) density dependence of survival and fecundity. If these parameters can be estimated from mark-recapture data, then this single data source can be used on its own to parameterize stage-structured population models.

Fecundity is a measure of breeding performance of a population and in the perspective of a PVA can be defined as per-capita number of offspring that survives to the next time step (Akçakaya et al. 1999). When no explicit data on fecundity are available, models such as Jolly-Seber and even some IPMs (for example, Ahrestani et al. 2017) estimate recruitment rate as the total number of individuals added to a population in a given time period (usually, immigrants and newborn individuals). Recruitment rate, in this regard, is not useful for making stage-structured population projections, because density dependence and stochasticity in fecundity cannot be explicitly accounted for in this parameter alone. A PVA that employs a stage-structured matrix model requires both the number of juveniles born per individual and the juvenile survival to be explicitly estimated (for example, Ryu et al., 2016).

Density dependence (DD) is the phenomenon of population vital rates being dependent

on population size. Along with exogenous factors, it is one of the main natural processes that

regulates population dynamics. There are examples of estimating the strength of density

dependence of vital rates from mark-recapture data. For example, Ryu et al. (2016) provided a

framework for estimation of fecundity and density dependence from robust design mark-

recapture data. However, their framework is a mixture of frequentist and Bayesian approaches of

mark-recapture models and it requires multiple models to be fit sequentially. As a result Ryu et

al. (2016)'s different model components do not inform one another during estimation, which

prevents making full use of the data at hand and propagating uncertainty in a hierarchical

manner.

Here, we present an update on Ryu et al. (2016)'s framework. We use both simulated and

real data to show that; 1) fecundity can be estimated alongside adult survival and capture

probability in a single Bayesian framework using only robust design mark-recapture data; 2)

estimating fecundity reduces uncertainty in juvenile survival estimates; and, 3) when these vital

rates estimates are combined with density dependence, resulting stage-structured population

models are useful for calculating conservation-relevant metrics.

Material and Methods

Model specifications

Two basic parameters are estimated in a standard CJS model (Cooch and White 2016):

survival and capture probability. Robust design mark-recapture data separates these two

processes to primary capture occasions (for example years), and secondary capture occasions

within primary occasions (for example months). Populations are assumed to be closed

(individuals don't die or leave the population) among secondary occasions within a primary

occasion. Capture probabilities are estimated for each secondary occasion, which can then be

used to estimate the population size for a given primary occasion. Populations are assumed to be

open among primary occasions, so individuals can leave the population or die. Survival is

estimated with information coming across primary occasions. Below, for simplicity in

presentation, we assume primary occasions are years and secondary occasions are months.

We denote $p_{x,k,t,h}$ as the monthly capture probability of a stage x individual at population

k, year t, and month h; where, $x = 1,2,3 \dots, X$; $k = 1,2,3\dots, K$; $t = 1,2,3\dots, T$; and $h =$

$1,2,3\dots, H$. We can use this monthly capture probability to calculate a yearly capture probability:

$$P_{x,k,t} = 1 - \prod_{h=1}^{H}\left(1 - p_{x,k,t,h}\right) \#(1)$$

83   Then, we can use the heuristic estimator of populations size with a correction for years

84   with no captures (Dail and Madsen 2011) to estimate the expected abundance of stage x

85   individuals at population k, year t:

$$N_{x,k,t} = \frac{n_{x,k,t}}{P_{x,k,t}} + \frac{\left(1 - P_{x,k,t}\right)}{P_{x,k,t}} \#(2)$$

86   Where, $n_{x,k,t}$ is the number of captured stage x individuals in population k and year t. Using the

87   expected abundance time series of each population we calculate a density index:

$$D_{k,t} = \frac{\sum_{x=1}^{X} N_{x,k,t}}{\frac{\sum_{t=1}^{T} \sum_{x=1}^{X} N_{x,k,t}}{T}} \#(3)$$

where, $D_{k,t}$ is the density index at population k, year t. The numerator in equation 3 is the total

abundance of population k at year t across all stages, and denominator is the average expected

total abundance across T years. D is an index for the deviation of population abundance in a

given year from the long term average population abundance and it can be considered as relative

population density. We use D as a covariate for estimating density dependence strength of

fecundity and survival.

All three parameters estimated by equations 1 to 3, which are yearly capture

probability, expected abundance, and density index, respectively, can be used to estimate

fecundity and its density dependence:

$$\log(F_{k,t}) = \theta + \zeta \cdot D_{k,(t-1)} + \omega_{k,t} \#(4a)$$

$$\omega_{k,t} \sim \text{Normal}(0, \sigma_f^2) \#(4b)$$

where, θ is the fecundity in log scale at 0 density; ζ is the change in fecundity in log scale with

one unit change in population density index; $\omega_{k,t}$ is the spatio-temporal random effect at

population k and time t; and σ_f^2 is the spatio-temporal variance of fecundity at log scale. We link

the fecundity estimate to the number of captured juveniles in a given year and population by

using expected abundances for adults calculated in equation 2. Below we present a simple case

for two stages where $x = 1$ are juveniles and $x = 2$ are adults:

$$N'_{1,k,t} = N_{2,k,t} \cdot F_{k,t} \#(5a)$$

$$n_{1,k,t} \sim Poisson(P_{1,k,t} \cdot N'_{1,k,t}) \#(5b)$$

The expected number of juveniles is estimated twice in this framework: once as a derived

variable $(N_{1,k,t})$ using the heuristic population size estimator, and a second time $(N'_{1,k,t})$ as a

function of density index and expected number of adults. We discuss this double estimation and

our choice of Poisson distribution in detail in discussion section below.

So far, we have only used captured number of juveniles and adults as a data source.

However, this form of a capture history is not enough to estimate parameters used in the above 5

equations. So, we link these 5 equations with a more typical CJS model where survival and

capture probability are used to model capture histories of individuals:

$$\text{logit}(\phi_{x,k,t}) = \alpha_x + \beta \cdot D_{k,t} + \epsilon_{k,t} \quad \#(6a)$$

$$\epsilon_t \sim \text{Normal}(0, \sigma_s^2) \quad \#(6b)$$

where, $\phi_{x,k,t}$ is the apparent survival probability of a stage x individual at population k and year

t; α_x is the survival probability of a stage x individual on logit scale at 0 density; β is the change

in survival in logit scale with one unit change in population density index; $\epsilon_{k,t}$ is the spatio-

temporal random effect at population k and time t; and σ_f^2 is the spatio-temporal variance of

survival at logit scale.

Apparent survival changes the latent states of individuals in a population between time

steps, from t to $t + 1$; this latent state indicates whether an individual is alive and in the

population ($Z = 1$), or it is dead or left the population ($Z = 0$). The latent state of the ith

individual, then, is determined by its state at time t and its survival to time $t + 1$.

$$Z_{i,(t+1)} \sim \text{Bernoulli}\left(Z_{i,t} \cdot \phi_{(S_{i,t}),(g_i),t}\right) \quad \#(7)$$

where, $\phi_{(S_{i,t}),(g_i),t}$ is the apparent survival probability at the breeding stage and population of the

ith individual from year t to $t + 1$; $S_{i,t}$ is a matrix indicating the breeding stage of the ith

individual in year t, and the vector g_i indicates the population that the ith individual is in. Latent

state, Z, also determines the potential capture of an individual; dead ones cannot be captured.

Hence, every element of the capture history, $y_{i,t,h}$ (1 if an individual is captured, 0 if it is not), is

a Bernoulli random variable with a monthly capture probability, $p_{(S_{i,t}),(g_i),t,h}$, conditional on the

individual being alive and in the population, $Z_{i,t}$.

$$y_{i,t,h} \sim \text{Bernoulli}\left(Z_{i,t} \cdot p_{(S_{i,t}),(g_i),t,h}\right) \quad \#(8)$$

For simplicity in referring to this framework, we named it CJS-pop; pop extension comes from

the fact that it can estimate necessary parameters to build a stage-based population model:

Fecundity, staged-based survival, density dependence, and process variance.

Simulated data

We simulated several sets of capture histories in order to test CJS-pop's ability to

correctly retrieve true parameter values, and to uncover any inherent biases, especially when

quantifying density dependence strength. We set up a simulation scheme where we explored the

effect of sample size on quantifying density dependence strength. We set the time series length to

17 years, which is the maximum time series length for Brown Creeper data set we are using (see

below). We simulated three cases with 1, 5, and 10 populations, and three carrying capacities

(which controls population size): 50, 100, 150. For each combination of the number of

populations and carrying capacity, we generated capture history data sets using weak, moderate

and strong density dependence on survival and fecundity, which created 27 separate simulation

sets. For each simulation set, we generated 56 capture history data sets and fitted CJS-pop to

each one. See Appendix S2 for more detailed discussion of the simulation framework and

Appendix S3 for its code.

Empirical data: Brown Creeper

We applied CJS-pop to Brown Creeper (*Certhia americana*) data obtained from the

Mapping Avian Productivity and Survivorship (MAPS) program. Brown Creeper is a widespread

North American songbird species. We treated each MAPS location (a cluster of mist-netting and

banding stations) of Brown Creeper to be a separate population. We only included data from

populations which were located in the contiguous U.S., and which have been monitored for at

least 5 years. This resulted in a data set with 2931 individuals. We categorized any individual in

its first year as a juvenile (MAPS age codes 2 and 4), and older individuals as adults (MAPS age

codes 1, 5, 6, 7, and 8).

We made several adjustments and additions to basic CJS-pop framework presented above

when applying to Brown Creeper data. First, we accounted for potentially transient individuals in

the data set. In a CJS model, estimated survival rates are said to be "apparent" because the

estimated survival rate cannot distinguish between dead individuals and the ones that just left the

population. This can bias survival estimates to be lower than their true values. Accounting for

transients is a partial way to correct for this bias and it is a frequently used technique in CJS

literature (for example, Ahrestani et al. 2017).

Second, we used the priors and population modelling structure of CJS-pop to our

advantage to estimate a juvenile survival rate with less uncertainty. We use information from

adult survival and fecundity estimates, and the fact that they are density dependent, to estimate

juvenile survival:

$$S_A + S_J \cdot F = 1 \ \#(9)$$

164   where, S_A and S_J are survival rates of adults and juveniles at mean population size (when $D = 1$)

165   and F is the fecundity at mean population size. This equation states that population growth rate

166   (λ) is equal to 1 when population abundance is at its long-term average. Using this equation, we

167   would only need priors for adult survival and fecundity, and we can calculate juvenile survival

168   as:

$$S_J = \frac{1 - S_A}{F} \ \#(10)$$

Third, we used a zero-inflated Poisson for modelling fecundity in equation 5b because

there were several years and populations with no juvenile captures. Fourth, we did not use

populations and years with no adult captures when modelling fecundity. While we account for no

capture years in equation 2, we believe that limiting fecundity estimation to years with adult

captures provides a more robust estimate.

Fifth, we changed the spatio-temporal variance structure of survival to be only temporal.

Spatio-temporal variance structure in survival for Brown Creeper proved to be problematic

because it made convergence harder for multiple parameters while also reducing the effective

sample size of their MCMC chains. Lastly, we standardized the density index (D), calculated in

equation 3, to be 0 at mean population size. This allowed for faster convergence of the MCMC

chains in JAGS. See Appendix S2 for details on these adjustments to basic CJS-pop framework

and goodness-of-fit testing of CJS-pop .

We fit three different CJS-pop models to Brown Creeper data: 1) density dependent, 2)

density independent, and 3) Density dependent without the residency model.

Population Projections

We ran population projections using a stage-structured population model with

environmental and demographic stochasticity in both survival and fecundity. We parameterized

these population models with 3 different parameter sets:

1) True simulation parameters that we used to generate capture history data.

2) Parameter estimates from density-dependent and density-independent CJS-pop fit to

simulated data. At each replication we randomly selected the parameters of CJS-pop fit to

one of the 56 data sets with this population and carrying capacity combination. To further

incorporate parameter uncertainty, we randomly used at each replication the 2.5%, 50%,

or 97.5% percentiles of the selected parameters.

3) Parameter estimates obtained from CJS-pop fit to the Brown Creeper data. We employed

the full posterior distribution of the estimated parameters. At each iteration of the

195   projections, parameter estimates were randomly selected from the posterior distributions

196   of all parameters with respect to their correlation structure.

197   Using each of these sets of parameters, we ran single-population projections, with a

198 carrying capacity of 1000 and an initial population of 500 adults and 500 juveniles. We ran the

199 projections with 1000 replicates, and each replicate for 20 years. In order to incorporate

200 environmental stochasticity, at each iteration and at each year we generated random temporal

201 effects separately for survival and fecundity using σ_s^2 and σ_f^2, respectively. We recorded the

202 minimum abundance of the population across 20 years for each iteration, and the distribution of

203 minimum abundance among iterations. The expected value of this distribution is called expected

204 minimum abundance (EMA) and it is a more informative metric than extinction risk, because the

205 latter often has a distribution restricted to near-zero or near-one values (McCarthy and

206 Thompson 2001).

207   *Software*

208   See appendix S2 for a list of R packages used in the analysis. We used JAGS (Plummer

209 2003) as the MCMC sampler when fitting CJS-pop to data. We ran the models with 4 chains,

210 50000 iterations, 20000 burn in, and a thinning rate of 10 for simulated data sets, and with 4

211 chains, 100000 iterations, 50000 burn in and a thinning rate of 20 for Brown Creeper data. We

212 checked convergence with R-hat values, and assumed chains were converged when $\hat{R}$ was <1.05.

213 See Appendix S4 for JAGS code of CJS-pop. Additionally, R and JAGS code of CJS-pop

214 analysis, example simulation data, and results of Brown Creeper analysis are accessible as Data

215 S1 and also at https://github.com/bilgecansen/CJS-pop. The source code and data has been

216 archived and made accessible in Zenodo (DOI: 10.5281/zenodo.3736702).

217   **Results**

Simulations

CJS-pop is able retrieve true parameter estimates without any apparent bias except for

density dependence (Appendix S1: Figs. S1-6). DD strength is estimated with no bias when

strength of DD used to generate capture history data is moderate. Strong DD in data simulation

leads to slight underestimation of DD strength. There is, however, considerable overestimation

of true DD parameters when capture history simulation was carried out with weak DD strength

(Appendix S1: Fig. S1).

Empirical Example: Brown Creeper

We detected weak DD on survival ($\beta = -0.27$, Fig. 1a), and on fecundity ($\zeta = -0.13$,

Fig. 1b) for Brown Creeper. Process variance estimations are low for survival, and high for

fecundity, ($\sigma_s = 0.23, \quad \sigma_f = 0.97$; Table S1). Survival at mean population size for adults and

juveniles were estimated at 0.42 and 0.31, respectively. Our estimate of fecundity at mean

population size was 1.94 (1.21 – 3.01) juveniles per adult (Appendix S1: Table S1). In addition,

Bayesian p-values for both the survival and fecundity components indicated good model fits

(0.24 and 0.50, respectively).

Projections

Density-independent projections parameterized with CJS-pop fit to Brown Creeper data

lead to frequent population extinctions and explosions, which is apparent in the population

trajectory (Fig. 1c) and bi-modal distribution of minimum abundances (Fig. 2). Density

dependence in population models lead to more regularized projections in which population

extinctions and explosions are less frequent (Fig. 1c and Fig. 2). The distribution of minimum

abundances from density-dependent projections of Brown Creeper show a generally similar

pattern to projections parameterized with CJS-pop fit to simulated data irrespective of the DD

strength of capture history simulations (Fig. 2).

Projections made with population models that are parameterized with density-dependent

CJS-pop fit to simulated data are close to projections made with true simulation parameters,

especially when true simulation parameters included moderate or strong DD. This

correspondence demonstrates, in a biologically relevant context, the ability of CJS-pop to fit

realistic models to data (Figs, 2c-d). In contrast, projections with density-independent CJS-pop

(Figs. 2b-d), and with CJS-pop fit to simulated capture history data with weak DD (Fig. 2b) were

not close to projections with the true simulation models. Overestimation of DD strength in CJS-

pop fit to weak DD data also results in overestimation of projected minimum abundances.

Density-independent projections tends to result in frequent population extinctions or explosions

(Figs. 2b-d).

Discussion

CJS-pop is ready to be applied to bird species captured in the MAPS program for making

population projetions. It can be extended to include weather, climate and other exogenous factors

in addition to population density. The true value of CJS-pop lies in its ability to use limited data

to parameterize population models that in turn can be used to predict changes in population sizes

and distributions. We believe this is especially important for developing countries that do not yet

have extensive bird banding and survey programs like MAPS and BBS. We don't, however,

think that whole CJS-pop framework needs to be used for the work we presented here to be

useful. Rather, we argue that the independent ideas explained in estimating fecundity, juvenile

survival and density dependence can be integrated to other CJS frameworks, potentially creating

models capable of population projections. Below we describe 4 of the main advances CJS-pop

provides to mark-recapture literature and discuss the trade-offs made when building the

framework.

1 – Fecundity Estimation

We estimated fecundity in the simulation data with no apparent bias (Appendix S1: Fig.

S4). The fecundity estimate for Brown Creeper ($F = 1.94$ ($1.21 - 3.01$); Table S1) is also

biologically realistic; Brown Creepers tend to have a single brood with a clutch size of 5 to 6

eggs in a breeding season. The main contribution here is that every vital rate (including

fecundity), capture probabilities, and nuisance parameters were estimated simultaneously in a

single model run. This allows for propagation of uncertainty among these parameters but also

makes it possible for parameters that are estimated with less data (juvenile survival) to be

informed by parameters estimated with more data (fecundity and adult survival).

2 – Juvenile Survival Estimation

We detected no biases in juvenile survival estimates from CJS-pop in the simulation data

(Appendix S1: Fig. S2). Juvenile survival estimates of Brown Creeper were similar between a

density-dependent CJS-pop, and a CJS model that did not include fecundity or density

dependence estimation but accounted for "transient" juveniles that leave their population in their

first year (0.32 and 0.30, respectively). However, using information from fecundity and adult

survival in setting the prior for juvenile survival reduced the estimation uncertainty considerably

in CJS-pop. The 95% credible interval for juvenile survival in the CJS model is $0.08 - 0.71$,

while in CJS-pop this interval is $0.19 - 0.48$.

3 – Density Dependence Strength Estimation

Density dependence strength in mark-recapture studies is usually estimated using

abundance directly as a covariate (for example, Nater et al. 2018). This approach can become

problematic with more than one population, especially when each population has different

habitat characteristics and therefore can support different number of individuals. If the goal is to

estimate a species-specific density dependence strength that is applicable across all populations

of the species, abundance of each population in each time step should be standardized with a

proxy for how many individuals each population can support (e.g. carrying capacity). Here, we

made this standardization using long-term abundance average of each population (equation 3).

The density dependence strength we estimate is minimally biased for capture histories generated

by moderate and strong density dependence (Appendix S1: Fig. S1). There is a more apparent

bias when capture histories are generated with weak density dependence (Appendix S1: Fig. S1).

However, the weak density dependence strength we used does not constitute a biologically

realistic scenario if we consider the intrinsic growth rate associated with the density dependence

strength from an allometric standpoint (See Appendix S2 for a detailed discussion about

parameterizing simulations). Most importantly, density dependence strength estimates of

survival and fecundity are ready to be used for population projections.

4 – Population Projections

Estimation of fecundity, juvenile survival, and density dependence strength makes it

possible to make population projections that are useful for conservation purposes. All of these

three vital rates and demographic parameters are required, in addition to what is estimated in

standard CJS models, to build a stage-structured population model. However, instead of

fecundity, a parameter called recruitment rate is frequently estimated from frameworks such as

IPM (for example, Ahrestani et al. 2017). This parameter cannot be used in stage-structured

population models because it combines information from both emigration rate and the number of

new-born individuals. Additionally, including only fecundity estimation in a CJS framework is

also not enough, because without density dependence, these population projections would not be

biologically meaningful. Population projections tend to explode in size or go extinct under

exponential growth when there is environmental stochasticity that is not regularized by density

dependence (Fig 1). The usefulness of this regularizing effect is also visible in the distribution of

minimum abundances from population projections. A density-independent model cannot capture

the minimum abundance distribution generated with a stochastic and density-dependent

simulation, even when density dependence is weak. (Fig 2).

Trade-offs

Using robust design mark-recapture data as the sole data set for the CJS-pop framework

requires several trade-offs. First, we estimate expected juvenile abundance twice (N_1 in equation

2 and N_1' in equation 5a). If we only use N_1', JAGS will give an error regarding the circular

structure of the model because N_1' would also have been used in the denominator of equation 3.

We see this as a minor issue because N_1' is used in the estimation of fecundity and we showed

that fecundity can be estimated without bias in this structure (Appendix S1: Fig. S4).

Second, we estimate population sizes as expected values rather than random variables in

CJS-pop framework. Modelling population sizes as random variables either requires informative

priors on population sizes themselves or another data set, such as population counts, to make the

model more stable and allow convergence (this is essentially what IPMs do). Here, however, we

wanted to show that the vital rates necessary for population models can be estimated using only

mark-recapture data. Using expected abundance, while not ideal, ensures that this framework

requires only a single data source.

Third, we use a Poisson distribution instead of a binomial in equation 5b. Number of

captured juveniles cannot be higher than the actual number of juveniles; this relationship is

explicitly modelled as such with a binomial distribution. However, because we are using

expected abundances in CJS-pop, there could be instances when there are more captured

individuals than the expected abundance. Poisson distribution, by allowing such instances to

occur, increases model stability and eases convergence. Because Bayesian p-value for brown

creeper data is 0.5, we can say that this structure can represent the data well (Bayesian p-values

close to 0.5 indicate better fit; Kéry and Royle 2016). Last but not least, the framework we

present here is complex and in JAGS it takes for about 20 hours for Brown Creeper model to

converge.

Acknowledgements

We thank Kevin Shoemaker for valuable input. This study was initiated under funding

from the NASA Biodiversity Program (NNH10ZDA001N-BIOCLIM). We thank the many

volunteers who have contributed to the MAPS program and the Institute for Bird Populations for

development and active curation of the MAPS dataset.

References

Ahrestani FS, Saracco JF, Sauer JR, et al (2017) An integrated population model for bird

monitoring in North America. Ecological Applications 27:916–924

Akçakaya HR, Burgman MA, Ginzburg LR (1999) Applied population ecology: principles and

computer exercises using RAMAS EcoLab 2.0. Sinauer Associates Sunderland, MA

Cooch EG, White GC (2016) Program Mark: A Gentle Introduction, 14th edn.

Dail D, Madsen L (2011) Models for Estimating Abundance from Repeated Counts of an Open

Metapopulation. Biometrics 67:577–587. https://doi.org/10.1111/j.1541-

0420.2010.01465.x

Ergon T, Gardner B (2014) Separating mortality and emigration: modelling space use, dispersal

and survival with robust-design spatial capture-recapture data. Methods in Ecology and

Evolution 5:1327–1336. https://doi.org/10.1111/2041-210X.12133

Kéry M, Royle JA (2016) Applied hierarchical modeling in ecology: analysis of distribution,

abundance and species richness in R and BUGS. Elsevier/AP, Academic Press is an

imprint of Elsevier, Amsterdam☐; Boston

Lebreton J-D, Burnham KP, Clobert J, Anderson DR (1992) Modeling Survival and Testing

Biological Hypotheses Using Marked Animals: A Unified Approach with Case Studies.

Ecological Monographs 62:67–118. https://doi.org/10.2307/2937171

McCarthy MA, Thompson C (2001) Expected minimum population size as a measure of threat.

Animal Conservation 4:351–355. https://doi.org/10.1017/S136794300100141X

Plummer M (2003) JAGS: A Program for Analysis of Bayesian Graphical Models using Gibbs

Sampling. 3rd International Workshop on Distributed Statistical Computing (DSC 2003);

Vienna, Austria 124:

Ryu HY, Shoemaker KT, Kneip É, et al (2016) Developing population models with data from

marked individuals. Biological Conservation 197:190–199.

https://doi.org/10.1016/j.biocon.2016.02.031

Schaub M, Abadi F (2011) Integrated population models: a novel analysis framework for deeper

insights into population dynamics. Journal of Ornithology 152:227–237.

https://doi.org/10.1007/s10336-010-0632-7

Schaub M, Royle JA (2014) Estimating true instead of apparent survival using spatial Cormack-

Jolly-Seber models. Methods in Ecology and Evolution 5:1316–1326.

https://doi.org/10.1111/2041-210X.12134

Williams BK, Nichols JD, Conroy MJ (2002) Analysis and management of animal populations:

modeling, estimation, and decision making. Academic Press, San Diego

Figure Legends

Figure 1: a) Relationship between adult survival and relative population density (D) of Brown Creeper as modeled by CJS-pop. b) Relationship between fecundity and relative population density (D) of Brown Creeper as modeled by CJS-pop. c) Solid line indicates the median trajectory of population size, across 12000 trajectories, of Brown Creeper projected by a stage-structured population model that was parameterized with a Density-Dependent (DD) or Density-Independent (DI) CJS pop. Shaded areas include 50% of the population trajectories. Carrying capacity was set to 1000 in the DD projections. Note the difference in scale in population size between DD and DI projections.

Figure 2: Distributions of minimum abundances resulting from population projections made with stage-structured population models that were parameterized with CJS-pop fit to Brown Creeper data (a), and with CJS-pop fit to simulation sets that was generated with different density-dependence strengths (b,c,d). Light blue represents parameterizations of population models with density-independent (DI) CJS-pop, orange represents density-dependent (DD) CJS-pop, and green (TRUE) represents population models that was parameterized with original simulation parameters that was used to generate capture history data. High probability density at 0 and 1000 indicates frequent population extinctions and explosions, respectively.

398   **Figure 1**

401   **Figure 2**

402