Cell growth model with stochastic gene expression helps understand the growth advantage of metabolic exchange and auxotrophy

Dibyendu Dutta [#], Supreet Saini [#]

Department of Chemical Engineering, Indian Institute of Technology Bombay, Mumbai, India

Running Title: How do cooperators grow faster than prototrophs

[#] **Corresponding Authors:**
Dibyendu Dutta   Email: dibyendu_dutta@iitb.ac.in
Supreet Saini   Email: saini@che.iitb.ac.in

Abstract:

During cooperative growth, microbes often experience higher fitness, due to sharing of resources by metabolic exchange and herd protection through biofilm structures. However, the trajectory of evolution of competitive species towards cooperation is not known. Moreover, existing models (based on optimisation of steady-state resources or fluxes) are often unable to explain the growth advantage for the cooperating species, even for simple reciprocally cross-feeding auxotrophic pairs. We present an abstracted model of cell growth that considers the stochastic burst-like gene expression of biosynthetic pathways of limiting biomass precursor metabolites, and directly connects their cellular levels to growth and division using a "metabolic sizer/adder" rule. Our model recapitulates Monod's law and yields the experimentally observed right-skewed long-tailed distribution of cell doubling times. The model further predicts the growth effect of secretion and uptake of metabolites, by linking it to changes in the internal metabolite levels. The model also explains why auxotrophs may grow faster when provided the metabolite they cannot produce, and why a pair of reciprocally cross-feeding auxotrophs can grow faster than prototrophs. Overall, our framework allows us to predict the growth effect of metabolic interactions in microbial communities and also sets the stage to study the evolution of these interactions.

Importance:

Cooperative behaviours are highly prevalent in the wild, but we do not understand how it evolves. Metabolic flux models can demonstrate the viability of metabolic exchange as cooperative interactions, but steady-state growth models cannot explain why cooperators grow faster. We present a stochastic model that connects growth to the cell's internal metabolite levels and quantifies the growth effect of metabolite exchange and auxotrophy. We show that a reduction in gene expression noise explains why cells that import metabolites or become auxotrophs can grow faster, and also why reciprocal cross-feeding of metabolites between complementary auxotrophs allow them to grow faster. Our framework can simulate the growth of interacting cells, which will enable us to understand the possible trajectories of the evolution of cooperation *in silico*.

Keywords:

Cross-feeding, mutualism, cooperation, microbial communities, stochastic growth model, metabolite exchange, adder, sizer, gene expression noise, division of labour, resource allocation

Introduction:

Unlike the pure cultures in a laboratory, in the wild, microbes live alongside many different species. To survive, they may either compete for limited resources or cooperate. Defying general evolutionary expectations, cooperative communities pervade different ecological niches (1–5), highlighting the purported selective advantage of cooperation (6). Cooperative interactions often involve secretions, ranging from extracellular enzymes, scavenging molecules (like siderophores) (7), to leaked metabolites – either toxic by-products (8, 9) or essential metabolites like amino acids (10–12). Cooperation may also involve living together in a biofilm, which protects the group from the effect of toxins, antibiotics, and predators (13).

To fully understand the advantages of cooperation in ecological communities (14) and to implement them in synthetic communities has been challenging (15–17). The focus has been on metabolic modelling to identify the keystone "currencies" of cooperation (9, 18–23). Knowing which metabolites are "traded" allows us to understand the mechanism of cooperation and how it enables survival in different environmental conditions. However, to emerge as the dominant strategy in any ecological niche, cooperative consortiums must grow faster than other free-living species. Metabolic modelling approaches, however, can only determine the metabolic viability and synergy, in terms of the consortium's productivity of metabolic flux exceeding the sum of individuals. Models usually do not yield the growth kinetics of the cooperative consortium, and hence it has not been possible to study theoretically, the evolution of cooperation using existing evolutionary frameworks like fitness landscapes.

Furthermore, there are several shortcomings in the existing modelling approaches. Microbial growth is often posed as an optimal resource allocation problem that operates on the steady-state values of intracellular components, to simplify computation. Growth is estimated using optimal steady-states values. However, often computation of the steady-state itself uses the growth rate as the rate of dilution due to cell division, which is assumed constant and not updated, leading to inaccurate estimates of growth. Often such models consider only a part of the cell and coarse-grain its dynamics to attempt maximisation of the energy generation while minimising the resource investment in catabolic enzymes and transporters (24, 25). More sophisticated whole-cell models like

bacterial growth laws and account for the shortcomings by computing a proxy measure

of growth rate using the steady-state value of ribosome fraction of the proteome (26–

29). Metabolic flux models entirely circumvent this by only computing the steady-state

fluxes to optimise the genome-wide flux contribution to biomass and uncover the

pathways or enzyme reactions that bottleneck growth (30–34).

These steady-state approaches, neglect the cell-to-cell differences due to stochastic gene

transcription (35–37), leading to differences in the numbers of proteins (depending on

the controlling promoter) (38), and in the cell's metabolic state (39). While such

variations are unimportant for prediction of the mean characteristics of the population,

it is critical to model the growth of a cooperative consortium with cell-to-cell

interactions such as the exchange of metabolites.

In this work, we develop a microbial growth model that not only captures the effects of

stochastic variation on growth of non-interacting cells but also is applicable for studying

the growth of cells that exchange metabolites. We implement stochastic burst-like gene

transcription using Monte Carlo methods following the framework proposed by Golding

et al. (40). To determine when a cell divides, we combined the concept of a biomass

objective function (41) with the empirical laws for bacterial cell growth and size

homeostasis: Adder and Sizer (42–45). We postulate that any cell size increment

corresponds to an equivalent amount and stoichiometric composition of biomass

precursor metabolites. Thus, the cell grows and divides only after producing the

required amounts of metabolites in the correct stoichiometry.

Our framework attempts to capture the dynamics of cooperative growth and study its

collective growth kinetics when cells exchange metabolites. We want to find out the

factors that lead to faster growth of the consortium, such that we can begin to

investigate the evolutionary relationships between these. We obtained the distribution

of cell doubling times, which mimics the experimentally observed distributions (46, 47)

obtained from single-cell growth experiments run on microfluidic devices (42, 48). Our

model successfully recapitulates the hyperbolic relationship between the substrate and

growth rate observed by Monod (49). We further demonstrate that gene expression

noise manifests as noise in metabolite levels, and hence delays cell division (i.e. the

noise is time additive). We demonstrate that by reducing this overall noise by reducing

the effective number of limiting metabolites, either by auxotrophy or by the direct

import of the metabolite can accelerate growth. Lastly, we demonstrate that reciprocal

cross-feeding between auxotrophs can be faster than the growth of prototrophs. In all,

we demonstrate how incorporating stochastic gene expression noise into a cellular

growth model allows us to capture the growth effect of auxotrophy and metabolite

exchange directly, and how it can help understand the evolution of cooperation.

Results:

1. Model Development

We develop a cellular growth model that avoids relying on cellular steady-states and also incorporates the stochastic variation in growth.

As discussed, recent studies have put forth the empirical size based laws for bacterial cell division: Adder and Sizer (42–45). Combining these with the idea of biomass objective functions, we argue that cell size increase depends upon the production of the equivalent amount of constituent metabolites, in the proper stoichiometry. We propose the idea as the "metabolic adder" and "metabolic sizer", and use it as the foundation of our proposed cell growth model, such that cell division is triggered only when the cell produces the minimum amounts of all the necessary metabolites (Fig – 1a). We demonstrate that our model approach recapitulates many of the known cellular phenomena and go on to extend the model to describe the kinetics of single-cell growth while they exchange metabolites among each other. We find the Adder and Sizer models very similar in their outcomes and proceed with Sizer for all the rest of the simulations in this work, primarily because it is found to be the better fit for the outcomes in case of poor media conditions, where growth in on minimal media with single substrates (44)(See Supplementary S4).

Now, while theoretically, a shortfall in any metabolite could prevent a cell from dividing, only a few metabolites at a time may act as bottlenecks to cell division. These may be required in substantial amounts, or the production of the respective enzymes may be variable and in low numbers (39), such that the metabolite demand cannot be met in time. Additionally, some metabolites may be prone to leakage (may serve as public goods) and may act as a bottleneck to growth (50). We assume the cell comprised of p anabolic pathways, whose products act as a bottleneck for cell division. All metabolite biosynthesis (anabolic) pathways are fed by the substrate supplied by the upstream catabolic pathways via different shunts from the Central Carbon Metabolism. We simplify the arrangement and assume that all anabolic pathways are fed a common

substrate, which we further assume to be supplied at a constant rate $p \cdot S_{in}$ per second

(Fig – 1b).

The distribution of cell doubling times across different growth conditions collapse to

one distribution when scaled by their mean (47). This leads us to believe that the

common downstream metabolic pathways like anabolism are responsible for the

observed noise in growth, since in different growth conditions, upstream catabolic

pathways may differ. Hence, our modelling approach focuses on quantifying the noise

from anabolic pathways alone. Furthermore, to circumvent the differences due to the

exact structure and kinetic properties of different anabolic pathways, we assume all p

pathways to be linear and comprised of n enzyme catalysed steps, all of which we

assume to be identical in kinetic properties and the transcription parameters.

We further assume that these n enzymes in each pathway are present on the same

operon, and hence, their transcriptional bursts are synchronised (Fig – 1b). To

incorporate the effect of stochasticity in our model, we incorporate stochastic burst-like

gene expression, using the findings of exponentially distributed durations of

transcription burst (t_{ON}) and the waiting time between successive bursts (t_{OFF}) (40).

We randomly sample OFF and ON durations iteratively from exponential distributions

of set mean, to generate the stochastic burst profile (Fig – 1c). Next, assuming no

shortage of RNA polymerases, transcription proceeds at a constant rate till the end of

t_{ON}, when any incomplete transcripts are terminated. Each mRNA produced is

randomly assigned a lifetime by sampling from an exponential distribution. During the

lifetime of an mRNA, one ribosome binds at a time and produces proteins, assuming

ribosomes are always sufficient. (Ribosome limitation only reduces the production of

proteins and represent a nutrient-poor condition, but will not affect the overall

dynamics). Any incomplete protein is terminated once the mRNA decays. Each protein

produced is randomly assigned a lifetime by sampling from an exponential distribution.

We use a system of coupled ODEs based on Michaelis-Menten kinetics for the enzyme

kinetics, with the stochastic protein profiles as time-varying parameters to compute the

production of metabolites (See Methods for the equations). We convert all quantities to

molecules per cell assuming a cellular volume of $1 \mu m^3$. When the production of all the p

metabolites crosses the set threshold based on the "metabolic adder or sizer" law used,

the cell divides. Post division, all the contents of the cell are divided between the two

daughter cells equally (51). The daughter cells start from the inherited values of the

mRNA, proteins and metabolites, but simulate unique stochastic burst profiles, going

forward.

2. Model properties and outcome:

Simulating our model, for a given set of parameters, we obtain the duration of time

between successive cell divisions or the cell generation time. The histogram of the

generation times for all simulated parameter sets reveals a right-skewed distribution

with a long tail, which concurs well with many experimental observations of bacterial

growth at a single-cell resolution (42, 44, 45, 47, 52, 53). These studies indicated that

the observed generation time distribution resembled a Gaussian on a logarithmic scale,

and hence may be a Gamma or log-Normal distribution (47). (46), on the other hand,

analysed the single-cell dataset extensively and made a distinction between the

quantities: cell generation time (T_μ) (or inter-division time) and the cell doubling

time (T_{div}), the time for the doubling of cell size.

$$Doubling\ time\ (T_\mu) = \frac{Generation\ time\ (T_{div})}{log_2\left(\dfrac{cell\ size\ at\ division}{cell\ size\ at\ birth}\right)} \tag{46}$$

He tested the fit of different distributions to the experimentally derived doubling time

distribution and found that a log-Fréchet distribution provided the best fit to the data.

Now, in our framework, cell division is triggered by exact size laws and division yields

perfect halves, hence the ratio of size at division and birth is always two and thus

generation time and doubling time are the same. The simulated generation times

obtained, give a long-tailed right-skewed distribution and is fit best by a Fréchet or

Generalised Extreme Value (GEV) Type – II distribution (Fig – 2a). Since the generation

time distribution obtained is the maximum of the production times of the p metabolites,

the obtained GEV distributions match with intuition derived from the Extreme Value

theory(as the limit distribution of the maxima of a sequence of independent and

identically distributed random variables) (54). We found that for all variations of p and

S_{in} tested, the generation time distribution was a Fréchet or GEV Type – II (shape

parameter: k > 0) (Supplementary S1).

The Fréchet distribution obtained from our simulations may appear to be at odds with

experimental data fitting to log-Fréchet. We thus looked for possible causes of

discrepancy and implicated the lack of DNA replication in our model (See

Supplementary S3). In our model, we assume ribosomes increase in sync with the

demand and are never limiting.

Now, if the number of bottleneck pathways (p) increases, the chance that one of the p

pathways would be slower due to the stochastic noise increases, and hence the

maximum of the first passage times, or generation time also increases. The distribution

of cell generation times, shifts to the right and reduces in skewness (Fig – 2b;

Supplementary S1). Moreover, upon scaling the data with the mean, we find that the

distributions visually overlap (Fig – 2b inset), but with increasing p, skewness and

kurtosis decreases (Supplementary S1).

Now, substrate flux per pathway (S_{in}) represents the available concentration of the

nutrients in the environment. The downstream anabolic pathways utilise the substrate

for metabolite production. When S_{in} is low, the rate of metabolite production is low and

cell generation times are longer but as S_{in} increases, metabolite production picks up,

and cell generation times are shorter, and we observe the distribution of cell generation

times shift to the left (Fig – 2c). Upon scaling the data with the mean, we see the

distributions visually overlap (Fig – 2c inset), but with increasing S_{in}, skewness and

kurtosis decrease (Supplementary S1). Next, when we plot the associated growth rate

against the corresponding S_{in}, we observe a hyperbolic relationship, for all values of p

simulated, which recapitulates Monod's observations between substrate concentration

and the cell's growth rate (Fig – 2d).

Substrate flux is known to be the primary determinant of growth rate in a cell, and

studies have hence focused on quantifying the noise in catabolic pathways as a measure

of substrate flux variation to estimate growth rate variations (36). However, during

balanced growth conditions, the substrate flux from catabolism is stable, and hence we

chose only to model the noise contribution from the downstream anabolic processes.

Thus, for all values of substrate flux, the variation in the generation time is the same, the

generation time distributions overlap when scaled by the mean.

Studies have shown that the cell division control is concerted by more than one factor

and limiting agent (55, 56), which in our model is represented by multiple bottleneck

pathways (p). However, the number of bottlenecks may depend on the cellular growth

conditions, and in case of poor media conditions (low S_{in}) it may increase. Although our

model is unable to perform such a switch, it nonetheless can still recapitulate

observations such as Monod's Law (Fig – 2d).

3. Effect of different modelling criteria on the nature of cell generation times observed

Imperfect partition of cells

As described before, the only source of stochastic noise we consider in our model is the

burst-like gene expression process. However, other sources can also add noise to the

cellular growth outcome such as variation in cellular partition ratio, protein translation,

and mRNA degradation. Nevertheless, their relative importance varies across different

growth rates, with gene transcription and cellular partition being the most important,

while mRNA degradation and protein translation are important only at very low rates of

growth (57). Hence, we discuss only the effect of imprecise partitioning.

Although the process of cell division in bacteria is highly robust and precise, the two

daughter cells obtained are not always of equal sizes. Discrepancies in partition ratio

are rare in simple rod-shaped cells, but higher in case of other cell shapes, and on an

average, the Coefficient of Variation (CV) in terms of cell length ranges from 3 – 15%

(58). Moreover, the profile of noise contribution due to unequal partitioning of the cell

has been shown to be virtually indistinguishable compared to stochastic gene

transcription (59). We incorporate a randomly sampled partition ratio from a Normal

distribution, with mean 0.5 and varying CV. The simulated outcomes are almost

indistinguishable from perfect partitioning, in case of a CV = 0.03 (Supplementary S6),

however for CV = 0.1, the additional noise delays the cell division process, and the

distribution of generation times is shifted to the right (Fig – 3a).

Effect of gene organisation as Operon

Bacterial genes that are co-expressed are mostly organised as a contiguous unit in the

genome called operon (60–65). Our model design thus assumes that all the biosynthetic

pathway enzymes occur in the same operon and share the same transcriptional burst

profile. If the enzymes were organised separately and transcribed independently, each

with its own burst profile, then the noisy lack of synchrony would not only lead to

wasted resources (66, 67), we also expect a delay in the metabolite production and

hence cell division. Simulating our model with independent expression shows the

distribution of generation times is shifted to the right, towards higher values (Fig – 3a).

Comparing the distributions when scaled to the means, a lowered skewness and

kurtosis is observed (Fig – 3a inset, Supplementary S1). In both the cases, GEV

distributions continue fitting the data, best. Our model thus demonstrates that operon

structure also leads to kinetic (growth rate) gains for the cell, in addition to the already

hypothesised economic gains (66).

The parameter space of transcription bursts

The parameter values of t_{ON} and t_{OFF} describe the burst-like gene expression in our

model (40). It is expected that a fast-growing cell would have lower t_{OFF} and higher t_{ON}

for anabolic pathways, however experimental evidence for the same does not exist. We

hence varied these parameters to explore its parameter space and its effect on the

model outcome. Reducing t_{OFF} and increasing t_{ON} both lead to an increase in the

number of mRNAs produced, due to longer and larger number of transcriptional burst

events. Consequently, the distribution of generation times shifts towards lower

generation times, along with a decrease in the width of the distribution (Fig – 3b,c). The

distributions scaled to the mean, reveal that reducing t_{OFF} is accompanied by a

reduction in skewness and kurtosis (Fig – 3b inset, Supplementary S1), but increasing

t_{ON} leads to an increase in skewness and kurtosis (Fig – 3c inset, Supplementary S1).

GEV distributions still fit the data, but with a lowered value of the shape parameter.

Alternately, an increase in t_{OFF} and a decrease in t_{ON} leads to a lower number of

transcripts, and hence the generation time distribution shifts its peak towards higher

values and becomes wider (Fig – 3b,c). The GEV distribution shape parameter however

increases. These results give us insight into how the internal parameters of a cell may

change and its associated growth effects. By anlysing experimentally obtained doubling

time distributions (42), we find skewness increases in richer media, which may indicate

that higher growth in rich media involves an increase in t_ON parameter rather than a

reducing in the t_OFF.

Varying metabolite threshold

Our model calls the minimum stoichiometric requirement of each metabolite to achieve

threshold size as the metabolite threshold. Varying the metabolite threshold in the

simulation thus corresponds to changing the size requirement for the cell division, given

the same set of intracellular parameters. Since we do not know how the metabolite

threshold varies relative to the internal expression parameters, we varied the threshold

while holding expression constant to understand if it has any bearing on observed

outcomes, which would help us derive better insights from experimental data. As

expected, we find that lowering the threshold leads to higher growth rates, while

increasing it reduces the growth rate (Fig – 3d). As threshold increases the skewness of

the distribution (scaled to mean) decreases, while CV increases (Supplementary S1).

4. Effect of metabolite uptake and secretion on simulated cells

Our primary goal is to develop a framework that can quantify the effect of nutrient

uptake and secretion on the growth of cells. Our model design allows us to capture the

effect of import of external metabolites and the export of internally produced

metabolites directly in terms of any change in the time required to meet the internal

metabolite thresholds, i.e. the generation time.

Till now, our simulations have considered cells were grown on minimal media, such that

only the substrate flux (S_{in}) can fuel the production of metabolites. We now simulate the

case when additional metabolites are present in the media (for example, media

supplemented with amino acids), and the cell can import them directly to meet the

requirement. To implement this, we consider one of the bottleneck metabolites is
imported into the cell at a constant rate, effectively lowering the threshold of a
metabolite that needed to be produced internally and thus cells on average divide
faster. Hence, with increasing rates of import of the limiting metabolite, the generation
times shift to the left (Fig – 4a(i)), and the growth rate increases (Fig – 4a(ii)). From our
model's perspective, import at a sufficiently high rate essentially decreases the number
of bottlenecks ($p \rightarrow p - 1$) and removes its noise contribution to cell division (Fig – 5d),
which implies that the advantage from the import of one metabolite has an upper
bound, and the advantage due to higher rate of import saturates quickly (Fig – 4a(iii)).

Next, we consider the effect of exporting metabolites from the cell. Some metabolites
are usually produced in excess and leak out from the cell (10–12, 50), without any
adverse effect on growth. However, if these metabolites are not useless by-products for
the cell, then excess secretion could turn it into an additional bottleneck and impede
growth. In case, one of the limiting metabolites is secreted, it leads to an effective
increase in the metabolite threshold, and hence the cell requires more time to divide.
We quantify the growth effect of secreting metabolites, by simulating secretion as a
fraction of the total metabolite produced per second, rather than a constant amount
secreted per unit time, which synchronises the secretion to the internal production of
the metabolite. It hence captures a more accurate estimate of the amount secreted (low
secretion after birth due to low enzyme, and higher secretion before division due to a
higher number of enzymes and transporters). As the secretion ratio increases, cell
generation times increase, and the distribution shifts to the right (Fig – 4b(i)). The
magnitude of the difference is however much more pronounced in case of a nutrient-
poor media (low S_{in}). With the distributions scaled to the mean, we further see that
increased secretion leads to lowered skewness, and skewness decreases more in case of
the nutrient-poor media. Unlike the case with nutrient uptake, the effect of secretion is
unbounded, and increased secretion further lowers growth rate (Fig – 4b(ii),(iii)). All of
the above represent testable predictions from our model, which may be tested by
growing appropriate strains at single-cell resolution in appropriate media.

5. Growth advantage of auxotrophy

We have already demonstrated how direct import of a limiting metabolite can increase

growth, and how the advantage is capped. We described this phenomenon as the partial

alleviation of the noise contribution from one of the bottleneck metabolites, which

corresponds to $(p \rightarrow p - 1)$ in our model. If the modelled prototrophic cell mutates to

lose one of the biosynthetic pathways for a limiting metabolite (and becomes an

auxotroph), then the same outcome may be obtained. Interestingly, this is a common

phenomenon, seen not only in endosymbiotic bacteria (68) but also in some free-living

bacteria (69, 70). Black Queen Hypothesis proposes that this genome reduction is

driven by a resulting selective advantage (50, 71). However, such gene loss makes the

survival of the cell contingent on sufficient external supply of the metabolite.

For the simulation, we consider a cell with three bottleneck pathways (p), and then

delete one of the concurrent biosynthetic pathways, while simultaneously allowing the

cell to meet the requirement for the missing pathway by direct import. At low import

rates, this missing metabolite determines the cell generation time, however at higher

import rates the missing metabolite is no longer the bottleneck, and the generation time

distributions overlap with the distribution for $p - 1$ bottlenecks (Fig – 5a), as expected.

It is interesting to note that when an auxotroph is grown in poor media conditions,

represented by the low substrate flux (S_{in}) in our simulations, a sufficient rate of import

of the limiting metabolite, allows the cells to grow much faster (Fig – 5b,c), relative to

the growth advantage observed for high substrate flux conditions. From the perspective

of our model, the available internal substrate flux ($p \cdot S_{in}$) originally meant for p

pathways is now redistributed among the remaining $p - 1$ pathways which in turn

allows these auxotrophs to reach their metabolite requirement thresholds faster and

hence divide faster.

Our model presents a novel alternative explanation for the nature of growth benefit or

selective advantage obtained due to gene loss (50, 71), leading to the spontaneous

evolution of auxotrophs (72), compared to the usual modelling arguments based on

saved resources. Our model does not explicitly incorporate any effect due to saved

resources. If it did, the auxotroph is likely to grow even faster, with additional

production of enzymes due to the reinvestment of the saved resources.

It is imperative to note that the high growth advantage demonstrated here, corresponds

to a low number of bottlenecks (low p). If a large number of metabolites are bottlenecks

simultaneously (large p), then the relative growth advantage observed due to loss of
one pathway or direct import of one limiting metabolite, will be significantly lower (Fig
– 5d). While it is known that for different nutrient media, different metabolites act as
bottlenecks for growth (73), it is not immediately clear which biomass precursor
metabolites act as a bottleneck, how their numbers vary, and their relative limiting
effect on growth (For our model, we considered all bottlenecks of equal strength).

6. Towards reciprocal metabolic cross-feeding

As discussed, the survival of auxotrophs is contingent on the external availability of the
metabolite they cannot produce internally. These cells invade ecological niches where
other cells secrete the necessary metabolites, either as by-products or public goods. In a
public-goods scenario, the invading cells act as "cheaters" and reduce community
productivity (74–76). However, if the metabolite is a by-product and its accumulation
toxic, then its clearance from the environment could induce the producer cells to secrete
more, and improve community productivity (8, 77, 78).

Analysis of available microbial genomes from various niches has revealed that most of
them are auxotrophic for at least a few essential metabolites, such as amino acids (71,
79). Although surprising, it explains why ecological examples of metabolite cross-
feeding are so commonplace (1–5). During the process of metabolite cross-feeding, the
metabolites secreted by the microbes diversify the nutrient composition of the
environment, which enables survival and further evolutionary diversification of the
group, by allowing for spontaneous evolution of auxotrophies (72).

However, prototrophic species may invade such cross-feeding populations, and by
importing the secreted metabolites "cheat" to grow faster and outcompete the
consortium (80). To avoid such a fate, cross-feeding must also enable fast growth,
enough to surpass the competing species, in addition to enabling the survival of the
group. Nevertheless, we do not understand how the division of labour may allow
cooperating cells to improve productivity enough to exhibit growth that is faster than
prototrophs.

We have demonstrated that the import of limiting metabolites accelerates growth, but

on the other hand, secretion of such limiting metabolites at high rates can also slow

down growth. In a cross-feeding interaction, cells reciprocally exchange metabolites, by

secreting one metabolite and importing another at the same time. How must cellular

resources be balanced between these processes to achieve a net positive outcome is not

understood, even for a simple symmetric exchange.

Let us consider two reciprocal auxotrophs, each overexpressing the metabolite the

other needs for survival. These auxotrophs due to loss of one of the metabolic

biosynthesis pathways, grow faster because they can redirect the saved resources from

the pathway towards growth, and also reduce the effect of gene expression noise on

growth (as we demonstrate using our model). However, the need to supply metabolites

to each other requires the cells to invest the saved resources towards metabolite

overproduction, taking away the growth advantage. Assuming, metabolites of similar

costs are exchanged, intuitively, we may expect that the reinvestment of saved

resources may enable a doubling of the metabolite production, and hence allow the cells

to exhibit a growth rate as high as an equivalent prototroph. Which is precisely the

result we obtain when we simulate a simplified steady-state version of our model (Fig -

6a, See Supplementary S2 for details of the steady-state model simulation). That too,

when not considering the possible losses and delays due to transport of the metabolites.

Thus, it is hard to imagine that cross-feeding can enable such growth advantages.

Pande *et al.* used synthetic reciprocal amino acid auxotrophs that also overproduced the

amino acids to study this phenomenon, and found a large number of cross-feeding pairs

exhibiting growth rates faster than the parental prototroph (11). However, only a few

pairs could outcompete the prototroph in a competitive co-culture, where the

prototroph could cheat on the secreted metabolites. Here, we use our stochastic growth

framework in an attempt to explain these observations.

We start with an auxotroph ($p = 2$) originally a prototroph ($p = 3$). Next, we need to

implement over-expression of the secreted metabolite by reinvesting the resources

saved. However, it is unclear how a real cell handles the over-expression of a pathway.

While in a linear framework, it is merely a matter of doubling the enzyme (considering

equal enzyme costs), in a stochastic framework, over-expression involves changing the

transcription burst frequency, and it would be hard to get precisely "double" the

enzyme. Hence, as a benchmarking test case, we double the kinetic constant of the

enzyme keeping the same stochastic burst profile. We also obtain burst frequency

parameters that double the enzyme produced, by trial and error, and compare the

results obtained. We considered secretion at different ratios of the amount of

metabolite produced per unit time. We further varied the direct import rate of the

missing metabolite. We obtain the resulting generation times, growth rates and

secretion rates (Supplementary S5).

Comparing the benchmark case with the case with altered transcriptional frequency, we

find both of them behaving similarly in terms of the growth outcomes (Supplementary

Fig - 6). For both cases, we find the cross-feeding cells demonstrate growth rates higher

than the prototroph (Fig – 6a(i)(ii)). Even when we compare the growth rates to that of

the prototroph fed one of the metabolites at the same rate as the cross-feeding cell, we

find multiple conditions where the model cross-feeding cells grow faster. It is

interesting to note that these fast-growing cross-feeding cells are also able to produce

the more of the secreted metabolites than what they consume per unit time (Fig –

6b(i)(ii)), indicating increased metabolite productivity. We call these "feasible

solutions" to the problem of the cross-feeding cells being able to exhibit faster growth

while maintaining higher overall metabolic productivity.

We find that the feasible solutions are obtained only when substrate flux is sufficiently

high (S_{in} > 10,000/sec), i.e. values beyond which growth rate saturated (Fig – 2d).

Without the availability of sufficient substrate flux, overproduction of metabolites is not

possible. Thus such kinetic advantage of cross-feeding consortiums may only be

observed in case sufficiently concentrated substrate (rich media) is available. Moreover,

the feasible solutions were observed only for metabolite import rates ($P_{in} \sim$ 7,500 to

10,000 per sec), which marks the values after which improvement in growth rate

saturates (Fig – 5c). Thus, it may be possible for cross-feeding to be beneficial across a

broad range of parameters.

The results we obtained here assume that only a few metabolites act as bottlenecks for

growth at a time, for a cell. For a large number of bottlenecks, relative advantage due to

gene loss is lower (Fig – 5d) and may not be enough to outcompete prototrophs. The

evolutionary success of cooperative microbes may thus be determined by the number of

simultaneous metabolic bottlenecks in the cell.

We demonstrate that even in the case with minimal headroom for improvement of

growth rate (i.e. when the competing bottleneck pathways are of an equal threshold),

feasible solutions for cross-feeding can be found. In cases where the subsequent

bottlenecks are of a lower threshold, the cell would have a higher headroom to improve

its growth with higher import rates. However, lower growth durations may also lead to

lower secretion. In a real cell, however, faster growth also implies faster DNA

replication, which would then allow higher protein production, which would lead to

higher metabolite production.

Discussion

Using our model framework, we have demonstrated that accounting for stochastic gene expression noise (40), allows us to capture the variations in the rates of production of essential cellular metabolites (39) and hence differences in cell growth and division kinetics. Our model is applicable for non-interacting cells, as well as cells that interact via metabolite exchange. Our model recapitulates fundamental relationships like Monod's law and also yields the experimentally observed right-skewed long-tailed distribution of cell doubling times. Our model further demonstrates the growth advantage of metabolite import and genome reduction, and also how reciprocal cross-feeding systems may enjoy a growth advantage, counter to the intuition derived from existing steady-state resource allocation models, that either rely on ribosome fractions (26–29) or changes in flux distribution (30–34).

Our approach to modelling cell growth shares similarities with Thomas *et al.*, where they attempted to understand the role of stochasticity in bacterial cell growth (57). They used a hybrid-stochastic approach to simulate the expression of ribosomes, enzymes, and transporters, which further regulate their own production, building on their previous steady-state model (26). A crucial difference between our approaches is, however, the way we determine cell division. Thomas *et al.* used the Cooper–Helmstetter model (44, 81), where cells divide after a constant C+D period from the initiation of DNA replication, determined using Donachie's constant initiation cell size per origin (82). We instead used a "metabolic sizer/adder", where cells divide after having produced all the biomass precursor metabolites in the required quantity and stoichiometry. Thomas *et al.* further considered cell mass as a proxy for cell size, and unlike our approach did not impose a strict requirement for the stoichiometry of the biomass components.

As previously stated, the goal of our model is to present a useful framework that directly yields growth differences between different cellular states that arise in the ecological niche as microbes compete and cooperate. The microbial cells may be prototrophs, or auxotrophs and may only import metabolites or may secrete them as well, at varying

rates, controlled by internal expression and transport parameters. The framework we

developed can determine the growth rate associated with these cellular states and how

it will change depending upon the metabolic interactions, such as the case of reciprocal

cross-feeding interactions between auxotrophs. Each of these represents intermediate

states towards the evolution of cooperation, and we intend to extend our framework to

study if such evolution towards cooperation is possible. Some studies have pitted

genome-scale flux models of microbes with different auxotrophies and metabolite

secretion rates (or leakiness) against each other using an evolutionary game theory

framework to derive intuition about the microbial combinations that would be

evolutionarily successful (83). However, it has been limited by its inability to account

for real-time changes in growth due to eco-evolutionary feedbacks. We aim to overcome

this by using the model we presented here along with a mutational framework and

provide new insights towards the evolution of cooperation.

Methods

System of coupled differential equations used to compute metabolite production

After generating the stochastic enzyme profile, we pass it as a time-varying parameter to a system of coupled ODEs, based on Michaelis-Menten kinetics.

The primary substrate S common to all the pathways is supplied at a constant rate $p \cdot S_{in}$ and each of the enzymes feeds off it. i represents the bottleneck pathways 1 to p.

$$\frac{dS}{dt} = p \cdot S_{in} - \sum_{i=1}^{p} k_{cat}^{i1} E_{i1} \frac{S}{S + K_M^{i1}}$$

The subsequent products except the final products are modelled as:

$$\frac{dP_{i,j}}{dt} = k_{cat}^{ij} E_{ij} \frac{P_{i-1,j}}{P_{i-1,j} + K_M} - k_{cat}^{ij} E_{i+1,j} \frac{P_{i,j}}{P_{i,j} + K_M}$$

j represents the linear enzyme steps in each pathway.

The final step of the biosynthetic pathway is modelled as:

$$\frac{dP_{i,n}}{dt} = k_{cat}^{in} E_{in} \frac{P_{i,n}}{P_{i,n} + K_M}$$

For our simulations, we considered each of the enzymes in the concurrent pathways equivalent in terms of their expression and kinetic parameters. Thus, $k_{cat}^{ij} = k_{cat}$ for all i, j.

Model setup and calibration

For our model, we take the kinetic parameters and gene lengths of an enzyme in amino acid biosynthesis pathway (histidine synthase).

We also directly use the stochastic gene expression parameters reported by Golding *et al.* 2005. However, the report is for a synthetic promoter and not a natural biosynthetic enzyme, and hence we have no corresponding measure of metabolite threshold. In order to estimate a threshold, for the given gene expression parameters, we firstly found the mean protein production trace from several Monte Carlo sampling of protein

production profiles. We next assumed that protein levels on average would roughly

double from the value at birth, before being halved again, post division. Using this, we

chose a 30-minute window on the mean protein trace where the protein roughly

doubles. We then compute the maximum amount of metabolite produced, using set

kinetic parameters and the mean protein profile. This value was then used as the

starting value of the metabolite threshold.

Having found an initial estimate of the threshold, we next varied threshold around the

value to find the effect of the same on the resultant distribution of cell generation times.

We also varied the gene expression parameters to find the major changes due to these

variations. We finally chose a threshold = 2e7 for Adder, and 4e7 for Sizer and

proceeded with the rest of the simulations.

Model Simulation

We start the simulation of the model with the obtained parameters, however for the

lack of knowledge of any initial conditions we start with all zero. As we start the

simulation, firstly stochastic mRNA and protein profiles are generated based upon the

set gene expression parameters, which are then used as time-dependent parameters

while solving the system of coupled ODE numerically using ode15s in MATLAB. When

all the p metabolites have met their production requirements, the events function is

triggered, which terminates the evaluation of the ODE, and marks the cell division. The

difference between the end and start times of the ODE solution gives us the growth

duration or generation time.

Before we start recording values from the simulation, we allow the system to saturate

by letting the cell run through 100 divisions or generations, following only one of the

daughter cells, starting from the zero initial values. Thereafter, we simulate an

exponential population growth, by simulation all the daughter cells, till 13 generations,

starting from 1 cell, and hence obtaining growth data for $2^{13} - 1 = 8191$ cells. We use

the generation times to compute the growth rate of the population, as described in the

next section. We repeat the simulations with 3 or 5 unique seed values to obtain

concordant observations.

Computing growth rate from generation times

From our simulations, we obtain the birth and division time of each simulated cell. If we sort the cells in the ascending order of their birth and consider each birth event as a unit increment to the population, it gives us the microbial population growth curve. The points appear as a straight line in a semi-log plot, which eventually saturates since there are no more cell divisions. To extract the growth rate, we consider only the first half of the dataset, to obtain the cells in a purely exponential state. We plot the best fit line on the semi-log plot and obtain the slope of the line, as the population growth rate. In order to improve the estimate and make use of the entire dataset, we permute the order of the cells during population growth and compute the growth rate. We iterate 100 times and take their average as the population growth rate. We report the final growth rate of the population for each simulated condition by computing the mean over all the unique seed values simulated.

Acknowledgements.

We are grateful to Christopher Marx for valuable discussions and feedback.

The authors also acknowledge the ICTS-ICTP Quantitative Systems Biology Schools 2015 and 2017 and the ICTS Bangalore School on Population Genetics and Evolution 2020, for the learning opportunity and exposure they provided.

D.D. was supported by funding from the Department of Science & Technology, Innovation in Science Pursuit for Inspired Research (DST-INSPIRE), Government of India.

We declare that we have no competing interests.

D.D. and S.S. conceptualised the study. D.D. and S.S. developed the theory. D.D. performed the simulations. D.D. wrote the manuscript with support from S.S.

References

1. Lipschultz F, Zafiriou OC, Wofsy SC, McElroy MB, Valois FW, Watson SW. 1981. Production of NO and N2O by soil nitrifying bacteria. Nature 294:641–643.

2. Falony G, Calmeyn T, Leroy F, De Vuyst L. 2009. Coculture Fermentations of Bifidobacterium Species and Bacteroides thetaiotaomicron Reveal a Mechanistic Insight into the Prebiotic Effect of Inulin-Type Fructans. Appl Environ Microbiol 75:2312–2319.

3. Rivière A, Gagnon M, Weckx S, Roy D, De Vuyst L. 2015. Mutual Cross-Feeding Interactions between Bifidobacterium longum subsp. longum NCC2705 and Eubacterium rectale ATCC 33656 Explain the Bifidogenic and Butyrogenic Effects of Arabinoxylan Oligosaccharides. Appl Environ Microbiol 81:7767–7781.

4. Moens F, Verce M, De Vuyst L. 2017. Lactate- and acetate-based cross-feeding interactions between selected strains of lactobacilli, bifidobacteria and colon bacteria in the presence of inulin-type fructans. Int J Food Microbiol 241:225–236.

5. Smith NW, Shorten PR, Altermann E, Roy NC, McNabb WC. 2019. The Classification and Evolution of Bacterial Cross-Feeding. Front Ecol Evol 7:153.

6. Hay ME, Parker JD, Burkepile DE, Caudill CC, Wilson AE, Hallinan ZP, Chequer AD. 2004. Mutualisms and Aquatic Community Structure: The Enemy of My Enemy Is My Friend. Annu Rev Ecol Evol Syst 35:175–197.

7. Smith P, Schuster M. 2019. Public goods and cheating in microbes. Curr Biol 29:R442–R447.

8. Harcombe W. 2010. Novel cooperation experimentally evolved between species. Evolution 64:2166–72.

9. Wintermute EH, Silver PA. 2010. Emergent cooperation in microbial metabolism. Mol Syst Biol 6:407.

10. Mee MT, Collins JJ, Church GM, Wang HH. 2014. Syntrophic exchange in synthetic

646   microbial communities. Proc Natl Acad Sci 111:E2149–E2156.

647   11.   Pande S, Merker H, Bohl K, Reichelt M, Schuster S, de Figueiredo LF, Kaleta C, Kost
648   C. 2014. Fitness and stability of obligate cross-feeding interactions that emerge
649   upon gene loss in bacteria. Isme J 8:953–962.

650   12.   Shou W, Ram S, Vilar JMG. 2007. Synthetic cooperation in engineered yeast
651   populations. Proc Natl Acad Sci 104:1877–1882.

652   13.   Kolter R, Greenberg EP. 2006. The superficial life of microbes. Nature 441:300–
653   302.

654   14.   Widder S, Allen RJ, Pfeiffer T, Curtis TP, Wiuf C, Sloan WT, Cordero OX, Brown SP,
655   Momeni B, Shou W, Kettle H, Flint HJ, Haas AF, Laroche B, Kreft J-U, Rainey PB,
656   Freilich S, Schuster S, Milferstedt K, van der Meer JR, Großkopf T, Huisman J, Free
657   A, Picioreanu C, Quince C, Klapper I, Labarthe S, Smets BF, Wang H, Soyer OS.
658   2016. Challenges in microbial ecology: building predictive understanding of
659   community function and dynamics. ISME J 10:2557–2568.

660   15.   Cavaliere M, Feng S, Soyer OS, Jiménez JI. 2017. Cooperation in microbial
661   communities and their biotechnological applications. Environ Microbiol 19:2949–
662   2963.

663   16.   Jagmann N, Philipp B. 2014. Design of synthetic microbial communities for
664   biotechnological production processes. J Biotechnol 184:209–218.

665   17.   Johns NI, Blazejewski T, Gomes ALC, Wang HH. 2016. Principles for designing
666   synthetic microbial communities. Curr Opin Microbiol 31:146–153.

667   18.   Stolyar S, Van Dien S, Hillesland KL, Pinel N, Lie TJ, Leigh JA, Stahl DA. 2007.
668   Metabolic modeling of a mutualistic microbial community. Mol Syst Biol 3:92.

669   19.   Zomorrodi AR, Segrè D. 2016. Synthetic Ecology of Microbes: Mathematical
670   Models and Applications. J Mol Biol 428:837–861.

671   20.   Khandelwal RA, Olivier BG, Röling WFM, Teusink B, Bruggeman FJ. 2013.
672   Community Flux Balance Analysis for Microbial Consortia at Balanced Growth.
673   PLoS One 8:64567.

21.   Zomorrodi AR, Maranas CD. 2012. OptCom: A Multi-Level Optimisation

Framework for the Metabolic Modeling and Analysis of Microbial Communities.
PLoS Comput Biol 8:e1002363.

22. Klitgord N, Segrè D. 2010. Environments that Induce Synthetic Microbial
Ecosystems. PLoS Comput Biol 6:e1001002.

23. Chan SHJ, Simons MN, Maranas CD. 2017. SteadyCom: Predicting microbial
abundances while ensuring community stability. PLOS Comput Biol 13:e1005539.

24. Kalisky T, Dekel E, Alon U. 2007. Cost–benefit theory and optimal design of gene
regulation functions. Phys Biol 4:229–245.

25. Semsey S, Jauffred L, Csiszovszki Z, Erdossy J, Steger V, Hansen S, Krishna S. 2013.
The effect of LacI autoregulation on the performance of the lactose utilisation
system in Escherichia coli. Nucleic Acids Res 41:6381–6390.

26. Weiße AY, Oyarzún DA, Danos V, Swain PS. 2015. Mechanistic links between
cellular trade-offs, gene expression, and growth. Proc Natl Acad Sci 112:E1038–
E1047.

27. Scott M, Gunderson CW, Mateescu EM, Zhang Z, Hwa T. 2010. Interdependence of
Cell Growth and Gene Expression: Origins and Consequences. Science (80-)
330:1099–1102.

28. Basan M, Hui S, Okano H, Zhang Z, Shen Y, Williamson JR, Hwa T. 2015. Overflow
metabolism in Escherichia coli results from efficient proteome allocation. Nature
528:99–104.

29. Scott M, Hwa T. 2011. Bacterial growth laws and their applications. Curr Opin
Biotechnol 22:559–565.

30. Edwards JS, Palsson BO. 2000. The Escherichia coli MG1655 in silico metabolic
genotype: Its definition, characteristics, and capabilities. Proc Natl Acad Sci
97:5528–5533.

31. Feist AM, Scholten JCM, Palsson BØ, Brockman FJ, Ideker T. 2006. Modeling
methanogenesis with a genome-scale metabolic reconstruction of
Methanosarcina barkeri. Mol Syst Biol 2.

32. Durot M, Bourguignon P-Y, Schachter V. 2009. Genome-scale models of bacterial

metabolism: reconstruction and applications. FEMS Microbiol Rev 33:164–190.

33.   Borodina I. 2005. Genome-scale analysis of Streptomyces coelicolor A3(2)

metabolism. Genome Res 15:820–829.

34.   Gu C, Kim GB, Kim WJ, Kim HU, Lee SY. 2019. Current status and applications of

genome-scale metabolic models. Genome Biol 20:121.

35.   Elowitz MB, Levine AJ, Siggia ED, Swain PS. 2002. Stochastic gene expression in a

single cell. Science (80-) 297:1183–1186.

36.   Kiviet DJ, Nghe P, Walker N, Boulineau S, Sunderlikova V, Tans SJ. 2014.

Stochasticity of metabolism and growth at the single-cell level. Nature 514:376–

379.

37.   Ferguson ML, Le Coq D, Jules M, Aymerich S, Radulescu O, Declerck N, Royer CA.

2012. Reconciling molecular regulatory mechanisms with noise patterns of

bacterial metabolic promoters in induced and repressed states. Proc Natl Acad Sci

109:155–160.

38.   Silander OK, Nikolic N, Zaslaver A, Bren A, Kikoin I, Alon U, Ackermann M. 2012. A

Genome-Wide Analysis of Promoter-Mediated Phenotypic Noise in Escherichia

coli. PLoS Genet 8:e1002443.

39.   Tonn MK, Thomas P, Barahona M, Oyarzún DA. 2019. Stochastic modelling reveals

mechanisms of metabolic heterogeneity. Commun Biol 2:108.

40.   Golding I, Paulsson J, Zawilski SM, Cox EC. 2005. Real-time kinetics of gene

activity in individual bacteria. Cell 123:1025–1036.

41.   Feist AM, Palsson BO. 2010. The biomass objective function. Curr Opin Microbiol

13:344–349.

42.   Taheri-Araghi S, Bradde S, Sauls JT, Hill NS, Levin PA, Paulsson J, Vergassola M,

Jun S. 2015. Cell-Size Control and Homeostasis in Bacteria. Curr Biol 25:385–391.

43.   Sauls JT, Li D, Jun S. 2016. Adder and a coarse-grained approach to cell size

homeostasis in bacteria. Curr Opin Cell Biol 38:38–44.

44.   Wallden M, Fange D, Lundius EG, Baltekin Ö, Elf J. 2016. The Synchronisation of

Replication and Division Cycles in Individual E. coli Cells. Cell 166:729–739.

45.   Campos M, Surovtsev I V., Kato S, Paintdakhi A, Beltran B, Ebmeier SE, Jacobs-
Wagner C. 2014. A Constant Size Extension Drives Bacterial Cell Size Homeostasis.
Cell 159:1433–1446.

46.   Pugatch R. 2015. Greedy scheduling of cellular self-replication leads to optimal
doubling times with a log-Frechet distribution. Proc Natl Acad Sci 112:2611–
2616.

47.   Kennard AS, Osella M, Javer A, Grilli J, Nghe P, Tans SJ, Cicuta P, Cosentino
Lagomarsino M. 2016. Individuality and universality in the growth-division laws
of single E. coli cells. Phys Rev E 93:012408.

48.   Wang P, Robert L, Pelletier J, Dang WL, Taddei F, Wright A, Jun S. 2010. Robust
Growth of Escherichia coli. Curr Biol 20:1099–1103.

49.   Monod J. 1949. The Growth of Bacterial Cultures. Annu Rev Microbiol 3:371–394.

50.   Morris JJ, Lenski RE, Zinser ER. 2012. The Black Queen Hypothesis: Evolution of
Dependencies through Adaptive Gene Loss. MBio 3:e00036-12-e00036-12.

51.   Osella M, Nugent E, Cosentino Lagomarsino M. 2014. Concerted control of
Escherichia coli cell division. Proc Natl Acad Sci 111:3431–3435.

52.   Iyer-Biswas S, Wright CS, Henry JT, Lo K, Burov S, Lin Y, Crooks GE, Crosson S,
Dinner AR, Scherer NF. 2014. Scaling laws governing stochastic growth and
division of single bacterial cells. Proc Natl Acad Sci 111:15912–15917.

53.   Soifer I, Robert L, Amir A. 2016. Single-Cell Analysis of Growth in Budding Yeast
and Bacteria Reveals a Common Size Regulation Strategy. Curr Biol 26:356–361.

54.   Coles S. 2001. An Introduction to Statistical Modeling of Extreme Values. Springer
London, London.

55.   Micali G, Grilli J, Osella M, Cosentino Lagomarsino M. 2018. Concurrent processes
set E. coli cell division. Sci Adv 4:eaau3324.

56.   Micali G, Grilli J, Marchi J, Osella M, Cosentino Lagomarsino M. 2018. Dissecting
the Control Mechanisms for DNA Replication and Cell Division in E. coli. Cell Rep

25:761-771.e4.

57.   Thomas P, Terradot G, Danos V, Weiße AY. 2018. Sources, propagation and
consequences of stochasticity in cellular growth. Nat Commun 9:4528.

58.   Trueba FJ. 1982. On the precision and accuracy achieved by Escherichia coli cells
at fission about their middle. Arch Microbiol 131:55–59.

59.   Huh D, Paulsson J. 2011. Non-genetic heterogeneity from stochastic partitioning
at cell division. Nat Genet 43:95–100.

60.   Jacob F, Monod J. 1961. Genetic regulatory mechanisms in the synthesis of
proteins. J Mol Biol 3:318–356.

61.   Zheng Y, Szustakowski JD, Fortnow L, Roberts RJ, Kasif S. 2002. Computational
Identification of Operons in Microbial Genomes. Genome Res 12:1221–1230.

62.   Price MN, Arkin AP, Alm EJ. 2006. The Life-Cycle of Operons. PLoS Genet 2:e96.

63.   Price MN, Huang KH, Arkin AP, Alm EJ. 2005. Operon formation is driven by co-
regulation and not by horizontal gene transfer. Genome Res 15:809–819.

64.   Rocha EPC. 2008. The Organization of the Bacterial Genome. Annu Rev Genet
42:211–233.

65.   Sabatti C, Rohlin L, Oh M-K, Liao JC. 2002. Co-expression pattern from DNA
microarray experiments as a tool for operon prediction. Nucleic Acids Res
30:2886–2893.

66.   Sneppen K, Pedersen S, Krishna S, Dodd I, Semsey S. 2010. Economy of Operon
Formation: Cotranscription Minimizes Shortfall in Protein Complexes. MBio 1.

67.   Ray JCJ, Igoshin OA. 2012. Interplay of Gene Expression Noise and Ultrasensitive
Dynamics Affects Bacterial Operon Organization. PLoS Comput Biol 8:e1002672.

68.   Nilsson AI, Koskiniemi S, Eriksson S, Kugelberg E, Hinton JCD, Andersson DI.
2005. Bacterial genome size reduction by experimental evolution. Proc Natl Acad
Sci 102:12112–12116.

69.   Dufresne A, Garczarek L, Partensky F. 2005. Accelerated evolution associated
with genome reduction in a free-living prokaryote. Genome Biol 6:R14.

70.   Giovannoni SJ, Tripp HJ, Givan S, Podar M, Vergin KL, Baptista D, Bibbs L, Eads J,
789   Richardson TH, Noordewier M, Rappé MS, Short JM, Carrington JC, Mathur EJ.
790   2005. Genome streamlining in a cosmopolitan oceanic bacterium. Science (80-)
791   309:1242–1245.

792   71.   D'Souza G, Waschina S, Pande S, Bohl K, Kaleta C, Kost C. 2014. Less is more:
793   Selective advantages can explain the prevalent loss of biosynthetic genes in
794   bacteria. Evolution (N Y) 68:2559–2570.

795   72.   D'Souza G, Kost C. 2016. Experimental Evolution of Metabolic Dependency in
796   Bacteria. PLOS Genet 12:e1006364.

797   73.   Boer VM, Crutchfield CA, Bradley PH, Botstein D, Rabinowitz JD. 2010. Growth-
798   limiting Intracellular Metabolites in Yeast Growing under Diverse Nutrient
799   Limitations. Mol Biol Cell 21:198–211.

800   74.   Katzianer DS, Wang H, Carey RM, Zhu J. 2015. "Quorum Non-Sensing": Social
801   Cheating and Deception in Vibrio cholerae. Appl Environ Microbiol 81:3856–
802   3862.

803   75.   Trivers RL. 1971. The Evolution of Reciprocal Altruism. Q Rev Biol 46:35–57.

804   76.   Travisano M, Velicer GJ. 2004. Strategies of microbial cheater control. Trends
805   Microbiol 12:72–78.

806   77.   Hillesland KL, Stahl DA. 2010. Rapid evolution of stability and productivity at the
807   origin of a microbial mutualism. Proc Natl Acad Sci 107:2124–2129.

808   78.   Morris BEL, Henneberger R, Huber H, Moissl-Eichinger C. 2013. Microbial
809   syntrophy: interaction for the common good. FEMS Microbiol Rev 37:384–406.

810   79.   D'Souza G, Waschina S, Kaleta C, Kost C. 2015. Plasticity and epistasis strongly
811   affect bacterial fitness after losing multiple metabolic genes. Evolution (N Y)
812   69:1244–1254.

813   80.   Sandoz KM, Mitzimberg SM, Schuster M. 2007. Social cheating in Pseudomonas
814   aeruginosa quorum sensing. Proc Natl Acad Sci 104:15876–15881.

815   81.   Cooper S, Helmstetter CE. 1968. Chromosome replication and the division cycle of
816   Escherichia coli. J Mol Biol 31:519–540.

817   82.   Donachie WD. 1968. Relationship between Cell Size and Time of Initiation of DNA
818   Replication. Nature 219:1077–1079.

819   83.   Zomorrodi AR, Segrè D. 2017. Genome-driven evolutionary game theory helps
820   understand the rise of metabolic interdependencies in microbial communities.
821   Nat Commun 8:1563.

Figure 1

Figure – 1

Model Schematic and behaviour of internal variables

a) Our model implements a "metabolite adder/sizer" model, where in we reinterpret
the empirical size-based adder and sizer laws, in terms of the metabolites necessary for
cell size increase. Such that cell division is triggered upon production of these
metabolites in required amounts and stoichiometry. Always a fixed amount in case of
Adder, and the balance amount in case of the Sizer.

b) Model Schematic: Bacteria uptakes the substrate molecules from the environment
and processes it via metabolic pathways to produce all the necessary metabolites for
survival. Some (p) metabolites (coloured squares) bottleneck cell growth, since
sufficient quantities cannot be produced in time. Our model assumes a simplified
arrangement for all these limiting metabolite anabolic pathways, as linear enzyme
cascades with n enzyme steps, all fed by a common substrate supplied at a constant rate.
We consider all enzymes are equivalent in terms of expression and kinetic properties
and are expressed in stochastic bursts.

c) Considering our model for p=2 limiting metabolites and the metabolic Sizer law. The
model implements stochastic burst like gene expression as a random process of
transcription bursts switching on and off. During the burst, mRNA are produced, which
act as template for protein translation. Both mRNA and proteins decay stochastically
but mRNA decay much faster. The varying protein enzymes allow the production of the
required metabolites at varying rates. According to Sizer, cell divides when both the
metabolites cross the threshold (dotted line). Post division, cellular contents are halved.

Figure -2

Figure – 2

Model properties

a) Distribution of cell Generation times obtained from model simulation is best fitted by
GEV (specifically Freschet) distributions.

b) Comparison of distribution of generation times for varying number of limiting
metabolites (p), at a high rate of substrate flux (S_{in} = 50000/sec). Inset shows the
distributions rescaled to their means. The distribution largely overlap with slight
changes in their statistical properties.

c) Comparison of distribution of generation times varying the substrate flux rate in the
simulations, for the case of $p = 3$ metabolite bottlenecks. Inset shows the distributions
rescaled to their means. The distribution largely overlap with slight changes in their
statistical properties.

d) Comparison of the mean growth rate of the simulated cells with varying number of
bottleneck metabolites (p), for different rates of substrate flux. The hyperbolic
relationship is reminiscent of Monod's Law.

Figure - 3

Figure – 3

Effect of different modelling criteria on the nature of cell generation times
observed

a) Comparison of distribution of generation times, while comparing between different
noise-generating modelling criteria: symmetric division, synchronised gene expression
and independent gene expression. Considering 3 bottleneck metabolites (p), and high
substrate flux ($S_{in} = 20000/sec$). Inset shows the distributions rescaled to the mean,
where they visually overlap, but the subtle statistical differences are revealed.

b) Comparison of distribution of generation times, for varying t_{OFF} parameter ($p =$
$3, and\ t_{ON} = 6\ min$). Inset shows the distributions rescaled to the mean. Skewness
increases with increasing t_{OFF}.

c) Comparison of distribution of generation times, for varying t_{ON} parameter ($p =$
$3, and\ t_{OFF} = 37\ min$). Inset shows the distributions rescaled to the mean. Skewness
increases with increasing t_{ON}.

d) Growth rates of simulated cells, when different combinations of gene expression
parameters are used, arranged in the ascending order of growth rate. Different
metabolite threshold levels are compared.

Figure -4

Figure – 4

Effect of metabolite uptake and secretion on simulated cells

a) Effect of metabolite uptake

i) Comparison of generation times for different rates of metabolite import. (Top) shows

low substrate flux case and (bottom) shows the high substrate flux case.

ii) Comparison of the mean growth rate of the simulated cells with varying rates of

metabolite import, for different rates of substrate flux.

iii) Comparison of the relative change in growth rate compared to the case of no feed,

due to varying metabolite feed rates, for different substrate flux (S_{in}).

b) Effect of metabolite secretion

i) Comparison of generation times for different ratios of metabolite secretion (fraction

of produced metabolite secreted). (Top) shows low substrate flux case and (bottom)

shows the high substrate flux case. Inset shows the distributions rescaled to their

means.

ii) Comparison of the mean growth rate of the simulated cells with varying ratios of

metabolite secretion, for different rates of substrate flux.

iii) Comparison of the relative change in growth rate compared to the case of no

secretion, due to varying metabolite secretion ratios, for different substrate flux (S_{in}).

Figure - 5

Figure – 5

Growth advantage of auxotrophy

a) Comparison of distribution of generation times, for an auxotroph that is fed the
missing metabolite at varying rates. (Top) shows the low substrate flux case, and
(bottom) shows the high substrate flux case.

b) Comparison of the mean growth rate of the simulated auxotroph (derived from
prototroph with $p = 3$) with varying feed rates for the missing metabolite, for different
rates of substrate flux (S_{in}). The grey and black dotted lines represent the prototrophs
with p = 3 and p = 2 bottleneck metabolites respectively.

c) Comparison of the relative change in the growth rate of the auxotrophs, when
compared with the original prototroph ($p = 3$), for varying rates of metabolite feed, and
substrate flux.

d) Comparison of the percentage increase in the growth rate upon loss of a bottleneck
pathway for different starting number of bottleneck pathways (p). Data for different
substrate flux rates is plotted.

Figure - 6

a)

b) i)

ii)

c) i)

ii)

Figure – 6

Reciprocal metabolic cross-feeding

a) Simplified Steady state version of our model framework. Comparison of the mean
generation times, against varying substrate flux, for the case of a prototroph, an
auxotroph and when the auxotroph produces twice the amount of one metabolite and
secretes half into the environment.

b) Kinetic advantage of auxotrophs over prototrophs for the benchmark case where
enzyme kinetic properties are doubled. c) Kinetic advantage of auxotrophs over
prototrophs for the case where gene expression parameters are altered to enable
doubling of protein production. Subplots i) and ii) represent the same kind of plots.

Plotted the mean secretion rate of metabolites (Y), for an input metabolite feed rate (X),
for different substrate flux rates (represented as different colours of the plotted
bubbles). The size of the circles represent the magnitude of the growth rate. In i) the
circles are filled if growth rate exceeds that of the prototroph ($p = 3$) when growth
without any metabolite feed. While in ii) circles are filled if the growth rate exceeds that
of the prototroph ($p = 3$), when fed with metabolite at the same rate as the auxotroph
being compared. The dotted diagonal line represents the line where secretion rate and
feed rates are equal.