Title:

CREB-binding protein gene, *HAC701*, negatively regulates WRKY45-dependent immunity
in rice

Running title:

HAC701-dependent immunity regulation

Authors:

Nino A. Espinas[1,3,*], Tu Ngoc Le[1], Saori, Miura[1], Yasuka Shimajiri[1,2], Ken Shirasu[3],
Hidetoshi Saze[1]

Affiliation:

[1]Plant Epigenetics Unit, Okinawa Institute of Science and Technology Graduate University,
Onna-son, Okinawa, Japan

[2]EditForce, Fukuoka, Japan

[3]RIKEN Center for Sustainable Resource Science, Yokohama, Kanagawa, Japan

Contact:

[*]Correspondence to: Nino A. Espinas (nino.espinas@riken.jp)

26 *Category:*

27 Biological Sciences, Plant Biology

SIGNIFICANCE

HAC701 is a member of CREB-binding protein (CBP) family that acts as transcriptional coactivator and acetyltransferase. However, little is known how it regulates innate immunity in plants. Herein we reported that rice *HAC701* suppresses WRKY45-dependent defense pathway. Our study showed that *HAC70* seemingly interacts genetically with *WRKY45* in rice to modulate immune responses against pathogens.

ABSTRACT

CREB-binding protein (CBP) is a known transcriptional coactivator and an acetyltransferase that functions in several cellular processes by regulating gene expression. However, how it functions in plant immunity remains unexplored. By characterizing *hac701*, we demonstrate that *HAC701* negatively regulates the immune responses in rice. *hac701* show enhanced disease resistance against a bacterial pathogen, *Pseudomonas syringae* pv. *oryzae* (*Pso*), which causes bacterial halo blight of rice. Our transcriptomic analysis revealed that rice *WRKY45*, one of the main regulators of rice immunity, is upregulated in *hac701* and possibly conferring the resistance phenotype against *Pso*. The morphological phenotypes of *hac701* single mutants were highly similar to *WRKY45* overexpression transgenic lines reported in previous studies. In addition, differentially expressed genes (DEGs) in *hac701* largely overlap with genes when *WRKY45* is overexpressed and chemically induced transiently. When we investigated for *cis*-elements found 1kb upstream of *WRKY45* gene and WRKY45-dependent DEGs, we found that *WRKY45* promoter contains the CRE motif, a possible target of HAC701-mediated regulation. Genome-wide H3K9 acetylation profiling showed depletion of acetylation at large set of genes in *hac701*. However, consistent with the upregulation of *WRKY45* gene expression, our ChIP-sequencing analysis demonstrated that regions of *WRKY45* promoter are enriched in H3K9 acetylation in *hac701* compared to the segregated wild type control. Finally, we show that *HAC701* may have roles in systemic immune signaling. We therefore propose that wild type *HAC701* negatively regulates *WRKY45* gene expression, thereby suppressing immune responses.

Keywords:

Oryza sativa, rice CREB-binding protein, *HAC701*, *WRKY45*, innate immunity,
Pseudomonas syringae pv. *oryzae* (*Pso*), *Magnaporthe*

INTRODUCTION

Histone acetyltransferases (HATs) are a diverse group of histone- and non-histone-modifying enzymes that contain multiple subunits to enable catalytic functions (1). These HATs primarily function as scaffolding bridges of proteins to form the transcriptional regulatory complex important for target recognition and subsequent substrate modification (1-3). The association of HATs to its interacting proteins and the existing state of the epigenome dictates the dual-switch functionality of either transcriptional activation or repression (1). They are considered integrators or adaptors as they were shown to interact with DNA-binding activators and the basal transcriptional machinery (4, 5).

In plant model Arabidopsis, HATs and histone deacetylases (HDACs) are grouped into four (Figure 1A) and three families, respectively (6). HATs consists of CBP, TAF1/TAFII250, GNAT, and MYST, while HDACs are RPD3/HDA1, HD2-like, and SIR2. A previous review of HATs and HDACs in plants listed about 12 putative HAT and 18 HDAC proteins exhibiting acetylation and deacetylation activities with differential site specificities (6). Work on rice (*Oryza sativa*), however, has just begun with a preliminary investigation of eight histone acetyltransferases (7, 8) and deacetylases (9).

Several studies have shown that histone acetylation and deacetylation control defense signaling in plants in response to phytohormone or pathogen application (10-12). More specifically, HATs and HDACs in rice model were shown to be responsive to abiotic stresses and can be modulated by phytohormones, thus implicating a role in biotic stresses (7, 13). Similar to Arabidopsis, HATs in rice are also classified into four families (Figure 1A) (7). In contrary, rice HDACs are represented in only two families, RPD3/HDA1-like and SIR2-like with no known rice member belonging to HD2 family (9).

OsHAC701 was reported (14) earlier as a putative rice CBP-related

acetyltransferase, herein after referred to as *HAC701*, of the p300/CBP acetyltransferase

(PCAT) family of proteins. The Arabidopsis genome contains five CBP genes, while rice

genome contains three (Figure 1A), having broad acetyltransferase specificity on histones (7).

Currently, there is no clear consensus on the similarity of biological functions of Arabidopsis

and rice homologs of CBP family proteins. In animals, p300 and CBP are paralogs and

originally described as transcriptional coactivator that exhibit histone acetyltransferase

activity on all four core histones specifically at H4 N-terminal tail sites K5, K8, K12, and

K16 (15, 16). Other sites include H3 sites K14, 23 (17), K18, K27 (18, 19), and K56 (20);

H2B sites are K12 and K15 (21). H3K9 is mainly acetylated by GCN5/PCAF (18), however

acetylation by p300/CBP was also reported (17, 22, 23). Its highly conserved function is

mainly found in multicellular organisms as it probably participates in complex physiological

processes acting as a limiting factor in various pathways due to its high cellular demand (14).

As HATs have been shown to be modulated by biotic stresses (24), we hypothesized

that it could be a potential system for studying rice immunity. Indeed, we show that *HAC701*

is involved in rice innate immunity. In this study, we have made CRISPR/Cas9 edited *hac701*

and showed that *HAC701* is apparently a negative regulator of immunity with the mutants

showing resistance against bacterial and fungal infection. Further transcriptomic investigation

showed that HAC701 possibly regulates *WRKY45* loci, one of the main regulators of rice

immunity pathways. In *hac701*, *WRKY45* and its direct and indirect gene targets were

upregulated upon pathogen inoculation. In addition, *hac701* phenocopies *WRKY45-*

overexpression (*WRKY45-ox*) lines, indicating a potential interaction between *HAC701* and

WRKY45-dependent immunity pathway in rice.

RESULTS

Characterization of a rice acetyltransferase mutant, *HAC701*.

We isolated null mutant lines by independently targeting two sites of *HAC701* gene (Fig. S1A-C; Fig. S1D-I) using CRISPR/Cas9 editing technology. From the 11 positive *HAC701* lines, we have isolated four biallelic homozygous mutant lines (Target2; Fig. S1D), which we utilized for experimentation including their segregating wild type siblings, 9-WT (Fig. 1B). The phenotype in terms of effective grain number and tiller number of mostly all biallelic homozygous mutants at T2 generation did not show impairment as compared to 9-WT (Fig. S1F). In addition, the T3 lines did not show any visible growth or developmental defect in height and photosynthesis (*e.g.* chlorosis phenotype) (Fig. S1G), respectively. However, the weights of 1000-grain T3 and T4 generations of mutants were increased compared to 9-WT (Fig. S1I). These results indicate that a loss of HAC701 does not cause severe developmental defects in rice, and that HAC701 may have a positive role in seed production.

orinline *HAC701* mutant rice is resistant to pathogen infection.

Our expression analysis of the rice HAT genes showed that HAC701 is significantly up-regulated by treatment of bacterial flg22 peptide, suggesting a potential role of HAC701 in plant defense against pathogens (Fig. S2). To investigate involvement of *HAC701* in basal defense in rice, we inoculated 9-WT and *HAC701* mutant lines, *9-12b$^{-/-}$* and *9-5$^{-/-}$* (hereafter collectively termed *hac701* unless otherwise specified) with mock (10 mM MgCl$_2$) and *Pseudomonas syringae* pv. *oryzae* (*Pso*) (OD = 0.2; resuspended in 10 mM MgCl$_2$) for 72 h (Fig. 1D). *Pso* is a causative agent of halo blight in rice characterized by brown lesions and yellow halo-like blotches on leaves (25). Infection of *Pso* showed that *hac701* (*9-12a$^{-/-}$*, *9-12b$^{-/-}$* and *9-5$^{-/-}$*) were resistant to *Pso*-treatment as compared to 9-WT (Fig. 1C, S3). These data suggest that *HAC701* might be involved in the regulation of rice innate immunity acting

in its capacity as a transcriptional coactivator and/or as an acetyltransferase. We further tested

whether *hac701* also show resistance phenotype when infected with a rice blast pathogen,

Magnaporthe oryzae. Although the data were not statistically significant, results showed a

resistant phenotype tendency in *hac701* (Fig. 1D). Overall, these indicate that *HAC701* plays

a negative role in rice innatee immunity.

Transcriptome profiling of the rice-*Pso* pathosystem.

To profile the genome-wide effect of *HAC701* mutation in rice innate immunity, we

performed RNA-sequencing on mock- and *Pso*-treated 9-WT and *hac701* using local leaf

tissues (Fig. S4; Table S2). We first characterized the effect of *Pso* infection on 9-WT and

analysis of the highly variable genes in mock- and *Pso*-treated 9-WT plants showed gene

clusters that were dependent on *Pso* induction alone (Fig. S4; Fig. S5). The top 10 highly

variable genes across the 9-WT samples showed features involved in tolerance and/or

resistance such as disease resistance and stress tolerance mostly by catalysis of primary and

secondary metabolism (Fig. S5). This includes well-documented genes involved in plant

resistance against pathogens, such as tryptophan decarboxylase 1 (Os08g0140300) that

catalyzes the production of serotonin via conversion of tryptophan to tryptamine in rice (26),

which has been implicated to confer resistance in rice infected with *Bipolaris oryzae* (27).

Another gene, naringenin 7-O-methyltransferase (Os12g0240900), that catalyzes the

production of rice phytoalexin sakuranetin has been reported to participate in rice defenses

via JA signaling (28, 29). Lignin and phytoalexin encoding genes for instance laccases (*e.g.*

Os12g0258700, Os11g0641500) and cytochrome P450s (*e.g.* Os07g0218700,

Os08g0508000) are metabolic products known to be involved in plant defenses as well (30-

32). Gene Ontology (GO) analysis of upregulated genes in *Pso*-infected 9-WT plants

revealed enrichment of genes involved in response to biotic stimulus (GO terms: Response to

wounding, Response to other organism, Cell wall macromolecule catabolic process, *etc.*)

(Fig. S6). Both up and downregulated genes showed enrichment of transcriptional machinery

regulation, a possible consequence of defense-related transcriptional reprogramming

processes during bacterial invasion. GO analysis also identified pathways associated with

upregulated gene networks involved in plant defense such as diterpene phytoalexin

biosynthetic pathway and chorismate biosynthesis. Thus, the rice-*Pso* pathosystem is a

functional system that can be utilized to analyze defense responses of *Pso*-infected rice as

well as *Pso* pathogenicity.

Potentiated WRKY45 gene in *hac701* provides resistance against *Pseudomonas* and

***Magnaporthe* infections.** To investigate the molecular basis of resistance phenotype

observed in *Pso*-infected *hac701*, we further analyzed our transcriptome data and found 141

upregulated genes in *hac701* (Fig. S7A; Table S3). We refer to these genes as *HAC701*-

repressed genes. GO analysis of *HAC701*-repressed genes showed enrichment of processes

involved in response to xenobiotic stimuli and defense response (GO terms: Response to

xenobiotic stimulus, Response to wounding, Regulation of defense response, Jasmonic acid

mediated signaling pathway, Defense response, *etc.*). These genes were normally not

upregulated in the 9-WT background. On the other hand, 336 upregulated genes in the 9-WT

background alone referred to as *HAC701*-dependent genes did not show a very distinct

enrichment of genes in defense responses rather showed a more general biological and

physiological processes (GO terms: Mitochondrial respiratory chain complex IV assembly,

Alanine transport, Cellular response to heat, *etc.*) (Fig. S7A; Table S4). We also looked into

the features of 31 downregulated genes in the mutant background referred to as *HAC701*-

enhanced genes (Fig. S7B; Table S3). These genes were not normally downregulated in the

9-WT background. *HAC701*-enhanced genes contained genes enriched in photosynthetic and

response to abiotic stress terms (GO terms: Glycerolipid biosynthetic pathway, Response to

cold, Photosynthesis, *etc.*). In addition, the remaining 46 downregulated genes in the 9-WT

background alone also showed a more general biological processes with no hint of

enrichment in defense response against pathogen infection (GO terms: Chloroplast

organization, Gluconeogenesis, Organelle fission, *etc.*) (Fig. S7B; Table S4).

We were intrigued what genes can be upregulated in *HAC701* mutant background

without *Pso*-treatment. Thus, we overlapped the list of genes derived from mock and *Pso*-

treated mutants, and found *WRKY45* (*OsWRKY45*) upregulated in the *hac701* background

(Fig. 2A; Table S5, S6). *WRKY45* was also identified as one of the 141 upregulated genes in

Pso-infected *HAC701* mutant (Fig. S7A). There was no candidate gene found downregulated

in *HAC701* mutant background alone (Fig. 2A). We then checked whether there were

differences in the upregulation strength of *WRKY45* in the *HAC701* mutation alone,

compared to *HAC701* mutation combined with *Pso*-treatment. Transcriptome data revealed

that *WRKY45* expression was increased by about 20% change upon *Pso*-treatment (Fig. 2B).

These results suggest that upregulation of *WRKY45* expression is partially dependent on

HAC701 expression. Additional *Pso*-treatment further upregulated *WRKY45* and this subtle

increase in expression allowed the co-expression of possibly direct and indirect genes

regulated by *WRKY45*.

***hac701* phenocopy *WRKY45* overexpression transgenic plants.**

To determine whether the upregulation of *WRKY45* in *hac701* is relevant to rice with

overexpressed *WRKY45* gene, we compared the differentially expressed genes (DEGs) in our

hac701 with *WRKY45* overexpression (*WRKY45-ox*) lines previously reported (in these

studies: 33, 34). Our results indicated a striking similarity for targeted genes from rice

samples despite the fact that they were inoculated with two different types of pathogens, a

bacterial (*Pso*; this project) and the fungal pathogen *Magnaporthe* (33, 34) and minor

differences in sampling time periods (Fig. 3). About 43.3% (286/660 genes) of *HAC701*

mutant-DEGs in *Pso* were overlapped with BTH (Benzothiadiazole)-responsive genes (*i.e.*

BTH is a synthetic analog of salicylic acid, SA) (Fig. 3A; Table S5; Table S7), while about

12.4% (82/660 genes) of the *HAC701* mutant-DEGs in *Pso* were overlapped with BTH- and

WRKY45-regulated genes (Fig. 3B; Table S8). In addition, genes upregulated in plants

expressing DEX-induced myc-tagged WRKY45 protein in rice were mostly found in

HAC701 mutant-DEGs in *Pso* (75% or 9/12 genes; Table S9). Our analysis also showed that

at most 9.1 % (60/660 genes) of the *HAC701* mutant-DEGs in *Pso* were regulated in BTH-

and NPR1-dependent manner (Fig. 3B; Table S10). We then analyzed the 72 *HAC701*

mutant-DEGs in *Pso* that were exclusively regulated by BTH- and WRKY45 as well as the

other 10 genes jointly regulated by WRKY45 and NPR1 under BTH treatment (Figure 3B-

D). We found that most of these *HAC701* mutant genes were genuine WRKY45-regulated

genes (Fig. 3C-D) (33, 34). Analysis of these WRKY45-dependent defense-related genes

showed that in the presence of *Pso*-infection, number of glutathione-S-transferase (GST)

gene, were more upregulated in 9-WT than in *hac701* (Fig. 3D). These results were also

consistent with the result that *OsWRKY62*, a known negative regulator and direct target of

OsWRKY45 (34), had lower transcription level in *HAC701* mutants (Fig. 3D). It is interesting

that genes in JA biosynthesis, signaling and perception tend to be more highly expressed in

HAC701 mutant than 9-WT (Fig. 3D). In addition, *HAC701* mutant genes that were found

regulated in BTH and *WRKY45* indeed contain W-box motifs (35), raising more the

possibility that these genes are direct WRKY45 targets (72/82 or 87.8%; Table S11).

Meanwhile, all 10 genes that were commonly regulated in BTH, *WRKY45*, and *NPR1* (Fig. 3)

contain W-box motifs as well (10/10 or 100%; Table S12). Taken together, these results

indicate that upregulation of *WRKY45* as an outcome of *HAC701* mutation resulted in

expression profile of genes almost identical to overexpressing *WRKY45* transgenic lines as

well as to DEX-inducible *WRKY45* expression system. These also suggest that a significant

fraction of *HAC701* mutant differentially expressed genes (DEGs) are probably WRKY45-

targeted, given the presence of W-box motif.

The upregulation of *WRKY45* in *hac701* did not result in any visible morphological

defects (Fig. S2G). Previous studies showed that *WRKY45-ox* had no obvious autoimmunity

phenotype, although there was a slight decrease in height observed as compared to wild type

in response to growth conditions (33, 34). In this study, WRKY45-dependent resistance

against *Pso* was observed in *hac701* (Fig. 1D, Fig. S3). Furthermore, these *HAC701* mutants

also showed resistant tendencies against *Magnaporthe* infection (Fig. 1E). *WRKY45-ox*

treated with *Magnaporthe* showed resistance, while a BTH-induction in knockdown *WRKY45*

mutants did not rescue the *Magnaporthe* resistance and remained susceptible (34). Together

these results suggest that morphological and disease phenotypes of *hac701* resembled that of

WRKY45-ox indicating that *HAC701* mutation indeed upregulates *WRKY45* gene that

prompts WRKY45-dependent defense pathway into action leading to resistance against

bacterial and fungal pathogens.

WRKY45 promoter contains CRE motif.

Rice HAC701 and two others, HAC703 and HAC704, belong to the CBP [Cyclic adenosine

monophosphate response element-binding protein (CREB) Binding Protein] family group of

proteins that is known to bind CREB transcription factor through its KIX domain (7, 36).

Therefore, we examined *in silico* the members of rice CBP family for the presence of KIX

domains using a KIX domain database (37). Our results showed that only HAC701 and

HAC703 proteins contain the highly conserved KIX domain (Fig. S8). Although we have not

identified CREB transcription factor candidate in rice so far, we further examined whether

WRKY45 promoter contains CRE motif that could possibly be targeted via KIX domain and

CREB transcription factor-dependent regulation. To our surprise, we found a full CRE motif

within 1 kb upstream of the start codon of *WRKY45* locus (Fig. S9A). Furthermore, we

checked the occurrence of CRE motif in *hac701* DEGs during infection and found that only

five genes including *WRKY45* (0.76%) contain full CRE motif (Fig. S9; Table S5). These

data show that *WRKY45* promoter could possibly be *cis*-regulated by HAC701 through its

KIX domain, while the occurrence of CRE motif is not necessarily associated with other

HAC701-regulated genes.

Genome-wide investigation of histone modifications and enriched gene targets in

***HAC701* mutant.**

The CREB-binding protein (CBP) is also known to regulate gene expression by acetylating

histone tails in both animals and plants (38). Thus, we performed chromatin

immunoprecipitation coupled to sequencing (ChIP-seq) analysis on histone markers

associated with active and repressive chromatin to examine the landscape of chromatin

modifications in *HAC701* mutant. We performed ChIP-seq analyses of H3K9 acetylation

(H3K9ac), H3K27 acetylation (H3K27ac), H3K9 di-methylation (H3K9me2), and H3K9 tri-

methylation (H3K9me3) on 9-WT and *9-12b*$^{-/-}$ mutant (Fig. S10; Table S13). Metaplot

profiles of these histone modifications were generated for genes, transposable elements

(TEs), and simple repeats (Fig. S11; Fig. S12). The enrichment profiles show that both

H3K9ac and H3K27ac accumulated in genes mainly along the transcriptional start site (TSS)

and tapering through the gene body regions. In contrast, we found that H3K9me2 and

H3K9me3 were highly depleted in the gene TSS region (Fig. S11A). As expected, H3K9 and

H3K27 acetylation marks were depleted in TEs, while H3K9 di- and tri-methylation marks

showed enrichment (Fig. S11B). In addition, enrichment analysis of simple repeats showed

similar profiles with TEs in both acetylation and methylation histone marks, although simple

repeats tend to be highly enriched in H3K9me2 at a narrow set of genomic loci (Fig. S12).

These results showed that our ChIP-seq profiles of histone modifications were largely

consistent with the previous report on rice ChIP-seq analysis (39).

The observed low enrichment of H3K9ac in *9-12b^{-/-}* mutant is a probable indication of

mutation on *HAC701* gene (Fig. S11A). Therefore, we further analyzed the significantly

enriched or depleted regions in *9-12b^{-/-}* by taking the enrichment ratio of 9-WT over *9-12b^{-/-}*

(wild type/*hac701*) in H3K9ac, H3K27ac, H3K9me2, and H3K9me3 modifications. We

found that H3K9ac fraction yielded peaks (265 regions) (Fig. 4A; Table S13). From the

metaplot analysis, it showed that *9-12b^{-/-}* has depleted H3K9ac enrichment in the 265 regions

identified containing putatively 263 genes (Fig. 4A; Table S14), which include *HAC704* and

HAC701 loci and a mediator gene, *MED11* (Fig. 4D; Fig. S13A). These 265 regions were not

associated with changes in other modifications in *9-12b-/-* mutant when compared to 9-WT

(Fig. 4B). These results suggest that HAC701 acetyltransferase may have a specificity to

H3K9 site. It also suggests that HAC701 probably autoacetylates its own promoter including

another member of CBP family, *HAC704*.

We then analyzed the GO enrichment features of the putative 263 genes found in

these 265 regions with H3K9ac changes (Fig. 4C; Table S14). The results indicate that these

genes function in processes involved in the flow of genetic information, DNA methylation in

the CHH-context, circadian rhythm, organellar transport, cell cycle and reproduction,

ubiquitination, and in a number of biosynthetic processes. It is also perhaps not surprising

that histone acetylation was also enriched. In addition, enrichment of genes involving

environmental responses included heat stress and innate immunity were observed, in which

about 9% (23/263) were possibly involved in defense responses. A large fraction of these

defense genes were identified as candidate sensors or decoys (e.g. Cyclin A1, Thioredoxin,

Jacalin-like lectin, *etc.*) that may act as negative regulators of rice effector-triggered

immunity (ETI-immunity) (40) (Fig. 4E; Fig. S13B). The distribution of these 265 regions in

H3K9ac ChIP revealed that a large portion were found in genes (73.58%) followed by

promoters (18.11%) (Fig. S14). It is also interesting to note that H3K9ac in these identified

regions was found in transposable elements (7.55%). Overall, these results demonstrate that

HAC701-dependent acetylation of H3K9 histone site might modulate transcription of decoy

genes that are monitored by NLR proteins in rice immunity. Additionally, H3K9ac assessed

in different regions of the rice genome showed that it is found mostly in genes and promoters

with high similarity in the previous study (39).

We proceeded by analyzing the general distribution of H3K9ac, H3K27ac, H3K9me2,

and H3K9me3 peaks in intergenic and intragenic regions of rice genome in 9-WT and *9-12b*⁻

⁻ mutant (Fig. 5A). A large proportion of H3K9ac and H3K27ac were concentrated on exonic

and intronic regions of the gene, while *HAC701* mutation resulted in the increase of

enrichment of H3K27ac genome-wide on all regions of the rice genome (Fig. 5A). It is also

worth mentioning that loss of *HAC701* increased the level of enrichment of these methylation

modifications in intergenic regions. These data suggest that the loss of *HAC701* leads to an

increase of a counter acetylation modification, H3K27ac, on a genome-wide scale.

To explain the disease resistance phenotype of *hac701* through WRK45-dependent

immune pathway, we investigated the genomic locus of rice *WRKY45* (Fig. 5B). The

upregulation of WRKY45 was accompanied by H3K9ac enrichment along the 1kb upstream

region of *WRKY45* gene in *9-12b*⁻/⁻ (Fig. 5B, C). As the CRE motif was specifically enriched

in the promoter region of *WRKY45* (Fig. S9A), we compared the enrichment of H3K9ac and

H3K27ac in *9-12b*⁻/⁻ *WRKY45* promoter. We found that H3K9ac, but not H3K27ac, was

highly enriched in this regulatory region (Fig. S15), while H3K27ac of *WRKY45* gene body

was less pronounced than H3K9ac. Overall, our ChIP-seq data are consistent with RNA-seq

data in showing that mutation in *HAC701* gene potentiated *WRKY45* locus by regulating the

enrichments of histone acetylation and methylation modifications on *WRKY45* promoter. We

also found that the CRE motif of *WRKY45* is modulated by H3K9ac and perhaps other

modifications, including those that are analyzed here.

Systemic gene expression in rice-*Pso* pathosystem. To explore the nature of systemic gene

expression in rice-*Pso* pathosystem, we performed RNA-sequencing on systemic or distal

tissues of 9-WT and *hac701* plants. The tissue samples were collected together with infected

samples (local samples) 72 hours after *Pso* infection without further bacterial treatment.

Genome-wide transcriptome analysis in systemic 9-WT samples showed that 24 genes were

differentially expressed, and among them were eight genes that were common to both local

and systemic tissues (Fig. S18A; Table S16). Comparative analysis also showed that although

there were substantial expression changes in systemic tissues, the number of DEGs was

highly reduced in systemic tissues than in local tissues. Among the eight DEGs, five of them

showed altered transcriptional expression from local to systemic tissues (Fig. S18B). These

genes were mannose-specific jacalin-related lectin/*OsJAC1* (Os12g0247700) (41), putative

cytochrome P450 (Os09g0275400), ATPase (Os07g0187400), GDP-L-galactose

phosphorylase (Os12g0190000), and terpene synthase (Os04g0344100), some with putative

functions in response to pathogen infection or stress. These results indicate that these genes

were mostly activated locally in the infection site and that any altered expression in distal

tissues may facilitate systemic form of resistance. To compare the effect of *HAC701* mutation

on the number of DEGs, we analyzed the MA-plots of systemic tissues of 9-WT and *hac701*.

Our results showed that the number of DEGs with significant expression (*i.e.* those that are in

red) was diminished in the mutant background implicating a possible role of *HAC701* in

systemic signaling in rice-*Pso* pathosystem (Fig. S19). Overall, these results indicate that

systemic tissues in *Pso*-challenged 9-WT plants have augmented transcriptional gene

expression that possibly aims to 'prime' distal tissues on the onset of secondary pathogen

attack. This also indicates that *HAC701* might potentially regulate systemic defenses through

an unknown mechanism at distal non-infected site in preparation for future infection

episodes.

DISCUSSION

Histone acetyltransferases or HATs and their complexes are involved in various biological

processes in the cell. Currently, studies have been focused on the roles of HATs specifically

members of the CREB-binding protein (CBP) family in regulating substrate specificity and

enzymatic activity such as acetylation (38). However, its involvement in molecular signaling

pathway leading to modulation of plant immunity remains to be explored. Therefore, it is

expected that the analysis of plant CBP will facilitate a broader understanding of regulation

of gene expression and protein function in plant immunity pathways.

In this study, we generated CRISPR/Cas9 mutants of the rice *HAC701* gene, a

member of the CBP family of HATs, since our results indicated its involvement in rice

immunity. By inoculating a rice compatible bacterial pathogen, *Pseudomonas syringae* pv.

oryzae (*Pso*), we showed that *hac701* are resistant to *Pso* proliferation and also showed

resistance tendency when infected with *Magnaporthe* (rice blast). These indicates that

HAC701 negatively regulates rice immunity responses to *Pso* and *Magnaporthe* pathogens.

When we analyzed our transcriptome data, it showed that this negative regulation of

immunity by *HAC701* is attributable to suppression of a major immunity pathway in rice

probably through WRKY45. In *hac701*, *WRKY45* gene expression is upregulated and it

seemed to be in a primed state enabling a further increase of expression upon bacterial

infection. In addition, both morphological phenotypes and target genes in *hac701* mimic

those in *WRKY45-ox* transgenic lines possibly indicating a genetic interaction of *HAC701* and

WRKY45 genes. Lastly, we showed in our ChIP-seq analysis that the promoter of *WRKY45* in

hac701 is enriched in H3K9ac coinciding with a *cis* regulatory CRE motif that might be

responsible for *WRKY45* upregulation and control.

In Arabidopsis, the p300/CBP acetyltransferase gene, *HAC1*, has been shown to

positively regulate defense priming only in repetitively abiotic-stressed plants (42). *hac1-1*

did not show any immunity related responses prior to abiotic stress applications indicating

that Arabidopsis *HAC1* requires several abiotic stress stimulations to activate its immunity.

However, this is not the case in *HAC701* in rice. We showed that the rice *HAC701* did not

need abiotic stress stimulation to be transcribed and to effect pathogen and disease specific

phenotypes (Fig. S1; Fig. 1D-E; Fig. S3). Rather, our results are consistent with the disease

phenotype exhibited by Arabidopsis mutant of GCN5, an acetyltransferase member of the

SAGA transcriptional coactivator complex catalyzing the H3K14 modification site and at the

same time influencing the acetylation of H3K9 (43). The *gcn5* showed upregulation of SA-

mediated immunity resulting in resistance against *Pseudomonas syringae* pv. *tomato*

infection. In our study, we found that WRKY45-dependent pathway confers resistance to

hac701. Consistent with the view that the rice hormone defense network does not support a

dichotomous role of salicylic acid (SA) and jasmonic acid/ethylene (JA/ET) phytohormones

in regulating rice immunity (44), we found no evidence in this study that SA-mediated

immunity played a central role in *hac701* resistance phenotype. However, we found JA-

related differential gene expression as components of WRKY45-dependent defense-related

genes (Fig. 3D). Indeed, JA biosynthesis is known as a downstream target of rice WRKY45

(45). In rice immunity model, both SA and JA/ET are considered effective in defense

responses against (hemi) biotrophic and necrotrophic pathogens (44, 46, 47). While SA-

mediated *AtNPR1* regulates majority, if not all, of Arabidopsis immune responses (48),

OsWRKY45 and *OsNPR1/NH1* are mostly independent of each other (34; Fig. S26).

Although, our results suggest that *hac701* DEGs partially overlapped with NPR1-dependent

pathways (Fig. 3B). Given that the rice system has a constitutive high levels of SA compared

to most plants including Arabidopsis and tobacco, and pathogen applications did not increase

its basal SA level (49, 50), our results are consistent with this report that SA in rice may not

directly act on pathogens, rather, it may function to prime endogenous defense pathways in

which *WRKY45* plays the central role. *OsWRKY45-ox* showed strong resistance against blast

pathogen, even though lacking the constitutive expression of defense genes prior to blast

infection (34). It could be that disruption of HAC701-dependent pathways through *HAC701*

mutation led to activation of WRKY45-dependent pathways in an already SA-primed rice

network. But, this needs to be further examined in detail. Our results are also consistent with

the disease phenotype presented by non-sense mutation in *Med15* gene, a subunit of Mediator

complex (51). The *med15b.D* wheat mutants were resistant to stem rust, and the authors

attribute a part of this to segmental coregulation in which a certain portion of the

chromosome containing NLR genes was differentially expressed in wild type compared to

mutant. While we did not observe any form of segmental coregulation in our analysis of

hac701, we found that mutations in *HAC701* resulted in downregulation of ETI components

notably the decoys/sensors (Fig. 4E).

OsHAC701, *AtHAC1*, *AtGCN5*, and *TaMed15*, play a central role in regulating the

transcriptional machinery. They likely do this regulation by associating themselves and their

complexes to RNA polymerase II and transcription factors as shown by their homologous

counterparts in animals (4, 14, 52, 53). This is possible because they are multidomain genes

that have the capacity to form and maintain large complexes. Os*HAC701*, *AtHAC1*, and

AtGCN5 are also involved in regulating abiotic stress responses suggesting that the

complexity and specificity of their activities are dependent on their interacting factors that

recruit them to the biological pathway itself (6, 7). Furthermore, majority of these genes,

OsHAC701, *AtHAC1*, and *TaMed15*, encode the KIX domain, a docking site for transcription

factors reported to be involved in plant immunity (54).

In line with the above results, we propose that HAC701 is a negative regulator of

WRKY45-dependent defense pathway in rice (Fig. S20). We suggest here two possible

mechanistic insights into HAC701 regulation of WRKY45 that need further experimental

evidence. First, similar to Med15, the HAC701-mediated immunity may require a loss or

truncation of HAC701 in regulating the transcription of *WRK45* locus. Given that the

promoter of *WRKY45* contains a rare CRE motif (Fig. S9), we assume that an unidentified

CREB-like transcription factor with regulatory potential is interacting with HAC701 through

its KIX domain (Fig. S8) and with the CRE motif of *WRKY45* promoter. This interaction may

also include other protein interactors specific to *WRKY45* locus regulation. From our results,

mutation of *HAC701* is enough to prime *WRKY45* and further addition of *Pso* enables *hac701*

to mount higher *WRKY45* transcription compared to 9-WT. These suggest to us that *WRKY45*

locus is poised for activation as suggested by our ChIP-seq analysis of untreated *hac701* (Fig.

5). The H3K9 acetylation we observed might not be catalyzed by HAC701, but may come

from other CBP family members (Fig. 1A) or other HATs. Although, we do not have ChIP-

seq data for pathogen inoculated *hac701*, our RNA-seq data suggest that the poised *WRKY45*

through *HAC701* mutation is enough to activate downstream defense likely through *WRK45*-

dependent signaling. Second, the decoys/sensors component that we found may act as

negative regulators as predicted by the Decoy model (Fig. 4E) (40, 55). It could be that

HAC701 maintains the acetylation of these decoys/sensors to equilibrium. Any disturbance in

their expression could possibly lead to activation of R genes monitoring these decoys/sensors,

which could in effect activate WRKY45-dependent immunity. Indeed, we found seven R

genes upregulated in *HAC701* mutation treated with *Pso* (Table S17). It is of course possible

that these two mechanisms are occurring simultaneously in conferring resistance, although

finer details of these mechanisms are still unknown.

479   The heterochromatic marker, H3K9me3, is common in animals, but rarely found in

Arabidopsis and in plants in general (56). Indeed, our analysis did not find the peaks in 9-WT

ChIP-seq, although we found 33 peaks in the *HAC701* mutant (Table S15). Most H3K9me3

enrichments were found in intergenic regions and in only two genes with intragenic

enrichments. Table S15 also shows genes nearest the H3K9me3 enrichment in the intergenic

regions. These results suggest that H3K9me3 is may be a rare modification and is possibly

modulated by HAC701 presence and activity in certain genomic regions.

486   Trade-off between growth and defense in plants is a common occurrence and often

considered inevitable (57, 58). However, we were surprised to find that *hac701* in the T2 up

until T4 generations do not have readily visible penalty yields (*e.g.* number of effective

grains and grain weight) (Fig. S2F-H). It could be that the primed state of *WRKY45* in *hac701*

has modulated the expression of resistance from affecting these agronomic traits.

Nonetheless, the mechanistic basis for this trade-off, if it truly exists, is not well understood

and needs further investigation.

493   We would like to emphasize that at the moment we do not have strong evidence that

HAC701 specifically catalyzes H3K9 acetylation. Although, the H3K9ac antibody used in

this study detected H3K9ac decrease of numerous loci in *hac701* compared to 9-WT that are

not present in H3K27ac, histone target-specificity of HAC701 still needs to be verified. As

H3K9ac and H3K14ac were seemed co-regulated and were directly correlated to each other

in regulating their common target genes (43), we assume that the H3K14ac in our *hac701*

would show the same correlation as well.

HAC701 appears to be an ancient protein in the CBP histone acetyltransferase family

clade (Fig. 1A). The fact that *HAC701* mutation is not lethal, suggests that other CBP

members and arguably members of other histone acetyltransferase families might have

redundant functions at some regulatory level. However, our results that *hac701* showed

enhanced disease resistance phenotype and reduction in acetylation demonstrate certain

functional specificities through HAC701-dependent pathways. Also, the fact that *hac701*

showed reduction in expressed genes at the systemic level (Fig. S19), indicates possible

broader roles of HAC701 other than localized modulation of rice innate immunity.

MATERIALS AND METHODS

Biological samples and plant growth conditions. *Oryza sativa* ssp. *japonica* cv.

Nipponbare (wild type) plants and mutant lines were grown in commercial soil (Kumiai, JA

Okinawa) at 30°C day/25°C night temperatures under a 12-h-light/12-h-dark photoperiod in

an incubator (BiOTRON, NK System). The lighting was supplied by white light at an

intensity of 31,000 lx. Relative humidity was at 70%. For embryonic rescue, rice embryos

were extracted from wild-type immature seeds 10-14 days after flowering. Embryos were

grown into plantlets in Murashige and Skoog (MS) agar medium before being transferred to

soil for further growth.

Isolation and screening of *HAC701* mutant lines. To generate *HAC701* CRISPR/Cas9

knockout mutant lines, two sgRNA *HAC701*-specific target sites were obtained from

CRISPR-P website and were used to synthesize primers for pRGEB31 (stable system) (59)

(Table S1). Briefly, the vectors were digested with BSA I, while primers were

phosphorylated and annealed to produce a DNA oligo duplex. The digested vectors were

ligated to DNA oligo duplex using T4 ligase. CRISPR/Cas9 vectors were introduced into

Agrobacterium EHA105 and rice calli were transformed using the standard *Agrobacterium-*

mediated transformation procedure. For CRISPR/Cas9 T0 and T1 screening, leaf samples

were collected and genomic DNA was extracted using a standard CTAB protocol. Then,

PCR-RFLP assay utilizing BseLI restriction enzyme (Thermo Fisher Scientific) was used to

detect the CRISPR/Cas9-engineered mutations on *HAC701* targets (Table S1). Positive lines

detected by PCR-RFLP assay were further analyzed by sequencing for INDEL mutations

using S2 and S5 forward primers (Table S1). Four DNA amplicons per line were cloned and

sequenced to determine the zygosity of the lines. Biallelic mutations were found in a few T0

and T1 lines. First exon (Target 1) was initially targeted in *HAC701* gene using a

CRISPR/Cas9 vector construct containing a single-guided RNA (S2) (Fig. S2A; Table S1).

The isolation of the first generation (T0) *CRISPR Cas9-HAC701-S2* lines yielded 90 positive

independent lines, and among these, randomly chosen representative lines were further

genotyped to characterize the identified DNA mutation. PCR and RFLP assays resulted in the

isolation of monoallelic lines characterized mostly by deletions and a few insertions

(INDELS) on or surrounding the targeted site of sgRNA (S2) (Fig. S2B-C; Table S1). The T1

HAC701-S2 generation as observed from seeds showed conservation of mutation directly

from parental lines (Fig. S2B). Then, mutant lines were isolated using the same technology

targeting the fifth exon (Target 2) of the *HAC701* gene using a construct that also contains a

single guide RNA (S5) (Fig. S2D; Table S1). The *CRISPR-Cas-HAC701-S5* (T0) lines

generated 11 positive independent lines of which three were genotyped for verification of

mutations (Fig. S2E). Second generation (T1) lines also showed conservation of mutations

(Fig. S2E). Similar to *HAC701-S2* lines, genotyping showed INDELS in proximity to or on

the target site.

Phenotyping. Positive lines containing biallelic homozygous mutations were phenotyped for

effective grain production by examining and counting the mature grains produced in each

panicle. Tiller number was counted one month after flowering. Chlorosis phenotype was used

as a proxy for any defect in photosynthesis and or apparatus. 1000-grain weight was

measured on seeds oven dried for 30-days at 37°C. Comparison of height and general

morphological structures were documented using photography.

flg22 leaf disc assay. flg22 peptide, a well-known inducer of plant innate immunity,

specifically of PTI was used to test which rice HATs respond to flg22 treatment. Leaf disc

assay (modified from 60, 61) was performed on fully expanded 30-d-old wild-type leaf

samples and treated with synthetic flg22 peptide (30-51 aa, Flic, *Pseudomonas aeruginosa*)

(ADI, Inc.) at different time periods and concentrations. Briefly, leaves were cut into about 5

mm sizes and floated on the water for 24-h in growth chamber to remove the symptoms of

wounding stress. Leaves were then treated with PAMP solution in water at 15 ml falcon tubes

with rotation (Corning Science). After treatment, paper towel-dried leaves were frozen in

liquid nitrogen.

Pathogenesis assay. Seeds were surface sterilized and imbibed in sterile water in the dark for

72 h before sowing on MS basal medium (Sigma Life Science). After 10 days, the plantlets

were transferred to soil and were grown for another 18 days until infection. *Pseudomonas*

syringae pv. *oryzae* (*Pso*) (MAFF No. 301530, NIAS Genebank) was grown in Luria-Bertani

(LB) broth (Sigma-Aldrich) at 28°C until OD = 0.2 and was resuspended in 10 mM $MgCl_2$.

The fourth leaf counting from the first true leaf was infected with *Pso* using needleless

syringe injected from the lower surface of the leaf. *Pso* leaf infiltration was performed 10 cm

from the tip of the leaf and was done 3x with approximately 1 cm space between the

infiltration sites. Infected plants were temporarily maintained outside the growth chamber for

2 h to allow drying of the infected sites before returning to the chamber. Infected leaf samples

were collected 3 days after *Pso* inoculation utilizing only the tissues comprising the spaces

between the infiltration sites. These tissues were grounded in sterile 10 mM $MgCl_2$ and the

grounded tissue suspensions were transferred to LB agar medium for incubation. The

infected local tissues were assayed with minor modifications as detailed in Liu, *et al*. (62).

For log cfu per leaf disc measurement, the grounded tissue suspensions were serially diluted

six times to be able to count with accuracy the *Pso* colonies on Luria-Bertani (LB) agar

medium. *Pso* colonies were counted from 4^{th} until 6^{th} serial dilution in several independent

and genotypically dissimilar *hac701* lines, *9-12b$^{-/-}$*, *9-5$^{-/}$*, and *9-12a$^{-/-}$*. The remaining samples

of the fourth leaf were used for RNA-sequencing analysis. For blast assay, the fifth leaf of

25-day-old plants were used for infection of *Magnaporthe oryzae* (MAFF No. 101511, NIAS

Genebank). Blast spores were incubated 10 cm from the tip of the leaf (1^{st} site) and at another

site of the same leaf 10 cm from the first site (2^{nd} site). Infected plants were incubated for 10

days inside an incubator (BiOTRON, NK System) and the length of blast infection was

measured.

Gene expression analysis. Total RNA extraction was performed using RNeasy Plant Mini

Kit (Qiagen) or Maxwell® 16 LEV Plant RNA Kit (Promega). cDNA was synthesized using

Primescript II 1^{st} Strand cDNA Synthesis Kit (Takara) according to manufacturer's

instructions. RT-qPCR assays were performed on three biologically independent samples or

as indicated. RT-qPCR was performed using SYBR Premix Ex Taq II (Tli RNaseH Plus)

(Takara) and was calculated following Pfaffl (63) by averaging the values relative to *ACT1*

control gene. Primer sequences are listed in Table S18.

RNA-sequencing. Total RNA was isolated from mock- and *Pseudomonas syringae* pv.

oryzae (*Pso*)-treated 9-WT wild type and *hac701* with Maxwell 16 LEV Plant RNA Kit

(Promega) run on the Maxwell 16 Instrument (Promega) and/or mirVana miRNA Isolation

Kit (Invitrogen by Thermo Fisher Scientific). The *hac701* lines, *9-12b$^{-/-}$* and *9-5$^{-/-}$*, were

lumped together as two independent biological samples during analysis. To remove the

contaminating genomic DNA from RNA samples isolated using the mirVana miRNA

Isolation Kit, RNA was treated with DNase I (RNA free) (Nippon Gene) following the

manufacturer's instructions. Samples were submitted to OIST Sequencing Center for RNA

quality checking, library preparation, and paired-end mRNA-sequencing (PE mRNA-seq).

Chromatin immunoprecipitation-sequencing.

ChIP analysis of histone modifications in wild type and mutants were performed as follows:

One-month-old mature leaves of wild type (9-WT) and *hac701* (*9-12b$^{-/-}$*) mutant were fixed

in a fixation buffer (10mM Tris-HCl (pH 7.5), 50 mM NaCl, 0.1 M sucrose, 1 %

formaldehyde) for 10 min, followed by quenching by 125 mM Glycine for 5min. Nuclei

isolation was performed as previously described (64). Immuno-precipitation was performed

for two replicates for each genotype (about 1 g tissue/IP) by SimpleChIP Plus Kit (Cell

Singaling Technology) according to the manufacturer's instructions. Anti-Histone H3

antibody (Abcam ab1791), Anti-acetyl Histone H3K9 antibody (Merck ABE18), Anti-

Dimethyl Histone H3K9 antibody (Merck 05-1249), Anti-Trimethyl Histone H3K9 antibody

(Merck 17-10242), Anti-acetyl Histone H3K27 antibody (Abcam ab4729) were used for IPs.

Dynabeads M-280 Sheep Anti-Rabbit IgG (Invitrogen 11203D) or Anti-Mouse IgG

(Invitrogen 11201D) was used for purification of chromatin-antibody complex. Precipitated

DNA samples were sequenced by Hiseq 4000 in 150 bp paired-end mode in OIST

Sequencing Center. For ChIP real-time PCR, we used the "Percent Input Method" (Thermo

Fisher Scientific) with 33% input value to normalize the data for enrichment calculation.

Three independent biological replicates were performed for each sample.

Data analysis. For RNA-sequencing analysis, high quality reads were trimmed in order to

remove sequencing bias and adapter effects. Trimmed reads were then mapped to the *Oryza*

sativa spp. *japonica* genome (Os-Nipponbare-Reference-IRGSP-1.0) using Tophat (65, 66).

Custom R scripts were used to generate the RNA count table necessary to analyze the

differentially expressed genes (DEGs). Differential expression analysis was performed using

DESeq2 package involving count normalization, dispersion estimation, and differential

expression test (67). DEGs were selected at Benjamini-Hochberg adjusted p value of < 0.01.

To perform Gene Ontology (GO) analysis, gene lists were submitted to Enrichment Analysis

(68) web tool of the Gene Ontology Consortium (69-71). To create the Venn diagram, gene

sets were submitted to InteractiVenn web tool using *unions by list* to obtain overlapping

components of input datasets (72). For ChIP-sequencing analysis, raw reads were trimmed

and were mapped to *Oryza sativa* spp. *japonica* genome (Os-Nipponbare-Reference-IRGSP-

1.0) using Bowtie (73). Read alignments were visualized using Integrated Genome Browser

(IGB) (74). ChIP-enriched peaks were called by deriving a quantitative reproducibility score

called "irreproducibility discovery rate" (IDR) (68). Comparative analyses and quantitation

of enrichments between 9-WT and *hac701* ($9\text{-}12b^{-/-}$) were analyzed using a suite of tools and

utilities found in deepTools (75) and BEDTools (76). Peaks were positioned to genomic

locations (intergenic, genes, transposable elements, simple repeats) and within genes

(promoter, exon, intron, TSS, TTS). Table S13 showed the number of peaks identified for

each histone modifications in 9-WT and *hac701* ($9\text{-}12b^{-/-}$). Peak-to-gene assignment was

performed using Homer (v4.11) (77). To create the phylogenetic tree of Arabidopsis and rice

histone acetyltransferases, we used the UniProt amino acid sequences and reconstructed the

phylogenetic relationships using Phylogeny.fr (78, 79). For KIX domain analysis, we used

KIXBASE to perform multiple sequence alignment (37). To identify *cis*-elements on DNA

sequences, we used cister prediction software that utilizes hidden Markov model (80).

Data visualization. Visualization of majority of the data was performed using Microsoft

Excel for Mac 2011 (version 14.7.1) and R packages: heatmaply, functions from *gplots*, and

ggplot2.

Data Repository. For RNA-sequencing and ChIP-sequencing, raw data will be deposited

shortly in the DDBJ Sequence Read Archive. Processed data will also be deposited in

 "Epigenomics Data site" maintained by Plant Epigenetics Unit

 (https://plantepigenetics.oist.jp/). All other data can be found in the Supplemental Tables 1-

 18 in this manuscript.

FIGURE LEGENDS

Figure 1 Disease phenotype of biallelic homozygous CRISPR/Cas9 lines targeting the fifth

exon of rice acetyltransferase gene, *HAC701*. (*A*) Phylogenetic relationships of members of

Arabidopsis and rice histone acetyltransferases. The numbers in red are branch support

values. All amino acid sequences are from UniProt database. (*B*) Upper panel: Alleles from

four lines of the second generation (T1) plants were identified by cloning and sequencing the

PCR products from *HAC701* target region using the primers found in Table S1. Lower panel:

PCR and RFLP assays of representative T1 generation lines. The mutant line *9-12a$^{-/-}$*

displaying a wild type-like band was due to unaltered sequence order recognized by BseLI

even after an insertion event, however we confirmed by sequencing that an insertion mutation

was present in this line. (*C*) *Pseudomonas syringae* pv. *oryzae* (*Pso*) infection assay of

hac701, *9-12b$^{-/-}$* and *9-5$^{-/-}$* (T2), using *9-5$^{+/+}$* segregated wild type line (9-WT) as control.

Bacterial density quantification uses values from 4th-6th serial dilutions. The mutant line *9-*

12b$^{-/-}$ was grown from the seeds of the third generation (T2), while *9-5$^{-/-}$* line was

embryonically rescued about 15 days post flowering from the second generation (T1)

parental plants. All mock measurements yielded zero bacterial growth for both wild type and

mutant lines. Bars represent the 95% confidence interval (CI) and compared to the wild type

using two-tailed Student's *t*-test at *P*<0.05 in *9-12b$^{-/-}$*; one-tailed Student's *t*-test at *P*<0.05 in

9-5$^{-/-}$. (*D*) *Magnaporthe oryzae* infection assay of *hac701*, *9-12b$^{-/-}$* and *9-5$^{-/-}$*, using *9-5$^{+/+}$*

segregated wild type line (9-WT) as control. Lesion length was measured on leaves after 10

dpi (days post infection). White length indicator bar on the left panel is 10 mm. Bars on the

right panel represent standard error (SE) and compared to the wild type using *F*-test for

variance and Student's *t*-test.

Figure 2 The rice *WRK45* gene expression is induced in *hac701* lines. (*A*) Overlap analysis

of upregulated and downregulated genes in *hac701* background under mock and pathogen

(*Pso*)-treatment conditions. *OsWRKY45* is the only differentially expressed gene in either the

absence or presence of pathogen. Transcriptome data were normalized using mock data sets

for each genotype and condition. (*B*) *OsWRKY45* expression when *HAC701* is mutated (+

sign). The addition of pathogen (+ sign) on *hac701* increased further the *OsWRKY45*

expression. Transcriptome data of *hac701*, *9-12b$^{-/-}$* and *9-5$^{-/-}$*, under mock or *Pso*-treated

conditions were lumped as two independent biological samples for analysis (See Table S2).

hpi, hours after infection.

Figure 3 *hac701* phenocopies *OsWRKY45* overexpression transgenic plants. (*A*) Overlap

analysis of differentially expressed genes (DEGs) in *hac701* background with

benzothiadiazole (BTH)-responsive genes (*i.e.* BTH is a salicylic acid analog) in wild type

rice. (*B*) Overlap analysis of DEGs in *hac701* background with BTH-inducible and

OsWRKY45-dependent or *OsNPR1*-dependent genes. (*C*) *OsWRKY45*-dependent defense

pathway components. (*D*) Gene expression of *OsWRKY45*-dependent defense genes in 9-WT

and *hac701*.

Figure 4 Genome-wide H3K9 acetylation is depleted in *hac701*. (*A*) Heatmap showing the

accumulation of H3K9 acetylation on 265 peaks by taking the enrichment ratio of 9-WT and

9-12b^{-/-} data (Wild type/mutant). (*B*) Heatmaps showing the accumulation of H3K27

acetylation, H3K9 di-methylation, and H3K9 tri-methylation on 265 peaks by taking the

same enrichment ratio as (*A*). Heatmaps in (*A*) and (*B*) are representatives of two biologically

independent data showing similar enrichment results and number of peaks identified. (*C*) GO

enrichment analysis showing only the "Biological Process" of the 263 genes found in the 265

peaks. Numbers in enrichment indicate Fold Enrichment Value using Fisher's Exact test with

results for uncorrected *P*<0.05. (*D*) Enrichment scores of seven representative genes showing

depletion of H3K9 acetylation in *hac701*, *9-12b^{-/-}*. (*E*) Enrichment scores of 10 representative

putative decoy/sensor genes (40) showing depletion of H3K9 acetylation in *hac701*, *9-12b^{-/-}*.

Scores in (*D*) and (*E*) are averages of two biologically independent samples with bars

showing standard deviation (SD).

Figure 5 *OsWRKY45* loci is enriched in H3K9 acetylation in *hac701*. (*A*) Enriched peaks on

histone modifications positioned to genomic locations (Intergenic, Promoter-TSS, Exon,

Intron, and TTS) in 9-WT and *hac701*, *9-12b^{-/-}*. All data were normalized to input. (*B*)

OsWRKY45 depth graph of RNA-seq and H3K9 acetylation ChIP-seq showing the number of

reads in RPKM (Reads Per Kilobase of transcript, per Million mapped reads) in 9-WT and

hac701. The red dotted box shows the 1kb upstream region of *OsWRKY45* with arrow

indicating the direction of transcription. (*C*) Upper panel: Gene map of *OsWRKY45* with

arrow showing the direction of transcription, gray bars are 5' and 3' untranslated regions

(UTRs), black bars are exons, and white bars are introns. Lower panel: 9-WT and *hac701* (*9-*

12b^{-/-}) mutant ChIP-qPCR showing enrichments of H3K9 acetylation and H3K9 di-

methylation in *OsWRKY45* locus. The significant difference in enrichment is computed using

two-tailed Student's *t*-test where asterisks: ***P<0.01, **P<0.05, and NS means Not

Significant.

TABLE LEGENDS

None

SUPPLEMENTARY FIGURE LEGENDS

Figure S1 Characterization of mutations generated by CRISPR/Cas9 editing on the first and fifth exon of rice acetyltransferase gene, *HAC701*. (*A*) Schematic showing an sgRNA targeted to the first exon of *HAC701* gene. (*B*) PCR and RFLP assays of representative T0 and T1 generation lines from leaf blade (Lb), panicle (Pa), and seed (S) DNA samples. +/+ and M represent the zygosity of the line, where +/+ refers to wild type and M refers to monoallelic. (*C*) Alleles from 12 T0 generation lines identified by cloning and sequencing the PCR products from *HAC701* target regions using the primers found in Table S1. Similar line number indicates that lines came from the same callus. For each line, four DNA amplicons were cloned and sequenced and the fraction indicates the number of times the type of mutations were found in each line. In case not indicated, it means wild type. The asterisk (*) indicates the most common mutation found within and across different lines. (*D*) Upper panel: Schematic showing an sgRNA targeted to the fifth exon of *HAC701*. Lower panel: Schematic showing the protein domains of HAC701. (*E*) PCR and RFLP assays of representative T0 and T1 generation lines from leaf blade DNA samples. +/+, M, B represent the zygosity of the line, where +/+ refers to wild type, M refers to monoallelic, and B refers to biallelic. T1 lines came from parental 2-3 line. (*F*) Phenotype of T2 lines showing effective grains and tiller number. (*G*) Images of T3 *hac701* lines compared to segregated wild type, 9-WT. (*H*) RT-qPCR levels of *HAC703* and *HAC704* genes in *HAC701* mutation line background. Samples were treated hydroponically with HDAC inhibitors (1μM Trichostatin A (TSA) and 100 μM Nicotinamide at final concentration) for three days. Bars represent

standard error (SE); n= 3. The significant difference in transcription is computed using two-

tailed Student's *t*-test where asterisks: *$P<0.01$. (*I*) Phenotype of T3 and T4 lines showing

grain weight.

Figure S2 flg22 induced the expression of rice pathogenesis-related and histone

acetyltransferase (HAT) gene, *HAC701*. Transcriptional levels of pathogenesis-related gene,

PR10a, and of eight HAT genes upon flg22 induction at concentrations in μM units. Data

shown are means ± SE; n= 3. The significant difference in transcription is computed using

two-tailed Student's *t*-test where asterisks: ***$P\leq0.01$, **$P\leq0.03$, * $P\leq0.05$.

Figure S3 *Pseudomonas syringae* pv. *oryzae* (*Pso*) infection assay in T2 *hac701* line, *9-12a$^{-/-}$*

. Values from 4th-6th serial dilutions were used for quantification. Segregated wild type line

was used as a control. Bars represent the 95% confidence interval (CI) and compared to wild

type using one-tailed Student's *t*-test at *$P<0.05$.

Figure S4 Sample distances of RNA-seq data after regularized-logarithm transformation

(rlog). (*A*) Heatmap showing the Euclidean sample distance matrix of mock- and

Pseudomonas syringae pv. *oryzae* (*Pso*)-treated samples in two biologically independent

replicates. (*B*) Principal component analysis (PCA) of the mock- and *Pso*-treated RNA-seq

samples. Principal component 1 (PC1) compares the variance between mock- versus *Pso*-

treated samples. Principal component 2 (PC2) compares the variance between biological

replicates within treatments.

Figure S5 Top 10 genes that are most highly variable in mock- and *Pso*-treated RNA-seq

samples. Gene descriptions were derived from Oryzabase: Integrated Rice Science Database

(NBRP) and Rice Genome Annotation Project (NSF). Two biologically independent RNA-

seq replicates are presented.

Figure S6 Rice-*Pseudomonas syringae* pv. *oryzae* (*Pso*) pathosystem. (*A*) Enriched pathways

in the rice-*Pso* pathosystem. (*B*) MA-plot of the rice-*Pso* pathosystem showing log fold

changes upon 72 hpi (hours post infection). *P* adjusted value is set at < 0.1. (*C*) Plot

dispersion estimates of genes in the rice-*Pso* pathosystem. The red line indicates the trend

showing the dispersion's dependence on the mean. The blue points are the final estimate of

each gene in reference to the red line. The blue circles are genes with high gene-wise

dispersion estimates and are considered outliers. Two biologically independent RNA-seq data

in mock and *Pso* treatments are presented where DEGs were selected at *P* adjusted value <

0.01.

Figure S7 Features of differentially expressed genes (DEGs) in *Pseudomonas syringae* pv.

oryzae (*Pso*)-treated 9-WT and *hac701* transcriptomes. Overlap and GO-analysis of

upregulated (*A*) and downregulated (*B*) genes in *hac701* under *Pso* treatment. Transcriptome

data were normalized using mock data sets for each genotype. GO enrichment analysis

showing only the "Biological Process" of upregulated *HAC701*-repressed genes (141) and

HAC701-dependent genes (336), downregulated *HAC701*-enhanced genes (31) and

HAC701-dependent genes (46). Fold Enrichment Value uses Fisher's Exact test with results

for uncorrected *P*<0.05.

Figure S8 Alignment of KIX domain found in different organisms. KIX domain in HAC701

(*A*) and HAC703 (*B*) proteins. Sc: *Saccharomyces cerevisiae*, Os: *Oryza sativa*, Hs: *Homo*

sapiens, Ce: *Caenorhabditis elegans*, Dm: *Drosophila melanogaster*, Mm: *Mus musculus*.

Figure S9 *OsWRKY45* promoter contains a rare CRE motif. (*A*) CRE motif cluster found in

1kb upstream sequences of *OsWRKY45* start site. The red line indicates the probability of

transcription factors binding to these CRE motif sites. CRE motif is found most probably on

the direct strand (+). (*B*) 5 genes out of 660 DEGs (0.76%) in the *hac701* contain the CRE

motif. Cister parameter settings: default settings were used.

Figure S10 Enrichment strength of ChIP signals in 9-WT input and histone modifications.

Figure S11 Heatmaps of genome-wide histone modification enrichments in genes and

transposable elements (TEs) of rice 9-WT and *hac701* (*9-12b$^{-/-}$*) mutant. Enrichment of H3K9

acetylation, H3K9 di-methylation, H3K9 tri-methylation, H3K27 acetylation, and H3 in

genes (*A*) and TE (*B*) regions. Data represent one of the two biologically independent

samples showing similar enrichment results.

Figure S12 Heatmaps of genome-wide histone modification enrichments in simple repeats of

rice 9-WT and *hac701* (*9-12b$^{-/-}$*) mutant. Enrichment of H3K9 acetylation, H3K9 di-

methylation, H3K9 tri-methylation, H3K27 acetylation, and H3 in simple repeat regions.

Data represent one of the two biologically independent samples showing similar enrichment

results.

Figure S13 Enrichment scores of additional representative genes showing differential enrichments of H3K9 acetylation in *hac701* (*9-12b$^{-/-}$*) mutant. (*A*) Transcriptional coactivator genes showing reduced or partially reduced H3K9 acetylation enrichments. (*B*) Putative guard genes showing differential enrichments of H3K9 acetylation. Scores are averages of two biologically independent samples with bars showing standard deviation (SD).

Figure S14 Distribution on genomic locations of 265 regions in H3K9 acetylation ChIP (wild type/mutant).

Figure S15 *OsWRKY45* depth graph of H3K9 and H3K27 acetylation ChIP-seq showing the number of reads in RPKM (Reads Per Kilobase of transcript, per Million mapped reads) in 9-WT and *hac701* (*9-12b$^{-/-}$*) mutant. The red dotted box shows the 1 kb upstream region of *OsWRKY45* with arrow indicating the direction of transcription. Data show a representative sample.

Figure S16 *OsWRKY45* depth graph of H3K9 acetylation and H3K9 di-methylation ChIP-seq showing the number of reads in RPKM (Reads Per Kilobase of transcript, per Million mapped reads) in 9-WT and *hac701* (*9-12b$^{-/-}$*) mutant. The red dotted box shows the 1 kb upstream region of *OsWRKY45* with arrow indicating the direction of transcription. Data show a representative sample.

Figure S17 *OsWRKY45* depth graph of H3K9 acetylation and H3K9 tri-methylation ChIP-seq showing the number of reads in RPKM (Reads Per Kilobase of transcript, per Million mapped reads) in 9-WT and *hac701* (*9-12b$^{-/-}$*) mutant. The red dotted box shows the 1 kb

upstream region of *OsWRKY45* with arrow indicating the direction of transcription. Data

show a representative sample.

Figure S18 Systemic gene expression analysis in rice-*Pseudomonas syringae* pv. *oryzae*

(*Pso*) pathosystem. (*A*) Differentially expressed genes (DEGs) in the local (870) and systemic

(24) tissues of plants infected with *Pso*. Eight DEGs were found to be common in both local

and systemic tissues. (*B*) The expression of eight DEGs shown in log2 fold changes. For

local tissue analysis, two independent RNA-seq samples in both treatments (mock and *Pso*)

are presented. For systemic tissue analysis, two independent RNA-seq samples in mock and

three in *Pso* treatment are presented. The number of reads are in RPKM (Reads Per Kilobase

of transcript, per Million mapped reads) in 9-WT and *hac701*.

Figure S19 MA-plot of differentially expressed genes (DEGs) in systemic tissues of 9-WT

and *hac701* infected with *Pseudomonas syringae* pv. *oryzae* (*Pso*). (*A*) and (*B*) plots showing

differences in measurements of differentially expressed genes (DEGs). *P* adjusted value is set

at <0.01. For 9-WT and *hac701* samples, two independent RNA-seq samples in mock

treatment and three in *Pso* treatment are presented.

Figure S20 *OsWRKY45*-dependent defense pathway is suppressed by *HAC701*. Defense-

related genes regulated directly and indirectly by *OsWRKY45* are negatively regulated by

HAC701 resulting in resistance phenotype of *hac701* through *OsWRKY45* upregulation.

SUPPLEMENTARY TABLE LEGENDS

Table S1 sgRNA oligonucleotides used for CRISPR/Cas9 mutagenesis and primers for

genotyping. Bold sequences in the guide RNAs are the Protospacer Adjacent Motif (PAM).

Table S2 Summary of total RNA-sequencing data in local tissues.

Table S3 *HAC701* repressed and enhanced genes.

Table S4 Wild type (9-WT) DEGs in *Pso*.

Table S5 *HAC701* mutation DEGs in *Pso* (*HAC701mut*-DEGs in *Pso*). The motifs were

identified 1kb upstream sequence of the start codon.

Table S6 *HAC701* mutation DEGs in Mock (*HAC701mut*-DEGs in Mock).

Table S7 BTH-responsive genes (24 hpi) in microarray experiments (34, 81) (Shimono et al.

2007, The Plant Cell & Sugano et al. 2010, Plant Molecular Biology). Overlap genes were

removed.

Table S8 BTH-inducible & WRKY45-dependent genes (33) (Nakayama et al. 2013, BMC

Plant Biology).

Table S9 Upregulated genes after DEX-induced *WRKY45* expression (33) (Nakayama et al.

2013, BMC Plant Biology).

Table S10 BTH-inducible genes affected by rice *NPR1* knockdown (81) (Sugano et al. 2010,

Plant Molecular Biology).

Table S11 *HAC701* mutation DEGs found in BTH-WRKY45 regulated genes that contain

W-box motifs. The motifs were identified 1 kb upstream sequence of the start codon.

Table S12 *HAC701* mutation DEGs found in BTH-WRKY45 and BTH-NPR1 regulated

genes that contain W-box motifs. The motifs were identified 1 kb upstream sequence of the

start codon.

Table S13 Rice ChIP-sequencing information of histone modifications.

Table S14 Identified peaks in H3K9ac and their corresponding genes in 9-WT/*9-12b*$^{-/-}$ (wild

type/mutant). Colored boxes indicate the repeated genes.

Table S15 Identified peaks in H3K9me3 and their corresponding genes in *9-12b*$^{-/-}$/Input

(mutant/Input). Colored boxes indicate repeated genes.

Table S16 Summary of total RNA-sequencing data in systemic tissues.

Table S17 List of R genes upregulated in *hac701* in *Pso*.

Table S18 Oligonucleotides used for RT-qPCRs (1) and ChIP-qPCRs (2).

ACKNOWLEDGEMENTS

We thank OIST SQC for RNA-seq and ChIP-seq analysis. This project benefited greatly from the comments of Drs. Yusuke Saijo, Hirofumi Nakagami, Justin Walley, and from the members of the Plant Epigenetics Unit (OIST) and Plant Immunity Research Group (RIKEN CSRS).

FUNDING

This work was supported in part by the Okinawa Institute of Science and Technology Graduate University (OIST). N.A.E. was also funded by Grant-in-Aid for JSPS Fellows (Grant No: 18F18085). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

AUTHOR CONTRIBUTIONS

NAE: Conceptualization, Data curation, Funding acquisition, Formal analysis, Investigation, Visualization, Writing-original draft, Writing-review & editing

LNT: Data curation, Formal analysis, Writing-review & editing

SM: Formal analysis, Investigation

YS: Formal analysis, Investigation

KS: Funding acquisition, Formal analysis, Supervision, Writing-review & editing

HS: Funding acquisition, Formal analysis, Investigation, Supervision, Writing-review & editing

COMPETING INTERESTS

The authors have declared that no competing interests exist.

REFERENCES

1.   Lee KK & Workman JL (2007) Histone acetyltransferase complexes: one size doesn't
fit all. *Nat. Rev. Mol. Cell Biol.* 8(4):284-295.

2.   Chan HM & La Thangue NB (2001) p300/CBP proteins: HATs for transcriptional
bridges and scaffolds. *J. Cell Sci.* 114(Pt 13):2363-2373.

3.   Kalkhoven E (2004) CBP and p300: HATs for different occasions. *Biochem.*
Pharmacol. 68(6):1145-1155.

4.   Janknecht R & Hunter T (1996) Transcription. A growing coactivator network.
Nature 383(6595):22-23.

5.   Goodrich JA & Tjian R (1994) TBP-TAF complexes: selectivity factors for
eukaryotic transcription. *Curr. Opin. Cell Biol.* 6(3):403-409.

6.   Pandey R, *et al.* (2002) Analysis of histone acetyltransferase and histone deacetylase
families of Arabidopsis thaliana suggests functional diversification of chromatin
modification among multicellular eukaryotes. *Nucleic Acids Res.* 30(23):5036-5055.

7.   Liu X, *et al.* (2012) Histone acetyltransferases in rice (Oryza sativa L.): phylogenetic
analysis, subcellular localization and expression. *BMC Plant Biol.* 12:145.

8.   Fang H, Liu X, Thorn G, Duan J, & Tian L (2014) Expression analysis of histone
acetyltransferases in rice under drought stress. *Biochem. Biophys. Res. Commun.*
443(2):400-405.

9.   Ma X, Lv S, Zhang C, & Yang C (2013) Histone deacetylases and their functions in
plants. *Plant Cell Rep.* 32(4):465-478.

10.   Song G & Walley JW (2016) Dynamic Protein Acetylation in Plant–Pathogen
Interactions. *Frontiers in plant science* 7.

11.   Ding B & Wang GL (2015) Chromatin versus pathogens: the function of epigenetics
in plant immunity. *Frontiers in plant science* 6:675.

12.   Zhu QH, Shan WX, Ayliffe MA, & Wang MB (2016) Epigenetic Mechanisms: An
Emerging Player in Plant-Microbe Interactions. *Molecular plant-microbe interactions*
: MPMI 29(3):187-196.

13.   Fu W, Wu K, & Duan J (2007) Sequence and expression analysis of histone
964   deacetylases in rice. *Biochem. Biophys. Res. Commun.* 356(4):843-850.

965   14.   Yuan LW & Giordano A (2002) Acetyltransferase machinery conserved in
966   p300/CBP-family proteins. *Oncogene* 21(14):2253-2260.

967   15.   Ogryzko VV, Schiltz RL, Russanova V, Howard BH, & Nakatani Y (1996) The
968   transcriptional coactivators p300 and CBP are histone acetyltransferases. *Cell*
969   87(5):953-959.

970   16.   Bannister AJ & Kouzarides T (1996) The CBP co-activator is a histone
971   acetyltransferase. *Nature* 384(6610):641-643.

972   17.   Henry RA, Kuo YM, & Andrews AJ (2013) Differences in specificity and selectivity
973   between CBP and p300 acetylation of histone H3 and H3/H4. *Biochemistry*
974   52(34):5746-5759.

975   18.   Jin Q*, et al.* (2011) Distinct roles of GCN5/PCAF-mediated H3K9ac and CBP/p300-
976   mediated H3K18/27ac in nuclear receptor transactivation. *EMBO J.* 30(2):249-262.

977   19.   Tie F*, et al.* (2009) CBP-mediated acetylation of histone H3 lysine 27 antagonizes
978   Drosophila Polycomb silencing. *Development* 136(18):3131-3141.

979   20.   Das C, Lucia MS, Hansen KC, & Tyler JK (2009) CBP/p300-mediated acetylation of
980   histone H3 on lysine 56. *Nature* 459(7243):113-117.

981   21.   Schiltz RL*, et al.* (1999) Overlapping but distinct patterns of histone acetylation by
982   the human coactivators p300 and PCAF within nucleosomal substrates. *J. Biol. Chem.*
983   274(3):1189-1192.

984   22.   Sakamoto S*, et al.* (2012) E2A and CBP/p300 act in synergy to promote chromatin
985   accessibility of the immunoglobulin kappa locus. *J. Immunol.* 188(11):5547-5560.

986   23.   Modak R*, et al.* (2013) Probing p300/CBP associated factor (PCAF)-dependent
987   pathways with a small molecule inhibitor. *ACS chemical biology* 8(6):1311-1323.

988   24.   Espinas NA, Saze H, & Saijo Y (2016) Epigenetic Control of Defense Signaling and
989   Priming in Plants. *Frontiers in plant science* 7.

990   25.   Kuwata H (1985) Pseudomonas syringiae pv. oryzae pv. nov., causal agent of
991   bacterial halo blight of rice. *Ann Phytopath. Soc. Japan* 51:212-218.

992   26.   Kang S, Kang K, Lee K, & Back K (2007) Characterization of rice tryptophan
993   decarboxylases and their direct involvement in serotonin biosynthesis in transgenic
994   rice. *Planta* 227(1):263-272.

995   27.   Ishihara A, *et al.* (2008) The tryptophan pathway is involved in the defense responses
996   of rice against pathogenic infection via serotonin production. *Plant J.* 54(3):481-495.

997   28.   Shimizu T, *et al.* (2012) Purification and identification of naringenin 7-O-
998   methyltransferase, a key enzyme in biosynthesis of flavonoid phytoalexin sakuranetin
999   in rice. *J. Biol. Chem.* 287(23):19315-19325.

29.   Ogawa S, *et al.* (2017) OsMYC2, an essential factor for JA-inductive sakuranetin
production in rice, interacts with MYC2-like proteins that enhance its transactivation
ability. *Scientific reports* 7:40175.

30.   Swaminathan S, Morrone D, Wang Q, Fulton DB, & Peters RJ (2009) CYP76M7 is
an ent-cassadiene C11alpha-hydroxylase defining a second multifunctional
diterpenoid biosynthetic gene cluster in rice. *Plant Cell* 21(10):3315-3325.

31.   Vanholme R, Demedts B, Morreel K, Ralph J, & Boerjan W (2010) Lignin
biosynthesis and structure. *Plant Physiol.* 153(3):895-905.

32.   Wang Q, *et al.* (2012) Characterization of CYP76M5-8 indicates metabolic plasticity
within a plant biosynthetic gene cluster. *J. Biol. Chem.* 287(9):6159-6168.

33.   Nakayama A, *et al.* (2013) Genome-wide identification of WRKY45-regulated genes
that mediate benzothiadiazole-induced defense responses in rice. *BMC Plant Biol.*
13(150).

34.   Shimono M, *et al.* (2007) Rice WRKY45 plays a crucial role in benzothiadiazole-
inducible blast resistance. *Plant Cell* 19(6):2064-2076.

35.   Cheng X, *et al.* (2019) Structural basis of dimerization and dual W-box DNA
recognition by rice WRKY domain. *Nucleic Acids Res.* 47(8):4308-4318.

36.   Parker D, *et al.* (1996) Phosphorylation of CREB at Ser-133 induces complex
formation with CREB-binding protein via a direct mechanism.pdf>. *Mol. Cell.*
Biol.:694-703.

37.   Yadav A, Thakur JK, & Yadav G (2017) KIXBASE: A comprehensive web resource
for identification and exploration of KIX domains. *Scientific reports* 7(1):14924.

38.   Dancy BM & Cole PA (2015) Protein Lysine Acetylation by p300/CBP. *Chemical*
reviews 115(6):2419-2452.

39.   Du Z, *et al.* (2013) Genome-wide analysis of histone modifications: H3K4me2,
H3K4me3, H3K9ac, and H3K27ac in Oryza sativa L. Japonica. *Molecular plant*
6(5):1463-1472.

40.　Sarris PF, Cevik V, Dagdas G, Jones JD, & Krasileva KV (2016) Comparative
analysis of plant immune receptor architectures uncovers host proteins likely targeted
by pathogens. *BMC Biol.* 14:8.

41.　Weidenbach D, *et al.* (2016) Polarized Defense Against Fungal Pathogens Is
Mediated by the Jacalin-Related Lectin Domain of Modular Poaceae-Specific
Proteins. *Molecular plant* 9(4):514-527.

42.　Singh P, *et al.* (2014) Environmental History Modulates Arabidopsis Pattern-
Triggered Immunity in a HISTONE ACETYLTRANSFERASE1-Dependent Manner.
Plant Cell.

43.　Kim S, *et al.* (2020) GCN5 modulates salicylic acid homeostasis by regulating
H3K14ac levels at the 5' and 3' ends of its target genes. *Nucleic Acids Res.*
48(11):5953-5966.

44.　De Vleesschauwer D, Gheysen G, & Hofte M (2013) Hormone defense networking in
rice: tales from a different world. *Trends Plant Sci.* 18(10):555-565.

45.　Vidhyasekaran P (2016) *Switching on plant innate immunity signaling systems*
(Springer International Publishing, Switzerland).

46.　De Vleesschauwer D, Xu J, & Hofte M (2014) Making sense of hormone-mediated
defense networking: from rice to Arabidopsis. *Frontiers in plant science* 5:611.

47.　Yang DL, Yang Y, & He Z (2013) Roles of plant hormones and their interplay in rice
immunity. *Molecular plant* 6(3):675-685.

48.　Yan S & Dong X (2014) Perception of the plant immune signal salicylic acid. *Curr.*
Opin. Plant Biol. 20:64-68.

49.　Silverman P, *et al.* (1995) Salicylic acid in rice (biosynthesis, conjugation, and
possible role). *Plant Physiol.* 108:633-639.

50.　Chen Z, Iyer S, Caplan A, Klessig DF, & Fan B (1997) Differential accumulation of
salicylic acid and salicylic acid-sensitive catalase in different rice tissues. *Plant*
Physiol. 114:193-201.

51.　Hiebert CW, *et al.* (2020) Stem rust resistance in wheat is suppressed by a subunit of
the mediator complex. *Nature communications* 11(1):1123.

52.　Brownell JE & Allis CD (1996) Special HATs for special occasions: linking histone
acetylation to chromatin assembly and gene activation. *Curr. Opin. Genet. Dev.*
6:176-184.

53.   Poss ZC, Ebmeier CC, & Taatjes DJ (2013) The Mediator complex and transcription
1060   regulation. *Crit. Rev. Biochem. Mol. Biol.* 48(6):575-608.

1061   54.   Canet JV, Dobon A, & Tornero P (2012) Non-recognition-of-BTH4, an Arabidopsis
1062   mediator subunit homolog, is necessary for development and response to salicylic
1063   acid. *Plant Cell* 24(10):4220-4235.

1064   55.   van der Hoorn RA & Kamoun S (2008) From Guard to Decoy: a new model for
1065   perception of plant pathogen effectors. *Plant Cell* 20(8):2009-2017.

1066   56.   Jackson JP*, et al.* (2004) Dimethylation of histone H3 lysine 9 is a critical mark for
1067   DNA methylation and gene silencing in Arabidopsis thaliana. *Chromosoma*
1068   112(6):308-315.

1069   57.   Karasov T, Chae E, Herman J, & Bergelson J (2017) Mechanisms to Mitigate the
1070   Tradeoff between Growth and Defense. *Plant Cell*.

1071   58.   Huot B, Yao J, Montgomery BL, & He SY (2014) Growth-defense tradeoffs in plants:
1072   a balancing act to optimize fitness. *Molecular plant* 7(8):1267-1287.

1073   59.   Xie K & Yang Y (2013) RNA-guided genome editing in plants using a CRISPR-Cas
1074   system. *Molecular plant* 6(6):1975-1983.

1075   60.   Heese A*, et al.* (2007) The receptor-like kinase SERK3/BAK1 is a central regulator of
1076   innate immunity in plants. *Proceedings of the National Academy of Sciences of the*
1077   *United States of America* 104(29):12217-12222.

1078   61.   Park CJ & Ronald PC (2012) Cleavage and nuclear localization of the rice XA21
1079   immune receptor. *Nature communications* 3:920.

1080   62.   Liu X*, et al.* (2015) Bacterial Leaf Infiltration Assay for Fine Characterization of
1081   Plant Defense Responses using the Arabidopsis thaliana-Pseudomonas syringae
1082   Pathosystem. *J Vis Exp* (104).

1083   63.   Pfaffl MW (2001) A new mathematical model for relative quantification in real-time
1084   RT-PCR. *Nucleic Acids Res.* 29(9):e45.

1085   64.   Saleh A, Alvarez-Venegas R, & Avramova Z (2008) An efficient chromatin
1086   immunoprecipitation (ChIP) protocol for studying histone modifications in
1087   Arabidopsis plants. *Nature protocols* 3(6):1018-1025.

1088   65.   Trapnell C, Pachter L, & Salzberg SL (2009) TopHat: discovering splice junctions
1089   with RNA-Seq. *Bioinformatics* 25(9):1105-1111.

1090   66.   Trapnell C*, et al.* (2012) Differential gene and transcript expression analysis of RNA-
1091   seq experiments with TopHat and Cufflinks. *Nature protocols* 7(3):562-578.

67.   Love MI, Huber W, & Anders S (2014) Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome biology* 15(12):550.

68.   Li Q, Brown JB, Huang H, & Bickel PJ (2011) Measuring reproducibility of high-
throughput experiments. *The Annals of Applied Statistics* 5(3):1752-1779.

69.   Consortium TGO (2017) Expansion of the Gene Ontology knowledgebase and
resources. *Nucleic Acids Res.* 45(D1):D331-D338.

70.   Consortium TGO (2000) Gene Ontology: tool for the unification of biology. *Nat.*
Genet. 25:25-29.

71.   Consortium TGO (2015) Gene Ontology Consortium: going forward. *Nucleic Acids*
Res. 43(Database issue):D1049-1056.

72.   Heberle H, Meirelles GV, da Silva FR, Telles GP, & Minghim R (2015)
InteractiVenn: a web-based tool for the analysis of sets through Venn diagrams. *BMC*
Bioinformatics 16:169.

73.   Langmead B, Trapnell C, Pop M, & Salzberg SL (2009) Ultrafast and memory-
efficient alignment of short DNA sequences to the human genome. *Genome biology*
10(3):R25.

74.   Freese NH, Norris DC, & Loraine AE (2016) Integrated genome browser: visual
analytics platform for genomics. *Bioinformatics* 32(14):2089-2095.

75.   Ramirez F, *et al.* (2016) deepTools2: a next generation web server for deep-
sequencing data analysis. *Nucleic Acids Res.* 44(W1):W160-165.

76.   Quinlan AR & Hall IM (2010) BEDTools: a flexible suite of utilities for comparing
genomic features. *Bioinformatics* 26(6):841-842.

77.   Heinz S, *et al.* (2010) Simple combinations of lineage-determining transcription
factors prime cis-regulatory elements required for macrophage and B cell identities.
Mol. Cell 38(4):576-589.

78.   Dereeper A, *et al.* (2008) Phylogeny.fr: robust phylogenetic analysis for the non-
specialist. *Nucleic Acids Res.* 36(Web Server issue):W465-469.

79.   Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and
high throughput. *Nucleic Acids Res.* 32(5):1792-1797.

80.   Frith M, Hansen U, & Weng Z (2001) Detection of cis-element clusters in higher
eukaryotic DNA. *Bioinformatics* 17:878-889.

81.   Sugano S, *et al.* (2010) Role of OsNPR1 in rice defense program as revealed by
1124   genome-wide expression analysis. *Plant Mol. Biol.* 74(6):549-562.

1125

1126

1127

1128

1129

1130

1131

1132

1133

1134

1135

1136

1137

1138

1139

1140

1141

1142

1143

1144

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure S1

Figure S2

Figure S3

Figure S4

Figure S5

Gene ID	Description
OS12G0491800	Kaurene synthase-like 10
OS12G0258700	Laccase 29
OS07G0218700	Cytochrome P450 family protein
OS08G0140300	Tryptophan decarboxylase 1
OS07G0218200	Terpene synthase 3
OS12G0240900	Naringenin 7-O-methyltransferase
OS10G0528300	Tau glutathione S-transferase 4
OS08G0508000	Cytochrome P450 76M2
OS04G0659300	Root meander curling
OS11G0641500	Laccase 23

Figure S6

A

Highly enriched terms upon *Pseudomonas syringiae pv. oryzae* infection in 9-WT rice		
Up-regulated terms	Fold Enrichment	raw P-value
Cell wall macromolecule catabolic process	54.94	4.22E-07
Adventitious root development	54.94	3.54E-02
Lignin biosynthetic process	54.94	3.54E-02
Sphingolipid biosynthetic process	54.94	3.54E-02
Gibberellin biosynthetic process	54.94	3.54E-02
Chitin catabolic process	54.94	4.22E-07
Response to wounding	54.94	3.54E-02
Response to other organism	54.94	2.70E-08
ent-kaurene oxidation to kaurenoic acid	54.94	3.54E-02
Diterpene phytoalexin precursor biosynthetic process pathway	41.21	1.89E-04
Shikimate biosynthetic process	36.63	3.08E-03
Chorismate biosynthetic process	36.63	3.08E-03
Carbohydrate transport	18.31	8.32E-03
Carbohydrate metabolic process	5.78	6.62E-03
Electron transport chain	5.49	1.17E-03
Regulation of transcription, DNA templated	2.79	2.43E-02

Down-regulated terms	Fold Enrichment	raw P-value
Regulation of transcription, DNA templated	10.26	1.69E-02

B

C

Figure S7

Figure S8

GAL11_Sc	....VQDKD......TLSNA........ERAKNVNGLLQVLMDINTLNGGSSDTADKIRIHAKNFEAALFAKSSSKKEYMDSMNEKVAVMR	73
HAC701_Os 37-106	QLDNMDQDTS...VVRNIIHRKIVEYLNERKEFCNFDLSFLMEI.......GKCIDRHLFEKADSKIKYMDIETLRTRLN.	70
ARC105_Hs	.GAMGQETDLRSTA.FRQKLVSQIED.........AMRKAGVA.......HSKSSKDMESHVFLKAKTRDEYLSLVARLIIHFR.	66
CBP1_Ce	.....DCTKEHHQVTKDLRNHLVGKLVK.....AIFPEPNQEAMNENRLKDLIAYARKVEKEMFESANDREEYYHLLAEKIYKIQ	76
CBP_Dm	.....DNEKDWRESVTADLRNHLVHKLVQ.....AIFPTSDPTTMQDKRMHNLVSYAEKVEKDMYBMAKSRSEYYHLLAEKIYKIQ	76
CBP_Hs/Mm	.....GVRKGCHEHVTQDLRSHLVHKLVQ.....AIFPTPDPAALKDRRMENLVAYAKKVEGDMYBSANSREYYHLLAEKIYKIQ	76
EP300 Hs/Mm	.....GIRKQLHEDITQDLRNHLVHKLVQ.....AIFPTPDPAALKDRRMENLVAYARKVEGDMYBSANNRAEYYHLLAEKIYKIQ	76
consensus		

GAL11_Sc	NTYNTRKNAVTA	85
HAC701_Os 37-106		70
ARC105_Hs	DIHNKKSQASV.	77
CBP1_Ce	KELQEKKNSRL.	87
CBP_Dm	KELEEKRLKR..	86
CBP_Hs/Mm	KELEEKRRSRL.	87
EP300 Hs/Mm	KELEKRRTRL..	86
consensus		

 non conserved

X ≥50% conserved

B

HAC703_Os 52-126	........LLMR...NTMRDRIFEYIGRKQSSTDWRRRLPELAMRLBEILYRKFLNKADYLNMMRGPVEPQLQFAIKTLSAQNQQN.	75
CBP1_Ce	...DCTKEMHHQVTKDLRNHLVGKLVKAIFPEPNQEAMNDNRLKDLIAYARKVEKEMFESANDREEYYHLLAEKIYKIQ....KELQEKKNSRL.	87
CBP_Dm	...DNEKDWRESVTADLRNHLVHKLVQAIFPTSDPTTMQDKRMHNLVSYAEKVEKDMYEMAKSRSEYYHLLAEKIYKIQ....KELEEKRLKR..	86
CBP_Hs/Mm	...GVRKGLHEHVTQDLRSHLVHKLVQAIFPTPDPAALKDRRMENLVAYAKKVEGDMYBSANSREYYHLLAEKIYKIQ....KELEEKRRSRL.	87
EP300 Hs/Mm	...GIRKQLHEDITQDLRNHLVHKLVQAIFPTPDPAALKDRRMENLVAYARKVEGDMYEBSANNRAEYYHLLAEKIYKIQ....KELEKRRTRL..	86
GAL11_Sc	...VQDKDTLSN..AERAKNVNGLLQVLMDINTLNGGSSDTADKIRIHAKNFEAALFAKSSSKKEYMDSMNEKVAVMR...NTYNTRKNAVTA	85
ARC105_Hs	GAMGQETDLRST..AFRQKLVSQIEDAMR...KAGV.....AHSKSSKDMESHVFLKAKTRDEYLSLVARLIIHFR....DIHNKKSQASV.	77
consensus		

X non conserved

X ≥50% conserved

Figure S9

Possible functional cis-elements

Type	Position	Strand	Sequence	Probability
CRE	303-314	+	cgtgacgtcatc	0.62
CRE	289-300	+	gctgacgtcgcc	0.56
CRE	317-328	-	gttgtcgtcccc	0.14
CRE	289-300	-	gctgacgtcgcc	0.12

Figure S10

Figure S11

Genes

Transposable elements

Figure S12

Simple Repeats

Figure S13

Figure S14

Figure S15

H3K9ac

H3K27ac

9-WT #1

9-12b-/- #1

9-WT #1

9-12b-/- #1

1kb

OsWRKY45

Figure S16

Figure S17

Figure S18

Figure S19

DEGs in *Pso*/Mock in wild type   DEGs in *Pso*/Mock in *HAC701-/-* mutant

Systemic tissues

Figure S20