Divergent DNA methylation signatures of X chromosome regulation in marsupials and eutherians

Devika Singh[1], Dan Sun[1], Andrew G. King[2], David E. Alquezar-Planas[2], Rebecca N. Johnson[2,3], David Alvarez-Ponce[4],*, and Soojin V. Yi[1],*

[1]School of Biological Sciences, Georgia Institute of Technology, Atlanta, Georgia, USA;
[2]Australian Museum Research Institute, Australian Museum, Sydney, New South Wales, Australia;
[3]National Museum of Natural History, Smithsonian Institution, Washington, DC, USA;
[4]Department of Biology, University of Nevada Reno, Reno, Nevada, USA.

*Corresponding Authors:
David Alvarez-Ponce
dap@unr.edu
775-682-5735

Soojin V. Yi
soojin.yi@biology.gatech.edu
404-385-6084

Abstract

X chromosome inactivation (XCI) mediated by differential DNA methylation between sexes is well characterized in eutherian mammals. Although XCI is shared between eutherians and marsupials, the role of DNA methylation in marsupial XCI remains contested. Here we examine genome-wide signatures of DNA methylation from methylation maps across fives tissues from a male and female koala (*Phascolarctos cinereus*) and present the first whole genome, multi-tissue marsupial "methylome atlas." Using these novel data, we elucidate divergent versus common features of marsupial and eutherian DNA methylation. First, tissue-specific differential DNA methylation in marsupials primarily occurs in gene bodies. Second, females show significant global reduction (hypomethylation) of X chromosome DNA methylation compared to males. We show that this pattern is also observed in eutherians. Third, on average, promoter DNA methylation shows little difference between male and female koala X chromosomes, a pattern distinct from that of eutherians. Fourth, the sex-specific DNA methylation landscape upstream of *Rsx*, the primary *lnc*RNA associated with marsupial XCI, is consistent with the epigenetic regulation of female- (and presumably inactive X chromosome-) specific expression. Finally, we utilize the prominent female X chromosome hypomethylation and classify 98 previously unplaced scaffolds as X-linked, contributing an additional 14.6 Mb (21.5 %) to genomic data annotated as the koala X chromosome. Our work demonstrates evolutionarily divergent pathways leading to functionally conserved patterns of XCI in two deep branches of mammals.

Keywords: X chromosome inactivation, marsupial, DNA methylation, RSX, koala, whole genome bisulfite sequencing (WGBS)

Introduction

X chromosome inactivation (XCI) is an iconic example of sex chromosome regulation in which one of the two X chromosomes in females is silenced, a mechanism thought to adjust the expression levels of X-linked genes (1). Although XCI occurs in both eutherian and marsupial mammals (2), there are several notable differences in XCI between the two lineages. First, in eutherians, the transcription of *Xist* RNA from the inactive X chromosome is essential for XCI (3-5). However, the *Xist* locus is not present in marsupials (6, 7). Instead, another *lnc*RNA gene, *Rsx*, drives marsupial XCI (8). Second, marsupials exhibit 'imprinted' XCI by selectively silencing the paternal X chromosome (9, 10). In contrast, XCI in eutherians occurs randomly between the maternally and paternally derived X chromosomes, although paternal XCI has been observed during early rodent development (11, 12). Third, while eutherian XCI involves the exclusion of active histone marks and the recruitment of repressive histone marks on the inactive X chromosome (13), marsupial X chromosomes do not show a consistent pattern (10, 14). These differences suggest that evolutionary pathways leading to XCI likely differ between eutherians and marsupials, and that novel insights into the mechanism of XCI can be gained from comparative studies.

The role of DNA methylation on marsupial XCI has been particularly controversial. Immunofluorescent labeling studies observed relative hypomethylation of the inactivate X chromosome in marsupials (15, 16). Other studies found little difference in DNA methylation between active and inactive marsupial X chromosomes (10, 17, 18). Recently, Waters et al. (19) analyzed reduced representation bisulfite sequencing (RRBS) data of a male and female opossum (*Monodelphis domestica*) and proposed that female X chromosomes in marsupials, but not in eutherians, exhibit hypomethylation near the transcription start sites (TSSs). Notably, these previous DNA methylation studies either examined a small number of CpGs or employed methodologies that over-represent promoters and CpG islands (in case of RRBS, (20)). Since patterns of DNA methylation vary greatly among distinctive genomic regions with different functional consequences, it is necessary to extend our knowledge to unbiased, whole-genome assays of DNA methylation.

We used the whole-genome bisulfite sequencing, which captures the DNA methylation state of every nucleotide in the genome, of the modern koala (*Phascolarctos cinereus*) to understand the role of DNA methylation on XCI in marsupials and to resolve previous

controversies. We compared DNA methylation maps of five diverse tissues from a male and a

female koala, providing the first multi-tissue, whole genome methylome resource of any marsupial

with information on tissue-specific variation of DNA methylation. Utilizing these novel resources,

we show distinctive impacts of DNA methylation on tissue-specific gene expression in marsupials,

as well as on XCI in eutherians and marsupials. We further classify previously undetected X-linked

regions from this key marsupial species using characteristic features of X chromosome DNA

methylation. Our findings provide new insights into the evolutionary pathways leading to

functionally convergent yet mechanistically divergent pathways of XCI and regulation of gene

expression in two deep branches of mammals.

Results

Genome-wide patterns of DNA methylation in the modern koala

We generated whole genome bisulfite sequencing (WGBS) data from five tissues (brain, lung,

kidney, skeletal muscle, and pancreas) from a male ("Ben," Australian Museum registration

M.45022) and female koala ("Pacific Chocolate," Australian Museum registration M.47723). The

mean depth of coverage fell between 9.9× and 14.6× (Supplementary Table 1). Autosome-linked

scaffolds displayed GC content ranging between 38.58 and 39.59%. Scaffolds linked to

chromosome X had higher GC contents compared to autosomal scaffolds (Supplementary Table

2), similar to (21). We performed a hierarchical clustering analysis and observed a clear clustering

of samples by tissue (Fig. 1A) with the pancreas samples exhibiting the most unique methylation

signature while the kidney and lung samples share the most similar methylation profiles.

Differential DNA methylation between tissues

We identified shared and tissue-specific differentially methylated regions (DMRs) using BSmooth

(22). Tissue-specific DMRs were defined as regions that were differentially methylated in a

particular tissue compared to all other tissues in a pairwise analysis, while shared DMRs were

those found in multiple tissues (Fig. 1B). Consistent with the results of the clustering analysis, the

pancreas samples contained the most tissue-specific DMRs followed by the skeletal muscle, brain,

kidney and lung (Fig. 1B). The majority (50-53%) of tissue-specific DMRs fell in gene bodies

(Fig. 1C, Supplementary Fig. 1, and Supplementary Table 3), a significant excess compared to

length and GC matched control regions (fold enrichment (FE) = 1.25~1.44, $p < 0.0001$ based on

10,000 bootstraps; Fig. 1C, Supplementary Fig. 1 and Supplementary Table 3). On the other hand,

DMRs were significantly depleted in intergenic regions compared to the control regions ($p < 0.05$

based on 10,000 bootstraps; Fig. 1C, Supplementary Fig. 1, and Supplementary Table 3). Genes

containing tissue-specific DMRs were enriched in specific biological functions (Table 1). For

example, brain specific DMRs were linked to genes associated with neural developmental

processes such as neurogenesis and central nervous system development.

We integrated our methylome data with a previously generated RNA-seq koala

transcriptomes (23). Of the 12 tissues surveyed in that study, three tissues (kidney, brain, and lung)

overlapped with the tissues examined here. Promoter DNA methylation and gene expression were

significantly negatively correlated across the entire genome (Table 2, Supplementary Fig. 2). In

contrast, the relationship between gene body DNA methylation and expression was complex; both

extremely hypomethylated and hypermethylated gene bodies showed high gene expression (Table

2, Supplementary Fig. 2). We considered differentially methylated genes (DMGs) containing

DMRs between the brain and kidney samples (n = 1944 genes from n = 4,615 DMRs) compared

with differentially expressed genes (DEGs) between brain and kidney RNA-seq samples.

Currently, available RNA-seq data from koalas do not include sufficient biological replicates. We

overcame this limitation by simulating replicates within each RNA-seq data set (NOISeq, (24))

and identified 600 putative DEGs (probability of differential expression > 95% according to the

NOISeq). We found that DMGs were significantly more likely to be differentially expressed than

genes that did not contain DMRs, exhibiting a 1.54-fold enrichment ($\chi^2 = 33.07$, $p < 0.0001$). In

addition, differential expression between tissues was significantly correlated with differential

promoter DNA methylation between tissues (Supplementary Fig. 3A). Both relative

hypomethylation and relative hypermethylation of gene bodies were associated with increased

expression (Supplementary Fig. 3B). In Fig. 1D and 1E, we show DNA methylation patterns of

two representative genes containing brain specific DMRs.

Global hypomethylation of female X chromosome in koalas

The koala genome project used cross-species chromosome painting data to identify 24 putative X

chromosome scaffolds and 406 putative autosomal scaffolds (25). As expected from 2:1 ratio of

X chromosomes in females compared to males, the median depth of coverage of CpGs on the

putative X scaffolds were consistently higher (approximately 2-fold) in female samples compared

to male samples ($p < 2.2 \times 10^{-16}$, Mann-Whitney U test, Supplementary Fig. 4A). Additionally, the

proportion of reads mapped to the putative X scaffolds showed a distinct bimodal distribution

whereby the male samples cluster close to 1.3% and the female samples cluster near 2.4%

(Supplementary Fig. 4B). In contrast, male and female samples were indistinguishable with respect

to read mapping to putative autosomes (Supplementary Fig. 4D).

The global DNA methylation level of the female X chromosome was lower than that of the

male X chromosome in all koala tissues examined (Fig. 2A, B and Supplementary Fig. 5, $p < 2.2$

$\times 10^{-16}$, Mann-Whitney U test). A comparison to autosomal DNA methylation indicated that the X

chromosome exhibits reduced DNA methylation in females. Consequently, we use the term

'female hypomethylation' (as opposed to male hypermethylation) (Fig. 2C) consistently in this

work. As a comparative basis of the DNA methylation patterns observed in eutherian mammals,

we performed parallel analyses on human DNA methylation data (methods). Human X

chromosomes were also globally hypomethylated in females compared to males (Fig. 2A, C).

Female hypomethylation was observed in all functional regions across the X chromosome (Fig.

2D), but most pronounced in gene bodies and intergenic regions. Promoters showed the least

difference between males and females. As expected, the autosomal scaffolds did not display a

significant variation between female and male methylation levels in any functional region (Fig.

2E). Fig. 2F depicts the DNA methylation difference between male and female X chromosomes

in humans. In contrast to the pattern observed in koalas, promoters on the human female X

chromosomes were hypermethylated compared to those on the male X chromosome, congruent

with previous studies (26-28).

Promoter DNA methylation is not a universal driver of sex-specific expression in koalas

To investigate the implications of the observed sex-specific DNA methylation, we once again

utilized the published RNA-seq koala transcriptomes (23). Of the total RNA-seq dataset, only one

tissue (kidney) had expression data from both a male and female koala that could be directly

compared to our methylome dataset. Consequently, the other tissues were not considered for

further analysis. We calculated the log-transformed fold change of female and male gene

expression values using NOISeq, capitalizing on its ability to simulate technical replicates. Of the

209 X-linked genes, 36 (17.2%) exhibit female overexpression while 11 (5.3%) show male

overexpression (probability of differential expression > 95% based on NOISeq, Supplementary

Fig. 6A). Although, on average, autosomal genes also exhibited slight female-bias of expression
(Supplementary Fig. 6B, C), the increase was more substantial in the X chromosome (mean
chromosome X female to male $\log_2$ fold change = 0.50, autosome female to male expression $\log_2$
fold change = 0.24). We calculated the female and male fractional methylation difference in X
chromosome-linked promoters and gene bodies correlated with gene expression difference (N =
209 gene bodies and N = 206 promoters, excluding 3 promoters with CpGs coverage < 3). In
promoters, no significant relationship was observed (Supplementary Fig. 7A). In fact, the
proportions of significantly female-over expressed genes were similar between female-
hypermethylated promoters and female-hypomethylated promoters (Supplementary Fig. 7B).
Interestingly, female and male DNA methylation difference in gene bodies showed an overall
negative correlation with gene expression (Spearman's rank correlation coefficient, ρ = -0.14, p =
0.04, Supplementary Fig. 7C). A deeper analysis of the relative DNA methylation and expression
levels revealed that both extreme hypomethylation and hypermethylation are associated with
increased expression (Supplementary Fig. 7C) attesting to the complexity and heterogeneity of the
relationship between gene body DNA methylation and gene expression.
174
The *Rsx* region displays a pattern suggesting methylation driven control of X chromosome
regulation in koalas
Sex-specific DNA methylation of a key regulator of XCI in other marsupials, the *lnc*RNA gene
Rsx, has been associated with its differential expression (8, 10). Based on the sequence homology
with the *Rsx* gene from the gray short-tailed opossum (*Monodelphis domestica*) (8), we identified
a 29.8 Kb candidate *Rsx* sequence (Methods), using PacBio long read sequencing by Johnson et
al. (25). Interestingly, we observed a female hypomethylated region containing two CpG islands
upstream of the candidate *Rsx* covering 101 CpGs exhibiting a 36% reduction in mean female
DNA methylation compared to mean male DNA methylation (mean sex difference: -0.36 ± 0.14,
Fig. 3). The mean DNA methylation difference between sexes within this region was in the top
23% of the distribution of differences across all X-linked scaffolds of the combined tissues. The
expression of the *lnc*RNA annotated within the candidate *Rsx* gene region was significantly greater
in females (mean normalized read count = 6987.1) than in males (mean normalized read count =
16, p < 0.05 from DeSeq2 using the Wald test). When considering a subset of tissue expression

data with matched male and female samples (spleen, kidney, and lung), the results remained robust

across different tools to measure differential gene expression (Table 3).

Identification of novel candidate X-linked scaffolds by sex-specific methylation patterns

We have demonstrated above that X-linked scaffolds exhibit several distinctive WGBS patterns in

koalas, including different depths of coverage per CpG, distinctive clustering based on the

proportion of mapped reads, and unique methylation distributions between females and males

(Supplementary Fig. 4). We thus quantified DNA methylation differences between females and

males to determine if additional candidate X scaffolds existed within the 6.7% of the koala

assembly that remained unclassified. We identified an additional 98 scaffolds that showed a clear

shift towards female hypomethylation (mean female-male 5mC for all candidate X scaffolds was

-0.25 ± 0.12) (Supplementary Tables 4,5). These candidate scaffolds contributed 14.6 Mb (21.5%)

to the total genomic region annotated as the koala X chromosome. All candidate scaffolds followed

the expected bimodal distribution seen in the putative X scaffolds; male samples clustered around

0.3% and the female samples clustered around 0.5% (Supplementary Fig. 4C). This clustering

pattern could be attributed to the 2:1 ratio of X chromosomes in females compared to males that

was not observed in putative autosomes.

207   **Discussion**

In this study, we analyzed novel nucleotide-resolution genomic DNA methylation maps of five

tissues of a male and a female koala. The overall DNA methylation levels of koala tissues are on

par with those in other mammals (27, 29, 30), and show heavy genome-wide DNA methylation

punctuated by hypomethylation of CpG islands (e.g., Fig. 1D,E). Tissue-specific differential DNA

methylated regions (DMRs) were significantly enriched in gene bodies. Gene body methylation is

an ancestral form of DNA methylation in animal genomes (e.g., (31, 32)), yet its role in regulation

of gene expression is less well understood than promoter DNA methylation owing in part to the

heterogeneity of gene body regions with respect to sequence length, composition and functionality

of introns and exons, and CpG density compared to promoter regions (31, 33). For example, DNA

methylation levels of the first exons/introns of genes are negatively correlated with gene

expression (34-36), and tend to be different from downstream genic regions (35). On the other

hand, high levels of cumulative gene body DNA methylation is positively correlated with gene

expression and may reduce spurious transcription of intragenic RNA (37, 38). These studies
illustrate the potential for both positive and negative regulatory control of gene expression by gene
body DNA methylation.

We show that female X chromosomes are globally hypomethylated compared to both the
male X chromosome and the autosomes of both sexes (Fig. 2). Even though it may appear
counterintuitive at the first glance, the hypomethylation of female X chromosome appears as a
common feature of eutherian and marsupial mammals. Hellmann and Chess (26) showed that the
inactive X chromosomes of humans had reduced gene body DNA methylation. Recent whole
genome bisulfite sequencing analyses of mouse (27) and humans (28) showed that the
hypomethylation of inactive X chromosome is pervasive in gene bodies and intergenic regions
(Fig. 4). Waters et al. (19) performed reduced representation bisulfite sequencing (RRBS) of
several species including mouse and opossum, and concluded that the reduction of methylation in
gene bodies was specific to marsupials, but not in eutherians (19). The reason why they did not
observe methylation difference in mouse might be due to the inherent bias of RRBS, which
disproportionately samples regions with high GC content (20). High GC-content regions tend to
be hypomethylated (39, 40) and show less variation of DNA methylation. For example, the degree
of methylation difference between sexes in koala promoters is small compared to other genomic
regions, at least partly due to high GC content of promoters (Supplementary Fig. 8). How
chromosome-wide DNA hypomethylation is linked to chromosome-wide gene silencing is
currently unknown. Interestingly, marsupial genomes harbor an additional copy of *DNMT1* (41),
which could lead to functional divergence between the mammalian lineages. Analyses of DMNT
expression however did not indicate significant differential expression of *DNMT*s between sexes
(probability of differential expression using NOISeq < 95%). Additional analysis with broad and
balanced sampling of both sexes is necessary to rigorously test the impact of the marsupial *DNMT1*
duplication on gene expression.

Interestingly, DNA methylation signatures of *Rsx,* the major player in XCI initiation in
marsupials (8), suggest that koala *Rsx* expression is regulated by promoter CpG island DNA
methylation (Fig. 3). Wang et al. (10) also showed differential DNA methylation of *Rsx* promoter
in opossum. Therefore, regulation of the key initiator of XCI via DNA methylation is another
parallel feature between eutherians and marsupials Notably, a recent study (42) found that *Xist* and

Rsx harbor non-linear sequence similarity. Consequently, their shared functionality may be
partially due to characteristics of tandem repeat regions.

In summary, we show that gene body DNA methylation is an important contributor to
differential expression between tissues in koala. We also show that the global hypomethylation of
female X chromosome is a conserved feature of X chromosome regulation in eutherian and
marsupial mammals. However, X chromosome promoter methylation and the subsequent effect
on the regulation of gene expression appear to be divergent between these two lineages.
Regulation of the *Rsx*, on the other hand, is supported by DNA methylation, which mirrors the
regulation of the eutherian *Xist* locus. Together, these conclusions illuminate the intricate
evolutionary pathways that have diverged and converged to influence X chromosome regulation,
XCI, and dosage compensation in eutherian and marsupial mammals.

Methods

Whole genome bisulfite sequencing and processing

All genomic DNA was extracted using a Bioline Isolate II Genomic DNA Extraction Kit (Cat#.
BIO-52067) following the recommended protocol with an additional DNAse free RNaseA
(100mg/ml) (Qiagen cat. #19101) treatment before column purification. 20mg tissue samples from
brain, kidney, lung, skeletal muscle, and pancreas from a female koala, "Pacific Chocolate"
(Australian Museum registration M.45022), and a male koala, "Ben" (Australian Museum
registration M.47723), were supplied to the Ramaciotti Centre for Genomics for methylome
sequencing. The bisulfite conversion was carried out by using the EX DNA Methylation-Lightning
Kit (Zymo cat. #D5030) and the WGBS libraries were constructed using the TruSeq DNA
methylation kit (Illumina cat.# EGMK81213). The libraries were sequenced on a NovaSeq6000
S2 (Illumina) using the 2 × 100bp PE option. Information on samples, coverage, and read counts
are provided in Supplementary Table 1. Processing of the WGBS data followed previous studies
(30). Bisulfite conversion rates were estimated for each WGBS sample following the methodology
of methPipe's bsrate (43) (Supplementary Table 1). Strand-specific methylation calls were
combined, and all samples were filtered to remove CpGs covered by fewer than three reads.

Analyses of tissue differentially methylated regions

A hierarchical clustering tree was drawn using the fractional methylation profiles of all ten samples
using hclust from R's stats package. The distance matrix was calculated using Euclidean distances
and the agglomeration method used was Ward's method. The data for the final tree was visualized
using R's dendextend package (44). Bismark generated CpG reports were filtered to remove
scaffolds that were less than 2 Mb in length, retaining 3.03×10^9 (94.8%) of the genome. DMRs
were called using BSmooth (22), with additional filtering of a minimum fractional methylation
difference of 0.3 (30%) and at least 5 CpG sites. DMRs were considered shared between tissues if
they overlapped by at least 50%. Using koala gene annotations from Ensembl
(Phascolarctos_cinereus.phaCin_unsw_v4.1.97 release), promoters were defined as regions
located 1000 bp upstream of the identified transcription start site (TSS). To test the enrichment of
tissue-specific DMRs in functional regions, we generated 10,000 length and GC content matched
controls for all unique DMRs. The lung and kidney samples shared the most similar methylation
profiles and consequently had few tissue-specific DMRs (Fig. 1C). Because of this feature, the
corresponding gene sets were combined for these two tissues. Functional annotation and GO term
enrichment analysis was performed utilizing the ToppGene Suite (45).
295
Differential DNA methylation between sexes
Johnson et al. (25) used cross-species chromosome painting data and linked 406 scaffolds spanning
2.9 Gb of sequence data to autosomal scaffolds from chromosomes 1-7 and 24 scaffolds covering
68 Mb of sequence data to chromosome X, leaving 6.7% of the genome unclassified. To explore
the same relative amount of genomic space, we randomly sampled a subset of the autosomal
scaffolds that were length matched with the X chromosome scaffolds, which we called the
"matched autosome" dataset. These scaffolds were divided into 10-kb bins and the difference in
mean fractional methylation at each 10-kb bin was compared between male and female samples
for all tissues. For the analysis of human data, we used WGBS fractional methylation reports from
a male brain (Epigenome ID: E071) and a female brain (Epigenome ID: E053) and the human
known gene annotations from Ensembl (hg19 release). Due to its similarity in size to the human
X chromosome, we used data from human chromosome 8 as our representative autosome in the
comparative analysis.
309
310 **Identification of candidate X-linked scaffolds**

311 To isolate candidate X-linked scaffolds from the 1,477 unclassified koala scaffolds, we binned the

312 unclassified scaffolds into 10-kb windows and calculated the mean fractional methylation of the

313 associated CpGs. We then determined the average female and male methylation differences across

314 the bins and plotted the density of the differences for all five tissues. SVY and DS proceeded to

315 independently select scaffolds that exhibited a statistically significant shift towards female

316 hypomethylation from zero. The scaffolds that showed significant female hypomethylation in all

317 five tissues and were selected by both SVY and DS were isolated (n = 98 covering 14.6 Mb of

318 sequence with mean female-male 5mC = -0.25 ± 0.12). As an external validation, the percent of

319 reads mapping to the putative X-linked and autosome-linked scaffolds over the total number of

320 mapped reads was computed for the male and female sample in all tissues.

Annotation of the koala *Rsx*

323 To annotate the genomic region around *Rsx*, we downloaded the published genome sequence fasta

324 files for the partial assembly of *Rsx* from the gray short-tailed opossum (8) and the complete

325 PacBio assembly of the koala *Rsx* (25, 42). We used BLASTN 2.2.29 (46) to align both sequences

326 to the koala reference genome (phaCin_unsw_v4.1) and obtained genomic coordinates. The entire

327 assembled koala *Rsx* sequence aligned with 100% identity and no gaps. Only one 30.4 kb

328 transcript, a novel *lnc*RNA, overlapped with the annotated *Rsx* region (overlap > 90% of transcript)

329 and was used to evaluate gene expression.

Analysis of differential gene expression

332 All RNA-seq expression data used in this analysis were obtained from the previously published

333 koala transcriptomes from eight females and seven males (23). To process the acquired data, we

334 followed the protocol outlined by Pertea et al. (47), using the koala GTF annotation from Ensembl

335 (Phascolarctos_cinereus.phaCin_unsw_v4.1.97.gtf.gz release) to assemble mapped reads into

336 transcripts using StringTie 2.0 (47) with the -e-b--A <gene_abund.tab> flags. We used StringTie's

337 functionality for *de novo* transcript assembly to identify candidate *Rsx* transcripts. An updated

338 GTF annotation was generated including novel transcripts using the --merge flag and then the

339 previously generated mapped reads were reassembled into transcripts guided by this GTF file.

340 DeSeq2 1.22.2 (48) was used to perform differential gene expression analysis between males and

341 females. NOISeq 2.26.1 (24) was used for differential expression analysis due to its ability to
342 simulate technical replicates within given RNA-seq data sets when no replicates are available.
343

344 **Data accessibility**

345 The raw and processed methylation datasets generated in this study have been deposited and
346 accessible through GEO Series accession number GSE149600 at
347 https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE149600.
348

349 **Competing interests**

350 The authors declare that they have no competing interests.
351

352 **Authors' contributions**

353 D.S., D.A.P. and S.V.Y. formed the research design; D.A.P., A.G.K., D.E.A.P, and R.N.J.
354 generated the data; D.S., D.S. and S.V.Y. performed the analysis and wrote the initial draft ; all
355 authors contributed to and approved the final manuscript.
356

Funding statement

This work was supported by a grant from the National Science Foundation (MCB 1818288) and a
Pilot Grant from the Smooth Muscle Plasticity COBRE of the University of Nevada, Reno (funded
by the National Institutes of Health grant 5P30GM110767-04) to DAP, grants by the National
Science Foundation (MCB 1615664) and the National Institute of Health (R01MH103517) to
SVY. DS was partially supported by the NIH Training Grant in Computational Biology and
Biomedical Genomics (T32 GM105490).

References

1.   Lyon MF. Gene Action in the X-chromosome of the Mouse (Mus musculus L.). Nature.
1961;190(4773):372-3.
2.   Shevchenko AI, Zakharova IS, Zakian SM. The evolutionary pathway of x chromosome
inactivation in mammals. Acta Naturae. 2013;5(2):40-53.
3.   Brown CJ, Hendrich BD, Rupert JL, Lafreniere RG, Xing Y, Lawrence J, et al. The human
XIST gene: analysis of a 17 kb inactive X-specific RNA that contains conserved repeats and is
highly localized within the nucleus. Cell. 1992;71(3):527-42.

4.   Heard E, Clerc P, Avner P. X-chromosome inactivation in mammals. Annu Rev Genet.
1997;31:571-610.
5.   Plath K, Mlynarczyk-Evans S, Nusinow DA, Panning B. Xist RNA and the mechanism of
X chromosome inactivation. Annu Rev Genet. 2002;36:233-78.
6.   Duret L, Chureau C, Samain S, Weissenbach J, Avner P. The Xist RNA gene evolved in
eutherians by pseudogenization of a protein-coding gene. Science. 2006;312(5780):1653-5.
7.   Ng K, Pullirsch D, Leeb M, Wutz A. Xist and the order of silencing. EMBO Rep.
2007;8(1):34-9.
8.   Grant J, Mahadevaiah SK, Khil P, Sangrithi MN, Royo H, Duckworth J, et al. Rsx is a
metatherian RNA with Xist-like properties in X-chromosome inactivation. Nature. 2012;487:254.
9.   Sharman GB. Late DNA Replication in the Paternally Derived X Chromosome of Female
Kangaroos. Nature. 1971;230(5291):231-2.
10.   Wang X, Douglas KC, Vandeberg JL, Clark AG, Samollow PB. Chromosome-wide
profiling of X-chromosome inactivation and epigenetic states in fetal brain and placenta of the
opossum, Monodelphis domestica. Genome Res. 2014;24(1):70-83.
11.   Huynh KD, Lee JT. Inheritance of a pre-inactivated paternal X chromosome in early mouse
embryos. Nature. 2003;426(6968):857-62.
12.   Okamoto I, Otte AP, Allis CD, Reinberg D, Heard E. Epigenetic dynamics of imprinted X
inactivation during early mouse development. Science. 2004;303(5658):644-9.
13.   Heard E. Delving into the diversity of facultative heterochromatin: the epigenetics of the
inactive X chromosome. Current opinion in genetics & development. 2005;15(5):482-9.
14.   Koina E, Chaumeil J, Greaves IK, Tremethick DJ, Graves JA. Specific patterns of histone
marks accompany X chromosome inactivation in a marsupial. Chromosome research : an
international journal on the molecular, supramolecular and evolutionary aspects of chromosome
biology. 2009;17(1):115-26.
15.   Ingles ED, Deakin JE. Global DNA Methylation patterns on marsupial and devil facial
tumour chromosomes. Mol Cytogenet. 2015;8:74-.
16.   Rens W, Wallduck MS, Lovell FL, Ferguson-Smith MA, Ferguson-Smith AC. Epigenetic
modifications on X chromosomes in marsupial and monotreme mammals and implications for
evolution of dosage compensation. Proceedings of the National Academy of Sciences.
2010;107(41):17657.
17.   Piper AA, Bennett AM, Noyce L, Swanton MK, Cooper DW. Isolation of a clone partially
encoding hill kangaroo X-linked hypoxanthine phosphoribosyltransferase: Sex differences in
methylation in the body of the gene. Somatic Cell and Molecular Genetics. 1993;19(2):141-59.
18.   Loebel D, Johnston PG. Methylation analysis of a marsupial X-linked CpG island by
bisulfite genomic sequencing. Genome Res. 1996;6(2):114-23.
19.   Waters SA, Livernois AM, Patel H, O'Meally D, Craig JM, Marshall Graves JA, et al.
Landscape of DNA Methylation on the Marsupial X. Molecular Biology and Evolution.
2018;35(2):431-9.
20.   Sun Z, Cunningham J, Slager S, Kocher J-P. Base resolution methylome profiling:
considerations in platform selection, data preprocessing and analysis. Epigenomics.
2015;7(5):813-28.
21.   Kasai F, O'Brien PCM, Pereira JC, Ferguson-Smith MA. Marsupial chromosome DNA
content and genome size assessed from flow karyotypes: invariable low autosomal GC content. R
Soc Open Sci. 2018;5(8):171539-.

22.   Hansen KD, Langmead B, Irizarry RA. BSmooth: from whole genome bisulfite sequencing
reads to differentially methylated regions. Genome Biol. 2012;13(10):R83.
23.   Hobbs M, Pavasovic A, King AG, Prentis PJ, Eldridge MDB, Chen Z, et al. A
transcriptome resource for the koala (Phascolarctos cinereus): insights into koala retrovirus
transcription and sequence diversity. BMC Genomics. 2014;15(1):786.
24.   Tarazona S, Furio-Tari P, Turra D, Pietro AD, Nueda MJ, Ferrer A, et al. Data quality
aware analysis of differential expression in RNA-seq with NOISeq R/Bioc package. Nucleic acids
research. 2015;43(21):e140.
25.   Johnson RN, O'Meally D, Chen Z, Etherington GJ, Ho SYW, Nash WJ, et al. Adaptation
and conservation insights from the koala genome. Nature Genetics. 2018;50(8):1102-11.
26.   Hellman A, Chess A. Gene body-specific methylation on the active X chromosome.
Science. 2007;315:1141-3.
27.   Keown CL, Berletch JB, Castanon R, Nery JR, Disteche CM, Ecker JR, et al. Allele-
specific non-CG DNA methylation marks domains of active chromatin in female mouse brain.
Proceedings of the National Academy of Sciences. 2017;114(14):E2882.
28.   Sun D, Maney DL, Layman TS, Chatterjee P, Yi SV. Regional epigenetic differentiation
of the Z Chromosome between sexes in a female heterogametic system. Genome Res.
2019;29(10):1673-84.
29.   Schultz MD, He Y, Whitaker JW, Hariharan M, Mukamel EA, Leung D, et al. Human body
epigenome maps reveal noncanonical DNA methylation variation. Nature. 2015;523(7559):212-
6.
30.   Mendizabal I, Shi L, Keller TE, Konopka G, Preuss TM, Hsieh TF, et al. Comparative
Methylome Analyses Identify Epigenetic Regulatory Loci of Human Brain Evolution. Mol Biol
Evol. 2016;33(11):2947-59.
31.   Zemach A, McDaniel IE, Silva P, Zilberman D. Genome-wide evolutionary analysis of
eukaryotic DNA methylation. Science. 2010;328(5980):916-9.
32.   Yi SV. Birds do it, bees do it, worms and ciliates do it too: DNA methylation from
unexpected corners of the tree of life. Genome Biol. 2012;13(10):174.
33.   Jones PA. Functions of DNA methylation: islands, start sites, gene bodies and beyond.
Nature reviews Genetics. 2012;13(7):484-92.
34.   Chuang T-J, Chen F-C, Chen Y-Z. Position-dependent correlations between DNA
methylation and the evolutionary rates of mammalian coding exons. Proceedings of the National
Academy of Sciences. 2012;109(39):15841.
35.   Brenet F, Moh M, Funk P, Feierstein E, Viale AJ, Socci ND, et al. DNA methylation of
the first exon is tightly linked to transcriptional silencing. PloS one. 2011;6(1):e14524.
36.   Anastasiadi D, Esteve-Codina A, Piferrer F. Consistent inverse correlation between DNA
methylation of the first intron and gene expression across tissues and species. Epigenetics &
Chromatin. 2018;11(1):37.
37.   Neri F, Rapelli S, Krepelova A, Incarnato D, Parlato C, Basile G, et al. Intragenic DNA
methylation prevents spurious transcription initiation. Nature. 2017;543(7643):72-7.
38.   Huh I, Zeng J, Park T, Yi SV. DNA methylation and transcriptional noise. Epigenetics &
chromatin. 2013;6(1):9-.
39.   Cotton AM, Price EM, Jones MJ, Balaton BP, Kobor MS, Brown CJ. Landscape of DNA
methylation on the X chromosome reflects CpG density, functional chromatin state and X-
chromosome inactivation. Hum Mol Genet. 2015;24(6):1528-39.

40. Elango N, Kim S-H, Program NCS, Vigoda E, Yi SV. Mutations of different molecular
origins exhibit contrasting patterns of regional substitution rate variation. PLoS Computational
Biology. 2008;4(2):e1000015.
41. Alvarez-Ponce D, Torres-Sanchez M, Feyertag F, Kulkarni A, Nappi T. Molecular
evolution of DNMT1 in vertebrates: Duplications in marsupials followed by positive selection.
PloS one. 2018;13(4).
42. Sprague D, Waters SA, Kirk JM, Wang JR, Samollow PB, Waters PD, et al. Nonlinear
sequence similarity between the Xist and Rsx long noncoding RNAs suggests shared functions of
tandem repeat domains. RNA. 2019;25(8):1004-19.
43. Song Q, Decato B, Hong EE, Zhou M, Fang F, Qu J, et al. A Reference Methylome
Database and Analysis Pipeline to Facilitate Integrative and Comparative Epigenomics. PLOS
ONE. 2013;8(12):e81148.
44. Galili T. dendextend: an R package for visualizing, adjusting and comparing trees of
hierarchical clustering. Bioinformatics. 2015;31(22):3718-20.
45. Chen J, Bardes EE, Aronow BJ, Jegga AG. ToppGene Suite for gene list enrichment
analysis and candidate gene prioritization. Nucleic acids research. 2009;37(Web Server
issue):W305-11.
46. Zhang Z, Schwartz S, Wagner L, Miller W. A greedy algorithm for aligning DNA
sequences. Journal of computational biology : a journal of computational molecular cell biology.
2000;7(1-2):203-14.
47. Pertea M, Kim D, Pertea GM, Leek JT, Salzberg SL. Transcript-level expression analysis
of RNA-seq experiments with HISAT, StringTie and Ballgown. Nature Protocols.
2016;11(9):1650-67.
48. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550.
49. Ogawa Y, Kakumoto K, Yoshida T, Kuwako K-i, Miyazaki T, Yamaguchi J, et al. Elavl3
is essential for the maintenance of Purkinje neuron axons. Scientific Reports. 2018;8(1):2722.

490
491

Figure Legends

Figure 1. Overview of DNA methylation patterns across the koala genome. (A) Hierarchical clustering of DNA methylation of five tissues. (B) Tissue-specific and shared differentially methylation regions (DMRs) between tissues. (C) Enrichments of DMRs in each functional region in comparison to length and GC matched control regions (***$p < 0.0001$, n.s. Not significant, from 10,000 bootstraps) from brain samples. Error bars depict standard deviation. (D-E) Examples of two brain-specific DMRs (highlighted in grey) with corresponding CpG fractional methylation for the two brain (red) and eight remaining tissues (blue). Line smoothing performed using local regression (LOESS). (D) *ELAVL3* is a neural specific RNA-binding protein linked to the maintenance of Purkinje neuron axons (49). A 1.84 kb region containing three CpG Islands (CGI) and overlapping the first exon of *ELAVL3* was hypermethylated in the brain compared to all other tissues, and this gene was up-regulated in the brain compared to the kidneys (probability of differential expression > 98% from NOISeq). (D) *ALDOC* encodes a catalytic enzyme responsible for the conversion of fructose-1,6-bisphosphate to glyceraldehyde 3-phosphate and dihydroxyacetone phosphate. A 945 bp, brain-specific hypomethylated DMR overlaps the promoter and part of *ALDOC*'s gene body. This gene was up-regulated in brain samples compared to kidney samples (probability of differential expression > 96% from NOISeq).

Figure 2. Global patterns of female and male DNA methylation (5mC) in eutherian and marsupial X chromosomes. (A) Comparison of chromosome-wide DNA methylation between sexes in human and koala brains. (B) Distributions of the DNA methylation difference between female and male koalas in autosomes and the X chromosome. (C) Comparing fractional DNA methylation between males and females demonstrate that the female X chromosomes are hypomethylated in both humans and koalas. (D-F) DNA methylation differences between males and females of (D) the X chromosome and (E) autosomes of koalas, and (F) the X chromosome of humans for all male (purple) and female (orange) samples. Line smoothing was performed using local regression (LOESS). (G) Average fractional methylation of CpGs in 100-bp sliding windows using a 10 bp step size in a 5 Kb region upstream and downstream of all chromosome X linked gene's transcription start sites (TSSs) across koala tissues.

Figure 3. Annotation of genomic of DNA methylation (5mC) around *Rsx*. The top panel

identifies CpG islands (CGI), the middle panel reports the absolute male (purple) and female

(orange) fractional methylation at each CpG, and the bottom panel shows the female and male

fractional methylation difference. Highlighted in grey across all panels is the female

hypomethylated region upstream of the *Rsx* TSS.

Figure 4. Model of DNA methylation (5mC) patterns across genomic functional regions for

eutherian mammals and marsupials. Eutherian mammals exhibit increased DNA methylation

of promoter regions and CpG islands coupled with a relative depletion of DNA methylation across

gene bodies and intergenic regions on the inactive X chromosome (X_I) compared to the active X

chromosome (X_A) in females. Marsupial mammals share similar DNA methylation depletion in

gene bodies and intergenic regions of the inactive X chromosome; however, they diverge from

eutherian mammals in their promoter methylation patterns. Marsupial promoters are modestly

hypomethylated in the female X chromosomes (X_A and X_I) compared to the male X chromosome

(X_M).

Tables

Table 1. Functional annotation of enriched biological processes associated with gene sets containing tissue-specific differentially methylated regions (DMRs).

Tissue	GO biological process term	Accession ID	p-value	q-value
Brain	central nervous system development	GO:0007417	5.79×10^{-13}	2.71×10^{-09}
Brain	generation of neurons	GO:0048699	1.47×10^{-12}	3.45×10^{-09}
Brain	head development	GO:0060322	2.84×10^{-12}	3.78×10^{-09}
Brain	Neurogenesis	GO:0022008	3.22×10^{-12}	3.78×10^{-09}
Brain	brain development	GO:0007420	4.98×10^{-12}	4.67×10^{-09}
Pancreas	response to endoplasmic reticulum stress	GO:0034976	2.14×10^{-11}	1.51×10^{-07}
Pancreas	oxoacid metabolic process	GO:0043436	1.58×10^{-10}	4.04×10^{-07}
Pancreas	organic acid metabolic process	GO:0006082	1.72×10^{-10}	4.04×10^{-07}
Pancreas	response to endogenous stimulus	GO:0009719	6.34×10^{-09}	1.12×10^{-05}
Pancreas	carboxylic acid metabolic process	GO:0019752	2.29×10^{-08}	3.24×10^{-05}
Skeletal Muscle	actin filament-based process	GO:0030029	1.26×10^{-09}	5.78×10^{-06}
Skeletal Muscle	actin cytoskeleton organization	GO:0030036	5.04×10^{-09}	1.16×10^{-05}
Skeletal Muscle	cytoskeleton organization	GO:0007010	1.50×10^{-08}	2.30×10^{-05}
Skeletal Muscle	cellular carbohydrate metabolic process	GO:0044262	2.19×10^{-08}	2.52×10^{-05}
Skeletal Muscle	embryonic morphogenesis	GO:0048598	2.37×10^{-07}	2.17×10^{-04}
Lung and Kidney	embryonic skeletal system morphogenesis	GO:0048704	1.50×10^{-10}	3.52×10^{-07}
Lung and Kidney	embryonic organ morphogenesis	GO:0048562	1.61×10^{-10}	3.52×10^{-07}
Lung and Kidney	embryonic skeletal system development	GO:0048706	5.24×10^{-10}	7.65×10^{-07}
Lung and Kidney	embryonic organ development	GO:0048568	1.13×10^{-08}	1.23×10^{-05}
Lung and Kidney	pattern specification process	GO:0007389	3.31×10^{-08}	2.90×10^{-05}

Gene ontology (GO) terms are presented for the top five most significantly enriched results of each tissue after correcting for multiple testing (FDR < 0.05). As the numbers of tissue-specific DMRs for lung (n=22) and kidney (n=119) samples were so few, the corresponding gene sets were combined for this analysis.

Table 2. Correlation analysis of mean promoter and gene body DNA methylation and ranked gene expression.

Tissue	Genomic Region	Gene Count	Rho (p-value)
Brain	Promoter	5,396	-0.08 (p $= 2.28 \times 10^{-9}$)
	Gene body	5,443	-0.16 (p $< 2.2 \times 10^{-16}$)
Kidney	Promoter	9,268	-0.12 (p $< 2.2 \times 10^{-16}$)
	Gene body	9,379	-0.12 (p $< 2.2 \times 10^{-16}$)
Lung	Promoter	9,192	-0.13 (p $< 2.2 \times 10^{-16}$)
	Gene body	9,265	-0.19 (p $< 2.2 \times 10^{-16}$)

Spearman's rank correlation coefficients (ρ) and associated significances are reported for all tissues with both whole genome bisulfite sequencing (WGBS) data and RNA-seq expression data.

Table 3. Sex-based differential expression of the *lnc*RNA *Rsx* utilizing different data subsets and expression quantification tools.

Expression Dataset	Tool	Female Count	Male Count	Significance
All Data (n = 15)	DeSeq2	6987.1	16	p-value $=$ 0.05
Matched Data (n = 6)	DeSeq2	6837.6	0	p-value $= 2.04 \times 10^{-30}$
Matched Data (n = 6)	NOISeq	7872.4	0.67	Probability $=$ 99.99%
Kidney Data (n = 2)	NOISeq	4074.4	0.68	Probability $=$ 99.99%

Normalized expression count values and significance of sex-based differential expression is shown for three data subsets using two expression quantification tools. All data refers to the dataset considering all 15 RNA-seq samples (7 male and 8 female). Matched data includes the tissues with both male and female RNA-seq samples (brain, kidney, and lung), and the kidney data is reported independently. DeSeq2 reports significance as an associated p-value from the Wald test while NOISeq reports a probability of differential expression threshold.

Figures

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Supplementary Tables

Supplementary Table 1. Overview of whole genome bisulfite sequencing (WGBS) data for all 10 koala samples.

Sample	Australian Museum registration	Name	Tissue	Mapped Reads	De-duplicated Reads	Total CpGs	Coverage (%)	Mean Depth	% Reads Cov > 3×	Bisulfite Conversion Rate
WGM145_01_S1	M.45022	Pacific Chocolate	Brain	602055711	228244836	16761785	97.5	13.99	93.4	98.2
WGM145_04_S2	M.47723	Ben	Brain	677593548	233893124	16761785	97.6	14.61	93.8	98.0
WGM145_06_S3	M.45022	Pacific Chocolate	Kidney	559911229	213274030	16761785	97.2	13.62	92.5	98.7
WGM145_08_S4	M.47723	Ben	Kidney	589121434	202094771	16761785	97.2	12.93	92.3	98.7
WGM145_09_S5	M.45022	Pacific Chocolate	Lung	637878987	210002505	16761785	97.5	13.8	93.2	98.6
WGM145_12_S6	M.47723	Ben	Lung	663204935	198555030	16761785	97.3	12.54	92.4	98.6
WGM145_14_S7	M.45022	Pacific Chocolate	Skeletal Muscle	592423486	168450992	16761785	96.7	10.93	90.0	98.7
WGM145_16_S8	M.47723	Ben	Skeletal Muscle	605979530	168220022	16761785	96.8	11.12	90.4	98.6
WGM145_19_S9	M.45022	Pacific Chocolate	Pancreas	563288200	159866998	16761785	96.2	9.857	88.0	98.6
WGM145_20_S10	M.47723	Ben	Pancreas	598508860	166573663	16761785	96.6	10.56	89.4	98.7

Supplementary Table 2. GC-content for all scaffolds linked to annotated autosomes (chromosomes 1-7) and the X chromosome.

Chromosome	Size (Mb)	Total GC Content (%)
1	730.67	38.72
2	595.75	38.58
3	480.35	39.35
4	412.55	39.38
5	299.29	39.56
6	248.35	38.66
7	250.28	39.20
X	67.91	42.02

Supplementary Table 3. Enrichment and significance of all tissue specific DMRs compared to length and GC matched control regions.

Tissue	Genomic Region	DMR Counts (%)	Enrichment	p-value
Pancreas	Promoter	52 (4.5%)	1.13	0.74
	Gene body	612 (52.5 %)	1.44	**< 0.0001**
	Intergenic	502 (43.0 %)	-1.40	**< 0.0001**
Brain	Promoter	17 (3.5%)	-1.35	0.25
	Gene body	256 (53.0 %)	1.38	**< 0.0001**
	Intergenic	210 (43.5%)	-1.34	**< 0.0001**
Skeletal Muscle	Promoter	19 (4.9 %)	-1.23	0.32
	Gene body	202 (51.8 %)	1.27	**< 0.0001**
	Intergenic	169 (43.3 %)	-1.26	**0.001**
Kidney	Promoter	5 (4.0 %)	-1.34	0.40
	Gene body	62 (50.0 %)	1.25	**< 0.0001**
	Intergenic	57 (46.0%)	-1.21	**0.04**
Lung	Promoter	1 (4.3 %)	-1.61	0.51
	Gene body	12 (52.2%)	1.37	**< 0.0001**
	Intergenic	10 (43.5%)	-1.31	0.14

Reported are the total counts of tissue specific differentially methylated regions (DMRs) falling within one of three annotated genomic regions: promoters, gene bodies, and intergenic regions. The enrichment of DMRs in each functional region is shown through a fold change comparison with a control dataset generated from 10,000 bootstraps using length and GC matched control regions. All significant p-values ($p < 0.05$) are highlighted in bold.

Supplementary Table 4. Mean and median sex-based DNA methylation difference calculated for all candidate X-scaffolds (n=98) by tissue.

Tissue	Median Male-Female 5mC	Mean Male-Female 5mC
Pancreas	-0.2282 ± 0.15	-0.2066 ± 0.12
Brain	-0.3273 ± 0.14	-0.2911 ± 0.12
Skeletal Muscle	-0.2748 ± 0.15	-0.2431 ± 0.12
Kidney	-0.2850 ± 0.15	-0.2567 ± 0.12
Lung	-0.2706 ± 0.14	-0.2437 ± 0.11
Combined Tissues	-0.2773 ± 0.15	-0.2484 ± 0.12

The female and male mean fractional DNA methylation (methylated reads/total reads per CpG) was calculated for all CpGs within 10 kb bins across candidate scaffold.

Supplementary Figures

Supplementary Figure 1. Enrichment of tissue specific differentially methylated regions (DMRs) falling within genomic functional regions.

Data shown for (A) pancreas, (B) skeletal muscle, (C) kidney, and (D) lung. The enrichment of DMRs in each functional region (promoter, gene body, and intergenic regions) is shown through a comparison with length and GC matched control regions (*** indicates $p < 0.0001$, ** indicates $p < 0.001$, * indicates $p < 0.05$, and non-significance is shown by n.s. based on 10,000 bootstraps). Error bars indicate standard deviation.

Supplementary Figure 2. Correlation of gene expression and DNA methylation (5mC) in CpGs across promoters and gene bodies.

Three tissues with both whole genome bisulfite sequencing (WGBS) DNA methylation data and RNA-seq gene expression data are shown, (A) brain (n = 5,396 promoters and n = 5,443 gene bodies), (B) kidney (n = 9,268 promoters and n = 9,379 gene bodies), and (C) lung (n = 9,192 promoters and n = 9,265 gene bodies). For A-C, TPM expression values were ranked from low to high for each gene and correlated with mean fractional DNA methylation (methylated reads/total reads per CpG site). Spearman's rank correlation coefficients and the associated p-values are reported.

Supplementary Figure 3. Correlation of tissue dependent DNA methylation (5mC) and gene expression from brain and kidney samples.

(A) The mean brain and kidney DNA methylation difference calculated for all CpGs across each gene promoter matched with the corresponding log-transformed ratio of brain to kidney expression. (B) The mean brain and kidney DNA methylation difference calculated for all CpGs across each gene body and matched with corresponding log-transformed ratio of brain to kidney expression. For A and B, Spearman's rank correlation coefficient and the associated p-value is reported. Blue dots indicate genes that are significantly differentially express between brain and kidney samples (probability of differential expression > 95% based on NOISeq) and red dots show all genes that are significantly similarly methylated in brain and kidney samples (probability of differential expression < 5% based on NOISeq).

Supplementary Figure 4. Sex-specific CpG depth of coverage and read mapping to autosomes and X chromosomes.

(A) For each of the five tissues, box-and-whisker plots of CpG depths across the X chromosome in male (purple) and female (orange) samples (*** indicates p < 2.2x10-16, Mann-Whitney U test). Histogram and distribution of sex-based read mapping per sample (n=10) to (B) X-linked scaffolds, (C) candidate X-linked scaffolds, and (D) a subset of autosome-linked scaffolds matched in length with all known X-linked scaffolds. For A-C, the percent of reads mapping to the scaffold category of interested over the total number of mapped reads in the genome was calculated for all male (n=5) and female samples (n=5). The known X-linked and candidate X-linked scaffolds show a bimodal distribution with an increase of read mapping to female samples expected from the 2:1 ratio X chromosomes in females to males. This bimodality is not observed in autosomes.

Supplementary Figure 5. The distribution of sex-based CpG fractional DNA methylation (5mC) differences across autosomes and X chromosomes.

The distribution of female and male mean fractional methylation difference from (A) brain, (B) lung, (C) kidney, (D) skeletal muscle and (E) pancreas samples across autosomes and X chromosomes. For A-E, the female and male mean fractional methylation (methylated reads/total reads per CpG) was calculated for all CpGs within 10 kb bins across each autosome- or X-linked scaffold. All tissues exhibited a significant shift towards female hypomethylation in the X chromosome compared to the autosome (*** indicates p < 2.2x10-16, Welch's t-test).

Supplementary Figure 6. Female and male gene expression across autosomes and the X chromosome using kidney RNA-seq data.

(A) The relative genomic location of all genes linked to X chromosome scaffolds aligned by scaffold length and the log-transformed female to male expression ratio generated by NOISeq. Orange dots indicate the 36 genes with significant female-biased expression and purple dots indicate the 11 genes with significant male-biased expression (probability of differential expression > 95% based on NOISeq). For all autosome-linked genes (n = 10,414) and chromosome X-linked genes (n= 209), a box-and-whisker plot (B) and density distribution (C) of the log-transformed female to male expression ratio (p = 0.14, Mann-Whitney U test).

Supplementary Figure 7. Correlation analysis of sex-based DNA methylation (5mC) and gene expression across chromosome X using kidney WGBS and RNA-seq data.

(A) The mean female and male fractional methylation difference calculated for all CpGs across each gene promoter matched with the corresponding log-transformed ratio of female to male expression. (B) The ratio of the number female hypermethylated and female hypomethylated gene promoters that show either significant male of female biased expression over the total number of genes in each category. (C) The mean female and male fractional methylation difference calculated for all CpGs across each gene body and matched with corresponding log-transformed ratio of female to male expression. (D) The ratio of the number female hypermethylated and female hypomethylated gene bodies that show either significant male of female biased expression over the total number of genes in each category. For A and C, Spearman's rank correlation coefficient and the associated p-value is reported. The Rsx gene was excluded from the correlation calculation. Orange dots indicate the 36 genes with significant female-biased expression and purple dots indicate the 11 genes with significant male-biased expression (probability of differential expression > 95% based on NOISeq).

Supplementary Figure 8. Sex-based DNA methylation (5mC) by GC-content across autosomes and the X chromosome.

For A and B, the mean female and male methylation difference calculated from CpGs in 1 Kb bins across (A) autosomes and (B) X chromosomes. For C and D, mean female and male methylation difference calculated from CpGs located in promoter regions (defined as regions 1 kb upstream of known gene TSSs) in (C) autosomes and (B) X chromosomes. For A-D, data from all five tissues (brain, kidney, lung, pancreas, and skeletal muscle) are reported. All plots are coloured by data density where blue represents low density regions and yellow represents high density regions.