Divergent DNA methylation signatures underlying X chromosome regulation in marsupials and eutherians

Devika Singh[1], Dan Sun[1], Andrew G. King[2], David E. Alquezar-Planas[2], Rebecca N. Johnson[2,3], David Alvarez-Ponce[4],*, and Soojin V. Yi[1],*

[1]School of Biological Sciences, Georgia Institute of Technology, Atlanta, Georgia, USA;

[2]Australian Museum Research Institute, Australian Museum, Sydney, New South Wales, Australia;

[3]National Museum of Natural History, Smithsonian Institution, Washington, DC, USA;

[4]Department of Biology, University of Nevada Reno, Reno, Nevada, USA.

*Corresponding Authors:

David Alvarez-Ponce

Department of Biology

University of Nevada, Reno, 1664 N. Virginia Street, Reno, NV 89557

dap@unr.edu

775-682-5735

Soojin V. Yi

School of Biological Sciences

Georgia Institute of Technology, 950 Atlantic Drive, Atlanta, GA 30332.

soojin.yi@biology.gatech.edu

404-385-6084

Abstract

1
The phenomenon of X chromosome inactivation (XCI) mediated by sex-specific differences in
DNA methylation is well characterized in eutherian mammals. Although XCI is shared between
eutherians and marsupials, the role of marsupial DNA methylation in this process remains
contested. Here we examine genome-wide signatures of DNA methylation from methylation maps
across fives tissues from a male and female koala (*Phascolarctos cinereus*) and present the first
whole genome, multi-tissue marsupial "methylome atlas." Using these novel data, we elucidate
divergent versus common features of marsupial and eutherian DNA methylation and XCI. First,
tissue-specific differential DNA methylation in marsupials primarily occurs in gene bodies.
Second, the marsupial X chromosome is significantly globally less methylated (hypomethylated)
in females compared to males. We show that this pattern is also observed in eutherian X
chromosomes. Third, on average, promoter DNA methylation shows little difference between male
and female koala X chromosomes, a pattern distinct from that of eutherians. Fourth, the sex-
specific DNA methylation landscape upstream of *Rsx*, the primary *lnc*RNA associated with
marsupial X chromosome inactivation, is consistent with the epigenetic regulation of female- (and
presumably inactive X chromosome-) specific expression. Finally, we utilize the prominent female
X chromosome hypomethylation and classify 98 previously unplaced scaffolds as X-linked,
contributing an additional 14.6 Mb (21.5 %) to genomic data annotated as the koala X
chromosome. Our work demonstrates evolutionarily divergent pathways leading to the
functionally conserved pattern of X chromosome inactivation in two deep branches of mammals.
21
22

23 **Keywords**: X chromosome inactivation, marsupial, DNA methylation, dosage compensation,
24 RSX, koala, whole genome bisulfite sequencing (WGBS)

Introduction

34
X chromosome inactivation (XCI) is an iconic example of sex chromosome regulation in which

one of the two X chromosomes in females is silenced, a mechanism thought to adjust the

expression levels of X-linked genes (Lyon 1961). Although XCI has been observed in both

eutherian and marsupial mammals (Shevchenko, et al. 2013), there are several notable differences

in XCI between the two lineages. First, in eutherians, the transcription of *Xist* RNA exclusively

from the inactive X chromosome is essential for the X chromosome inactivation process (Brown,

et al. 1992; Heard, et al. 1997; Plath, et al. 2002). However, the *Xist* locus is not present in

marsupials (Duret, et al. 2006; Ng, et al. 2007). Instead, another *lnc*RNA gene, *Rsx*, is associated

with the marsupial X chromosome inactivation (Grant, et al. 2012). Second, marsupials

demonstrate imprinted XCI by selectively silencing the paternal X chromosome (Sharman 1971;

Wang, et al. 2014). In contrast, XCI in eutherians occurs randomly between the maternally and

paternally derived X chromosomes, although paternal XCI has been observed during early

development in rodents (Huynh and Lee 2003; Okamoto, et al. 2004). Third, in eutherians, XCI

involves the exclusion of active histone marks and the recruitment of repressive histone marks on

the inactive X chromosome (Heard 2005). In comparison, the inactive marsupial X chromosome,

while depleted of the active histone marks, shows variable enrichment patterns of repressive

histone marks (Koina, et al. 2009; Wang, et al. 2014). Specifically, out of five repressive marks

examined in two marsupial studies, H3K9me3, H4K20me3, and HP1α were enriched (Koina, et

al. 2009) while H3K27me3 and H3K9me2 (Rens, et al. 2010) were not enriched on the inactive X

chromosome. By contrast, all these marks are enriched on the inactive X in eutherians. These

differences indicate complex evolutionary pathways leading to the parallel patterns of XCI in the

two mammalian lineages and that novel insights into the mechanism of XCI can be gained from

comparative studies.

Given that the patterns of repressive histone marks are diverged between eutherian and

marsupial X chromosome inactivation, it is of interest to examine DNA methylation marks. In

eutherians, active and inactive X chromosomes clearly exhibit differential DNA methylation

(Hellman and Chess 2007; Keown, et al. 2017; Sun, et al. 2019). Specifically, the inactive X

chromosome shows increased DNA methylation of promoters and decreased DNA methylation

across gene bodies and intergenic regions compared to the active X chromosome. Because

promoters account for a relatively small portion of all genomic regions, collectively these DNA

methylation patterns result in a global reduction of methylation (hypomethylation) of the inactive

X chromosome in eutherians (Sun, et al. 2019). It remains unclear whether marsupial X

chromosomes undergo similar differential DNA methylation. Interestingly, marsupial genomes

harbor an additional copy of *DNMT1*, the gene encoding the DNA methyltransferase responsible

for DNA methylation maintenance (Leonhardt, et al. 1992; Goll and Bestor 2005), through gene

duplication (Alvarez-Ponce, et al. 2018), which could lead to functional divergence between the

mammalian lineages. Several previous studies found little difference in DNA methylation between

active and inactive marsupial X chromosomes (Piper, et al. 1993; Loebel and Johnston 1996;

Wang, et al. 2014). These studies generally focused on promoter CpG islands. DNA methylation

differences between the active and inactive X chromosomes, however, is a chromosome-wide

phenomenon where the patterns differ depending on the nature of genomic regions (e.g. promoter,

gene body, or intergenic regions) (Hellman and Chess 2007; Keown, et al. 2017; Sun, et al. 2019).

Therefore, a comprehensive and unbiased investigation is necessary to understand the role of DNA

methylation on X chromosome inactivation in marsupials.

The modern koala (*Phascolarctos cinereus*) is an Australian marsupial well-known for its

unique biology and life history. This marsupial is the sole surviving member of the family

Phascolarctidae following the loss of the Late Pleistocene giant koala (*P. stirtoni*) (Price 2008;

Black, et al. 2014). The recently completed koala genome assembly (Johnson, et al. 2018) offers

an opportunity to investigate the relationship between DNA methylation epigenetic modifications

and regulation of gene expression on the sex chromosome in a marsupial in detail. A previous

study characterized DNA methylation across the opossum X chromosome using reduced

representation bisulfite sequencing (Waters, et al. 2018), a method that enriches for

hypomethylated regions and generally over-represents promoters and CpG islands (Sun, et al.

2015). To extend the survey of DNA methylation to another marsupial species across all genomic

regions independent of their CpG contents, here we generated unbiased, nucleotide-resolution

genomic DNA methylation maps of diverse tissues from a male and a female koala using whole

genome bisulfite sequencing.

Our data provide the first multi-tissue, whole genome methylome resource of any

marsupial with information on tissue-specific variation of DNA methylation. Utilizing these data,

here we show previously unknown patterns of chromosome-wide DNA methylation differences

between male and female X chromosomes, indicating distinctive impacts of DNA methylation on

X chromosome inactivation in marsupials. We further classify previously undetected X-linked
regions from a key marsupial species using characteristic features of X chromosome DNA
methylation. Our findings provide new insights into the evolutionary pathways leading to
functionally convergent yet mechanistically divergent pathways of X chromosome inactivation
and regulation of gene expression in eutherian and marsupial mammals.
101
Results

Genome-wide patterns of DNA methylation in the modern koala

To analyze global DNA methylation (5mC) patterns, we generated whole genome bisulfite
sequencing (WGBS) data across five tissues (brain, lung, kidney, skeletal muscle, and pancreas)
from a male ("Ben," Australian Museum registration M.45022) and female koala ("Pacific
Chocolate," Australian Museum registration M.47723). Of the total 16,761,785 CpG sites in the
currently assembled koala scaffolds, 88-94% of CpGs had at least $3\times$ coverage (Supplementary
Table 1). The mean depth of coverage fell between $9.9\times$ and $14.6\times$. The total GC content across
all 1907 scaffolds covering 3.2 Gb was 39.05%. Autosome-linked scaffolds displayed similar GC
content ranging between 38.58 and 39.59%. Scaffolds linked to chromosome X had higher GC
contents compared to autosomal scaffolds (Supplementary Table 2), as previously observed
(Kasai, et al. 2018). Fractional methylation values of all CpGs were used for a hierarchical
clustering analysis between tissues (Fig. 1A). We observed a clear clustering of samples by tissue
with the pancreas samples exhibiting the most unique methylation signature while the kidney and
lung samples share the most similar methylation profiles.
117

Differential DNA methylation between tissues

We identified shared and tissue specific differentially methylated regions (DMRs) using BSmooth
(Hansen, et al. 2012). Tissue specific DMRs were defined as regions that were differentially
methylated in a particular tissue compared to all other tissues in a pairwise analysis while shared
DMRs were those found in more than one tissue. Fig. 1B shows the total number of shared and
tissue specific DMRs. Consistent with the results of the clustering analysis, the pancreas samples
contained the greatest number of tissue specific DMRs (N=1,135) followed by the skeletal muscle,
brain, kidney and lung (Fig. 1B). Interestingly, across all tissues we found that the majority (50-
53%) of tissue specific DMRs fell in gene bodies (Fig. 1C, Supplementary Fig. 1, and

Supplementary Table 3), a significant excess compared to length and GC matched control regions
(fold enrichment (FE) = 1.25~1.44, p < 0.0001 based on 10,000 bootstraps; Fig. 1C,
Supplementary Fig. 1 and Supplementary Table 3). Even though we observed numerous DMRs in
intergenic regions, they were significantly depleted in intergenic regions compared to the control
regions (p < 0.05 based on 10,000 bootstraps; Fig. 1C, Supplementary Fig. 1, and Supplementary
Table 3).

Genes containing tissue specific DMRs were enriched in specific biological functions
(Table 1). Brain specific DMRs were linked to genes associated with neural developmental
processes such as neurogenesis and central nervous system development. DMRs in the skeletal
muscle samples were found in genes associated with actin and cytoskeleton organization. Lung
and kidney DMRs were merged due to the fact that their methylation profiles were highly similar
and consequently both tissues had very few unique DMRs (n=22 and n=119 respectively) and were
linked to genes associated with several terms involving embryonic processes including organ
development and morphogenesis. Pancreas specific DMRs overlapped the largest number of
genes, which showed enrichment for several metabolic processes.

To explore the consequences of tissue specific DMRs on gene expression levels, we
integrated our methylome data with a previously generated RNA-seq koala transcriptome
reference (Hobbs, et al. 2014). Of the 12 tissues surveyed in that study, three tissues (kidney, brain,
and lung) overlapped with the current methylome dataset. From all methylation and expression
linked samples, we found significant negative correlations between promoter DNA methylation
and gene expression across the entire genome, similar to the patterns observed in eutherian
mammals (Table 2, Supplementary Fig. 2). The relationship between gene body DNA methylation
and expression was complex. Specifically, both extremely hypomethylated and hypermethylated
gene bodies showed high gene expression, again consistent with studies in eutherian mammals and
other taxa (Table 2, Supplementary Fig. 2, and also (Lister, et al. 2009; Zemach, et al. 2010; Jjingo,
et al. 2012; Spainhour, et al. 2019)). To directly compare the DNA methylation and expression
datasets, we considered differentially methylated genes (DMGs) containing DMRs between the
brain and kidney samples (n = 1944 genes from n = 4,615 DMRs). We then identified differentially
expressed genes (DEGs) between brain and kidney RNA-seq samples. Currently, available RNA-
seq data from koala do not include sufficient biological replicates. We overcame this limitation by
simulating replicates within each RNA-seq data set (NOISeq, (Tarazona, et al. 2015)) and

identified 600 putative DEGs (probability of differential expression > 95% according to the
NOISeq software). We found that DMGs were significantly more likely to be differentially
expressed than genes that did not contain DMRs, exhibiting a 1.54-fold enrichment of observed to
expected DMGs that were also DEGs, ($\chi^2 = 33.07$, p < 0.0001). In addition, differential expression
between tissues was significantly correlated with differential DNA methylation between tissues in
promoters (Supplementary Fig. 3A). In agreement with the observation from the whole genome,
both relative hypomethylation and relative hypermethylation of gene bodies were associated with
increased expression (Supplementary Fig. 3B). The methylation patterns of two representative
genes containing brain specific DMRs are shown in Fig. 1D and 1E.
167
Global hypomethylation of female X chromosome in koalas
The koala genome project used cross-species chromosome painting data to identify 24 putative X
chromosome scaffolds and 406 putative autosomal scaffolds consisting of 68 Mb and 2.9 Gb of
sequences respectively (Johnson, et al. 2018). We utilized this classification to examine sex
specific DNA methylation on the X chromosome and autosomes. As expected from 2:1 ratio of X
chromosomes in females compared to males, the median depth of coverage of CpGs on the putative
X scaffolds were significantly higher (approximately 2-fold higher) in female samples compared
to male samples for all five tissues (p < 2.2×10^{-16}, Mann-Whitney U test, Supplementary Fig.
4A). In addition, the percent of total reads mapped to the putative X scaffolds out of all the mapped
reads showed a distinct bimodal distribution whereby the male samples cluster close to 1.3% and
the female samples cluster near 2.4% (Supplementary Fig. 4B). In contrast, male and female
samples were indistinguishable with respect to read mapping to putative autosomes
(Supplementary Fig. 4D).
Next, we characterized sex-based DNA methylation distributions across the autosomes and
X chromosomes. We found that the global DNA methylation level of the female X chromosome
was lower than that of the male X chromosome in all koala tissues examined (Fig. 2A, B and
Supplementary Fig. 5, p < 2.2×10^{-16}, Mann-Whitney U test). In contrast, autosomes were similarly
methylated between females and males (Fig. 2A, B). A comparison to autosomal DNA methylation
indicated that the X chromosome exhibits reduced DNA methylation in females. Consequently,
we use the term 'female hypomethylation' (as opposed to male hypermethylation) (Fig. 2C)
consistently in this work.

To gain better insights into divergent patterns between male and female X chromosomes,
we examined DNA methylation in different functional regions (promoters, exons, introns, and
intergenic regions). We found that, on average, the female samples were hypomethylated in all
functional regions across the X chromosome (Fig. 2D). This difference was most pronounced in
gene bodies and intergenic regions, while promoters showed the least sex-based difference. As
expected, the autosomal scaffolds did not display a significant variation between female and male
methylation levels in any functional region (Fig. 2E). Fig. 2F depicts the DNA methylation
difference between male and female X chromosomes in humans. In contrast to the pattern observed
in koalas, promoters on the human female X chromosomes were hypermethylated compared to
those on the male X chromosome, congruent with previous studies (Hellman and Chess 2007;
Keown, et al. 2017; Sun, et al. 2019).

Promoter DNA methylation is not a universal driver of sex-specific expression in koalas
To investigate the implications of the observed sex-specific DNA methylation, we once again
utilized the published RNA-seq koala transcriptome reference (Hobbs, et al. 2014). Of the total
RNA-seq dataset, only one tissue, kidney, had expression data from both a male and female koala
that could be directly compared to our methylome dataset. Consequently, the other tissues were
not considered for further analysis. We calculated the log-transformed fold change of female and
male gene expression values using NOISeq, capitalizing on its ability to simulate technical
replicates. This analysis suggested that of the 209 X-linked genes, 36 (17.2%) exhibit female
overexpression while 11 (5.3%) show male overexpression (probability of sex-based differential
expression > 95% based on NOISeq, Supplementary Fig. 6A). Although, on average, autosomal
genes also exhibited slight female-bias of expression (Supplementary Fig. 6B, C), the increase is
more substantial in X chromosome linked genes (mean chromosome X female to male $\log_2$ fold
change = 0.50, autosome female to male expression $\log_2$ fold change = 0.24). Using the matched
WGBS and RNA-seq data from the kidney samples, we calculated the female and male fractional
methylation difference in X chromosome linked promoters and gene bodies and correlated them
with gene expression difference (N = 209 gene bodies and N = 206 promoters, excluding 3
promoters with CpGs coverage < 3). In promoters, we found no significant relationship between
female and male DNA methylation difference and female-biased gene expression (Supplementary
Fig. 7A). In fact, the proportions of significantly female-over expressed genes (probability of sex-

based differential expression > 95% based on NOISeq) were similar between female-

hypermethylated promoters and female-hypomethylated promoters (Supplementary Fig. 7B).

Interestingly, female and male DNA methylation difference in gene bodies showed an overall

negative correlation with gene expression (Spearman's rank correlation coefficient, ρ = -0.14, p =

0.04, Supplementary Fig. 7C). However, a deeper analysis of the relative DNA methylation and

expression levels revealed that both extreme hypomethylation and hypermethylation are associated

with increased expression (Supplementary Fig. 7C) attesting to the complexity and heterogeneity

of the relationship between gene body DNA methylation and gene expression (Zemach, et al. 2010;

Jjingo, et al. 2012; Spainhour, et al. 2019).

The *Rsx* region displays a pattern suggesting methylation driven control of X chromosome

regulation in koalas

Above we have demonstrated that male and female differential methylation was the most notable

in gene bodies and intergenic regions on the X chromosome. Nevertheless, sex-specific

methylation of a key regulator of the X inactivation (XCI) in other marsupials, the long non-coding

RNA gene *Rsx*, has been associated with its differential expression (Grant, et al. 2012; Wang, et

al. 2014). To determine if differential DNA methylation is similarly associated with differential

regulation of *Rsx* in koalas, we isolated a 42 Kb region to examine on the koala X chromosome

based on the sequence homology with the partial assembly of the *Rsx* gene from the gray short-

tailed opossum, *Monodelphis domestica* (Grant, et al. 2012). We then identified a 29.8 Kb

candidate *Rsx* sequence within the isolated region from the alignment to the full koala *Rsx*

sequence assembled using PacBio long read sequencing by Johnson et al. (Johnson, et al. 2018).

The total examined region spanned 266 CpGs, of which 165 were covered by more than three

reads in all 10 samples from the WGBS dataset. The candidate *Rsx* region harbored 93 CpGs over

this read depth threshold. Because the WGBS data allows us to explore the entire genomic region

around the candidate gene, we used these CpGs to examine the DNA methylation difference

between male and female koala samples. Interestingly, we observed a female hypomethylated

region containing two CpG islands upstream of the candidate *Rsx* covering 101 CpGs exhibiting a

36% reduction in mean female DNA methylation compared to mean male DNA methylation (mean

sex difference: -0.36 ± 0.14, Fig. 3). The mean sex-based DNA methylation difference within this

region was in the top 23% of the distribution of differences across all X-linked scaffolds of the
combined tissues.

Having identified an upstream female hypomethylated hotspot, we tested whether *Rsx*
expression was exclusive to females, consistent with it being the X chromosome inactivation
center in koalas (Hobbs, et al. 2014; Johnson, et al. 2018). We used DESeq2 (Love, et al. 2014) to
determine differential gene expression on curated RNA-seq data from 8 female and 7 male koala
samples obtained from 12 tissues (Hobbs, et al. 2014). As expected, the expression of the *lnc*RNA
annotated within the candidate *Rsx* gene region was significantly greater in females (mean
normalized read count = 6987.1) than in males (mean normalized read count = 16, p < 0.05 from
DeSeq2 using the Wald test). When considering a subset of tissue expression data with matched
male and female samples (spleen, kidney, and lung), the results remained robust across different
tools to measure differential gene expression (Table 3).

Identification of novel candidate X-linked scaffolds by sex specific methylation patterns
We have demonstrated above that X-linked scaffolds exhibit several distinctive WGBS patterns in
koalas. Namely, putative X scaffolds show different depths of coverage per CpG, distinctive
clustering based on the proportion of mapped reads, and unique methylation distributions between
females and males (Supplementary Fig. 4). We thus quantified DNA methylation differences
between females and males to determine if additional candidate X scaffolds existed within the
6.7% of the koala assembly that remained unclassified. We identified an additional 98 scaffolds
which showed a clear shift towards female hypomethylation (mean female-male 5mC for all
candidate X scaffolds was -0.25 ± 0.12) (Supplementary Table 4). These candidate scaffolds
contributed 14.6 Mb (21.5%) to the total genomic region annotated as the koala X chromosome.
All candidate scaffolds followed the expected bimodal distribution seen in the putative X scaffolds
when we analyzed the percent of total reads mapped to the candidate X scaffolds out of all the
mapped reads. The male samples clustered around 0.3% and the female samples clustered around
0.5% (Supplementary Fig. 4C). This clustering pattern could be attributed to the 2:1 ratio of X
chromosomes in females compared to males that was not observed in putative autosomes.

Discussion

In this study, we investigated nucleotide-resolution genomic DNA methylation maps of five tissues of a male and a female koala. The overall DNA methylation levels of koala tissues are on par with the patterns observed in other eutherian mammals (Schultz, et al. 2015; Mendizabal, et al. 2016; Keown, et al. 2017), and show a clear pattern of heavy genome-wide DNA methylation punctuated by hypomethylation of CpG islands (e.g., Fig. 1D,E). We identified tissue-specific differential DNA methylated regions (DMRs), which were consistently and significantly enriched in gene bodies. Previous studies of human tissues also observed that DMRs tended to occur in gene bodies (e.g. (Mendizabal and Yi 2015; Peters, et al. 2015; Schultz, et al. 2015)). Gene body methylation is an ancestral form of DNA methylation in animal genomes (e.g., (Zemach, et al. 2010; Yi 2012)), yet its role in regulation of gene expression is less well understood than promoter DNA methylation (Zemach, et al. 2010; Jones 2012). Integrating relative DNA methylation differences and relative gene expression differences, we observe that tissue-specific hypo- and hyper methylation of gene bodies both contribute to increased gene expression (Supplementary Fig. 3B) in koalas, a characteristic previously observed in eutherian mammals and in other taxa (Lister, et al. 2009; Zemach, et al. 2010; Jjingo, et al. 2012; Mendizabal, et al. 2016). Collectively, these observations suggest that gene body methylation may be an important component of regulation of gene expression in koala.

The patterns of DNA methylation across the marsupial X chromosome show significant differences between sexes. Specifically, the female X chromosomes are globally hypomethylated compared to both the male X chromosome and the autosomes of both sexes (Fig. 2). Most of the literature of X chromosome inactivation emphasizes that DNA methylation is increased in promoters of the inactive X chromosomes compared to those of the active X chromosome. However, we show that the hypomethylation of the female X chromosome observed in koala is in fact consistent with DNA methylation data from eutherian mammals. We summarized the current knowledge from eutherian mammals and our proposed model for marsupials in Fig. 4. Concisely, in eutherians, while DNA methylation of promoter regions and CpG islands increase on the inactive X chromosome, gene bodies and intergenic regions of the inactive X chromosome are relatively depleted of DNA methylation compared to the active X chromosome (Fig. 4). Hellmann and Chess (Hellman and Chess 2007) demonstrated the difference between promoter and gene body DNA methylation of active and inactive X chromosomes in humans. An array based DNA

methylation study of humans (Cotton, et al. 2015) and whole genome bisulfite sequencing analyses

of mouse (Keown, et al. 2017) and humans (Sun, et al. 2019) showed that the hypomethylation of

inactive X chromosome is pervasive in gene bodies and intergenic regions.

In marsupials, Rens et al. (Rens, et al. 2010) observed hypomethylation of the inactive X

chromosome through immuno-staining using antibodies against methyl-cytosines in two species,

common brushtail possum (*Trichosurus vulpecula*) and long-nosed potoroo (*Potorous tridactylus*).

Waters et al. (Waters, et al. 2018) have also observed female hypomethylation of the X

chromosome in gray-tailed opossum (*Monodelphis domestica*) using reduced representation

bisulfite sequencing (RRBS). In contrast to Keown et al. (Keown, et al. 2017), Waters et al.

(Waters, et al. 2018) did not observe substantial hypomethylation of mouse X chromosomes. They

thus concluded that the reduction of methylation in gene bodies was specific to marsupials (Waters,

et al. 2018). The discrepancy between Keown et al. (Keown, et al. 2017) and Waters et al. (Waters,

et al. 2018) could be due to the inherent bias of the RRBS method, used in the latter study, which

disproportionately samples regions with high GC content (Sun, et al. 2015). High GC-content

regions tend to be hypomethylated (Elango, et al. 2008; Cotton, et al. 2015) and show less variation

of DNA methylation between males and females (Cotton, et al. 2015), which we also demonstrate

below (Supplementary Fig. 7).

It was previously suggested that differential DNA methylation does not play substantial

roles in marsupial X chromosome gene regulation (Piper, et al. 1993; Loebel and Johnston 1996;

Wang, et al. 2014) due to the lack of DNA methylation difference in promoters and CpG islands

(also in (Waters, et al. 2018)). Here we show that promoters are, on average, clearly

hypomethylated in female tissues compared to male tissues in koalas, although the degree of

methylation difference is small compared to other genomic regions. This modest sex-based

difference may at least partly be attributed to the fact that promoters tend to be GC-rich

(Supplementary Fig. 7). Due to the negative correlation between GC-content and DNA

methylation, these lowly methylated regions are inherently less variable and would subsequently

show subtle differences between sexes ((Cotton, et al. 2015) and Supplementary Fig. 7).

Summarizing these findings, we observe that the hypermethylation of promoters on the inactive X

chromosome appears to be restricted to eutherian mammals, while chromosome-wide

hypomethylation of female X chromosome (Fig. 4), consistent with the hypomethylation of the

inactive X chromosome, is conserved between eutherian and marsupial mammals. How

chromosome-wide DNA hypomethylation is linked to chromosome-wide gene silencing is
currently unknown, and likely to involve other epigenetic marks. Analyses of DMNT expression,
including that of a newly duplicated *DNMT1* (Alvarez-Ponce, et al. 2018), did not indicate
significant differential expression between sexes (Results not shown). However, additional
analysis of data with broad and balanced sampling of both sexes is necessary to rigorously test the
impact of the marsupial *DNMT1* duplication on sex specific gene expression.

Interestingly, despite the fact that gene regulation by sex-specific promoter DNA
methylation is not a global phenomenon in marsupials, the *lnc*RNA *Rsx,* the major player in XCI
initiation in marsupials (Grant, et al. 2012), presents an exception. Similar to the mechanism of
activation of the *Xist* RNA in eutherian mammals, *Rsx* is exclusively transcribed from and coats
the inactive X chromosome initiating the XCI process through which the inactive X chromosome
adopts a specific chromatin configuration within cells (Galupa and Heard 2018). Observed
signatures unique to female koala samples suggest that *Rsx* expression is regulated by promoter
DNA methylation, while gene body methylation of *Rsx* is similar between males and females (Fig.
3). Wang et al. (Wang, et al. 2014) showed that the *Rsx* promoter in opossum is also regulated by
differential DNA methylation of CpG islands. Therefore, regulation of the key initiator of XCI via
DNA methylation is another parallel feature between eutherians and marsupials, although utilizing
distinctive components. In addition, a recent study (Sprague, et al. 2019) found that *Xist* and *Rsx*
harbor non-linear sequence similarity. Consequently, their shared functionality may be partially
due to characteristics of tandem repeat regions.

Therefore, we show that the overarching hypomethylation of female X chromosome is a
conserved feature of X chromosome regulation in eutherian and marsupial mammals. However, X
chromosome promoter methylation and the subsequent effect on the regulation of gene expression
between males and females appear to be divergent between these two lineages. Regulation of the
Rsx, on the other hand, is supported by DNA methylation, which mirrors the regulation of the
eutherian *Xist* locus. Together, these conclusions illuminate the intricate evolutionary pathways
that have diverged and converged to influence X chromosome regulation, XCI, and dosage
compensation in eutherian and marsupial mammals.

371
372
373

Materials and Methods

Whole Genome Bisulfite Sequencing Data Generation

All genomic DNA was extracted using a Bioline Isolate II Genomic DNA Extraction Kit (Cat#. BIO-52067) following the recommended protocol with an additional DNAse free RNaseA (100mg/ml) (Qiagen cat. #19101) treatment before column purification. 20mg tissue samples from brain, kidney, lung, skeletal muscle, and pancreas from a female koala, "Pacific Chocolate" (Australian Museum registration M.45022), and a male koala, "Ben" (Australian Museum registration M.47723), were supplied to the Ramaciotti Centre for Genomics for methylome sequencing. The bisulfite conversion was carried out by using the EX DNA Methylation-Lightning Kit (Zymo cat. #D5030) and the WGBS libraries were constructed using the TruSeq DNA methylation kit (Illumina cat.# EGMK81213). The libraries were sequenced on a NovaSeq6000 S2 (Illumina) using the 2×100bp PE option. Information on samples, coverage, and the numbers of reads are provided in Supplementary Table 1.

WGBS Data Processing

Paired-end raw fastq files were processed for quality control and adapter trimming using Trim Galore! (version 0.3.7) with a Phred score threshold of 20. Trimmed reads were filtered for phi-X174 (NC_001422.1) to remove the spike-in control before being mapped to the koala reference assembly (phaCin_unsw_v4.1; (Johnson, et al. 2018)) using Bismark with the Bowtie 2 setting and deduplicated. The current koala genome reference assembly was generated from the same female koala ("Pacific Chocolate," Australian Museum registration M.45022) that we used to obtain the female WGBS data. Bisulfite conversion rates were estimated for each WGBS sample following the methodology of methPipe's bsrate (Song, et al. 2013) (Supplementary Table 1). Strand-specific methylation calls were combined, and all samples were filtered to remove CpGs covered by fewer than three reads. See Supplementary Table 1 for an overview of the dataset.

Analyses of tissue differentially methylated regions

A hierarchical clustering tree was drawn using the fractional methylation profiles of all ten samples representing five tissues using *hclust* from R's stats package. The distance matrix was calculated using Euclidean distances and the agglomeration method used was Ward's method. The data for the final tree was visualized using R's dendextend package (Galili 2015).

Before identifying differentially methylated regions (DMRs), all Bismark generated CpG reports

were filtered to remove scaffolds that were less than 2 Mb in length. Following this step, we

retained 3.03×10^9 positions for downstream analyses, covering 94.8% of the genome. DMRs

were called using BSmooth (Hansen, et al. 2012) in 10 pairwise comparisons of all five tissues.

Detected DMRs with a minimum fractional methylation difference of 0.3 (30%) between each pair

of compared tissues that also contained at least 5 CpG sites were retained. Shared DMRs were

defined as those that overlapped at least 50% in multiple tissues. The distribution of common and

tissue specific DMRs located among different genomic functional regions (promoters, gene bodies,

and intergenic regions) were identified based on koala gene annotations from Ensembl

(Phascolarctos_cinereus.phaCin_unsw_v4.1.97 release). Promoters were defined as regions

located 1000 bp upstream of the identified transcription start site (TSS). To test the enrichment of

tissue specific DMRs in the previously specified genomic regions, we generated 10,000 length and

GC content matched control regions for all unique DMRs, and calculated the p-values from the

ratio of the number of simulations with values at least as extreme as the observed ones over the

total number of simulations.

The genes containing tissue specific DMRs were extracted to make a set of differentially

methylated genes unique to each analyzed tissue. As mentioned above, the lung and kidney

samples shared the most similar methylation profiles and consequently had few tissue specific

DMRs (Fig. 1C). Because of this feature, the corresponding gene sets were combined for these

two tissues. Functional annotation and GO term enrichment analysis was performed utilizing the

ToppGene Suite (Chen, et al. 2009).

Quantifying sex differences in DNA methylation across chromosomes and functional regions

Johnson et al. (Johnson, et al. 2018) used cross-species chromosome painting data and linked 406

scaffolds spanning 2.9 Gb of sequence data to autosomal scaffolds from chromosomes 1-7 and 24

scaffolds covering 68 Mb of sequence data to chromosome X, leaving 6.7% of the genome

unclassified. To explore the same relative amount of genomic space, we randomly sampled a

subset of the autosomal scaffolds that were length matched with the X chromosome scaffolds,

which we called the "matched autosome" dataset. Separately, all matched autosome and

chromosome X scaffolds were divided into 10-kb bins and the mean fractional methylation levels

of the corresponding CpGs were calculated for each of the 5 tissue samples. The difference in
mean fractional methylation at each 10-kb bin was computed between male and female samples
for all tissues. For our comparative analysis with human sex-specific DNA methylation, we
downloaded publicly available WGBS fractional methylation reports from a male brain
(Epigenome ID: E071) and a female brain (Epigenome ID: E053) generated by the Roadmap
Epigenomics Consortium (Kundaje, et al. 2015) and used CpGs with minimum $3\times$ coverage. Due
to its similarity in size to the human X chromosome, we used data from human chromosome 8 as
our representative autosome in the comparative analysis.

For the analysis of sex-based fractional DNA methylation across different genomic
functional regions, we used the previously defined koala gene annotations from Ensembl
(Phascolarctos_cinereus.phaCin_unsw_v4.1.97 release) and human known gene annotations from
Ensembl (hg19 release). Individual annotated promoters, exons, introns, and intergenic regions
were evenly divided into 20 bins and the mean fractional methylation of CpGs within each bin was
computed.
450

Identification of candidate X-linked scaffolds
To isolate candidate scaffolds potentially located on the X chromosome from the 1,477
unclassified scaffolds covering 6.7% of the koala genome, we binned the remnant scaffolds into
10-kb windows and calculated the mean fractional methylation of the associated CpGs. We then
determined the average female and male methylation differences across the bins and plotted the
density of the differences for all five tissues. SVY and DS proceeded to independently select
scaffolds that exhibited a statistically significant shift towards female hypomethylation from zero
based on an initial visual examination. The scaffolds that showed significant female
hypomethylation in all five tissues and were selected by both SVY and DS were then further
analyzed (n = 98 covering 14.6 Mb of sequence with mean female-male 5mC = -0.25 ± 0.12). As
a baseline for an external validation, the percent of reads mapping to the putative X-linked and
autosome-linked scaffolds over the total number of mapped reads was computed for the male and
female sample in all tissues. The expectation was that the female samples would show and increase
in reads mapped to the X-linked scaffolds compared to the male samples due to the 2:1 ratio of X
chromosomes in females to males. This sex-biased separation is not expected across length
matched autosome-linked scaffolds as autosome counts are balanced in both sexes. Once this

baseline analysis was confirmed, we calculated the percent of reads mapping to the candidate X-

linked scaffolds over the total number of mapped reads from all 10 samples.

Annotation of the *Rsx* Region

To annotate the genomic region around *Rsx*, we downloaded the published genome sequence fasta

files for the partial assembly of *Rsx* from the gray short-tailed opossum (Grant, et al. 2012) and

the complete PacBio assembly of the koala *Rsx* (Johnson, et al. 2018; Sprague, et al. 2019). We

then used BLASTN 2.2.29 (Zhang, et al. 2000) to align both sequences to the koala reference

genome (phaCin_unsw_v4.1) and obtained genomic coordinates information. As expected, both

sequences aligned to a scaffold previously linked to the X chromosome and overlapped with one

another. The entire assembled koala *Rsx* sequence aligned with 100% identity and no gaps.

RNA-seq Data Processing

All RNA-seq expression data used in this analysis were obtained from the previously published

koala transcriptome reference (Hobbs, et al. 2014). This dataset included eight tissues from a

female koala, spleen (SRR1203868), liver (SRR1205138), uterus (SRR1205176), kidney

(SRR1205998), lung (SRR1205218), heart (SRR1205223), brain (SRR1205222), adrenal glands

(SRR1205224), and seven tissues from a male koala, salivary gland (SRR1207973), kidney

(SRR1207974), testes (SRR1207975), bone marrow (SRR1106690), lymph node (SRR1106707),

liver (SRR1121764), and spleen (SRR1122141). To process the acquired data, we followed the

protocol outlined by Pertea et al. (Pertea, et al. 2016). Briefly, raw reads were filtered for low

quality and trimmed of adapter sequences by Trim Galore! (version 0.3.7) before alignment to the

reference koala genome (phaCin_unsw_v4.1) using HISAT2 (Kim, et al. 2015). We used the koala

GTF annotation from Ensembl (Phascolarctos_cinereus.phaCin_unsw_v4.1.97.gtf.gz release) to

assemble mapped reads into transcripts using StringTie 2.0 (Pertea, et al. 2016) with the -e-b--A

<gene_abund.tab> flags. We chose to use StringTie's functionality for an additional *de novo*

transcript assembly to quantify transcripts not currently annotated in the koala reference,

specifically those associated with the *lnc*RNA *Rsx*. First, a new GTF annotation was generated

including novel transcripts using the --merge flag and then the previously generated mapped reads

were reassembled into transcripts guided by the novel GTF file.

Analysis of sex-based differential gene expression

Gene expression for the previously annotated and novel transcripts was measured for all fifteen

samples available. Using the prepDE.py script from Stringtie 2.0 (Pertea, et al. 2016), we generated

a raw expression count matrix containing the data from all male and female samples. Subsequently,

we inputted the count matrix into DeSeq2 1.22.2 (Love, et al. 2014), which uses the median of

ratios method to normalize raw expression data for sequencing depths and RNA composition to

perform differential gene expression analysis. Considering all eight female and seven male

samples, we first sought to specifically test the male and female differential expression of the

annotated *Rsx* region from the novel transcript analysis. Only one 30.4 kb transcript, a novel

*lnc*RNA, overlapped with the annotated *Rsx* region (overlap > 90% of transcript) and was used to

evaluate gene expression. From DeSeq2's output, we collected data on the normalized male and

female expression counts, the log transformed fold change ($\log_2$[female expression/male

expression]), and the p-value associated with the sex-based differential expression. We also

performed a permutation of this analysis by only considering the tissues that had both a male and

female sample, namely the kidney, lung, and spleen.

NOISeq 2.26.1 (Tarazona, et al. 2015) was used for differential expression analysis due to

its ability to simulating technical replicates within given RNA-seq data sets when no replicates are

available. Using the raw count matrix from annotated transcripts, we generated sex-specific

differentially methylated genes for one dataset considering only the male and female kidney data

and another dataset considering the three tissues with male and female samples (kidney, lung, and

spleen). We compared the gene expression outputs for the three tissue datasets generated by both

DeSeq2 and NOISeq and found highly consistent results when considering the female over male

expression change values for all genes (Spearman's rank correlation coefficient, $\rho = 0.997$, $p < 2.2$

$\times 10^{-16}$).

Availability of data and materials

523   The raw and processed methylation datasets generated in this study have been deposited in the

524   NCBI Gene Expression Omnibus (Edgar, et al. 2002). They are accessible through GEO Series

525   accession number GSE149600 (reviewer's token: wdqhkaoyhxqdfoz) at

526   https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE149600.

Acknowledgements

This work was supported by a grant from the National Science Foundation (MCB 1818288) and a Pilot Grant from the Smooth Muscle Plasticity COBRE of the University of Nevada, Reno (funded by the National Institutes of Health grant 5P30GM110767-04) to DAP, grants by the National Science Foundation (MCB 1615664) and the National Institute of Health (R01MH103517) to SVY. In addition, DS was partially supported by the NIH Training Grant in Computational Biology and Biomedical Genomics (T32 GM105490).

References

Alvarez-Ponce D, Torres-Sanchez M, Feyertag F, Kulkarni A, Nappi T. 2018. Molecular evolution of DNMT1 in vertebrates: Duplications in marsupials followed by positive selection. PLOS ONE 13.

Black KH, Price GJ, Archer M, Hand SJ. 2014. Bearing up well? Understanding the past, present and future of Australia's koalas. Gondwana Research 25:1186-1201.

Brown CJ, Hendrich BD, Rupert JL, Lafreniere RG, Xing Y, Lawrence J, Willard HF. 1992. The human XIST gene: analysis of a 17 kb inactive X-specific RNA that contains conserved repeats and is highly localized within the nucleus. Cell 71:527-542.

Chen J, Bardes EE, Aronow BJ, Jegga AG. 2009. ToppGene Suite for gene list enrichment analysis and candidate gene prioritization. Nucleic Acids Res 37:W305-311.

Cotton AM, Price EM, Jones MJ, Balaton BP, Kobor MS, Brown CJ. 2015. Landscape of DNA methylation on the X chromosome reflects CpG density, functional chromatin state and X-chromosome inactivation. Hum Mol Genet 24:1528-1539.

Duret L, Chureau C, Samain S, Weissenbach J, Avner P. 2006. The Xist RNA gene evolved in eutherians by pseudogenization of a protein-coding gene. Science 312:1653-1655.

Edgar R, Domrachev M, Lash AE. 2002. Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. Nucleic Acids Res 30:207-210.

Elango N, Kim S-H, Program NCS, Vigoda E, Yi SV. 2008. Mutations of different molecular origins exhibit contrasting patterns of regional substitution rate variation. PLoS Computational Biology 4:e1000015.

Galili T. 2015. dendextend: an R package for visualizing, adjusting and comparing trees of hierarchical clustering. Bioinformatics (Oxford, England) 31:3718-3720.

Galupa R, Heard E. 2018. X-Chromosome Inactivation: A Crossroads Between Chromosome Architecture and Gene Regulation. Annual Review of Genetics 52:535-566.

Goll MG, Bestor TH. 2005. Eukaryotic Cytosine Methyltransferases. Annual Review of Biochemistry 74:481-514.

Grant J, Mahadevaiah SK, Khil P, Sangrithi MN, Royo H, Duckworth J, McCarrey JR, VandeBerg JL, Renfree MB, Taylor W, et al. 2012. Rsx is a metatherian RNA with Xist-like properties in X-chromosome inactivation. Nature 487:254.

Hansen KD, Langmead B, Irizarry RA. 2012. BSmooth: from whole genome bisulfite sequencing reads to differentially methylated regions. Genome biology 13:R83.

569   Heard E. 2005. Delving into the diversity of facultative heterochromatin: the epigenetics of the
570   inactive X chromosome. Curr Opin Genet Dev 15:482-489.
571   Heard E, Clerc P, Avner P. 1997. X-chromosome inactivation in mammals. Annu Rev Genet
572   31:571-610.
573   Hellman A, Chess A. 2007. Gene body-specific methylation on the active X chromosome. Science
574   315:1141-1143.
575   Hobbs M, Pavasovic A, King AG, Prentis PJ, Eldridge MDB, Chen Z, Colgan DJ, Polkinghorne
576   A, Wilkins MR, Flanagan C, et al. 2014. A transcriptome resource for the koala
577   (Phascolarctos cinereus): insights into koala retrovirus transcription and sequence
578   diversity. BMC Genomics 15:786.
579   Huynh KD, Lee JT. 2003. Inheritance of a pre-inactivated paternal X chromosome in early mouse
580   embryos. Nature 426:857-862.
581   Jjingo D, Conley AB, Yi SV, Lunyak VV, Jordan IK. 2012. On the presence and role of human
582   gene-body DNA methylation. Oncotarget 3:462.
583   Johnson RN, O'Meally D, Chen Z, Etherington GJ, Ho SYW, Nash WJ, Grueber CE, Cheng Y,
584   Whittington CM, Dennison S, et al. 2018. Adaptation and conservation insights from the
585   koala genome. Nature Genetics 50:1102-1111.
586   Jones PA. 2012. Functions of DNA methylation: islands, start sites, gene bodies and beyond. Nat
587   Rev Genet 13:484-492.
588   Kasai F, O'Brien PCM, Pereira JC, Ferguson-Smith MA. 2018. Marsupial chromosome DNA
589   content and genome size assessed from flow karyotypes: invariable low autosomal GC
590   content. Royal Society open science 5:171539-171539.
591   Keown CL, Berletch JB, Castanon R, Nery JR, Disteche CM, Ecker JR, Mukamel EA. 2017.
592   Allele-specific non-CG DNA methylation marks domains of active chromatin in female
593   mouse brain. Proceedings of the National Academy of Sciences 114:E2882.
594   Kim D, Langmead B, Salzberg SL. 2015. HISAT: a fast spliced aligner with low memory
595   requirements. Nature methods 12:357-360.
596   Koina E, Chaumeil J, Greaves IK, Tremethick DJ, Graves JA. 2009. Specific patterns of histone
597   marks accompany X chromosome inactivation in a marsupial. Chromosome Res 17:115-
598   126.
599   Krueger F, Andrews SR. 2011. Bismark: a flexible aligner and methylation caller for Bisulfite-Seq
600   applications. Bioinformatics (Oxford, England) 27:1571-1572.
601   Kundaje A, Meuleman W, Ernst J, Bilenky M, Yen A, Heravi-Moussavi A, Kheradpour P, Zhang
602   Z, Wang J, Ziller MJ, et al. 2015. Integrative analysis of 111 reference human epigenomes.
603   Nature 518:317-330.
604   Leonhardt H, Page AW, Weier H-U, Bestor TH. 1992. A targeting sequence directs DNA
605   methyltransferase to sites of DNA replication in mammalian nuclei. Cell:865-873.
606   Lister R, Pelizzola M, Dowen RH, Hawkins RD, Hon G, Tonti-Filippini J, Nery JR, Lee L, Ye Z,
607   Ngo Q-M. 2009. Human DNA methylomes at base resolution show widespread
608   epigenomic differences. Nature 462:315-322.
609   Loebel D, Johnston PG. 1996. Methylation analysis of a marsupial X-linked CpG island by
610   bisulfite genomic sequencing. Genome research 6:114-123.
611   Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-
612   seq data with DESeq2. Genome biology 15:550.
613   Lyon MF. 1961. Gene Action in the X-chromosome of the Mouse (Mus musculus L.). Nature
614   190:372-373.

Mendizabal I, Shi L, Keller TE, Konopka G, Preuss TM, Hsieh TF, Hu E, Zhang Z, Su B, Yi SV.
2016. Comparative Methylome Analyses Identify Epigenetic Regulatory Loci of Human
Brain Evolution. Mol Biol Evol 33:2947-2959.
Mendizabal I, Yi SV. 2015. Whole-genome bisulfite sequencing maps from multiple human
tissues reveal novel CpG islands associated with tissue-specific regulation. Human
Molecular Genetics 25:69-82.
Ng K, Pullirsch D, Leeb M, Wutz A. 2007. Xist and the order of silencing. EMBO reports 8:34-
39.
Ogawa Y, Kakumoto K, Yoshida T, Kuwako K-i, Miyazaki T, Yamaguchi J, Konno A, Hata J,
Uchiyama Y, Hirai H, et al. 2018. Elavl3 is essential for the maintenance of Purkinje
neuron axons. Scientific Reports 8:2722.
Okamoto I, Otte AP, Allis CD, Reinberg D, Heard E. 2004. Epigenetic dynamics of imprinted X
inactivation during early mouse development. Science 303:644-649.
Pertea M, Kim D, Pertea GM, Leek JT, Salzberg SL. 2016. Transcript-level expression analysis of
RNA-seq experiments with HISAT, StringTie and Ballgown. Nature Protocols 11:1650-
1667.
Peters TJ, Buckley MJ, Statham AL, Pidsley R, Samaras K, V Lord R, Clark SJ, Molloy PL. 2015.
De novo identification of differentially methylated regions in the human genome.
Epigenetics & Chromatin 8:6.
Piper AA, Bennett AM, Noyce L, Swanton MK, Cooper DW. 1993. Isolation of a clone partially
encoding hill kangaroo X-linked hypoxanthine phosphoribosyltransferase: Sex differences
in methylation in the body of the gene. Somatic Cell and Molecular Genetics 19:141-159.
Plath K, Mlynarczyk-Evans S, Nusinow DA, Panning B. 2002. Xist RNA and the mechanism of
X chromosome inactivation. Annu Rev Genet 36:233-278.
Price GJ. 2008. Is the modern koala (Phascolarctos cinereus) a derived dwarf of a Pleistocene
giant? Implications for testing megafauna extinction hypotheses. Quaternary Science
Reviews 27:2516-2521.
Rens W, Wallduck MS, Lovell FL, Ferguson-Smith MA, Ferguson-Smith AC. 2010. Epigenetic
modifications on X chromosomes in marsupial and monotreme mammals and implications
for evolution of dosage compensation. Proceedings of the National Academy of Sciences
107:17657.
Schultz MD, He Y, Whitaker JW, Hariharan M, Mukamel EA, Leung D, Rajagopal N, Nery JR,
Urich MA, Chen H, et al. 2015. Human body epigenome maps reveal noncanonical DNA
methylation variation. Nature 523:212-216.
Sharman GB. 1971. Late DNA Replication in the Paternally Derived X Chromosome of Female
Kangaroos. Nature 230:231-232.
Shevchenko AI, Zakharova IS, Zakian SM. 2013. The evolutionary pathway of x chromosome
inactivation in mammals. Acta naturae 5:40-53.
Song Q, Decato B, Hong EE, Zhou M, Fang F, Qu J, Garvin T, Kessler M, Zhou J, Smith AD.
2013. A Reference Methylome Database and Analysis Pipeline to Facilitate Integrative and
Comparative Epigenomics. PLOS ONE 8:e81148.
Spainhour JCG, Lim HS, Yi SV, Qiu P. 2019. Correlation Patterns Between DNA Methylation
and Gene Expression in The Cancer Genome Atlas. Cancer Informatics
18:1176935119828776.

Sprague D, Waters SA, Kirk JM, Wang JR, Samollow PB, Waters PD, Calabrese JM. 2019.
Nonlinear sequence similarity between the Xist and Rsx long noncoding RNAs suggests
shared functions of tandem repeat domains. RNA 25:1004-1019.
Sun D, Maney DL, Layman TS, Chatterjee P, Yi SV. 2019. Regional epigenetic differentiation of
the Z Chromosome between sexes in a female heterogametic system. Genome research
29:1673-1684.
Sun Z, Cunningham J, Slager S, Kocher J-P. 2015. Base resolution methylome profiling:
considerations in platform selection, data preprocessing and analysis. Epigenomics 7:813-
828.
Tarazona S, Furio-Tari P, Turra D, Pietro AD, Nueda MJ, Ferrer A, Conesa A. 2015. Data quality
aware analysis of differential expression in RNA-seq with NOISeq R/Bioc package.
Nucleic Acids Res 43:e140.
Wang X, Douglas KC, Vandeberg JL, Clark AG, Samollow PB. 2014. Chromosome-wide
profiling of X-chromosome inactivation and epigenetic states in fetal brain and placenta of
the opossum, Monodelphis domestica. Genome research 24:70-83.
Waters SA, Livernois AM, Patel H, O'Meally D, Craig JM, Marshall Graves JA, Suter CM, Waters
PD. 2018. Landscape of DNA Methylation on the Marsupial X. Molecular Biology and
Evolution 35:431-439.
Yi SV. 2012. Birds do it, bees do it, worms and ciliates do it too: DNA methylation from
unexpected corners of the tree of life. Genome biology 13:174.
Zemach A, McDaniel IE, Silva P, Zilberman D. 2010. Genome-wide evolutionary analysis of
eukaryotic DNA methylation. Science 328:916-919.
Zhang Z, Schwartz S, Wagner L, Miller W. 2000. A greedy algorithm for aligning DNA sequences.
J Comput Biol 7:203-214.

683
684

 Figure Legends

Figure 1. Overview of DNA methylation patterns across the koala genome. (A) Hierarchical

clustering of all 10 Whole Genome Bisulfite (WGBS) CpG reports representing five tissues using

Euclidean distances and Ward's agglomeration method. (B) Matrix reporting tissue-specific and

shared differentially methylation regions (DMRs) based on the pairwise comparison of DMRs

between tissues. (C) Considering data from brain, the total count of tissue specific DMRs falling

within one of three annotated genomic regions: promoters, gene bodies, and intergenic regions.

The enrichment of DMRs in each functional region is shown through a comparison with length

and GC matched control regions (***$p < 0.0001$, n.s. Not significant, from 10,000 bootstraps).

Error bars depict standard deviation. (D-E) Examples of two brain-specific DMRs (highlighted in

grey) with the corresponding CpG fractional methylation values reported for all WGBS samples

including the two brain (red) and eight remaining tissues (blue). Line smoothing performed using

local regression (LOESS). (D) *ELAVL3* is a neural specific RNA-binding protein highly expressed

in adult mammalian brains and linked to the maintenance of Purkinje neuron axons (Ogawa, et al.

2018). A 1.84 kb region containing three CpG Islands (CGI) and overlapping the first exon of

ELAVL3 was highly methylated (hypermethylated) in the brain samples compared to all other

tissues, and this gene was up-regulated in the brain compared to the kidneys (log-transformed brain

to kidney ratio = 15.4, probability of differential expression > 98% from NOISeq). (D) *ALDOC*

encodes a catalytic enzyme responsible for the conversion of fructose-1,6-bisphosphate to

glyceraldehyde 3-phosphate and dihydroxyacetone phosphate. A 945 bp, brain-specific lowly

methylated (hypomethylated) DMR overlaps the promoter and part of *ALDOC*'s gene body, and

this gene was up-regulated in koala brain samples compared to the kidney samples (log-

transformed brain to kidney ratio = 5.45, probability of differential expression > 96% from

NOISeq).

Figure 2. Global patterns of female and male DNA methylation (5mC) across eutherian and

marsupial X chromosomes. (A) Comparison of chromosome-wide sex-based DNA methylation

differences in human brain and koala brain samples. The distribution of differences in female and

male mean CpG fractional methylation is shown for autosomes and X chromosomes in human and

koala data (B) The combined distribution of female and male differences of mean CpG fractional

methylation from koala brain, lung, kidney, skeletal muscle and pancreas samples. For A-B, the

difference of female and male mean fractional methylation was calculated for 10 kb bins across
each autosome- or X-linked scaffold (*** indicated $p < 2.2 \times 10^{-16}$, Mann-Whitney U test). (C)
Mean female and male fractional DNA methylation across autosomes and X chromosomes from
human brain and koala brain samples. For D-F, mean fractional methylation levels across X-linked
or autosome-linked gene regions in male and female whole genome bisulfite (WGBS) samples
across genomic functional regions (promoters, exons, introns, and intergenic regions). Functional
regions were divided into 20 equally spaced bins and the mean fractional methylation of the
corresponding CpGs were calculated for all male (purple) and female (orange) samples. Line
smoothing was performed using local regression (LOESS). (D) Average fractional methylation of
CpGs in 100-bp sliding windows using a 10 bp step size in a 5 Kb region upstream and downstream
of all chromosome X linked gene's transcription start sites (TSSs) across all five koala tissues.

Figure 3. Annotation of genomic of DNA methylation (5mC) around *Rsx*. The genomic
coordinates are shown above the figure in grey isolating a 57 kb window containing the annotated
Rsx region (blue gene model below). The top panel identifies CpG islands (CGI), the middle panel
reports the absolute male (purple) and female (orange) fractional methylation at each CpG, and the
bottom panel show the female and male fractional methylation difference. Highlighted in grey
across all panels is the identifies female hypomethylated hotspot containing 101 CpGs and two
CGIs upstream of the Rsx transcription start site.

Figure 4. Model of DNA methylation (5mC) patterns across genomic functional regions for
eutherian mammals and marsupials. In summary, eutherian mammals exhibit increased DNA
methylation of promoter regions and CpG islands coupled with a relative depletion of DNA
methylation across gene bodies and intergenic regions on the inactive X chromosome (X_I)
compared to the active X chromosome (X_A) in females. Marsupial mammals share similar DNA
methylation depletion in gene bodies and intergenic regions of the inactive X chromosome;
however, they diverge from eutherian mammals in their promoter methylation patterns. Marsupial
promoters are modestly hypomethylated in the female X chromosomes (X_A and X_I) compared to
the male X chromosome (X_M).

Tables

Table 1. Functional annotation of enriched biological processes associated with gene sets containing tissue specific differentially methylated regions (DMRs).

Tissue	GO biological process term	Accession ID	p-value	q-value
Brain	central nervous system development	GO:0007417	$5.79{\times}10^{-13}$	$2.71{\times}10^{-09}$
Brain	generation of neurons	GO:0048699	$1.47{\times}10^{-12}$	$3.45{\times}10^{-09}$
Brain	head development	GO:0060322	$2.84{\times}10^{-12}$	$3.78{\times}10^{-09}$
Brain	Neurogenesis	GO:0022008	$3.22{\times}10^{-12}$	$3.78{\times}10^{-09}$
Brain	brain development	GO:0007420	$4.98{\times}10^{-12}$	$4.67{\times}10^{-09}$
Pancreas	response to endoplasmic reticulum stress	GO:0034976	$2.14{\times}10^{-11}$	$1.51{\times}10^{-07}$
Pancreas	oxoacid metabolic process	GO:0043436	$1.58{\times}10^{-10}$	$4.04{\times}10^{-07}$
Pancreas	organic acid metabolic process	GO:0006082	$1.72{\times}10^{-10}$	$4.04{\times}10^{-07}$
Pancreas	response to endogenous stimulus	GO:0009719	$6.34{\times}10^{-09}$	$1.12{\times}10^{-05}$
Pancreas	carboxylic acid metabolic process	GO:0019752	$2.29{\times}10^{-08}$	$3.24{\times}10^{-05}$
Skeletal Muscle	actin filament-based process	GO:0030029	$1.26{\times}10^{-09}$	$5.78{\times}10^{-06}$
Skeletal Muscle	actin cytoskeleton organization	GO:0030036	$5.04{\times}10^{-09}$	$1.16{\times}10^{-05}$
Skeletal Muscle	cytoskeleton organization	GO:0007010	$1.50{\times}10^{-08}$	$2.30{\times}10^{-05}$
Skeletal Muscle	cellular carbohydrate metabolic process	GO:0044262	$2.19{\times}10^{-08}$	$2.52{\times}10^{-05}$
Skeletal Muscle	embryonic morphogenesis	GO:0048598	$2.37{\times}10^{-07}$	$2.17{\times}10^{-04}$
Lung and Kidney	embryonic skeletal system morphogenesis	GO:0048704	$1.50{\times}10^{-10}$	$3.52{\times}10^{-07}$
Lung and Kidney	embryonic organ morphogenesis	GO:0048562	$1.61{\times}10^{-10}$	$3.52{\times}10^{-07}$
Lung and Kidney	embryonic skeletal system development	GO:0048706	$5.24{\times}10^{-10}$	$7.65{\times}10^{-07}$
Lung and Kidney	embryonic organ development	GO:0048568	$1.13{\times}10^{-08}$	$1.23{\times}10^{-05}$
Lung and Kidney	pattern specification process	GO:0007389	$3.31{\times}10^{-08}$	$2.90{\times}10^{-05}$

Gene ontology (GO) terms are presented for the top five most significantly enriched results of each tissue after correcting for multiple testing (FDR < 0.05). As the numbers of tissue specific DMRs for lung (n=22) and kidney (n=119) samples were so few, the corresponding gene sets were combined for this analysis.

782 **Table 2.** Correlation analysis of mean promoter and gene body DNA methylation and ranked gene expression.
783

Tissue	Genomic Region	Gene Count	Rho (p-value)
Brain	Promoter	5,396	-0.08 (p = 2.28×10^{-9})
	Gene body	5,443	-0.16 (p < 2.2×10^{-16})
Kidney	Promoter	9,268	-0.12 (p < 2.2×10^{-16})
	Gene body	9,379	-0.12 (p < 2.2×10^{-16})
Lung	Promoter	9,192	-0.13 (p < 2.2×10^{-16})
	Gene body	9,265	-0.19 (p < 2.2×10^{-16})

784
785 Spearman's rank correlation coefficients (ρ) and associated significances are reported for all tissues with both whole genome bisulfite
786 sequencing (WGBS) data and RNA-seq expression data.
787
788
789
790
791 **Table 3**. Sex-based differential expression of the *lnc*RNA *Rsx* utilizing different data subsets and expression quantification tools.
792

Expression Dataset	Tool	Female Count	Male Count	Significance
All Data (n = 15)	DeSeq2	6987.1	16	p-value = 0.05
Matched Data (n = 6)	DeSeq2	6837.6	0	p-value = 2.04×10^{-30}
Matched Data (n = 6)	NOISeq	7872.4	0.67	Probability = 99.99%
Kidney Data (n = 2)	NOISeq	4074.4	0.68	Probability = 99.99%

793
794 Normalized expression count values and significance of sex-based differential expression is shown for three data subsets using two
795 expression quantification tools. All data refers to the dataset considering all 15 RNA-seq samples (7 male and 8 female). Matched data
796 includes the tissues with both male and female RNA-seq samples (brain, kidney, and lung), and the kidney data is reported
797 independently. DeSeq2 reports significance as an associated p-value from the Wald test while NOISeq reports a probability of
798 differential expression threshold.
799
800
801
802

Figures

Figure 1.

820 **Figure 2.**

822
823
824
825

826   **Figure 3.**

827
828

Figure 4.

Supplementary Tables

Supplementary Table 1. Overview of whole genome bisulfite sequencing (WGBS) data for all 10 koala samples.

Sample	Australian Museum registration	Name	Tissue	Mapped Reads	De-duplicated Reads	Total CpGs	Coverage (%)	Mean Depth	% Reads Cov > 3×	Bisulfite Conversion Rate
WGM145_01_S1	M.45022	Pacific Chocolate	Brain	602055711	228244836	16761785	97.5	13.99	93.4	98.2
WGM145_04_S2	M.47723	Ben	Brain	677593548	233893124	16761785	97.6	14.61	93.8	98.0
WGM145_06_S3	M.45022	Pacific Chocolate	Kidney	559911229	213274030	16761785	97.2	13.62	92.5	98.7
WGM145_08_S4	M.47723	Ben	Kidney	589121434	202094771	16761785	97.2	12.93	92.3	98.7
WGM145_09_S5	M.45022	Pacific Chocolate	Lung	637878987	210002505	16761785	97.5	13.8	93.2	98.6
WGM145_12_S6	M.47723	Ben	Lung	663204935	198555030	16761785	97.3	12.54	92.4	98.6
WGM145_14_S7	M.45022	Pacific Chocolate	Skeletal Muscle	592423486	168450992	16761785	96.7	10.93	90.0	98.7
WGM145_16_S8	M.47723	Ben	Skeletal Muscle	605979530	168220022	16761785	96.8	11.12	90.4	98.6
WGM145_19_S9	M.45022	Pacific Chocolate	Pancreas	563288200	159866998	16761785	96.2	9.857	88.0	98.6
WGM145_20_S10	M.47723	Ben	Pancreas	598508860	166573663	16761785	96.6	10.56	89.4	98.7

Supplementary Table 2. GC-content for all scaffolds linked to annotated autosomes (chromosomes 1-7) and the X chromosome.

Chromosome	Size (Mb)	Total GC Content (%)
1	730.67	38.72
2	595.75	38.58
3	480.35	39.35
4	412.55	39.38
5	299.29	39.56
6	248.35	38.66
7	250.28	39.20
X	67.91	42.02

Supplementary Table 3. Enrichment and significance of all tissue specific DMRs compared to length and GC matched control regions.

Tissue	Genomic Region	DMR Counts (%)	Enrichment	p-value
Pancreas	Promoter	52 (4.5%)	1.13	0.74
	Gene body	612 (52.5 %)	1.44	**< 0.0001**
	Intergenic	502 (43.0 %)	-1.40	**< 0.0001**
Brain	Promoter	17 (3.5%)	-1.35	0.25
	Gene body	256 (53.0 %)	1.38	**< 0.0001**
	Intergenic	210 (43.5%)	-1.34	**< 0.0001**
Skeletal Muscle	Promoter	19 (4.9 %)	-1.23	0.32
	Gene body	202 (51.8 %)	1.27	**< 0.0001**
	Intergenic	169 (43.3 %)	-1.26	**0.001**
Kidney	Promoter	5 (4.0 %)	-1.34	0.40
	Gene body	62 (50.0 %)	1.25	**< 0.0001**
	Intergenic	57 (46.0%)	-1.21	**0.04**
Lung	Promoter	1 (4.3 %)	-1.61	0.51
	Gene body	12 (52.2%)	1.37	**< 0.0001**
	Intergenic	10 (43.5%)	-1.31	0.14

Reported are the total counts of tissue specific differentially methylated regions (DMRs) falling within one of three annotated genomic regions: promoters, gene bodies, and intergenic regions. The enrichment of DMRs in each functional region is shown through a fold change comparison with a control dataset generated from 10,000 bootstraps using length and GC matched control regions. All significant p-values ($p < 0.05$) are highlighted in bold.

Supplementary Table 4. Mean and median sex-based DNA methylation difference calculated for all candidate X-scaffolds (n=98) by tissue.

Tissue	Median Male-Female 5mC	Mean Male-Female 5mC
Pancreas	-0.2282 ± 0.15	-0.2066 ± 0.12
Brain	-0.3273 ± 0.14	-0.2911 ± 0.12
Skeletal Muscle	-0.2748 ± 0.15	-0.2431 ± 0.12
Kidney	-0.2850 ± 0.15	-0.2567 ± 0.12
Lung	-0.2706 ± 0.14	-0.2437 ± 0.11
Combined Tissues	-0.2773 ± 0.15	-0.2484 ± 0.12

The female and male mean fractional DNA methylation (methylated reads/total reads per CpG) was calculated for all CpGs within 10 kb bins across candidate scaffold.

Supplementary Figures

Supplementary Figure 1. Enrichment of tissue specific differentially methylated regions (DMRs) falling within genomic functional regions.

Data shown for (A) pancreas, (B) skeletal muscle, (C) kidney, and (D) lung. The enrichment of DMRs in each functional region (promoter, gene body, and intergenic regions) is shown through a comparison with length and GC matched control regions (*** indicates $p < 0.0001$, ** indicates $p < 0.001$, * indicates $p < 0.05$, and non-significance is shown by n.s. based on 10,000 bootstraps). Error bars indicate standard deviation.

Supplementary Figure 2. Correlation of gene expression and DNA methylation (5mC) in CpGs across promoters and gene bodies.

Three tissues with both whole genome bisulfite sequencing (WGBS) DNA methylation data and RNA-seq gene expression data are shown, (A) brain (n = 5,396 promoters and n = 5,443 gene bodies), (B) kidney (n = 9,268 promoters and n = 9,379 gene bodies), and (C) lung (n = 9,192 promoters and n = 9,265 gene bodies). For A-C, TPM expression values were ranked from low to high for each gene and correlated with mean fractional DNA methylation (methylated reads/total reads per CpG site). Spearman's rank correlation coefficients and the associated p-values are reported.

Supplementary Figure 3. Correlation of tissue dependent DNA methylation (5mC) and gene expression from brain and kidney samples.

(A) The mean brain and kidney DNA methylation difference calculated for all CpGs across each gene promoter matched with the corresponding log-transformed ratio of brain to kidney expression. (B) The mean brain and kidney DNA methylation difference calculated for all CpGs across each gene body and matched with corresponding log-transformed ratio of brain to kidney expression. For A and B, Spearman's rank correlation coefficient and the associated p-value is reported. Blue dots indicate genes that are significantly differentially express between brain and kidney samples (probability of differential expression > 95% based on NOISeq) and red dots show all genes that are significantly similarly methylated in brain and kidney samples (probability of differential expression < 5% based on NOISeq).

Supplementary Figure 4. Sex-specific CpG depth of coverage and read mapping to autosomes and X chromosomes.

(A) For each of the five tissues, box-and-whisker plots of CpG depths across the X chromosome in male (purple) and female (orange) samples (*** indicates p < 2.2x10-16, Mann-Whitney U test). Histogram and distribution of sex-based read mapping per sample (n=10) to (B) X-linked scaffolds, (C) candidate X-linked scaffolds, and (D) a subset of autosome-linked scaffolds matched in length with all known X-linked scaffolds. For A-C, the percent of reads mapping to the scaffold category of interested over the total number of mapped reads in the genome was calculated for all male (n=5) and female samples (n=5). The known X-linked and candidate X-linked scaffolds show a bimodal distribution with an increase of read mapping to female samples expected from the 2:1 ratio X chromosomes in females to males. This bimodality is not observed in autosomes.

Supplementary Figure 5. The distribution of sex-based CpG fractional DNA methylation (5mC) differences across autosomes and X chromosomes.

The distribution of female and male mean fractional methylation difference from (A) brain, (B) lung, (C) kidney, (D) skeletal muscle and (E) pancreas samples across autosomes and X chromosomes. For A-E, the female and male mean fractional methylation (methylated reads/total reads per CpG) was calculated for all CpGs within 10 kb bins across each autosome- or X-linked scaffold. All tissues exhibited a significant shift towards female hypomethylation in the X chromosome compared to the autosome (*** indicates p < 2.2x10-16, Welch's t-t

Supplementary Figure 6. Female and male gene expression across autosomes and the X chromosome using kidney RNA-seq data.

(A) The relative genomic location of all genes linked to X chromosome scaffolds aligned by scaffold length and the log-transformed female to male expression ratio generated by NOISeq. Orange dots indicate the 36 genes with significant female-biased expression and purple dots indicate the 11 genes with significant male-biased expression (probability of differential expression > 95% based on NOISeq). For all autosome-linked genes (n = 10,414) and chromosome X-linked genes (n= 209), a box-and-whisker plot (B) and density distribution (C) of the log-transformed female to male expression ratio (p = 0.14, Mann-Whitney U test).

Supplementary Figure 7. Correlation analysis of sex-based DNA methylation (5mC) and gene expression across chromosome X using kidney WGBS and RNA-seq data.

B Promoter

	Male Over Expressed / Total Genes	Female Over Expressed / Total Genes
Female Hypermethylated	$\frac{5}{32} = 0.16$	$\frac{14}{44} = 0.32$
Female Hypomethylated	$\frac{6}{57} = 0.11$	$\frac{22}{73} = 0.30$

D Gene Body

	Male Over Expressed / Total Genes	Female Over Expressed / Total Genes
Female Hypermethylated	$\frac{3}{11} = 0.27$	$\frac{7}{15} = 0.47$
Female Hypomethylated	$\frac{8}{79} = 0.10$	$\frac{29}{103} = 0.28$

(A) The mean female and male fractional methylation difference calculated for all CpGs across each gene promoter matched with the corresponding log-transformed ratio of female to male expression. (B) The ratio of the number female hypermethylated and female hypomethylated gene promoters that show either significant male of female biased expression over the total number of genes in each category. (C) The mean female and male fractional methylation difference calculated for all CpGs across each gene body and matched with corresponding log-transformed ratio of female to male expression. (D) The ratio of the number female hypermethylated and female hypomethylated gene bodies that show either significant male of female biased expression over the total number of genes in each category. For A and C, Spearman's rank correlation coefficient and the associated p-value is reported. The Rsx gene was excluded from the correlation calculation. Orange dots indicate the 36 genes with significant female-biased expression and purple dots indicate the 11 genes with significant male-biased expression (probability of differential expression > 95% based on NOISeq).

Supplementary Figure 7. Sex-based DNA methylation (5mC) by GC-content across autosomes and the X chromosome.

For A and B, the mean female and male methylation difference calculated from CpGs in 1 Kb bins across (A) autosomes and (B) X chromosomes. For C and D, mean female and male methylation difference calculated from CpGs located in promoter regions (defined as regions 1 kb upstream of known gene TSSs) in (C) autosomes and (B) X chromosomes. For A-D, data from all five tissues (brain, kidney, lung, pancreas, and skeletal muscle) are reported. All plots are coloured by data density where blue represents low density regions and yellow represents high density regions.