# Natural enemies have inconsistent impacts on the coexistence of competing species

J. Christopher D. Terry*† [1] 0000-0002-0626-9938 c.terry@qmul.ac.uk

J. Chen* [2] 0000-0002-8435-0897 jinlin.chen@zoo.ox.ac.uk

O. T. Lewis [2] 0000-0001-7935-6111 owen.lewis@zoo.ox.ac.uk

*: Joint first authorship. †Corresponding Author
1. School of Biological and Chemical Sciences, Queen Mary University of London
2. Department of Zoology, University of Oxford

## Key Words:

*Modern coexistence theory, competition, natural enemies, Bayesian, model fitting, Drosophila, parasitoid*

## Author Contributions

JCDT conceptualised the experiment, led the analysis and wrote the initial draft. JC led the design and execution of the experimental work. Experimental work was conducted by JC and JCDT. All authors contributed to the development of ideas and writing of the manuscript.

Funding: The work was supported by NERC grant NE/N010221/1 to OTL and a grant from the China Scholarship Council to JC. JCDT was also funded through NERC grant NE/T003510/1

## Data Accessibility:

All raw observation data and code to replicate the analyses is available at https://github.com/jcdterry/TCL_DrosMCT. If accepted for publication, a copy will also be archived at Zenodo.

# Abstract:

1. The role of natural enemies in promoting coexistence of competing species has generated substantial debate. Modern coexistence theory provides a detailed framework to investigate this topic, but there have been remarkably few empirical applications to the impact of natural enemies.

2. We tested experimentally the capacity for a generalist enemy to promote coexistence of competing insect species, and the extent to which any impact can be predicted by trade-offs between reproductive rate and susceptibility to natural enemies.

3. We used experimental mesocosms to conduct a fully-factorial pairwise competition experiment for six rainforest *Drosophila* species, with and without a generalist pupal parasitoid. We then parameterised models of competition and examined the coexistence of each pair of *Drosophila* species within the framework of modern coexistence theory.

4. We found idiosyncratic impacts of parasitism on pairwise coexistence, mediated through changes in fitness differences, not niche differences. There was no evidence of an overall reproductive rate – susceptibility trade-off. Pairwise reproductive rate – susceptibility relationships were not useful shortcuts for predicting the impact of parasitism on coexistence.

5. Our results exemplify the value of modern coexistence theory in multi-trophic contexts and the importance of contextualising the impact of natural enemies. In the set of species investigated, competition was affected by the higher trophic level, but the overall impact on coexistence cannot be easily predicted just from knowledge of relative susceptibility. Methodologically, our Bayesian approach highlights issues with the separability of model parameters within modern coexistence theory and shows how using the full posterior parameter distribution improves inferences. This method should be widely applicable for understanding species coexistence in a range of systems.

# Introduction

Species compete for limited resources and (in almost all cases) are consumed by other species. Understanding how the impacts of competitors and natural enemies interact to influence species coexistence is a long-standing challenge. Natural enemies are regularly cited as a driver of coexistence, for example by suppressing species that would otherwise be competitively dominant (Paine, 1966), but their impact can be highly variable (Chase et al., 2002).

Analyses investigating the effect of natural enemies often focus on 'reduction in competition' (e.g. (Gurevitch et al., 2000), as measured by negative effects between focal species; however consumers can affect competition and coexistence in convoluted ways (Chase et al., 2002; Pringle et al., 2019). Modern coexistence theory (MCT) focuses on the property of mutual invasibility (Grainger, Levine, et al., 2019), and allows the precise framing of questions regarding pairwise species coexistence as a balance between niche differences promoting stabilisation and fitness differences fostering exclusion of one species by another (Chesson, 2000; Letten et al., 2017). Through the lens of modern coexistence theory, interactions mediated through competition for resources and through natural enemies are conceptually equivalent (Chesson, 2018; Chesson & Kuang, 2008) and a body of theory has developed to integrate consumer effects within the coexistence framework (Chesson & Kuang, 2008; McPeek, 2019). This theoretical work has emphasised that impacts of any process on coexistence can only be determined with information on both intrinsic growth rates and interaction

terms. Alongside the theory, empirical studies using the modern coexistence framework to
investigate the impact of consumers on coexistence is now emerging (Bartomeus & Godoy, 2018).

Natural enemies can affect both fitness differences and niche differences between species. A
frequently-invoked mechanism is that they reduce fitness differences, mediated by a trade-off
between population growth rate and susceptibility to attack (Kraaijeveld & Godfray, 1997; Paine,
1966). However, such a trade-off is in itself insufficient to enable long-term coexistence, since some
kind of stabilising niche-difference is required to mitigate any fitness difference (Chesson, 2000).
Even with zero fitness difference, stochastic drift would preclude coexistence in the absence of
stabilising mechanisms (Adler et al., 2007). Natural enemies can also potentially influence niche
differences through modifying the overall competitive pressure exerted between competitors.
Enemies may cause competitors to change their behaviour, or natural enemies may change their
foraging behaviour as the relative abundances of the competing species change (Abrams, 2010).

Most experimental work within a formal MCT framework has been conducted on plant or microbial
systems (Bartomeus & Godoy, 2018; Grainger, Letten, et al., 2019) including investigation of the
effects of seed predation (Nottebrock et al., 2017; Petry et al., 2018), and seed pathogens
(Mordecai, 2013). Experimental studies of insect communities have considerable potential to further
our understanding of the role of natural enemies in species coexistence. Since parasitoids kill their
insect hosts as they develop in or on them, each attack is directly interpretable in terms of host
fitness (Hassell, 2000). Although ecologists have been competing insect species (especially
*Drosophila*) against each other for decades (Davis et al., 1998; Gilpin et al., 1986; Pearl, 1932;
Worthen, 1989), very few studies have explicitly tested for mutual-invasibility in insect systems
(Siepielski et al., 2018), rather than particular conditions necessary for coexistence (Godwin et al.,
2020; Spaak & de Laender, 2020).

Here we present an analysis parameterising the impact of consumers on pairwise species
coexistence, focusing on a community of six rainforest *Drosophila* species. Our experiment was
designed specifically to capture the key parameters of our focal species and their competitive
interactions. We examine how niche and fitness differences are influenced by the presence of the
parasitoid. We use a Bayesian approach to propagate uncertainty in model parameters through to
niche and fitness differences, a method that should be widely applicable for understanding species
coexistence processes in a range of systems. We show that parasitoids strongly influence the fitness
differences between species and have inconsistent impacts on pairwise coexistence that would not
be identifiable solely from reproductive rate – susceptibility relationships.

# Methods

## Study Species

We investigated a system of six naturally co-occurring *Drosophila* vinegar flies (*D. birchii* (BIR), *D.*
*pallidifrons* (PAL), *D. pandora* (PAN), *D. pseudoananassae* (PSA), *D. simulans* (SIM), *and D.*
*sulfurigaster* (SUL)) from a well-characterised Australian rainforest community (Hangartner et al.,
2015; Jeffs et al., 2020; O'Brien et al., 2017), along with a co-occurring generalist parasitoid wasp
from the genus *Trichopria* (Hymenoptera: Diapriidae, lab strain 66LD, SI 1). This parasitoid attacks
*Drosophila* species at their immobilized pupal stage. Successful development of a parasitoid results
in the death of its host. *Drosophila* can mount a variety of physiological defences to parasitoid
attack, which can be costly (Fellowes & Godfray, 2000).

*Drosophila* species have long been used in experimental ecology since they are straightforward to
culture under laboratory conditions, allowing high levels of replication and standardisation. The set
of *Drosophila* species used here represents a significant fraction of the diversity of *Drosophila*
species co-occurring at rainforest sites in northern Queensland, Australia (Jeffs *et al.* 2020). The six
species span a range of body sizes (0.94 mg – 2.64 mg, mean wet mass of female adults) and
phylogenetic divergences (SI 1, Table S1.1), have similar generation times, and can be distinguished
visually (at least in the case of males). *Drosophila* are generalist feeders (Valadão et al., 2019) and in
natural populations they experience high levels of parasitism, including from a suite of highly
generalist natural enemies (Jeffs et al., 2020).

## Experimental design

We conducted a large, single-generation pairwise competition assay using a response-surface design
following the guidelines of Hart *et al.* (2018), and in the presence or absence of pupal parasitoids.
We used 25mm-diameter standard *Drosophila* vials (VWR International) containing 4 ml of
cornflour-yeast-agar medium. This creates conditions where *Drosophila* compete for space to lay
eggs, for larval food, and for pupation sites. Equal numbers of male and female *Drosophila* were
introduced to each vial, so we refer to the number of founder pairs. Each two-species combination
(15 combinations in total) was tested at different founding densities in a factorial design: (3 pairs of
species A, 3 pairs of species B), (6A,3B), (9A,3B), (12A,3B), (3A,6B), (3A,9B) and (3A, 12B), with each
combination replicated three times in each parasitism treatment. We also included monocultures of
each species, with 12 replicates of a single founding pair, 6 replicates of 3, 9 or 12 pairs within each
treatment, making a total of 990 experimental vials (360 single-species, 630 two-species). The entire
experiment was split over three blocks, each with identical sets of combinations, but staggered by
two days. The experiment was conducted in a single large incubator maintained at 26°C, with the
location of trays within the incubator regularly rotated. Further details are given in SI 1.

## Initiation of treatments

Founding adults were offspring of adults drawn from mass-bred line cages and were raised in vials at
moderate density (approximately 100 adults per vial) at a constant 23°C. These adults emerged 8-10
days before the start of the experiment, and so could be assumed to be sexually mature and mated.
On the day preceding the start of the experiment the founding flies were lightly anesthetised under
$CO_2$ to be separated by sex. Males were added to the experimental vials immediately. Females were
maintained at moderate density (~50 adults per vial) in holding vials with fly medium and were
transferred under light $CO_2$ anaesthesia to experimental vials the following day, minimising the time
interval between the addition of each species in multi-species assays. They were able to lay eggs and
re-mate for 24 hours before being removed. Fig. 1 provides an overview of the experimental
protocol.

**Figure 1** Summary of the experimental regime showing the founding process, treatment and data collection procedures described in the main text.

## Parasitism

At the onset of pupation, 5 days after the founder flies were removed, the fly vials were moved into Perspex parasitism boxes (35x20x14cm) containing up to 88 vials (SI 1). Vial positions within each box were randomised. The vial plugs were removed, and 50 female and 50 male mature and mated parasitoid wasps were added. This ratio of parasitoids to *Drosophila* was selected based on prior experience to result in a parasitism rate in line with field-parasitism levels of around 15% or greater (Jeffs et al 2020). The wasps were free to move between vials within each box. While it is possible that late-pupating *Drosophila* larvae could cross from one vial to another during this period, we only observed a handful of cases where 1-2 individuals of the 'wrong' species emerged from a vial, and very few pupae (<5 per box) were developed on the outside of the vials, suggesting that movement was negligible. After 30 hours, before the emergence of the fastest-developing *Drosophila* species, all wasps were removed with an aspirator, vial plugs were replaced, and the vials were returned to their original trays.

## Counting

We removed adult flies emerging from all vials every three days, minimising the risk that emerged adults started a subsequent generation. Flies in single-species vials were sexed and counted at each removal time, while flies in multi-species vials were sexed and identified at the end of the experiment (Fig. 1). There was a highly consistent 1:1 sex ratio across the single-species trials and in those 2-species trials where females of different species could be reliably distinguished (SI 2 Fig. S2.1). In the five of the combinations (BIR-PAN, BIR-PSA, BIR-SIM, PSA-PAN and PAL-SUL) where it was not possible to identify female flies to species we counted the males and inferred the total number of flies of each species assuming a 1:1 sex ratio.

## Modelling

### Determining growth rates

The larval competition of *Drosophila* often occurs within individual fruits that represent a discrete and temporary resource suitable for only a single generation of flies (Atkinson & Shorrocks, 1981; Shorrocks, 1991). We take as our underlying model of growth rates the Beverton-Holt model, without any carry-over between generations:

$$N_{i,t+1} = \frac{R_{0i} N_{i,t}}{1 + \alpha_{ii} N_{i,t} + \alpha_{ij} N_{j,t}} \quad (1)$$

where $N_{i,t}$ and $N_{i,t+1}$ are the size of the population of species $i$ in the founding and emerging
generations, respectively. $R_{0i}$ is a species-specific generational reproduction term, $\alpha_{ii}$ represents
the intra-specific competitive effect of species $i$ and $\alpha_{ij}$ are competition coefficients terms
representing the effect of species $j$ on $i$. It is important to emphasise that these $\alpha$ terms capture *all*
impacts between species, including those through indirect means. The same function has been used
for annual plants that have been the focus of most recent coexistence work (Bartomeus & Godoy,
2018; Levine & HilleRisLambers, 2009; Pérez-Ramos et al., 2019).

The mean number of adults emerging per founder was consistently lower in vials with a solitary
founding pair compared to those with three founders (SI 2, Fig. S2.2). *Drosophila* use cues from
other females when determining oviposition sites (Duménil et al., 2016; Wertheim et al., 2002),
which could be causing this Allee effect. It is not possible to incorporate such positive density-
dependence effects directly within the standard MCT framework without additional assumptions
(Barabás et al., 2018; Schreiber et al., 2019). Since solitary founders are quite possibly a rather
artificial scenario, we excluded data from all singleton pair trials from the model fitting.

## Inferring model parameters

The models were fit within a Bayesian framework and posterior distributions of the parameters were
obtained from Hamiltonian Monte-Carlo sampling using STAN (Carpenter et al., 2017). We assumed
a negative binomial error distribution and fitted separate overdispersion terms $\phi$ for each parasitism
treatment. Since we counted both species, each two-species experiment contributed two data
points. Although this contributes a small component of non-independence, the gain in total sample
size is significant. Since we excluded the single-founder vials, our effective total 'n' across both
treatments was 1476, to estimate a maximum of 86 free parameters in our most complex model
(including overdispersion terms).

The competition $\alpha$ terms were each fit with an underlying Gaussian prior (mean=0, $\sigma$ =1) and
constrained to be positive, implying only competitive interspecific interactions, with no facilitation
possible. Growth rate terms $R_0$ were constrained to be positive, and fit with a weak Gaussian prior
(mean=10, $\sigma$ =10). Tests fitting $\alpha$ values with different constraints and priors did not give notably
different results (SI 3).

We fitted and compared a sequence of candidate models of increasing simplicity. In the first,
separate $R_0$ and $\alpha$ terms were fitted for each treatment. In the second, a single set of $\alpha$ terms, but
separate $R_0$ terms were fitted across both treatments (i.e. parasitism affects emergence rate, but
not competition coefficients). In the third model a single set of parameters were fitted across both
treatments (i.e., parasitism has no effect). The three models were compared based on expected log
pointwise predictive density (ELPD, a measure of out-of-sample predictive capacity), computed using
the *loo* R package (Vehtari et al., 2017). This approach allows an assessment of whether there is
sufficient evidence in the data to separate the parameters assigned to the treatment groups in a
comparable manner to other information criteria such as AIC (Burnham & Anderson, 2002). All STAN
model code and data to fit the models are available online. Posterior predictive checks and
diagnostics are presented in SI 4.

Pairwise coexistence criteria

The invasibility criterion for species coexistence requires each species to have a positive long-term
low-density growth rate in the presence of the rest of the community (Chesson 2000). Following
Godoy & Levine (2014), coexistence of a pair of species is determined to be possible, following
Beverton-Holt model dynamics, when:

$$\sqrt{\frac{\alpha_{ij}\alpha_{ji}}{\alpha_{jj}\alpha_{ii}}} \; < \; \frac{R_{0i}-1}{R_{0j}-1}\sqrt{\frac{\alpha_{ji}\alpha_{jj}}{\alpha_{ij}\alpha_{ii}}} \; < \; \sqrt{\frac{\alpha_{jj}\alpha_{ii}}{\alpha_{ij}\alpha_{ji}}}. \quad (2)$$

221 Equation (2) expresses the coexistence criterion as a balance between niche overlap ($\rho$) and fitness
222 differences ($\frac{\kappa_i}{\kappa_j}$) where:

$$\rho = \sqrt{\frac{\alpha_{ij}\alpha_{ji}}{\alpha_{jj}\alpha_{ii}}} \quad (3)$$

223 and

$$\frac{\kappa_i}{\kappa_j} = \frac{R_{0i}-1}{R_{0j}-1}\sqrt{\frac{\alpha_{ji}\alpha_{jj}}{\alpha_{ij}\alpha_{ii}}} \quad (4)$$

Following Chesson (2000) and Godoy & Levine (2014), we define the niche difference between two
species as $1 - \rho$. Certain $\alpha$ values can result in a niche overlap $\rho > 1$, implying a 'negative niche
difference'. This does not have a clear ecological meaning, and some previous authors have elected
to define such cases as 0. However, since these negative $1 - \rho$ values can be ecologically interpreted
as potentially identifying priority effect situations (Ke & Letten, 2018), we present the values as
calculated. For both negative and zero values, coexistence is not expected under the MCT
framework.

The *Drosophila* species we examined vary in their time to maturation. However, it is important to
note that the mathematical expression of the invasibility criteria in a constant environment,
equation (2), is not dependent on generation time. That is to say, the sole criterion for the invader
species to expand its population while the resident is at an equilibrium is that its population size in
its next generation will be larger, regardless of how long a generation takes.

For each pair of species, we calculated the niche and fitness differences across 3000 draws from the
posterior distributions of parameter estimates. We partitioned the posterior to infer the support
given by the data to each of the four outcomes of competition: coexistence, exclusion of either
species or a situation of priority effects (see Fig. 3a). Note that the dynamics described by our
models are fundamentally deterministic – these values express uncertainty in outcome given the
data, rather than necessarily stochastic outcomes.

# Results

Based on model comparison, there was strong support for the expectation that parasitism reduces
the growth rate of each species, but little support for differences in competition coefficients
between parasitism and no-parasitism treatments. Our second model (with different $r$ values for

parasitism and no-parasitism treatments but common α values) performed best (ΔELPD vs Model 1 =
-43.02, vs Model 3= -34.03, SI4 Table S4.1).

The impact of the parasitoids on coexistence was inconsistent across species pairs (Fig. 2c). In three
pairs of species the fitness difference was so large there was essentially no support for coexistence
under either treatment. Of the remaining twelve pairs, five had higher support for coexistence under
the parasitism treatment, while seven had less support (Table 1). In only two pairs (*D. sulfurigaster*
and *D. pallidifrons, D. simulans* and *D. pallidifrons*) did the parasitism treatment cause the median
estimate of fitness difference to cross the boundary delineating coexistence from competitive
exclusion. In both cases the parasitism was associated with a moderate reduction in support for
coexistence (Table 1). In the pair showing the largest change in support for coexistence between
parasitism and no-parasitism treatments (+ 32 percentage points, *D. sulfurigaster* and *D. birchii,* Fig
2b), the median value did not cross the boundary. Strong positive correlations between fitted
parameters, led by relationships between $R_{0i}$ and $\alpha_{ii}$ terms, resulted in non-independence of
estimates of the fitness and niche differences and non-eliptical posterior distributions (Fig. 2b,c).

**Figure 2.** The likelihood of coexistence of *Drosophila* species pairs responds idiosyncratically to parasitism. a) Guide to interpretation of coexistence plane diagrams. The region of coexistence (dark grey, bordered by red line) defines the region where either species could invade from rare. The pale grey area denotes the region of 'priority effects', where neither species could invade the other. The white area is where fitness difference is large enough that the dominant species can competitively exclude the other species despite their niche difference. b) Full posterior distribution of niche and fitness differences of one example species pair (*D. sulfurigaster* and *D. birchii*) using the best-fitting model. Note the irregular, non-circular, shape ultimately derived from correlations between fitted vales in the posterior distribution and alpha terms contributing to both niche and fitness differences. c) Equivalent contour plots for all 15 species combinations. The median of the posterior distribution of parameters are shown with diamonds. Table 1 details the support assigned to each region for each pair under each treatment. Note that because the best model did not include varying competition coefficients ( ) between treatments, only the fitness difference (y-axis) shifts between treatments.

| Species $i$ | Species $j$ | Parasitoids Present | Median Niche Difference $(1-\rho)$ | Median Fitness Difference $\left(\frac{\kappa_i}{\kappa_j}\right)$ | Coexistence under Median Estimate | Percentage posterior support for regions of coexistence plane | | | | Change in support for coexistence |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | | | | Species $i$ Wins | Species $j$ Wins | Coexistence | Priority Effect | |
| pallidifrons | birchii | No | 0.03 | 3.64 | | 99.93 | 0.00 | 0.00 | 0.07 | |
| | | Yes | 0.03 | 3.07 | | 99.73 | 0.00 | 0.00 | 0.27 | 0 |
| pandora | birchii | No | 0.49 | 6.27 | | 99.80 | 0.00 | 0.18 | 0.02 | |
| | | Yes | 0.49 | 4.75 | | 99.23 | 0.00 | 0.70 | 0.07 | 0.5 |
| pandora | pallidifrons | No | 0.36 | 0.84 | TRUE | 9.32 | 20.97 | 67.68 | 2.03 | |
| | | Yes | 0.36 | 0.76 | TRUE | 5.13 | 30.18 | 62.62 | 2.07 | -5.1 |
| pseudoananassae | birchii | No | 0.70 | 1.17 | TRUE | 3.03 | 0.60 | 96.37 | 0.00 | |
| | | Yes | 0.70 | 1.00 | TRUE | 1.35 | 1.32 | 97.33 | 0.00 | 1 |
| pseudoananassae | pallidifrons | No | 0.21 | 0.25 | | 0.00 | 99.75 | 0.03 | 0.22 | |
| | | Yes | 0.21 | 0.26 | | 0.00 | 99.78 | 0.05 | 0.17 | 0 |
| pseudoananassae | pandora | No | 0.02 | 0.70 | | 6.93 | 61.57 | 20.15 | 11.35 | |
| | | Yes | 0.02 | 0.79 | | 10.92 | 49.73 | 23.58 | 15.77 | 3.4 |
| simulans | birchii | No | -0.18 | 3.77 | | 99.82 | 0.00 | 0.00 | 0.18 | |
| | | Yes | -0.18 | 3.79 | | 99.88 | 0.00 | 0.00 | 0.12 | 0 |
| simulans | pallidifrons | No | 0.50 | 1.74 | TRUE | 33.28 | 0.00 | 66.72 | 0.00 | |
| | | Yes | 0.50 | 2.07 | | 53.97 | 0.00 | 46.03 | 0.00 | -20.7 |
| simulans | pandora | No | 0.64 | 0.96 | TRUE | 2.25 | 3.30 | 94.45 | 0.00 | |
| | | Yes | 0.64 | 1.28 | TRUE | 8.35 | 0.87 | 90.73 | 0.05 | -3.7 |
| simulans | pseudoananassae | No | 0.07 | 2.25 | | 95.67 | 0.13 | 2.62 | 1.58 | |
| | | Yes | 0.07 | 2.63 | | 98.20 | 0.07 | 1.05 | 0.68 | -1.6 |
| sulfurigaster | birchii | No | 0.33 | 2.48 | | 95.47 | 0.00 | 4.38 | 0.15 | |
| | | Yes | 0.33 | 1.63 | | 61.45 | 0.73 | 36.42 | 1.40 | 32 |
| sulfurigaster | pallidifrons | No | 0.32 | 0.81 | TRUE | 1.37 | 28.87 | 69.20 | 0.57 | |
| | | Yes | 0.32 | 0.63 | | 0.10 | 60.47 | 39.43 | 0.00 | -29.8 |
| sulfurigaster | pandora | No | 0.44 | 1.04 | TRUE | 6.57 | 11.27 | 81.52 | 0.65 | |
| | | Yes | 0.44 | 0.90 | TRUE | 2.30 | 17.48 | 79.92 | 0.30 | -1.6 |
| sulfurigaster | pseudoananassae | No | 0.25 | 3.40 | | 99.80 | 0.00 | 0.03 | 0.17 | |
| | | Yes | 0.25 | 2.61 | | 98.58 | 0.00 | 0.48 | 0.93 | 0.4 |
| sulfurigaster | simulans | No | 0.61 | 1.17 | TRUE | 0.37 | 0.62 | 99.02 | 0.00 | |
| | | Yes | 0.61 | 0.77 | TRUE | 0.00 | 7.83 | 92.17 | 0.00 | -6.9 |

**Table 1.** Summary of the posterior distribution across the coexistence plane between the two treatments for each species pair. The percentage of posterior draws that fall into the four major regions of the coexistence plane can provide a summary of the confidence in the outcome of a particular competition, given the data.

The parasitism treatment reduced population growth rates, but we did not identify a reproduction
rate – susceptibility trade-off across our six species (Fig. 3). The pairwise relationship between
reproductive rate and susceptibility to parasitism (proportional reduction in $R_0$) did not provide a
useful heuristic in identifying the impact of parasitism on reducing fitness differences. Positive and
negative reproduction rate – susceptibility relationships were both associated with increases and
decreases in fitness differences.

**Figure 3** No clear reproductive rate – susceptibility trade-off across the six *Drosophila* species. Median value
for each species (large black points) are plotted alongside a posterior distribution point cloud and fitted ellipse
(60%, fitted assuming a multivariate-t distribution) to show the distribution of uncertainty for each species.

# Discussion

Our results demonstrate an inconsistent impact of natural enemies on the pairwise coexistence of a
guild of competing species. *Drosophila* species showed differential susceptibility to parasitism, which
manifests in shifts of the fitness differences in the coexistence plane. Natural enemies shift the
fitness difference in favour of the species that is less susceptible, but this does not necessarily
enhance coexistence. Across our six species, no general reproductive rate – susceptibility trade-off
was observable and the presence of parasitoids did not lead to a consistent shift towards (or away
from) a state of potential coexistence. At the pairwise level, reproductive rate – susceptibility
relationships did not provide a useful guide to the impact of parasitoids on coexistence. This is in line
with results of Petry et al (2018) where seed predators reshuffled the results of competition and
emphasises the insufficiency of fecundity as a substitute for competitive ability in practice. However,
in contrast to Petry et al (2018) we did not observe any cases of 'overshooting' the zone of
coexistence and reversing the victor of competition.

Fully quantifying pairwise competitive ability requires considerable effort (Hart et al., 2018), that
scales quadratically with the number of species to be assessed. Since an increased fecundity
increases competitive ability, taking shortcuts based on direct measured of fecundity and
susceptibility to determine the role of natural enemies is appealing. However, our results provide
further evidence that this represents a dangerous trap, and there is no substitute to full
characterisation of interspecific competition. Predation can only favour coexistence when the more
susceptible species is otherwise competitively dominant, not just more fecund. This underscores the

importance and challenges of accurately measuring competition terms ($\alpha$) to understand the

consequences of any given factor's influence on coexistence.

## Impact of enemies on competition coefficients

In natural systems, both direct effects of mortality and impacts on the interaction between species

will determine the impact of consumers on species coexistence (Pringle et al., 2019). Our study

design allowed parasitoids to travel between vials, opening the prospect of either associational

susceptibility or resistance mediated through parasitoid foraging behaviour (Abrams, 2010; Holt &

Kotler, 1987). If these competitive interaction modification effects were strong enough to be

detectable, they would manifest as shifts in the fitted $\alpha$ values and change the niche differences

between parasitism and no-parasitism treatments. However, our model selection procedure

suggested that modelling changes to the competition coefficients matrix between treatments leads

to overfitting i.e. there were no identifiable impact of parasitism on pairwise competition

coefficients. Resolving the relative importance of direct effects of predators versus their indirect

effects on other species is a long standing challenge (Werner & Peacor, 2003). Our study suggests

that in this case the direct effects are dominant. However, mechanisms that can be modelled as

shifts in single-species parameters (e.g. $R_0$) are inherently more identifiable statistically than

changes to multi-species processes (e.g. $\alpha_{ij}$) as they can be determined more directly from simpler

experiments.

Nevertheless, our model selection test is a rather blunt all-or-nothing approach. It may well be that

only particular competition coefficients are strongly affected by the parasitoid. Indeed, in our Model

1, while most of the between-treatment $\alpha$ differences cluster tightly together, several show large

divergences (SI 4 Fig. S4.14). There is no clear predictor for the terms that show these large shifts,

occurring between different species. A more complex model-fitting approach incorporating

regularisation could potentially determine whether such cases are identifiable effects rather than

model-fitting noise.

Our single-generation design meant that density-mediated apparent competition derived from an

increased total population of natural enemies was not directly detectable. If we had run a multi-

generational experiment, the results could be expected to be complex: Sommers & Chesson (2019)

show that the relative balance between apparent and resource competition is influenced by prey

defensive actions. Methods to compare resource competition and predation effects directly in

parameterised models are being developed (Shoemaker et al., 2020) and will be an exciting area for

future research.

## Model fitting considerations within coexistence theory

Recent meta-analyses have emphasised the importance of parameterised models for understanding

mechanisms of coexistence (Broekman et al., 2019), particularly since large-scale global

environmental changes will create novel communities (Alexander et al., 2016). Modern coexistence

theory defines a sharp line between coexistence and the inability to invade from rare. This

mathematical clarity and precision represents a major advantage, but poses challenges to the

incorporation of inevitable experimental uncertainty, even before comparison to field contexts. It is

important to reiterate that the uncertainty of coexistence we discuss should be interpreted as

statements of uncertainty in the underlying data, rather than implying a probabilistic outcome. The

precision of estimates of parameters can differ between treatments, which can have the effect of

changing the support for coexistence, even if the central estimate does not change. Furthermore,

these values should not be directly interpreted as some measure of 'distance from coexistence',
although the concepts may be related. The framework of structural stability is better equipped to
address such questions (Godoy et al., 2018; Saavedra et al., 2017).

Nonetheless, parameter fitting can have high data demands, and key ecological model parameters
are often not fully separable (Bolker, 2008). For example, in a simple model of competition similar
predictions could be made by lowering the growth rate term and increasing the competition term.
With a well-designed experiment and sufficient data, uncertainty can be minimised, but in any
realistically-sized competition experiment, separability of parameters can be a significant but largely
underacknowledged issue. This problem is particularly pertinent when working within the
framework of modern coexistence theory, where conclusions are drawn from comparisons between
ratios composed of multiple parameters from the underlying model (Barabás et al., 2018; Song et al.,
2019). Failing to address these issues could dramatically affect the confidence with which
conclusions are drawn from experimental data.

Understanding how the interrelationships between parameters affect uncertainty in the derived
quantities based upon them is difficult *a priori,* since the contribution of different correlations will be
contingent on both the underlying ecology and the experimental design. In the context of modern
coexistence theory, even a moderate confounding of underlying growth rates and intra-specific
competition terms may have a significant effect on the conditions for coexistence. In our study, the
posteriors were very often curved - posterior draws generating large niche differences were
associated with larger relative fitness differences. This has the beneficial effect of often increasing
the confidence with which pairs of species are located within regions of the coexistence plane.
Although our individual estimates of model parameters may be rather uncertain, we can often be
comparatively confident in the result of competition.

This effect can be attributed to indeterminacy, and hence correlation in the posterior, between
growth rates and intra-specific competition terms. One consequence of this can be seen clearly
when the condition for coexistence (equation 2) is re-arranged in a simplified form as:

$$\frac{\alpha_{ji}}{\alpha_{ii}} < \frac{r_j - 1}{r_i - 1} < \frac{\alpha_{jj}}{\alpha_{ij}} \qquad (5)$$

Where (as in our data) there are strong positive correlations in the posterior draws of $\alpha_{ii}$ and $r_i$
(shown in bold), this would tend to affect both sides of the inequality simultaneously, increasing the
tendency that it maintains its value.

Fitting parameterising models with ecological data is almost always both challenging and
illuminating in equal measure. Because the fitted values of the parameters inevitably correlate to
some degree, propagating parameter uncertainty from a multi-parameter posterior distribution
gives a better understanding of the uncertainty associated with the key quantities of modern
coexistence theory (Mordecai, 2013; Uricchio et al., 2019). Multi-dimensional posteriors can pose
considerable challenges both to intuition and to visualise. However, approaches that directly
propagate standard errors associated with individual parameters (e.g. Matías et al. 2018) or
estimate uncertainty by bootstrapping observations (e.g. García-Callejas et al. (2020) are unable to
capture these effects, and risk misrepresenting the confidence that can be placed in results.
Although we took an explicitly Bayesian approach using weak priors, similar methods could equally
be followed using multi-dimensional likelihood surfaces computed with a frequentist approach.

## Limitations of simple laboratory experiments

The modern coexistence theory approach is agnostic with respect to the particular factors involved but will always be limited by the scope of the empirical data available. Experiments such as this one can rarely replicate full natural lifecycles of animals or capture the full range of conditions that occur throughout their habitat (Hart et al., 2018). Impacts on one life stage can mitigate differences at another (Moll & Brown, 2008) and our method does not capture differences in reproductive lifespan. Equally, our experiment uses a highly simplified representation of the consumer pressure experienced by natural communities of *Drosophila*. We included a single pupal parasitoid species; in their natural habitats these species exist in a complex trophic network (Jeffs et al., 2020) including specialist parasitoids and larval parasitoids (which may directly influence larval competitive performance).

Coexistence may only be possible at particular spatial scales (Hart et al., 2017), or through spatial mechanisms not reflected in our experiment. The laboratory experiments cannot capture environmental heterogeneities hypothesised to aid coexistence (Chesson, 2000; Kuang & Chesson, 2010). Despite these limitations, we believe that our experiment is still instructive in isolating particular aspects for further investigation.

The *Drosophila* species we investigated co-occur in their natural habitat of Australian rainforest (Jeffs et al., 2020), although it is important to note that this is not necessarily evidence that they can coexist in the sense of mutual invasibility (Siepielski & McPeek, 2010). Furthermore, pairwise coexistence is not the same as community-level coexistence, and coexistence or otherwise between pairs does not necessarily translate to the multispecies community (Song et al., 2019), although they are likely to be at least somewhat related (Broekman et al., 2019; Friedman et al., 2017). A diversity of mechanisms can allow multispecies coexistence where pairwise coexistence is not possible (Letten & Stouffer, 2019; Levine et al., 2017; McPeek, 2019).

## Conclusion

Our experiment demonstrates that natural enemies can change the result of competition between species pairs, but that this effect is not consistent or necessarily predictable from pairwise reproduction rate - susceptibility relationships, reiterating the value of analysing parametrised models (Siepielski & McPeek, 2010). Further work will be needed to investigate whether trait differences impact the niche and fitness differences observed in our system, and more experimental studies of the impact of natural enemies on coexistence are needed to form the basis for a more general understanding of their impacts beyond plants (Viola et al., 2010). Modern coexistence theory is emerging as a powerful tool to investigate a range of biotic and abiotic impacts on coexistence experimentally (Lanuza et al., 2018; Matías et al., 2018). However, since the precision with which growth rates and competition coefficients can be estimated is limited (even within large studies such as ours), the fundamental inseparability of niche and fitness differences can impact the likelihood assigned to coexistence between any given species pair. Although these challenges are significant, we hope our approach demonstrates that they can be readily and transparently addressed.

## Acknowledgements:

We thank the Hrček lab and the Oxford Fly group for advice and use of facilities, Chia-Hua Lue for advice on parasitoid taxonomy, František Sládeček and Jamal Holle for assistance maintaining cultures, and Mark Wong for constructive comments on the manuscript.

## References

Abrams, P. A. (2010). Implications of flexible foraging for interspecific interactions: lessons from simple models. *Functional Ecology, 24*(1), 7–17. https://doi.org/10.1111/j.1365-2435.2009.01621.x

Adler, P. B., HilleRisLambers, J., & Levine, J. M. (2007). A niche for neutrality. *Ecology Letters, 10*(2), 95–104. https://doi.org/10.1111/j.1461-0248.2006.00996.x

Alexander, J. M., Diez, J. M., Hart, S. P., & Levine, J. M. (2016). When Climate Reshuffles Competitors: A Call for Experimental Macroecology. *Trends in Ecology and Evolution, 31*(11), 831–841. https://doi.org/10.1016/j.tree.2016.08.003

Atkinson, W. D., & Shorrocks, B. (1981). Competition on a divided and ephemeral resource: a simulation model. *Journal of Animal Ecology, 50*(2), 461–471.

Barabás, G., D'Andrea, R., & Stump, S. M. (2018). Chesson's coexistence theory. *Ecological Monographs, 88*(3), 277–303. https://doi.org/10.1002/ecm.1302

Bartomeus, I., & Godoy, O. (2018). Biotic controls of plant coexistence. *Journal of Ecology, 106*(5), 1767–1772. https://doi.org/10.1111/1365-2745.13040

Bolker, B. M. (2008). *Ecological Models and Data in R.* Princeton University Press.

Broekman, M. J. E., Muller-Landau, H. C., Visser, M. D., Jongejans, E., Wright, S. J., & Kroon, H. (2019). Signs of stabilisation and stable coexistence. *Ecology Letters, 22*(11), 1957–1975. https://doi.org/10.1111/ele.13349

Burnham, K. P., & Anderson, D. R. (2002). *Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach.* Springer Science & Business Media.

Carpenter, B., Gelman, A., Hoffman, M. D., Lee, D., Goodrich, B., Betancourt, M., Brubaker, M., Guo, J., Li, P., & Riddell, A. (2017). Stan⊠: A Probabilistic Programming Language. *Journal of Statistical Software, 76*(1). https://doi.org/10.18637/jss.v076.i01

Chase, J. M., Abrams, P. A., Grover, J. P., Diehl, S., Chesson, P., Holt, R. D., Richards, S. A., Nisbet, R. M., & Case, T. J. (2002). The interaction between predation and competition: a review and synthesis. *Ecology Letters, 5*(2), 302–315. https://doi.org/10.1046/j.1461-0248.2002.00315.x

Chesson, P. (2000). Mechanisms of Maintenance of Species Diversity. *Annual Review of Ecology, Evolution, and Systematics, 31*, 343–366.

Chesson, P. (2018). Updates on mechanisms of maintenance of species diversity. *Journal of Ecology, 106*(5), 1773–1794. https://doi.org/10.1111/1365-2745.13035

Chesson, P., & Kuang, J. J. (2008). The interaction between predation and competition. *Nature, 456*(7219), 235–238. https://doi.org/10.1038/nature07248

Davis, A. J., Jenkinson, L. S., Lawton, J. H., Shorrocks, B., & Wood, S. (1998). Making mistakes when predicting shifts in species range in response to global warming. *Nature, 391*(6669), 783–786. https://doi.org/10.1038/35842

Duménil, C., Woud, D., Pinto, F., Alkema, J. T., Jansen, I., van der Geest, A. M., Roessingh, S., & Billeter, J. C. (2016). Pheromonal Cues Deposited by Mated Females Convey Social Information about Egg-Laying Sites in Drosophila Melanogaster. *Journal of Chemical Ecology, 42*(3), 259–269. https://doi.org/10.1007/s10886-016-0681-3

Fellowes, M. D. E., & Godfray, H. C. J. (2000). The evolutionary ecology of resistance to parasitoids by Drosophila. *Heredity, 84*(1), 1–8. https://doi.org/10.1046/j.1365-2540.2000.00685.x

Friedman, J., Higgins, L. M., & Gore, J. (2017). Community structure follows simple assembly rules in microbial microcosms. *Nature Ecology & Evolution, 1*(5), 0109. https://doi.org/10.1038/s41559-017-0109

García-Callejas, D., Godoy, O., & Bartomeus, I. (2020). cxr⊠: A toolbox for modelling species coexistence in R. *Methods in Ecology and Evolution, 1*(1),

2041-210X.13443. https://doi.org/10.1111/2041-
210X.13443

Gilpin, M. E., Carpenter, M. P., & Pomerantz, M. J.
(1986). The assembly of a laboratory community:
multi-species competition in Drosophila. In J. M.
Diamond & T. J. Case (Eds.), *Community Ecology*.
Harper and Row.

Godoy, O., Bartomeus, I., Rohr, R. P., & Saavedra, S.
(2018). Towards the Integration of Niche and
Network Theories. *Trends in Ecology & Evolution*,
*33*(4), 287–300.
https://doi.org/10.1016/j.tree.2018.01.007

Godoy, O., & Levine, J. M. (2014). Phenology effects on
invasion success: Insights from coupling field
experiments to coexistence theory. *Ecology*,
*95*(3), 726–736. https://doi.org/10.1890/13-
1157.1

Godwin, C. M., Chang, F., & Cardinale, B. J. (2020). An
empiricist's guide to modern coexistence theory
for competitive communities. *Oikos*, *129*(8),
1109–1127. https://doi.org/10.1111/oik.06957

Grainger, T. N., Letten, A. D., Gilbert, B., & Fukami, T.
(2019). Applying modern coexistence theory to
priority effects. *Proceedings of the National
Academy of Sciences*, *116*(13), 6205–6210.
https://doi.org/10.1073/pnas.1803122116

Grainger, T. N., Levine, J. M., & Gilbert, B. (2019). The
Invasion Criterion: A Common Currency for
Ecological Research. *Trends in Ecology &
Evolution*, *34*(10), 925–935.
https://doi.org/10.1016/j.tree.2019.05.007

Gurevitch, J., Morrison, J. A., & Hedges, L. v. (2000). The
interaction between competition and predation:
A meta-analysis of field experiments. *American
Naturalist*, *155*(4), 435–453.
https://doi.org/10.1086/303337

Hangartner, S. B., Hoffmann, A. A., Smith, A., & Griffin, P.
C. (2015). A collection of Australian Drosophila
datasets on climate adaptation and species
distributions. *Scientific Data*, *2*(1), 150067.
https://doi.org/10.1038/sdata.2015.67

Hart, S. P., Freckleton, R. P., & Levine, J. M. (2018). How
to quantify competitive ability. *Journal of Ecology*,
*106*(5), 1902–1909.
https://doi.org/10.1111/1365-2745.12954

Hart, S. P., Usinowicz, J., & Levine, J. M. (2017). The
spatial scales of species coexistence. *Nature
Ecology & Evolution*, *1*(8), 1066–1073.
https://doi.org/10.1038/s41559-017-0230-7

Hassell, M. P. (2000). *The Spatial and Temporal
Dynamics of Host-Parasitoid Interactions*. Oxford
University Press.

Holt, R. D., & Kotler, B. P. (1987). Short-term apparent
competition. *American Naturalist, 130*(3), 412–
430. https://doi.org/10.1086/284718

Jeffs, C. T., Terry, J. C. D., Higgie, M., Jandová, A.,
Konvičková, H., Brown, J. J., Lue, C.-H., Schiffer,
M., O'Brien, E. K., Bridle, J., Hrček, J., & Lewis, O.
T. (2020). Molecular analyses reveal consistent
food web structure with elevation in rainforest
Drosophila - parasitoid communities. *BioRxiv*, 1-
16 https://doi.org/10.1101/2020.07.21.213678.
https://doi.org/https://doi.org/10.1101/2020.07.
21.213678

Ke, P.-J., & Letten, A. D. (2018). Coexistence theory and
the frequency-dependence of priority effects.
*Nature Ecology & Evolution, 2*(11), 1691–1695.
https://doi.org/10.1038/s41559-018-0679-z

Kraaijeveld, A. R., & Godfray, H. C. J. (1997). Trade-off
between parasitoid resistance and larval
competitive ability in Drosophila melanogaster.
*Nature, 389*(1464), 259–261.
https://doi.org/10.1098/rspb.2000.1354

Kuang, J. J., & Chesson, P. (2010). Interacting
coexistence mechanisms in annual plant
communities: Frequency-dependent predation
and the storage effect. *Theoretical Population
Biology, 77*(1), 56–70.
https://doi.org/10.1016/j.tpb.2009.11.002

Lanuza, J. B., Bartomeus, I., & Godoy, O. (2018).
Opposing effects of floral visitors and soil
conditions on the determinants of competitive
outcomes maintain species diversity in
heterogeneous landscapes. *Ecology Letters, 21*(6),
865–874. https://doi.org/10.1111/ele.12954

Letten, A. D., Ke, P. J., & Fukami, T. (2017). Linking
modern coexistence theory and contemporary
niche theory. *Ecological Monographs, 87*(2), 161–
177. https://doi.org/10.1002/ecm.1242

Letten, A. D., & Stouffer, D. B. (2019). The mechanistic
basis for higher-order interactions and non-
additivity in competitive communities. *Ecology
Letters, 22*(3), 423–436.
https://doi.org/10.1111/ele.13211

Levine, J. M., Bascompte, J., Adler, P. B., & Allesina, S.
(2017). Beyond pairwise mechanisms of species
coexistence in complex communities. *Nature*,

*546*(7656), 56–64.
https://doi.org/10.1038/nature22898

Levine, J. M., & HilleRisLambers, J. (2009). The importance of niches for the maintenance of species diversity. *Nature, 461*(7261), 254–257. https://doi.org/10.1038/nature08251

Matías, L., Godoy, O., Gómez-Aparicio, L., & Pérez-Ramos, I. M. (2018). An experimental extreme drought reduces the likelihood of species to coexist despite increasing intransitivity in competitive networks. *Journal of Ecology, 106*(3), 826–837. https://doi.org/10.1111/1365-2745.12962

McPeek, M. A. (2019). Mechanisms influencing the coexistence of multiple consumers and multiple resources: resource and apparent competition. *Ecological Monographs, 89*(1), e01328. https://doi.org/10.1002/ecm.1328

Moll, J. D., & Brown, J. S. (2008). Competition and coexistence with multiple life-history stages. *The American Naturalist, 171*(6), 839–843. https://doi.org/10.1086/587517

Mordecai, E. A. (2013). Despite spillover, a shared pathogen promotes native plant persistence in a cheatgrass-invaded grassland. *Ecology, 94*(12), 2744–2753. https://doi.org/10.1890/13-0086.1

Nottebrock, H., Schmid, B., Treurnicht, M., Pagel, J., Esler, K. J., Böhning-Gaese, K., Schleuning, M., & Schurr, F. M. (2017). Coexistence of plant species in a biodiversity hotspot is stabilized by competition but not by seed predation. *Oikos, 126*(2), 276–284. https://doi.org/10.1111/oik.03438

O'Brien, E. K., Higgie, M., Reynolds, A., Hoffmann, A. A., & Bridle, J. R. (2017). Testing for local adaptation and evolutionary potential along altitudinal gradients in rainforest Drosophila: beyond laboratory estimates. *Global Change Biology, 23*(5), 1847–1860. https://doi.org/10.1111/gcb.13553

Paine, R. T. (1966). Food Web Complexity and Species Diversity. *The American Naturalist, 100*(910), 65–75. https://doi.org/10.1086/282400

Pearl, R. (1932). The influence of density of population upon egg production in Drosophila melanogaster. *Journal of Experimental Zoology, 63*(1), 57–84. https://doi.org/10.1002/jez.1400630103

Pérez-Ramos, I. M., Matías, L., Gómez-Aparicio, L., & Godoy, Ó. (2019). Functional traits and phenotypic plasticity modulate species coexistence across contrasting climatic conditions. *Nature Communications, 10*(1), 2555. https://doi.org/10.1038/s41467-019-10453-0

Petry, W. K., Kandlikar, G. S., Kraft, N. J. B. B., Godoy, O., & Levine, J. M. (2018). A competition-defence trade-off both promotes and weakens coexistence in an annual plant community. *Journal of Ecology, 106*(5), 1806–1818. https://doi.org/10.1111/1365-2745.13028

Pringle, R. M., Kartzinel, T. R., Palmer, T. M., Thurman, T. J., Fox-Dobbs, K., Xu, C. C. Y., Hutchinson, M. C., Coverdale, T. C., Daskin, J. H., Evangelista, D. A., Gotanda, K. M., A. Man in 't Veld, N., Wegener, J. E., Kolbe, J. J., Schoener, T. W., Spiller, D. A., Losos, J. B., & Barrett, R. D. H. (2019). Predator-induced collapse of niche structure and species coexistence. *Nature, 570*(7759), 58–64. https://doi.org/10.1038/s41586-019-1264-6

Saavedra, S., Rohr, R. P., Bascompte, J., Godoy, O., Kraft, N. J. B., & Levine, J. M. (2017). A structural approach for understanding multispecies coexistence. *Ecological Monographs, 87*(3), 470–486. https://doi.org/10.1002/ecm.1263

Schreiber, S. J., Yamamichi, M., & Strauss, S. Y. (2019). When rarity has costs: coexistence under positive frequency-dependence and environmental stochasticity. *Ecology, 100*(7), e02664. https://doi.org/10.1002/ecy.2664

Shoemaker, L. G., Barner, A. K., Bittleston, L. S., & Teufel, A. I. (2020). Quantifying the relative importance of variation in predation and the environment for species coexistence. *Ecology Letters, 23*(6), 939–950. https://doi.org/10.1111/ele.13482

Shorrocks, B. (1991). Competition on a divided and ephemeral resource: a cage experiment. *Biological Journal of the Linnean Society, 43*(3), 211–220. https://doi.org/10.1111/j.1095-8312.1991.tb00594.x

Siepielski, A. M., Hasik, A. Z., & Ousterhout, B. H. (2018). An ecological and evolutionary perspective on species coexistence under global change. *Current Opinion in Insect Science, 29*, 71–77. https://doi.org/10.1016/j.cois.2018.06.007

Siepielski, A. M., & McPeek, M. A. (2010). On the evidence for species coexistence: a critique of the coexistence program. *Ecology, 91*(11), 3153–3164. https://doi.org/10.1890/10-0154.1

Sommers, P., & Chesson, P. (2019). Effects of predator
avoidance behavior on the coexistence of
competing prey. *American Naturalist, 193*(5),
E132–E148. https://doi.org/10.1086/701780

Song, C., Barabás, G., & Saavedra, S. (2019). On the
consequences of the interdependence of
stabilizing and equalizing mechanisms. *American
Naturalist, 194*(5), 627–639.
https://doi.org/10.1086/705347

Spaak, J. W., & de Laender, F. (2020). Intuitive and
broadly applicable definitions of niche and fitness
differences. *Ecology Letters, 23*(7), 1117–1128.
https://doi.org/10.1111/ele.13511

Uricchio, L. H., Daws, S. C., Spear, E. R., & Mordecai, E. A.
(2019). Priority effects and nonhierarchical
competition shape species composition in a
complex grassland community. *American
Naturalist, 193*(2), 213–226.
https://doi.org/10.1086/701434

Valadão, H., Proença, C. E. B., Kuhlmann, M. P., Harris, S.
A., & Tidon, R. (2019). Fruit-breeding drosophilids
(Diptera) in the Neotropics: playing the field and
specialising in generalism? *Ecological Entomology,
44*(6), 721–737.
https://doi.org/10.1111/een.12769

Vehtari, A., Gelman, A., & Gabry, J. (2017). Practical
Bayesian model evaluation using leave-one-out
cross-validation and WAIC. *Statistics and
Computing, 27*(5), 1413–1432.
https://doi.org/10.1007/s11222-016-9696-4

Viola, D. v., Mordecai, E. A., Jaramillo, A. G., Sistla, S. A.,
Albertson, L. K., Gosnell, J. S., Cardinale, B. J., &
Levine, J. M. (2010). Competition-defense
tradeoffs and the maintenance of plant diversity.
*Proceedings of the National Academy of Sciences,
107*(40), 17217–17222.
https://doi.org/10.1073/pnas.1007745107

Werner, E. E., & Peacor, S. D. (2003). A Review of Trait-
Mediated Indirect Interactions in Ecological
Communities. *Ecology, 84*(5), 1083–1100.
http://www.jstor.org/stable/3107918

Wertheim, B., Marchais, J., Vet, L. E. M., & Dicke, M.
(2002). Allee effect in larval resource exploitation
in Drosophila: an interaction among density of
adults, larvae, and micro-organisms. *Ecological
Entomology, 27*, 608–617.
http://doi.wiley.com/10.1046/j.1365-
2311.2002.00449.x

Worthen, W. B. (1989). Predator-mediated coexistence
in laboratory communities of mycophagous
Drosophila (Diptera: Drosophilidae). *Ecological
Entomology, 14*(1), 117–126.
https://doi.org/10.1111/j.1365-
2311.1989.tb00761.x