Original Article

Ecological fitness cost associated with the AHAS Trp574Leu mutation in feral *Raphanus*

sativus

Roman B. Vercellino[1*], Fernando Hernández[1], Claudio E. Pandolfo[1], Miguel Cantamutto[2],

Alejandro Presotto[1]

[1]*CERZOS, Dpto. Agronomía, Universidad Nacional del Sur (UNS)-CONICET, San Andrés 800,*

CP 8000, Bahía Blanca, Argentina, [2]*Estación Experimental Agropecuaria Hilario Ascasubi,*

Instituto Nacional de Tecnología Agropecuaria, Ruta 3 Km 794, CP 8142, Hilario Ascasubi,

Argentina.

Fitness cost in AHAS-resistant *Raphanus sativus*

ORCIDs:

Roman B. Vercellino: https://orcid.org/0000-0002-9313-3501

Fernando Hernández: https://orcid.org/0000-0001-5158-2029

Claudio E. Pandolfo: https://orcid.org/0000-0002-7089-7138

Miguel Cantamutto: https://orcid.org/0000-0002-6854-7497

Alejandro Presotto: https://orcid.org/0000-0002-9085-9673

* **Correspondence**. Roman B Vercellino, Departamento de Agronomía, Universidad Nacional

del Sur, San Andrés 800, 8000 Bahía Blanca, Argentina. *E-mail rbvercellino@cerzos-*

conicet.gob.ar

Abstract

BACKGROUND: Gene mutations endowing herbicide resistance may have negative pleiotropic

effects on plant fitness. Quantifying these effects is critical for predicting the evolution of

herbicide resistance and developing management strategies for herbicide-resistant weeds. This

study reports the effects of the acetohydroxyacid synthase (AHAS) Trp574Leu mutation

throughout the life cycle of the weed feral radish (*Raphanus sativus* L.).

RESULTS: Resistant and susceptible biotypes responded differently to light and water

treatments in relation to germination and emergence. Under light exposure, the resistant biotype

showed higher germination and emergence, but no differences were found in seed dormancy,

germination in darkness and emergence from buried seeds or pods. The resistant biotype

showed delayed and reduced seedling emergence relative to the susceptible biotype under

rainfed conditions, but these differences between the biotypes were not detected in irrigated soil.

The biotypes showed similar relative growth rates and vegetative biomass. However, under

wheat interference, resistant plants had 36–46% less total above-ground biomass, 26–47% less

seeds per plant, and 36–53% less plant yield than susceptible ones, and these differences were

more evident at higher plant density.

CONCLUSION: This study provides a better understanding of the ecological fitness cost

associated with the AHAS Trp574Leu mutation in feral *R. sativus*. The fitness costs could

reduce the frequency of the resistant allele in areas untreated with AHAS inhibiting herbicides.

Keywords: cost of resistance, pleiotropic effects, feral radish, herbicide resistance evolution,

AHAS herbicides.

1. INTRODUCTION

Acetohydroxyacid synthase (AHAS), also known as acetolactate synthase (ALS), is the first enzyme in the biosynthesis of the branched chain amino acids valine, leucine and isoleucine, and the common target-site of AHAS-inhibiting herbicides (hereafter referred to as AHAS herbicides)[1]. Repeated use of AHAS herbicides in agricultural fields has resulted in the evolution of AHAS-resistant biotypes in at least 165 weed species worldwide [2]. Target-site resistance (TSR) is the most common mechanism resulting in resistant to AHAS herbicides[2,3]. In the AHAS gene, at least 29 resistance-endowing gene mutations in eight conserved amino acids were identified in field-evolved AHAS-resistant weed biotypes[4]. Some of them may alter normal enzymatic activity, resulting in a whole plant fitness cost [5].

Mutations conferring herbicide resistance, which enable adaptation to a new environment, may have negative pleiotropic effects, also called fitness costs [6,7]. These fitness costs could influence the persistence and evolution of the adaptive alleles in the absence of the selective agent[5,8]. It has been reported that fitness costs associated with TSR to AHAS herbicides can be manifested in different life-history traits (i.e. dormancy, germination, early growth and reproductive outputs)[6,7,9] and differentially expressed depending on the specific mutation[10], weed species[11,12] and environmental conditions[5,13,14]. There is also a general idea that fitness costs may be more evident under stressful environmental conditions[6,7], but this prediction has not always been identified[13,14]. A case-by-case analysis is thus required to study the expression of the specific gene AHAS-resistant mutation in fitness costs. Quantifying fitness costs associated with herbicide resistant alleles throughout the entire life cycle is necessary to predict the dynamic of the evolution of herbicide resistance and to develop management strategies for herbicide-resistant weeds[7,8,15].

Raphanus sativus L. (radish) is an ancient root crop from the Brassicaceae family. Spontaneous populations of radish found in Europe, North America, South America and Japan have been classified as de-domesticated (feral) forms derived from the cultivated biotype [16]. Radish and its feral forms are self-incompatible and insect-pollinated with an annual or biennial life cycle [16]. Feral radish is a widespread problematic winter weed in temperate zones of the Americas that has developed resistance to AHAS herbicides in Brazil, Chile and Argentina [2,17]. In Argentina, feral radish is one of the most noxious weeds in the south of Buenos Aires province, the main winter cereal producing area[18], and the Trp574Leu mutation has been identified in several AHAS-resistant populations [17,19].

In a previous study, AHAS-resistant feral radish carrying the Trp574Leu mutation showed negative pleiotropic effects on AHAS activity and reproductive traits under intraspecific interference[19]. However, the lack of information of fitness costs associated with the Trp574Leu

mutation in different life-history traits of feral radish and under crop interference limits the

development and implementation of evolution-knowledge-based resistance management

strategies [7]. The comparison of resistant and susceptible individuals with similar genetic

backgrounds is crucial for the quantification of fitness costs associated with herbicide resistant

alleles [6,7,20]. Here, we compare recently evolved AHAS-resistant Trp574Leu homozygous

individuals with susceptible individuals from a single population to minimize the effect of

genetic background[15,20,21]. The aims of this study were to investigate the effect of the Trp574Leu

mutation on (1) seed dormancy, germination and emergence under contrasting conditions of

temperature, light, water and burial depth, and on (2) relative growth rate, and vegetative and

reproductive traits, under intra- and interspecific interference. This study will provide a better

understanding of the fitness costs of the AHAS Trp574Leu mutation in feral *R. sativus*.

2. MATERIALS AND METHODS

2.1 Plant material

The first AHAS-resistant feral *R. sativus* population reported in Argentina was used in this

study[2,19,22]. Recently evolved AHAS–resistant plants carrying the Trp574Leu mutation were

collected in the south of Buenos Aires province, in a field planted with IMI–resistant oilseed

rape after imazethapyr application by the farmer[19,22]. From these plants, a purified homozygous

sub-population with the Trp574Leu mutation (hereafter R biotype) was generated by selecting

at least 15 homozygous resistant plants using a specific Trp574Leu mutation CAPS marker,

following the procedure described in Pandolfo *et al.*[17]. Because susceptible plants within field

were killed by imazethapyr and they could not be isolated from the progeny of plants collected

within the field, their tested AHAS–susceptible counterparts collected in the field margin

(adjacent to the crop, at less than 50 m away from resistant plants), which remained without

imazethapyr application, were used to minimize differences in genetic background[19,22]. These

plants were multiplied once in a common garden and were used as the susceptible wild-type

(hereafter S biotype). The obtention of R and S accessions from the original population is

shown in Fig. S1. In this field, feral radish plants constitute a continuum, invading cultivated

fields, field margins and fence lines since several years prior to our collection, and because feral

radish is a self-incompatible and insect-pollinated species with high levels of gene flow[23], we

can assume similar genetic backgrounds in R and S [20,21].

2.2 Control of maternal environmental effects

To increase the amount of seed and to minimize the maternal environmental effects, at least 15

plants of each biotype (R and S) were used to produce a new generation. To prevent cross-

pollination, each biotype was isolated in an individual tent built with a pollen-proof mesh. Open

pollination was allowed within each biotype (each tent) by adding hives of honeybees (*Apis*

mellifera). At the end of the growing season, mature pods were collected and seeds of a portion

of pods were extracted by hand or using a mortar. Seeds and intact pods were stored at room

temperature and less than 10% moisture until use. Two-hundred progeny individuals were

randomly selected from within each sub-population (R and S), and AHAS-resistance and -

susceptibility was verified by applying metsulfuron-methyl at double the recommended rate

(X=6 g a.i. ha^{-1}).

2.3 Effects of dry storage, temperature and light on seed germination

In order to determine the effect of the Trp574Leu mutation on seed dormancy and germination,

isolated seeds and seeds inside intact pods of R and S biotypes stored under laboratory

conditions for 0 (freshly harvested), 2, 6 and 12 months were tested for germination in 9-cm-

diameter Petri dishes incubated at 14/4, 21/9 and 28/14 °C in light/darkness (12/12 h) or in

constant darkness for 15 days, according to Vercellino *et al.*[24]. The 14/4, 21/9 and 28/14°C

regimes represent the average winter, spring–autumn and summer temperatures in the south of

Buenos Aires province. At the end of the experiment, the pods were cut open, and the

germinated and non-germinated seeds were counted. A seed was considered germinated when

the radicle had emerged. The viability of non-germinated seeds was tested using a 1 % (w/v)

tetrazolium chloride solution. As the tetrazolium test on the non-germinated seeds showed

>99% viability, all non-germinated seeds were considered as alive. The experiment was

conducted as a completely randomized design with four replicates.

2.4 Effect of burial depth on seedlings emergence

To determine the effect of the Trp574Leu mutation on seeding emergence at different burial

depths, twenty-five isolated seeds and five intact pods (29.8 ± 1.2 seeds per replicate) of R and

S biotypes (dry-stored for 2 months) were sown at 0, 1, 8, 12 and 16 cm-depths in 10 L (18 cm

diameter × 24 cm height) plastic pots containing a sandy-loam soil with 1.1% organic matter

and pH 7.7 (the typical soil in the southwest of Buenos Aires province). Six pots per treatment

[two seed treatments (isolated seeds and seed inside intact pods) × two biotypes × five soil-

depth] were established in a greenhouse (20 ± 5 °C, natural light), and watered regularly.

Seedling emergence was evaluated, and the seedlings were removed periodically (3–4 days

intervals) for 85 days. For each experiment, the proportion of emerged seedlings was calculated

as the ratio between the sown seeds and the emerged seedlings at the end of the experiment. To

determine the percentage of emergence in the seeds inside pod treatments, six replicates of ten

pods of each biotype were threshed and their seeds were counted. The experiment was

conducted as a completely randomized design with six replicates.

2.5 Seedling emergence pattern

An outdoor experiment (38°41′38″S, 62°14′53″W) was conducted in the Agronomy Department's experimental field at the Universidad Nacional del Sur, Bahía Blanca, Argentina, to evaluate the seedling emergence pattern of the R and S biotypes. In mid-summer (6 Feb 2018), 50 isolated seeds and 10 intact pods of R and S biotypes were sown at a 0.5 cm-depth in 10 L plastic pots containing the same soil that the previous experiment, under two soil irrigation. Six pots per treatment [two seed treatments (isolated seeds and seed inside intact pods) × two biotypes × two irrigation (irrigated and non-irrigated)] were established. In the irrigated treatment, the soil was watered to field capacity every 3 – 4 days throughout the experiment, while in the non-irrigated treatment the soil only received water from rainfall. Seedling emergence was monitored weekly for one year. Daily minimum and maximum air temperature and rainfall data were obtained from the National Research Council of Argentina (CONICET), Bahía Blanca (38° 39' 60'' S, 62° 13' 58'' W) (Fig. S2). For each experiment, the proportion of emerged seedlings was calculated as above. The experiment was conducted as a completely randomized design with six replicates.

2.6 Vegetative growth

In a greenhouse in Bahía Blanca, Argentina, three experiments were performed to evaluate the relative growth rate (RGR) and competitive response of R and S biotypes during early vegetative growth. Both eco-physiological parameters are widely used to evaluate the expression of herbicide resistance fitness costs [11,12,25]. Growth experiments were conducted with isolated plants (physiological fitness cost) and plants growing under two interference environments, intraspecific interference and interspecific interference with wheat (*Triticum aestivum* L.) (ecological fitness cost). In all three experiments, the plants were grown in pots (15 cm diameter × 12 cm height) containing the same soil as above, in a greenhouse (20 ± 5 °C, natural light), watered regularly and fertilized with 6 L ha^{-1} of liquid fertilizer (Xilonen$^®$, grade 10-13-10) and 30 kg ha^{-1} urea (grade 46-0-0) until they reached the four-leaf stage. The pots were rearranged regularly to randomize any environmental differences within the greenhouse and distributed spatially to avoid competition for light.

2.6.1 Growth of plants without competition

Seeds of the R and S biotypes were germinated in Petri dishes at 14/28 °C in darkness for 48 h [24]. At least 50 germinated seedlings of uniform-size (< 1 cm) of R and S were transplanted into individual pots (one plant per pot). The above-ground biomass of 25 individual plants of each biotype were harvested at 18 and 32 days after transplanting (DAT), when plants were at two and four-leaf stage, respectively. The biomass was oven dried for a week at 60 °C and then weighed for dry biomass estimation. Pots were arranged in a completely randomized design with 25 replicates.

2.6.2 Growth of plants under intraspecific competition

A replacement-series design was used to evaluate which of R or S biotypes was the most

competitive. Uniform-size germinated seedlings (< 1 cm) of R and S were transplanted

equidistantly at four seedlings per pot. R and S seedlings were grown alone or in mixtures with

ratios of (R:S) 4:0 (100:0 %), 3:1 (75:25 %), 2:2 (50:50 %), 1:3 (25:75 %) and 0:4 (0:100 %).

The above-ground dry biomass from R and S plants was determined at 35 DAT, when plants

were at the four-leaf stage, as above. The pots were arranged in a randomized block design with

six replicates.

2.6.3 Growth of plants under interspecific competition

A target-neighbourhood design was used to evaluate the competitive responses of R and S

biotypes in competition with wheat[12]. One feral radish plant per pot was subjected to

competition from increasing densities of neighboring wheat plants (0, 57, 113, 226, 340 and 453

plants m^{-2}, which resulted in 0, 1, 2, 4, 6, and 8 plants pot^{-1}). Feral radish seedlings (<1 cm)

were transplanted into pots when the wheat plants were at the two-leaf stage[12,25]. The above-

ground dry biomass of R and S biotypes and wheat plants from each experimental unit (pot) was

determined at 38 DAT, when feral radish control plants (0 wheat plants m^{-2}) were at the four-

leaf stage, as above. The experiments were arranged in a completely randomized design with six

replicates.

2.7 Reproductive traits under field condition

Field experiments were conducted to evaluate the total above-ground dry biomass and

reproductive traits of R and S biotypes grown under interference with wheat. As feral radish is

considered a facultative species with two different cohorts[16], the experiments were performed on

two representative extreme planting dates. The facultative wheat was planted at the end of

autumn [mid-May 2018 (winter wheat), cultivar ACA 360 at 200 plants m^{-2}] and at the end of

winter [early August 2018 (spring wheat), cultivar KLEIN PROTEO at 350 plants m^{-2}]. Each

experimental unit was composed of nine rows 1.5 m in length and spaced at 0.2 m apart.

Seedlings of the R and S biotypes were established in plastic trays containing potting mix

(Grow Mix Terrafertil®), grown in a greenhouse (20 ± 5 °C, natural light) and watered daily. At

the two-three wheat leaf stage, feral radish seedlings (two to three-leaf stage) of each biotype (R

and S) were transplanted into the experimental field between the wheat plants according to the

planned weed density. The feral radish densities studied were 0 (control), 4, and 12 plants m^{-2}.

The trials were drip irrigated complementarily and fertilized with 90 kg ha^{-1} diammonium

phosphate at planting and 200 kg ha^{-1} urea at the five–leaf stage. The non-target weeds were

regularly removed by hand. The experiment was arranged in a split-plot design with the planting

date as the main plot, and the biotype and feral radish densities as sub-plots. Within each

planting date, experimental units were arranged in a randomized block design with four

replicates.

At the end of the growing season, plant height and branch number were measured in four

neighbouring feral radish plants located in the centre of each experimental unit. Then, the

above-ground biomass from the feral radish plants were hand-harvested, dried at 60 °C to

constant weight and weighed to determine the total dry biomass per plant. After that, pods per

plant, seeds per pod, seed weight, seeds per plant and plant yield were estimated following the

procedure described in Vercellino *et al.*,[19]. In each experimental unit, the wheat yield was

estimated by hand-harvesting and hand-threshing three replicates of 0.5 m long in the middle of

the three central rows. The wheat yield was standardized to 13.5 % moisture content. Data from

feral radish and wheat of each experimental unit were averaged for the statistical analysis.

2.8 Statistical analysis

Germination and emergence data were analyzed using generalized linear models (GLM) based

on maximum likelihood estimation (ML) with PROC GLIMMIX in SAS software (SAS

University Edition, SAS Institute Inc., Cary, NC, USA). For experiment '*2.3*', a binomial linear

model was fitted with a logit link to germination as the response variable. The models included

five fixed effects, i.e. light, storage time, temperature, seed treatment (isolated seeds and seeds

inside pods) and biotype, and all the interactions.

Two beta models were adjusted due to the non-normal natural distribution of the emergence

proportion data. For the experiment '*2.4*', the model included three fixed effects, i.e. seed

treatment, burial depth and biotype, and all the interactions. For experiment '*2.5*', the

emergence was evaluated at five different times after the experiment began (1, 3, 6, 9 and 12

months) and the model included three fixed effects, i.e. irrigation (irrigated and non-irrigated),

seed treatment and biotype, and all the interactions.

For experiment '*2.6.1*', the RGR was estimated using the unbiased formula proposed by

Hoffmann and Poorter[26]. The variance (*V*) of RGR was estimated with Venus and Causton[27].

The RGR was compared between R and S biotypes using the Student's t-test with PROC

TTEST in SAS. In addition, *ANOVA* was performed to compare the above-ground dry biomass

of the R and S biotypes using PROC GLM. The model included harvest time (18 and 32 DAT),

biotype, and their interaction. Data were log-transformed to improve the normality and

homogeneity of variances before analysis.

For experiment '*2.6.2*', the relative crowding coefficient (RCC) for above-ground dry biomass

was calculated according to the formula proposed by Novak *et al.*,[28] following the procedure

described by Tardif *et al*., [11]:

$$RCC = \left(\left(M_S^{3:1}/M_R^{3:1}\right) + \left(M_S^{2:2}/M_R^{2:2}\right) + \left(M_S^{1:3}/M_R^{1:3}\right)/3\right) / \left(M_S^{4:0}/M_R^{4:0}\right)$$

where M_S n:n and M_R n:n were the dry biomass per plant for the S and R biotypes at densities

n:n, respectively. An RCC value of 1 indicates that both biotypes have equal competitiveness.

RCC < 1 shows superior competitiveness of S over R biotype and RCC > 1 indicates the

opposite. The RCC values were compared to 1 using the Student's t-test with PROC TTEST. In

addition, *ANOVA* was performed to compare the above-ground dry biomass in the 4:0 (S) and

the 0:4 (R) treatments with PROC GLM.

For experiment '*2.6.3*', data for the above-ground dry biomass of feral radish grown in

competition with wheat was expressed as a percentage of that trait in the absence of

competition, and it was analyzed following the procedure described in Li *et al.*,[12] and Vila-Aiub

et al.,[25]. The response of R and S biotypes to competition by neighbouring wheat was analyzed

using a hyperbolic non-linear model, which described the response of the feral radish plants to

increasing biomass of neighbour wheat plants [29]:

$$Y = \frac{a}{1+bx}$$

where Y is the feral radish dry biomass at the neighbour biomass x, a is the feral radish dry

biomass in the absence of competitors (neighbours) (x = 0), and b the slope of the regression.

Slope of the non-linear regression model (b parameter) was compared between the R and S

biotypes using the Student's t-test with PROC TTEST. The hyperbolic non-linear model was

fitted after log-transforming data to improve the assumptions of the regression analysis.

In the experiment '*2.7*', *ANOVA* with PROC GLM was performed to investigate differences in

the total above-ground dry biomass per plant and per area (m^{-2}), plant height, branch number,

pods per plant, seeds per pod, seeds per plant, seed weight, plant yield and partitioning of the

dry biomass (stems and branches, pods and seeds) between the S and R biotypes. The model

included planting date, weed density (4 and 12 plants m^{-2}), biotype, all their interactions and the

block within planting date. In addition, *ANOVA* was performed to evaluate the effect of the

interference of two densities of S and R plants on wheat yield. Means were compared using

Fisher's LSD test ($P \leq 0.05$).

3. RESULTS

3.1 Effects of dry storage, temperature and light on germination

Analysis of the germination at varying times after harvest revealed significant effects of light

($F_{1;288} = 102.68$; $P < 0.0001$), seed treatment ($F_{1;288} = 77.57$; $P < 0.0001$), temperature ($F_{2;288} =$

31.05; $P < 0.0001$), biotype ($F_{1;288} = 6.70$; $P = 0.0101$) and the light by seed treatment interaction

($F_{1;288} = 23.97$; $P < 0.0001$). However, storage time effect, the five-order interaction, the four-

order interactions, the three-order interactions and the rest of the two-order interactions were not

significant (Table S1). In light, the R biotype showed higher seed germination than the S

throughout the experiment, both inside pods and isolated, especially in the 9/21°C and 14/28°C

treatments (Fig. 1A and C). These differences between the biotypes were not detected at low

temperatures (4/14 °C), nor in seed germination under darkness, either inside the pods or

isolated (Fig. 1B and D).

Figure 1. Final germination percentages (means ± 95 % confidence intervals) of isolated seeds

under light/dark (A), isolated seeds under constant darkness (B), seeds inside pods under

light/dark (C) and seeds inside pods under constant darkness (D) of AHAS herbicide-resistant

Trp475Leu (R) and susceptible wild-type (S) feral *Raphanus sativus* biotypes, incubated at three

alternating temperature regimes following 0, 2, 6 and 12 months of dry storage under laboratory

conditions.

3.2 Effect of burial depth on seedlings emergence

The analysis of the total seedling emergence revealed significant effects of the seed treatment ×

burial depth × biotype three-order interaction ($F_{5;120}$ = 2.58; P = 0.0299). Therefore, we

compared the differences in seedling emergence between the R and S biotypes for each seed

treatment and burial depth. On the soil surface, isolated seeds of the R biotype showed 56 % (25

vs. 14 %) higher germination than the S ($F_{1;10}$ = 8.73; P = 0.0144) (Fig. 2A). Inside pods, less

than 1% of the seeds germinated on the soil surface in both biotypes (Fig. 2A). As the burial depth increased, seedling emergence from both the isolated seeds and intact pods decreased, except for seeds sown on the soil surface. Seedling emergence from isolated seeds was higher than from the intact pods in both biotypes at all the depths evaluated (Fig. 2), except at 16 cm where less than 1% emergence was detected in both the isolated seeds and the intact pods (data not shown). However, no significant differences were found between the biotypes in cumulative seedling emergence when isolated seeds and intact pods were below the soil surface at up to 16 cm deep (Fig. 2B, C and D).

Figure 2. Dynamics of seedling emergence from isolated seeds and seeds inside pods, buried at 0 (A), 1 (B), 8 (C) and 12 (D) cm depth, of AHAS herbicide-resistant Trp475Leu (R) and susceptible wild-type (S) feral *Raphanus sativus* biotypes. Vertical bars represent the standard error of the mean.

3.3 Seedling emergence pattern

The mean annual temperature was 15.4 °C and mean monthly maximum and minimum temperatures of the coldest (July) and hottest (January) months were 12.0 and 3.0 °C (mean 7.1 °C) and 32.6 and 16.2 °C (mean 23.3 °C), respectively (Fig. S2). Annual precipitation was 592 mm, about 25% (152 mm) in autumn, 11% (65 mm) in winter, 39% (229 mm) in spring and the

remaining 25% (89 and 58 mm) in summer (6 Feb 2018 – 20 Mar 2018 and 21 Dec 2018 – 6

Feb 2019, respectively) (Fig. S2).

Analysis of seedling emergence revealed significant effects of the irrigation by seed treatment

and irrigation by biotype interactions throughout the study period (Table S2). Therefore, we

compared the differences in seedling emergence between the R and S biotypes for each

irrigation condition (Table S3). The biotype by seed treatment interaction was not significant for

any irrigation condition throughout the experiment (Table S3). The presence of the pericarp

strongly reduced seedling emergence in both the irrigation conditions and biotypes throughout

the study period (Fig. 3, Table S3).

Under rainfed conditions, the R biotype delayed and reduced seedling emergence relative to the

S (Fig. 3, Table S3). Seedling emergence of the R biotype was 13% (76 vs. 87%) and 76% (4

vs. 17%) lower than of the S biotype, from isolated seeds and intact pods, respectively, during

the first month (Fig. 3A). The R biotype showed a 6% additional emergence from isolated seeds

at the end of winter (Fig. 3A). The R and S biotypes showed 16 and 20% emergence during the

autumn and 28 and 20% emergence at the end of winter and in early spring, respectively (Fig.

3A). At the end of the experiment, the R biotype showed 7% (91 vs. 85%) and 15% (59 vs.

50%) lower seedling emergence than the S from isolated seeds and intact pods, respectively

(Fig. 3A, Table S3).

No differences were detected between the R and S biotypes in irrigated soil (Fig. 3B, Table S3).

Under this condition, about 90% of seedlings emerged from isolated seeds and only 40-45%

from intact pods during the first month with no differences between the biotypes (Fig. 3B).

Seedling emergence from intact pods exceeded 50% during autumn (mid-May) and raised an

additional 25% between late winter and spring (Fig. 3B). At the end of the experiment, the total

seedling emergence was more than 90% in the isolated seeds and 75% in the intact pods, with

no differences between the biotypes (Fig. 3B, Table S3).

Figure 3. Dynamics of seedling emergence from isolated seeds and seeds inside pods, buried at 1 cm depth in non-irrigated (A) and irrigated (B) soil conditions, of AHAS herbicide-resistant Trp475Leu (R) and susceptible wild-type (S) feral *Raphanus sativus* biotypes. Vertical bars represent the standard error of the mean.

3.4 Vegetative growth under isolation, intra and interspecific competition

No significant differences ($t = 0.76$; $P = 0.45$) were found in the RGR between R (0.1206 ± 0.0047 mg mg^{-1} día^{-1}) and S (0.1258 ± 0.0048 mg mg^{-1} día^{-1}) biotypes. There was no significant harvest time by biotype interaction ($F_{1;96} = 0.29$; $P = 0.59$). Dry biomass of the R biotype at 18 (295.02 ± 15.35 mg plant^{-1}) and 32 (1672.55 ± 144.65 mg plant^{-1}) DAT did not show any significant differences ($F_{1;96} = 3.23$; $P = 0.07$) compared to the S plants at 18 (322.37 ± 17.4 mg plant^{-1}) and 32 (1935.36 ± 143.73 mg plant^{-1}) DAT.

In the absence of inter-biotype competition, treatments 4:0 and 0:4, no significant differences ($F_{1;11} = 0.64$; $P = 0.44$) were found in the dry biomass between S and R (Fig. 4). However, when R and S biotypes were grown in a mixture, the RCC value was 1.25 ($t = 2.96$; $P = 0.0253$), indicating a slight competitive advantage of S relative to R (Fig. 4).

Figure 4. Replacement series diagram for the above-ground vegetative dry biomass of AHAS
herbicide-resistant Trp475Leu (R) and susceptible wild-type (S) feral *Raphanus sativus*
biotypes 35 days after transplanting. Vertical bars represent the standard error of the mean.

The hyperbolic model adequately explained variations in the feral radish dry biomass to
increasing biomass of neighboring wheat plants ($R^2 = 0.57 – 0.58$, $P < 0.0001$). This led to a
reduction in the feral radish dry biomass with increasing wheat competition (Fig. 5).
Competitive responses, evaluated by comparing estimates of regression slopes between the R
and S biotypes in the presence of wheat, did not show any significant differences ($P > 0.95$),
indicating similar reductions in feral radish dry biomass in the R and S biotypes with increasing
wheat biomass (Fig. 5).

Figure 5. Variations in the above-ground vegetative dry biomass of target AHAS herbicide–
resistant Trp475Leu (R) and susceptible wild-type (S) feral *Raphanus sativus* individuals under
increasing biomass of wheat neighbour plants. Comparison of regression slopes (*b* parameter)
determines differences in competitive responses of the target plants to wheat neighbour plants.

3.5 Evaluation of reproductive traits under field conditions

There were no significant interactions between planting date and biotype, planting date and

density, biotype and density and the third-order interaction for all evaluated traits, therefore data

from the two planting dates were pooled (Table S4, Fig. 6). Significant differences between S

and R biotypes were found for the total aboveground biomass ($F_{1;31} = 37.0$; $P < 0.0001$), pod

number ($F_{1;31} = 16.2$; $P = 0.0008$), seed number ($F_{1;31} = 27.5$; $P < 0.0001$), seed weight ($F_{1;31} =$

19.0; $P = 0.0004$) and plant yield ($F_{1;31} = 51.05$; $P < 0.0001$), but no significant differences were

found in plant height, branch number and seeds per pod (Table S4). The R biotype had 36–46%

less total dry biomass, 20–48% fewer pods per plant, 10–11% less seed weight, 26–47% fewer

seeds per plant, and 36–53% lower plant yield than its counterpart S, for both planting dates and

weed densities (Table S4, Fig. 6).

Analysis of biomass allocation to reproductive traits (i.e. proportion of seed biomass relative to

total dry biomass) revealed significant effects of biotype by density, and biotype by planting

date, interactions (Table S4). Therefore, the data were separated by planting date and weed

density (Fig. S3). At 4 weed plants m^{-2}, no significant differences were found in the biomass

allocation to reproductive structures between the R and S biotypes, neither in winter ($F_{1;\,7} =$

3.06; $P = 0.18$) or in spring ($F_{1;7} = 0.63$; $P = 0.49$). However, at 12 weed plants m^{-2}, the R plants

showed 19 % ($18,8 \pm 0,5$ vs. $15,2 \pm 0,6$ %) and 9 % ($20,4 \pm 0,6$ vs. $18,5 \pm 0,6$ %) lower biomass

allocation to reproduction structures than the S plants in both winter ($F_{1;7} = 24.75$; $P = 0.0156$)

and spring ($F_{1;7} = 14.20$; $P = 0.0327$) (Fig. S3).

The wheat yield was significantly reduced ($F_{4;39} = 58.41$; $P > 0.0001$) by feral radish

interference at both weed densities and biotypes. No significant differences were found between

planting dates ($F_{1;39} = 2.83$; $P = 0.11$), block (inside each planting date) ($F_{6;39} = 1.97$; $P = 0.11$)

and the interaction planting date by treatment (S and R biotypes at 0, 4 and 12 weed density)

($F_{4;39} = 0.98$; $P = 0.44$). The interference of the R biotype reduced the wheat yield less than the

S biotype at both feral radish densities (Fig. S4). The interference of R and S feral radish

biotypes reduced the wheat yield by 38 and 49 % at 4 weed plants m^{-2} and 54 and 66 % at 12

weed plants m^{-2}, respectively (Fig. S4).

Figure 6. Total above-ground dry biomass per plant (A) and per area (B), pods per plant (C), seeds per plant (D), seed biomass (E) and plant yield (F) of AHAS herbicide-resistant Trp475Leu (R) and susceptible wild-type (S) feral *Raphanus sativus* biotypes. Data are the average of two planting dates. Each point (●;●) is the average of four plants. Vertical bars represent the standard error of the mean.

4. DISCUSSION

In the present study, we evaluated several life-history traits defining the overall fitness costs associated with the AHAS Trp574Leu mutation in feral radish. In relation to germination and emergence, no differences were found between the R and S biotypes in seed dormancy and germination in darkness, or in seedling emergence from buried seeds/pods, but the R biotype showed higher germination and emergence than the S under light exposure (Fig. 1 and 2). The R biotype showed delayed and reduced seedling emergence relative to the S under rainfed conditions, but these differences were not detected under irrigation (Fig. 3). Finally, both

biotypes showed similar relative growth rates and vegetative biomass. However, the R biotype showed lower competitive ability, total above-ground dry biomass and reproductive outputs than the S (Fig. 4 and 6). All of these suggest that the pleiotropic effects imposed by the Trp574Leu mutation can be expressed throughout the life-history of feral radish and in response to several conditions, such as light exposure and water availability, highlighting the importance of considering both the whole life-history and environmental variation in fitness cost studies.

4.1 Seed germination and seedling emergence

We previously reported that fresh isolated seeds of feral radish were non-dormant, but that light and the pericarp reduced germination, especially under low temperatures[24], as was observed here (Fig. 1). This is consistent with the high emergence from buried isolated seeds and reduced emergence from seeds inside pods and seeds/pods on soil surface (light exposure) (Fig. 2). When seeds and pods were buried, seedlings emergence decreased with increasing burial depth, and this effect was greater when the seeds were inside pods. In darkness, no differences were found in seed germination between the R and S biotypes, isolated or inside pods, or in the emergence from buried seeds/pods. So, the darkness condition together with the release of the seeds from the pod by breaking the pericarp – e.g. during sowing or tillage up to 12-cm-deep – would promote germination and emergence of feral radish[24,30]. Under these conditions, there would be no differences in seedling emergence between the biotypes.

However, the light and irrigation conditions showed differential responses between the R and S biotypes. Light stimulated an increase in seed germination, whether isolated or inside pods, in the R biotype. Accordingly, the R biotype showed higher emergence than the S in isolated seeds placed on the soil surface. The increased emergence may have an adaptive importance in agricultural systems resembling no-till systems, widely used in Argentina (>90%)[31], where weed seeds are often exposed on the soil surface, e.g. after pod scarification by the harvester. The seeds inside pods of both biotypes germinated <2% on the soil surface at least five months after the beginning of the experiment (data not shown), suggesting the development of secondary dormancy, which could be crucial for seed persistence in the soil seedbank [24,32].

Under outdoor conditions, seedling emergence from isolated seeds was concentrated at the end of the summer, soon after seed or pod dispersal and the presence of the pericarp (seeds inside pods) reduced, spread and/or prolonged the seedling emergence period, as was suggested previously [24]. Seedling emergence from intact pods was mainly concentrated in autumn and spring in both biotypes, independent of water availability, possibly associated with medium/high temperatures, and with higher precipitations under rainfed conditions. In addition, high humidity and medium/high temperatures could have increased the growth potential of the embryo and/or reduced the mechanical resistance of the pericarp, due to its softening or

degradation by water and/or fungal activity [24,33,34]. No differences were found between the

biotypes in the seedling emergence pattern under irrigation, whereas the R biotype showed

delayed and reduced seedling emergence relative to the S under rainfed conditions. It has been

suggested that selection for late germinating seed cohorts is associated with herbicide resistance

(reviewed in Darmency *et al.*[9]), and although the genetic or physiological basis of this trait is

unknown, it could be linked to an increase in or development of dormancy [9,32]. For example, it

has been shown an increase in seed dormancy selected along with herbicide resistance in high-

intensity cropping systems[35,36]. The possibility that the effects observed here were caused by

environmental selection, such as cropping field vs. field margin[37], instead of the pleiotropic

effects of the Trp574Leu mutation cannot be ruled out. Additional experiments would be

necessary to check our inferences and the biological, ecological and evolutionary significances

of these results.

4.2 Plant fitness: Vegetative and reproductive traits

In the present study, R and S isolated individuals, or those under intra-/interspecific

competition, had similar relative growth rates and vegetative biomass. These results are in

agreement with two previous studies showing similar relative growth rates and competitive

resource response associated with the Trp574Leu mutation in *Raphanus raphanistrum*[12] and

Lolium rigidum[10]. Conversely, the R individuals showed a slight competitive disadvantage

during intraspecific competition with the S wild-type individuals. A similar result was found in

AHAS-resistant *Amaranthus powellii* with the Trp574Leu mutation[11]. However, in *A. powellii*

the relative crowding coefficient value was >6, much higher than that observed in our

experiment (RCC = 1.25). *A. powellii* with the Trp574Leu mutation showed strong pleiotropic

effects in leaf morphology and anatomy, which resulted in a significant reduction in biomass[11].

This alteration in leaf morphology and anatomy could occur because the Trp574Leu mutation

interferes with the normal C4 expression in developing leaves[11]. Nevertheless, *R. sativus* is a C3

species and to date, there is no evidence that the Trp574Leu mutation affects the expression of

leaf development.

Under field conditions, the R individuals grown with wheat interference had a lower above-

ground total biomass than the S wild-type with similar flowering time (pers. obs.), plant height

and resource allocation to reproductive traits (this latter was only slightly affected at the highest

weed density). This resulted in R individuals with a lower number of pods and seeds per plant,

lower seed biomass and less yield per plant compared with the S ones. These results are in

agreement with our previous study carried out in the absence of crop interference [19]. However,

the fitness cost on feral radish reproductive traits found under wheat interference was slightly

higher than that found in the absence of crop interference [19]. These studies reveal that fitness

costs associated with AHAS Trp574Leu mutant *R. sativus* plants are evident under a wide range of environmental conditions. The physiological mechanism responsible for fitness costs associated with the Trp574Leu mutation in *R. sativus* is likely to be associated with the adverse effect of this mutation on AHAS activity relative to the S wild-type [5,19]. Consequently, reduced AHAS activity due to the Trp574Leu mutation could be expressed under resource limiting conditions or in advanced stages of weed life-history. Accordingly, S individuals showed higher wheat yield losses than the R at both weed densities and planting dates. The Trp574Leu mutation also showed fitness cost in *A. powellii*[11] and *Amaranthus tuberculatus*[38].

4.3 Eco-evolutionary implications of AHAS herbicide resistance

The ecological and evolutionary implications of an adaptive mutation, e.g. the AHAS Trp574Leu mutation, depends on multiple interactions between operational factors related to herbicide application, genetic factors related to fitness benefits/costs and biological factors focused on weed ecology and life-history traits[5]. In AHAS herbicide-treated areas, it is expected that the Trp574Leu mutation increases their frequency [5,17]. However, our studies[19] suggest negative pleiotropic effects on the reproductive traits of feral radish imposed by the Trp574Leu mutation, which would reduce the frequency of the R allele in both ruderal and agrestal environments without AHAS herbicide selection[5,8,21]. In cross pollinated species, such as feral radish[16], gene flow between R and S populations could modify the dynamics of R alleles[39,40]. Gene flow from populations with null/low resistance level could act as a refuge, delaying resistance evolution in populations with a high level of herbicide selection and/or decreasing the frequency of R alleles in herbicide untreated areas. Accordingly, the importance of generating lower selection pressure is highlighted and also maintaining fence lines and fields margin without any herbicide application[40].

The eco-evolutionary consequences of the differential germination and emergence responses between R and S biotypes are environmentally dependent and complex to predict. It is the ecological environment that will determine the advantage or disadvantage of one biotype over the other during seed germination and seedling emergence (see Section 4.1). On the other hand, a persistent feral radish seedbank could act as a refuge, delaying the evolution of resistance or reducing R alleles. All this knowledge could be incorporated into herbicide resistance evolution models to explore the consequences of our results on the dynamics of resistance to AHAS herbicides and identify possible management strategies to prevent, delay or reverse weed resistance evolution.

These results should not be generalized because there could be an interaction between the Trp574Leu mutation and the genetic background[5,41], although a previous study comparing several R and S populations showed similar results[19]. The possibility that the effects observed

here were caused by genetic linkage or phenotypic correlations with other traits instead of the

pleiotropic effects of the Trp574Leu mutation and only correlated with it, cannot be completely

ruled out [8,9]. To reinforce these results, future research should focus on monitoring the

frequency of the AHAS-R Trp574Leu mutation for several years under field conditions, in the

absence of AHAS herbicide selection, using R and S biotypes.

ACKNOWLEDGEMENTS

This work was supported by grant ANPCYT-PICT 2012-2854. We thank the National Research

Council of Argentina (CONICET) for a fellowship to FH and RBV.

REFERENCES

Duggleby RG, McCourt JA, and Guddat LW, Structure and mechanism of inhibition of plant acetohydroxyacid synthase, *Plant Physiol Biochem* **46**:309–324 (2008).

Heap I, *The International Survey of Herbicide Resistant Weeds* [Online]. Available: http://www.weedscience.org [Accessed date: 5 February 2020].

Yu Q and Powles SB, Resistance to AHAS inhibitor herbicides: Current understanding, *Pest Manag Sci* **70**:1340–1350 (2014).

Tranel PJ, Wright T, and Heap IM, *Mutations in herbicide-resistant weeds to ALS inhibitors* [Online]. Available: http://www.weedscience.org [Accessed date: 5 February 2020].

Vila-Aiub MM, Fitness of herbicide-resistant weeds: Current knowledge and implications for management, *Plants* **8**:469 (2019).

Vila-Aiub MM, Neve P, and Powles SB, Fitness costs associated with evolved herbicide resistance genes in plants, *New Phytol* **184**:751–767 (2009).

Cousens RD and Fournier-Level A, Herbicide resistance costs: what are we actually measuring and why?, *Pest Manag Sci* **74**:1539–1546 (2018).

Baucom RS, Evolutionary and ecological insights from herbicide-resistant weeds: what have we learned about plant adaptation, and what is left to uncover?, *New Phytol* **223**:68–82 (2019).

Darmency H, Colbach N, and Le Corre V, Relationship between weed dormancy and herbicide rotations: implications in resistance evolution, *Pest Manag Sci* **73**:1994–1999 (2017).

Yu Q, Han H, Vila-Aiub MM, and Powles SB, AHAS herbicide resistance endowing mutations: Effect on AHAS functionality and plant growth, *J Exp Bot* **61**:3925–3934 (2010).

Tardif FJ, Rajcan I, and Costea M, A mutation in the herbicide target site acetohydroxyacid synthase produces morphological and structural alterations and reduces fitness in *Amaranthus powellii*, *New Phytol* **169**:251–264 (2006).

Li M, Yu Q, Han H, Vila-Aiub M, and Powles SB, ALS herbicide resistance mutations in *Raphanus raphanistrum*: Evaluation of pleiotropic effects on vegetative growth and ALS activity, *Pest Manag Sci* **69**:689–695 (2013).

Ashigh J and Tardif FJ, An amino acid substitution at position 205 of acetohydroxyacid synthase reduces fitness under optimal light in resistant populations of *Solanum ptychanthum*, *Weed Res* **49**:479–489 (2009).

Panozzo S, Scarabel L, Rosan V, and Sattin M, A new Ala-122-Asn amino acid change confers decreased fitness to ALS-resistant *Echinochloa crus-galli*, *Front Plant Sci* **8**:2042 (2017).

Vila-aiub MM, Gundel PE, Preston C, Experimental methods for estimation of plant fitness costs associated with herbicide-resistance genes, *Weed Sci* **63**:203–216 (2015).

Snow AA and Campbell LG, Can feral radishes become weeds?, in *Crop Ferality and Volunteerism: A Threat to Food Security in the Transgenic Era?,* ed. by Gressel J. Taylor & Francis, Boca Raton, FL, pp 193–207 (2005).

Pandolfo CE, Presotto A, Moreno F, Dossou I, Migasso JP, Sakima E, *et al.*, Broad resistance to acetohydroxyacid-synthase-inhibiting herbicides in feral radish (*Raphanus sativus* L.) populations from Argentina, *Pest Manag Sci* **72**:354–361 (2016).

Scursoni JA, Gigón R, Martín AN, Vigna M, Leguizamón ES, Istilart C, *et al.*, Changes in weed communities of spring wheat crops of Buenos Aires province of Argentina, *Weed Sci* **62**:51–62 (2014).

Vercellino RB, Pandolfo CE, Breccia G, Cantamutto M, and Presotto A, AHAS Trp574Leu substitution in *Raphanus sativus* L.: screening, enzyme activity and fitness cost, *Pest Manag Sci* **74**:1600–1607 (2018).

Vila-Aiub MM, Neve P, and Roux F, A unified approach to the estimation and interpretation of resistance costs in plants, *Heredity* **107**:386–394 (2011).

Keshtkar E, Zand E, Sasanfar H, Kudsk P, Dayan FE, Abdolshahi R, *et al.*, Assessing fitness costs from a herbicide-resistance management perspective: A review and insight, *Weed Sci* **67**:137–148 (2019).

Pandolfo CE, Presotto A, Poverene M, and Cantamutto M, Limited occurrence of resistant radish (*Raphanus sativus*) to AHAS-inhibiting herbicides in Argentina, *Planta Daninha* **31**:657–666 (2013).

Ellstrand NC, Devlin B, and Marshall DL, Gene flow by pollen into small populations: Data from experimental and natural stands of wild radish, *Proc Natl Acad Sci* **86**:9044–9047 (1989).

Vercellino RB, Pandolfo CE, Cerrota A, Cantamutto M, and Presotto A, The roles of

light and pericarp on seed dormancy and germination in feral *Raphanus sativus* (Brassicaceae), *Weed Res* **59**:396–406 (2019).

Vila-Aiub MM, Yu Q, Han H, and Powles SB, Effect of herbicide resistance endowing Ile-1781-Leu and Asp-2078-Gly ACCase gene mutations on ACCase kinetics and growth traits in *Lolium rigidum*, *J Exp Bot* **66**:4711–4718 (2015).

Hoffmann WA and Poorter H, Avoiding bias in calculations of relative growth rate, *Ann Bot* **90**:37–42 (2002).

Venus JC and Causton DR, Plant growth analysis: a re-examination of the methods of calculation of relative growth and net assimilation rates without using fitted functions, *Ann Bot* **43**:633–638 (1979).

Novak MG, Higley LG, Christianssen CA, and Rowley WA, Evaluating larval competition between *Aedes albopictus* and *A. triseriatus* (Diptera: Culicidae) through replacement series experiments, *Environ Entomol* **22**:311–318 (1993).

Weiner J, A neighborhood model of annual-plant interference., *Ecology* **63**:1237–1241 (1982).

Chauhan BS, Gill GS, and Preston C, Tillage system effects on weed ecology, herbicide activity and persistence: A review, *Aust J Exp Agric* **46**:1557–1570 (2006).

Scursoni JA, Vera ACD, Oreja FH, Kruk BC, and De La Fuente EB, Weed management practices in Argentina crops, *Weed Technol* **33**:459–463 (2019).

Baskin CC and Baskin JM, *Seeds: ecology, biogeography, and evolution of dormancy and germination, 2nd edn.* Elsevier Academic Press, San Diego, CA, USA (2014).

Sperber K, Steinbrecher T, Graeber K, Scherer G, Clausing S, Wiegand N, *et al.*, Fruit fracture biomechanics and the release of Lepidium didymum pericarp-imposed mechanical dormancy by fungi, *Nat Commun* **8**:1–11 (2017).

Steinbrecher T and Leubner-Metzger G, The biomechanics of seed germination, *J Exp Bot* **68**:765–783 (2017).

Owen MJ, Michael PJ, Renton M, Steadman KJ, and Powles SB, Towards large-scale prediction of *Lolium rigidum* emergence. II. Correlation between dormancy and herbicide resistance levels suggests an impact of cropping systems, *Weed Res* **51**:133–141 (2011).

Owen MJ, Goggin DE, and Powles SB, Intensive cropping systems select for greater seed dormancy and increased herbicide resistance levels in *Lolium rigidum* (annual

ryegrass), *Pest Manag Sci* **71**:966–971 (2015).

37 Kleemann SGL and Gill GS, Seed dormancy and seedling emergence in ripgut brome (*Bromus diandrus*) populations in Southern Australia, *Weed Sci* **61**:222–229 (2013).

38 Wu C, Davis AS, and Tranel PJ, Limited fitness costs of herbicide-resistance traits in *Amaranthus tuberculatus* facilitate resistance evolution, *Pest Manag Sci* **74**:293–301 (2018).

39 Roux F, Paris M, and Reboud X, Delaying weed adaptation to herbicide by environmental heterogeneity: a simulation approach, *Pest Manag Sci* **64**:16–29 (2008).

40 Maino JL, Renton M, Hoffmann AA, and Umina PA, Field margins provide a refuge for pest genes beneficial to resistance management, *J Pest Sci* **92**:1017–1026 (2019).

41 Paris M, Roux F, Bérard A, and Reboud X, The effects of the genetic background on herbicide resistance fitness cost and its associated dominance in *Arabidopsis thaliana*, *Heredity* **101**:499–506 (2008).

Legend:
- Resistant / Susceptible: 4/14 °C, 9/21 °C, 14/28 °C

A - Isolated seeds - Light/dark

Germination (%) vs Time (months)

B - Isolated seeds - Dark

Germination (%) vs Time (months)

C - Seeds inside pods - Light/dark

Germination (%) vs Time (months)

D - Seeds inside pods - Dark

Germination (%) vs Time (months)

Legend:
- R - Isolated seeds
- S - Isolated seeds
- R - Seeds inside pods
- S - Seeds inside pods

A - Soil surface (0 cm)

B - 1-cm depth

C - 8-cm depth

D - 12-cm depth

Days from planting

Cumulative emergence (%)

$Slope_S = 0.00151$
$Slope_R = 0.00148$
$P = 0.95$