Phosphate Induces a Morphological Shift that Enhances Vascular Dissemination of *Cryptococcus neoformans*

Steven T. Denham[1], Brianna Brammer[1], Krystal Y. Chung[1], Morgan A. Wambaugh[1], Joseph Bednarek[1], Li Guo[1], and Jessica C.S. Brown[1,2*]

[1]Division of Microbiology and Immunology, Department of Pathology, University of Utah, Salt Lake City, UT 84112, USA

[2]Lead Contact

[*]Correspondence: jessica.brown@path.utah.edu

SUMMARY:

Phenotypic heterogeneity is a common microbial strategy to improves fitness in fluctuating environments. The fungal pathogen *Cryptococcus neoformans* exhibits dramatic size heterogeneity: it varies from 2-100 μm in diameter during mammalian infection. Following pulmonary invasion, cells enlarge to >30 μm diameter, then decrease over disease course. Extrapulmonary organs, particularly the brain, contain uniformly small cells, implying that that morphotype is important for dissemination. To test this hypothesis, we isolated size-based *ex vivo* cell populations directly from mouse lungs. Small *ex vivo* cells readily disseminated compared with other *ex vivo* populations, small beads, and *in vitro*-grown small cells. The latter two groups are close in size to small *ex vivo* cells, suggesting that while size is important, fungal-specific elements also drive extrapulmonary dissemination. We found that mannose exposure facilitates host cell interaction and organ uptake. Phosphate induces small cell formation. This demonstrates how environmental cues shift phenotypic heterogeneity to drive pathogenesis.

INTRODUCTION:

Phenotypic heterogeneity within microbial populations promotes fitness in fluctuating and complex host environments (Weigel and Dersch, 2018). Heterogeneity arises among isogenic microbial populations as cells communicate, divide labor, and stochastically alter gene expression. Microbial responses to host microenvironments can also lead to phenotypic heterogeneity, even within single foci of infection (Davis, 2018). For example, microbial cells at the perimeter of infection foci may be more open to attack from immune cells, while the core may be depleted of nutrients and oxygen (Davis et al., 2015; Kowalski et al., 2019). Phenotypic heterogeneity can also influence which cells in a localized population are more likely to spread to new sites within a host, a phenomenon that significantly impacts the likelihood and trajectory of cancer metastasis (San Juan et al., 2019). However, due to the bottlenecks involved, we

understand little about how phenotypic heterogeneity influences the spread of infectious

microbes within a host.

In this study, we used the model human fungal pathogen *Cryptococcus neoformans* to

address the role that phenotypic heterogeneity plays in intra-host dissemination. A ubiquitous

saprophytic fungus associated with trees and bird guano (Emmons, 1955; Lazéra et al., 1996),

C. neoformans is generally harmless, with human exposure thought to be as high as 70% by

age five (Goldman et al., 2001). Infection follows inhalation of a fungal cell or spore (Maziarz

and Perfect, 2016), and in immunocompromised individuals *C. neoformans* disseminates via the

bloodstream and/or lymphatics to almost any organ in the body (Lin et al., 2015; Maziarz and

Perfect, 2016). *C. neoformans* cells rapidly proliferate in the central nervous system, and the

resulting fungal meningoencephalitis drives cryptococcosis patient mortality, killing ~180,000

people each year worldwide (Rajasingham et al., 2017).

C. neoformans is an excellent model organism to study the impact of phenotypic

heterogeneity on intra-host dissemination. Two of its primary virulence attributes—cell and

capsule size—vary considerably during infection and are readily observable via microscopy and

flow cytometry (Li and Nielsen, 2017). Isolates belonging to the *Cryptococcus* genus exhibit a

high degree of cell and capsule size diversity, and morphological heterogeneity strongly impacts

C. neoformans virulence: increased capacity for morphological variation among isolates

correlates with worse clinical outcome (Fernandes et al., 2018).

Cell and capsule size are finely tuned to a multitude of environmental signals and impact

stress resistance, immune recognition, and overall pathogenesis (Casadevall et al., 2019;

O'Meara and Alspaugh, 2012). Cells grown in nutrient replete laboratory media are

morphologically homogenous (5–10 µm in total (cell + capsule) diameter) and poorly

encapsulated (O'Meara and Alspaugh, 2012), while fungal cells proliferating in the lungs are

morphologically heterogeneous (1-100 µm in total diameter) and encapsulated, although

capsule thickness varies considerably (Wang and Lin, 2015; Zaragoza, 2011). In the host, the

polysaccharide capsule shields immunostimulatory microbe-associated molecular patterns in

the fungal cell wall, suppresses phagocytosis, and protects against oxidative stress (Casadevall

et al., 2019; O'Meara and Alspaugh, 2012).

The best understood *C. neoformans* morphotype is the polyploid (≥4C) "titan" cell,

measuring 10-100 µm in cell body diameter (Dambuza et al., 2018; Hommel et al., 2018;

Okagaki and Nielsen, 2012b; Okagaki et al., 2011; Trevijano-Contador et al., 2018; Zaragoza et

al., 2010; Zhou and Ballou, 2018). Titans are more resistant to phagocytosis and oxidative

stress and skew immunity toward a non-protective Th2 response (Crabtree et al., 2012; Okagaki
and Nielsen, 2012b; Okagaki et al., 2010).
While titans contribute considerably to cryptococcal pathogenesis, they are not the only
morphotype present during infection, accounting for a minority of cryptococcal cells within the
lungs (Zhou and Ballou, 2018). Tissue-specific differences in cryptococcal morphology also
manifest in human patients and animal infection models (Denham et al., 2017; Rivera et al.,
1998; Xie et al., 2012). Titan cells are rarely observed outside of the lungs, and overall cell and
capsule sizes tend to be substantially smaller in extrapulmonary sites, including the brain
(Charlier et al., 2005; Denham et al., 2017; Rivera et al., 1998; Xie et al., 2012). We previously
observed that median cell and capsule size within the lungs of mice inoculated with the highly
virulent KN99 strain of *C. neoformans* decreased from >30 μm to ~10 μm over the course of
infection, correlating with extrapulmonary dissemination (Denham et al., 2017). Thus, we
hypothesize that smaller morphotypes are more likely to disseminate.
Most infection models—including intravenous dissemination models—use *C.*
neoformans yeast cells collected from nutrient replete laboratory media as the inoculum, failing
to capture the phenotypic heterogeneity generated during lung infection (Mukaremera et al.,
2019; Ngamskulrungroj et al., 2012b; Sabiiti et al., 2012). In this study, we accounted for
phenotypic heterogeneity generated during lung infection by isolating cells directly from infected
lung tissue and sorted them according to size. We then intravenously challenged naïve mice
with those freshly isolated cryptococcal cells to determine the influence of morphology on
bloodstream dissemination.
90
91 **RESULTS:**
92
***Cryptococcus* strain-dependent morphological transitions during pulmonary infection**
correlate with dissemination.
Cryptococcus spp. isolates vary in their capacity for morphological variation (Fernandes
et al., 2018). To more broadly assess morphological shifts during infection, we inoculated ~8-
week-old female B6 mice with a panel of common *Cryptococcus* spp. reference strains (2.5×10^4
cells / mouse). At 3 days post-inoculation (dpi) and 17 dpi, we homogenized lungs and
measured fungal cell body and capsule size (**Figures 1A and 1B**). Morphogenesis varied
widely among *Cryptococcus* isolates. While KN99 (*C. neoformans*) and Bt63 (*C. neoformans*)
cells shifted smaller between 3 and 17 dpi, R265 (*C. deuterogattii*) and WM276 (*C. gattii*)
continued to increase in size—largely due to increased capsule thickness. Smaller median cell

size at 17 dpi correlated with increased extrapulmonary burden (**Figures 1C and 1D**). These
data suggest that among *Cryptococcus* isolates, the emergence of small cells (<10 μm) within
the lungs is a strong predictor of extrapulmonary dissemination.

We also measured cell and capsule size in the blood, liver, spleen, and brains of mice
inoculated with KN99 cells (**Figure 1E**). The range of cell size in the blood and liver was
surprisingly broad considering that murine microcapillaries can be as small as 3.5 μm. Although
rare, we did find fungal cells in the blood measuring 50 μm across. However, median total
diameter in the blood at 17 dpi was 12.72 μm, which was noticeably smaller than earlier
timepoints in the lungs (27.21 μm at 3 dpi and 18.71 μm at 14 dpi) indicating that entry into the
blood may still be a size-limiting bottleneck. In contrast, cell size in the spleen and brain was far
more restricted. The vast majority of cells in the spleen were <15 μm, while cells in the brain
rarely exceeded 10 μm in total diameter. These data suggest that the cell/capsule size profile
generated within the lungs is a strong predictor of extrapulmonary dissemination, and that
multiple bottlenecks may restrict vascular dissemination.

**Inoculum size alters the rate at which *C. neoformans* populations shift toward smaller
cell size in the lungs and disseminate to the brain.**

Lower initial cell density increases the frequency of titan cell formation *in vitro* (Trevijano-
Contador et al., 2018) and *in vivo* (Okagaki et al., 2010). We hypothesized that reducing the
number of cells in the inoculum would increase the frequency of larger cells early in lung
infection and slow the rate of small cell formation (**Figures S1A, S1B, and S1C**). We inoculated
mice with 10^3, 10^4, or 10^5 KN99-mCherry cells / mouse and employed the flow cytometry
forward scatter-area (FSC-A) parameter as a high-throughput assay to estimate relative
changes in cell size at 3, 7, 10, 14, and 17 dpi.

Consistent with our hypothesis, inoculating mice with fewer fungal cells increased fungal
cell size at 3 dpi, decreased the rate of cell size reduction in the lungs (**Figures S1A, S1B, and
S1C**), decreased lung fungal burden (**Figure S1D**), and decreased dissemination to the brain at
17 dpi (**Figure S1E**). Therefore, lower initial fungal cell density slows the rate at which fungal
cells shift smaller in the lungs and disseminate to the brain.

**Populations of cells isolated from the lungs according to size display differences in their
cell surface features and are differentially bound by soluble host factors.**

When we compared the labor-intensive microscopy-based measurements of cell size
with flow cytometry-based measurements, we still observed a fungal size shift over the course

of infection (**Figure S1A**). Thus, we sought to isolate KN99-mCherry cells from infected lung

tissue (14 dpi) via fluorescence-activated cell sorting (FACS) according to size (FSC-A),

characterize them, and directly test their ability to disseminate through the vasculature. We

sorted cross-sections of the fungal cell size distribution corresponding to the smallest (small *ex*

vivo: median total diameter of 7.56 μm), middle (mid *ex vivo*: median total diameter of 14.04

μm), and largest (large *ex vivo*: median total diameter of 21.80 μm) ~20% of the population

(**Figure 2A**). We further confirmed that we isolated distinct populations based on cell and

capsule size (**Figure 2B**) and that fungal cells sorted from each gate were also equivalently

viable (**Figure 2C**).

In addition to the differences in cell and capsule size among the three populations, we

observed differences in DNA ploidy measured by 4′,6-diamidino-2-phenylindole (DAPI) staining

(**Figure 2D**). Haploid cells cultured in minimal yeast nitrogen base (YNB) medium +/- benomyl

served as controls for 1C and 2C DNA content; benomyl inhibits cell division, trapping cells at

2C. Cells with >2C content are considered titan cells. Only the large *ex vivo* population

contained an appreciable number of titan cells (mean: 21.4% >2C) (**Figure S2**). Mid *ex vivo*

cells averaged 74.2% 1C and 22.8% 2C, while small *ex vivo* cells averaged 89.2% 1C and

10.0% 2C.

Morphological changes in *C. neoformans* are often accompanied by alterations in cell

surface architecture and immune recognition. We screened the binding rate (% total fungal cells

bound) of a panel of soluble receptors to small, mid, and large *ex vivo* cells (**Figure 2E**). Large

ex vivo cells were more likely to be bound by complement protein C3 than mid, or small *ex vivo*

cells. In contrast, small and mid *ex vivo* cells were more likely to be bound by lung surfactant

protein D (SPD). SPD binding to *C. neoformans* is deleterious for the host, potentially because

SPD agglutinates fungal cells and disrupts fungal clearance by macrophages (Geunes-Boyer et

al., 2012). We saw uniform and low-level binding of IgA and IgM class antibodies, and little to no

binding by mannose-binding lectins.

Differences in host factor binding suggested that cell surface architecture may differ

among the sorted populations. We used two lectins to probe the exposure of microbial features

in different layers of the cell wall: wheat germ agglutinin (WGA), which recognizes chitin in the

inner-most layers of the cryptococcal cell wall and concanavalin A, which binds mannose on

mannoproteins in the outer cell wall and inner capsule. There were no significant differences in

chitin exposure as measured by flow cytometry (**Figure 2F**). However, mannose was

significantly more exposed on small *ex vivo* cells compared with mid and large *ex vivo* cells and

cells grown in non-capsule inducing minimal YNB medium (**Figure 2G**).

We visualized mannose exposure pattern via microscopy and characterized staining as
negative, punctate (one or more distinct staining puncta), diffuse (staining covers the observable
cell surface), or both punctate and diffuse (**Figure 2H**). Mannose was more likely to be exposed
both at specific puncta and diffusely across the surface of small *ex vivo* cells. Mannose
exposure can correspond to recent bud scars, which may explain the punctate staining in our
analysis (Panepinto et al., 2007). The high prevalence of diffuse mannose exposure on small
cells may be due to their relatively thin capsules. Altogether, these results suggest that cell and
capsule size heterogeneity correlates with heterogeneity in cell surface architecture and
immune recognition.
180
Small *ex vivo* cells disseminate to extrapulmonary organs at a higher rate than mid and
large *ex vivo* cells.
To directly address the impact of fungal cell and capsule size on dissemination, we
intravenously inoculated naïve mice with small, mid, and large *ex vivo* cells (10^5 cells / mouse).
With blood flow from the tail vein running directly to the heart and then lungs for reoxygenation,
the lungs harbor the first major bed of microcapillaries fungal cells are likely to encounter
following tail vein inoculation. Therefore, sorting fungal cells from infected lung tissue prior to
tail vein inoculation roughly replicates the fate of fungal cells had they entered the bloodstream
from the lungs. Consistent with our hypothesis, a greater number of small *ex vivo* cells reached
the liver, spleen, and brain by 3 hours post-inoculation (hpi) than mid and large *ex vivo* cells
(**Figure 3A**). In contrast, the majority of mid and large *ex vivo* cells remained in the lungs at 3
hpi, possibly because they failed to navigate the lung microcapillaries as well as small *ex vivo*
cells. We observed the same trends in fungal burden at 3 dpi, indicating that small *ex vivo* cells
are well-capable of proliferating in extrapulmonary tissues (**Figure 3A**).
Although brain fungal burden was equivalent in mice inoculated with mid and large *ex*
vivo cells at 3 hpi, there were more mid *ex vivo* cells in the brain by 3 dpi. We asked whether the
reduced brain fungal burden in mice inoculated with large *ex vivo* cells was due to inefficient
blood-brain barrier crossing or reduced proliferation in the brain. To estimate blood-brain barrier
crossing *in vivo*, we intracardially perfused mice at 3 hpi to drive out circulating fungal cells and
normalized the fungal burden in perfused mice to nonperfused mice inoculated on the same
day. While perfusion did not significantly affect brain fungal burden in mice inoculated with small
or mid *ex vivo* cells, perfusion drove nearly all of the large *ex vivo* cells from the brain (**Figure**
3B). In contrast, fungal burden post-intracranial inoculation was independent of initial fungal cell

size (**Figure 3C**), suggesting that while large *ex vivo* cells are poorer at reaching the brain and

crossing the blood-brain barrier, they are not deficient at proliferating in the brain.

Macrophages, but not platelets or neutrophils, clear small *ex vivo* cells from the

vasculature to reduce brain dissemination.

As small *ex vivo* cells most efficiently disseminated to extrapulmonary organs, we

examined the mechanisms governing their dissemination profile more closely. Most small *ex*

vivo cells that made it past the lungs at 3 hpi were cleared in the liver and spleen (**Figure 3A**).

The liver plays a major role in clearing microbes and microbial components from the blood

(Macpherson et al., 2016). Recently, intravital imaging revealed that liver macrophages clear *C.*

neoformans cells from the blood stream (Sun et al., 2019). However, the inoculum for those

studies consisted of *C. neoformans* cells cultured *in vitro*. Our cell surface characterization

revealed that small *ex vivo* cells differ from cells cultured *in vitro* (YNB medium), mid *ex vivo*,

and large *ex vivo* cells, which may influence liver macrophage recognition of *C. neoformans*

cells.

To address the role of macrophages in clearing small *ex vivo* cells from the bloodstream,

we depleted macrophages from the liver, spleen, and bone marrow of mice with clodronate

liposomes 48 hours prior to inoculation (**Figures S3A and S3B**). Macrophage depletion reduced

small *ex vivo* cell fungal burden in the liver, while increasing fungal burden in the lungs, kidneys,

and brain at 1 dpi (**Figure 3D**). These results corroborate the findings of Sun *et al*, in that liver

macrophages play a critical role in filtering *C. neoformans* from the blood (Sun et al., 2019).

However, we also found that small *ex vivo* cells grow rapidly in the liver compared with *in vitro*-

grown cells or other *ex vivo* populations (**Figure 3E**), suggesting that attempts by the host to

clear *C. neoformans* cells by sequestering them in the liver may backfire.

Besides macrophages, platelets and neutrophils are among the most abundant immune

cell-types present in the blood that contribute to microbial clearance. Platelets can recognize

either complement protein bound to microbes or microbial features directly to initiate microbial

clearance, especially in the spleen (Broadley et al., 2016). Platelets don't appear to interact with

encapsulated *C. neoformans*, or mediate the clearance of *in vitro*-grown *C. neoformans* cells

from the blood (Sun et al., 2019). We also did not see any effect on fungal dissemination when

platelets were depleted prior to inoculation with small *ex vivo* cells (**Figures S4A, S4B, and**

S4C).

It's unclear whether neutrophils play a predominantly protective or detrimental role

during cryptococcosis, though there is evidence neutrophils can clear *C. neoformans* from the

blood (Sun et al., 2016; Zhang et al., 2016). In an intravenous infection model using *in vitro*-

grown fungal cells as the inoculum, neutrophil depletion greatly increased fungal burden in the

lungs and brain (Sun et al., 2016; Zhang et al., 2016). However, neutrophil-depletion prior to

inoculation with small *ex vivo* cells slightly increased lung fungal burden, but did not affect

fungal burden in other organs, indicating that the small *ex vivo* cells are more resistant to

neutrophil-mediated clearance than typical *in vitro*-grown fungal cells (**Figures S4D, S4E, and**

S4F).

Macrophage recognition of small *ex vivo* cells is partially mediated by mannose

exposure.

We predicted that liver macrophages may recognize the high levels of exposed

mannose on the surface of small *ex vivo* cells. We isolated liver macrophages from healthy ~8-

week-old B6 mice (Li et al., 2014) and co-cultured them with 6 µm plastic beads, minimal (YNB)

medium-grown cells, and sorted *ex vivo* populations at a multiplicity of infection of 2:1. Half of

the wells also received pretreatment with exogenous mannan, with the prediction that it would

reduce macrophage association when mannan recognition is critical. After 4 hours we washed

away free fungal cells, leaving only the fungal cells bound or internalized by macrophages,

which we define as "macrophage association".

Approximately 60% of macrophages had associated with 6 µm plastic beads by 4 hours,

indicating that the liver macrophages still displayed phagocytic capacity *ex vivo* (**Figure 3F**).

Additionally, 17.67% of liver macrophages associated with small *ex vivo* cells, which was more

than twice the frequency of mid and large *ex vivo* cells. Pretreatment with exogenous mannan

prior to infection reduced macrophage association with small *ex vivo* cells by ~40%, but did not

affect association with mid and large *ex vivo* cells, YNB-grown cells, or beads. These data

suggest that liver macrophages recognize small *ex vivo* cells in part through exposed mannose,

and that mannose exposure could contribute to fungal clearance in the liver.

Liver macrophages and other macrophages that sample the bloodstream are subject to

shear force, which can affect recognition and phagocytosis of microbes and other foreign matter

(Broadley et al., 2016). We assessed whether mannose exposure on small *ex vivo* cells affects

vascular clearance *in vivo* by intravenously injecting either 400 µg of exogenous mannan or

vehicle (PBS) 2-3 minutes prior to intravenous inoculation with 10^5 small *ex vivo* cells. 10

minutes later, we perfused mice to drive out non-recirculating fungal cells and plated organs for

colony forming units to estimate fungal burden. If mannose recognition contributed to fungal

clearance in a given organ, we expected to see reduced fungal burden in mice that received

exogenous mannan, compared with the vehicle control. We repeated this experimental design

with additional fungal cell wall and capsule components (GXM, β-glucan, chitin), as well as

polyinosinic acid (poly(I)), which is a scavenger receptor ligand (Pearson et al., 1993).

Although mannan did reduce fungal clearance in the liver and kidneys by ~20% and in

the spleen by ~75% (**Figure 3G**), it was not the only fungal ligand that influenced vascular

clearance. Poly(I) reduced clearance to a strikingly similar extent as mannan. The purified

capsule polysaccharide GXM reduced clearance in a similar extent in the liver, spleen, kidneys

at roughly the same level (~40% reduction). β-glucan and chitin, on the other hand, did not

appreciably affect fungal clearance, possibly because of their buried position beneath the

capsule. These data indicate that clearance of small ex vivo cells is not dependent on a single

cell surface ligand and that there are tissue-specific dependencies on various fungal features

that mediate vascular clearance.

Enhanced small *ex vivo* cell dissemination is not entirely dependent on size.

We next assessed whether small *ex vivo* cells disseminate more efficiently due to their

size alone, or if there were other contributing factors. We inoculated mice via the tail vein with

either large *ex vivo* cells, small *ex vivo* cells or size-matched inert polystyrene beads. Large *ex*

vivo cells disseminated similarly to size-matched 25 μm beads; the majority resided in the lungs

at 3 hpi (**Figure 4A**). Additionally, more 6 μm beads disseminated past the lungs to reach the

liver and spleen compared with 25 μm beads. However, small *ex vivo* cells reached the liver and

spleen at an even higher frequency, indicating that absolute size influences fungal

dissemination through the vasculature, but it is not the only contributing factor.

We next examined *in vitro* conditions that could reproduce the small *ex vivo* cell

morphotype and dissemination profile. Previously, we observed that cells grown in capsule-

inducing medium (CAP) shifted toward smaller cell and capsule size if they were supplemented

with conditioned medium (CM) from cells grown in non-capsule inducing medium (yeast

nitrogen base + 2% glucose, or YNB) (Denham et al., 2017). YNB medium contains

vitamins, nitrogen, and glucose but not amino acids. YNB-grown cells suppress capsule

formation, and are typically <10 μm in diameter. CAP medium consists of 10% Sabouraud's

dextrose broth—made from digested animal tissue—buffered to pH 7.4. CAP medium induces

capsule growth and is sufficient for cell division. We collected CM by filtering supernatant from

saturated YNB-grown cultures. In this case, CM consists of unspent nutrients and soluble fungal

factors secreted during growth in non-capsule inducing conditions.

We compared the dissemination profile of small *ex vivo* cells to cells grown in three *in*
vitro populations: (1) YNB-grown cells (2) CAP-grown cells, and (3) CAP-grown cells
supplemented with CM (CAP+CM). We also subjected the *in vitro*-grown populations to the
same sorting process as *ex vivo* cells by sorting YNB, and CAP+CM-grown cells from the
"small" FSC-A gate (YNB cells: 5.64 μm median total diameter; CAP+CM cells: 6.48 μm median
total diameter; few YNB or CAP+CM cells fell in the "mid" or "large" FSC-A gates) and sorting
the CAP-grown cells from the "mid" FSC-A gate (12.11 μm median total diameter; few CAP-
grown cells fell into the "small" or "large" FSC-A gates) (**Figure 4B**).
The prior culture conditions and morphology of cells in the inoculum heavily impacted
dissemination. Only CAP+CM cells replicated the dissemination profile of small *ex vivo* cells in
terms of fungal burden at 3 hours and 3 days post-inoculation (**Figure 4C**). Despite being
roughly the same size as small *ex vivo* cells, YNB-grown cells displayed similar fungal burden to
CAP-grown cells at 3 hours, and lower fungal burden in extrapulmonary organs than CAP,
CAP+CM, and small *ex vivo* cells at 3 days. Therefore, both absolute fungal cell size, and the
physiological state of cells entering the blood influence the outcome of vascular dissemination.
320

Cells sorted from infected host tissue or cells grown in host-like medium survive better
in the blood than cells grown in nutrient replete medium.
The caveat to using fungal burden as a readout for dissemination is that we can't easily
distinguish between the ability of cells to reach certain organs, and their ability to survive. We
hypothesized that YNB-grown cells may not be as primed for survival in the bloodstream as
cells sorted from infected host-tissue or cells grown under stressful host-like conditions (CAP
medium).
To address that hypothesis, we examined the survival of *in vitro*- and *ex vivo*-sorted cells
in whole mouse blood, liver macrophage co-culture (multiplicity of infection of 1:1), and DMEM
as a medium control (96-well plates, 37 °C, 5% CO_2, static). We assessed fungal cell survival as
a percentage of input at 0-, 24-, and 48-hour timepoints. Liver macrophages suppressed *in vitro*
and *ex vivo* cell growth equivalently (**Figures 4D and 4E**). However, the *ex vivo* populations and
in vitro populations from host-like conditions survived better in blood than YNB-grown cells
(**Figure 4F**), with large and mid-*ex vivo* populations faring the best. These results support the
idea that size and physiology are critical for bloodstream dissemination: small *ex vivo* cells
combine the stress resistance required for increased survival in the blood with the smaller cell
size required for widespread dissemination via microcapillaries.
338

Phosphate acquisition genes are upregulated in capsule-inducing medium and suppressed following conditioned medium supplementation.

CAP+CM-grown cells resemble small *ex vivo* cells in terms of cell + capsule size and vascular dissemination profile, so we hypothesized that they might suitably model small cell formation in the lungs. We analyzed the transcriptome of cells grown in YNB, CAP, and CAP+CM medium to identify genes that might facilitate the transition to small cell size in host-like capsule inducing conditions. We also analyzed the transcriptome of small, mid and large *ex vivo* cells.

Comparing the transcriptome of CAP, CAP+CM, and the *ex vivo* populations to YNB-grown cells demonstrated that the YNB-grown transcriptome is vastly different from each of these populations (**Figure 5A**). Core sets of 699 and 291 genes were up- and down-regulated respectively among the *ex vivo* populations and CAP/CAP+CM populations relative to YNB-grown cells. There were also 1085 up-regulated and 1297 downregulated genes shared only among the *ex vivo* populations relative to YNB-grown cells, demonstrating that the transcriptome within the host environment is still quite distinct from even the host-like *in vitro* conditions. Among the *ex vivo* populations, it is notable that small *ex vivo* cells were the most transcriptionally distinct (**Figure S5A**). 928 genes were differentially expressed in small vs mid *ex vivo* cells, and 2,365 genes were differentially expressed in small vs large *ex vivo* cells. For comparison, only 93 genes were differentially expressed in mid vs large *ex vivo* cells (**Figure S5A**).

Gene ontology term (GO-term) enrichment analysis revealed that phosphate ion transport was upregulated in CAP-grown cells vs YNB-grown cells (**Figure 5C**), and downregulated in CAP+CM vs CAP-grown cells (**Figure 5D**). In other words, phosphate acquisition genes are induced when cells enter capsule-inducing conditions (CAP medium), and then suppressed following supplementation with CM.

We observed the same trends when analyzing the expression of specific genes involved in phosphate acquisition and storage (**Figure 5E**). When *C. neoformans* cells are starved of phosphate, the phosphate acquisition transcription factor, Pho4, translocates to the nucleus where it promotes its own gene expression and the expression of genes involved in phosphate acquisition and storage. These include phosphate transporters: *PHO84*, *PHO89*, and *PHO840*; phosphatases: *APH1* (secreted/vacuolar acid phosphatase) and *APH4* (predicted intracellular acid phosphatase); and *VTC4* (vacuolar transport chaperone involved in processing polyphosphate). Pho4-dependent genes were upregulated in CAP vs YNB-grown cells and then suppressed in following CM supplementation—although many phosphate acquisition genes still

remained more highly expressed in CAP+CM-grown cells compared with YNB-grown cells.

Phosphate acquisition genes were also upregulated *in vivo* compared with YNB-grown cells,

suggesting that phosphate acquisition could be similarly linked to morphological shifts in the

host (**Figure 5E**).

Phosphate acquisition genes were also upregulated in the *ex vivo*-sorted populations

relative to YNB-grown cells, suggesting that the host lung environment is phosphate-limiting to a

degree. However, there were no differences in phosphate acquisition gene expression among

the three *ex vivo* populations. We limited our transcriptomic profiling of ex vivo cells to just one

timepoint (14 days post-inooculation). Had we profiled gene expression over time, we may have

observed changes more reflective of available phosphate concentrations in the lungs.

Phosphate and acidic pH are sufficient to drive *C. neoformans* populations toward

smaller morphotypes following growth in alkaline and host-like conditions.

We hypothesized that phosphate might be a critical nutrient that facilitates the

emergence of small cells in CAP+CM media. Conditioned medium consists of unspent YNB

medium and soluble fungal cell derived factors (Albuquerque et al., 2014). YNB medium is

defined, so we screened every component individually (including phosphate) to determine which

components are sufficient to induce small cell formation when cells are grown in CAP medium.

We also tested two soluble fungal-derived factors that accumulate during growth in YNB

medium: exo-GXM (Denham et al., 2017) and the quorum sensing-like peptide Qsp1 (Homer et

al., 2016; Lee et al., 2007).

We grew cells for 24 hours in YNB medium, then sub-cultured 10^5 cells/mL into CAP

medium for another 24 hours. We then sub-cultured CAP-grown cells 1:1 into fresh CAP

medium with 10% of the final volume being a YNB medium component, fungal-derived factor, or

water as the vehicle control. After a final 24-hour period, we measured cell size (FSC-A) by flow

cytometry.

Conditioned YNB medium and fresh YNB medium were sufficient to stimulate a shift

toward smaller cell size (**Figures 6A and 6B**). Among the YNB medium components, only

phosphate was sufficient to stimulate a shift toward smaller cell size (**Figure S5A**), and did so in

a concentration-dependent manner (**Figures 6A and 6B**). Phosphate-supplemented cells

decreased both total (cell+capsule) diameter (**Figure 6C**) as well as cell (**Figure S6A**) and

capsule (**Figure S6B**) size individually. Phosphate was also sufficient to induce a shift toward

smaller size for cells grown in titan cell-inducing medium (5% Sabouraud's, 10% fetal calf

serum, 15 µM sodium azide, pH 7.4, 5% CO_2, 37 °C) (**Figure S5B**).

We previously reported that exo-GXM may trigger the formation of smaller cells

(Denham et al., 2017). However, we were unable to reproduce those results (**Figure S5A**),

possibly due to differences in exo-GXM preparation, but more likely because we previously

solubilized exo-GXM in phosphate buffered saline. Qsp1 is a cell-density dependent signaling

peptide that suppresses the formation of titan cells (Trevijano-Contador et al., 2018). However,

the addition of Qsp1 did not affect cell size in this model (**Figures S5A and S5B**).

The Pho84 and Pho840 phosphate transporters are phosphate/H^+ symporters, and *C.*

neoformans therefore requires a proton gradient to efficiently transport phosphate into the cell

(Kretschmer et al., 2014). In alkaline pH, *C. neoformans* struggles to efficiently import

phosphate and upregulates phosphate acquisition machinery (Kretschmer et al., 2014; Lev et

al., 2017). *C. neoformans* releases metabolites that acidify the local microenvironment and

increase nutrient uptake (Himmelreich et al., 2003; Wright et al., 2002). For instance,

cryptococcomas in the brain acidify to levels as low as pH 5.5 (Himmelreich et al., 2003; Wright

et al., 2002).

We measured lung pH in mice inoculated with *C. neoformans* throughout infection, and

found that pH in the lungs decreased from ~7.4 to as low as ~6.3 between 3 and 17 dpi (**Figure**

6D). Phosphate did not induce significant shifts in FSC-A-measured cell size when CAP

medium was buffered to 6.3, potentially because phosphate availability in CAP medium buffered

to pH 6.3 is not as limiting as pH 7.4 (**Figures 6A and 6B**). However, we did measure a modest,

but significant decrease in total (cell+capsule) diameter when pH 6.3 CAP-grown cells were

supplemented with phosphate (**Figure 6C**), which was mostly due to a decrease in capsule

thickness (**Figure S5**). These data indicate that cell body and capsule size enlarge when

phosphate is limiting due to starvation, alkaline pH, or both. When phosphate availability

increases due to higher extracellular concentrations or decreased extracellular pH, *C.*

neoformans cells shift toward smaller morphotypes.

Limiting phosphate acquisition genetically suppresses morphogenesis *in vivo*.

Since phosphate was sufficient to trigger cell and capsule size reduction *in vitro*, we

hypothesized that limiting phosphate acquisition *in vivo* would alter morphogenesis, perhaps

delaying the appearance of small morphotypes. To test this hypothesis, we intranasally

inoculated mice with wild-type KN99, or either of two independently constructed KN99:*pho4Δ*

strains, which lack the gene encoding the transcription factor *pho4*. Without *pho4*, *C.*

neoformans cells fail to adequately upregulate genes involved in phosphate acquisition when

available phosphate is limiting, and become hypersensitive to alkaline conditions. The *pho4Δ*

mutant is attenuated for growth in the lungs but more severely attenuated for dissemination to
the brain, due to its poor survivability in alkaline blood (Lev et al., 2017). Our findings replicated
the work from Lev *et al.*, in that the *pho4Δ* mutant cells were attenuated for proliferation in the
lungs **(Figure 6E)** and dissemination to the brain **(Figure 6F)**. We also found that cell and
capsule size of *pho4Δ* mutant cells populating the lungs was static and less heterogeneous
relative to wild-type cells **(Figures 6G, S6C, and S6D)**. Most *pho4Δ* mutant cells remained
between 10 and 20 µm in total (cell + capsule) diameter over 17 days **(Figure 6G)**, with many of
them also exhibiting irregular budding **(Figures 6H)**. Total median wild-type (cell + capsule)
diameter was far larger than the *pho4Δ* mutant at day 3 (wild-type: 26.13 µm; *pho4Δ* #1: 14.55
µm; *pho4Δ* #2: 14.17 µm) and smaller than the *pho4Δ* mutant at day 17 (wild-type: 10.25 µm;
pho4Δ #1: 13.87 µm; *pho4Δ* #2: 15.10 µm), indicating that phosphate acquisition mediated by
pho4 is required for the full range of fungal morphogenesis during infection, including the shift
toward smaller morphotypes **(Figure 6G)**.
454
DISCUSSION:
We found that phenotypic heterogeneity is dynamic and supports *C. neoformans*
dissemination from the lungs. Capsule growth and titan cell formation during early stages of
lung infection enhance fungal resiliency, and as fungal proliferation continues, smaller cells with
greater potential for extrapulmonary dissemination emerge and/or survive. These smaller
morphotypes are phenotypically and transcriptionally distinct and can more easily navigate
microcapillaries to reach sites such as the brain. To our knowledge, these experiments were the
first to directly test the influence of *in vivo*-generated phenotypic heterogeneity on *C.*
neoformans dissemination, and the first to analyze the transcriptome of discrete *C. neoformans*
subpopulations infecting the lungs.
C. neoformans morphology is important at multiple stages of disease progression. For
example, inhalation of spores rather than yeast cells may accelerate dissemination, because
dendritic cells more readily phagocytose spores and quickly traffic them to draining lymph nodes
(Walsh et al., 2019). Spores exhibit elevated pathogenicity compared to genetically identical
yeast cells (Walsh et al., 2019). On yeast cells, the degree of encapsulation also altered the
trajectory of disease progression through initial interactions with macrophages in a zebrafish
infection model (Bojarczuk et al., 2016). Titan cells, in addition to being resistant to
phagocytosis, decrease the phagocytosis of smaller cells, potentially facilitating smaller cells'
dissemination (Okagaki and Nielsen, 2012a).

The dynamics of phenotypic heterogeneity *in vivo* varied widely among *Cryptococus*
isolates. Isolates that shifted toward smaller morphotypes (KN99 and Bt63) exhibited greater
extrapulmonary fungal burden than isolates that remained morphologically static (52D) or those
that shifted toward larger morphotypes as disease progressed (R265 and WM276). Absolute
size is likely not the sole reason for the difference in dissemination among these isolates. For
instance, R265 cells shifted larger over the course of lung infection, but R265 cells also die
more quickly in blood than KN99 cells (Ngamskulrungroj et al., 2012b). The capacity for
morphological heterogeneity is likely more important than any given morphological feature, such
as capsule thickness, cell body size, or titan formation, as different morphologies will hold
advantages at different stages of disease progression (Fernandes et al., 2018), allowing for a
bet hedging strategy during infection.
Our data also indicate that size alone does not explain the disseminating ability of small
ex vivo cells. Namely, cells originating from more stressful environments such as host lungs or
host-like *in vitro* conditions were better primed for survival and dissemination *in vivo*. Poorly
encapsulated small cells grown in nutrient replete YNB medium survived worse in blood and
failed to disseminate as well as small encapsulated *ex vivo* cells or small encapsulated cells
generated *in vitro* (CAP+CM) (**Fig. 4**). This suggests that efficient bloodstream dissemination
requires that cryptococcal cells adopt smaller cell and capsule morphologies, along with stress
resistance that will prime them for survival in the blood (Alanio et al., 2015; Ngamskulrungroj et
al., 2012a). These findings will be important to consider when optimizing future models of fungal
dissemination.
Cell size heterogeneity correlates with heterogeneity in cell surface architecture. Smaller
cell body size correlated with thinner capsule size and greater mannose exposure *in vivo*. Prior
data supporting the importance of mannose recognition by the host during cryptococcosis is
contradictory. Mannose receptor deficient mice are more susceptible to *C. neoformans* infection
(Dan et al., 2008). However, mannose recognition was dispensable for murine, but not primary
human, macrophage phagocytosis of *C. neoformans and C. gattii* cells *in vitro* (Lim et al., 2018).
There is also conflicting evidence as to whether mannose-binding lectin (MBL) deficiency
predisposes individuals to cryptococcal infection (Eisen et al., 2008; Fang et al., 2015). Despite
the increased mannose exposure on small *ex vivo* cells, we did not see significant MBL-binding
to cryptococcal cells. This may be due to the fact that MBL is most abundant in serum, and we
analyzed fungal cells from perfused lung tissue. We did find increased surfactant protein D
(SPD) binding of small and mid *ex vivo* cells compared with large *ex vivo* cells, and SPD can
bind to mannose-rich microbial features (Sahly et al., 2002). Our results indicate that the

importance of mannose recognition depends on a given fungal cell's morphotype: the addition of
exogenous mannan reduced macrophage association with small *ex vivo* cells, but not large *ex*
vivo cells or small YNB-grown cells. Increased mannose exposure may actually present a fungal
cell fitness cost that comes with reduced size. Smaller capsules and more recent bud scarring
may leave more mannose (and potentially other) residues exposed for recognition by
phagocytes.
Nutrient availability, particularly phosphate, is sufficient to shift toward smaller cell and
capsule morphologies. Out of all the YNB medium components, only phosphate was sufficient
to induce the shift toward smaller cell and capsule size *in vitro*. Also, suppressing phosphate
acquisition by deleting the *pho4* gene limited phenotypic heterogeneity *in vivo*: neither large nor
small morphotypes appeared in mouse lungs. Phosphate is essential for cell homeostasis and
growth, and its acquisition is tightly controlled in microbes that absorb fluctuating levels of
phosphate from their surroundings (Köhler et al., 2020; Lev and Djordjevic, 2018). Phosphate is
also critical for *Candida albicans* morphogenesis (hyphal formation), stress resistance, and
tissue invasion (Ikeh et al., 2016; Liu et al., 2018).
Phosphate acquisition is also essential for survival in the blood and dissemination to the
brain (Lev et al., 2017). *Cryptococcus* relies on a proton gradient to import many different
nutrients, including phosphate, into the cell (Lev and Djordjevic, 2018). Thus, *C. neoformans*
survives poorly in the alkaline (pH 7.4) bloodstream when its ability to upregulate phosphate
acquisition machinery is impaired (Lev et al., 2017). Our results suggest that increased
phosphate availability also impacts dissemination at an even earlier stage by mediating the
emergence of small *C. neoformans* morphotypes within the lungs. We observed an order of
magnitude decrease in lung pH over the course of infection, which could also enhance
acquisition of phosphate and other nutrients (Lev et al., 2017) or independently repress capsule
growth (Farhi et al., 1970; O'Meara and Alspaugh, 2012).
Since *pho4* mutants cannot form either large or small cells *in vivo*, it leads to the
question of whether small cells are daughters of large cells such as titans. Zhou *et al* (Zhou et
al., 2020) reported small cells, termed "titanides," that are titan daughter cells. We can obtain
small cells *in vitro* without inducing titan cells beforehand (**Fig. 6**). Since phosphate-induced
small cell formation is pH-dependent (**Fig. 6A-C**), there could be multiple pathways for small cell
formation. Moreover, cell division is required for small cell formation *in vitro*, since if we do not
add fresh growth medium when inducing small cells *in vitro*, the population size does not shift
(**Fig. 6B**). In other ways, titanides are similar to our "small" cells, particularly their hypothesized
roll in dissemination. Small cells are also about the same size as "drop" cells (Alanio et al.,

2015). These cells are of similar size as small cells but were isolated due to high non-
cytoplasmic levels of the glutathione stain CMFDA, which measures oxidative stress response,
and are not viable (Alanio et al., 2015).

Many studies identify fungal cell ploidy as a key driver of adaptation, in-host survival,
pathogenicity (Gerstein et al., 2015; Okagaki and Nielsen, 2012b; Selmecki et al., 2015). Of our
ex vivo populations, only large cells contain >2N cells (**Fig. S2**), and even within the large cell
population, the majority of cells are 2N, which could be either cells in the G2 phase of mitosis or
smaller titan cells. Medium cells display the DNA content profile of actively growing haploid
yeast cells (Todd et al., 2018). Thus, we think that comparisons between small and medium *ex*
vivo cells are likely independent of ploidy.

There are some notable limitations to our study. For example, we did not define the
subpopulations within the lungs based on clear biological criteria, but rather sorted cross-
sections of the population of fungal cells proliferating within the lungs. Small *ex vivo* cells
correspond with the size of disseminated cells in the brain, but medium and large cells are
based on arbitrary size cutoffs designed to distinguish them from small cells. Clearer definitions
of fungal subpopulations within the lungs and analysis of their dissemination would greatly
enhance our understanding of systemic cryptococcosis. For example, multiple groups have also
documented the appearance of unusually small "micro" (~1 μm), sometimes tear-dropped
shaped cells *in vivo* that are somewhat similar in shape to the small ex *vivo* cells in this study
(Feldmesser et al., 2001; Fernandes et al., 2018; Zaragoza, 2011). Properties of these "micro-
cells" are largely unknown and they could be very young, metabolically dormant, or aneuploid
cells (Dambuza et al., 2018; Feldmesser et al., 2001; Fernandes et al., 2018; Zaragoza, 2011).
Despite limited characterization, they likely represent the smallest cryptococcal cells seen
during infection. Though some of the cells sorted from infected lung tissue did present these
phenotypes, we did not attempt to further classify micro-cells.

Furthermore, *C. neoformans* faces multiple bottlenecks during dissemination, including
escaping the lungs. *C. neoformans* can escape the lung mucosa to access the bloodstream by
crossing lung epithelial cells, or via the Trojan horse mechanism within macrophages (Denham
and Brown, 2018). We focused on extrapulmonary organ entry rather than examining lung
escape by smaller cells directly. However, our small cells are more likely to be phagocytosed,
which may lead to increased Trojan horse dissemination. We also find that in late infection (17
dpi), the total diameter of cells in the blood shows a size distribution similar to the lungs (**Fig.**
1E), suggesting that pulmonary escape may be less of a bottleneck than previously thought.

Phosphate was sufficient to induce size reduction of *C. neoformans* cells. Since
phosphate is released from damaged cells (Gabow et al., 1982), we propose that small cells
could form as a progressing *C. neoformans* infection damages the host tissue, thus priming the
invading fungi for dissemination as tissue barriers are breaking down. Conditioned and fresh
YNB medium induced a greater size shift than phosphate alone, particularly at pH 6.3.
Combinations of different nutrients may also be sufficient to induce a size shift. Shifts toward
smaller cell and capsule size can be mediated by signals in addition to nutrient availability. For
example, Zhao *et al*. recently demonstrated that alleviation of genotoxic stress, such as that
seen during immune-mediated oxidative bursts, activates meiosis genes in polyploid titan cells
such that they begin to bud smaller haploid cells (Zhao et al., 2020).
Overall, we determine that a small, phosphate-inducible morphotype impacts a critical
step in *C. neoformans* pathogenesis: dissemination from the lungs to the brain. Our results are
consistent with observations in humans (Fernandes et al., 2018), in that phenotypic
heterogeneity strongly influences clinical outcome. Small *C. neoformans* cells are found in the
brains of human patients with cryptococcal meningitis (Xie et al., 2012). The emergence and/or
survival of smaller morphotypes within the lungs results in a subpopulation of fungal cells
primed for vascular dissemination that survival in the blood and are better able to enter organs
than their larger counterparts..
593

ACKNOWLEDGEMENTS:
This work was supported by research grant R01AI130248 from the National Institutes of
Health to J.C.S.B. S.T.D. was supported by grant T32AI055434 from the National Institutes of
Health. Flow cytometry work was supported by the University of Utah Flow Cytometry Facility
through the National Cancer Institute Award Number 5P30CA042014-24 and by the National
Center for Research Resources of the National Institutes of Health under Award Number
1S10RR026802-01. RNA sequencing and analysis was performed by the High-Throughput
Genomics and Bioinformatics Analysis Facility, a part of the Huntsman Cancer Institute at the
University of Utah. Generation of the *pho4Δ* #1 clone was constructed in the Madhani laboratory
with support from National Institutes of Health funding (R01AI100272). We would also like to
thank Dr. Kyla Ost for her valuable input regarding the analysis of opsonin-binding via flow
cytometry. We would also like to thank Dr. Matthew Rondina and Dr. Alicia Eustes for their
assistance with *in vivo* platelet depletion.

607
AUTHOR CONTRIBUTIONS:

S.T.D. and J.C.S.B. conceived and designed the experiments. S.T.D., B.B., K.Y.C., and

M.A.W. performed the experiments. S.T.D., B.B., and J.C.S.B. analyzed the data. S.T.D. and

J.C.S.B. wrote the paper. S.T.D., B.B., K.Y.C., M.A.W., and J.C.S.B. edited the paper.

DECLARATION OF INTERESTS:

The authors have no competing interests to declare.

Figure 1: Morphological transitions during pulmonary infections correlate with fungal

dissemination. (A) Total (cell+capsule) diameter, cell body diameter, and capsule thickness

measurements at 3 and 17 days post-intranasal inoculation (dpi) of mice (Mann-Whitney U test;

N=4 mice per timepoint, 57-65 fungal cells measured per mouse at 3 dpi and 117-266 fungal

cells per mouse at 17 dpi). Solid lines in the violin plots indicate the median and dotted lines

mark quartiles. **(B)** Representative India ink images of fungal cells quantified in 1A. **(C)** Fungal

burden in mice at 17 dpi (One-way ANOVA and uncorrected Fisher's LSD). The bar graph

represents the mean. **(D)** Nonlinear regression of mean extrapulmonary (liver+spleen+brain)

fungal burden and median total (cell+capsule) diameter. **(E)** Total diameter in KN99-infected

lungs at 3, 10, and 14 dpi and total diameter in KN99-infected organs at 17 dpi (Mann-Whitney

U-test; N=4 mice per timepoint; fungal cells measured per mouse in the lungs: 275, blood: 38,

liver: 72, spleen: 43, brain: 72). Solid lines in the violin plots indicate the median and dotted lines

mark quartiles. All panels in Figure 1: ns: not significant *$p<0.05$; **$p<0.01$; ***$p<0.001$;

****$p<0.0001$.

Figure 2: Sorting and characterization of fungal populations from infected lungs. (A)

FACS gating scheme for sorting KN99-mCherry cells from infected mouse lungs into

populations of "small", "mid", and "large" cells using forward scatter-area (FSC-A) to

approximate size. **(B)** Representative India ink images of cells sorted from "small", "mid", and

"large FSC-A gates and cell/capsule size measurements (N=97-120 fungal cells). Solid lines in

the violin plots indicate the median and dotted lines mark quartiles. **(C)** Cell viability post-sort

quantified by colony-forming unit (CFU) count (N=4). **(D)** Representative flow plot of DNA-

content measured with DAPI staining. YNB (minimal YNB medium). **(E)** Percentage of total cells

bound by complement (C3), surfactant protein D (SPD), mannose binding lectins A and C (MBL-

A; MBL-C), IgA, and IgM. **(F)** Representative flow plots and quantification of wheat-germ

agglutinin (WGA) stained cells to determine chitin exposure. **(G)** Representative flow plots and

quantification of concanavalin A (ConA) stained cells to determine mannose exposure. **(H)**

Representative images and characterization of mannose exposure pattern after staining with

ConA (N=50 fungal cells). All comparisons in Figure 2 were analyzed using one-way ANOVA

and uncorrected Fisher's LSD (ns: not significant; *$p<0.05$; **$p<0.01$; ***$p<0.001$; ****$p<0.0001$).

All bar graphs display the mean.

Figure 3: Small *ex vivo* cells disseminate to extrapulmonary organs at a higher rate than

larger cells, with mannose recognition by macrophages contributing to their clearance

from the vasculature. (A) Fungal burden, measured by colony forming unit counts (CFU), at 3

hours post-inoculation (hpi) and 3 days post-inoculation (dpi) in mice intravenously-inoculated

with small, mid, or large *ex vivo* cells (One-way ANOVA and uncorrected Fisher's LSD). **(B)**

Fungal burden in mice inoculated with sorted *ex vivo* fungal cells and perfused at 3 hpi. Fungal

burden was normalized to inoculated, but non-perfused control mice (Mann-Whitney U-test). **(C)**

Fungal burden at 3 dpi in mice intracranially inoculated with sorted *ex vivo* fungal cells (One-

way ANOVA and uncorrected Fisher's LSD). **(D)** Fungal burden at 1 dpi in mice pretreated with

either clodronate or vehicle control (PBS) liposomes and then inoculated with small *ex vivo* cells

and (Unpaired t test). **(E)** Fungal burden in the livers of mice (N=4-6) inoculated with *ex vivo*

fungal cells or fungal cells harvested from minimal YNB (YNB) medium at 3 and 72 hpi. Points

indicate the mean and the error bars indicate standard deviation. **(F)** Percentage of primary

cultured murine liver macrophages with physically associated plastic beads or fungal cells

(stained with Direct Yellow 96) after 4 hours. Dashed lines indicate 10-minute pretreatment with

exogenous mannan (One-way ANOVA and uncorrected Fisher's LSD; N=2263-3197 total

macrophages counted across 3 independent replicates). **(G)** Fungal burden in mice

intravenously treated with indicated ligand 2-3 minutes prior to intravenous inoculation with

small *ex vivo* fungal cells. Mice were perfused 10 minutes post-inoculation and fungal burden

was normalized to vehicle (PBS) treated controls (Unpaired t-test; N=6 mice per group). All

panels: ns: not significant *$p<0.05$; **$p<0.01$; ***$p<0.001$; ****$p<0.0001$. All bar graphs display

the mean.

Figure 4: Small cells generated under host-like conditions *in vitro* replicate the

dissemination profile of small *ex vivo* cells. (A) Percentage of total plastic beads or live cells

in the indicated organ 3 hours post-intravenous inoculation (hpi). **(B)** Representative India ink

images and quantification of cell and capsule size (N=100 fungal cells). YNB (minimal YNB

medium); CAP (capsule inducing medium); CAP+CM (capsule inducing medium supplemented

with 10% conditioned medium). **(C)** Fungal burden, measured by colony forming units (CFU), in

mice at 3 hpi. and 3 days post-inoculation (dpi). **(D)** Fungal cell survival in Dulbecco's modified

Eagle's medium (DMEM) and **(E)** liver macrophage co-culture (multiplicity of infection of 1:1)

measured as a percentage of input. **(F)** Fungal cell survival in mouse blood measured as a

percentage of input. All comparisons in Figure 4 were analyzed using one-way ANOVA and

uncorrected Fisher's LSD (*p<0.05; **p<0.01; ***p<0.001; ****p<0.0001). All bar graphs display

the mean.

Figure 5: Expression of phosphate acquisition genes correlate with cell and capsule size

changes *in vitro* and are upregulated *in vivo*. (A) Venn diagram of genes upregulated or

downregulated in different conditions relative to YNB-grown cells (YNB: minimal YNB medium;

CAP: capsule-inducing medium; CAP+CM: capsule inducing medium supplemented with 10%

conditioned medium). **(B)** Comparisons of the total number of differentially regulated genes

among *in vitro*-grown populations. **(C)** Top-ten most significantly enriched gene ontology (GO)

terms representing genes upregulated in CAP vs YNB-grown cells (-log10 (p-value)). Phosphate

ion transport was #46. **(D)** Top-ten most significantly enriched GO-terms representing genes

downregulated in CAP+CM vs CAP-grown cells (-log10 (p-value)). **(E)** Select phosphate

acquisition and storage RNA-seq gene expression relative to YNB-grown cells (adjusted p

values; *p<0.05; **p<0.01; ***p<0.001; ****p<0.0001). All bar graphs display the mean and error

bars the standard deviation.

Figure 6: Phosphate is sufficient to drive *C. neoformans* toward smaller morphotypes in

alkaline host-like conditions and is required for morphogenesis *in vivo*. (A) Cell size

measured using flow cytometry forward scatter (FSC-A) after growth in CAP-medium and

supplementation with the indicated factor (YNB: minimal YNB medium; CAP: capsule-inducing

medium; CAP+CM: capsule inducing medium supplemented with 10% conditioned medium).

FSC-A median fluorescence intensity (MFI) was normalized to control (10% H_2O)-treated cells

(one-way ANOVA and uncorrected Fisher's LSD). Bar graphs display the mean. **(B)**

Representative flow cytometry plots for conditions in Figure 6A. **(C)** Total (cell+capsule)

diameter measurements representing conditions in Figure 6A (Mann-Whitney U-test; N=200

fungal cells). Solid lines in the violin plots indicate the median and dotted lines mark quartiles.

(D) pH in *C. neoformans* strain KN99-infected and mock-infected mouse lungs at 3, 10, and 17

days post-inoculation (dpi) (N=4 mice; one-way ANOVA and uncorrected Fisher's LSD; error

bars represent standard deviation). All panels: ns: not significant *p<0.05; **p<0.01; ***p<0.001;

****p<0.0001. **(E)** Lung fungal burden in mice at 3, 10, and 17 dpi (One-way ANOVA and

uncorrected Fisher's LSD). The bar graph represents the mean. **(F)** Brain fungal burden in mice

at 17 dpi (One-way ANOVA and uncorrected Fisher's LSD). The bar graph represents the

mean. **(G)** Total (cell+capsule) diameter measurements at 3, 10, and 17 dpi of mice (Mann-

Whitney U test; N=6 mice per timepoint; 50, 100, and 150 wild-type cells measured per mouse

at days 3, 10, and 17 respectively. 50, 50, and 100 *pho4Δ* mutant cells measured per mouse at

days 3, 10, and 17 respectively). **(H)** Representative India ink images of fungal cells quantified

in 6G.

Supplemental Figure 1: Inoculum size alters the rate at which *C. neoformans* populations

shift toward smaller cell size in the lungs and disseminate to the brain. (A) We inoculated

mice with the indicated number of KN99-mCherry cells. The line graph shows the forward

scatter-area (FSC-A) measurements at the indicated days post-inoculation (dpi), which

estimates relative fungal size in the lungs (N=6 mice per timepoint). Points indicate the mean

and the error bars indicate standard deviation. **(B)** Representative flow cytometry plots from

mice inoculated with 10^4 KN99-mCherry cells / mouse. **(C)** Quantification of the area under the

curve for the curves displayed in Figure S1A. Bar graphs display the mean fluorescence

intensity and error bars indicate standard deviation. **(D)** Lung fungal burden in the same mice

described in Figure S1A and quantified by colony forming units (CFU). **(E)** Brain fungal burden

in mice at 17 dpi quantified by CFUs. Bar graphs display the mean. All comparisons in Figure

S1 were analyzed using one-way ANOVA and uncorrected Fisher's LSD (ns: not significant;

*p<0.05; **p<0.01; ***p<0.001; ****p<0.0001).

Supplemental Figure 2: DNA content of *ex vivo C. neoformans* populations. Percentage of

cells with 1C, 2C, and >2C DNA content in the indicated population. Values correspond to the

representative flow cytometry plot in Figure 2D. Bar graphs display the mean.

Supplemental Figure 3: Macrophage depletion efficiency in the liver and spleen. (A)

Representative flow cytometry plots displaying the frequency of CD45$^+$, F4/80$^+$ macrophages in

the liver and spleen after treatment with either PBS (control) liposomes or clodronate liposomes.

(B) Quantification of macrophage depletion measured as the ratio of macrophages in

clodronate-treated mice to PBS-treated mice.

Supplemental Figure 4: Depletion of platelets or neutrophils does not affect the

dissemination of small *ex vivo* cells. (A) Fungal burden in mice, measured by colony forming

units (CFU), following treatment with PBS (control) or anti-GPlbα antibody to deplete platelets

prior to intravenous inoculation with 10^5 small *ex vivo* cells / mouse. **(B)** Platelet count (K: units)

in the blood before and after treatment with PBS (control) or anti-GPIbα antibody. **(C)**

Quantification of platelet depletion in Figure S4B. **(D)** Fungal burden in mice treated with PBS

(control) or anti-Ly6G antibody to deplete neutrophils prior to intravenous inoculation with 10^5

small *ex vivo* cells / mouse. **(E)** Representative flow cytometry plots displaying the frequency of

CD45$^+$, Ly6C$^+$, Ly6G$^+$ neutrophils in the blood after treatment with either PBS (control) or anti-

Ly6G antibody. **(F)** Quantification of neutrophil depletion measured as the ratio of neutrophils in

anti-Ly6G antibody-treated mice to PBS-treated mice. All comparisons in Figure S4 were

analyzed using unpaired t-tests (ns: not significant *p<0.05; **p<0.01; ***p<0.001;

****p<0.0001). All bar graphs display the mean.

Supplemental Figure 6: Screen for factors that induce cell size reduction. (A) Fungal cell

size measured by forward scatter-area (FSC-A). Cells were cultured in capsule inducing

medium (CAP medium) buffered to pH 7.4 or 6.3 before subculture (1:1) in fresh medium with

10% of the final volume being the indicated supplement solubilized in H_2O (CM: conditioned

medium; GXM: glucuronoxylomannan; YNB: minimal YNB medium; PBS: phosphate buffered

saline). **(B)** Fungal cell size measured by forward scatter-area (FSC-A) of cells cultured in titan

cell medium and subcultured as described in Figure S5A. All comparisons in Figure S5 were

analyzed using one-way ANOVA and uncorrected Fisher's LSD (ns: not significant; *p<0.05;

**p<0.01; ***p<0.001; ****p<0.0001). All bar graphs display the mean and error bars indicate

standard deviation.

Supplemental Figure 5: Phosphate acquisition genes are upregulated in *ex vivo*-sorted

cells. (A) Comparisons of the total number of differentially regulated genes among *ex vivo*-

sorted populations. **(B)** Select phosphate acquisition and storage RNA-seq gene expression

relative to YNB-grown cells (adjusted p values; *p<0.05; **p<0.01; ***p<0.001; ****p<0.0001). All

bar graphs display the mean and error bars the standard deviation.

Supplemental Figure 7: Phosphate is sufficient to induce cell body diameter and capsule

thickness reduction in alkaline capsule-inducing medium and is required for

morphogenesis *in vivo*. (A) Cell body diameter and **(B)** capsule thickness of cells cultured in

capsule inducing medium (CAP medium) buffered to pH 7.4 or 6.3 before subculture (1:1) in

fresh medium with 10% of the final volume being the indicated supplement solubilized in H_2O

(CM: conditioned medium; YNB: minimal YNB medium). Solid lines in the violin plots indicate

the median and dotted lines mark quartiles. All comparisons in Figure S6 were analyzed using

Mann-Whitney U-tests; N=200 fungal cells (ns: not significant; *p<0.05; **p<0.01; ***p<0.001;

****p<0.0001). **(C)** Cell body diameter and **(D)** capsule thickness measurements at 3, 10, and

17 days post-intranasal inoculation (dpi) of mice (Mann-Whitney U test; N=6 mice per timepoint;

50, 100, and 150 wild-type cells measured per mouse at days 3, 10, and 17 respectively. 50, 50,

and 100 *pho4Δ* mutant cells measured per mouse at days 3, 10, and 17 respectively).

METHODS:

Fungal strains and growth media

The following strains were used in this study: KN99α (Brown lab stock which was also

the background for mutant strains), Bt63 (Brown lab stock, NCBI:txid1295841), 52D (ATCC

24067), R265 (ATCC MYA4093), WM276 (ATCC MYA-4071).

The following growth media were used in this study: YNB medium (yeast nitrogen base

without amino acids [Difco catalog no. 291940], 2% glucose), CAP medium (10% Sabouraud's

dextrose [Difco catalog no. 238230], buffered with 50 mM HEPES (pH 7.4) or MES (pH 6.3)),

Titan cell medium (5% Sabouraud's dextrose, 10% fetal calf serum (GenClone catalog no. 25-

550), 15 µM sodium azide, 50 mM HEPES (pH 7.4)).

To screen for factors that facilitated shifts in cell size, we picked single colonies from *C.*

neoformans cells streaked on YPAD agar and cultured them overnight (12-18 hours) in YNB

medium at 30 °C. We then sub-cultured those cells into YNB medium, CAP medium, or titan cell

medium for 24 hours at an initial cell density of 10^5 cells/mL. YNB- and CAP-grown cells were

cultured at 37 °C, while titan cell medium-grown cells were cultured at 37 °C and 5% CO_2. At

this stage, conditioned medium (CM) was collected from the 24-hour YNB cultures by pelleting

the fungal cells and filtering (0.22 µm pore size) the supernatant. 450 µl of the CAP and titan

cell media cultures were sub-cultured into 450 µl fresh media and 50 µl (10% of the final

volume) of one of the following supplements solubilized in H_2O: 10% H_2O (vehicle control), CM,

1% CM, 50 µg GXM, 50 µM Qsp1 peptide (Peptide 2.0: NFGAPGGAYPW), 50 µM Qsp1

scrambled peptide (Peptide 2.0: AWAGYFPGPNG), 10% 1XPBS (phosphate-buffered saline),

10% YNB medium; the following YNB medium supplements were added to a final concentration

that equaled their concentrations in YNB medium: 2% glucose, 4.531 nM folic acid, 8.186 nM

biotin, 250.6 nM copper sulfate pentahydrate, 531.4 nM riboflavin, 602.4 nM potassium iodide,

839.5 nM calcium pantothenate, 971.3 nM Sodium molybdate dihydrate, 1.186 µM thiamine

hydrochloride, 1.233 µM ferric chloride hexahydrate, 1.458 µM p-aminobenzoic acid, 1.945 µM

pyridoxine hydrochloride, 2.477 µM zinc sulfate heptahydrate, 2.649 µM manganese sulfate

monohydrate, 3.249 µM niacin, 8.087 µM boric acid, 11.10 µM inositol, 901.1 µM calcium

chloride dihydrate, 1.711 mM sodium chloride, 4.154 mM magnesium sulfate heptahydrate,

7.348 mM potassium phosphate monobasic, and 37.8mM ammonium sulfate. The

supplemented cultures were incubated for another 24 hours at 37 °C, following which 400 µl

was aliquoted to estimate cell size by flow cytometry and another 400 µl was aliquoted to

measure cell and capsule size (see "fungal cell size measurements") in India ink (Higgins

catalog no. 44201).

Mouse infection models and fungal burden analysis

825   For intranasal inoculations, ~8-week-old female C57BL/6NJ mice (Jackson Laboratory)

mice were anesthetized with ketamine/dexmedetomidine hydrochloride (Dexdomitor) delivered

intraperitoneally. They were then suspended by their front incisors on a horizontal strand of

thread. Unless otherwise indicated, mice were inoculated intranasally with 2.5×10^4

Cryptococcus cells in 50 µl of 1XPBS using a micropipette. The inoculum was placed dropwise

onto a nasal flare before being inhaled by mice. Ten minutes later, mice were intraperitoneally

administered the reversal agent atipamezole (Antisedan).

832   For intravenous inoculations, ~8-week-old female C57BL/6NJ mice (Jackson

Laboratory) were warmed under a heat lamp before being placed in a restraint. Mice were

inoculated via the tail vein with 10^5 *C. neoformans* cells or beads in 200 µl of 1XPBS using 28-

gauge x 12.7 mm syringes. In order to competitively inhibit host interactions with fungal

components *in vivo*, we administered 400 µg of GXM (see "GXM isolation), mannan (Sigma-

Aldrich catalog no. M7504), β-glucan (Millipore Sigma catalog no. 1048288), the chitin monomer

N-acetyl glucosamine (Vector Laboratories S-9002), or polyinosinic acid (Poly(I)) (Sigma-Aldrich

catalog no. 26936-41-4) in 200 µl of 1XPBS 2-3 minutes prior to inoculation with fungal cells.

840   For intracranial inoculations, ~6-week-old female C57BL/6NJ mice (Jackson

841   Laboratories) were anesthetized with ketamine/dexmedetomidine hydrochloride as described

842   above. They were inoculated intracranially with 10^3 *C. neoformans* cells in 30 µl of 1XPBS via a

843   26-gauge 1/2-inch needle. Following inoculation, mice were intraperitoneally administered the

844   reversal agent atipamezole (Antisedan; ~0.0125 mg/g).

845   Nonperfused mice were euthanized by CO2 asphyxiation and cervical dislocation. Mice

that were intracardially perfused at time of death were anesthetized with isoflurane and perfused

in a nonrecirculating fashion before cervical dislocation. Fungal burden was assessed by

excising organs and homogenizing them in 5 mL 1XPBS, washing the probe in between

samples: 30 seconds 10% bleach, 45 seconds 70% EtOH, and 10 seconds sterile H_2O. Ten-fold

serial dilutions of organ homogenate were plated on Sabouraud's agar containing 10 mg/ml

gentamicin and 100 mg/ml carbenicillin and stored at 30°C for 3 days before counting CFUs.

All animal procedures were approved by the University of Utah Institutional Animal Care

and Use Committee.

Fungal cell size measurements

C. neoformans cells harvested from laboratory growth medium or infected mouse lungs

were fixed in 2% paraformaldehyde for 20 minutes. To visualize capsule, 4 μl of India ink with 4

μl of cell suspension on a microscope slide. At least 10 successive images were taken starting

at one edge of the coverslip and moving across to the opposite side, as smaller cells tended to

drift to the edge of the coverslips. Total diameter was measured from one edge of the capsule to

the other. Cell body diameter was measured from one edge of the cell wall to the other. Capsule

thickness was calculated as follows: capsule thickness = (total diameter – cell body diameter) /

2. The total number of cells counted for a given experiment is indicated in the figure legends.

FACS isolation of fungal cells

Mice were intranasally inoculated with *C. neoformans* KN99-mCherry (see "Mouse

infection models"). At 14 dpi, lungs from 3 mice were excised and placed in 15 mL of 1XPBS.

The three lungs were homogenized together using a mechanical tissue homogenizer. The

homogenate was filtered first through a 70 μM cell strainer and then through a 40 μM cell

strainer to prevent downstream clogging of the flow cytometer. Cells were centrifuged at 2195xg

for 10 min. The supernatant was discarded and the pellet was resuspended in ~1mL 1XPBS.

The suspension was filtered once more through a 70 μM cell strainer and was resuspended at

~1x10^7 cells/mL in 1XPBS for FACS (BD FACS Aria). The FACS gating scheme is represented

in **Figure 2**.

Isolation of polystyrene beads from mouse organs

Mice were intravenously administered (see "Mouse infection models") 10^5 green

fluorescent polystyrene beads (Polysciences, Inc. catalog no. 17156-2 and 18241-2), and 3

hours later the indicated organs were harvested. Organs were homogenized with a mechanical

tissue homogenizer in 5 mL tissue lysis buffer (100 mM Tris-Cl pH 8.0, 200mM NaCl, 5mM

EDTA, 0.2% (w/v) SDS), washing the probe in between samples: 15 seconds H_2O, 15 seconds

70% EtOH, and 15 seconds fresh H_2O.

Beads from each organ were concentrated via differential centrifugation. The

homogenates were poured into a fresh 50 mL conical. The homogenization tubes were washed

with 30 mL of H_2O and combined with the respective homogenized sample. The samples were

filtered through cell strainers (70 µM pore size), after which another 10 mL of H_2O was passed

through each strainer. The samples were centrifuged at 4480xg for 40 minutes. The supernatant

was aspirated to the 5 mL mark and the samples were transferred to fresh 15 mL tubes

containing 2 mL of 100% Percoll (GE Healthcare catalog no. 17-0891-01). Water was added to

the 10 mL mark. The tubes were vortexed to mix and allowed to settle for 5 minutes before

centrifugation at 1120xg for 45 minutes. The supernatant was aspirated to the 2 mL mark. 10

mL of H2O was added to the samples before centrifuging at 4480xg for 20 minutes. The

supernatant was aspirated and resuspended in 200-1000 µl 1XPBS for flow cytometry. Entire

samples were analyzed to determine the number of beads per organ.

Fungal cell staining and flow cytometry analysis

Between 10^5 and 10^6 *C. neoformans* cells harvested from laboratory growth medium or

infected mouse lungs were pelleted and fixed in 2% paraformaldehyde for 20 minutes. Fixed

cells were pelleted and washed twice with 1XPBS. Cells were then resuspended in 1XPBS and

stained with the desired antibodies/lectins/fluorescent dye.

To estimate DNA content, cells were stained for 10 minutes (room temperature) in 0.3

µg/mL 1XPBS+0.1% triton X-100. Mid-log phase fungal cells grown in YNB medium +/- 80

µg/mL benomyl (Agilent catalog no. PST-1245) for 24 hours were used to set 1C and 2C gates.

Untreated mid-log phage fungal cells display two DAPI peaks (1C ad 2C), while benomyl

prevents cell division and traps cells at 2C.

To estimate fungal microbial associated molecular pattern exposure, cells were stained

at room temperature with 5 µg/mL fluorescein-conjugated wheat-germ agglutinin (WGA, Vector

Laboratories catalog no. FL-1021) for 15 minutes to detect exposed chitin or 50 µg/mL

fluorescein-conjugated concanavalin A (ConA, Vector Laboratories catalog no. FL-1001) for 5

minutes to detect exposed mannose.

To estimate the percentage of cells bound by soluble host factors, we used the following

antibodies: α-complement C3 (primary: mAb 11H9, ThermoFisher catalog no. MA1-40046;

secondary: Mouse anti-Rat IgG2a, FITC, Thermofisher catalog no. 11-4817-82), α-surfactant

protein D (primary: polyclonal, Abcam catalog no. ab203309; secondary: Donkey anti-Rabbit

IgG, AlexaFluor 405, Abcam catalog no. ab175651), α-mannose binding lectin A (primary: mAb

8G6, Hycult catalog no. HM1035; secondary: Mouse anti-Rat IgG2a, FITC, Thermofisher

catalog no. 11-4817-82), α-mannose binding lectin C (primary: mAb 14D12, Abcam catalog no.

ab106046; secondary: Mouse anti-Rat IgG2a, FITC, Thermofisher catalog no. 11-4817-82), α-

IgM (mAb II/41, FITC, BD Biosciences catalog no. 553437), α-IgA (mAb 11-44-2, FITC,

SouthernBiotech catalog no. 1165-02). Cells were stained with primary antibodies (1:50 dilution)

for 30 minutes on ice. When a secondary antibody was required, primary antibody labeled cells

were pelleted, washed twice with 1XPBS and stained with the secondary antibody (1:50 dilution)

for 20 minutes on ice.

After staining, cells were pelleted and washed twice with 1mL 1XPBS before being

resuspended in 200-400 μl of 1XPBS for flow cytometry.

GXM isolation

GXM was isolated as described previously (Wozniak and Levitz, 2009) with slight

modifications. *C. neoformans* (KN99) cells were grown in 100 mL for 3 days at 37°C in YNB

medium (YNB+2% glucose). Cells were centrifuged for 15 minutes at 12,000xg. The

supernatant was filtered (0.45 μM pore size) and 50 mL of supernatant was transferred to a 250

mL bottle. ~150 mL of 95% EtOH was added to the samples in 4-5 aliquots, mixing after each

addition. Samples were stored at 4°C overnight to precipitate GXM. Samples were centrifuged

for 30 minutes at 15,000xg, 4°C. The pellets were resuspended in 0.2M NaCl to a concentration

of about 10 mg/mL (~50 mL per 250 mL bottle). Samples were transferred to a clean beaker

stirred until samples were completely resuspended. 3 mg hexadecyltrimethylammonium

bromide (CTAB) per 1mg precipitate was slowly added with stirring and low heat. Once

solubilized, the samples were removed from heat and 250 mL of 0.05% CTAB was slowly

added with stirring. The samples were moved to fresh 250 mL bottles and centrifuged at for 2

hours at 11,000xg, 4°C. The supernatant was discarded and the pellet was washed in 150 mL

of 10% EtOH. The samples were centrifuge for 20 minutes at 15,000xg, room temperature. The

supernatant was discarded and the pellets were resuspended in 50 mL of 1M NaCl. The

samples were transferred to a clean beaker and stirred until completely resuspended.

Approximately 2 volumes of 95% EtOH was added dropwise with stirring to precipitate the GXM

while leaving the CTAB in solution. The samples were centrifuged in fresh 250 mL bottles for 20

minutes at 11,000xg, room temperature. The supernatant was discarded and the precipitate

was dissolved in 2M NaCl. The resulting solution was placed in a snake-skin dialysis cassette

with a 3,500 molecular weight cutoff (Thermo Fisher Scientific catalog no. 68035) and dialyzed

overnight against 1M NaCl. The samples were subsequently dialyzed against distilled H_2O for

14 days at 4°C. The distilled H_2O was changed every 1-2 hours for the first 2 days, and then 4-5

times a day for the next 6 days. For the last 2-6 days, the water was changed 2-3 times a day.

The dialyzed samples were lyophilized and stored at -80 °C.

Cell-type specific depletion in mice

Hepatic and splenic macrophages were depleted from mice using clodronate liposomes.

Mice were intravenously administered 200 µl of clodronate liposomes (Liposoma BV catalog no.

C-010) or PBS control liposomes (Liposoma BV catalog no. P-010) 48 hours prior to inoculation

with *C. neoformans* cells. Macrophage depletion efficiency in the liver and spleen was assessed

on the day of inoculation via flow cytometry. Macrophages were identified as CD45+ (anti-

CD45-efluor450, eBiosciences catalog no. 48-0451-82) and F4/80+ (anti-F4/80-APC,

eBiosciences catalog no. 17-4801-80).

Neutrophils were depleted from mice using anti-Ly6G antibodies. Mice were

intraperitoneally administered 200 ug of anti-Ly6G (clone 1A8 BioXcell catalog no. BP0075-1)

antibodies or PBS control at 24 and 2 hours prior to inoculation with *C. neoformans* cells.

Neutrophil depletion efficiency in the blood was assessed on the day of inoculation via flow

cytometry. Neutrophils were identified as CD45+ (anti-CD45-efluor450, eBiosciences catalog

no. 48-0451-82), CD11b+ (anti-CD11b-APC, eBiosciences catalog no. 17-0112-82), and Ly6G+

(anti-Ly6G-FITC, eBiosciences catalog no. 11-5931-82).

Platelets were depleted from mice using anti-GPIbα antibodies. Mice were intravenously

administered 80 µl of anti-GPIbα antibodies (Emfret catalog no. R300) or PBS control 24 hours

prior to inoculation with *C. neoformans* cells. Platelet depletion efficiency was determined by

Hemavet 950FS (Drew Scientific Group) analysis of blood collected by cheek bleed 10 minutes

prior to anti-GPIbα administration and on the day of inoculation.

Liver macrophage isolation and infection

Liver macrophages were isolated from 8- to 10-week-old female C57BL/6NJ mice

(Jackson Laboratory) according to established procedures (Li et al., 2014). Isolated liver

macrophages were resuspended in complete DMEM (Dulbecco's Modified Eagle's Medium,

GenClone catalog no. 25-500) supplemented with 10% fetal bovine serum (FBS, (GenClone

catalog no. 25-550) and 100 U/mL Penicillin/Streptomycin and seeded into a T-25 flask at a

density of 8-10 × 10^6 cells/flask. Macrophages were allowed to settle and adhere for 4 hours in a

mammalian tissue culture incubator (37 °C, 5% CO_2). Non-adherent cells were then removed

from the dish by gently washing 3 times with 1XPBS (without divalent cations), leaving adherent

macrophages. The media was replaced with 5 mL complete DMEM, and the cells were rested

for 4 days. The media was replaced on the day after initial seeding, and then as needed.

Fungal infection and association with liver macrophages was assessed using the

following procedures. The media was aspirated from rested liver macrophages and the cells

were washed once with 1XPBS. Macrophages were lifted from the plate by treating them with

1mL Accutase (Corning catalog no. 25-058-CI) for ~5 minutes, tapping the plate to dislodge

cells. 1 mL complete DMEM was added to the Accutase-treated cells, which were then

centrifuged at 300xg for 5 minutes. Liver macrophages were resuspended cells in 0.5-1 mL

complete DMEM and live cells were counted by mixing trypan blue (HyClone catalog no.

SV30084.01) with cell suspension at a 1:1 ratio and counting clear cells on a hemocytometer.

40,000 live cells/well were seeded into 96-well plates in complete DMEM. The liver

macrophages were allowed to settle and adhere overnight.

The next day, the inoculum was prepared by resuspending *C. neoformans* cells in 500 µl

of 0.01% Direct Yellow 96 (AK Scientific 61725-08-4), vortexing briefly to mix, and incubating for

5 minutes on the benchtop. *C. neoformans* cells stained with Direct Yellow 96 were pelleted,

washed once with 1 mL 1XPBS, and resuspended at a concentration of $8x10^6$ cells/mL in

serum-free DMEM. The supernatant was aspirated from the macrophages seeded in the 96-well

plates and was replaced with 200 µl of serum-free DMEM. 10 µl of the inoculum ($8x10^4$ cells)

was added to each well, resulting in a multiplicity of infection of 1:1. Macrophages and *C.

neoformans* cells were co-cultured for 4 hours in a mammalian tissue culture incubator (37 °C,

5% CO_2). The medium was then aspirated from each well, and the cells were washed three

times with 100 µl of 1XPBS to remove non-macrophage associated fungal cells. The remaining

cells were fixed in 100 µl of 4% paraformaldehyde for 10 min. The paraformaldehyde was then

removed and the cells were washed twice with 100 µl of 1XPBS and stained with100 µl of DAPI

(Sigma Aldrich catalog no. 28718-90-3) working solution (600 nM DAPI; 0.1% Triton X-100; in

1XPBS) for 5 minutes. The DAPI working solution was then removed and cells were washed

twice with 100 µl of 1XPBS. Finally, 100 µl of 1XPBS was added to the wells to prevent

desiccation. Cells were imaged on a Nikon widefield microscope, and the number of

macrophages with associated *C. neoformans* cell(s) were counted (green-fluorescent due to

Direct Yellow 96). Six images were taken per well in the same relative positions and every

imaged macrophage was scored as having a physically associated fungal cell(s) or not,

resulting in 2263-3197 macrophages counted per experimental condition across three

independent replicates.

In order to asses liver macrophage killing of *C. neoformans* cells, liver macrophages

were seeded in 96-well plates and infected with *C. neoformans* cells as described above. The

plates containing *C. neoformans* and macrophages were incubated for 24 or 48 hours. After the

desired time, the supernatant was collected from each well and transferred to a microcentrifuge

tube. 200 μl of sterile distilled water was then added to each well in order to lyse the

macrophages for 40 minutes at 37°C. The contents of each well were then mixed to collect

every *C. neoformans* cell and combined with the respective supernatant now in a

microcentrifuge tube. The wells were rinsed with 200 μl sterile PBS and added to the respective

microcentrifuge tube. Serial dilutions of each of the well contents were plated on YPAD agar

and incubated at 30°C for 2-3 days before counting CFUs. CFUs quantified at each timepoint

were normalized to the CFUs in the inoculum in order to calculate the "% of input". Each

macrophage killing experiment was repeated on three different days with each experimental

condition being performed in duplicate wells.

1030

Fungal survival in blood

We collected whole blood from 8- to 10-week-old female C57BL/6NJ mice

(Jackson Laboratory) to assess *C. neoformans* survival in blood. Mice were anesthetized with

isoflurane, and whole blood was collected by cardiac puncture using 28-gauge x 12.7 mm

syringes pre-coated with 0.5M EDTA. Blood was immediately transferred to tubes containing

heparin to achieve a final concentration of 30 units of heparin per mL of blood. 100 μl of blood

was added to individual wells of a 96-well plate. 100 *C. neoformans* cells in 5 ul of 1XPBS were

used to inoculate each well. The plates were incubated in a mammalian tissue culture incubator

(37 °C, 5% CO_2) without shaking. At the indicated timepoint, the well contents were transferred

to fresh microcentrifuge tubes. The wells were washed with 100 ul of 0.05M EDTA, which was

then added to the respective microcentrifuge tube. The entire contents of the microcentrifuge

tubes were plated on YPAD agar and incubated at 30°C for 2-3 days before counting CFUs.

CFUs quantified at each timepoint were normalized to the CFUs in the inoculum in order to

calculate the "% of input". Each experiment was repeated on three different days with each

experimental condition being performed in duplicate wells.

1046

RNA isolation, sequencing, and analysis

RNA sequencing data is publicly available at NCBI GEO (Accession number:

GSE152784). *C. neoformans* cells were harvested from laboratory media and from infected

mouse lungs for RNA isolation and sequencing. Three replicates were sequenced for each

sample with replicates being defined as follows. Approximately 10^7 cells were harvested from

the indicated *in vitro* growth condition on three separate days. At least 10^6 *ex vivo* cells were

isolated per sort gate each day (see "FACS isolation of fungal cells"). *Ex vivo* cells were isolated

from the pooled lung homogenate of three mice. *Ex vivo* sample populations were isolated on

two consecutive days and pooled for RNA isolation and sequencing, with the end result being

that each sequenced *ex vivo* replicate represents fungal cells isolated from pools of six mice.

RNA was isolated using phenol-chloroform extraction, followed by Qiagen RNeasy

column-based cleanup. Fungal cells were pelleted, flash frozen in liquid nitrogen, and

lyophilized until completely dry. Lyophilized cells were resuspended in 750 µl of TRIzol

(ThermoFisher Scientific catalog no. 15596026) inside screw-cap microcentrifuge tubes.

Approximately 50 µl of 1-mm and 150 µl of 0.5mm zirconium beads (BioSpec Products catalog

no. 11079110z and 11079105z) were added to each sample and placed in a Biospec Products

mini bead beater to lyse (12 x 2 min beating pulses, using sample blocks stored at -20 °C to

prevent overheating). 150 µl of chloroform was added to the lysed samples and incubated for 2-

3min at room temperature. Samples were centrifuged for 15 minutes at 12,000 x g, 4°C. The

aqueous phase was transferred to a fresh tube and combined with 5-10 µg RNase-free

glycogen as a carrier. 0.375 mL of isopropanol was added to each sample before incubating at

least ~20 minutes on ice. The samples were centrifuged for 15 minutes at 12,000 x g, 4°C. The

supernatant was discarded, and the isopropanol was removed by pipetting. The samples were

then briefly air-dried before resuspension in 350 µl of RLT buffer (Qiagen RNeasy) and 350 µl

70% EtOH. At this junction, samples were run through Qiagen's RNAeasy kit (Qiagen catalog

no. 74104) with on-column DNase treatment.

cDNA libraries were generated with the NEBNext Ultra II Directional RNA Library Prep

with rRNA Depletion Kit (New England BioLabs catalog no. E7760) by the High Throughput

Genomics Core at the University of Utah. Sample libraries were sequenced on the Illumina

NovaSeq 6000. One set of samples (primarily the *ex vivo* small, medium, and large cells) were

stored by the High Throughput Genomics Core at -80°C for six months prior to library

construction and showed more degradation products than replicates stored at -80°C for one

month. We therefore limited our analysis and interpretation of the data from these samples.

The *Cryptococcus neoformans* var. grubii H99 genome and gene feature files were

downloaded from Ensembl Fungi release 46 and the reference database was created using

STAR version 2.7.2c with splice junctions optimized for 150 base pair reads (Dobin et al., 2013).

Optical duplicates were removed from the paired end FASTQ files using clumpify v38.34 and

reads were trimmed of adapters using cutadapt 1.16 (Martin, 2011). The trimmed reads were

aligned to the reference database using STAR in two pass mode to output a BAM file sorted by

coordinates. Mapped reads were assigned to annotated genes in the gene feature file using

featureCounts version 1.6.3 (Liao et al., 2013). The output files from cutadapt, FastQC, Picard

CollectRnaSeqMetrics, STAR and featureCounts were summarized using MultiQC to check for

any sample outliers (Ewels et al., 2016). Differentially expressed genes were identified using a

5% false discovery rate with DESeq2 version 1.24.0 (Love et al., 2014). Differentially expressed

genes were run through FungiDB Gene Ontology (GO) Enrichment analysis, selecting for

"biological process" with a p-value cutoff of 0.05 (Basenko et al., 2018; Stajich et al., 2012).

Venn Diagrams were made with software from the Van de Peer group at Ghent University

(http://bioinformatics.psb.ugent.be/webtools/Venn/).

1095

Lung pH measurements

To measure lung tissue pH, mouse lungs were excised and immediately pierced with a

micro pH electrode (Orion (ThermoFisher Scientific) catalog no. 9863BN). The electrode was

completely embedded in the lung tissue before taking pH readings. Two readings were taken

per mouse. One reading was taken from the left lobe of the lungs and a second reading was

taken from one of the right lobes. These two readings were averaged to estimate lung pH for a

given mouse.

1103

Construction of the KN99:*pho4*Δ strains

The KN99:*pho4*Δ (*CNAG_06751*Δ) strains were constructed via homologous

recombination-based replacement of the *pho4* genomic sequence with a nourseothicin

resistance marker. The KN99:*pho4*Δ #1 strain was obtained from the KN99 gene deletion library

(Fungal Genetics Stock Center, 2015 Madhani plates). The KN99:*pho4*Δ #2 strain was

generated by PCR amplification of the nourseothicin resistance marker and ~1,000 base pair

flanking regions from the KN99:*pho4*Δ #1 strain (primers used to amplify the knock-out

construct: Forward: 5' AAAACGGCTGAAGGCTCGTTCT3', Reverse:

CTTCTGCAAGGTGAAGTTCACG). The resulting pho4 knock-out cassette was used to

transform wild-type KN99 cells via biolistic transformation as described previously (Davidson et

al., 2000). Putative transformants were PCR-verified by confirming the absence of the *pho4*

open reading frame (Primers to verify the absence of the open reading frame: Forward:

5'CCATCTCAGATACCAACTCGCC', Reverse: 5'CAATTTGCTGAGAGCCATAGGC3') and the

integration of the knock-out construct at the correct site (Primers to verify the correct 5'

integration site: Forward: 5'AGAGCTATATGGTATGACGAAC3', Reverse:

5'TGTGCTGATCATCCGATGCCAC3' and the correct 3' integration site: Forward

5'TGTGGAGGATGGTGGGGAATAG3', Reverse: 5'GGACGTGAGCCAATAAGTTCCT3'.

1121

Statistics

All statistical analyses, with the exception of RNA sequencing analysis, were performed

with GraphPad Prism 8 software. Hypothesis-driven experimental comparisons were analyzed

by unpaired t-test or one-way ANOVA and uncorrected Fisher's LSD if the data held a Gaussian

distribution and a Mann-Whitney U-test if the data were not Gaussian. *In vitro* experiments were

replicated three times unless otherwise stated. *In vivo* (murine infection) experiments were

replicated twice unless otherwise stated.

1129

REFERENCES:

Alanio, A., Vernel-Pauillac, F., Sturny-Leclère, A., and Dromer, F. (2015). *Cryptococcus*
neoformans Host Adaptation: Toward Biological Evidence of Dormancy. mBio *6*, e02580-
02514.
Albuquerque, P., Nicola, A.M., Nieves, E., Paes, H.C., Williamson, P.R., Silva-Pereira, I., and
Casadevall, A. (2014). Quorum sensing-mediated, cell density-dependent regulation of growth
and virulence in *Cryptococcus neoformans*. mBio *5*, e00986-00913.
Basenko, E.Y., Pulman, J.A., Shanmugasundram, A., Harb, O.S., Crouch, K., Starns, D.,
Warrenfeltz, S., Aurrecoechea, C., Stoeckert, C.J., Jr., Kissinger, J.C*., et al.* (2018). FungiDB: an
integrated bioinformatic resource for fungi and oomycetes. J Fungi *4*, 39.
Bojarczuk, A., Miller, K.A., Hotham, R., Lewis, A., Ogryzko, N.V., Kamuyango, A.A., Frost,
H., Gibson, R.H., Stillman, E., May, R.C*., et al.* (2016). *Cryptococcus neoformans* intracellular
proliferation and capsule size determines early macrophage control of infection. Sci Rep *6*,
21489.
Broadley, Steven P., Plaumann, A., Coletti, R., Lehmann, C., Wanisch, A., Seidlmeier, A., Esser,
K., Luo, S., Rämer, Patrick C., Massberg, S*., et al.* (2016). Dual-track clearance of circulating
bacteria balances rapid restoration of blood sterility with induction of adaptive immunity. Cell
Host Microbe *20*, 36-48.
Casadevall, A., Coelho, C., Cordero, R.J.B., Dragotakes, Q., Jung, E., Vij, R., and Wear, M.P.
(2019). The capsule of *Cryptococcus neoformans*. Virulence *10*, 822-831.
Charlier, C., Chrétien, F., Baudrimont, M., Mordelet, E., Lortholary, O., and Dromer, F. (2005).
Capsule structure changes associated with *Cryptococcus neoformans* crossing of the blood-brain
barrier. Am J Pathol *166*, 421-432.
Crabtree, J.N., Okagaki, L.H., Wiesner, D.L., Strain, A.K., Nielsen, J.N., and Nielsen, K. (2012).
Titan cell production enhances the virulence of *Cryptococcus neoformans*. Infect Immun *80*,
3776-3785.
Dambuza, I.M., Drake, T., Chapuis, A., Zhou, X., Correia, J., Taylor-Smith, L., LeGrave, N.,
Rasmussen, T., Fisher, M.C., Bicanic, T*., et al.* (2018). The *Cryptococcus neoformans* titan cell
is an inducible and regulated morphotype underlying pathogenesis. PLoS Pathog *14*, e1006978.
Dan, J.M., Kelly, R.M., Lee, C.K., and Levitz, S.M. (2008). Role of the mannose receptor in a
murine model of *Cryptococcus neoformans* infection. Infect Immun *76*, 2362-2367.

Davidson, R.C., Cruz, M.C., Sia, R.A.L., Allen, B., Alspaugh, J.A., and Heitman, J. (2000). Gene disruption by biolistic transformation in serotype D strains of Cryptococcus neoformans. Fungal Genetics and Biology *29*, 38-48.

Davis, K.M. (2018). All *Yersinia* are not created equal: phenotypic adaptation to distinct niches within mammalian tissues. Front Cell Infect Microbiol *8*, 261-261.

Davis, K.M., Mohammadi, S., and Isberg, R.R. (2015). Community behavior and spatial regulation within a bacterial microcolony in deep tissue sites serves to protect against host attack. Cell Host Microbe *17*, 21-31.

Denham, S.T., and Brown, J.C.S. (2018). Mechanisms of pulmonary escape and dissemination by *Cryptococcus neoformans*. J Fungi *4*, 25.

Denham, S.T., Verma, S., Reynolds, R.C., Worne, C.L., Daugherty, J.M., Lane, T.E., and Brown, J.C.S. (2017). Regulated release of cryptococcal polysaccharide drives virulence and suppresses immune infiltration into the central nervous system. Infect Immun.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15-21.

Eisen, D.P., Dean, M.M., O'Sullivan, M.V.N., Heatley, S., and Minchinton, R.M. (2008). Mannose-binding lectin deficiency does not appear to predispose to cryptococcosis in non-immunocompromised patients. Med Mycol *46*, 371-375.

Emmons, C.W. (1955). Saprophytic sources of *Cryptococcus neoformans* associated with the pigeon (COLUMBA LIVIA)1. Am J Epidemiol *62*, 227-232.

Ewels, P., Magnusson, M., Lundin, S., and Käller, M. (2016). MultiQC: summarize analysis results for multiple tools and samples in a single report. Bioinformatics *32*, 3047-3048.

Fang, W., Fa, Z., and Liao, W. (2015). Epidemiology of *Cryptococcus* and cryptococcosis in China. Fungal Genet Biol *78*, 7-15.

Farhi, F., Bulmer, G.S., and Tacker, J.R. (1970). *Cryptococcus neoformans* IV. The not-so-encapsulated yeast. Infect Immun *1*, 526-531.

Feldmesser, M., Kress, Y., and Casadevall, A. (2001). Dynamic changes in the morphology of *Cryptococcus neoformans* during murine pulmonary infection. Microbiol *147*, 2355-2365.

Fernandes, K.E., Brockway, A., Haverkamp, M., Cuomo, C.A., van Ogtrop, F., Perfect, J.R., and Carter, D.A. (2018). Phenotypic variability correlates with clinical outcome in *Cryptococcus* isolates obtained from Botswanan HIV/AIDS patients. mBio *9*, e02016-02018.

Gabow, P.A., Kaehny, W.D., and Kelleher, S.P. (1982). The spectrum of rhabdomyolysis. Medicine (Baltimore) *61*, 141-152.

Gerstein, A.C., Fu, M.S., Mukaremera, L., Li, Z., Ormerod, K.L., Fraser, J.A., Berman, J., and Nielsen, K. (2015). Polyploid Titan Cells Produce Haploid and Aneuploid Progeny To Promote Stress Adaptation. mBio *6*.

Geunes-Boyer, S., Beers, M.F., Perfect, J.R., Heitman, J., and Wright, J.R. (2012). Surfactant protein D facilitates *Cryptococcus neoformans* infection. Infect Immun *80*, 2444.

Goldman, D.L., Khine, H., Abadi, J., Lindenberg, D.J., Pirofski, L.-a., Niang, R., and Casadevall, A. (2001). Serologic evidence for *Cryptococcus neoformans* infection in early childhood. Pediatrics *107*, e66-e66.

Himmelreich, U., Allen, C., Dowd, S., Malik, R., Shehan, B.P., Mountford, C., and Sorrell, T.C. (2003). Identification of metabolites of importance in the pathogenesis of pulmonary cryptococcoma using nuclear magnetic resonance spectroscopy. Microb Infect *5*, 285-290.

Homer, C.M., Summers, D.K., Goranov, A.I., Clarke, S.C., Wiesner, D.L., Diedrich, J.K.,
Moresco, J.J., Toffaletti, D., Upadhya, R., Caradonna, I., *et al.* (2016). Intracellular action of a
secreted peptide required for fungal virulence. Cell Host Microbe *19*, 849-864.
Hommel, B., Mukaremera, L., Cordero, R.J.B., Coelho, C., Desjardins, C.A., Sturny-Leclère, A.,
Janbon, G., Perfect, J.R., Fraser, J.A., Casadevall, A., *et al.* (2018). Titan cells formation in
Cryptococcus neoformans is finely tuned by environmental conditions and modulated by positive
and negative genetic regulators. PLoS Pathog *14*, e1006982.
Ikeh, M.A.C., Kastora, S.L., Day, A.M., Herrero-de-Dios, C.M., Tarrant, E., Waldron, K.J.,
Banks, A.P., Bain, J.M., Lydall, D., Veal, E.A., *et al.* (2016). Pho4 mediates phosphate
acquisition in *Candida albicans* and is vital for stress resistance and metal homeostasis. Mol Biol
Cell *27*, 2784-2801.
Köhler, J.R., Acosta-Zaldívar, M., and Qi, W. (2020). Phosphate in virulence of *Candida*
albicans and *Candida glabrata*. J Fungi *6*, 40.
Kowalski, C.H., Kerkaert, J.D., Liu, K.-W., Bond, M.C., Hartmann, R., Nadell, C.D., Stajich,
J.E., and Cramer, R.A. (2019). Fungal biofilm morphology impacts hypoxia fitness and disease
progression. Nat Microbiol *4*, 2430-2441.
Kretschmer, M., Reiner, E., Hu, G., Tam, N., Oliveira, D.L., Caza, M., Yeon, J.H., Kim, J.,
Kastrup, C.J., Jung, W.H., *et al.* (2014). Defects in phosphate acquisition and storage influence
virulence of *Cryptococcus neoformans*. Infect Immun *82*, 2697-2712.
Lazéra, M.S., Pires, F.D.A., Camillo-Coura, L., Nishikawa, M.M., Bezerra, C.C.F., Trilles, L.,
and Wanke, B. (1996). Natural habitat of *Cryptococcus neoformans var. neoformans* in decaying
wood forming hollows in living trees. J M Vet Mycol *34*, 127-131.
Lee, H., Chang, Y.C., Nardone, G., and Kwon-Chung, K.J. (2007). TUP1 disruption in
Cryptococcus neoformans uncovers a peptide-mediated density-dependent growth phenomenon
that mimics quorum sensing. Mol Microbiol *64*, 591-601.
Lev, S., and Djordjevic, J.T. (2018). Why is a functional PHO pathway required by fungal
pathogens to disseminate within a phosphate-rich host: a paradox explained by alkaline pH-
simulated nutrient deprivation and expanded PHO pathway function. PLoS Pathog *14*, e1007021.
Lev, S., Kaufman-Francis, K., Desmarini, D., Juillard, P.G., Li, C., Stifter, S.A., Feng, C.G.,
Sorrell, T.C., Grau, G.E.R., Bahn, Y.-S., *et al.* (2017). Pho4 is essential for dissemination of
Cryptococcus neoformans to the host brain by promoting phosphate uptake and growth at
alkaline pH. mSphere *2*, e00381-00316.
Li, P.-z., Li, J.-z., Li, M., Gong, J.-p., and He, K. (2014). An efficient method to isolate and
culture mouse Kupffer cells. Immunol Lett *158*, 52-56.
Li, Z., and Nielsen, K. (2017). Morphology changes in human fungal pathogens upon interaction
with the host. J Fungi *3*, 66.
Liao, Y., Smyth, G.K., and Shi, W. (2013). featureCounts: an efficient general purpose program
for assigning sequence reads to genomic features. Bioinformatics *30*, 923-930.
Lim, J., Coates, C.J., Seoane, P.I., Garelnabi, M., Taylor-Smith, L.M., Monteith, P., Macleod,
C.L., Escaron, C.J., Brown, G.D., Hall, R.A., *et al.* (2018). Characterizing the mechanisms of
nonopsonic uptake of Cryptococci by macrophages. J Immunol *200*, 3539-3546.
Lin, Y.-Y., Shiau, S., and Fang, C.-T. (2015). Risk factors for invasive *Cryptococcus neoformans*
diseases: a case-control study. PloS One *10*, e0119090-e0119090.
Liu, N.-N., Uppuluri, P., Broggi, A., Besold, A., Ryman, K., Kambara, H., Solis, N., Lorenz, V.,
Qi, W., Acosta-Zaldívar, M., *et al.* (2018). Intersection of phosphate transport, oxidative stress
and TOR signalling in *Candida albicans* virulence. PLoS Pathog *14*, e1007076.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550.
Macpherson, A.J., Heikenwalder, M., and Ganal-Vonarburg, S.C. (2016). The liver at the nexus
of host-microbial interactions. Cell Host Microbe *20*, 561-571.
Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet *17*, 10-12.
Maziarz, E.K., and Perfect, J.R. (2016). Cryptococcosis. Infect Dis Clin North Am *30*, 179-206.
Mukaremera, L., McDonald, T.R., Nielsen, J.N., Molenaar, C.J., Akampurira, A., Schutz, C.,
Taseera, K., Muzoora, C., Meintjes, G., Meya, D.B.*, et al.* (2019). The mouse inhalation model
of *Cryptococcus neoformans* infection recapitulates strain virulence in humans and shows that
closely related strains can possess differential virulence. Infect Immun *87*, e00046-00019.
Ngamskulrungroj, P., Chang, Y., Sionov, E., and Kwon-Chung, K.J. (2012a). The Primary
Target Organ of Cryptococcus gattii Is Different from That of Cryptococcus neoformans in a
Murine Model. *3*, e00103-00112.
Ngamskulrungroj, P., Chang, Y., Sionov, E., and Kwon-Chung, K.J. (2012b). The primary target
organ of *Cryptococcus gattii* is different from that of *Cryptococcus neoformans* in a murine
model. mBio *3*, e00103-00112.
O'Meara, T.R., and Alspaugh, J.A. (2012). The *Cryptococcus neoformans* capsule: a sword and a
shield. Clin Microbiol Rev *25*, 387-408.
Okagaki, L.H., and Nielsen, K. (2012a). Titan Cells Confer Protection from Phagocytosis in
Cryptococcus neoformans Infections. *11*, 820-826.
Okagaki, L.H., and Nielsen, K. (2012b). Titan cells confer protection from phagocytosis in
Cryptococcus neoformans infections. Eukaryot Cell *11*, 820-826.
Okagaki, L.H., Strain, A.K., Nielsen, J.N., Charlier, C., Baltes, N.J., Chrétien, F., Heitman, J.,
Dromer, F., and Nielsen, K. (2010). Cryptococcal cell morphology affects host cell interactions
and pathogenicity. PLoS Pathog *6*, e1000953.
Okagaki, L.H., Wang, Y., Ballou, E.R., O'Meara, T.R., Bahn, Y.-S., Alspaugh, J.A., Xue, C., and
Nielsen, K. (2011). Cryptococcal titan cell formation is regulated by G-protein signaling in
response to multiple stimuli. Eukaryot Cell *10*, 1306-1316.
Panepinto, J.C., Komperda, K.W., Hacham, M., Shin, S., Liu, X., and Williamson, P.R. (2007).
Binding of serum mannan binding lectin to a cell integrity-defective *Cryptococcus neoformans*
ccr4Delta mutant. Infect Immun *75*, 4769-4779.
Pearson, A.M., Rich, A., and Krieger, M. (1993). Polynucleotide binding to macrophage
scavenger receptors depends on the formation of base-quartet-stabilized four-stranded helices. J
Biol Chem *268*, 3546-3554.
Rajasingham, R., Smith, R.M., Park, B.J., Jarvis, J.N., Govender, N.P., Chiller, T.M., Denning,
D.W., Loyse, A., and Boulware, D.R. (2017). Global burden of disease of HIV-associated
cryptococcal meningitis: an updated analysis. Lancet Infect Dis *17*, 873-881.
Rivera, J., Feldmesser, M., Cammer, M., and Casadevall, A. (1998). Organ-dependent variation
of capsule thickness in *Cryptococcus neoformans* during experimental murine infection. Infect
Immun *66*, 5027-5030.
Sabiiti, W., May, R.C., and Pursall, E.R. (2012). Experimental models of cryptococcosis. Int J
Microbiol *2012*, 626745-626745.
Sahly, H., Ofek, I., Podschun, R., Brade, H., He, Y., Ullmann, U., and Crouch, E. (2002).
Surfactant protein D binds selectively to lipopolysaccharides containing mannose-rich O-
antigens. J Immunol *169*, 3267.

San Juan, B.P., Garcia-Leon, M.J., Rangel, L., Goetz, J.G., and Chaffer, C.L. (2019). The complexities of metastasis. Cancers (Basel) *11*, 1575.

Selmecki, A.M., Maruvka, Y.E., Richmond, P.A., Guillet, M., Shoresh, N., Sorenson, A.L., De, S., Kishony, R., Michor, F., Dowell, R.*, et al.* (2015). Polyploidy can drive rapid adaptation in yeast. Nature *519*, 349-352.

Stajich, J.E., Harris, T., Brunk, B.P., Brestelli, J., Fischer, S., Harb, O.S., Kissinger, J.C., Li, W., Nayak, V., Pinney, D.F.*, et al.* (2012). FungiDB: an integrated functional genomics database for fungi. Nucleic Acids Res *40*, D675-D681.

Sun, D., Sun, P., Li, H., Zhang, M., Liu, G., Strickland, A.B., Chen, Y., Fu, Y., Xu, J., Yosri, M.*, et al.* (2019). Fungal dissemination is limited by liver macrophage filtration of the blood. Nat Commun *10*, 4566.

Sun, D., Zhang, M., Liu, G., Wu, H., Li, C., Zhou, H., Zhang, X., and Shi, M. (2016). Intravascular clearance of disseminating *Cryptococcus neoformans* in the brain can be improved by enhancing neutrophil recruitment in mice. Eur J Immunol *46*, 1704-1714.

Todd, R.T., Braverman, A.L., and Selmecki, A. (2018). Flow Cytometry Analysis of Fungal Ploidy. Curr Protoc Microbiol *50*, e58-e58.

Trevijano-Contador, N., de Oliveira, H.C., García-Rodas, R., Rossi, S.A., Llorente, I., Zaballos, Á., Janbon, G., Ariño, J., and Zaragoza, Ó. (2018). *Cryptococcus neoformans* can form titan-like cells in vitro in response to multiple signals. PLoS Pathog *14*, e1007007.

Walsh, N.M., Botts, M.R., McDermott, A.J., Ortiz, S.C., Wüthrich, M., Klein, B., and Hull, C.M. (2019). Infectious particle identity determines dissemination and disease outcome for the inhaled human fungal pathogen *Cryptococcus*. PLoS Pathog *15*, e1007777-e1007777.

Wang, L., and Lin, X. (2015). The morphotype heterogeneity in *Cryptococcus neoformans*. Curr Opin Microbiol *26*, 60-64.

Weigel, W.A., and Dersch, P. (2018). Phenotypic heterogeneity: a bacterial virulence strategy. Microb Infect *20*, 570-577.

Wozniak, K.L., and Levitz, S.M. (2009). Isolation and purification of antigenic components of *Cryptococcus*. Methods Mol Biol *470*, 71-83.

Wright, L., Bubb, W., Davidson, J., Santangelo, R., Krockenberger, M., Himmelreich, U., and Sorrell, T. (2002). Metabolites released by *Cryptococcus neoformans var. neoformans* and *var. gattii* differentially affect human neutrophil function. Microb Infect *4*, 1427-1438.

Xie, S., Sao, R., Braun, A., and Bottone, E.J. (2012). Difference in *Cryptococcus neoformans* cellular and capsule size in sequential pulmonary and meningeal infection: a postmortem study. Diagn Microbiol Infect Dis *73*, 49-52.

Zaragoza, O. (2011). Multiple disguises for the same party: the concepts of morphogenesis and phenotypic variations in *Cryptococcus neoformans*. Front Microbiol *2*, 181-181.

Zaragoza, O., García-Rodas, R., Nosanchuk, J.D., Cuenca-Estrella, M., Rodríguez-Tudela, J.L., and Casadevall, A. (2010). Fungal cell gigantism during mammalian infection. PLoS Pathog *6*, e1000945.

Zhang, M., Sun, D., Liu, G., Wu, H., Zhou, H., and Shi, M. (2016). Real-time *in vivo* imaging reveals the ability of neutrophils to remove *Cryptococcus neoformans* directly from the brain vasculature. J Leukoc Biol *99*, 467-473.

Zhao, Y., Wang, Y., Upadhyay, S., Xue, C., and Lin, X. (2020). Activation of meiotic genes mediates ploidy reduction during cryptococcal infection. Curr Biol.

Zhou, X., and Ballou, E.R. (2018). The *Cryptococcus neoformans* titan cell: from *in vivo* phenomenon to *in vitro* model. Curr Clin Microbiol Rep *5*, 252-260.

1344   Zhou, X., Zafar, H., Sephton-Clark, P., Mohamed, S.H., Chapuis, A., Makarova, M.,
1345   MacCallum, D.M., Drummond, R.A., Dambuza, I.M., and Ballou, E.R. (2020). Host
1346   environmental conditions induce small fungal cell size and alter population heterogeneity in
1347   *Cryptococcus neoformans*. bioRxiv, 2020.2001.2003.894709.
1348

Figure 1

Figure 2

A. Gating scheme:

C. neoformans-mCherry

Single Cells

FSC-A Sort Gates
small: 21.6% mid: 23.9% large: 20%

C. Cell viability
ns
small *ex vivo*
mid *ex vivo*
large *ex vivo*

B.
small *ex vivo*   mid *ex vivo*   large *ex vivo*
50µm   50µm   50µm

Total (cell+capsule) diameter
****
small *ex vivo*
mid *ex vivo*
large *ex vivo*

Cell body diameter
****

Capsule thickness
****

D.
1C 2C >2C
YNB-grown
YNB+benomyl
small *ex vivo*
mid *ex vivo*
large *ex vivo*
DAPI

E. Opsonization
***
small *ex vivo*
mid *ex vivo*
large *ex vivo*
C3 SPD MBL-A MBL-C IgA IgM

F. Chitin exposure
unstained
YNB-grown
small *ex vivo*
mid *ex vivo*
large *ex vivo*
WGA (FITC) / FSC-A

G. Mannose exposure
unstained
YNB-grown
small *ex vivo*
mid *ex vivo*
large *ex vivo*
ConA (FITC) / FSC-A

H. ConA-FITC / DIC merge
small *ex vivo* — diffuse
mid *ex vivo* — punctate
large *ex vivo*
10 µm

Chitin exposure
ns

Mannose exposure
**

Mannose exposure pattern
both
diffuse
punctate
negative
small *ex vivo*
mid *ex vivo*
large *ex vivo*

Figure 4

Figure 5

A.

Upregulated vs MIN

Downregulated vs MIN

B.

C.

Upregulated in CAP vs YNB

D.

Downregulated in CAP+CM vs CAP

E.

Figure 6