Original Article

Interference of feral radish (*Raphanus sativus*) resistant to AHAS-inhibiting herbicides in

oilseed rape, wheat and sunflower crops

Roman B. Vercellino[1], Claudio E. Pandolfo[1], Miguel Cantamutto[2], Alejandro Presotto[1]

[1]CERZOS, Dpto. Agronomía, Universidad Nacional del Sur (UNS)-CONICET, San Andrés 800,

CP 8000, Bahía Blanca, Argentina, [2]Estación Experimental Agropecuaria Hilario Ascasubi,

Instituto Nacional de Tecnología Agropecuaria, Ruta 3 Km 794, CP 8142, Hilario Ascasubi,

Argentina.

Raphanus sativus **interference in crops**

ORCiDs

Roman B. Vercellino: https://orcid.org/0000-0002-9313-3501

Claudio E. Pandolfo: https://orcid.org/0000-0002-7089-7138

Miguel Cantamutto: https://orcid.org/0000-0002-6854-7497

Alejandro Presotto: https://orcid.org/0000-0002-9085-9673

Author for correspondence: Roman B. Vercellino, Email: rbvercellino@cerzos-conicet.gob.ar or

Alejandro Presotto, Email: apresotto@uns.edu.ar

Abstract

Raphanus sativus (feral radish), a cosmopolitan weed, has developed resistance to acetohydroxyacid synthase (AHAS) inhibitor herbicides in several countries of South America. This study reports the effects of season-long interference of several feral radish densities on grain yield and yield components of oilseed rape, wheat and sunflower, and on feral radish traits under field conditions. Feral radish density treatments consisted of 0, 2, 4, 8 and 16 plants m^{-2} in oilseed rape, 0, 4 and 12 plants m^{-2} in wheat, and 0, 1.6, 4, 8 and 16 plants m^{-2} in sunflower. The number of inflorescences per area, seeds per inflorescence and the seed biomass of crops were reduced with increasing feral radish densities. The rectangular hyperbola model revealed yield losses by up to 100 %, 74.4 % and 12.2 % in oilseed rape, wheat and sunflower, respectively. Feral radish seed production ranged from 4,300 to 31,200, and 1000 to 4,700 seeds m^{-2} in winter crops and sunflower, respectively. Season-long feral radish interference can result in serious economic losses in oilseed rape, wheat and sunflower. The adverse impact of feral radish on the yield of winter and summer crops and the high feral radish seed and pods production suggests the need for the development and implementation of diverse and effective long-term weed management practices.

Keywords: weed interference, weed resistant, yield losses, seed production, weed seed bank

Introduction

Weeds are considered the major constraint to food production in agricultural systems around the

world (Oerke 2006). Chemical weed management is currently the most adopted and the most

effective approach for controlling weeds all over the world (Bajwa 2014, Scursoni et al. 2019).

Repeated applications of herbicides in arable fields have resulted in the rapid evolution of resistant

populations in at least 262 weed species worldwide (Heap 2020), possibly favoured by the

increasing adoption of the no-till system together with a reduction in crop rotations (Kraehmer et al.

2014). Understanding the potential crop loss from herbicide resistant weeds provides useful

information for developing and/or optimizing integrated weed management strategies and

understanding the strengths and weaknesses of target weeds.

Raphanus sativus L. (radish) is an ancient root crop from the Brassicaceae family. Weedy radish

populations are considered as de-domesticated (feral) forms derived from the cultivated radish.

Feral radish grows naturally in Europe, North America, South America and Japan, and it is a

problematic winter weed in temperate zones of the Americas (Snow and Campbell 2005, Theisen

2008, Vercellino et al. 2018). Its probable wild ancestor, *Raphanus raphanistrum* (wild radish), is a

successful invader and is one of the worst weeds worldwide (Snow and Campbell 2005). In

Australia, the southern United States and southern Brazil, *R. raphanistrum* is considered one of the

most troublesome weeds in winter crops (Lamego et al. 2013, Snow and Campbell 2005, Warwick

and Francis 2005). Both species have developed resistance to herbicides, including resistance to

acetohydroxyacid synthase (AHAS) inhibitor herbicides (Heap 2020). AHAS-resistant feral radish

has been found in Brazil, Chile and Argentina (Heap 2020, Pandolfo et al. 2016, Theisen 2008), and

wild radish in Australia, Brazil and South Africa (Heap 2020). Feral radish and wild radish are self-

incompatible and insect-pollinated with an annual or biennial life cycle (Snow and Campbell 2005).

High seed production, a darkness requirement for germination, the presence of a pericarp covering

the seeds, and seed germination in a broad range of temperatures are some of attributes that favour

the emergence and establishment of feral radish (Vercellino et al. 2018, 2019), as in the wild

relative *R. raphanistrum* (Cheam 1986, Malik et al. 2010).

Feral radish is well distributed throughout the Argentinean agricultural landscape and is one of the

most noxious weeds in the southeast of Buenos Aires province, the main winter cereal producing

area (Scursoni et al. 2014). In that area, we reported the occurrence of several AHAS-resistant feral

radish populations infesting winter and summer crops (Pandolfo et al. 2013, 2016, Vercellino et al.

2018). Since the initial report in 2011, the spread of feral radish resistant to AHAS herbicides has

rapidly increased, and currently, its presence has been reported in Buenos Aires, Tucumán and Salta

provinces (AAPRESID 2019).

AHAS-resistant feral radish must be controlled in both winter and summer crops to prevent yield

losses. In Argentina, since more than 90 % of agricultural area is sown under no-till system

(ReTAA 2020), weed control depends primarily on herbicides (Scursoni et al. 2019). In some crops,

pre-emergence herbicides may provide effective control of AHAS-resistant feral radish (Pandolfo et

al. 2016). On the other hand, post-emergent broadleaf weed control in broadleaf crops is not easy.

Farmers have a few (e.g. aclonifen) or no herbicide options for post-emergence control of AHAS–

resistant feral radish in sunflower and oilseed rape crops, respectively (Pandolfo et al. 2016). In

winter cereals (i.e. wheat and barley), post-emergent control of AHAS-resistant feral radish is

limited to phenoxy (2,4-D and MCPA) and photosystem-II inhibitor (bromoxynil) herbicides. For

these reasons, several extensive populations of AHAS-resistant feral radish, due to the Trp574Leu

mutation, have been found severely infesting both winter and summer crops (Pandolfo et al. 2016,

Theisen 2008, Vercellino et al. 2018).

Limited information exists on the effect of feral radish interference on both winter and summer

crops. In one study in Brazil, feral radish interference at densities of >40 plants m^{-2} reduced

soybean yield by 15 % (Bianchi et al. 2011). Interference studies on AHAS-resistant feral radish in

oilseed rape, wheat and sunflower will provide useful information on the importance of controlling

this species using integrated weed management strategies. The aims of this study were: (1) to estimate the impact of the interference of several densities of AHAS-resistant feral radish on the yield and yield components of oilseed rape, wheat and sunflower crops, and (2) to estimate feral radish dry biomass and seed production at several densities in oilseed rape, wheat and sunflower crops. Feral radish densities were 0, 2, 4, 8 and 16 plants m^{-2} in oilseed rape, 0, 4 and 12 plants m^{-2} in wheat, and 0, 1.6, 4, 8 and 16 plants m^{-2} in sunflower. This is the first report of feral radish interference in oilseed rape, wheat and sunflower.

Material and methods

Plant material

Two AHAS-resistant feral radish populations (RSBA10 and RSBA3-R) carrying the Trp574Leu mutation, originating from southern Buenos Aires province (Argentina) and previously characterized by Pandolfo et al. (2016) and Vercellino et al. (2018), were used in these studies. RSBA10, with 95 % of individuals resistant to AHAS herbicides carrying the Trp574Leu mutation, was collected in an IMI-resistant sunflower field treated with imazapyr near Pieres, Lobería county (S 38° 24', W 58° 35'). RSBA3 was collected in an IMI-resistant oilseed rape field treated with imazethapyr (RSBA3) near Balcarce (S 37° 35', W 58° 31'). A purified homozygous population with the AHAS Trp574Leu mutation (RSBA3-R) was generated from RSBA3 by selecting 15 homozygous resistant plants using a specific Trp574Leu mutation CAPS marker and following the procedure described in Pandolfo et al. (2016). In a previous study, no differences were found in the phenology and pods and seeds production between RSBA10 and RSBA3 (Vercellino et al. 2018). We generally used the RSBA10 accession in interference experiments.

Field experiments

Field experiments evaluating feral radish interference in oilseed rape, wheat and sunflower were conducted between 2016 and 2018 in the Agronomy Department experimental field at the

Universidad Nacional del Sur, Bahía Blanca, Argentina (38°41'46''S, 62°14'55''W). The site is

characterized as semi-arid with a temperate climate, with a mean annual temperature of 14.9 °C; the

mean temperatures of the coldest (July) and the hottest (January) month are 7.6 and 23.0 °C,

respectively. Average annual precipitation is 641 mm (National Meteorological Service 2019,

https://www.smn.gob.ar/). The soil has a well-drained loamy sand texture with 1.1 % organic matter

and pH 7.7 (a typical soil in the southwest of Buenos Aires province). Experiments were drip

irrigated as necessary for optimal growing conditions.

Under farm conditions, we observed the emergence of feral radish in the planting row probably

because the seeder disc broke the pods and released and buried their seeds during cultivation, as

reported by Cheam (1986) in the wild relative *R. raphanistrum*. For this reason, we evaluated the

interference of feral radish established at the same time as the crops. We selected the weed densities

according to our previous surveys under field conditions. In each experiment, the target feral radish

densities were achieved by manually sowing 2–3 seeds per hole and 2–3 cm deep, spaced according

to the density plan. At the four–leaf stage of each crop (i.e. oilseed rape, wheat and sunflower), the

target weed stand was adjusted by hand thinning. Hand weeding was carried out periodically to

avoid interference from other weeds. Three different set of experiments were performed:

a. Oilseed rape

The oilseed rape cultivar Hyola 575 CL was planted by hand in 0.2-m rows at 50 plants m^{-2} on May

2017 and 2018, during the winter growing season. Experimental units (plots) were seven rows wide

spaced at 0.2 m by 2 m long. Simultaneously, feral radish seeds were planted, as was described

above, to achieve 0 (control), 2, 4, 8 and 16 plants m^{-2}. Experiments were fertilized with 90 kg ha^{-1}

diammonium phosphate at planting and 200 kg ha^{-1} urea at the four–six-leaf stage. The

experimental design was a randomized complete block with four replications. The experiments

were performed in the same manner in 2017 and 2018.

At oilseed rape maturity, the plant height, number of branches, and pods per main inflorescence

were measured in ten successive oilseed rape plants in the middle of the three central rows of each

experimental unit. Then, the plants were hand-harvested and dried at 60 °C to constant weight to

homogenize treatments. After that, we quantified seeds per pod, seed weight and yield per plant.

Seeds per pod was estimated by averaging the number of seeds per pod of five pods per plant. Each

plant was threshed by hand and its seeds were cleaned and weighed to obtain the plant yield. The

seed weight was estimated by averaging the weight of three replicates of 100 seeds per plant. Data

from the ten oilseed rape plants per experimental unit were averaged for the statistical analysis.

Oilseed rape seed weight and yield were standardized to 8 % moisture content.

b. Wheat

Wheat cultivars ACA 360 and KLEIN PROTEO were planted in mid-May 2018 (winter) and in

early August 2018 (spring), respectively. Crop density was adjusted to 200 and 350 plants m^{-2} for

winter and spring wheat, respectively. Plots were nine rows wide spaced at 0.2 m by 1.5 m long.

Simultaneously, feral radish seeds were planted, as was described above, to achieve 0 (control), 4

and 12 plants m^{-2}. Experiments were fertilized with 90 kg ha^{-1} diammonium phosphate at planting

and 200 kg ha^{-1} urea at the five–leaf stage. The experimental design was a randomized complete

block with four replications. The experiments were performed in the same way in both growing

seasons.

At wheat maturity, plant height was measured on three replicates of five successive plants, and

three replicates of 0.5 m long in the middle of the three central rows of each experimental unit were

hand-harvested and dried at 60 °C to constant weight to homogenize treatments and the growing

seasons. In each replicate, we also evaluated the number of spikes m^{-2}, spikelets per spike, grains

per spikelet, grain weight and yield. Spikelets per spike and grains per spikelet were obtained by

averaging five spikes, and grain weight by averaging the weight of four replicates of 100 grains.

Data from wheat plants of each experimental unit were averaged for the statistical analysis. Wheat

grain weight and yield were standardized to 13.5 % moisture content.

c. Sunflower

The sunflower cultivar SYN 3970 CL was planted at 6.4 plants m^{-2} on Nov 2016, during the

summer growing season. Plots were five rows wide spaced at 0.52 m by 2.1 m long.

Simultaneously, feral radish seeds were planted, as was described above, to achieve 0 (control), 1.6,

4, 8 and 16 plants m^{-2}. The experimental plots were fertilized with 90 kg ha^{-1} diammonium

phosphate at planting and 150 kg ha^{-1} urea at the four–six-leaf stage. Experimental design was a

randomized complete block with four replications.

At the end of crop flowering, the plant height, width of the three middle leaves and the number of

green leaves were measured on three successive sunflower plants in the middle of the central row of

each experimental unit. The leaf area was estimated according to Aguirrezábal et al. (1996) from

the leaf width, and the leaf area was obtained by averaging the leaf area of three leaves per plant. At

sunflower maturity, the heads of the three evaluated plants were hand-harvested and dried at 60 °C

to constant weight. After that, we measured the head diameter, seeds per plant, seed weight and

yield per plant in the three plants harvested per experimental unit. The seed weight was estimated

by averaging the weight of four replicates of 100 seeds. Data from the sunflower plants in each

experimental unit were averaged for the statistical analysis. Sunflower seed weight and yield were

standardized to 11 % moisture content.

Weed traits

At the experimental level, we measured the plant height in four successive feral radish plants

located in the centre of each experimental unit at weed maturity. Then, plants were hand-harvested,

dried at 60 °C to constant weight and weighed to determine the dry biomass per weed plant. After

that, pods from these plants were counted and crushed by hand or using a mortar. Seeds were

cleaned and weighed to obtain the plant yield. The seed weight was estimated by averaging the

weight of three replicates of 100 seeds per plant. The number of seeds per plant was estimated by

dividing the plant yield by the seed weight for each plant. Seeds per pod was estimated by dividing

the seeds per plant by the number of pods per plant. Data from the feral radish plants per

experimental unit were averaged for the statistical analysis. Weed traits were measured in the same

way for all three different set of experiments.

Statistical analysis

The relationship between the crop yield and feral radish density was analysed for each crop and

year (i.e. oilseed rape) or growing season (i.e. wheat) with the rectangular hyperbolic model

(Cousens 1985) using PROC NLIN procedure in SAS/STAT software (SAS University Edition,

SAS Institute Inc., Cary, NC, USA).

The yield loss relationship is described by Equation 1:

$YL = ID/(1 + ID/A)$ (1)

where *YL* is the percentage crop yield loss relative to the weed-free treatment, *I* is the percentage

crop yield loss per weed as the weed density approaches zero, *D* is the weed density, and *A* is the

percentage crop yield loss as the weed density approaches infinity. The model was constrained to an

asymptotic percentage yield loss, A lying between 0 and 100% because yield loss cannot exceed

100%. The parameter *I* is used as an indicator of potential weed competitiveness (Cousens 1985).

As no other crop data fitted a logical/appropriate regressions model, crop data were analysed using

general linear models (GLMs) with PROC GLM in SAS. Feral radish density and year (i.e. oilseed

rape) or growing season (i.e. wheat) were considered as fixed factors, to test for significant main

effects and interactions. ANOVA and Fisher's Protected LSD tests were used to determine the

variation between treatments.

To analyse the relationships between feral radish traits (e.g. dry biomass, pod number and seed

production) and plant density for each experiment, we performed regression analyses using Prism

software (Graphpad Prism 7.0; GraphPad Software, San Diego, California, USA). These

relationships were established from the best fits of the experimental data to the appropriate

functions, and the coefficients of determination (R^2) were reported. ANOVA was used to determine

the variation between treatments in weed plant height, seeds per pod and seed weight.

The two-parameter hyperbolic regression model (Equation 2) was used to describe the density-

dependent effects of feral radish on the dry biomass of feral radish, the pod number and seed

production per meter square:

$Y = ab / (b + D)$ (2)

where Y is feral radish dry biomass, pod number and seed production per square meter, a is the

asymptote or estimate of maximum feral radish dry biomass, pod number and seed production, b is

the estimate of the feral radish density at which 50 % of maximum feral radish dry biomass, pod

number and seed production, and D is weed density.

The one-phase decay regression model (Equation 3) was used to describe the density-dependent

effects of feral radish on feral radish dry biomass, pod number and seed production per plant:

$Y = b + (a - b)^{-cD}$ (3)

where Y is feral radish dry biomass, pod number and seed production per plant, b is the Y value at

infinite feral radish density, expressed in the same units as Y, a is the Y value when D approaches

zero, and D is weed density.

Results and Discussion

Oilseed rape

In both years, the oilseed rape yield decreased substantially as the feral radish density increased.

Analysis of oilseed rape traits revealed no significant interactions between weed density and year in

the plant height, number of branches, pods per main inflorescence and seeds per pod, therefore data

from the two growing seasons were pooled (Table 1 and S1). However, there was a significant

interaction in the seed weight, therefore this data was analysed separately by year (Table 1 and S1).

Feral radish interference had a significant effect on plant height, number of branches, pods per main

inflorescence and seeds per pod (Table 1 and S1). The plant height was reduced at 4 feral radish m^{-2}

and increasing weed density from 4 to 16 plants m^{-2} reduced the plant height by up to 15.6 %. The

number of branches was reduced at 2 weeds m^{-2} and increasing feral radish density from 2 to 16

plants m^{-2} decreased the number of branches by up to 35.7 % (Table 1). Pods per main inflorescence

and seeds per pod were only reduced at the high weed density (16 plants m^{-2}), and the reduction was

20.6 and 18.4 %, respectively (Table 1). Similarly, the number of branches, the pods per

inflorescence and the seeds per pod were the most affected yield components in oilseed rape during

interference in several studies (Rondanini et al. 2017, Holman et al. 2004). Seed weight was not

affected by feral radish interference in 2017, but there were significant reductions (19.8 to 28.5%) at

the densities ≥ 8 feral radish m^{-2} in 2018 (Table 1).

Table 1. Oilseed rape traits under interference of five feral radish densities

Weed m^{-2}	Plant height (cm)	Number of branches	Pods per main inflorescence	Seeds per pod	Seed biomass (mg) 2017	Seed biomass (mg) 2018
0	99.2 ± 4.4 a	5.86 ± 0.24 a	27.6 ± 1.4 ab	17.4 ± 0.4 ab	3.27 ± 0.06	3.46 ± 0.08 a
2	97.6 ± 5.9 ab	5.30 ± 0.18 b	29.2 ± 2.0 a	17.8 ± 0.4 a	3.27 ± 0.16	3.25 ± 0.13 a
4	91.8 ± 3.9 bc	4.56 ± 0.30 c	26.6 ± 1.0 ab	17.4 ± 0.3 ab	3.20 ± 0.05	3.22 ± 0.11 a
8	92.4 ± 5.0 ab	4.66 ± 0.21 c	25.1 ± 1.6 b	16.1 ± 0.4 b	3.05 ± 0.12	2.77 ± 0.14 b
16	84.9 ± 3.2 c	3.77 ± 0.21 d	21.9 ± 1.4 c	14.2 ± 0.5 c	3.12 ± 0.10	2.46 ± 0.02 c
ANOVA	**	**	**	**	ns	**

Observations are means ± SE of two years, except for the variable "seed weight" with showed

density x year interaction, therefore means ± SE are shown for each year (2017 and 2018). For each

trait, a different letter indicates significant differences (*$P \leq 0.05$ Fisher's LSD; **$P \leq 0.01$ Fisher's

LSD).

The adjusted rectangular hyperbolic functions (R^2 = 0.95 and 0.93) showed that feral radish

decreased the oilseed rape yield per unit by 16.8 – 15.7 % at low weed density (parameter *I*), and by

100 % when the density approaches infinity (parameter *A*) for 2017 (F = 622.87; P < 0.0001) and

2018 (F = 374.18; P < 0.0001), respectively (Fig. 1). Oilseed rape yields with the weed-free

treatment were 6706 ± 340 and 4647 ± 302 kg ha^{-1} in 2017 and 2018, respectively. Differences

observed between years were probably related to prevailing environmental conditions during the

growing season. Despite the differences in oilseed rape yield between the two years, oilseed rape

yield losses were very similar (Fig. 1). Our results are similar to those obtained by Blackshaw et al.

(2002) in the wild relative *R. raphanistrum*, although they used a wider range of weed densities (0

to >60 wild radish plants m^{-2}), which could indicate that feral radish is more competitive than wild

radish. High level of interference in oilseed rape was also found as a result of the interference of the

Brassicaceae species *Sinapis arvensis* (Naderi and Ghadiri 2011). Because seeds of *R. sativus* have

about 32.6 % erucic acid (Mandal et al. 2002) and 149 μM g^{-1} glucosinolates (Daxenbichler et al.

1991), seeds of feral radish could also reduce oilseed rape quality by increasing the erucic acid

content of oilseed rape oil and the glucosinolate levels in oilseed rape meal that is marketed as

livestock feed, as demonstrated in *R. raphanistrum* (Blackshaw et al. 2002).

Figure 1. Relationship between feral radish density (plants m^{-2}) and yield loss (%) of oilseed rape

(2017 and 2018) crops described with a rectangular hyperbola model. Vertical bars indicate means

$\pm$ SE. Data fit the model $YL = 16.82D/(1 + 16.82D/112.10)$ (oilseed rape 2017) and $YL = 15.72D/(1$

$+ 15.72D/123.70)$ (oilseed rape 2018).

Wheat

In both growing seasons, as the feral radish density increased the wheat yield also decreased

considerably. Analysis of the wheat traits revealed no significant interaction between the weed

density and growing season in the plant height, spikes m^{-2}, spikelets per spike, grains per spikelet

and grain weight, therefore data from the two growing seasons were pooled (Table 2 and S2).

Interference from feral radish did not affect wheat plant height. However, spikes m^{-2}, spikelets per

spike, grains per spikelet and the grain weight were significantly reduced at 4 weeds m^{-2}, and

increasing feral radish density from 4 to 12 plants m^{-2} reduced the spikes m^{-2} by up to 36.3 %, the

spikelets per spike by up to 7.5 %, the grains per spikelet by up to 20.1 % and the grain weight by

up to 12.3 % (Table 2). Similar results were observed due to the interference of the grass weeds,

Avena fatua and *Lolium persicum* (Holman et al. 2004, Guillen-Portal et al. 2006). However, the

Brassicaceae species *Rapistrum rugosum* reduce wheat yield mainly due to a reduction in the

number of spikes per area and grains per spike but, unlike our results, the weed did not affect the

grain weight (Manalil and Chauhan 2019).

Table 2. Wheat traits under interference of three feral radish densities.

Weed m^{-2}	Plant height (cm)	Spikes m^{-2}	Spikelets per spike	Seeds per spikelet	Seeds biomass (mg)
0	95.7 $\pm$ 1.2	716 $\pm$ 36 a	16.0 $\pm$0.2 a	2.29 $\pm$ 0.06 a	43.7 $\pm$ 1.3 a
4	95.9 $\pm$ 0.8	526 $\pm$ 40 b	15.4 $\pm$ 0.1 b	2.07 $\pm$ 0.07 b	39.6 $\pm$ 1.1 b
12	96.6 $\pm$ 0.8	456 $\pm$ 29 c	14.8 $\pm$ 0.2 c	1.83 $\pm$ 0.04 c	38.3 $\pm$ 1.7 b
ANOVA	ns	**	**	**	*

Observations are means $\pm$ SE of two growing seasons. For each trait, a different letter indicates

significant differences (*$P \leq 0.05$ Fisher's LSD; **$P \leq 0.01$ Fisher's LSD).

The adjusted rectangular hyperbolic functions (R^2 = 0.93 and 0.86) showed that feral radish reduced

the wheat yield per unit by 46.3 and 14.5 % at low weed density (parameter *I*), and by 62.2 and 74.4

% when the density approached infinity (parameter *A*) for winter (*F* = 204.15; *P* < 0.0001) and

spring (*F* = 83.83; *P* < 0.0001) wheat, respectively (Fig. 2). Although, the low number of weed

densities could result in over or underestimation of the parameter *I* (Fig. 2), the use of this model

allowed us to have an estimation of the effect of feral radish interference on wheat yield at mid or

high densities (situation commonly found under field conditions because weeds grow in patches)

(Fig. 2). Wheat yields with the weed-free treatment were 8049 ± 299 and 8538 ± 204 kg ha^{-1} in

winter and spring growing season, respectively. Differences observed between growing seasons

were probably related to differences in the competitive capacity of wheat cultivars and prevailing

environmental conditions during the growing season. Our results were similar to those obtained by

Eslami et al. (2006) and Manalil and Chauhan (2019) in *R. raphanistrum* and *R. rugosum*,

respectively, although they used a wider range of weed densities (0 to >60 wild radish plants m^{-2}

and 0 to >45 *R. rogosum* plants m^{-2}), which could indicate that feral radish is more competitive than

wild radish and *R. rugosum*, two important Brassicaceae weeds.

Figure 2. Relationship between feral radish density (plants m^{-2}) and yield loss (%) of wheat (winter

and spring) crops described with a rectangular hyperbola model. Vertical bars indicate means ± SE.

Data fit the model *YL* = 46.26D/(1 + 46.26D/62.19) (winter wheat) and *YL* = 14.54D/(1 +

14.54D/74.44) (spring wheat).

Sunflower

Sunflower yield was also affected by feral radish interference. The interference of feral radish

reduced the sunflower plant height, green leaves per plant and head diameter. The plant height and

number of green leaves were reduced by 6.5 and 11.0 %, only at the high weed density (16 plants

m^{-2}). The head diameter was reduced by 7.2% with 8 feral radishes m^{-2} (Table 3 and S3). However,

the interference of feral radish did not significantly affect the number of seeds per head and the seed

weight (Table 3 and S3). The rectangular hyperbolic functions (R^2 = 0.57) showed that feral radish

decreased (*F* = 48.36; *P* < 0.0001) the sunflower yield per unit by 5.4 % at low weed density

(parameter *I*), and by 12.2 % when the density approached infinity (parameter *A*) (Fig. 3). The

estimated weed-free sunflower yield was 3728 ± 236 kg ha^{-1}. A similar result was found in the

season-long interference of feral radish on soybean (Bianchi et al. 2011). The higher interference

caused by feral radish in oilseed rape and wheat crops compared to sunflower and soybean (Bianchi

et al. 2011) may possibly be because feral radish is a winter weed (Snow and Campbell 2005), and

therefore it had larger plant size in winter than in summer (see below) (Vercellino et al. 2018). In

another conditions, Pandolfo (2016) observed sunflower yield losses greater than 70% due to feral

radish interference at farm level. This indicate that environmental conditions, as well as other

factors (e.g. soil type and the timing of weed emergence relative to the crop), can play a key role in

weed interference (Swanton et al. 2015).

Table 3. Sunflower traits under interference of feral radish under experimental and farm conditions.

Weed m^{-2}	Plant height (cm)	Number of green leaves	Leaf area (cm^2)	Head diameter	Seeds per head	Seed biomass

					(cm)		(mg)
0	156.3 ± 2.0 a	16.4 ± 0.7 a	361.1 ± 6.3	13.8 ± 0.4 a	1238 ± 19	47.3 ± 2.2
1.6	151.8 ± 2.7 ab	16.2 ± 0.1 ab	313.2 ± 23.5	13.6 ± 0.6 ab	1167 ± 68	47.3 ± 1.6
4	151.1 ± 1.7 ab	15.9 ± 0.6 ab	286.6 ± 5.6	13.0 ± 0.4 ab	1113 ± 39	47.3 ± 1.5
8	150.5 ± 1.8 ab	15.2 ± 0.1 ab	314.2 ± 31.5	12.8 ± 0.5 b	1099 ± 108	47.4 ± 1.2
16	146.1 ± 3.3 b	14.6 ± 0.6 b	239.9 ± 28.4	13.0 ± 0.5 ab	1094 ± 86	45.7 ± 0.8
ANOVA	*	*	ns	*	ns	ns

Observations are means ± SE. For each trait and field condition, a different letter indicates

significant differences (*$P \leq 0.05$ Fisher's LSD; **$P \leq 0.01$ Fisher's LSD).

Figure 3. Relationship between feral radish density (plants m^{-2}) and yield loss (%) of sunflower

crop described with a rectangular hyperbola model. Vertical bars indicate means ± SE. Data fit the

model $YL = 5.43D/(1 + 5.43D/12.19)$.

Clearly, high-yielding oilseed rape, wheat and sunflower crops are sensitive to interference from a

competitive broadleaf weed such as feral radish. These results showed that oilseed rape, wheat and

sunflower yields tended to decrease as the feral radish density increased. In general, feral radish

interference mainly reduced the number of inflorescences per area (represented by spikes m^{-2} in

wheat and the number of primary branches by plant density in oilseed rape), followed by a

reduction in the inflorescence size and the number of seeds per inflorescence (represented by

spikelets per spike and by grains per spikelet in wheat, pods per inflorescence and by seeds per pod

in oilseed rape, and head diameter and seeds per head in sunflower), and finally the seed weight.

These results are in agreement with those previous that evaluate the effect of the weed interference

in oilseed rape, wheat and sunflower crops (Manalil and Chauhan 2019, Guillen-Portal et al. 2006,

Holman et al. 2004).

Feral radish traits

All feral radish in these studies assumed an annual life cycle. Feral radish traits were more affected

in the sunflower experiment than in either the oilseed rape or wheat experiments. The feral radish

plant height was not affected by weed density, which ranged from 118.5 ± 7.9 cm to 149.0 ± 3.3 cm

in both the oilseed rape and wheat experiments. However, the feral radish plant height was $65.4 \pm$

1.8 cm in the sunflower experiment. Interference of feral radish may result in indirect yield losses at

harvest because it matures late in the growing season and farmers may avoid harvesting densely

infested areas. The presence of mature feral radish plants at the same height as the crop

inflorescences may also interfere with harvest operations and compromise the quality by

contaminating the seeds with weed seeds and foreign matter, lowering the value of the harvested

grains (Varanasi et al. 2016). This may be particularly important in the oilseed rape and wheat crops

because the feral radish plants were taller than the crops; however, it would not be important in

sunflower because the feral radish height was much lower (>50 cm) than the crop.

Feral radish dry biomass per plant and per area were very similar in oilseed rape and wheat

experiments, and at least seven times higher than in the sunflower experiment (Fig. 4 A and B).

Feral radish dry biomass combined over both oilseed rape and wheat experiments approximately

triplicated, from about $265 – 380$ g m^{-2} to $850 – 1050$ g m^{-2}, as weed density increased at least six-

fold (from 2 to >12 plants m^{-2}) (Fig. 4 B). At similar densities, feral radish dry biomass in sunflower

experiment increased from about 35 to 115 g m^{-2} (Fig. 4 B). The rapid stem elongation and leaf

expansion characteristic of sunflower (Presotto et al. 2017) could have shaded the feral radish plants

and reduced their growth. These results were consistent with higher yield losses in oilseed rape and

wheat than in sunflower. At similar weed densities, Blackshaw et al. (2002) and Eslami et al. (2006)

reported slightly lower values of dry biomass in wild radish under interference with oilseed rape

and wheat, respectively. Conversely, feral radish seeds per pod and seed weight were not affected

by weed density in any experiment (data not shown).

Preventing weed seed production in herbicide-resistant populations is a major concern among

farmers around the world in order to improve weed management and/or crop production

(Norsworthy et al. 2012). Feral radish pod and seed production in the oilseed rape and wheat

experiments were at least four times higher than in the sunflower experiment (Fig. 4C, D, E and F).

Feral radish produced around 4,300 to 6,700 and 13,800 to 31,200 seeds m^{-2} at low (2 plants m^{-2})

and high (>12 plants m^{-2}) densities, respectively, in both the oilseed rape and wheat experiments.

Although feral radish seed production was only about 1,000 to 4,700 seeds m^{-2} at low and high

weed densities in the sunflower experiment (Fig. 4F), it is a considerable amount of seeds that can

enriches the seedbank. However, feral radish has high seed retention at maturity (i.e. the indehiscent

pods remain firmly attached to the plant) (Snow and Campbell 2005), which could help in

managing this weed through seed capture and destruction (Walsh et al. 2013).

Figure 4. Relationship between feral radish density and dry matter per plant (A), dry matter per square meter (B), pod production per plant (C), pod production per square meter (D), seed production per plant (E) and seed production per square meter (E) of feral radish in oilseed rape (2017 and 2018), wheat (winter and spring) and sunflower experiments described with one-phase decay or hyperbola models. Vertical bars indicate means ± SE.

The darkness requirement and the indehiscent pericarp may have the potential to prevent, delay and spread the germination of a cohort over an extended period (Vercellino et al. 2019), which therefore results in the formation of a persistent seedbank and hinders weed control due to the lack of synchronization in germination (Chauhan et al. 2006), what make the need to destroy the feral radish seeds/pods before to dispersion even more evident.

The importance of maintaining a weed free environment in the early growth stages of crops to

prevent yield losses has been stablished in several crops (Harker et al. 2008; Page et al. 2012). Few

options to control AHAS-resistant feral radish in pre- and post-emergence of sunflower, wheat and

oilseed rape are effective. Since there are no chemical alternatives for pre- and post-emergence

control of AHAS-resistant feral radish in oilseed rape, its sowing is not recommended in

environments with presence of AHAS-resistant broadleaf weeds; at least in countries where

glyphosate-resistant oilseed rape is forbidden. In sunflower, Pandolfo et al. (2016) showed effective

control of AHAS-resistant feral radish with the pre-emergence mixture of acetochlor and

fluorochloridone, and with aclonifen in post-emergence. In winter cereals, phenoxy (2,4-D and

MCPA) and photosystem-II inhibitor (bromoxynil) herbicides are frequently used to control AHAS-

resistant feral radish in post-emergence. But, the repeated use of these herbicides to control AHAS-

resistant feral radish could led to the evolution of multiple-herbicide resistant populations as it has

happened in wild radish (Walsh et al. 2004, Lu et al. 2019).

This study shows that the interference of feral radish can cause considerable yield losses in oilseed

rape, wheat and sunflower. The high seed production, and the prolonged and irregular emergence

pattern offers an adaptive potential to feral *R. sativus* to become a dominant weed. However, high

seed retention at maturity provides an opportunity for controlling feral radish through harvest weed

seed control tactics. The use of all cultural, mechanical and herbicidal options available for effective

feral radish control must be considered in order to minimize crop yield losses and avoid/reduce

weed-resistant seed production (Norsworthy et al. 2012, Bajwa 2014). Future research should be

conducted to determine the interference and seed production of feral radish emerging at different

times after crop sowing. These studies will provide critical information for timely management of

feral radish.

Acknowledgements

This work was supported by the grant ANPCYT-PICT 2012-2854. We thank the National Research

Council of Argentina (CONICET) for a fellowship to RBV. No conflicts of interest have been

declared.

References

AAPRESID (Argentine No-Till Farmers Association) (2019) Mapa de malezas. Available:

http://www.aapresid.org.ar/rem/mapas-rem/

Aguirrezábal LAN, Orioli G, Hernández L, Pereyra V, Miravé J (1996) Girasol. Aspectos

fisiológicos que determinan el rendimiento. Unidad Integrada Balcarce. Buenos Aires.

Argentina.

Bajwa AA (2014) Sustainable weed management in conservation agriculture. Crop Prot 65:105–

113.

Bianchi M, Fleck N, Agostinetto D, Rizzardi M (2011) Interferência de *Raphanus sativus* na

produtividade de cultivares de soja. Planta Daninha 29:783–792.

Blackshaw RE, Lemerle D, Young KR (2002) Influence of wild radish on yield and quality of

canola. Weed Sci 50:344–349.

Chauhan BS, Gill GS, Preston C (2006) Tillage system effects on weed ecology, herbicide activity

and persistence: A review. Aust J Exp Agric 46:1557–1570.

Cheam AH (1986) Seed production and seed dormancy in wild radish (*Raphanus raphanistrum* L.)

and some possibilities for improving control. Weed Res 26:405–413.

Cousens R (1985) A simple model relating yield loss to weed density. Ann Appl Biol 107:239–252.

Daxenbichler ME, Spencer GF, Carlson DG, Rose GB, Brinker AM, Powell RG (1991)

Glucosinolate composition of seeds from 297 species of wild plants. Phytochemistry 30:2623–

2638.

Eslami SV, Gill GS, Bellotti B, McDonald G (2006) Wild radish (*Raphanus raphanistrum*) interference in wheat. Weed Sci 54:749–756.

Guillen-Portal FR, Stougaard RN, Xue Q, Eskridge KM (2006) Compensatory mechanisms associated with the effect of spring wheat seed size on wild oat competition. Crop Sci 46: 935–945.

Harker KN, O'Donovan JT, Clayton GW, Mayko J (2008) Field-scale time of weed removal in canola. Weed Technol 22:747–479.

Heap I (2020) The International Survey of Herbicide Resistant Weeds. Online, May 2, 2019. Available: http://www.weedscience.org.

Holman JD, Bussan AJ, Maxwell BD, Miller PR, Mickelso JA (2004) Spring wheat, canola, and sunflower response to persian darnel (*Lolium persicum*) interference. Weed Technol 18: 509–520.

Kraehmer H, van Almsick A, Beffa R, Dietrich H, Eckes P, Hacker E, Hain R, Strek HJ, Stuebler H, Willms L (2014) Herbicides as weed control agents: state of the art: II. Recent achievements. Plant Physiol 166:1132–1148.

Lamego FP, Ruchel Q, Kaspary TE, Gallon M, Basso CJ, Santi AL (2013) Habilidade competitiva de cultivares de trigo com plantas daninhas. Planta Daninha 31:521–531.

Lu H, Yu Q, Han H, Owen MJ, Powles SB (2019) A novel psbA mutation (Phe-274-Val) confers resistance to PSII herbicides in wild radish (*Raphanus raphanistrum*). Pest Manag Sci 75:144 –151.

Malik MS, Norsworthy JK, Riley MB, Bridges W (2010) Temperature and light requirements for wild radish (*Raphanus raphanistrum*) germination over a 12-month period following maturation. Weed Sci 58:136–140.

Manalil S, Chauhan BS (2019) Interference of turnipweed (*Rapistrum rugosum*) and Mexican pricklepoppy (*Argemone mexicana*) in wheat. Weed Sci 67: 666–672.

Mandal S, Yadav S, Singh R, Begum G, Suneja P, Singh M (2002) Correlation studies on oil content and fatty acid profile of some Cruciferous species. Genet Resour Crop Ev 49:551–556.

Naderi R, Ghadiri H (2011) Competition of wild mustard (*Sinapis arvense* L.) densities with rapeseed (*Brassica napus* L.) under different levels ofnitrogen fertilizer. J Agric Sci Technol 13:45–51.

Norsworthy JK, Ward SM, Shaw DR, Llewellyn RS, Nichols RL, Webster TM, Bradley KW, Frisvold G, Powles SB, Burgos NR, Witt WW, Barrett M (2012) Reducing the risks of herbicide resistance: Best management practices and recommendations. Weed Sci 60:31–62.

Oerke EC (2006) Crop losses to pests. J Agric Sci 144:31–43.

Page ER, Cerrudo D, Westra P, Loux M, Smith K, Foresman C, Wright H, Swanton CJ (2012) Why early season weed control is important in maize. Weed Sci 60:423–430.

Pandolfo CE, Presotto A, Poverene M, Cantamutto M (2013) Limited occurrence of resistant radish (*Raphanus sativus*) to AHAS-inhibiting herbicides in argentina. Planta Daninha 31:657–666.

Pandolfo CE (2016) Caracterización agroecológica de poblaciones ferales brasicáceas con resistencia a herbicidas. PhD thesis. Universidad Nacional del Sur, Bahía Blanca, Argentina. pp 83–118.

Pandolfo CE, Presotto A, Moreno F, Dossou I, Migasso JP, Sakima E, Cantamutto M (2016) Broad resistance to acetohydroxyacid-synthase-inhibiting herbicides in feral radish (*Raphanus sativus* L.) populations from Argentina. Pest Manag Sci 72:354–361.

Presotto A, Hernández F, Díaz M, Fernández-Moroni I, Pandolfo CE, Basualdo J, Cuppari S, Cantamutto M, Poverene M (2017) Crop-wild sunflower hybridization can mediate weediness

throughout growth-stress tolerance trade-offs. Agric Ecosyst Environ 249:12–21.

[ReTAA] Relevamiento de Tecnología Agrícola Aplicada de la Bolsa de Cereales (2020) http://www.bolsadecereales.com/imagenes/retaa/2020-02/199-juanbriheters2020190220.pdf. Accessed: June 2020

Rondanini DP, Menendez YC, Gomez NV, Miralles DJ, Botto JF (2017) Vegetative plasticity and floral branching compensate low plant density in modern spring rapeseed. Field Crops Res 210: 104–113.

Scursoni JA, Gigón R, Martín AN, Vigna M, Leguizamón ES, Istilart C, López R (2014) Changes in weed communities of spring wheat crops of Buenos Aires province of Argentina. Weed Sci 62:51–62.

Scursoni JA, Duarte Vera ACD, Oreja FH, Kruk BC, de la Fuente EB (2019) Weed management practices in Argentina crops. Weed Technol 33:459–463.

Snow AA, Campbell LG (2005) Can feral radishes become weeds?, pp. 193–208 in Gressel, J. (Ed.), Crop ferality and volunteerism. CRC Press, Boca Raton, FL, USA.

Swanton CJ, Nkoa R, Blackshaw RE (2015) Experimental methods for crop–weed competition studies. Weed Sci 63:2–11.

Theisen G (2008) Aspectos botânicos e relato da resistência de nabo silvestre aos herbicidas inibidores de ALS, pp. 1–24, Embrapa Clima Temperado, Pelotas, Brazil.

Varanasi A, Prasad PVV, Jugulam M (2016) Impact of climate change factors on weeds and herbicide efficacy. Advances in Agronomy 135:107–146.

Vercellino RB, Pandolfo CE, Breccia G, Cantamutto M, Presotto A (2018) AHAS Trp574Leu substitution in *Raphanus sativus* L.: Screening, enzyme activity and fitness cost. Pest Manag Sci 74:1600–1607.

Vercellino RB, Pandolfo CE, Cerrota A, Cantamutto M, Presotto A (2019) The role of light and the

pericarp on seed dormancy and germination in feral *Raphanus sativus* (Brassicaceae). Weed

Res 59: 396–406.

Walsh M, Newman P, Powles S (2013) Targeting weed seeds in-crop: a new weed control paradigm

for global agriculture. Weed Technol 27:431–436

Walsh MJ, Powles SB, Beard BR, Parkin BT, Porter SA (2004) Multiple-herbicide resistance across

four modes of action in wild radish (*Raphanus raphanistrum*). Weed Sci 52:8–13.

Warwick SI, Francis A (2005) The biology of Canadian weeds. 132. *Raphanus raphanistrum*. L.

Can J Plant Sci 85:709–733.