A novel role for the C-terminal domain of the influenza A virus host shutoff

endoribonuclease PA-X in mediating rapid turnover of the PA-X protein

Rachel Emily Levene[a,b], Shailab D. Shrestha[a,b], Marta Maria Gaglia[a,b #]

[a]Graduate Program in Molecular Microbiology, Tufts University Graduate School of Biomedical

Sciences, Boston, MA, USA

[b]Department of Molecular Biology and Microbiology, Tufts University School of Medicine,

Boston, MA, USA

[#]Send correspondence to:

Marta Maria Gaglia (marta.gaglia@tufts.edu)

Key Words: influenza A virus, PA-X, host shutoff, protein turnover

Running Title: The influenza A virus RNase PA-X is unstable

Abstract Word Count: 190

Main Text Word Count: 4,963

ABSTRACT:

The influenza A endoribonuclease PA-X regulates virulence and transmission of the virus by

reducing host gene expression and thus regulating immune responses to influenza A virus.

Despite this key function in viral biology, the levels of PA-X protein remain markedly low

during infection, and previous results suggest that these low levels are not solely the result of

regulation of the level of translation and RNA stability. How PA-X is regulated post-

translationally remains unknown. We now report that the PA-X protein is rapidly turned over.

This instability is conserved across multiple viral strains, although the half-life of PA-X from

different strains varies, ranging from 30 minutes to a few hours. Moreover, sequences in the

variable PA-X C-terminal domain regulate PA-X half-life. Interestingly, we find that the PA-X

from the 2009 pandemic H1N1 strain has a longer half-life compared to the other variants we

tested. This PA-X isoform has been reported to have a higher host shutoff activity, suggesting a

role for protein turnover in regulating PA-X activity. Collectively, this study reveals a novel

regulatory mechanism of PA-X protein levels that may impact host shutoff activity during

influenza A virus infection.

IMPORTANCE

The PA-X protein from influenza A virus reduces host immune responses to infection through

suppressing host gene expression, including genes encoding the antiviral response. Thus, it plays

a central role in influenza A virus biology. Despite its key function, PA-X was only discovered

in 2012 and much remains to be learned including how PA-X activity is regulated to promote

optimal levels of viral infection. In this study we reveal that PA-X protein levels are very low

because of rapid turnover. We show that instability is a conserved property among PA-X variants

from different strains of influenza A virus, but that the half-lives of PA-X variants differ.

Moreover, the longer half-life of PA-X from the 2009 pandemic H1N1 strain correlates with its

reported higher activity. Therefore, PA-X stability may be a way to regulate its activity and may

contribute to the differential virulence of influenza A virus strains.

INTRODUCTION:

Viruses employ multiple strategies to block host antiviral defenses. In many viruses, this
inhibition is partly achieved through a blockade of host gene expression, a process termed "host
shutoff". One of the ways influenza A virus induces host shutoff is through the activity of a viral
endoribonuclease (RNase), PA-X (1–3). The RNase activity of PA-X results in global changes in
the cellular transcriptome (3, 4), while viral mRNAs and genomic RNAs escape PA-X
degradation (5). Influenza A viruses engineered to lack PA-X trigger a more potent innate
immune and inflammatory response in mice, chickens, and pigs, presumably because of
decreased host shutoff (1, 6–10). These results indicate that PA-X controls immune responses *in
vivo.* More recently, a role for PA-X in transmission has also been established (11, 12). Despite
its clear *in vivo* role, much remains to be determined about PA-X biology during infection. PA-X
was discovered in 2012 and studies to date have primarily focused on dissecting its mechanism
of action, including how subsets of RNAs, including viral RNAs, escape PA-X degradation.
Whether and how PA-X activity itself is regulated has not been explored in detail. One study
found that co-translational N-terminal acetylation is required for PA-X host shutoff activity (13),
but whether there are other co- and post-translational regulatory mechanisms remains unknown.
Interestingly, prolonged PA-X expression may be cytotoxic, as shown in yeast by Oishi *et al.*
(13, 14). While this is not surprising given its widespread downregulation of RNAs, this
observation underscores the idea that PA-X activity may need to be regulated during infection to
prevent premature death of the infected cells.

One way to regulate or limit PA-X activity could be to keep pools of active PA-X protein low in
the infected cell. Indeed, PA-X protein levels are remarkably low during infection. Metabolic

labeling of nascent proteins in influenza A virus infected cells showed that the levels of PA-X

are barely above the limit of detection and lower than other influenza proteins, including the

well-known host shutoff factor NS1 (1). These low levels are in part due to how PA-X is

produced. PA-X production requires a programmed +1 ribosomal frameshift during translation of

segment 3, which encodes the polymerase acidic (PA) subunit of the viral RNA dependent RNA

polymerase (RdRp) (1, 15). Only a small percentage of translational runs result in frameshifting

and produce PA-X, which has the same N-terminal RNase domain as PA (PA-X amino acids (aa)

1-191) and a unique C-terminal domain known as the "X-ORF" (PA-X aa 192-232 or 192-252

depending on the strain) (1). Of note, despite this non-canonical production mechanism, PA-X is

present in all influenza A virus isolates (16) demonstrating its importance in influenza A virus

biology. However, the mechanism of PA-X production does not entirely explain the low levels of

PA-X. Indeed, PA-X levels are also low or undetectable when PA-X is expressed ectopically

from a construct in which a single nucleotide has been deleted to mimic the frameshift and all

translational runs produce PA-X (17). The low levels are also not because PA-X degrades and

limits the levels of its own mRNA, as mutating the PA-X catalytic residues D108 or K134 in this

ectopic expression system does not increase PA-X levels dramatically (3, 17). This suggests that

PA-X levels may be regulated at the level of protein stability.

Regulation of stability is a well-established post-translational mechanism of protein regulation.

For example, Kosik *et. al* established a role for protein turnover in the regulation of the

immunomodulatory activity of PB1-F2, another influenza A virus virulence factor (18). PB1-F2

inhibits induction of the antiviral cytokine interferon beta (IFN-β) and increases the activity of

the viral RdRp (19–21). Mutating the residues required for PB1-F2 ubiquitination and rapid

turnover increases PB1-F2 protein levels, which results in a stronger inhibition of IFN-β

induction (18). Similarly, regulation of PA-X at the level of protein stability could modulate the

host shutoff activity of influenza A virus.

Previous observations suggest that the X-ORF may regulate the levels of the PA-X protein, as

ectopic expression of the N-terminal domain of PA-X results in much higher protein levels than

ectopic expression of either wild type (wt) or catalytically inactive PA-X (17). This is not due to

changes in frameshifting, as this ectopic expression construct has a nucleotide deletion to mimic

the frameshift and only encodes PA-X. A role for the X-ORF in post-translational control of PA-

X levels and activity would explain why almost all influenza A strains encode a PA-X protein

that has a 61 or 41-aa X-ORF (16) even though only the first 15 aa of the X-ORF are required for

PA-X shutoff activity in cells (3, 17, 22). Moreover, most of the sequence variation among PA-X

from different strains occurs after the 15th amino acid of the X-ORF, and the strength of PA-X

activity has been reported to vary among strains (17, 23, 24). Therefore, we hypothesized that the

later portion of the X-ORF and the sequence variation in this region have a regulatory effect on

PA-X levels and host shutoff activity.

Here we report the first investigation of post-translational regulation of PA-X protein levels. We

find that PA-X is markedly unstable and identify a region of the X-ORF that regulates PA-X

turnover. Additionally, as the X-ORF sequences vary among influenza A virus strains, we find

that the half-life of PA-X differs among strains, even though all the PA-X variants tested have

generally short half-lives. Interestingly, of the variants we tested, PA-X from the 2009 pandemic

H1N1 strain had the longest half-life, which may explain its reported higher host shutoff activity

(17, 23, 24). Thus, modulation of PA-X stability may be a mechanism used to regulate PA-X

host shutoff activity in different strains and may contribute to influenza A pathogenesis.

RESULTS

PA-X is unstable

Several observations suggested to us that PA-X may be unstable. PA-X is expressed at low levels

not only during infection (1), but also during ectopic overexpression from a high expression

CMV promoter [(17), our unpublished observations]. Furthermore, ectopic overexpression of the

N-terminal RNase domain from the same high expression CMV promoter results in much higher

protein levels (17). While the RNase domain in isolation is inactive in cells (3, 5, 17, 22), the

differences in protein levels are not due to the loss of activity, because expressing a catalytically

inactive version of PA-X still results in markedly low PA-X expression levels (3, 17). Thus, we

sought to directly test if PA-X is rapidly turned over. To determine the half-life of the PA-X

protein, we performed a cycloheximide chase in human embryonic kidney (HEK) 293T cells

transfected with a C-terminally myc tagged PA-X from the influenza A virus strain A/Puerto

Rico/8/1934 H1N1 (PR8) containing the D108A mutation in the RNase active site. This mutation

abolishes PA-X catalytic activity and host shutoff function (3, 5). We chose to use catalytically

inactive PA-X for this experiment and throughout this study to eliminate any confound from

changes in host gene expression or PA-X mRNA levels as a result of PA-X activity.

Additionally, to make sure any differences in PA-X protein levels are due to changes in turnover

and not frameshifting, in this experiment and throughout this study we used PA-X expression

constructs where all translational runs produce PA-X, as a single nucleotide was deleted to

mimic the frameshift. After treating PA-X-expressing cells with cycloheximide to inhibit

translation, we analyzed PA-X protein levels at 0, 15, 30, 60, and 120 minutes after

cycloheximide addition by western blotting. Strikingly, PA-X was almost completely gone one

hour after the addition of cycloheximide (Fig 1A and 1B), suggesting that PA-X is markedly

unstable. By contrast, levels of PA-X mRNA did not decrease throughout the cycloheximide

time course (Fig 1C), confirming that the decrease in PA-X protein levels is not due to changes

in transcript levels. Based on regression analysis, we calculated a half-life of 86 minutes for PR8

PA-X (Fig 1B, Table 1). This is low compared to the half-life of other influenza proteins. For

example, the host shutoff factor NS1 from PR8 has a half-life of over 24 hours (25) and PR8

PB1-F2 has a half-life of 6.5 hours (26). In addition, human proteins have a median half-life of

8.7 hours (27). Together, these data show that PA-X is rapidly turned over compared to other

viral and host proteins.

The X-ORF mediates PA-X instability

Because ectopic overexpression of the N-terminal domain of PA-X results in higher protein

levels than overexpression of full length PA-X (17), we decided to test whether the X-ORF

controls PA-X stability. We thus measured the half-life of a truncated version of the catalytically

inactive PR8 PA-X that ended after aa 16 of the X-ORF ("PA-X16", Fig 2A). We chose to test

PA-X16 instead of the RNase domain alone because truncation of the whole X-ORF results in

loss of nuclear localization of PA-X (5, 17). In contrast, PA-X mutants retaining at least 15 aa of

the X-ORF are primarily localized to the nucleus, like full-length PA-X (17). Additionally, we

and others have previously shown that the first 15 residues of the X-ORF are necessary and

sufficient for host shutoff activity (3, 17, 22). Thus, using the PA-X16 truncation should exclude

potential confounding factors, such as changes in subcellular localization or in protein-protein

interactions needed for activity. In contrast to the rapid decrease in the levels of full-length PA-X

(PA-X61) after cycloheximide addition, the protein levels of PA-X16 remained constant

throughout the two-hour time course (Fig 2B and 2C). This result shows that PA-X16 is more

stable than the full-length protein. There were also no significant changes in the mRNA levels of

PA-X16 throughout the time course (Fig 2D). Together, these results indicate that sequences in

the X-ORF after the 16th residue are responsible for the instability of PA-X. Since the first 15

amino acids of the X-ORF are necessary and sufficient for host shutoff activity (3, 5, 17, 22),

these data also suggest that the portions of the X-ORF involved in mediating host shutoff activity

and stability are different.

The half-life of PA-X varies among strains

Although all influenza A virus strains are predicted to encode PA-X, the sequence of PA-X

varies across strains, particularly in the unique C-terminal X-ORF (16). To make sure that the

instability we observed is not specific to the X-ORF sequence of PR8 PA-X, we sought to

determine if this phenotype was conserved among PA-X from other strains of influenza A virus.

We created catalytically inactive D108A mutants in the PA-X from a 2009 pandemic H1N1

strain, A/California/7/2009pdm (pH1N1), a lab-adapted H3N2 strain, A/Udorn/307/1972

(Udorn), a seasonal H3N2 strain, A/Perth/16/2009 (Perth), and a pre-pandemic seasonal H1N1

strain, A/Hong Kong/218847/2006 (HK06), all with a C-terminal myc tag. Tagging all variants

with the same tag allows us to compare among strains since many antibodies against influenza

proteins only recognize a particular strain or serotype. PA-X from Udorn and HK06 were as or

more unstable than PR8 PA-X, with half-lives of 34 and 83 minutes, respectively (Fig 3A, 3B,

3D, 3E, Table 1). In contrast, PA-X from Perth was slightly more stable, with a calculated half-

life of 250 minutes (4.2 hrs, Fig 3G, 3H, Table 1). PA-X from pH1N1 was significantly more

stable, as its levels remained constant throughout the cycloheximide time course (Fig 3J, 3K,

Table 2). That said, after four hours of cycloheximide treatment, pH1N1 PA-X protein levels

decreased by half (Fig 3M and 3N), suggesting it may also have a shorter half-life than the

average human protein (~9 hrs) (27). In all cases, there were no significant decreases in mRNA

levels of PA-X from any of the strains (Fig 3C, 3F, 3I, 3L, 3O), again confirming that the

decrease in PA-X protein levels are not due to changes in transcript levels. Together, these data

show that instability is a conserved property among PA-X variants, but that there are differences

in PA-X half-life among strains. Moreover, the pH1N1 PA-X is substantially longer lived than

other variants.

The length difference between the two major variants of the PA-X X-ORF does not account

for differences in PA-X instability

There are two main variants of PA-X in influenza A strains distinguishable by their protein

lengths: one is 252-aa long, with an X-ORF of 61 aa, and the other is 232-aa long, with an X-

ORF of 41 aa (16). The PA-X variants in PR8, Perth, Udorn and HK06 all have a 61-aa X-ORF,

like most human strains, while the pH1N1 PA-X has a 41-aa X-ORF, which is otherwise found

in a subset of swine and canine strains (16). Although some studies have suggested that the

length of the X-ORF may regulate activity (12, 28, 29), it is still unclear exactly what the

function of the additional 20 aa is. Since pH1N1 PA-X has a much longer half-life than most of

the other PA-X isoforms we tested (Fig 3K, Table 1), we wondered if the shorter X-ORF

accounted for the difference in stability. We thus tested the half-life of a truncated version of

catalytically inactive PR8 PA-X that ended after the 41st amino acid (Fig 4A), which we termed

"PA-X41". Despite the truncation, PR8 PA-X41 protein levels rapidly decreased after translation

inhibition (Fig 4B and 4C), with a half-life of 99 minutes. This is not significantly different from

that of full length PR8 PA-X (PA-X61) (p = 0.8118) (Fig 4C). We note that we consistently

detected a double band in cells expressing PR8 PA-X41. While the reason for this doublet is

currently unknown, we note that a similar doublet was also apparent in the staining for pH1N1

PA-X in Hayashi *et al.* (17). There were no significant decreases in the mRNA levels of PR8

PA-X41 (Fig 4D), confirming that the decrease in PR8 PA-X41 protein levels is not due to

changes in transcript levels. Thus, the distal 20 residues likely do not play a role in protein

stability, and other sequence differences between PR8 and pH1N1 PA-X control their turnover

rates.

PA-X host shutoff activity in infected cells is reduced upon translational inhibition

To confirm that PA-X is unstable in the context of influenza A virus infection we first attempted

to measure its half-life in PR8-infected cells using an antibody against PA that has been

previously used to detect PR8 PA-X (30). However, like Rigby *et al.* (30)*,* we observed a large

number of bands in the section of the blot corresponding to the predicted molecular weight of

PA-X, and were unable to reliably and reproducibly identify the band corresponding to PA-X.

Unfortunately, we cannot tag PA-X in the context of the virus. The C-terminus cannot be tagged

because it overlaps with the coding sequence for the C-terminus of PA (1, 15). N-terminal tags

reduce PA activity and viral replication (31, 32), and are also expected to reduce the activity of

PA-X because of the importance of its N-terminal acetylation (13). Therefore, we sought another

way to look for evidence of PA-X instability during infection and used PA-X activity in the

presence of cycloheximide as a proxy for rapid changes in PA-X protein levels. We reasoned that

if levels of PA-X are quickly depleted after halting translation with cycloheximide, there will be

less PA-X host shutoff activity in cells. To confirm this, we first tested PA-X host shutoff

activity in the presence of cycloheximide in A549 cells that inducibly overexpress either wt PR8

PA-X or the catalytically inactive PR8 PA-X D108A mutant. These cells allow us to measure

changes in the RNA levels of endogenous genes (3, 5). Following overnight induction of PA-X

expression and a 2-hour cycloheximide treatment, we analyzed levels of G6PD and TAF7 RNA,

representative PA-X targeted and resistant RNAs, respectively (3). Levels of G6PD significantly

decreased upon PA-X expression, but this decrease was not observed in PA-X-expressing cells

treated with cycloheximide (Fig 5A). In contrast, cycloheximide had no effect on the levels of

G6PD in cells expressing catalytically inactive PA-X or levels of the PA-X resistant RNA TAF7

(Fig 5A), confirming that the effect of cycloheximide on RNA levels was linked to PA-X

activity. Furthermore, PA-X mRNA levels did not decrease with cycloheximide, confirming that

any changes in PA-X host shutoff activity are not the result of changes in PA-X transcript levels

(Fig 5A).

We then performed the same experiment in A549 cells infected with either wt PR8 or PA-X-

deficient PR8 virus (ΔX). PR8 ΔX carries mutations in the frameshifting site in segment 3 that

reduce PA-X production, as well as a nonsense mutation at PA-X aa 201, which abolishes

activity of any residual PA-X (3). At 24 hours post-infection, we analyzed levels of G6PD and

TAF7 mRNA after a 2-hour cycloheximide treatment. As expected, G6PD levels were

significantly lower in wt PR8-infected cells compared to PR8 ΔX-infected cells due to PA-X

mediated host shutoff (Fig 5B), while the levels of the PA-X-resistant RNA TAF7 were similar.

However, the difference in G6PD levels was not detected in infected cells treated with

cycloheximide, suggesting that cycloheximide treatment prevents PA-X-mediated host shutoff,

presumably due to rapid loss of PA-X expression (Fig 5B). Conversely, PA-X had no effect on

TAF7 levels. There were no significant changes in the RNA levels of the viral gene HA during

cycloheximide treatment, confirming that the decrease in PA-X activity is not the result of

changes in viral replication (Fig 5B). Thus, while we cannot confirm that PA-X protein levels

significantly decrease in infected cells after a translational block, we do observe markedly

reduced PA-X activity. Taken together with the decrease in both PA-X protein levels (Fig 1) and

PA-X activity (Fig 5A) during ectopic overexpression, our results indicate that the reduction in

PA-X activity during infection is likely the result of decreased PA-X protein levels. Therefore, it

is likely that PA-X is rapidly turned over during infection.

DISCUSSION

The PA-X protein is present at low levels in influenza A virus-infected cells, which has been

generally assumed to be due to its production mechanism based on ribosomal frameshifting.

However, we have found that PA-X protein levels are also regulated at the post-translational

level, and that PA-X is rapidly turned over (Fig 1, 3, 5). We show that PA-X aa 208-232 (aa 17-

41 of the X-ORF) contain a stability-regulating sequence, as truncating PA-X at aa 232 does not

change its half-life (Fig 4), while truncating it at aa 207 significantly increases it (Fig 2). Our

results also indicate that instability is a conserved property of PA-X protein, but that differences

in PA-X sequence among the strains lead to different half-lives (Fig 3). We propose that this

post-translational regulation results in low steady-state levels of PA-X protein in infected cells,

which in turn may limit the active pool of PA-X. Given that PA-X is already made at low levels

during infection, the instability of this protein suggests that limiting PA-X activity is of

paramount importance for viral replication.

It is still unknown how PA-X is degraded by the cell. We previously identified interactions

between the X-ORF of both PR8 and pH1N1 PA-X and host proteins involved in protein

turnover, including three components of the proteasome, PSME3, PSMD3, and PSMC5, and a

ubiquitin E3 ligase, HUWE1 (3). While components of the proteasome are often found as

contaminants in proteomics studies, the identification of an E3 ligase as an interactor increases

confidence in the identification of the proteasome and suggests a role for proteasomal

degradation. Understanding the role of HUWE1 and the ubiquitin-proteasome system in

regulation of PA-X protein levels is a current avenue of investigation.

It is attractive to think that by rapidly turning over the pool of active PA-X in the cell, instability

could serve as a mechanism to control PA-X host shutoff activity. Prolonged PA-X activity

could in theory have catastrophic consequences not only for the cell but also for successful viral

infection. We have observed that prolonged, high level PA-X expression kills human cells (data

not shown), as reported by Oishi *et al.* in yeast (13, 14). This cytopathic effect could be due to

the activation of cellular stress response pathways as a response to prolonged widespread RNA

degradation, or to a direct effect of the RNA degradation on the levels of unstable proteins that

keep the cell alive, like the Bcl-2 family member MCL-1 (33). During infection, this toxicity

could prematurely kill the infected cell before progeny virions are made, which would curtail

viral replication. In support of this, our group previously found that the X-ORF from PR8 PA-X

truncated to 16 amino acids may have stronger activity on a co-transfected reporter compared to

full length PR8 PA-X, although the difference was not statistically significant (3). Additionally,

the Takimoto lab showed that truncation of the pH1N1 PA-X X-ORF to 15 residues in the virus

(through a premature stop codon) increased host shutoff during infection (17). Thus, the X-ORF

may curtail PA-X activity by reducing its protein levels, although the link between PA-X

stability, host shutoff activity and virulence remains to be rigorously tested during infection.

Our study implicates the middle portion of the X-ORF in the regulation of PA-X stability. With

the exception of a few equine H7N7 and bat influenza strains, all influenza A strains have PA-X

with X-ORFs of 41 or 61 amino acids (16). However, there are other locations where a stop

codon could have evolved without changes to the PA reading frame (16), and only the first 15

amino acids of the X-ORF are required for host shutoff activity (3, 17, 22). Therefore, the fact

that sequences beyond aa 15 are nonetheless retained through evolution suggests that these

additional sequences have a necessary function despite being dispensable for activity. Our study

implies that the rest of the X-ORF may be needed to regulate PA-X, and in particular to ensure

that PA-X is rapidly turned over (Fig 2). A post-translational mechanism to control PA-X protein

levels may be important to limit activity, because in the context of infection PA-X does not

cleave its own transcript and thus does not autoregulate its expression (5). This is in contrast to

other viral host shutoff RNases like the alphaherpesvirus protein vhs, which is thought to be able

to target its own transcript (34). If hyperactive PA-X is detrimental to viral fitness, there may be

selective pressure to maintain the instability of PA-X, and thus retain the later portion of the X-

ORF even though it is not required for host shutoff activity.

It was surprising that the protein decay rate of pH1N1 PA-X was dramatically different from that

of most other strains (Fig 3). This is particularly striking given that the biggest difference

between pH1N1 and the other tested strains is the absence of the 20 distal amino acids, but we

find that this is not responsible for the reduced degradation (Fig 4). PA-X from strains with avian

origin PA segments, like that of pH1N1 (35), has been reported to have higher host shutoff

activity compared to PA-X from strains with human-adapted PA segments (23). This has raised

the possibility that PA-X host shutoff activity is linked to virulence and pathogenicity (36).

Given that pH1N1 PA-X accumulates to higher levels in the cell when compared to its

counterparts from PR8, Udorn, and HK06 (Fig 1 and 3), which are human-adapted, it is possible

that strain-specific differences in PA-X activity may be the result of strain-specific differences in

stability. This is an interesting relationship, especially given the higher virulence of a mouse

adapted pH1N1 strain when compared to PR8 (37). It will be important to assess the functional

consequences of PA-X stability and what it means for PA-X host shutoff activity and viral

pathogenesis.

Collectively our results reveal for the first time that PA-X is markedly unstable. The stability of

different PA-X variants may control host shutoff activity, which could have consequences for

virulence and pathogenesis. Since the region of PA-X regulating stability is maintained despite

not being required for host shutoff, having an unstable PA-X may be advantageous for the virus

and be selected for during evolution. Further elucidation of the mechanism of PA-X turnover in

the infected cell and its functional consequences will uncover important aspects of PA-X biology

and regulation of host shutoff, allowing us to further link PA-X activity with influenza A virus

virulence and pathogenesis.

ACKNOWLEDGEMENTS

We thank Dr. Seema Lakdawala, Dr. Jesse Bloom and Dr. Richard Webby for constructs, and

members of the Moore and Bohm lab and Malavika Raman for suggestions and feedback. We

also thank Dr. Denys Khaperskyy, Dr. Craig McCormick and members of the Gaglia lab for

suggestions, feedback, and critical reading of the manuscript. This work was supported by NIH

grant R01 AI137358 to M.M.G. R.E.L. was supported by NIH training grant T32 AI007422.

S.D.S. is a David J. Cosman, Ph.D. fellow in the Tufts Molecular Microbiology program.

MATERIALS AND METHODS

Plasmids

pCR3.1-PR8-PA-X-D108A-myc was previously described (5, 38). pCR3.1-PA-X D108A-myc

constructs for A/California/7/2009pdm, A/Udorn/307/1972, and A/Hong Kong/218847/2006

were generated from the corresponding previously described wild-type plasmids (3, 5) by site-

directed mutagenesis. The indicated PA-X coding regions were PCR-amplified using primers in

Table 2 and inserted into the SalI and XhoI sites of a pCR3.1-C-terminal-myc backbone.

pCR3.1-PA-X-D108A-myc constructs for A/Perth/16/2009 (Perth) was generated by subcloning

the PA-X coding region from the plasmid from segment 3 for this strain (a kind gift from Dr.

Seema Lakdawala). The Perth PA-X coding region was PCR-amplified using primers in Table 2

to delete a base (which mimics the frameshift) and insert the D108A mutation. It was then

inserted into the SalI and MluI sites of a pCR3.1-C-terminal-myc backbone. pCR3.1-PR8-PA-

X16-D108A-myc and pCR3.1-PR8-PA-X41-D108A-myc were generated by PCR-amplifying

the truncated coding region from pCR3.1-PA-X-PR8-D108A-myc using primers in Table 2. The

truncated regions were then inserted into the SalI and MluI sites of a pCR3.1-C-terminal-myc

backbone. Gibson cloning using HiFi assembly mix (New England Biolabs) was used to make

all constructs.

Cell lines, transfections, and drug treatments

Human embryonic kidney (HEK293T) cells (ATCC), Madin-Darby Canine Kidney (MDCK)

cells (ATCC), A549 (ATCC) cells and derivatives were cultured in Dulbecco's modified Eagle's

medium (DMEM; Life Technologies) supplemented with 10% fetal bovine serum (Hyclone) and

maintained at 37°C in 5% CO_2 atmosphere. A549 cell lines expressing PA-X under a

doxycycline-inducible promoter (A549-iPA-X_PR8_wt#1 and A549-iPA-X_PR8_D108A#8)

were previously described (5). For all 2-hour half-life experiments, HEK293T cells were plated

in 10-cm dishes and transfected with 12 μg total DNA (including 6 μg of PA-X construct) using

JetPrime (Polyplus transfection). 6 hours post transfection, cells were transferred to 6-well

plates, and one well was set up per indicated time point. For the indicated 4-hour cycloheximide

experiments, HEK293T cells were plated in 2 6-wells for each indicated strain and transfected

with 2 μg of DNA (including 1 μg of PA-X construct) per well using JetPrime (Polyplus

transfection). 6 hours post transfection, the 2 wells for each construct were trypsinized, mixed

together, and replated into 2 wells. 24 hours post transfection, cells were treated with 200 μg/mL

of cycloheximide (Sigma Aldrich) or dH$_2$O as vehicle control, and collected for RNA and

protein isolation at the indicated time points. For experiments using iPA-X cells, cells were

treated with 0.5 μg/mL doxycycline (Thermo Fisher) for 14 hours to induce PA-X expression

and then 200 μg/mL of cycloheximide (or vehicle control, dH$_2$O) for 2 hours prior to RNA

collection.

Viruses and Infections

Wild type influenza A virus A/Puerto Rico/8/1934 H1N1 (PR8) and the mutant PR8 ΔX were

generated as previously described in (3) using the 8-plasmid reverse genetic system (39). Viral

stocks were produced in MDCK cells and infectious titers were determined by plaque assay in

MDCKs using a 1.2% Avicel overlay described in Matrosovich *et al.* (40). A549 monolayers

were mock infected or infected with the wild-type or PA-X mutant viruses at a MOI = 5 for 1 hr

at 37°C. Monolayers were washed with PBS, and fresh infection media (0.5% Bovine Serum

Albumin (BSA; Sigma Aldrich) in DMEM supplemented with 0.5 μg/mL of L-(tosylamido-2-

phenyl) ethyl chloromethyl ketone (TPCK)-treated trypsin) (Sigma Aldrich) was added and cells

were incubated at 37°C in 5% CO_2 atmosphere for 24 hours prior to cycloheximide treatment.

RNA purification, cDNA preparation and RT-qPCR

RNA was extracted from cells and purified using the Quick-RNA Miniprep kit (Zymo Research),

following manufacturer's protocol. In all cases, the RNA was treated with Turbo DNase (Life

Technologies), then reverse transcribed using iScript supermix (Bio-Rad) per manufacturer's

protocol. qPCR was performed using iTaq Universal SYBR Green supermix (Bio-Rad), on the

Bio-Rad CFX Connect Real-Time System qPCR and analyzed with Bio-Rad CFX Manager 3.1

Program. The primers used are listed in Table 2.

Protein extraction, western blotting and antibodies

Total cellular protein was collected in protein lysis buffer (50 mM Tris-HCl pH 7.4, 150 mM

NaCl, 2 mM EDTA, 0.5% sodium deoxycholate, 0.5% SDS, 1% NP-40) supplemented with

cOmplete EDTA-free protease inhibitor cocktail (Roche) and 1 U/mL benzonase nuclease

(Sigma Aldrich). The benzonase is included to degrade DNA and promote release of PA-X from

the nucleus. Proteins were separated on 12% SDS-PAGE gels and transferred onto 0.22 μm

PVDF membranes (Millipore). Western Blots were performed in PBST with 0.5% milk using

antibodies against the Myc tag (1:1000, Cell Signaling Technologies 2276), and human β-tubulin

(1:1000, Cell Signaling Technologies 2128). Secondary antibodies conjugated to HRP were

purchased from Southern Biotechnologies (rabbit, OB4030-05; mouse, OB1031-05) and used at

1:5,000 dilution. Membranes were developed using Pierce ECL Western blotting substrate

(Thermo Fisher) and imaged with a Syngene G:Box Chemi XT4 gel doc system. Images were

quantified using GeneTools version 4.3.8.0

Statistical Analysis

All statistical analysis was performed using GraphPad Prism version 8.0 for Mac OS X,

(GraphPad Software, San Diego, CA, USA). For half-lives experiments, statistical significance

was determined by performing linear regression and comparing slopes and intercepts. For other

experiments, statistical significance was determined by Student's *t*-test, or analysis of variance

(ANOVA) followed by a *post-hoc* multiple comparisons test (Sidak's or Tukey's) as indicated. A

minimum of three biological replicates were conducted for all experiments.

References

1.   Jagger BW, Wise HM, Kash JC, Walters K-A, Wills NM, Xiao Y-L, Dunfee RL,

Schwartzman LM, Ozinsky A, Bell GL, Dalton RM, Lo A, Efstathiou S, Atkins JF, Firth

AE, Taubenberger JK, Digard P. 2012. An Overlapping Protein-Coding Region in

Influenza A Virus Segment 3 Modulates the Host Response. Science 337:199–204.

2.   Levene RE, Gaglia MM. 2018. Host Shutoff in Influenza A Virus: Many Means to an End.

Viruses 10:475.

3.   Gaucherand L, Porter BK, Levene RE, Price EL, Schmaling SK, Rycroft CH, Kevorkian Y,

McCormick C, Khaperskyy DA, Gaglia MM. 2019. The Influenza A Virus

Endoribonuclease PA-X Usurps Host mRNA Processing Machinery to Limit Host Gene

Expression. Cell Reports 27:776-792.e7.

4.   Chaimayo C, Dunagan M, Hayashi T, Santoso N, Takimoto T. Specificity and functional

interplay between influenza virus PA-X and NS1 shutoff activity 26.

5.   Khaperskyy DA, Schmaling S, Larkins-Ford J, McCormick C, Gaglia MM. 2016. Selective

Degradation of Host RNA Polymerase II Transcripts by Influenza A Virus PA-X Host

Shutoff Protein. PLOS Pathogens 12:e1005427.

6.   Gao H, Sun Y, Hu J, Qi L, Wang J, Xiong X, Wang Y, He Q, Lin Y, Kong W, Seng L-G,

Sun H, Pu J, Chang K-C, Liu X, Liu J. 2015. The contribution of PA-X to the virulence of

pandemic 2009 H1N1 and highly pathogenic H5N1 avian influenza viruses. Scientific

Reports 5:8262.

7.   Gong X-Q, Sun Y-F, Ruan B-Y, Liu X-M, Wang Q, Yang H-M, Wang S-Y, Zhang P,

Wang X-H, Shan T-L, Tong W, Zhou Y-J, Li G-X, Zheng H, Tong G-Z, Yu H. 2017. PA-X

protein decreases replication and pathogenicity of swine influenza virus in cultured cells

and mouse models. Veterinary Microbiology 205:66–70.

8.   Hayashi T, MacDonald LA, Takimoto T. 2015. Influenza A Virus Protein PA-X

Contributes to Viral Growth and Suppression of the Host Antiviral and Immune Responses.

Journal of Virology 89:6442–6452.

9.   Hu J, Mo Y, Wang X, Gu M, Hu Z, Zhong L, Wu Q, Hao X, Hu S, Liu W, Liu H, Liu X,

Liu X. 2015. PA-X Decreases the Pathogenicity of Highly Pathogenic H5N1 Influenza A

Virus in Avian Species by Inhibiting Virus Replication and Host Response. Journal of

Virology 89:4126–4142.

10.   Xu G, Zhang X, Liu Q, Bing G, Hu Z, Sun H, Xiong X, Jiang M, He Q, Wang Y, Pu J, Guo

X, Yang H, Liu J, Sun Y. 2017. PA-X protein contributes to virulence of triple-reassortant

H1N2 influenza virus by suppressing early immune responses in swine. Virology 508:45–

53.

11.   Sun Y, Hu Z, Zhang X, Chen M, Wang Z, Xu G, Bi Y, Tong Q, Wang M, Sun H, Pu J,

Iqbal M, Liu J. 2020. R195K mutation in the PA-X protein increases the virulence and

transmission of influenza A virus in mammalian hosts. J Virol JVI.01817-19,

jvi;JVI.01817-19v1.

12.   Liu L, Song S, Shen Y, Ma C, Wang T, Tong Q, Sun H, Pu J, Iqbal M, Liu J, Sun Y. 2020.

Truncation of PA-X Contributes to Virulence and Transmission of H3N8 and H3N2 Canine

Influenza Viruses in Dogs. J Virol 94:e00949-20, /jvi/94/15/JVI.00949-20.atom.

13.   Oishi K, Yamayoshi S, Kozuka-Hata H, Oyama M, Kawaoka Y. 2018. N-Terminal

Acetylation by NatB Is Required for the Shutoff Activity of Influenza A Virus PA-X. Cell

Reports 24:851–860.

14.   Oishi K, Yamayoshi S, Kawaoka Y. 2018. Identification of novel amino acid residues of

influenza virus PA-X that are important for PA-X shutoff activity by using yeast. Virology

516:71–75.

15.   Firth AE, Jagger BW, Wise HM, Nelson CC, Parsawar K, Wills NM, Napthine S,

Taubenberger JK, Digard P, Atkins JF. 2012. Ribosomal frameshifting used in influenza A

virus expression occurs within the sequence UCC_UUU_CGU and is in the +1 direction.

Open Biology 2:120109.

16.   Shi M, Jagger BW, Wise HM, Digard P, Holmes EC, Taubenberger JK. 2012. Evolutionary

Conservation of the PA-X Open Reading Frame in Segment 3 of Influenza A Virus. Journal

of Virology 86:12411–12413.

17.   Hayashi T, Chaimayo C, McGuinness J, Takimoto T. 2016. Critical Role of the PA-X C-

Terminal Domain of Influenza A Virus in Its Subcellular Localization and Shutoff Activity.

J Virol 90:7131–7141.

18. Košík I, Práznovská M, Košíková M, Bobišová Z, Hollý J, Varečková E, Kostolanský F,

Russ G. 2015. The Ubiquitination of the Influenza A Virus PB1-F2 Protein Is Crucial for

Its Biological Function. PLoS ONE 10:e0118477.

19. Varga ZT, Palese P. 2011. The influenza A virus protein PB1-F2. Virulence 2:542–546.

20. Varga ZT, Ramos I, Hai R, Schmolke M, García-Sastre A, Fernandez-Sesma A, Palese P.

2011. The Influenza Virus Protein PB1-F2 Inhibits the Induction of Type I Interferon at the

Level of the MAVS Adaptor Protein. PLoS Pathog 7:e1002067.

21. Varga ZT, Grant A, Manicassamy B, Palese P. 2012. Influenza Virus Protein PB1-F2

Inhibits the Induction of Type I Interferon by Binding to MAVS and Decreasing

Mitochondrial Membrane Potential. J Virol 86:8359–8366.

22. Oishi K, Yamayoshi S, Kawaoka Y. 2015. Mapping of a Region of the PA-X Protein of

Influenza A Virus That Is Important for Its Shutoff Activity. Journal of Virology 89:8661–

8665.

23. Desmet EA, Bussey KA, Stone R, Takimoto T. 2013. Identification of the N-Terminal

Domain of the Influenza Virus PA Responsible for the Suppression of Host Protein

Synthesis. Journal of Virology 87:3108–3118.

24. Nogales A, Martinez-Sobrido L, Chiem K, Topham DJ, DeDiego ML. 2018. Functional

Evolution of the 2009 Pandemic H1N1 Influenza Virus NS1 and PA in Humans. Journal of

Virology 92:e01206-18.

25.   Santos A, Pal S, Chacón J, Meraz K, Gonzalez J, Prieto K, Rosas-Acosta G. 2013.

504   SUMOylation Affects the Interferon Blocking Activity of the Influenza A Nonstructural

505   Protein NS1 without Affecting Its Stability or Cellular Localization. Journal of Virology

506   87:5602–5620.

507   26.   Cheng Y-Y, Yang S-R, Wang Y-T, Lin Y-H, Chen C-J. 2017. Amino Acid Residues 68–71

508   Contribute to Influenza A Virus PB1-F2 Protein Stability and Functions. Front Microbiol

509   8:692.

510   27.   Chen W, M. Smeekens J, Wu R. 2016. Systematic study of the dynamics and half-lives of

511   newly synthesized proteins in human cells. Chemical Science 7:1393–1400.

512   28.   Gao H, Sun H, Hu J, Qi L, Wang J, Xiong X, Wang Y, He Q, Lin Y, Kong W, Seng L-G,

513   Pu J, Chang K-C, Liu X, Liu J, Sun Y. 2015. Twenty amino acids at the C-terminus of PA-

514   X are associated with increased influenza A virus replication and pathogenicity. Journal of

515   General Virology 96:2036–2049.

516   29.   Feng KH, Sun M, Iketani S, Holmes EC, Parrish CR. 2016. Comparing the functions of

517   equine and canine influenza H3N8 virus PA-X proteins: Suppression of reporter gene

518   expression and modulation of global host gene expression. Virology 496:138–146.

519   30.   Rigby RE, Wise HM, Smith N, Digard P, Rehwinkel J. 2019. PA-X antagonises MAVS-

520   dependent accumulation of early type I interferon messenger RNAs during influenza A

521   virus infection. Sci Rep 9:7216.

522   31.   Hara K, Schmidt FI, Crow M, Brownlee GG. 2006. Amino Acid Residues in the N-

523   Terminal Region of the PA Subunit of Influenza A Virus RNA Polymerase Play a Critical

Role in Protein Stability, Endonuclease Activity, Cap Binding, and Virion RNA Promoter

Binding. Journal of Virology 80:7789–7798.

32. Maier HJ, Kashiwagi T, Hara K, Brownlee GG. 2008. Differential role of the influenza A

virus polymerase PA subunit for vRNA and cRNA promoter binding. Virology 370:194–

204.

33. Adams KW, Cooper GM. 2007. Rapid Turnover of Mcl-1 Couples Translation to Cell

Survival and Apoptosis. J Biol Chem 282:6192–6200.

34. Elgadi MM, Hayes CE, Smiley JR. 1999. The Herpes Simplex Virus vhs Protein Induces

Endoribonucleolytic Cleavage of Target RNAs in Cell Extracts. J Virol 73:7153–7164.

35. Garten RJ, Davis CT, Russell CA, Shu B, Lindstrom S, Balish A, Sessions WM, Xu X,

Skepner E, Deyde V, Okomo-Adhiambo M, Gubareva L, Barnes J, Smith CB, Emery SL,

Hillman MJ, Rivailler P, Smagala J, Graaf M de, Burke DF, Fouchier RAM, Pappas C,

Alpuche-Aranda CM, López-Gatell H, Olivera H, López I, Myers CA, Faix D, Blair PJ, Yu

C, Keene KM, Dotson PD, Boxrud D, Sambol AR, Abid SH, George KS, Bannerman T,

Moore AL, Stringer DJ, Blevins P, Demmler-Harrison GJ, Ginsberg M, Kriner P,

Waterman S, Smole S, Guevara HF, Belongia EA, Clark PA, Beatrice ST, Donis R, Katz J,

Finelli L, Bridges CB, Shaw M, Jernigan DB, Uyeki TM, Smith DJ, Klimov AI, Cox NJ.

2009. Antigenic and Genetic Characteristics of Swine-Origin 2009 A(H1N1) Influenza

Viruses Circulating in Humans. Science 325:197–201.

36. Lutz MM, Dunagan MM, Kurebayashi Y, Takimoto T. 2020. Key Role of the Influenza A

Virus PA Gene Segment in the Emergence of Pandemic Viruses. 4. Viruses 12:365.

37.   Meunier I, Morisseau O, Garneau É, Marois I, Cloutier A, Richter MV. 2015. Infection

with a Mouse-Adapted Strain of the 2009 Pandemic Virus Causes a Highly Severe Disease

Associated with an Impaired T Cell Response. PLoS One 10.

38.   Khaperskyy DA, Emara MM, Johnston BP, Anderson P, Hatchette TF, McCormick C.

2014. Influenza A Virus Host Shutoff Disables Antiviral Stress-Induced Translation Arrest.

PLOS Pathogens 10:e1004217.

39.   Hoffmann E, Neumann G, Kawaoka Y, Hobom G, Webster RG. 2000. A DNA transfection

system for generation of influenza A virus from eight plasmids. Proceedings of the National

Academy of Sciences 97:6108–6113.

40.   Matrosovich M, Matrosovich T, Garten W, Klenk H-D. 2006. New low-viscosity overlay

medium for viral plaque assays. Virol J 3:63.

 FIGURE AND TABLE LEGENDS

TABLE 1: HALF-LIVES OF PA-X FROM PR8, pH1N1, UDORN, HK06, AND PERTH. A

linear regression was carried out on the quantification plots in Figure 3. The slope of the linear

regression line and the calculated half-life are reported. The half-life was calculated as the

negative of the inverse of the slope in minutes. Where the 1/slope value yielded a half-life over 2

hours, the calculated values are expected to be less accurate. The slope and half-life for PR8 PA-

X was calculated based on regression analysis from Figure 1B.

TABLE 2: PRIMERS USED IN THIS STUDY. Primers used in this study for qPCR and

cloning.

FIGURE 1: PA-X is unstable.

(A-C) Protein and RNA samples were collected from HEK293T cells expressing the catalytically

inactive PR8 PA-X at the indicated time points after treatment with 200 μg/mL of cycloheximide

(CHX) to inhibit translation. (-) indicates vehicle-treated cells collected at 120 minutes. (A)

Representative western blot using anti-myc antibodies to detect myc-tagged PA-X and tubulin as

a loading control. (B) Plot of the relative levels of PA-X after cycloheximide treatment. For each

replicate, PA-X protein levels were normalized to tubulin protein levels and reported as the $\log_2$

transformed ratio relative to time 0. The line represents the linear regression and the shaded

region its 95% confidence interval. n = 7 (C) Levels of PA-X mRNA were measured by RT-

qPCR. After normalization to 18S rRNA, levels were plotted as amounts relative to time 0. n = 3.

FIGURE 2: The X-ORF mediates PA-X instability

(A) Schematic diagram of PA-X domains and PR8 PA-X16 truncation used. (B-D) Protein and

RNA samples were collected from HEK293T cells expressing the catalytically inactive full-

length or truncated C-terminally myc-tagged PA-X at the indicated time points after treatment

with 200 μg/mL of cycloheximide (CHX) to inhibit translation. (-) indicates vehicle-treated cells

collected at 120 minutes. (B) Representative western blot using anti-myc antibodies to detect

myc-tagged PA-X and tubulin as a loading control. (C) Plot of the relative levels of PA-X after

cycloheximide treatment. For each replicate, PA-X protein levels were normalized to tubulin

protein levels and reported as the $\log_2$ transformed ratio from time 0. A linear regression was

performed for each dataset and the *p* value calculated relative to PR8 PA-X61 (full length PA-

X). The lines represent the linear regression and the shaded regions indicate the 95% confidence

interval (PA-X61: slope = -0.72, confidence interval = -1.04 – -0.39; PA-X16: slope = 0.34,

confidence interval = 0.04 – 0.64). The half-life ($t_{1/2}$) for PA-X61 was calculated as the negative

inverse of the slope and is reported in minutes. As the slope for PA-X16 is positive, the half-life

of this protein cannot be calculated and is therefore reported simply as longer than the time

course of the experiment. n ≥ 3 (D) Levels of PA-X16 mRNA were measured by RT-qPCR.

After normalization to 18S rRNA, levels are plotted as amounts relative to time 0. n > 3.

FIGURE 3: The stability of PA-X varies among influenza A virus strains.

Protein and RNA samples were collected from HEK293T cells expressing the catalytically

inactive C-terminally myc tagged PA-X from indicated strain at the indicated time points (A-L)

or 4 hours (M-O) after treatment with 200 μg/mL of cycloheximide (CHX) to inhibit translation.

(-) indicates vehicle-treated cells collected at 120 minutes (A-L) or 4 hours (M-O). (A, D, G, J,

M) Representative western blots using anti-myc antibodies to detect myc-tagged PA-X and

tubulin as a loading control. (B, E, H, K) Plot of the relative levels of PA-X after cycloheximide

treatment. For each replicate, PA-X protein levels were normalized to tubulin protein levels and

reported as the $\log_2$ transformed ratio from time zero. The line represents the linear regression

and the shaded region its 95% confidence interval. n = 3 for each variant. In K, the PR8 PA-X

dataset from Figure 1 is included for comparison and the *p* value relative to PR8 PA-X is

reported. Two of the three pH1N1 PA-X replicates were performed in parallel with PR8 PA-X.

(N) Plot of average change in PA-X levels after 4-hour cycloheximide treatment. For each

replicate, protein levels of PA-X were normalized to tubulin protein levels and reported as the

fold change vs. vehicle-treated PA-X expressing cells. * = $p < 0.05$, Student's *t*-test. n = 3 . (C, F,

I, L, O) Levels of PA-X mRNA were measured by RT-qPCR. After normalization to 18S rRNA,

levels are plotted as amounts relative to time 0 (C, F, I, L) or as arbitrary units (O). n = 3.

FIGURE 4: The length difference between the two major variants of the PA-X X-ORF does

not account for differences in PA-X instability.

(A) Schematic diagram of PA-X domains and PR8 PA-X41 truncation used. (B-D) Protein and

RNA samples were collected from HEK293T cells expressing the catalytically inactive truncated

or full-length C-terminally myc-tagged PA-X at the indicated time points after treatment with

200 μg/mL of cycloheximide (CHX) to inhibit translation. (-) indicates vehicle-treated cells

collected at 120 minutes. (B) Representative western blot using anti-myc antibodies to detect

myc-tagged PA-X and tubulin as a loading control. (C) Plot of the relative levels of PA-X after

cycloheximide treatment. For each replicate, PA-X protein levels were normalized to tubulin

protein levels and reported as the $\log_2$ transformed ratio from time zero. A linear regression was

performed for each dataset and the *p* value calculated relative to PR8 PA-X61 (full length PA-

X). The lines represent the linear regression and the shaded region indicates the 95% confidence

interval (PA-X61: slope = -0.58; confidence interval: -0.87 – -0.28; PA-X41: slope = -0.61,

confidence interval = -0.84 – -0.38). $n \geq 3$. The half-life (t1/2) values were calculated as the

negative inverse of the slope and are reported in minutes. (D) Levels of PA-X41 mRNA were

measured by RT-qPCR. After normalization to 18S rRNA, levels are plotted as amounts relative

to time 0. $n > 3$.

FIGURE 5: Translation inhibition reduces PA-X-mediated host shutoff, suggesting PA-X is

likely unstable during infection

(A) Levels of the indicated endogenous RNAs were measured by RT-qPCR in A549 "iPA-X"

cells, which express wild-type (wt) PR8 PA-X or a PR8 PA-X catalytic mutant (D108A) under a

doxycycline inducible promoter. RNA samples were collected after 14 hours of PA-X induction

with doxycyline followed by a 2-hour treatment with 200 μg/mL of cycloheximide (CHX) or

vehicle control. After normalization to 18S rRNA, mRNA levels are plotted relative to

uninduced wt iPA-X cells (G6PD, TAF7) or relative to induced wt iPA-X cells (PA-X) on a

logarithmic scale. $n \geq 3$. *** = $p < 0.001$, ns = $p > 0.05$ 3-way ANOVA followed Tukey's

multiple comparisons test. (B) Levels of the indicated endogenous human mRNAs (G6PD and

TAF7) and viral RNAs (HA) were measured by RT-qPCR in cells infected with wt PR8 or PR8

ΔX at a MOI of 5 for 24 hours prior to a 2-hour treatment with 200 μg/mL of cycloheximide

(CHX) to inhibit translation or a vehicle control. After normalization to 18S rRNA, RNA levels

are plotted relative to mock infected cells (G6PD, TAF7) or as amounts normalized to cellular

18S rRNA (HA) on a logarithmic scale. The dashed lines indicate RNA levels in mock infected

cells. n = 3. * = $p < 0.05$, ns = $p > 0.05$, 2-way ANOVA followed by Sidak's multiple

comparisons test.

Table 1: Half-lives of PA-X from PR8, pH1N1, Udorn, HK06, and Perth

PA-X	Slope of linear regression	Half-Life in Minutes	95% confidence interval on slope
A/Puerto Rico/8/1934	-0.70	86	-0.90 to -0.50
A/California/7/2009pdm pH1N1	-0.11	546	-0.35 to 0.12
A/Udorn/307/1973 H3N2	-1.77	34	-2.41 to -1.14
A/Perth/16/2009 H3N2	-0.24	250	-0.78 to 0.30
A/Hong Kong/218847/2006 H1N1	-0.73	83	-1.07 to -0.38

654   **Table 2: Primers used in this study**

Purpose	Sequence (5' - 3')
18S qPCR Forward	GTAACCCGTTGAACCCCATT
18S qPCR Reverse	CCATCCAATCGGTAGTAGCG
PA-X qPCR Forward- amplifies BGH 3' UTR region on the vector (Fig 1-4)	GAGTCTAGAGGGCCCGTTTA
PA-X qPCR Reverse- amplifies BGH 3' UTR region on the vector (Fig 1-4)	GAGGGGCAAACAACAGATGG
PA-X qPCR Forward - Fig 5	GCGACAATGCTTCAATCCGA
PA-X qPCR Reverse - Fig 5	TTGACTCGCCTTGCTCATTG
HA qPCR Forward	CTGGACCTTGCTAAAACCCG
HA qPCR Reverse	TCTGGAAAGGGAGACTGCTG
G6PD qPCR Forward	TGAGCCAGATAGGCTGGAA
G6PD qPCR Reverse	TAACGCAGGCGATGTTGTC
TAF7 qPCR Forward	TAGCACTGATCCTAAAGCAAGC
TAF7 qPCR Reverse	GGCAGAGTAATTCCGTGGTTC
Clone PA-X Variants -amplifies CDS starting	CTTAAGCTTGGTACCGGAGTACTGG

from 5' end vector sequences to clone into pCR3.1 SalI site	
Clone PA-X Variants -amplifies CDS at BGH 3' UTR to clone into pCR3.1 XhoI site	GTTTAAACGGGCCCTCTAGACTCGA
Make D108A A/HongKong/218847/2006 - mutates active site (Fwd)	CAGAGCTGGCAGAAACTTTGGTTTCTC AGCT
Make D108A A/HongKong/218847/2006 - mutates active site (Rev)	GCCAGCTCTGTATGATTACAAAGAGAA TAGA
Make D108A A/California/7/2009pdm- mutates active site (Fwd)	GTCATACAAAGCAGGAAGAAATTTAG GCTTC
Make D108A A/California/7/2009pdm- mutates active site (Rev)	TTTCTTCCTGCTTTGTATGACTACAAAG AG
Make D108A A/Udorn/307/1972-mutates active site (Fwd)	ATCATACAAAGCTGGCAGAAACTTCGG
Make D108A A/Udorn/307/1972-mutates active site (Rev)	GTTTCTGCCAGCTTTGTATGATTACAA GG
Clone A/Perth/16/2009 PA-X (wt and D108A) - amplifies from 5' UTR to clone into pCR3.1 SalI site	TAAACTTAAGCTTGGTACCGGAGTACT GGTCGACCCTCCG
Make D108A A/Perth/16/2009 - mutates active site (Rev)	AATCATACAAAGCTGGTAGAAACTTCG G
Make D108A A/Perth/16/2009 - mutates active site (Fwd)	TTTCTACCAGCTTTGTATGATTACAAGG AG
Clone A/Perth/16/2009 PA-X (wt and D108A) - inserts deletion for frameshifting (Rev)	GGACTGACGAAGGAATCCCATAGG
Clone A/Perth/16/2009 PA-X (wt and D108A) - inserts deletion for frameshifting (Fwd)	CCTATGGGATTCCTTCGTCAGTCC
Clone Make D108A A/Perth/16/2009 PA-X (wt and D108A) - amplifies at end of CDS to clone into pCR3.1 MluI site	CTGAGATCAGCTTCTGCTCCACGCGTGA CTTCTTTGGACATTTGAGAAAGCTTGC
Make PR8 PA-X15 - amplifies at 3' end of truncated CDS to clone into pCR3.1 MluI site	ATCAGCTTCTGCTCCACGCGTGATTTC AAACCTTTCTTCAATTG
Make PR8 PA-X41 - amplifies at 3' end of truncated CDS to clone into pCR3.1 MluI site	ATCAGCTTCTGCTCCACGCGTGACATA GGCTCTAAAATTTTCAA

Figure 1

A

A/Puerto Rico/8/1934 (PR8)

Minutes post CHX: 0   15   30   60   120   (-)

PA-X-myc

Tubulin

B   *A/Puerto Rico/8/1934 (PR8)*   **C**   *PR8 PA-X mRNA Levels*

FIGURE 1: PA-X is unstable. (A-C) Protein and RNA samples were collected from HEK293T cells expressing the catalytically inactive PR8 PA-X at the indicated time points after treatment with 200 µg/mL of cycloheximide (CHX) to inhibit translation. (-) indicates vehicle-treated cells collected at 120 minutes. (A) Representative western blot using anti-myc antibodies to detect myc-tagged PA-X and tubulin as a loading control. (B) Plot of the relative levels of PA-X after cycloheximide treatment. For each replicate, PA-X protein levels were normalized to tubulin protein levels and reported as the log2 transformed ratio relative to time 0. The line represents the linear regression and the shaded region its 95% confidence interval. n = 7 (C) Levels of PA-X mRNA were measured by RT-qPCR. After normalization to 18S rRNA, levels were plotted as amounts relative to time 0. n = 3.

Figure 2

FIGURE 2: The X-ORF mediates PA-X instability
(A) Schematic diagram of PA-X domains and PR8 PA-X16 truncation used. (B-D) Protein and RNA samples were collected from HEK293T cells expressing the catalytically inactive full-length or truncated C-terminally myc-tagged PA-X at the indicated time points after treatment with 200 μg/mL of cycloheximide (CHX) to inhibit translation. (-) indicates vehicle-treated cells collected at 120 minutes. (B) Representative western blot using anti-myc antibodies to detect myc-tagged PA-X and tubulin as a loading control. (C) Plot of the relative levels of PA-X after cycloheximide treatment. For each replicate, PA-X protein levels were normalized to tubulin protein levels and reported as the log2 transformed ratio from time 0. A linear regression was performed for each dataset and the p value calculated relative to PR8 PA-X61(full length PA-X). The lines represent the linear regression and the shaded regions indicate the 95% confidence interval (PA-X61: slope = -0.72, confidence interval = -1.04 – -0.39; PA-X16: slope = 0.34, confidence interval = 0.04 –0.64). The half-life (t1/2) for PA-X61 was calculated as the negative inverse of the slope and is reported in minutes. As the slope for PA-X16 is positive, the half-life of this protein cannot be calculated and is therefore reported simply as longer than the time course of the experiment. n ≥ 3 (D) Levels of PA-X16 mRNA were measured by RT-qPCR. After normalization to 18S rRNA, levels are plotted as amounts relative to time 0. n > 3.

Figure 3

FIGURE 3: The stability of PA-X varies among influenza A virus strains.
Protein and RNA samples were collected from HEK293T cells expressing the catalytically inactive C-terminally myc tagged PA-X from indicated strain at the indicated time points (A-L) or 4 hours (M-O) after treatment with 200 μg/mL of cycloheximide (CHX) to inhibit translation. (-) indicates vehicle-treated cells collected at 120 minutes (A-L) or 4 hours (M-O). (A, D, G, J, M) Representative western blot using anti-myc antibodies to detect myc-tagged PA-X and tubulin as a loading control. (B, E, H, K) Plot of the relative levels of PA-X after cycloheximide treatment. For each replicate, PA-X protein levels were normalized to tubulin protein levels and reported as the log2 transformed ratio from time zero. The line represents the linear regression and the shaded region its 95% confidence interval. n = 3 for each variant. In K, the PR8 PA-X dataset from Figure 1 is included for comparison and the reported p value is relative to PR8 PA-X. Two of the three pH1N1 PA-X replicates were performed in parallel with PR8 PA-X. (N) Plot of average change in PA-X levels after 4-hour cycloheximide treatment. For each replicate, protein levels of PA-X were normalized to tubulin protein levels and reported as the fold change vs. vehicle-treated PA-X expressing cells. * = p < 0.05, Student's t-test. n = 3. (C, F, I, L, O) Levels of PA-X mRNA were measured by RT-qPCR. After normalization to 18S rRNA, levels are plotted as amounts relative to time 0 (C, F, I, L) or as arbitrary units (O). n = 3.

Figure 4

FIGURE 4: The length difference between the two major variants of the PA-X X-ORF does not account for differences in PA-X instability.

(A) Schematic diagram of PA-X domains and PR8 PA-X41 truncation used. (B-D) Protein and RNA samples were collected from HEK293T cells expressing the catalytically inactive truncated or full-length C-terminally myc-tagged PA-X at the indicated time points after treatment with 200 μg/mL of cycloheximide (CHX) to inhibit translation. (-) indicates vehicle-treated cells collected at 120 minutes. (B) Representative western blot using anti-myc antibodies to detect myc-tagged PA-X and tubulin as a loading control. (C) Plot of the relative levels of PA-X after cycloheximide treatment. For each replicate, PA-X protein levels were normalized to tubulin protein levels and reported as the log2 transformed ratio from time zero. A linear regression was performed for each dataset and the p value calculated relative to PR8 PA-X61 (full length PA-X). The lines represent the linear regression and the shaded region indicates the 95% confidence interval (PA-X61: slope = -0.58; confidence interval: -0.87 – -0.28; PA-X41: slope = -0.61, confidence interval = -0.84 – -0.38). n ≥ 3. The half-life (t1/2) values were calculated as the negative inverse of the slope and are reported in minutes. (D) Levels of PA-X41 mRNA were measured by RT-qPCR. After normalization to 18S rRNA, levels are plotted as amounts relative to time 0. n > 3.

Figure 5

A *iPA-X cell line:*

FIGURE 5: Translation inhibition reduces PA-X-mediated host shutoff, suggesting PA-X is likely unstable during infection. (A) Levels of the indicated endogenous RNAs were measured by RT-qPCR in A549 "iPA-X" cells, which express wild-type (wt) PR8 PA-X or a PR8 PA-X catalytic mutant (D108A) under a doxycycline inducible promoter. RNA samples were collected after 14 hours of PA-X induction with doxycyline followed by a 2-hour treatment with 200 µg/mL of cycloheximide (CHX) or vehicle control. After normalization to 18S rRNA, mRNA levels are plotted relative to uninduced wt iPA-X cells (G6PD, TAF7) or relative to induced wt iPA-X cells (PA-X) on a logarithmic scale. n ≥ 3. *** = $p < 0.001$, ns = $p > 0.05$ 3-way ANOVA followed Tukey's multiple comparisons test. (B) Levels of the indicated endogenous human mRNAs (G6PD and TAF7) and viral RNAs (HA) were measured by RT-qPCR in cells infected with the wt PR8 or PR8 ΔX at a MOI of 5 for 24 hours prior to a 2-hour treatment with 200 µg/mL of cycloheximide (CHX) to inhibit translation or a vehicle control. After normalization to 18S rRNA, RNA levels are plotted relative to mock infected cells (G6PD, TAF7) or as amounts normalized to cellular 18S rRNA (HA) on a logarithmic scale. The dashed lines indicate RNA levels in mock infected cells. n = 3. * = $p < 0.05$, ns = $p > 0.05$, 2-way ANOVA followed by Sidak's multiple comparisons test.