Breadth and function of antibody response to acute SARS-CoV-2 infection in humans

Kuan-Ying A. Huang[1,2†], Tiong Kit Tan[3#], Ting-Hua Chen[4], Chung-Guei Huang[1,5], Ruth Harvey[6], Saira Hussain[6], Cheng-Pin Chen[7], Adam Harding[8], Javier Gilbert-Jaramillo[8], Xu Liu[8], Michael Knight[8], Lisa Schimanski[3,9], Shin-Ru Shih[1,5], Yi-Chun Lin[10], Chien-Yu Cheng[7], Shu-Hsing Cheng[10], Yhu-Chering Huang[2], Tzou-Yien Lin[2], Jia-Tsrong Jan[4], Che Ma[4], William James[8], Rodney S. Daniels[6], John W. McCauley[6], Pramila Rijal[3,9#], Alain R. Townsend[3,9†]

[1] Research Center for Emerging Viral Infections, College of Medicine, Chang Gung University, Taoyuan, Taiwan

[2] Division of Pediatric Infectious Diseases, Department of Pediatrics, Chang Gung Memorial Hospital, Taoyuan, Taiwan

[3] MRC Human Immunology Unit, Weatherall Institute of Molecular Medicine, University of Oxford, John Radcliffe Hospital, Oxford, UK

[4] Genomics Research Center, Academia Sinica, Taipei, Taiwan

[5] Department of Laboratory Medicine, Chang Gung Memorial Hospital, Taoyuan, Taiwan

[6] Worldwide Influenza Centre, The Francis Crick Institute, 1 Midland Road, London NW1 1AT, UK

[7] Department of Infectious Diseases, Taoyuan General Hospital, Ministry of Health

and Welfare, Taoyuan, and National Yang-Ming University, Taipei, Taiwan

[8] Sir William Dunn School of Pathology, University of Oxford, Oxford, UK

[9] Centre for Translational Immunology, Chinese Academy of Medical Sciences

Oxford Institute, University of Oxford, Oxford, UK

[10] Department of Infectious Diseases, Taoyuan General Hospital, Ministry of Health

and Welfare, Taoyuan, and Taipei Medical University, Taipei, Taiwan

Tiong   Kit   Tan   and   Pramila   Rijal   contributed   equally.

† Correspondence: Kuan-Ying A. Huang (arthur1726@cgmh.org.tw), or Alain R.

Townsend (alain.townsend@imm.ox.ac.uk)

Abstract

Serological and plasmablast responses and plasmablast-derived IgG monoclonal antibodies (MAbs) have been analysed in three COVID-19 patients with different clinical severities. Potent humoral responses were detected within 3 weeks of onset of illness in all patients and the serological titre was elicited soon after or concomitantly with peripheral plasmablast response. An average of 13.7% and 13.0% of plasmablast-derived MAbs were reactive with virus spike glycoprotein or nucleocapsid, respectively. A subset of anti-spike (10 of 32) and over half of anti-nucleocapsid (19 of 35) antibodies cross-reacted with other betacoronaviruses tested and harboured extensive somatic mutations, indicative of an expansion of memory B cells upon SARS-CoV-2 infection. Fourteen of 32 anti-spike MAbs, including five anti-RBD, three anti-non-RBD S1 and six anti-S2, neutralised wild-type SARS-CoV-2 in independent assays. Anti-RBD MAbs were further grouped into four cross-inhibiting clusters, of which six antibodies from three separate clusters blocked the binding of RBD to ACE2 and five were neutralising. All ACE2-blocking anti-RBD antibodies were isolated from two patients with prolonged fever, which is compatible with substantial ACE2-blocking response in their sera. At last, the identification of non-competing pairs of neutralising antibodies would offer potential templates for the development of prophylactic and therapeutic agents against SARS-CoV-2.

Introduction

In late 2019, a novel coronavirus emerged and was identified as the cause of a cluster of respiratory infection cases in Wuhan, China. It spread quickly around the world. In March of 2020 a pandemic was declared by the World Health Organization, the virus was formally named as Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) and the resulting disease was named COVID-19. As of 1 October 2020, there have been over 33 million confirmed cases of SARS-CoV-2 infection with 1,010,986 deaths (World Health Organization, https://covid19.who.int/).

There is no fully effective drug or licenced vaccine for COVID-19. Antibodies neutralise SARS-CoV-2 *in vitro*, offering hope that a vaccine may induce a protective response, and antibodies may be an effective treatment for COVID-19 in clinical practice. Convalescent plasma is being tested in ongoing clinical trials as a therapy for COVID-19 (1, 2), and was previously used in the treatment of SARS (3). The virus spike glycoprotein is a target of neutralising antibodies, which makes it a key candidate for both vaccine development and immunotherapy (4). B cell responses in COVID-19 patients have been detected concomitantly with follicular helper T cell responses from week one after illness onset (5). In SARS patients, B cell responses typically arise first against the nucleocapsid protein then, within four to eight days after symptom onset, antibody responses to spike glycoprotein have been found; neutralising antibody responses begin to develop by week two, and most patients develop neutralising antibodies by week three (6). Two serological studies of COVID-19 patients showed anti-SARS-CoV-2 IgG seroconversion at week three after onset and some cross-reactivity to nucleocapsid of SARS (7, 8).

Antibodies may play a role in protection against SARS-CoV-2 infection. The

underlying B-cell response leading to the rapid production of plasmablasts (antibody-

secreting cells) that secrete antibodies upon natural SARS-CoV-2 exposure/infection

is only beginning to be understood (5). Here, we characterised the infection-induced

serological and plasmablast responses and the derived IgG anti-SARS-CoV-2 spike

glycoprotein and nucleocapsid monoclonal antibodies (MAbs) from adult patients

with laboratory-confirmed COVID-19. The antigenic specificity and breadth of

antibodies and the sequence of their variable domains have been characterised in

detail. Virus neutralising antibodies were detected that bound epitopes on receptor-

binding domain (RBD), non-RBD regions of the S1 polypeptide, and the S2

polypeptide of the spike glycoprotein.

Results

Serological response and anti-spike glycoprotein and anti-nucleocapsid IgG antibodies from circulating plasmablasts

Serum IgG antibodies to the virus spike glycoprotein and the isolated RBD were analysed by indirect enzyme-linked immunosorbent and flow cytometry assays in three patients with laboratory-confirmed COVID-19. The clinical characteristics of the three patients studied are shown in Supplemental Table 1. Antibodies to spike glycoprotein and RBD were detected in all three patients after week 3 of illness onset (Figure 1a). Case A showed a robust response to spike glycoprotein and the isolated RBD by day (D) 22 (D22). Longitudinal sera from case B and C showed lower anti-spike glycoprotein and RBD IgG titres at week 1 or the beginning of week 2 and an elevated titre that peaked at the end of week 2 through week 3 (a peak 50% effective dilution (ED_{50}) titre to RBD 1:1,051 at D18 in case B and 1:588 at D14 in case C). Case B had prolonged fever and developed pneumonia at the end of week 2, which was followed by a robust increase of anti-spike glycoprotein and RBD IgG titres at week 3 (Figure 1a, Supplemental Table 1). By contrast, case C experienced a two-day course of febrile illness and reduction of all symptoms within the first week, followed by anti-spike glycoprotein and RBD IgG titres that peaked earlier at the end of week 2 (Figure 1a, Supplemental Table 1).

An increased frequency of circulating plasmablasts was detected in all three patients (healthy adults baseline less than 1%) (5) (Figure 1b). In case A, a plasmablast response containing a substantial IgG subset was detected at the end of week 2 (D14), followed by a high serological titre at week 3 (D22). Case B had a robust plasmablast response at the end of week 2 (D14) but the IgG plasmablast subset continued to rise,

dominated at week 3 (D18), but then subsided at the beginning of week 4 (D22),

which is compatible with high anti-spike glycoprotein and RBD IgG serological titres

at week 3. Case C produced a significant early plasmablast response at the end of

week 1 (D6), the IgG plasmablast subset dominated at the same time, and both the

plasmablast response and its IgG subsets subsided at the end of week 2 (D14).

Circulating plasmablasts were identified and used to generate human IgG monoclonal

antibodies (MAbs) from the three patients (Figure 2a). A total of 219 plasmablast-

derived IgG MAbs were produced, of which 67 (10 of 50 from case A, 48 of 131 from

case B, 9 of 38 from case C) were shown to bind spike glycoprotein or nucleocapsid

antigens of SARS-CoV-2 by one or more of the following: staining of spike

glycoprotein and RBD-expressing cells, ELISA, or immunofluorescence and specific

virus neutralisation. Averages of 13.7±6.8% (6.0-18.4%) and 13.0±7.3% (5.3-19.9%)

of plasmablast-derived IgG MAbs were reactive with virus spike glycoprotein and

nucleocapsid, respectively.

Genetic and phenotypic characteristics of anti-spike glycoprotein antibodies

Among 32 anti-spike MAbs, 10 bound to the RBD, 13 to non-RBD S1, and the other

9 to the S2 subunit of the SARS-CoV-2 spike glycoprotein (Figure 2a and Table 1).

Twenty-four of these MAbs bound to virus antigens as assessed by

immunofluorescence of SARS-CoV-2-infected Vero E6 cells (Supplemental Figure 1),

suggesting that the majority of anti-spike glycoprotein human antibodies recognise

complex conformational epitopes on the virus glycoprotein. Moderate binding was

observed for a subset (4 of 9) of anti-S2 MAbs with full-length spike glycoprotein

ectodomain in the indirect ELISA but bound strongly to the isolated S2 subunit

(Figure 2b).

Ten of 32 anti-SARS-CoV-2 spike glycoprotein MAbs cross-reacted with the

glycoproteins of other betacoronaviruses, including SARS, MERS or human common

cold coronavirus OC43 in ELISA (Table 1), suggesting the presence of conserved

epitopes on the spike glycoproteins of betacoronaviruses.

Each of 32 anti-spike glycoprotein MAbs was encoded by a unique set of heavy chain

VDJ and light chain VJ rearrangements in the variable domain (Supplemental Table

2). Fourteen of 32 SARS-CoV-2 spike-reactive MAb genes possessed low numbers of

somatic mutations resulting in 0 or 1 amino acid substitutions suggesting a *de novo* B

cell response to the SARS-CoV-2 virus in humans. Six of the 32 MAb genes

possessed ≥ 20 nucleotide mutations, and these cross-reacted on other beta-

coronaviruses, including OC43. Of the nine anti-S2 antibodies five cross-reacted on

OC43 virus and three of these also cross-reacted on MERS (Table 1). All five cross-

reactive anti-S2 antibodies had high rates of somatic mutation (25±5), indicating a

memory phenotype, and three of the five were neutralising to a moderate level (half

maximal effective concentration, EC_{50}, 36-133.33 nM, Table 1).

The CDR3 length varied among anti-spike glycoprotein antibodies (Supplemental

Table 2). No significant differences were found between anti-S2 and anti-S1 or anti-

RBD subsets. Among anti-S2 MAbs, a significantly longer heavy chain CDR3 length

was found in the cross-reactive group compared to the specific group (Cross-reactive

20±2 versus Specific 12±4, p= 0.02, two-tailed Mann-Whitney test; Figure 2c),

indicating that a long CDR3 may play a role in antigen binding, which is also found in

several broadly reactive human MAbs against human immunodeficiency virus and

influenza virus (9, 10).

The binding activities of 10 anti-RBD MAbs were further characterised in detail.

Using MDCK-SIAT1 cells transduced to express the RBD and flow cytometry,

binding activities of the anti-RBD MAbs were shown to vary with 50% binding

concentration from 0.10 to 1.83 µg/ml (Supplemental Figure 2). The MAbs with

strong anti-RBD binding have a relatively long heavy chain CDR3 length (50%

binding concentration <0.5 µg/ml versus >0.5 µg/ml, p=0.03, two-tailed Mann-

Whitney test; Supplemental Figure 3).

Neutralisation by anti-spike glycoprotein antibodies

The optimum method for detecting neutralisation of SARS-CoV-2 has not been

established. For example, different laboratories have obtained varying results for the

well-defined monoclonal antibody CR3022 (11-13). We have therefore obtained

results from neutralisation assays performed in independent laboratories: a systematic

survey of the thirty two anti-spike MAbs at the Francis Crick Institute (London), with

follow up for a selection of MAbs at the Sir William Dunn School (Oxford) and

Chang Gung Memorial Hospital (Taiwan).

The 32 anti-spike glycoprotein MAbs were systematically examined by plaque

reduction neutralisation (PRNT) assay for neutralisation of wild type SARS-CoV-2

virus (see methods; summarised in Table 1). A total of 14 neutralising antibodies

distributed between different regions of the spike glycoprotein were identified: 5 of 10

to RBD, 3 of 13 to S1 (non-RBD), 6 of 9 to S2. The EC_{50} concentrations, as a

measure of potency, ranged from 0.05 to ~133 nM (8 ng/ml - ~20 µg/ml).

Neutralisation of anti-RBD antibodies was corroborated by a microneutralisation test

that measured a reduction in fluorescent focus-forming units (see methods,

Supplemental Figure 4). MAbs were also tested by a PCR-based neutralisation assay

(see methods): inhibition of virus replication was measured by quantitative PCR in the

supernatant bathing the infected cells. This results corroborated that anti-RBD FD

11A, anti-RBD FI 3A, anti-RBD FD 5D, anti-RBD EY 6A and anti-S2 EW 9C, as

crude culture supernatants, reduced the virus signal from ~56- to ~10,085-fold

(Supplemental Figure 5).

ACE2 blockade by anti-RBD antibodies

Potent neutralising antibodies to the RBD of SARS-CoV-2 spike glycoprotein were

identified and we thus analyse the blockade of the ACE2-RBD interaction by anti-

RBD antibodies in two assays (Figure 3, Table 1), including an assay that we reported

previously (14, 15). In the first assay unlabelled MAbs in at least 10-fold excess were

mixed with biotin labelled ACE2-Fc and binding to the RBD, displayed on RBD-VLP

(16) bound to an ELISA plate (Figure 3a, Supplemental Figure 6). Anti-RBD

neutralising antibodies FI 3A and FI 1C, as well as the unlabelled ACE2-Fc, strongly

inhibited the binding of labelled ACE2-Fc to RBD. A partial inhibition was detected

by anti-RBD antibodies FD 11A, FD 5D and FI 4A. Another unlabelled nanobody

VHH72-Fc in excess inhibited the binding of labelled ACE2-Fc to RBD in this assay.

The structure of VHH72-Fc bound to RBD is known (17) and its footprint on the

RBD does not overlap that of ACE2, so inhibition is thought to occur by steric

hindrance.

In the second assay, we employed MDCK-SIAT1 cells overexpressing full-length

human ACE2 as a transmembrane protein. Unlabelled antibodies or ACE2-Fc were

mixed in excess with biotinylated RBD, and binding of RBD was detected with

Streptavidin-HRP in ELISA (Figure 3b). The results of this assay mostly mirrored

those of the first assay and confirmed that in this orientation anti-RBD neutralising

antibodies FD 11A and FD 5D competed in excess with soluble RBD for binding to

ACE2 (Figure 3b). In addition, anti-RBD neutralising antibody EY 6A competed with

RBD for ACE2 binding. The binding pattern of EY 6A is analogous to a previously

described antibody CR3022 (Table 1) (11). These two antibodies are known to bind to

the same region of RBD away from the ACE2 binding site, but they influence the

binding kinetics of RBD to ACE2, presumably through steric effects (15).

Division of anti-RBD antibodies into cross-inhibiting groups

The ten anti-RBD MAbs were then divided into cross-inhibiting groups as described

for human MAbs to Ebola (18) by assessing competition of unlabelled antibodies at

10-fold (or greater) excess over a biotin labelled target antibody by ELISA. Included

as controls were the VHH72-Fc (17) and H11-H4-Fc (14) nanobodies linked to the

hinge and Fc region of human IgG1, CR3022 and S309 human MAbs (11, 19)

reconstituted as an IgG1 antibody. These four control molecules have characterised

binding footprints on the RBD defined by crystal structures (12, 14, 17, 19), as does

EY 6A (15). Also included was the protease domain (residues 18-615) of ACE2 linked

to the Fc region of human IgG1 (ACE2-Fc dimer).

The ten antibodies formed four cross-inhibiting clusters (Table 2), represented by

antibodies EY 6A (cluster 1, which included CR3022), FI 3A (cluster 2, which

included H11-H4), FD 11A (cluster 3, which included S309) and FJ 10B (cluster 4).

The strongest inhibitors of ACE2-Fc binding were in clusters 2 and 3 (Tables 1 and 2).

Neutralising antibodies were detected in clusters 1, 2 and 3, with the strongest

antibodies FI 3A and FD 11A being in clusters 2 and 3 (Tables 1 and 2).

Neutralising antibodies to RBD and relationship to ACE2 blockade

Five neutralising anti-RBD MAbs partially or completely blocked the interaction

between RBD and ACE2 (Tables 1 and 2, Figure 3). The most potent neutralising

antibodies were ACE2 blockers (FI 3A in cluster 2, and FD 11A in cluster 3), and

bound independently of each other to the RBD (Figure 3c). MAb EY 6A has been

shown to alter the binding kinetics of the interaction without full inhibition (15) and it

had a moderate effect on ACE2 binding in the assay where ACE2 was expressed at the

cell surface. These three MAbs bound independently of each other indicating the

existence of at least three neutralisation-sensitive epitopes within the RBD (Figure

3c). All five neutralising MAbs to the RBD (EY 6A, FI 3A, FI 1C, FD 11A, FD 5A)

had V gene sequences close to germline (Supplemental Table 2), indicating the

development of *de novo* anti-RBD antibodies upon natural SARS-CoV-2 infection in

humans.

The ability of convalescent sera to block soluble RBD for binding to cell-expressed

ACE2 was assessed. Both convalescent sera from case A (D22) and case B (D26)

exhibited detectable ACE2-blocking activities (Figure 3d). This serological activity is

compatible with the isolation of potent ACE2-blocking antibodies from both patients

(Table 1). By contrast, the D18 serum from case C showed minimal ACE2-blocking

activity. Both cases A and B had prolonged fever and the development of pneumonia

during hospitalization (Supplemental Table 1), which suggests that the development

of ACE2-blocking antibody response in the convalescent stage is likely associated

with clinical severity after infection.

Neutralising antibodies to S2

Six of nine MAbs specific for S2 showed moderate neutralisation in the PRNT assay

(Table 1). The antibodies FB 1E, FJ 4E and EW 9C, are moderately neutralising (EC_{50}

36-133.33 nM), cross-react on the spike glycoprotein from the common cold

betacoronavirus OC43, and show sequence characteristics of memory cells with high

numbers of somatic mutations. This indicates that memory B cells, likely primed by

an endemic or epidemic betacoronavirus related to OC43, can give rise to antibodies

that neutralise SARS-CoV-2, albeit modestly. The other three neutralising antibodies

specific for S2, FD 10A, FG 7A and FM 1A were close to germline in sequence

(Supplemental Table 2) and did not cross-react strongly with other betacoronaviruses

(Table 1). FD 10A exhibits the most potent neutralising activity in the PRNT assay

and also completely inhibits SARS-CoV-2-induced cytopathic effect (see methods) at

8.33 nM.

Neutralising antibodies to non-RBD S1

Thirteen MAbs were defined that bound the S1 region and three, close to germline in

sequence, were neutralising. FJ 1C showed strong neutralisation (EC$_{50}$ 55.5 nM),

whilst FD 11E (EC$_{50}$ 70 nM) and FD 1E (EC$_{50}$ 110 nM) were moderately neutralising

(Table 1). We investigated the specificity of these antibodies using flow cytometry

with MDCK-SIAT1 cells expressing the N-terminal domain (NTD) of spike

glycoprotein linked to the transmembrane and cytoplasmic region of influenza

haemagglutinin (see methods). FJ 1C (neutralising) and FD 7C bound strongly to this

NTD construct defining them as NTD-specific (Table 1, Supplemental Figure 7).

Characterisation of anti-nucleocapsid antibodies

A set of 35 anti-SARS-CoV-2 nucleocapsid MAbs were derived from circulating

plasmablasts in the naturally infected subjects (Table 3, Supplemental Figure 8); 19 of

these strongly cross-react with at least one of the betacoronaviruses tested, 17 with

SARS CoV, 14 with OC43 virus, and 13 with MERS CoV in ELISA. This suggested

extensive cross-reactivity of the antibody response to epitopes in nucleocapsids of

betacoronaviruses following natural infection with SARS-CoV-2.

The 35 MAbs were evolved from 33 clonal groups defined by their heavy chain VDJ

and light chain VJ rearrangements (Supplemental Table 3). Those cross-reactive with

other betacoronaviruses carried considerably more amino acid substitutions than

SARS-CoV-1-specific MAbs in their variable domains (CR heavy 22±15 versus

specific heavy 9±11, p=0.008, two-tailed Mann-Whitney test; Figure 2d), supporting

the interpretation that approximately 50% of the anti-nucleocapsid antibody response

in these three patients was derived from memory B cells, presumably primed by

betacoronaviruses endemic to humans and related to OC43.

Discussion

We detected a robust and rapid plasmablast response encoding diverse anti-spike glycoprotein and anti-nucleocapsid antibody populations within 3 weeks of onset of illness in COVID-19 patients. The concomitant serologic response occurred as early as the first week after illness onset, with gradually increasing levels of SARS-CoV-2 spike glycoprotein- and RBD-binding IgG antibodies from the second to third week after symptom onset. The kinetics of plasmablast and virus-specific serologic responses was observed to vary between subjects. Consecutive samples from donor B showed a highly expanded plasmablast subset at the time of progression into pneumonia, followed by class-switching to an IgG-predominant plasmablast response until week three of illness. On the other hand, donor C presented with mild symptoms and produced an early plasmablast response with class-switching at the end of first week of symptom onset. Thevarajan et al demonstrated a similar kinetic of B cell response in a mild COVID-19 case, in whom the peripheral plasmablast response and the SARS-CoV-2-binding IgG antibodies peaked at day eight, soon after the disappearance of fever (5). An early class-switching phenotype of the humoral response was also noted within the first week of onset in paediatric patients who mainly experienced mild illness after SARS-CoV-2 infection, although the immunological basis for this phenotype is unclear (20). Both peak plasmablast frequency and serologic titre for spike glycoprotein were higher in donor B, who presented with severe symptoms, than in donor C. Similar observations on the serologic titre and clinical severity have been reported by others (7, 21).

In this study, a substantial subset of plasmablast-derived anti-S2 (five of nine, 56%) and anti-nucleocapsid antibodies (14 of 35, 40%) cross-reacted with human

betacoronavirus OC43. Cross-reactivity among betacoronaviruses has also been

reported in polyclonal sera (22, 23). Human coronavirus OC43, discovered in the

1960s, is one of the major betacoronaviruses that cause common colds in the

community (24, 25), and severe respiratory infections in elderly and

immunocompromised individuals (23, 26). Epidemiologic surveys reveal that OC43

infection can occur in early childhood, and that OC43 seropositivity reaches nearly

90% in adults (27, 28). The S2 component of SARS-CoV-2 spike glycoprotein shares

43~89% amino acid identity with SARS, MERS and OC43, and, similarly, the amino

acid sequence of SARS-CoV-2 nucleocapsid protein of is 34~90% homologous to

those of other betacoronaviruses (Supplemental Figures 9 and 10), suggesting the

presence of conserved epitopes on these antigens.

The presence of pre-existing immune memory to betacoronavirus that cross-react with

SARS-CoV-2 is supported by the accumulation of somatic mutations in the genes

encoding cross-reactive antibodies isolated from COVID-19 patients (Figures 2c and

2d, Supplemental Tables 2 and 3). This situation is reminiscent of re-exposure to

immunogenic epitopes shared by closely related viruses leading to induction of

broadly cross-reactive antibodies in patients infected with influenza, dengue or Zika

viruses (29-31).

The 32 MAbs that bound to the spike glycoprotein were systematically tested for

neutralisation (summarised in Table 1). Results established that neutralising epitopes

were present on the RBD, S1-NTD, S1-non NTD/RBD, and S2 regions of the spike

glycoprotein. The range of neutralisation EC_{50} titres against live SARS-CoV-2

reported in the literature for human and murine monoclonal antibodies span at least

three orders of magnitude, from ng/ml to µg/ml (32, 33). Our results reflect the range

of values in *in vitro* potency. The relationship between the EC_{50} value in neutralisation

assays and therapeutic potential is not established, although antibodies to the RBD

with EC_{50} in the nanogram range have shown therapeutic activity in small animal

models (34-36). The majority of the most strongly neutralising antibodies were close

to germline in sequence, showing that full affinity maturation is not required in order

to achieve SARS-CoV-2 virus neutralisation. We made a similar observation for

antibodies induced by an Ebola vaccine (18), and this has been observed in other

collections of monoclonal antibodies to SARS-CoV-2 (4, 33-40).

The RBDs of SARS and SARS-CoV-2 are known to contain neutralising epitopes (15,

32-42), and vaccines based on the RBD of SARS and SARS-CoV-2 induce strong

neutralising antibodies that are protective in animal models (43-45). There is a

tendency for the most potent neutralising antibodies to be those that block the binding

of the RBD to its receptor ACE2 (32-42, 46-48). Previous structural studies

demonstrated that potent antibodies with footprints overlapping that of ACE2 are

composed of two major subgroups, one with representative Vh3-53 gene usage and

short CDR3 binding to the RBD in an up conformation and the other binding RBDs in

both up and down conformations and even adjacent RBDs (49, 50). Potent MAb FI

3A shares similar characteristics with the first subgroup described above. Occasional

antibodies that do not occupy the ACE2 footprint or weakly block ACE2 binding, i.e.

S309, CR3022 and EY6A, can be almost as potent, perhaps through spike trimer

cross-linking or triggering a conformational change in the spike glycoprotein that

renders it non-functional (4, 15, 19, 40).

We characterise ten MAbs targeting the RBD that can be arranged into four cross-

inhibiting clusters (Table 2). Three of these MAb clusters (represented by EY 6A, FI

3A and FD 11A) demonstrate neutralisation of SARS-CoV-2 to some level, while

those in the fourth (represented by FM 7B) did not neutralise. This suggests that three

neutralising antibodies may be able to bind to the RBD simultaneously. The most

potent neutralising MAbs fall in two clusters and interfere strongly with ACE2

binding, which may be exploited therapeutically. For instance, antibodies FI 3A in

cluster 2 (EC_{50} 8.67 nM) and FD 11A in cluster 3 (EC_{50} 0.05 nM) could be combined

to limit possible selection of neutralisation-resistant variants. This principle has been

demonstrated in recent studies (36, 38, 51).

Neutralisation by human MAbs to the S1-NTD of SARS-CoV-2 has been described

(35, 52), but their mechanism of action is not known. The range of neutralising EC_{50}

titres (0.09-51.1 nM) was overlapping with those targeting the RBD, similar to our

NTD-specific antibody FJ 1C. Cocktails of antibodies that include a representative to

the NTD would further reduce the likelihood of selecting neutralisation-resistant

viruses. A second strong binder to the NTD, FD 7C, was not neutralising. Structural

comparisons of these two antibodies bound to spike glycoprotein may provide insight

into the neutralising action of FJ 1C.

We detected a subset of six MAbs to the S2 region of the spike glycoprotein that

neutralised moderately (EC_{50} 36-133 nM). Three of these were clearly derived from a

memory population; showing significant accumulation of somatic mutations in the

MAb encoding genes and cross-reactivity for the OC43 common cold virus spike

glycoprotein (Figure 2c). Further investigation is required to ascertain whether such

antibodies, that may be weak- or non-neutralising and cross-reactive with common

cold viruses, are beneficial or detrimental with respect to COVID-19 disease.

Our results have significance for serologic tests employing the N and S2 antigens of

SARS-CoV-2. Serologic surveys, some of which are based on these antigens, with

sera from donors infected with SARS-COV-2 during the spring and summer months

have shown very high specificity (as judged by comparing convalescent sera from

COVID-19   patients   versus   sera   collected   in   the   pre-COVID-19

period)(www.gov.uk/government/publications/covid-19-laboratory-evaluations-of-

serological-assays). However, if many individuals have memory B cells to S2 and N

antigens of circulating betacoronaviruses that cross-react with SARS-CoV-2,

concurrent winter infections with these viruses might erode the specificity of

serologic tests for SARS-CoV-2 based on these two antigens. Cross-reactivity of

antibodies targeting the S1 subunit is relatively lower, so using this antigen is more

likely to give a true indication of SARS-CoV-2 specificity.

In   summary,   COVID-19   patients   developed   strong   anti-SARS-CoV-2   spike

glycoprotein and nucleocapsid plasmablast and antibody responses. A panel of IgG

MAbs targeted a diverse spectrum of epitopes on the RBD, S1-NTD, non-NTD/RBD

S1 and S2 regions of the spike glycoprotein, of which 14 neutralised wild type virus

with EC_{50}s in the range 0.05 to ~133 nM. Neutralising activities of the majority of

anti-RBD MAbs were linked to ACE2-binding blockade, and non-competing pairs of

such MAbs, perhaps combined with a neutralising MAb to the NTD, offer potential

formulations for the development of prophylactic and therapeutic agents against

SARS-CoV-2. Antibody responses to nucleocapsid and the S2 component of spike

glycoprotein confirm marked cross-reactivity with a common cold virus.

Methods

Study design

This study was designed to isolate SARS-CoV-2 antigen-specific MAbs from peripheral plasmablasts of humans infected with SARS-CoV-2 and to characterise the antigenic specificity and phenotypic activities of the MAbs. Diagnosis of SARS-CoV-2 infection was based on positive real-time reverse transcriptase polymerase chain reaction results of respiratory samples. The study protocol and informed consent were approved by the ethics committee at the Chang Gung Medical Foundation and the Taoyuan General Hospital, Ministry of Health and Welfare. Each patient provided signed informed consent. The study and all associated methods were carried out in accordance with the approved protocol, the Declaration of Helsinki and Good Clinical Practice guidelines.

Staining and sorting of plasmablasts

Freshly separated peripheral blood mononuclear cells (PBMCs) or thawed PBMCs were stained with fluorescent-labelled antibodies to cell surface markers purchased from BD Biosciences, USA; Pacific blue anti-CD3 (clone UCHT1, Cat. No. 558117, BD), Fluorescein isothiocyanate anti-CD19 (clone HIB19, Cat. No. 555412, BD), Phycoerythrin-Cy7 anti-CD27 (clone M-T271, Cat. No. 560609, BD), Allophycocyanin-H7 anti-CD20 (clone L27, Cat. No. 641396, BD), Phycoerythrin-Cy5 anti-CD38 (clone HIT2, Cat. No. 555461, BD) and Phycoerythrin anti-human IgG (clone G18-145, Cat. No. 555787, BD). The $CD3^{neg}CD19^{pos}CD20^{neg}CD27^{hi}CD38^{hi}IgG^{pos}$ plasmablasts were gated and isolated in chamber as single cells as previously described (53).

Production of human IgG 1 monoclonal antibodies

Sorted single cells were used to produce human IgG MAbs as previously described (53). Briefly, the variable region genes from each single cell were amplified in a reverse transcriptase polymerase chain reaction (RT-PCR: QIAGEN, Germany) using a cocktail of sense primers specific for the leader region and antisense primers to the Cγ constant region for heavy chain and Cκ and Cλ for light chain. The RT-PCR products were amplified in separate polymerase chain reactions for the individual heavy and light chain gene families using nested primers to incorporate restriction sites at the ends of the variable gene as previously described (53). These variable genes were then cloned into expression vectors for the heavy and light chains. Plasmids were transfected into the HEK293T cell line for expression of recombinant full-length human IgG MAbs in serum-free transfection medium. A selected panel of MAbs were further expanded and purified.

To determine the individual gene segments employed by VDJ and VJ rearrangements and the number of nucleotide mutations and amino acid replacements, the variable domain sequences were aligned with germline gene segments using the international ImMunoGeneTics (IMGT) alignment tool (http://www.imgt.org/IMGT_vquest/input).

Enzyme-linked immunosorbent assay (ELISA)

ELISA plates (Corning® 96-well Clear Polystyrene High Bind Stripwell™ Microplate, USA) were coated with 8 µg/ml SARS-CoV-2 antigens (spike glycoprotein extracellular or receptor-binding domains, or nucleocapsid: Sino Biological, China) or SARS antigen (spike glycoprotein S1 subunit: Sino Biological, China) or Middle East Respiratory Syndrome coronavirus (MERS) antigen (spike glycoprotein extracellular domain: Sino Biological, China) or human coronavirus

OC43 antigen (spike glycoprotein extracellular domain: Sino Biological, China) at

4°C overnight. Plates were washed with phosphate-buffered saline containing 0.05%

Tween-20 and blocked with 3% bovine serum albumin (BSA) at room temperature for

1 hour on a shaker. Serial dilutions of MAb-containing cell culture supernatant or

purified MAb were added and plates were incubated at 37°C for 1 hour. Plates were

washed and incubated with horseradish peroxidase-conjugated rabbit anti-human IgG

secondary antibody (Rockland Immunochemicals, USA). Plates were washed and

developed with TMB substrate reagent (BD Biosciences, USA). Reactions were

stopped with 0.5M hydrochloric acid and absorbances was measured at 450nm on a

microplate reader. Non-transfected cell culture supernatant, anti-influenza H3 human

IgG MAb BS 1A (in house), anti-SARS spike glycoprotein MAb CR3022 and

convalescent serum were used as controls for each experiment. Reaction yielding an

absorbance value above three times the mean absorbance of the negative control BS

1A were considered positive.

Flow-cytometry based binding assay

MDCK-Spike cells were produced by stably transfecting parental MDCK-SIAT1 cells

with cDNA expressing full-length SARS-CoV-2 spike glycoprotein. MDCK-RBD

cells were produced by, stably transducing MDCK-SIAT1 cells with a Lentiviral

vector encoding a cDNA expressing RBD amino acids 340-538 (NITN.GPKK

underlined) fused via a short linker to the transmembrane domain of haemagglutinin

H7 (A/Hong Kong/125/2017) (EPI977395) at the C-terminus for surface expression

(sequence:

MNTQILVFALIAIIPTNA/DKIGSGSNITNLCPFGEVFNATRFASVYAWNRKRISN

CVADYSVLYNSASFSTFKCYGVSPTKLNDLCFTNVYADSFVIRGDEVRQIAPG

QTGKIADYNYKLPDDFTGCVIAWNSNNLDSKVGGNYNYLYRLFRKSNLKPFE

RDISTEIYQAGSTPCNGVEGFNCYFPLQSYGFQPTNGVGYQPYRVVVLSFELL

HAPATVCGPKKTGSGGSGKLSSGYKDVILWFSFGASCFILLAIVMGLVFICVKN

GNMRCTICI*). MDCK-NTD cells were produced by stably transfecting parental

MDCK-SIAT1 cells with cDNA expressing SARS-CoV-2 NTD.

Both MDCK-Spike and MDCK-RBD cells were then FACS sorted for highly

expressing cells using the CR3022 antibody. MDCK-Spike or MDCK-RBD cells

were prepared and resuspended. Cells were probed with purified MAbs in 3% BSA.

Bound primary antibodies were detected with FITC-conjugated anti-IgG secondary.

The binding activities were analyzed by BD FACSCanto™ II flow cytometer (BD

Biosciences, USA).

Immunofluorescence assay

Under biosafety level 3 (BSL-3) conditions, Vero E6 cells were infected with 100

$TCID_{50}$ (median tissue culture infectious dose) SARS-CoV-2 (hCoV-

19/Taiwan/CGMH-CGU-01/2020, EPI_ISL_411915). Infected cells were placed on

coverslips and, and fixed with acetone at room temperature for 10 minutes. After

blocking with 1% BSA at room temperature for 1 hour and washing, fixed cells were

incubated with MAb-containing cell culture supernatant. The anti-influenza human

monoclonal antibody BS 1A, anti-SARS spike glycoprotein MAb CR3022 and

convalescent serum were used as antibody controls for each experiment. Following

incubation and wash, cells were stained with FITC-conjugated anti-human IgG

secondary antibody and Evans blue dye as counterstain. Antibody-bound infected

cells demonstrated an apple-green fluorescence against a background of red

fluorescing material stained by the Evans Blue counterstain. Images were acquired

with original magnification 40x, scale bar 20 μm.

Plaque reduction neutralisation assay (Francis Crick Institute)

Confluent monolayers of Vero E6 cells in 96-well plates were incubated with ~14 plaque forming units (PFU) of SARS CoV-2 (hCoV-19/England/02/2020, EPI_ISL_407073) and antibodies in a 2-fold dilution series (triplicates) for 3 hours at room temperature. Inoculum was then removed, and cells were overlaid with plaque assay overlay. Cells were incubated at 37°C, 5% CO_2 for 24 hours prior to fixation with 4% paraformaldehyde at 4°C for 30 minutes. Fixed cells were then permeabilised with 0.2% Triton-X-100 and stained with a horseradish peroxidase conjugated-antibody against virus protein for 1 hour at room temperature. TMB substrate was then added to visualise virus plaques as described previously for influenza virus (54). Convalescent serum from COVID-19 patients was used as a control.

Fluorescent focus-forming units microneutralisation assay (FMNT) (Oxford)

In brief, this rapid, high-throughput assay determines the concentration of antibody that produces a 50% reduction in infectious focus-forming units of authentic SARS-CoV-2 in Vero cells, as follows. Triplicate serial dilutions of antibody are pre-incubated with a fixed dose of SARS-CoV-2 (Australia/VIC01/2020, GenBank MT007544) (55) in triplicate before incubation with Vero cells. A carboxymethyl cellulose-containing overlay is used to prevent satellite focus formation. Twenty hours post-infection, the monolayers are fixed with paraformaldehyde and stained for N antigen using MAb EY 2A. After development with a peroxidase-conjugated antibody and substrate, foci are enumerated by enzyme-linked immune absorbent spot reader. Data are analysed using four-parameter logistic regression (Hill equation) in GraphPad Prism 8.3.

Quantitative PCR-based neutralisation assay

Neutralisation activity of MAb-containing supernatant was measured using SARS-CoV-2 (hCoV-19/Taiwan/CGMH-CGU-01/2020, EPI_ISL_411915) infected Vero E6 cells. Briefly, Vero E6 cells were pre-seeded in a 96 well plate at a concentration of 10^4 cells per well. The following day, MAb-containing supernatants were mixed with equal volumes of 100 TCID$_{50}$ virus preparation and incubated at 37°C for 1 hour, then mixtures were added to seeded Vero E6 cells and incubated at 37°C for 5 days. Cell, virus and virus back-titration controls were setup for each experiment. At day 5 the culture supernatant was harvested from each well, and virus RNA was extracted and quantified by real-time RT-PCR targeting the E gene of SARS-CoV-2 as previously described. The cycle threshold values of real-time RT-PCR were used as indicators of the copy number of SARS-CoV-2 RNA in samples with lower cycle threshold values corresponding to higher virus copy numbers.

CPE-based neutralisation assay

Vero E6 cells in Dulbecco's Modified Eagle's Medium containing 10% FBS were added into 96-well plates and incubated at 37°C with 5% CO_2 overnight to reach confluence. After washing with virus growth medium (VGM: Dulbecco's Modified Eagle's Medium containing 2% FBS), two-fold serially diluted MAbs in VGM starting at 100 µg/ml were added to each duplicated well. The plates were immediately transferred to a BSL-3 laboratory and 100 TCID$_{50}$ SARS-CoV-2 (hCoV-19/Taiwan/4/2020, EPI_ISL_411927) in VGM was added. The plates were further incubated at 37°C with 5% CO_2 for three days and the cytopathic morphology of the cells was recorded using an ImageXpress Nano Automated Cellular Imaging System.

Competitive binding assays

Competitive binding assays were performed as described previously (18) with slight

modifications for epitope mapping of the anti-RBD MAbs. Briefly, 0.5 μg/ml of

RBD-VLP were coated on NUNC plates (50 μl per well) overnight at 4ºC, washed

and blocked with 300 μl of 5% (w/v) dried skimmed milk in PBS for 1 hour at room

temperature prior to the assays. Antibody was biotinylated using EZ-Link Sulfo-NHS-

LC-biotin (21237; Life Technologies) and then mixed with competing MAb (in at

least 10-fold excess) and transferred to the blocked NUNC plates for 1 hour. A second

layer Streptavidin-HRP (S911, Life Technologies) diluted 1:1,600 in PBS/0.1% BSA

(37525; Thermo Fisher Scientific) was then added and incubated for another 1 hour.

Plates were then washed, and signal was developed by adding POD substrate

(11484281001, Roche) for 5 minutes before stopping the reaction with 1 M H_2SO_4.

Absorbance (OD_{450}) was measured using a Clariostar plate reader (BMG, Labtech).

Mean and 95% confidence interval of 4 replicate measurements were calculated.

Competition was measured as: (X-minimum binding/(maximum binding-minimum

binding), where X is the binding of the biotinylated MAb in the presence of

competing MAb. Minimum binding is the self-blocking of the biotinylated MAb or

background binding. Maximum binding is binding of biotinylated MAb in the

presence of non-competing MAb (anti-influenza N1 neuraminidase MAb).

ACE2 blocking assays

Two assays were used to determine the blocking of binding of ACE2 to RBD by

MAbs. RBD was anchored on the plate in the first assay whereas ACE2 was anchored

for the second assay.

In the first ACE2 blocking assay, RBD-VLP (Spycatcher-mi3 VLP-particles

conjugated with Spytagged-RBD recombinant protein) (16) was coated on ELISA

plates as described for the competitive binding assay. Recombinant ACE2-Fc (18-615)

protein expressed in Expi293F (Life Technologies) cells was chemically biotinylated

using EZ-link Sulfo-NHS-Biotin (A39256; Life Technologies) and buffer exchanged

to PBS using a Zebaspin desalting column (Thermo Fischer). MAbs were titrated in

duplicate or triplicate as half-log serial dilution, 8-point series starting at 1 µM in 30

µl volume with PBS/0.1% BSA buffer. 30 µl of biotinylated ACE2-Fc at approx. 0.2

nM (40 ng/ml) was added to the antibodies. 50 µl of the mixture was transferred to the

PBS-washed RBD-VLP coated plates and incubated for 1 hour at room temperature.

Secondary Streptavidin-HRP antibody (S911, Life Technologies) diluted to 1:1600

was then added to the PBS-washed plates and incubated for 1 h at room temperature.

Plates were then washed four times with PBS and signal was developed by adding

POD substrate (11484281001, Roche) for 5 minutes before stopping with 1 M H_2SO_4.

OD450 was measured using a Clariostar plate reader (BMG, Labtech). The control

antibody (a non-blocking anti-influenza N1 MAb) or ACE2-Fc without antibody used

to obtain the maximum signal and wells with PBS/BSA buffer only were used to

determine the minimum signal. Graphs were plotted as % binding of biotinylated

ACE2 to RBD. Binding % = {(X - Min)/(Max - Min)}*100 where X = measurement

of the antibody, Min = buffer only, Max = biotinylated ACE2-Fc alone. 50%

inhibitory concentrations of the antibodies against ACE2 was determined using non-

linear regression curve fit using GraphPad Prism 8.

The second ACE2 blocking assay was performed as described previously (14, 15).

Briefly, MDCK-SIAT1 cells were stably transfected to overexpress codon-optimised

human ACE2 cDNA (NM_021804.1) using lentiviral vector and FACS sorted

(MDCK-ACE2). Cells (3 x 10^4 per well) were seeded on a flat-bottomed 96-well plate

the day before the assay. RBD-6H (340-538; NITN.GPKK) was chemically

biotinylated using EZ-link Sulfo-NHS-Biotin (A39256; Life Technologies). Serial

half-log dilutions (starting at 1 µM) of antibodies and controls were performed in a U-

bottomed 96 well plate in 30 µl volume. 30 µl of biotinylated RBD (25 nM) were

mixed and 50 µl of the mixture was then transferred to the MDCK-ACE2 cells. After

1 hour a second layer Streptavidin-HRP antibody (S911, Life Technologies) diluted

1:1,600 in PBS/0.1% BSA (37525; Thermo Fisher Scientific) was added and

incubated for another 1 hour. Plates were then washed four times with PBS and signal

was developed by adding POD substrate (11484281001, Roche) before stopping with

1 M H_2SO_4 after 5 minutes. OD450 was measured using a Clariostar plate reader

(BMG, Labtech). The control antibody (a non-blocking anti-influenza N1 antibody)

was used to obtain maximum signal and PBS only wells were used to determine

background. Graphs were plotted as % binding of biotinylated RBD to ACE2. The

50% inhibitory concentration of the blocking antibody was determined as described

above.

Data availability

Antibodies are available (by contacting Kuan-Ying A. Huang

[arthur1726@cgmh.org.tw] from the Chang Gung Memorial Hospital and Chang

Gung University and Alain R. Townsend [alain.townsend@imm.ox.ac.uk] from

Oxford University) for research purposes only under an MTA, which allows the use of

the antibody for non-commercial purposes but not their disclosure to third parties. The

data that support the findings of this study are available from the corresponding

authors on request.

Author contributions

K.-Y.A.H. conceived and designed the study of MAb isolation and characterisation. K.-Y.A.H. produced and characterised plasmablast-derived MAbs. A.R.T. conceived and designed the study of MAb characterisation at Oxford. P.R, T.K.T, L.S generated cell lines, expressed proteins and antibodies, and performed experiments. S.H., R.H. R.S.D and J.W.M. designed and performed neutralisation assay at Crick Institute. A. H., J. G-J., X. L., M. K. and W.J. designed and performed neutralisation assay at Oxford. T-H.C., C-G.H., C-P.C., S-R.S, Y-C.L., C-Y.C., S-H.C., Y-C.H., T-Y.L., J-T.J. and C.M helped prepare materials, perform experiments and analyse data. All authors read and approved the manuscript.

Acknowledgements

We acknowledge the BD FACSAria™ cell sorter service provided by the Core Instrument Center of Chang Gung University. Plasmablast sorting, production and characterisation of human MAbs were supported by the Chang Gung Memorial Hospital (BMRPE22). We would like to acknowledge Paul Sopp and Craig Waugh in the flow cytometry facility at the MRC WIMM for providing cell sorting services. The facility is supported by the MRC HIU, MRC MHU (CC_UU_12009); NIHR Oxford BRC; Kay Kendall Leukaemia Fund (KKL1057), John Fell Fund (131/030 and 101/517), the EPA fund (CF182 and CF170) and by the MRC WIMM Strategic Alliance awards G0902418 and MC_UU_12025. Production of antibodies was funded by the Fast Grant Application given to A.R.T., P.R., and T.K.T. P.R., L.S. and A.R.T. are funded by the Chinese Academy of Medical Sciences (CAMS) Innovation Fund for Medical Science (CIFMS), China (grant no. 2018-I2M-2-002). T.K.T. is funded by

the EPA Cephalosporin Fund and The Townsend–Jeantet Charitable Trust (charity no.

1011770). The work done at the Crick Worldwide Influenza Centre was supported by

the Francis Crick Institute receiving core funding from Cancer Research UK

(FC001030), the Medical Research Council (FC001030) and the Wellcome Trust

(FC001030). The Oxford work was funded in part through the generous support of

philanthropic donors to the University of Oxford's COVID-19 Research Response

Fund.

References

1. Bloch, E. M. et al. Deployment of convalescent plasma for the prevention and

treatment of COVID-19. *J Clin Invest* 130, 2757-2765 (2020).

2. Li, L. et al. Effect of Convalescent Plasma Therapy on Time to Clinical

Improvement in Patients With Severe and Life-threatening COVID-19: A

Randomized Clinical Trial. *JAMA* e2010044 (2020).

3. Cheng, Y. et al. Use of convalescent plasma therapy in SARS patients in Hong Kong.

Eur J Clin Microbiol Infect Dis 24, 44-46 (2005).

4. Walls, A. C. et al. Structure, Function, and Antigenicity of the SARS-CoV-2 Spike

Glycoprotein. *Cell* 181, 281-292.e6 (2020).

5. Thevarajan, I. et al. Breadth of concomitant immune responses prior to patient

recovery: a case report of non-severe COVID-19. *Nat Med* 26, 453-455 (2020).

6. Nie, Y. et al. Neutralizing antibodies in patients with severe acute respiratory

syndrome-associated coronavirus infection. *J Infect Dis* 190, 1119-1126 (2004).

7. Long, Q. X. et al. Antibody responses to SARS-CoV-2 in patients with COVID-19.

Nat Med 26, 845-848 (2020).

8. Lou, B. et al. Serology characteristics of SARS-CoV-2 infection since exposure and

post symptom onset. *Eur Respir J* 2000763 (2020).

9. Ekiert, D. C. et al. Cross-neutralization of influenza A viruses mediated by a single

antibody loop. *Nature* 489, 526-532 (2012).

10. West, A. P. Jr et al. Structural insights on the role of antibodies in HIV-1 vaccine and

therapy. *Cell* 156, 633-648 (2014).

11. ter Meulen, J. et al. Human monoclonal antibody combination against SARS

coronavirus: synergy and coverage of escape mutants. *PLoS Med* 3, 237 (2006).

12. Huo, J. et al. Neutralization of SARS-CoV-2 by Destruction of the Prefusion Spike.

Cell Host Microbe S1931-3128(20)30351-6 (2020).

13. Yuan M, et al. A highly conserved cryptic epitope in the receptor binding domains of

SARS-CoV-2 and SARS-CoV. *Science* 368, 630-633 (2020).

14. Huo, J. et al. Neutralizing nanobodies bind SARS-CoV-2 spike RBD and block

interaction with ACE2. *Nat Struct Mol Biol* 10.1038/s41594-020-0469-6 (2020).

15. Zhou, D. et al. Structural basis for the neutralization of SARS-CoV-2 by an antibody

from a convalescent patient. *Nat Struct Mol Biol* 10.1038/s41594-020-0480-y

(2020).

16. Bruun, T. U. J., Andersson, A. C., Draper, S. J. & Howarth, M. Engineering a

Rugged Nanoscaffold To Enhance Plug-and-Display Vaccination. *ACS Nano* 12,

8855-8866 (2018).

17. Wrapp, D. et al. Structural Basis for Potent Neutralization of Betacoronaviruses by

Single-Domain Camelid Antibodies. *Cell* 181, 1004-1015.e15 (2020).

18. Rijal, P. et al. Therapeutic Monoclonal Antibodies for Ebola Virus Infection Derived

from Vaccinated Humans. *Cell Rep* 27, 172-186.e7 (2019).

19. Pinto, D. et al. Cross-neutralization of SARS-CoV-2 by a human monoclonal SARS-

CoV antibody. *Nature* 583, 290-295 (2020).

20. Zhang, Y. et al. Protective humoral immunity in SARS-CoV-2 infected pediatric

patients. *Cell Mol Immunol* 17, 768-770 (2020).

21. Zhao, J. et al. Antibody responses to SARS-CoV-2 in patients of novel coronavirus

disease 2019. *Clin Infect Dis* ciaa344 (2020).

22. Hicks, J. et al. Serologic cross-reactivity of SARS-CoV-2 with endemic and seasonal

Betacoronaviruses. Preprint. *medRxiv* 2020.06.22.20137695 (2020).

23. Patrick, D. M. et al. An Outbreak of Human Coronavirus OC43 Infection and

Serological Cross-reactivity with SARS Coronavirus. *Can J Infect Dis Med*

Microbiol 17, 330-336 (2006).

24. McIntosh, K., Dees, J. H., Becker, W. B., Kapikian, A. Z. & Chanock, R. M.

Recovery in tracheal organ cultures of novel viruses from patients with respiratory

disease. *Proc Natl Acad Sci U S A* 57, 933-940 (1967).

25. Nickbakhsh, S. et al. Epidemiology of Seasonal Coronaviruses: Establishing the

Context for the Emergence of Coronavirus Disease 2019. *J Infect Dis* 222, 17-25

(2020).

26. Morfopoulou, S. et al. Human Coronavirus OC43 Associated with Fatal

Encephalitis. *N Engl J Med* 375, 497-498 (2016).

27. Severance, E. G. et al. Development of a nucleocapsid-based human coronavirus

immunoassay and estimates of individuals exposed to coronavirus in a U.S.

metropolitan population. *Clin Vaccine Immunol* 15, 1805-1810 (2008).

28. Dijkman, R. et al. The dominance of human coronavirus OC43 and NL63 infections

in infants. *J Clin Virol* 53, 135-139 (2012).

29. Beltramello, M. et al. The human immune response to Dengue virus is dominated by

highly cross-reactive antibodies endowed with neutralizing and enhancing activity.

Cell Host Microbe 8, 271-283 (2010).

30. Rogers, T. F. et al. Zika virus activates de novo and cross-reactive memory B cell

responses in dengue-experienced donors. *Sci Immunol* 2, eaan6809 (2017).

31. Wrammert, J. et al. Broadly cross-reactive antibodies dominate the human B cell

response against 2009 pandemic H1N1 influenza virus infection. *J Exp Med* 208,

181-193 (2011).

32. He, Y., Lu, H., Siddiqui, P., Zhou, Y. & Jiang, S. Receptor-binding domain of severe

acute respiratory syndrome coronavirus spike protein contains multiple

conformation-dependent epitopes that induce highly potent neutralizing antibodies. *J

Immunol* 174, 4908-4915 (2005).

33. Robbiani, D. F. et al. Convergent antibody responses to SARS-CoV-2 in

convalescent individuals. *Nature* 10.1038/s41586-020-2456-9 (2020).

34. Cao, Y. et al. Potent Neutralizing Antibodies against SARS-CoV-2 Identified by

High-Throughput Single-Cell Sequencing of Convalescent Patients' B Cells. *Cell*

182, 73-84.e16 (2020).

35. Liu, L. et al. Potent neutralizing antibodies against multiple epitopes on SARS-CoV-

2 spike. *Nature* 10.1038/s41586-020-2571-7 (2020).

36. Wu, Y. et al. A noncompeting pair of human neutralizing antibodies block COVID-

19 virus binding to its receptor ACE2. *Science* 368, 1274-1278 (2020).

37. Brouwer, P. J. M. et al. Potent neutralizing antibodies from COVID-19 patients

define multiple targets of vulnerability. *Science* 369, 643-650 (2020).

38. Hansen, J. et al. Studies in humanized mice and convalescent humans yield a SARS-

CoV-2 antibody cocktail. *Science* eabd0827 (2020).

39. Rogers, T. F. et al. Isolation of potent SARS-CoV-2 neutralizing antibodies and

protection from disease in a small animal model. *Science* eabc7520 (2020).

40. Ju, B. et al. Human neutralizing antibodies elicited by SARS-CoV-2 infection.

Nature 584, 115-119 (2020).

41. Wang, C. et al. A human monoclonal antibody blocking SARS-CoV-2 infection. *Nat

Commun* 11, 2251 (2020).

42. Chen, X. et al. Human monoclonal antibodies block the binding of SARS-CoV-2

spike protein to angiotensin converting enzyme 2 receptor. *Cell Mol Immunol* 17,

647-649 (2020).

43. Zhu, X., Liu, Q., Du, L., Lu, L. & Jiang, S. Receptor-binding domain as a target for

developing SARS vaccines. *J Thorac Dis* 5 Suppl 2, S142-S148 (2013).

44. Quinlan, B. D. et al. The SARS-CoV-2 receptor-binding domain elicits a potent

neutralizing response without antibody-dependent enhancement. *bioRxiv*

2020.04.10.036418 (2020).

45. Yang, J. et al. A vaccine targeting the RBD of the S protein of SARS-CoV-2 induces

protective immunity. *Nature* 10.1038/s41586-020-2599-8 (2020).

46. Shang, J. et al. Structural basis of receptor recognition by SARS-CoV-2. *Nature* 581,

221-224 (2020).

47. Lan, J. et al. Structure of the SARS-CoV-2 spike receptor-binding domain bound to

the ACE2 receptor. *Nature* 581, 215-220 (2020).

48. Wang, Q. et al. Structural and Functional Basis of SARS-CoV-2 Entry by Using

Human ACE2. *Cell* 181, 894-904.e9 (2020).

49. Barnes, C. O. et al. SARS-CoV-2 neutralizing antibody structures inform therapeutic

strategies. *Nature* 2020 Oct 12. doi: 10.1038/s41586-020-2852-1 (2020).

50. Greaney, A. J., et al. Complete mapping of mutations to the SARS-CoV-2 spike

receptor-binding domain that escape antibody recognition. *bioRxiv*

2020.09.10.292078 (2020).

51. Baum, A. et al. Antibody cocktail to SARS-CoV-2 spike protein prevents rapid

mutational escape seen with individual antibodies. *Science* eabd0831 (2020).

52. Chi, X. et al. A neutralizing human antibody binds to the N-terminal domain of the

Spike protein of SARS-CoV-2. *Science* 369, 650-655 (2020).

53. Huang, K. A. et al. Structure-function analysis of neutralizing antibodies to H7N9

influenza from naturally infected humans. *Nat Microbiol* 4, 306-315 (2019).

54. Lin, Y. et al. Optimisation of a micro-neutralisation assay and its application in

antigenic characterisation of influenza viruses. *Influenza Other Respir Viruses* 9,

331-340 (2015).

55. Caly, L. et al. Isolation and rapid sharing of the 2019 novel coronavirus (SARS-CoV-

2) from the first patient diagnosed with COVID-19 in Australia. *Med J Aust* 212,

459-462 (2020).

Case	Onset date	Plasmablasts of CD3^{neg}CD19^{pos} B cells (%)	IgG Plasmablasts	
			of B cells(%)	of Plasmablasts (%)
A	D14	1.99	0.34	16.95
B	D14	37.48	1.67	4.47
	D18	1.13	0.64	56.93
	D22	0.20	0.05	23.33
C	D2	2.56	0.11	4.38
	D6	2.09	0.97	46.39
	D14	0.83	0.09	11.11

Figure 1. The IgG serology and plasmablast response to acute SARS-CoV-2 infection among enrolled patients. (a) The binding activity of post-infection sera IgG with SARS-CoV-2 spike glycoprotein in an ELISA and SARS-CoV-2 RBD assessed by flow cytometry on transfected cells, among enrolled patients. Each experiment was repeated twice. Values are presented as mean ± standard error of the

mean. Two sera from healthy adults (one collected at day 28 post 2018-19 influenza

vaccination and one collected from an influenza-infected patient 9 days after

symptom onset) in 2018 were included as controls. Linear regression was used to

determine the 50% end-point dilution (ED$_{50}$). **(b)** The gating strategy used for

peripheral total B cells, plasmablasts and IgG plasmablasts in flow cytometry. The

frequency of circulating plasmablasts (percentage of total B cells) among enrolled

cases was measured by flow cytometry. Onset date (D = Day).

Figure 2. Plasmablast-derived IgG monoclonal antibodies from three COVID-19

patients. (a) A total of 219 IgG monoclonal antibodies were produced from COVID-

19 patients (50 from case A, 131 from case B, 38 from case C). An average of

13.7±6.8% and 13.0±7.3% of antibodies were reactive with spike glycoprotein (S)

and nucleocapsid (N) antigens of SARS-CoV-2, respectively. The data are presented

as specificity, number of antibodies, and the percentage of total antibodies isolated

from each patient. **(b)** The binding activity of anti-SARS-CoV-2 MAbs with spike

glycoprotein, RBD and the S2 subunit in ELISA. Anti-influenza H3 MAb BS-1A and

anti-SARS RBD CR3022 were included as controls. Each experiment was repeated

twice. The OD_{450} values are presented as mean ± standard error of the mean. Panels

(c) and **(d)** show numbers of variable domain mutations in MAb genes and variation

in MAb CDR3 lengths among anti-S2 and anti-N MAbs, respectively. Antibodies that

strongly cross-react with at least one betacoronavirus (SARS or MERS or OC43)

were defined as cross-reactive MAbs. CDR3 length and mutation numbers are

presented as mean ± standard error of the mean (anti-S2, specific, n=4 versus cross-

reactive, n=5; anti-N, specific, n=16 versus cross-reactive, n=19). The two-tailed

Mann-Whitney test was performed to compare the mutations between two groups. * p

< 0.05, ** p < 0.01 ; D, =Day ; ns, non-significant ; CR, cross-reactive.

Figure 3. ACE2-blocking activities with anti-RBD antibodies and convalescent

sera. The analyses were performed with **(a)** RBD anchored and **(b)** ACE2 anchored

on plates (see methods). Anti-SARS-CoV-2 RBD nanobody VHH72 linked to the

hinge and Fc region of human IgG1 and ACE2-Fc were included as controls.

Experiments were performed in duplicate and repeated twice. **(c)** Mapping of

neutralising anti-RBD antibodies on the SARS-CoV-2 RBD structure (PDB 6ZCZ)

based on competitive binding and ACE2-blocking analyses. The RBD was colored in

green. The epitopes recognized by EY 6A, CR3022 and VHH72 (cluster 1 MAb) (11,

15, 17) were colored in magenta. The epitopes recognized by ACE2 and H11-H4

(cluster 2 MAb) (14) were overlapping and colored in blue and light blue. The

epitopes recognized by S309 (cluster 3 MAb) (19) were colored in yellow. **(d)**

Convalescent sera were analysed in the ACE2-blocking (ACE2 anchored) assay.

Experiments were performed in triplicate. Anti-RBD antibody FD 11A and anti-

influenza H3 antibody BS 1A were included as controls. Data are presented as mean $\pm$

standard error of the mean. IC_{50}, 50% inhibitory concentration.

Table 1. The antigenic specificity, cross-reactivity and function of 32 anti-SARS-CoV-2 spike antibodies derived from COVID-19 patients.

RBD-specific

Antibody	Case	Domain	SARS-CoV-2[a]				Cross-reactivity[a]			PRNT[b]	ACE2-block[c]	
			IFA	Spike	RBD	S2	SARS	MERS	OC43	EC_{50} (nM)	RBD anchored	ACE2 anchored
FD 11A	B	RBD	pos	1.64	1.17	0.13	0.10	0.10	0.12	0.05	+	++
FI 3A	B	RBD	pos	1.60	1.12	0.12	0.08	0.08	0.09	8.67	++++	++++
FI 1C	B	RBD	pos	1.88	1.22	0.16	0.15	0.15	0.18	16.67	++	++
FD 5D	B	RBD	pos	1.83	1.19	0.13	0.09	0.10	0.09	133.33	+	+++
EY 6A	A	RBD	pos	1.75	1.18	0.13	1.82	0.12	0.11	133.33	-ve	++
FI 4A	B	RBD	pos	1.68	1.12	0.12	0.08	0.09	0.10	-ve	+	-ve
EZ 7A	B	RBD	pos	1.66	1.12	0.33	2.05	1.00	1.05	-ve	-ve	-ve
FJ 10B	B	RBD	-ve	1.48	1.22	0.13	1.26	0.10	0.14	-ve	-ve	-ve
FM 7B	C	RBD	pos	1.80	1.27	0.12	2.25	0.09	0.10	-ve	-ve	-ve
FN 12A	C	RBD	pos	2.24	1.33	0.30	0.36	0.29	0.38	-ve	-ve	-ve

S1-non-RBD

Antibody	Case	Domain	SARS-CoV-2[a]				Cross-reactivity[a]			PRNT[b]
			IFA	Spike	RBD	S2	SARS	MERS	OC43	EC_{50} (nM)
FJ 1C	B	NTD	pos	1.61	0.27	0.15	0.23	0.15	0.16	55.50
FD 11E	B	non-NTD S1	pos	1.45	0.11	0.12	0.11	0.12	0.14	70.00
FD 1E	B	non-NTD S1	pos	1.45	0.11	0.13	0.12	0.12	0.14	110.00
EW 8B	B	non-NTD S1	-ve	1.61	0.11	0.13	0.11	0.11	0.12	-ve
FD 11D	B	NTD	pos	1.42	0.18	0.22	0.42	0.25	0.29	-ve
FD 11C	B	non-NTD S1	pos	1.20	0.14	0.20	0.11	0.11	0.12	-ve
FD 7D	B	non-NTD S1	-ve	1.44	0.12	0.14	0.11	0.11	0.12	-ve
FD 8B	B	non-NTD S1	-ve	1.10	0.14	0.12	0.08	0.10	0.09	-ve
FD 7C	B	NTD	pos	1.90	0.15	0.15	0.13	0.12	0.13	-ve
FG 12C	A	non-NTD S1	pos	1.74	0.14	0.11	0.08	0.08	0.10	-ve
FN 8C	C	non-NTD S1	-ve	0.54	0.16	0.16	0.11	0.09	0.12	-ve
FD 5E	B	non-NTD S1	pos	0.34	0.16	0.16	0.13	0.11	0.11	-ve
EW 9B	B	non-NTD S1	-ve	0.22	0.17	0.11	0.09	0.10	0.09	-ve

S2-specific

Antibody	Case	Domain	SARS-CoV-2[a]				Cross-reactivity[a]			PRNT[b]
			IFA	Spike	RBD	S2	SARS	MERS	OC43	EC_{50} (nM)
FD 10A	B	S2	pos	1.78	0.15	1.32	0.22	0.36	0.39	111.13
FB 1E*	C	S2	pos	1.40	0.11	1.20	0.12	1.01	2.34	36.00
FJ 4E*	B	S2	-ve	0.26	0.18	0.91	0.18	1.08	1.68	75.33
EW 9C*	B	S2	pos	1.22	0.18	1.17	0.14	0.26	1.82	133.33
FG 7A	A	S2	pos	0.28	0.17	1.05	0.12	0.13	0.16	133.33
FM 1A	C	S2	-ve	0.25	0.17	0.88	0.12	0.09	0.14	133.33
FB 9D*	C	S2	pos	1.43	0.12	1.27	0.10	1.50	2.13	-ve
FD 1D	B	S2	pos	0.40	0.15	1.06	0.16	0.22	0.25	-ve
FN 2C*	C	S2	pos	1.77	0.14	1.15	0.09	0.10	2.20	-ve

Controls

Antibody	Case	Domain	IFA	Spike	RBD	S2	SARS	MERS	OC43	EC_{50} (nM)	RBD anchored	ACE2 anchored
CR3022		RBD	pos	1.08	1.31	0.10	2.24	0.11	0.11	42.00	-ve	++
BS 1A		Flu H3	-	0.07	0.09	0.11	0.10	0.09	0.11	-ve	-ve	-ve

[a] A sample (10 µg/ml) was considered positive when the measured extinction is at least 3 times the OD
value of the negative control in the ELISA. CR3022 is an anti-SARS RBD human MAb and BS-1A is
an anti-influenza H3 human MAb. The OD value ≥ 1.00, 0.50-0.99, or ≤ 0.49 is highlighted in deep
green, green and light green, respectively.
[b] The PRNT assay was performed with wild type SARS-CoV-2 at Francis Crick Institute (see methods)

and the 50% effective concentration (EC_{50}) was determined using linear regression analysis.
^{c} ACE2-blocking activity of anti-RBD antibody compared to ACE2-Fc (see methods): +, partial; ++,
IC_{50} > ACE2-Fc; +++, IC_{50} ~= ACE2-Fc; ++++, IC_{50} < ACE2-Fc.
* Memory phenotype.
Abbreviations: IFA, immunofluorescence; RBD, receptor-binding domain; PRNT, plaque reduction
neutralisation assay; ACE2, Angiotensin-Converting Enzyme 2.

Table 2. Competitive binding analysis of anti-SARS-CoV-2 RBD antibodies.[a]

Antibody	Vh	Neut[b]	Cluster
FI 4A	3-21	-	1
EY 6A	3-30	+	1
CR3022*		+	1
VHH72-Fc*		+	1
ACE2-Fc*		+	
FI 3A	3-53	+	2
FI 1C	3-11	+	2
H11-H4-Fc*		+	2
FN 12A	1-69	-	3
FD 5D	3-33	+	3
FD 11A	3-48	+	3
S309*		+	3
FJ 10B	5-10	-	4
FM 7B	1-3	-	4
EZ 7A	5-51	-	4
-ve controls			
EW 8B		-	
EW 9C		+	
Z3B2*		-	

(Column headers for the grid: FI 4A-Bio, EY 6A-Bio, CR3022-Bio, VHH72-Fc-Bio, ACE2-Fc-Bio, FI 3A-Bio, FI 1C-Bio, H11-H4-Fc-Bio, FN 12A-Bio, FD 5D-Bio, FD 11A-Bio, FJ 10B-Bio, FM 7B-Bio, EZ 7A-Bio)

[a] Competitive inhibition:values are shown for percentage inhibition and those with $\geq$ 75% blocking, 50-
74% blocking, and < 50% blocking are highlighted in red, orange and green, respectively.
[b] Neutralisation of antibody against wild type SARS-CoV-2 was analysed in the PRNT assay (+ :
positive; - : negative).
* SARS and SARS-CoV-2 cross-reactive anti-RBD MAb CR3022 and S309 were included as positive
controls. SARS and SARS-CoV-2 cross-reactive anti-RBD nanobodies VHH72 and H11-H4 linked to
the hinge and Fc region of human IgG1 were included as positive controls. ACE2-Fc was included as a
positive control. Anti-influenza MAb Z3B2 was included as a negative control.

Table 3. The antigenic specificity and cross-reactivity of 35 anti-SARS-CoV-2 nucleocapsid antibodies derived from COVID-19 patients.

Antibody	Case	SARS-CoV-2[a]	Cross-reactivity[a]			Antibody	Case	SARS-CoV-2[a]	Cross-reactivity[a]		
			SARS	MERS	OC43				SARS	MERS	OC43
EY 12B	A	1.23	1.24	1.10	1.24	EZ 9B	B	1.28	0.66	0.72	0.91
EY 9C	A	1.42	1.12	1.07	1.39	FD 9B	B	1.11	0.66	0.74	0.47
EY 5A	A	1.40	1.30	1.32	1.32	FD 3E	B	1.14	0.69	0.75	0.81
EY 12A	A	1.34	1.37	1.39	1.45	FD 5B	B	1.06	0.28	0.58	0.42
EY 2A	A	1.28	1.13	0.31	0.33	EZ 7B	B	1.01	0.29	0.29	0.33
EY 3B	A	1.22	1.00	0.22	0.26	EZ 4A	B	1.03	0.37	0.36	0.45
EY 8A	A	0.65	0.63	0.66	0.57	FJ 3C	B	0.98	1.28	1.16	0.86
						FD 4B	B	0.98	0.43	0.44	0.46
EZ 8C	B	1.34	1.11	1.07	1.44	FD 4E	B	0.93	0.53	0.63	0.65
EZ 8B	B	1.38	1.28	1.24	1.12	FD 4C	B	0.84	0.47	0.59	0.54
FJ 6D	B	1.30	1.30	1.24	1.05	FD 8C	B	0.84	0.46	0.47	0.56
EW 1A	B	1.37	1.36	1.31	1.28	EZ 4C-1	B	0.60	0.47	0.26	0.33
EW 9A	B	1.45	1.35	1.24	1.23	EZ 4C-2	B	0.71	0.59	0.58	0.34
EW 5A	B	1.37	1.12	1.04	1.30	EW 10C	B	0.56	0.62	0.57	0.59
EZ 9A	B	1.51	1.28	1.29	1.13	EZ 11A	B	0.53	0.46	0.46	0.23
EW 4C	B	1.24	1.23	1.24	1.31						
FD 6D	B	1.73	1.19	0.12	0.21	FB 9B	C	1.27	1.28	0.13	0.21
EZ 11C	B	1.46	0.73	0.72	1.16	FL 9B	C	0.92	0.60	0.68	0.94
EZ 9C	B	1.53	0.87	0.87	1.27						
Controls											
CR3022		0.07	0.06	0.07	0.17						
BS-1A		0.08	0.07	0.07	0.15						

845   [a] A sample (10 µg/ml) is considered positive when the measured extinction is at least 3 times the OD_{450}
846   value of the negative control in the ELISA. CR3022 is an anti-SARS RBD human MAb and BS-1A is
847   an anti-influenza H3 human MAb. OD values of ≥ 1.00 or 0.50-0.99 or ≤ 0.49 are highlighted in deep
848   orange, orange and light orange, respectively.