Genetic diversity analysis of Chinese plum (*Prunus salicina* L.) based on whole-genome resequencing

Xiao Wei[1,#], Fei Shen[2,#], Qiuping Zhang[1,*], Ning Liu[1], Yuping Zhang[1], Ming Xu[1], Shuo Liu[1], Yujun Zhang[1], Xiaoxue Ma[1], Weisheng Liu[1,*]

1 *Liaoning Institute of Pomology, Yingkou 115009*

2 *Beijing Agro-biotechnology Research Center, Beijing Academy of Agriculture and Forestry Sciences, Beijing 100193*

[#] Authors that contributed equally to this work.

[*] Corresponding author: Qiuping Zhang (e-mail: lbzhangqiuping2@163.com) and Weisheng Liu (e-mail: wsliulaas@163.com)

Abstract

Chinese plum (*Prunus salicina* L.), also known as Japanese plum, is gaining

importance due to their extensive genetic diversity and nutritional attributes beneficial

for human health. Single-nucleotide polymorphisms (SNPs) are the most abundant

form of genomic polymorphisms and are widely used in population genetics research.

Here, we construed high-density haplotype map by whole-genome resequencing of 67

Prunus accessions with a depth of $\sim$20$\times$ to evaluate the genome-level diversity and

population structure. The phylogenetic analysis, the principal component analysis, and

the population structure profiling, indicated that the 67 plum accessions could be

classified into four groups corresponding to their origin location, the southern cultivar

group (SCG), the northern cultivar group (NCG), the foreign cultivar group (FG), and

the mixed cultivar group (MG). Some cultivars from South China were clustered with

the other three groups. The genetic diversity indices including the private allele

number, the observed heterozygosity, the expected heterozygosity, and the nucleotide

diversity of the SCG were higher than those of the NCG. The gene flow from the

SCG to the FG was detected as well. We concluded that the origin center of Chinese

plum was at the Yangtze River Basin in South China. This study provided genetic

variation features and population structure of Chinese plum cultivars, laying a

foundation for breeders to use diverse germplasm and allelic variants for improving

Chinese plum varieties.

Keywords: *Prunus salicina* Lindl.; whole-genome resequencing; population structure;

genetic introgression

1. Introduction

Plum is one of the most important fruit crops in the world, possessing extensive genetic diversity and high economic value. The largest plum producer is China, with an annual production of 6,801,187 metric tons in 2018, accounting for 53.9% of the world's total [1]. Within the genus, *Prunus*, Chinese plum (*Prunus salicina* L., 2n = 2x = 16), also known as Japanese plum, is widely grown for fresh market consumption and the canning industry, which includes both pure Chinese plum and its hybrids with other diploid plum species, such as *Prunus simonii* Carr., *Prunus cerasifera* Ehrh., *Prunus americana* Marsh., and others [2].

Chinese plum possible originates from the Yangtze River Basin, and now there are abundant plum germplasm resources in China [3]. With its long growing history and extensive geographical distribution, there are more than 800 indigenous plum cultivars in China derived from *Prunus salicina* L. alone. Over 600 of them are now preserved at National Germplasm Repository for Plums and Apricots (NGRPA) located in Xiongyue, Liaoning Province, China [4,5]. Yu et al. [6] carried out comprehensive phenotyping of 405 Chinese plum cultivars and its hybrids from NGRPA, totally 32 morphological and agronomic characters were investigated, of which their coefficients of variations (CVs) were calculated ranging from 14.85% to 47.09%, suggesting that the genetic variability of Chinese plums distributed widely.

Genetic variability is the prerequisite for any plant breeding program. Learning the extent and structure of genetic variation in germplasm collections is a crucial step for the efficient conservation and utilization of biodiversity in cultivated crops, and using

diverse plum resources to broaden the genetic base of worldwide plum cultivars is the

critical objective for plum breeders [2,7]. Previous efforts have been made to

understand the genetic diversity of plum better. Zhang and Zhang [8] took the lead in

collecting plum germplasm resources and evaluating their genetic diversity on the

basis of morphological traits and isozyme polymorphisms, whereas, morphological

traits were highly susceptible to environmental factors, so the estimates of genetic

diversity were not precise; while isozymes had a low degree of polymorphism and

hence were not efficient enough for the characterization of germplasm genetic

diversity [9]. Overcoming the above-mentioned problems, DNA-based markers

became widely applied in plum genetic diversity analysis, including random amplified

polymorphic DNAs (RAPDs) [5,10], simple sequence repeats (SSRs) [11-13],

inter-simple sequence repeats (ISSRs) [2,14]. In addition, benefiting from the

advances in next-generation sequencing (NGS) technologies, genotyping by

sequencing (GBS), provides a great wealth of information that makes it possible to

identify thousands of single nucleotide polymorphisms (SNPs), which, after adequate

filtering, allow us to carry out detailed genetic diversity studies [7,15-17].

By conducting DNA-based marker assessments, Chinese plums were generally

classified into two major groups, the southern cultivar group (SCG) and the northern

cultivar group (NCG) [18-20]. The *P. salicina* system in South China was rather

primitive than that in North China basing on a palynology study, and the spread of *P.*

salicina was from the south to the north [21]. However, simply cultivar clustering

works are inadequate for the integrated study of genetic diversity of Chinese plum. As

described in peach, large scale SNP data-based diversity analysis boosted the

deciphering of the evolution and domestication of peach germplasm resources

[22,23].

In the present study, we used SNP markers generated from high depth whole-genome

resequencing data (an average depth of ~20✕) for the first time to elucidate the

pattern of genetic diversity, population structure, and domestication of a diverse *P.*

salicina collection. An in-depth understanding of their genetic relationships could

benefit plum germplasm conservation and utilization, and lead to new cultivar

improvement.

2. Results

2.1 Markers development based on whole-genome resequencing

We collected 67 plum accessions representing different geographic and morphological

characteristics (Table S1). The 67 plum accessions were sequenced using the Illumina

HiSeq 2500 platform with a sequence depth higher than 20×; a totally 462.6 Gb clean

data were retained with an average of 6.9 Gb for each accession after filtering out

low-quality reads. The quality of the sequencing data was high with a Q30 > 85%,

and a GC content slightly fluctuated around 38.0%. The mapping rate to the peach

genome was range from 85.73% to 88.57%, with a average of 87.49% (Table S1). We

called SNPs with the unique mapped reads, and a total of 16,600,033 SNPs and

2,107,664 INDELs were identified across the 67 accessions using SAMtools v1.4

software. After filtering out these in low quality, we obtained 14,549,234 SNPs which

could ensure the accuracy and reliability of subsequent genetic diversity and

population structure analyses (Figure 1). About 43.81% of the SNPs were located in

intergenic regions, and 11.88% were in coding regions. The non-synonymous to

synonymous substitution ratio (dN/dS) for the SNPs in the coding regions was 15.92.

Transitions were found in a proportion of 59.58% (10315764/16600033), with a

transition/transversion ratio (Ts/Tv ratio) of 1.47 (Figure 1, Table 1).

Figure 1. The statistics of markers generated from the whole-genome resequencing of

the 67 plum accessions. a) The chromosome-scale SNP distribution; b) The

distribution of transition/transversion ratio (Ts/Tv ratio); c) The counts of different

types of transitions and transversions; d) The counts of genome-wide insertions and

deletions.

Table 1. Statistics of variations across different chromosomes

Categories	Numbers	Ratio
Raw SNPs	16,600,033	88.73%
InDels	2,107,664	11.27%
Filtered SNPs	14,549,234	87.65%
Intergenic regions SNPs	7,272,007	43.81%
Coding regions SNPs	1,971,748	11.88%
Transition	10,315,764	59.58%
Transversion	6,999,126	40.42%
Transition/Transversion	--	147.00%
Non-synonymous SNPs	1,855,201	94.09%
Synonymous SNPs	116,547	5.91%

2.2 Phylogenetic analysis

To exploit the relationship among the Chinese plum, a Neighboring-joining

phylogenetic tree of the 67 accessions was constructed with all SNPs (Figure 2a). The

phylogenetic tree classified the accessions into four main groups which are

corresponding to their origin location: (1) the southern cultivar group (SCG) was

comprised of the plum cultivars from Sichuan, Guizhou, Yunnan, Guangdong,

Guangxi, Zhejiang, and Fujian Province; (2) the northern cultivar group (NCG)

comprised of plum cultivars mainly from Hebei, Henan, Shandong, Shaanxi Province,

and the south part of Liaoning Province; (3) the foreign cultivar group (FG) included

plum cultivars from the USA and Japan; and (4) the mixed cultivar group (MG)

consisted of newly bred cultivars from the NGRPA and several cultivars originated

SCG and FG. Notably, we fould 'Saozouli' and 'Shuili' cultivars collected from south

China were clustered with NCG, 'Zaohuangli' and 'Jinshali' with FG, and 'Wanshu

huanai', 'Huahongli' and 'Abazhoumeiguili' with MG, suggesting that SCG is more

diverse than any other group.

Figure 2. The population structure of the 67 plum accessions. a) Neighboring-joining

phylogenetic tree constructed using SNPs at fourfold degenerate site. Each group was

color coded; b) Bayesian model-based clustering of the 67 plum accessions with the

number of ancestry kinship (*K*) from 2 to 10; c) Principle Component Analysis (PCA)

of the 67 plum accessions. FG, the foreign cultivar group; MG, the mixed cultivar

group; SCG, the southern cultivar group; NCG, the northern cultivar group.

2.3 Population structure analysis

Population genetic structure analysis was performed based on high-quality SNPs.

We employed 5-fold cross validation to inference the number of ancestor population,

K (Figure 2b). The NCG was divergent from SCG when the K >2, suggesting the

NCG was derived from SCG. We observed that SCG includes individuals with similar

or different genetic structure to these from other groups like NCG and FG, further

indicating that it has higher diversity than others. The observation was further

supported by genetic diversity analysis with π (Figure 3b). Thus, other groups

including FG and NCG potentially associated with SCG during plum evolution.

Figure 3. The genetic diversity of different groups. a) The decay of linkage

disequilibrium (LD) measured as the squared correlation coefficient (r^2) by pairwise

physical distance; b) The nucleotide diversity (π) of different groups; c) The genetic

differentiation analysis between groups. The values between pairs indicate population

divergence (F_{ST}). FG, the foreign cultivar group; MG, the mixed cultivar group; SCG,

the southern cultivar group; NCG, the northern cultivar group.

2.4 Principal component analysis

To further confirm the relationship among the cultivars, we performed principal

component analysis (PCA) toward the 67 plum accessions (Figure 2c). Based on the

principal component plot with first two principal components, the NCG exhibited a

relatively close relationship with SCG, which was consistent with the phylogenetic

tree and population structure analysis. We deduced that the NCG could be considered

as a more independent subgroup of SCG and may be derived from a specific ecotype.

Furthermore, samples from SCG showed some degree of correlation, suggesting the

close relationship between SCG and others. The results further confirm that other

groups potentially originate from SCG.

2.5 Genetic diversity analysis

As shown in Table 2, the expected heterozygosity (He) of the *Prunus* populations

varied between 0.234 and 0.304; the observed heterozygosity (Ho) of the *Prunus*

populations ranged between 0.328 and 0.429; Wright's F-statistic (F_{IS}) of the *Prunus*

populations varied between –0.241 and –0.146; the number of private alleles (A_P) in

the *Prunus* populations varied between 546 and 31,285; the nucleotide diversity (π)

ranged between 0.00358 and 0.00467. The mean nucleotide variation of *P. salicina*

was higher than that of other perennial crops, such as peach ($\pi = 0.0015$) [24], cassava

($\pi = 0.0026$) [25], and apricot ($\pi = 0.0027$) [26], but lower than that of date palm ($\pi =$

0.0092) [27]. The Tajima's D values of the four groups were all tested positive (1.002

– 1.497) and significantly different from zero. Thus, the null hypothesis of neutral

evolution could be rejected. As shown in Figure 3a, the LD decay rate was fastest in

the SCG and was lowest in the FG, suggesting a lower frequency of genetic

recombination in the FG, as they included artificially hybridized cultivars.

Table 2.The statistical values of genetic diversity within different populations.

Groups	Private allele number (Ap)	Observed heterozygosity (Ho)	Expected heterozygosity (He)	Nucleotide diversity (π)	Wright's F-statistic (F_{IS})	Tajima's D
FG	31,285	0.4290	0.3036	0.00467	-0.2411	1.4974
MG	30,112	0.3658	0.2846	0.00433	-0.1560	1.0094
SCG	27,287	0.3365	0.2606	0.00395	-0.1460	1.0022
NCG	546	0.3280	0.2336	0.00358	-0.1829	1.3211

2.6 Genetic differentiation analysis

The F_{ST} between the four groups varied between 0.0485 and 0.1303 (Fig. 3). The F_{ST}

value of MG-FG, MG-SCG, and SCG-NCG pairs was relatively lower among all the

group-pairs analyzed, indicating that the genetic differences within populations were

higher than those between populations and the possible genetic exchange between

populations. The F_{ST} value between the NCG and FG groups were the largest among

all the groups analyzed, which may have been due to geographical isolation, low gene

flow between populations, and significant genetic differences.

2.7 Gene flow analysis and inferred evolutionary path

We analyzed the gene flow between the four geographic groups (Figure 4b, 4c).

Allowing one or two migration events (m = 1 or 2), we observed that gene flow

occurred between the SCG and FG accessions, which likely reflected their many

shared genomic components due to hybridization in their domestication and breeding

histories. Considering the geographical locations of SCG and NCG, though these two

ecological groups were close to each other, we found no gene flow between them,

indicating a relatively independent domestication process.

Figure 4. The inferred evolutionary path (a) and gene flow analysis between

populations as inferred by TreeMix using a model with one (b) and two (c) admixture

events. Admixtures are colored according to their weight. FG, the foreign cultivar

group; MG, the mixed cultivar group; SCG, the southern cultivar group; NCG, the

northern cultivar group.

3. Discussion

SNPs have become tremendously important as markers for genetics research in plants

because they have been found to be in high frequency, display a lower mutation rate

compared to SSR-based markers, and they are uniformly distributed across the

genome [28]. NGS is a kind of low-cost technologies to discover large numbers of

SNPs for extensive genetic studies, including within-species diversity analysis,

linkage map construction, and genome-wide association studies (GWAS), which is

allowing significant advances in plant genetics and breeding [6,22,23,29]. The genus

Prunus has shown conserved intraspecific and intragenic collinearity in the *Rosaceae*

family, and peach has been considered as a model species for the genus *Prunus*

regarding multiple genetic types of research [30-33]. On these bases, for the first time,

we conducted high-depth (~20×) whole-genome resequencing of 67 plum accessions

using the peach genome sequence as a reference, obtaining an average of 6.9 Gb

high-quality data for each acquisition, which ensured the accurate and comprehensive

population genetic analysis.

Based on the high-quality SNPs, the 67 plum accessions in this study were divided

into four groups, the SCG, NCG, FG, and MG, supported by the phylogenetic tree, the

population structure analysis, and the PCA results, which were partly in accordance

with the previously obtained results through RAPD, ISSR, and SSR marker assessing

[18-20]. We noticed that several cultivars from the south part of China distributed in

other three groups (Figure 2a), combining with the fact that the *P. salicina* system in

South China was rather primitive than that in North China basing on a palynology

study, we speculated that, for SCG, the genetic background was relatively broader, the

gene exchange was more frequent with other groups, and the possible origin center of

Chinese plum was at the Yangtze River Basin in South China [3,20,21,34]. For the

five newly bred cultivars from the NGRPA in MG, one of their parents was from the

USA ('Blackamber', 'Friar') or Japan ('Akihime') and the other parent was native to

China, and the gene flow between SCG and FG was detected (Figure 4). Thus, we

assumed that those cultivars in MG were genetically related. As mentioned in other

studies, most of the Chinese plum cultivars from Japan and the improved Chinese

plum hybrids from the USA were found distributed across Chinese indigenous plum

group through dendrogram analysis [2,5,20], which was according with the *P.*

salicina-predominant genetic components of improved Chinese plum hybrids [35],

and in agreement with the fact that China-originated plum cultivars were initially

introduced to Japan, and then imported to the USA in 1870 from Japan by Luther

Burbank [3].

Southwest China was speculated as a primitive domestication center of plum as there

were wild plum populations discovered in Yunnan, Sichuan, and Guizhou Province

[20]. Suggesting by the high Shannon index of diversity, the high value of average

sufficient allele, and the high expected heterozygosity, the Chinese plums cultivated in

southwest China were tested with high genetic diversity using SSR markers [20].

Additionally, by analyzing the phenotyping variation of 405 plum accessions

distributed in different geographical regions in China, the fruit weight of cultivars in

South China exhibited the most significant coefficient of variation [6]. The π, Ho, and

He of the SCG calculated with SNPs in our study were higher than those of the NCG,

corresponding with the higher genetic diversity of the SCG; these genetic diversity indices of the SCG were lower than that of the FG and MG, presumably due to the artificial hybridized cultivars in the FG and MG.

Generally, the genetic differentiation between populations is considered moderate when the F_{ST} value is higher than 0.05 and is considered highly differentiated when the F_{ST} value is higher than 0.15 [36]. In our study, the NCG-FG pair possessed the highest F_{ST} value, following by the SCG-FG couple. In contrast, the F_{ST} value of the MG-FG was significantly lower as there were five newly selected plum cultivars by NGRPA in the MG with foreign lineage from the FG. Human intervention via the artificial selection of favorable phenotypic traits to enhance production and improve desirable agronomic traits can both reduce the levels of genetic variability and skew allele frequencies [37].

4. Materials and Methods

4.1 Plant materials

A diverse collection consisting of 67 *Prunus* spp. accessions, including 65 *P. salicina* accessions and two *P. simonii* accessions, were selected for the whole-genome resequencing study (Table S1). All these trees were held at NGRPA located in Xiongyue county, Liaoning Province, China (40°18′ N, 122°16′ E) in a planting density of 3.0 × 4.0 m, trained to the open vase system with three or four main branches, and maintained under conventional management and pest control operations.

4.2 DNA extraction, library preparation, and sequencing

Genomic DNAs were isolated from the young and healthy leaf samples of the 67

accessions using a modified CTAB protocol [38]. The quality and integrity of the

DNAs were examined using a NanoDrop® spectrophotometer (ND-1000, Thermo

Fisher Scientific Inc., USA) followed by electrophoresis on 1% agarose gels.

Quantification of the DNA samples was assessed with Qubit™ (2.0 Fluorometer,

Invitrogen, Carlsbad, CA, USA).

High molecular weight DNA aliquots with 230/260 and 260/280 ratios ranging

between 1.8 – 2.0 and 1.8 – 2.2, respectively, were then sent to BGI (Shenzhen, China)

for library construction and sequencing. The insert-size of the libraries was 500 bp,

and the length of pair-end reads was 150 bp. All the libraries were sequenced using

the Illumina HiSeq 2500 platform (Illumina, San Diego, CA, USA).

4.3 Reads mapping, SNP calling, and SNP annotation

The qualified paired-end reads of each accession were aligned against the peach

reference genome v2.0 [24] using BWA v0.7.12-r1039 with the parameters mem -t 4

-k 32 –M [39], and SNPs were identified using SAMtools v1.4 [40]. Low-quality

SNPs were filtered out by minimum minor allele frequency (mnMAF < 0.01) and

missing data per site (MDpS > 10%), and finally converted into Variant Call Format

file (VCF). Gene-based SNP annotation was performed using the ANNOVAR

package v2018-04-16 [41]. Based on the reference genome annotation, SNPs were

categorized as occurring in exonic regions (overlapping with a coding exon), intronic

regions (overlapping with an intron), upstream and downstream regions (within a 1 kb

region upstream or downstream from the transcription start site), or intergenic regions.

Additionally, the heterozygosity was calculated using VCFtools v0.1.14 [42].

4.4 Population structure analysis

TreeBeST v1.9.2 software was used to calculate the distance matrix [43]. RAxML

v8.2.12 was used to construct the ML phylogenetic tree, and 1000 bootstrap replicates

were used [44]. The obtained phylogenetic tree was visualized using MEGA v5.05

[45].

The principal component analysis was performed using the PLINK v1.07 software

with default parameters [46].

We used ADMIXTURE v1.23 to infer the population structure [47]. To identify the

best genetic clusters K, cross-validation error was tested for each K value from 2 to 10.

The termination criterion was 1e-6 (stopping when the log-likelihood increased by

less than 1e-6 between iterations).

4.5 Linkage disequilibrium (LD) analysis

Linkage disequilibrium (LD) was calculated using SNPs with MAF greater than 0.05

through PLINK v1.07 software with the following setting: --file --r2 --ld-window

99999 --ld-window-kb 200 --out. LD decay was calculated based on the squared

correlation coefficient (r^2) values between two SNPs and the physical distance

between the two SNPs [46].

4.6 Gene flow analysis

The TreeMix v1.13 software was used to evaluate the gene flow among different

groups with the parameters -se-bootstrap-k 1000 -m, where the number (-m) varied

from one to three [48].

4.7 Genetic diversity and differentiation analysis

To evaluate the genetic diversity and differentiation, we used a 100-kb sliding window

with a step size of 10 kb to calculated the number of private alleles (A_P), the observed

heterozygosity (Ho), the expected heterozygosity (He), the nucleotide diversity (π),

the Tajima's D value, Wright's F-statistic (F_{IS}), and the population-differentiation

statistics (Fixation index, F_{ST}) using VCFtools v0.1.14 [42].

5. Conclusions

In the current study, using high-quality SNPs generated from high-depth

whole-genome resequencing of 67 *Prunus* accessions, we concluded that the origin

center of Chinese plum was at the Yangtze River Basin in South China. This study

also provided genetic variation features of plum cultivars and dissected their genetic

diversity and population structure, laying a foundation for breeders to use diverse

germplasm and allelic variants towards developing improved Chinese plum varieties.

Data availability: All of the raw reads of the plum accessions generated in this study

have been deposited in the public database of National Center of Biotechnology

Information under PRJNA659814.

Supplementary Materials: Supplementary material can be found at http://www.

Authors' contributions: ZQP and LWS conceived and designed the experiments.

WX and SF performed the experiments. WX, SF and ZQP analyzed the data and

wrote the paper. LN, ZYP, XM, LS, MXX and ZYJ collected samples and managed

the materials. LWS revised the manuscript. All authors have read and approved the

final manuscript.

Funding: This work was supported by National Natural Science Foundation of China

(31401826), the Program of Conservation and Utilization of Crop Germplasm

Resources (2018NWB003) and National Crop Germplasm Resources Platform of

China (NICGR2018-056).

Conflict of interest: The authors declare no conflict of interest.

References

1. Food and Agriculture Organization of the United Nations. World production of

    plums and sloes. Available online: http://www.fao.org/faostat/zh/#data/QC

    (accessed on 23 April. 2020).

2. Liu W., Liu D., Zhang A., et al. Genetic diversity and phylogenetic relationships

    among plum germplasm resources in China assessed with inter-simple sequence

    repeat markers. *J. Am. Soc. Hortic. Sci.* **2007**, 132, 619-628.

3. Hartmann W. and Neumulle M. Plum breeding. In Breeding Plantation Tree

    Crops: Temperate Species; Jain, S.M., Priyadarshan, P.M., Eds.; Springer

    Science Business Media: New York, NY, USA, 2009; pp. 161-231.

4. Zhang J. Report on the investigation for national plum and apricot resources.

    China Fruits. **1990**, 4, 29-34.

5. Liu W., Liu D., Feng C., et al. Genetic diversity and phylogenetic relationships

in plum germplasm resources revealed by RAPD markers. *J. Hortic. Sci.*

Biotechnol. **2006**, 81, 242-250.

6. Yu X., Zhang Q., Liu W., et al. Genetic diversity analysis of morphological and

agronomic characters of Chinese plum (*Prunus salicina* L.) germplasm. *J. Plant*

Genet. Resour. **2011**, 12, 402-407.

7. Urrestarazu J., Errea P., Miranda C., et al. Genetic diversity of Spanish Prunus

domestica L. germplasm reveals a complex genetic structure underlying. *PLoS*

One 2018, 13, 1-21.

8. Zhang J. and Zhou E. China fruit-plant monographs: plum flora; China Forestry

Press: Beijing, China, 1998; pp.10-67.

9. Khush G. *Molecular genetics-plant breeder's perspective*; In Molecular

techniques in crop improvement; Jain S.M., Brar D.S., and Ahloowalia B.S.,

Eds.; Kluwer Academic Publishers: Dordrecht, The Netherlands, 2002, pp. 1-8.

10. Ben Tamarzizt H., Ben Mustapha S., Baraket G., et al. Assessment of genetic

diversity and relationships among wild and cultivated Tunisian plums (*Prunus*

spp.) using random amplified microsatellite polymorphism markers. *Genet. Mol.*

Res. **2015**, 14, 1942-1956.

11. Pop R., Hârța M., Szabo K., et al. Genetic diversity and population structure of

plum accessions from a Romanian germplasm collection assessed by simple

sequence repeat (SSR) markers. *Not. Bot. Horti. Agrobot. Cluj-Napoca*. **2018**,

46, 90-96.

12. Zhang Q., Wei X., Liu W., et al. The genetic relationship and structure of some

natural interspecific hybrids in *Prunus* subgenus *Prunophora*, based on nuclear

and chloroplast simple sequence repeats. *Genet. Resour. Crop Evol.* **2018**, 65,

625-636.

13. Acuña C.V., Rivas J.G., Brambilla S.M., et al. Characterization of Genetic

Diversity in Accessions of *Prunus salicina* Lindl: Keeping fruit flesh color

ideotype while adapting to water stressed environments. *Agronomy* **2019**, 9,

1-12.

14. Wu W., Chen F., Yeh K., Chen J. ISSR analysis of genetic diversity and

structure of plum varieties cultivated in Southern China. *Biology* (Basel) **2019**,

8, 1-13.

15. Elshire R.J., Glaubitz J.C., Sun Q., et al. A robust, simple genotyping-by-

sequencing (GBS) approach for high diversity species. *PLoS One* **2011**, 6, 1-10.

16. Salazar J.A., Pacheco I., Silva C., et al. Development and applicability of GBS

approach for genomic studies in Japanese plum (*Prunus salicina* L.). *J. Hortic.*

Sci. Biotechno. **2019**, 1 94, 284-294.

17. Zhebentyayeva T., Shankar V., Scorza R., et al. Genetic characterization of

worldwide *Prunus domestica* (plum) germplasm using sequence-based

genotyping. Hortic. Res. 2019, 6, 12.

18. Qiao Y. Establishment of RAPD, ISSR and SSR reactions system and analysis

of genetic diversity of Japanese plum cultivars. Doctor of thesis, Nanjing

Agricultural University, Nanjing, China, 2003.

19. Liu W. Studies on the genetic diversity among plum germplasm resources and

the phylogenetic relationships of main plum species. Doctor of thesis, China

Agricultural University, Beijing, China, 2005

20. Wei X., Zhang Q., Liu N., et al. Genetic diversity of the *Prunus salicina* L. from

different sources and their related species. *Sci. Agric. Sin.* **2019**, 52, 568-578

21. Guo Z. Collection and utilization of *Prunus salicina* germplasm resource in

South China . Doctor of thesis, Nanjing Agricultural University, Nanjing, China,

2006.

22. Cao K., Zheng Z., Wang L., et al. Comparative population genomics reveals the

domestication history of the peach, *Prunus persica*, and human influences on

perennial fruit crops. *Genome Biol.* **2014**, 15, 1-15.

23. Li Y., Cao K., Zhu G., et al. Genomic analyses of an extensive collection of wild

and cultivated accessions provide new insights into peach breeding history.

Genome Biol. **2019**, 20, 1-18.

24. Verde I., Abbott A.G., Scalabrin S., et al. The high-quality draft genome of

peach (*Prunus persica*) identifies unique patterns of genetic diversity,

domestication and genome evolution. *Nat. Genet.* **2013**, 45:487-494.

25. Kawuki R.S., Ferguson M., Labuschagne M., et al. Identification,

characterisation and application of single nucleotide polymorphisms for

diversity assessment in cassava (*Manihot esculenta* Crantz). *Mol. Breed.* **2009**,

23, 669-684.

26.   Li W., Liu L., Wang Y., et al. Genetic diversity, population structure, and

relationships of apricot (*Prunus armeniaca*) based on restriction site-associated

DNA sequencing. Hortic. Res. **2020**, 7, 69.

27.   Hazzouri K.M., Flowers J.M., Visser H.J., et al. Whole genome resequencing of

date palms yields insights into diversification of a fruit tree crop. *Nat. Commun.*

2015, 6, 8824.

28.   Dirlewanger E., Cosson P., Tavaud M., et al. Development of microsatellite

markers in peach [*Prunus persica* (L.) Batsch] and their use in genetic diversity

analysis in peach and sweet cherry (*Prunus avium* L.). *Theor. Appl. Genet.* **2002,**

105, 127-138.

29.   Pinosio S., Marroni F., Zuccolo A., et al. A draft genome of sweet cherry

(*Prunus avium* L.) reveals genome wide and local effects of domestication.

Plant. J. **2020**, 1-13.

30.   Arús P., Verde I., Sosinski B., et al. The peach genome. *Tree. Genet. Genomes.*

2012, 8, 531-547.

31.   Carrasco B., González M., Gebauer M., et al. Construction of a highly saturated

linkage map in Japanese plum (*Prunus salicina* L.) using GBS for SNP marker

calling. *PLoS. One.* **2018**, 13, 1-14.

32.   Marti A.F.I., Saski C.A., Manganaris G.A., et al. Genomic sequencing of

Japanese plum (*Prunus salicina* Lindl.) mutants provides a new model for

rosaceae fruit ripening studies. *Front. Plant. Sci.* **2018**, 9, 21.

33. Zhang Q., Wei X., Liu N., et al. Construction of an SNP-based high-density

genetic map for Japanese plum in a Chinese population using specific length

fragment sequencing. *Tree. Genet. Genomes*. **2020**, 16, 18.

34. Topp B., Russell D., Neumuller M., et al. Plum. In Fruit Breeding; Badenes M.,

Byrne D.. Eds.; Science Business Media: New York, NY, USA, 2012, pp.

571-621.

35. Faust M. and Suranyi D. Origin and dissemination of plum. *Horticultural*

Reviews **1999**, 23, 179-231.

36. Xu G. Plant population genetics; Science Press: Beijing, China, 2009; pp.

24-78.

37. Tajima F. Statistical method for testing the neutral mutation hypothesis by DNA

polymorphism. *Genetics* **1989**, 123, 585-595.

38. Doyle J.J. and Doyle J.L. Isolation of plant DNA from fresh tissue. *Focus* **1990**,

12, 13-15.

39. Li H. and Durbin R. Fast and accurate long-read alignment with

Burrows-Wheeler transform. *Bioinformatics* **2010**, 26, 589-595.

40. Li H., Handsaker B., Wysoker A., et al. The Sequence Alignment/Map format

and SAMtools. *Bioinformatics* **2009**, 25, 2078-2079.

41. Wang K., Li M., Hakonarson H. ANNOVAR: Functional annotation of genetic

variants from high-throughput sequencing data. *Nucleic Acids. Res.* **2010**, 38,

1-7.

42.   Danecek P., Auton A., Abecasis G., et al. The variant call format and VCFtools.

463   *Bioinformatics* **2011**, 27, 2156-2158.

464   43.   Vilella A.J., Severin J., Ureta-Vidal A., et al. EnsemblCompara GeneTrees:

465   Complete, duplication-aware phylogenetic trees in vertebrates. *Genome Res.*

466   **2009**, 19, 327-335.

467   44.   Stamatakis A. RAxML version 8: A tool for phylogenetic analysis and

468   post-analysis of large phylogenies. *Bioinformatics* **2014**, 30, 1312-1313.

469   45.   Tamura K., Peterson D., Peterson N., et al. MEGA5: Molecular evolutionary

470   genetics analysis using maximum likelihood, evolutionary distance, and

471   maximum parsimony methods. *Mol. Biol. Evol.* **2011**, 28, 2731-2739.

472   46.   Purcell S., Neale B., Todd-Brown K., et al. PLINK: A tool set for whole-genome

473   association and population-based linkage analyses. *Am. J. Hum. Genet.* 2007, 81,

474   559-575.

475   47.   Alexander D.H., Novembre J., Lange K. Fast model-based estimation of

476   ancestry in unrelated individuals. *Genome Res.* 2009, 19, 1655-1664.

477   48.   Pickrell J.K. and Pritchard J.K. Inference of population splits and mixtures from

478   genome-wide allele frequency data. *PLoS Genet.* 2012, 8(11), e1002967.

479