Title: Rest is Required to Learn an Appetitively-Reinforced Operant Task in *Drosophila*

Running Title: Learning of Reward Requires Rest

Author Names: Timothy D. Wiggin 1, Yung-Yi Hsiao 1#, Jeffrey B. Liu 1#, Robert

Huber 2,3, Leslie C. Griffith 1*

1. Department of Biology, National Center for Behavioral Genomics and Volen Center

for Complex Systems, Brandeis University, 415 South St. MS008, Waltham, MA 02454

2. Radcliffe Institute for Advanced Studies, Harvard University, 10 Garden St,

Cambridge, MA 02138

3. Juvatech, 1112 N Reynolds Rd., Toledo, OH 43615

# These authors contributed equally

*Corresponding author:

Leslie C. Griffith

Brandeis University

Department of Biology MS008

415 South Street

Waltham, MA, 02454-9110, USA

griffith@brandeis.edu

Conflict of Interest:

The authors declare no competing financial interests.

ABSTRACT

Maladaptive operant conditioning contributes to development of neuropsychiatric disorders. Candidate genes have been identified that contribute to this maladaptive plasticity, but the neural basis of operant conditioning in genetic model organisms remains poorly understood. The fruit fly *Drosophila melanogaster* is a versatile genetic model organism that readily forms operant associations with punishment stimuli. However, operant conditioning with a food reward has not been demonstrated in flies, limiting the types of neural circuits that can be studied. Here we present the first sucrose-reinforced operant conditioning paradigm for flies. In the paradigm, flies walk along a Y-shaped track with reward locations at the terminus of each hallway. When flies turn in the reinforced direction at the center of the track, sucrose is presented at the end of the hallway. Only flies that rest early in training learn the reward contingency normally. Flies rewarded independently of their behavior do not form a learned association but have the same amount of rest as trained flies, showing that rest is not driven by learning. Optogenetically-induced sleep does not promote learning, indicating that sleep itself is not sufficient for learning the operant task. We validated the sensitivity of this assay to detect the effect of genetic manipulations by testing the classic learning mutant *dunce. Dunce* flies are learning impaired in the Y-Track task, indicating a likely role for cAMP in the operant coincidence detector. This novel training paradigm will provide valuable insight into the molecular mechanisms of disease and the link between sleep and learning.

INTRODUCTION

Learning is a broadly conserved, highly regulated, and health relevant function of the nervous system. Learning updates the frequency of behaviors to reflect stimulus predictability in an animal's environment. The associative forms of learning transfer the value of an innately valued stimulus (an unconditioned stimulus or US) to an associated predictor, either a behavior or cue (Fanselow and Wassum, 2015). US association with internally-generated behavior (e.g. locomotion, static posture, lever press) produces "operant conditioning" across a wide range of animal species (Skinner, 1948; Kimble et al., 1955; Susswein et al., 1986). Operant conditioning allows the animal to modify its behavior to increase the likelihood of obtaining rewarding stimuli and decrease the likelihood of encountering aversive stimuli.

Operant conditioning to reward or relief from punishment incorporates a positive feedback loop – learning increases the generation of the behavior, which in turn increases reward frequency, which strengthens the learned association. This type of positive feedback loop is hypothesized to contribute to diverse neuropsychiatric disorders including childhood anxiety, compulsive behaviors, and chronic pain (Ollendick et al., 2001; Korff and Harvey, 2006; Chóliz, 2010; Gatzounis et al., 2012). Genome-wide association studies have identified candidate genes that increase susceptibility to these operant conditioning-associated disorders (Smith et al., 2016; Levey et al., 2020; Smit et al., 2020). However, despite the relevance to human health, the neural basis of operant conditioning in genetic model organisms remains incompletely understood. It is not currently possible to trace a neural circuit of operant conditioning in animals more complex than *Aplysia californica* (Nargeot and Simmers, 2011), nor has there been a genetic screen for molecular components of operant learning in model organisms. A promising system to address this gap in knowledge is the fruit fly *Drosophila melanogaster*. Much of the known molecular machinery underlying learning and memory was first discovered using genetics in the fly and these molecules have subsequently been shown to be essentially identical in humans (Greenspan and Dierick, 2004). Furthermore, a draft map of the neural connections in a fruit fly hemi-brain has been recently published which, along with advanced genetic

tools, greatly facilitates mapping complex neural circuits (Pfeiffer et al., 2010; Xu et al.,
2020).

Operant conditioning has been studied extensively in flies, but only limited
progress has been made in understanding circuit-level mechanisms. There have been
many operant conditioning paradigms reported in flies: geotaxis training (Murphey,
1967), leg position conditioning (Booker and Quinn, 1981), proboscis extension
suppression (DeJianne et al., 1985), flight simulator heat avoidance (Wolf and
Heisenberg, 1991), conditioned place preference (Wustmann et al., 1996), social
freezing (Kamyshev et al., 1997), and left-right navigation in tethered ball-walking
(Nuwal et al., 2012). However, a pair of landmark publications (Brembs and Plendl,
2008; Brembs, 2009) demonstrated that when predictive sensory cues are available,
flies preferentially learn these sensory cues and block the formation of operant
conditioning. This finding dramatically compromises a number of paradigms that
intended to test operant learning in flies, since predictive sensory information present
during training may have inhibited operant learning. The remaining purely operant
learning paradigms that are routinely used in flies, flight simulator heat avoidance and
proboscis extension suppression, have two important limitations. First, they use
restrained fly preparations, which unavoidably alter animal behavior (Stowers et al.,
2017). Second, they use an aversive US which may not recruit the full repertoire of US
pathway neurons (Liu et al., 2012) and may use neurons outside the brain for learning
(Booker and Quinn, 1981).

In order to extend the range of operant conditioning paradigms in flies, we
developed a positively reinforced, self-paced, operant training task for untethered flies,
which we call the Y-Track. Surprisingly, we found that this operant training paradigm
only produces a major change in behavioral frequency in a relatively small subset of
experimental animals that rest during training. This surprising finding further reinforces
the importance of rest for learning (Maquet, 2001) and opens a new avenue for
measuring this link in a single-session paradigm.

MATERIALS AND METHODS

Experimental Animals

Flies were raised on cornmeal-dextrose-yeast food in bottles at room temperature or in a 25 C incubator with a 12 hour:12 hour light:dark cycle. Wild type (WT) flies were from the Canton-Special (CS) background. Transgenic flies were obtained from the Bloomington Drosophila Stock Center (BDSC) and Vienna Drosophila Resource Center (VRDC) as follows: *P{VT058968-GAL4}attP2* (VT058968-GAL4, VDRC# 204550), *P{w[+mW.hs]=GawB}104y* (104y-Gal4, BDSC# 81014), *PBac{y[+mDint2] w[+mC]=UAS-ChR2.XXL}VK00018* (UAS-ChR2.XXL, BDSC# 58374), and *dnc[1]* (BDSC# 6020). Flies with Gal4 and UAS transgene insertions were outcrossed to a CS background for several generations because we found that *white* knock-out backgrounds may be learning deficient in this task (*data not shown*).

Design of the Y-Track Apparatus

The Y-Track conditioning apparatus was designed as a 4 layered structure (Figure 1). The first (top) layer of the structure was a 3D printed holder for a USB camera (ELP-USBFHD01M) and 3.6mm S-mount lens facing downward toward the track. The second layer was a mount for a red filter (Tiffen #25 Red) to block light blue and green light from optogenetic activation and light landmark experiments. These top layers are supported by four 3D pillars on each side of the apparatus. Red LEDs (630nm, Vishay VLDS1235G) were attached to each pillar and illuminated the Y-Track area. The third layer of the apparatus was the Y-Track itself. Two versions of the Y-Track were tested: a square-walled track and a curved floor track. In the square-walled track, the width of the hallways was 3.5 mm and the height of the hallways was 2.5 mm. In the curved floor track, the track surface was described by a circular arc with a diameter of 9 mm. The width of the top of the hallways was 6.7 mm and the height at the middle of the hallways was 1.5 mm. In both tracks, each of the three hallways that made up the Y shape was 20mm long, the hallways met in the middle of the track, and the hallways had 120° radial spacing. At the end of each arm of the maze was a circular plastic holder for reward filter paper ("food circle") securely screwed to a servomotor (Towerpro MG91). Each food circle had two filter paper slots, one for a sucrose-soaked

filter paper and the other for a water-soaked filter paper. The top of the Y-Track was

covered by a clear acrylic plate with a small hole for aspirating flies, machined by the

Brandeis University Machine Shop. A small 3D frame with 3 RBG LEDs (Broadcom

HSMF-C114) was superglued below the Y-Track to deliver optogenetic stimulation and

landmark location cues. The fourth (bottom) layer of the Y-Track apparatus is a frame

that positions the servomotors correctly relative to the Y-Track and secures the entire

apparatus to the base. Modelling of the 3D printed components was done using

Autodesk Fusion 360 (San Rafael, CA). These components were fabricated from

Polylactic Acid (PLA) filament in the Brandeis MakerLab.

This apparatus was controlled by a custom Java program running on an Udoo

X86 Advanced Plus single-board computer. The JavaGrinders library was used to

interface with the camera and servomotors (Donelson et al., 2012). Servomotors were

controlled via a Phidget Advanced Servo (Phidgets, Calgary, CA). Red LEDs

illuminating the Y-Track were powered by a BuckPuck (LuxDrive, 03021-D-E-700). The

Y-Track apparatus and electronics were securely mounted inside a custom-built particle

board box to provide environmental isolation. The internal walls of the box were painted

white to reduce visual cues and a 120mm low-noise ventilation fan was installed to

prevent overheating. Each Y-Track single-board computer was connected to a central

control computer and controlled remotely via Virtual Network Computing (VNC). Code

and 3D models are available on GitHub (https://github.com/Griffith-Lab).

Learning Assay

Flies were collected when 0-1 days old and housed in mixed-sex vials for 24

hours to allow mating. The flies were then screened under light CO_2 anesthesia and

stored in single-sex vials of up to 20 flies each. Flies were housed in a 25 C incubator

that was only accessed during the lights-on period for 7 days prior to the experiment to

ensure circadian entrainment. Each vial of flies was flipped onto fresh food at 5 days old

(48 hours before training), flipped onto a food-deprivation vial at 6 days old (24 hours

before training), and trained in the Y-Track at 7 days old. Food-deprivation vials were

made by inserting a kimwipe soaked with 1mL of tap water into an empty vial.

Prior to introducing the flies into the Y-Track, filter paper was prepared for the

food circles by pipetting 30µL of 2M sucrose solution (reward stimulus) or tap water

(neutral stimulus) and allowing the paper to dry overnight. The dried filter papers were

securely placed into the food circles, and the positioning of the servomotors was

adjusted to ensure that the flies could access only the intended stimulus and were not

able to escape. Following the final positioning of the motors, a reference image of the Y-

Track without a fly present was captured for image background subtraction during the

experiment. Finally, the reward direction for the experiment and the sex of the

experimental animal was chosen based on the experimental design. In the square-wall

experiment, half of the flies were rewarded for turning right and the other half were

rewarded for turning left. We found no difference between training efficacy between the

reward directions, so subsequent experiments used left-turn rewards for all animals. In

the square-wall experiment, half of the animals were male and half were female. We did

not find a significant difference in learning between male and female flies in this

experiment, but we found that males were more susceptible to starvation stress than

females, consistent with previous literature (Jang and Lee, 2015). In order to reduce

inter-individual variability, subsequent experiments used only female flies. Training was

performed during the lights-on period of the fly's circadian day, Zeitgeber Time (ZT) 0-9.

At the beginning of a standard training session, a single fly was aspirated out of

the food deprivation vial into a Y-Track apparatus, and the lid of the track was secured

in place to prevent escape. The Java control program was initialized to run the

remainder of the experimental protocol, as follows: 1) The fly was given five minutes to

acclimate to the maze with no sucrose presented. This acclimation time was a fixed

interval and not dependent upon fly locomotion. 2) Block 0 began (Trials 1-20) and

left/right turn decisions were recorded. No sucrose was presented. Block 0 was a

locomotion-dependent acclimation period to ensure the fly is navigating the track. 3)

Block 1 began (Trial 21). All servomotors turn to present the sucrose-soaked filter paper

to the fly. A trial was initiated when the fly came within 6 mm of the center of the arena

("center zone"). If the fly back-tracked into the arm of arena it previously occupied, the

servomotor turned and presented the water-soaked filter paper until the fly re-entered

the center zone, but the trial continued. If the fly turned in the unrewarded direction, the

servomotor turned and presented the water-soaked filter paper, ending the trial. If the fly

turned in the rewarded direction, the servomotor did not turn, and the fly was given

access to the sucrose-soaked filter paper, ending the trial. After a rewarded trial, the fly

was given 10 seconds to consume sucrose. If the fly did not initiate a new trial by

entering the center zone within 10 seconds, the servomotor turned and presented the

water-soaked filter paper until the fly initiated a new trial. 4) After Trial 160, the fly was

removed from the Y-Track.

In the open-loop, yoked-control experiment (Figure 3), acclimation and trials were

defined exactly as in the standard protocol. However, instead of trials being rewarded or

unrewarded based on turn direction, trial outcome was determined by the reward/non-

reward sequence of a previously run fly. In the visual land-mark experiment (Figure 4),

the training protocol was the same as the standard experiment but a single green LED

(525 nm peak) was illuminated under the Y-Track. In optogenetics experiments, flies

were fed food supplemented with either 1.6 mM all trans retinal (ATR) dissolved in

ethanol (4% final concentration), or ethanol alone as a Vehicle control. Food deprivation

vials were also supplemented with ATR or Vehicle in the same concentration as the

food. During ATR supplementation flies were housed in the dark to prevent premature

activation of ChR2.XXL expressing neurons. During training, blue LEDs (470 nm peak)

were illuminated for 5 minutes after the initiation of Trial 50 (Block 2).

Quantification of Activity and Rest

During training, the frame-by-frame coordinates of each fly (acquired at 20Hz),

trial times, and trial outcomes were recorded. Coordinates were processed following

training to remove incorrect detections, which were identified by fly coordinates outside

the Y-Track region or a change in position faster than a fly could plausibly execute

(Mendes et al., 2013). Gaps in data introduced by this error checking were filled by a

linear interpolation of fly position. The position of the fly over time was used to

determine when the fly was active: activity episodes were continuous periods of

movement greater than 1 px / frame (approx. 50 ms), in which the fly also exceeded 2

px / frame at least once. The activity/inactivity sequence was used to find rest episodes,

which were then used to classify flies as drowsy (3 minutes or more of rest in Block #1-

3), late-resting (9 minutes or more minutes of rest in Block #4-6), or restless. Finally, the

error corrected sequence of left/right turn directions was compared to the real-time turn

direction determined during training. If more than 10% of the trial outcomes differed

between the real-time and *post-hoc* methods, the data from the fly was excluded from

further analysis due to suspected tracking errors. Analysis code implementing this

process is available on GitHub (https://github.com/Griffith-Lab).

Simulation of Turn Behavior with Turn-Direction and Alternation Bias

The interaction of turn-direction preference and alternation preference was

determined using a simulation designed to be analogous to the Y-Track. Simulated, or

in silico, flies occupied one of three positions: *P1*, *P2*, or *P3*. Like the arms of the Y-

Track, these positions were arranged clockwise in a circle, so that *P1* was to the "left" of

P2, *P2* was to the left of *P3*, and *P3* was to the left of *P1*. At each simulation step, the *in*

silico fly moved from its current position to a new position. The *in silico* fly was given a

pre-determined preference for/against alternation and for/against left-hand turns in the

range -2 to 2, where 0 was indifference. The choice of destination was based on a

biased random process based on the previous location occupied and the turn-direction

preference. Alternation and turn-direction preference could combine additively or offset

one another, depending on the sequence of positions visited. These preferences were

combined into *net bias = 2^(alternation preference + left-turn preference)* or

2^(alternation preference - left-turn preference), for the additive and offsetting case,

respectively. This net bias was used to find the turn probability: *P = 1 – 2^(-net bias)*.

Each simulation was run for 1000 iterations to create a statistically representative

sample of *in silico* behavior. Code implementing this simulation is available on GitHub

(https://github.com/Griffith-Lab).

Confocal Imaging of Fly Brains

To measure the expression pattern of Gal4 driver lines, adult fly brains were

dissected at 2-5 days old and prepared for imaging. Dissection, fixation, staining, and

mounting were performed according to the FlyLight protocol (Meissner et al., 2018).

Primary antibodies were rabbit anti-GFP (Invitrogen, RRID:AB_221569) and mouse

anti-Bruchpilot (nc82, RRID:AB_2314866) (Wagh et al., 2006). Samples were mounted

in DPX (dibutyl phthalate in xylene, Sigma) and imaged with a Zeiss LSM 880 confocal

microscope.

Experimental Design and Statistical Analysis

Experiments were designed with change in Preference Index (ΔPI) as the

primary measure of learning. Preference Index (PI) was defined as the preference for

the rewarded turn direction and equal to (#correct turns - #incorrect turns)/#total turns.

ΔPI was defined as the difference between the PI in Block 7 and Block 1. Learning was

measured by performing a right-side, one-sample bootstrap test testing the null

hypothesis that mean ΔPI is less than or equal to zero. The two-sample, two-sided

bootstrap test was used to compare the magnitude of learning in groups of flies.

Hypothesis testing was performed using 50,000 bootstrap samples; the minimum

resolvable p-value using this method is approximately 0.0001. Comparisons of behavior

between flies in square-walled and curved-floor Y-Tracks were performed using Two-

Factor ANOVAs and Tukey-procedure protected *post-hoc* tests. Within figures, groups

that are not statistically different are identified by the same letter assignment.

Coordinate data and statistics were calculated using MATLAB (MathWorks, Natick, MA)

and 0.05 was used as the p-value for statistical significance. The analysis code used to

calculate bootstrap statistical tests is available on GitHub (https://github.com/Griffith-

Lab).

RESULTS

Flies That Rest Learn the Operant Contingency in a Novel Sucrose-seeking Task

We used an ethology-informed approach to design a positive-valence operant

conditioning paradigm. Flies locomote spontaneously while awake (Martin et al., 1999),

forage for food in open fields (Hughson et al., 2018), and are adept at navigational tasks

(Warren et al., 2019). We therefore used food-seeking and navigation as the central

features of the learning paradigm (Figure 1A). Flies are individually loaded into a Y-

shaped track (Y-Track). At the terminus of each arm of the track is a reward location

that can be switched between a food reward and a neutral stimulus. Food reward is only

available when the flies turn in the in the rewarded direction (*i.e.* left or right) relative to

their previous location in the track. Because the rewarded choice is defined relative to

the location of the animal, the location of the next rewarded location changes based on

the previous behavioral choice - no single arm of the track is preferentially rewarded.

Over many left/right choices ("trials"), the turn preference index (PI) is calculated for

blocks of 20 trials as *PI = (# correct turns – # incorrect turns) / # total turns.* Because

baseline left/right turn preference is idiosyncratic to individual flies (Buchanan et al.,

2015), learning is measured as the change in PI (ΔPI) across training to determine if the

flies increase their preference for turning in the direction of food reward.

Implementing this task in a physical apparatus required satisfying several design

constraints (Figure 1B,C). First, the animal must be alert, healthy, and active to engage

in spontaneous locomotion and learning. We included a loading port in a tightly fitted Y-

Track lid that allowed us to load and remove flies without anesthesia using gentle

aspiration. Second, the apparatus must include a detector element that records the

performance of the reinforced behavior in real time. We used JavaGrinders real-time

video tracking to measure locomotor behavior and turn choices (Donelson et al., 2012).

Third, the apparatus must be able to actuate reward delivery based on the behavioral

contingency. We used closed-loop control to allow the real-time tracker to activate

servomotors at the terminus of each arm of the Y-Track and present either 10 seconds

of access to a food reward (filter paper pre-soaked with 2M sucrose) or a neutral

stimulus (plain filter paper). Importantly, the servomotors are positioned to present

sucrose at both termini while the fly is in the center of the Y-Track. The flies are not able

to determine which arm is rewarded simply by smelling or seeing reward. In trials where

the fly turns in the non-reinforced direction, the servomotor is actuated rapidly so that

the fly is never able to actually obtain food.

In order to validate the sensitivity of the real-time tracking, we compared long-

term recordings of living flies and dead flies (*i.e.* flies that have no genuine locomotion;

$n = 2$ per group; Figure 1D). We found that the tracked position of the dead flies was

contained within a radius of 1 pixel over several hours. Locomotor episodes were

therefore defined as continuous sequences of frames in which the fly moved at least

one pixel, with the requirement that the fly must exceed a speed of 2 pixels/frame (0.38

mm/s) for at least one frame. Flies frequently paused between locomotor episodes,

sometimes for extended periods of time. We defined pauses of greater than 1 minute as

"rest" (Figure 1E).

Throughout the prototyping process, we evaluated the effectiveness of our

apparatus in shaping wild type (WT) fly behavior. In pilot experiments (n = 15, WT flies,

mixed sex), we found a small training effect of making sucrose available contingent

upon turn direction in the center of the Y-Track. Interestingly, change in turn direction

preference was correlated with time spent resting during training (Pearson's R = 0.50).

In order to rigorously test the hypothesis that rest is correlated with learning in the Y-

Track, we trained a large cohort of flies (n = 85 female, 87 male; Figure 1F-I). Of this

cohort, 11 (7%) rested early in training (3 minutes or more in Block #1-3). An additional

16 (9%) had high rest late in training (9 minutes or more in Block #4-6) and were

excluded because WT flies reduce food seeking behavior during high sleep times

(Donelson et al., 2012). Flies that did not rest (139 of 172) showed no increase in turn

preference in the direction of reward, indicating that they did not learn the task (One-

sample bootstrap test; p = 0.894). However, consistent with our pilot results, flies with

early rest (11 of 172) had a significantly increased likelihood of turning in the direction of

reward (One-sample bootstrap test; p = 0.031; Figure 1G-I). These results indicate that

WT flies learn a sucrose-rewarded operant contingency only when they rest in the first

half of the training trials. Because of the behavioral importance of these rest-defined

groups, we will refer to flies that rest early in training as "drowsy" flies, and flies with low

rest as "restless" flies.

Y-Track Geometry Significantly Affects Thigmotaxis and Spontaneous Alternation

Operant conditioning paradigms designed for flies can be confounded by sensory

cues; when presented with both an operant contingency and a classical prediction cue,

flies preferentially attend to the classical cue (Brembs and Plendl, 2008). No classical

cues were intentionally introduced into the Y-Track, but *post-hoc* examination of

locomotor behavior of the in the apparatus revealed strong thigmotaxis behavior (Figure

2A). This is consistent with the behavior of flies in open-arenas (Simon and Dickinson,

2010), but it is potentially problematic for the Y-Track task for three reasons. First, if flies

maintain contact with the wall through the vertex of the Y-Track, turn direction is

correlated with a unilateral touch stimulus, which may act as a classical predictor.

Second, it is unclear where the "choice point" for choosing a turn direction is located –

presumably at whatever track location the flies "attach" to one of walls. Third,

thigmotaxis may contribute to spontaneous alternation (Lewis et al., 2017), another

behavior typical of unmanipulated WT flies. Spontaneous alternation would not

independently result in flies preferring the rewarded turn direction, but, in simulated

behavior, spontaneous alternation magnifies small turn biases into large turn preference

indices (Figure 2C). If the effect of early rest is to modulate spontaneous alternation, it

may be that restless and drowsy flies have the same mild change in "true" turn bias and

the difference in turn preference index is due to changes in alternation.

To address the sensory-motor confounds of thigmotaxis, we designed a second

iteration of the Y-Track with gently curved floor, similar to open-field arenas (Simon and

Dickinson, 2010). Thigmotaxis was dramatically reduced in the curved floor track

compared to the square wall track, including as the flies pass through the vertex of the

track ($n = 103$ WT female; Figure 2B). Median distance from the wall is significantly

smaller in the square-wall design than it is in the curved floor design for both drowsy

and restless flies (Two-way ANOVA; $F_{(1,1)} = 2788$; $p < 0.0001$; Figure 2D). There was

no difference in thigmotaxis between drowsy and restless flies within experiments (Post-

hoc test; all $p > 0.17$).

To quantify any link between thigmotaxis, spontaneous alternation, and revealed

turn preference, we measured spontaneous alternation at the end of training for each of

our groups of WT flies. Track geometry significantly affected alternation rate (Two-way

ANOVA; $F_{(1,1)} = 11$; $p = 0.001$; Figure 2E) and drowsy flies have significantly lower

alternation than restless flies (Post-hoc test; all $p < 0.005$). Although spontaneous

alternation may be related to thigmotaxis and rest, spontaneous alternation changes in

the wrong direction to confound the observed interaction between learning and rest.

Learning to Turn Toward Sucrose is Independent of Track Geometry but Dependent
upon an Informative Operant Contingency

The curved floor track geometry dramatically reduces thigmotaxis (Figure 2), so
we repeated Y-Track conditioning in curved floor apparatus to determine if the sensory-
motor experience of thigmotaxis contributes to Y-Track learning (Figure 3A). Learning in
the curved floor track ($n = 103$ WT female flies) was similar to learning in the square
wall track. The percentage of drowsy flies was not different between the square wall and
curved floor tracks (Fisher's Exact Test; $p = 0.8$). Drowsy flies in the curved floor track
(8 of 103) significantly increased their likelihood of turning toward reward (One-sample
bootstrap test; $p < 0.0001$) while restless flies (94 of 103) had no increase in turn
preference (One-sample bootstrap test; $p = 0.918$; Figure 3B-D).

In order to control for any confounding effect of time spent in the experimental
apparatus on turn direction preference, we also performed an open-loop, "yoked"
control in the curved floor track ($n = 110$ WT female flies). In the yoked control flies, the
reward sequence of a previously trained fly was presented to a naïve fly independent of
the turning behaviors of the naïve fly. The yoked control flies therefore had the same
amount of sucrose/reward access as trained flies, but there was no behavioral
contingency. As predicted, neither drowsy (8 of 110) nor restless (101 of 110) yoked
control flies significantly increase turn preference in the pseudo-rewarded direction
(One-sample bootstrap test; $p = 0.476$ and $p = 0.392$, respectively; Figure 3E-H).
Together, these results allow us to conclude that the change in turn preference is due to
learning of the operant contingency and is not dependent upon sensory-motor feedback
from thigmotaxis. The curved floor Y-Track was used for all subsequent experiments
because learning is equivalent to the square wall track, without the potential
confounding effect of thigmotaxis.

A World Orientation Cue does not Facilitate Learning the Operant Task

The curved floor Y-Track experiment (Figure 3) eliminated the potential for an
egocentric sensory-motor confounding cue. We next considered the possibility that the
flies were attending to an inadvertently introduced world-orientation cue rather than the
operant contingency. In order to test the hypothesis that flies are able to use orientation

information to learn the task, we trained flies in the presence of a strong orientation cue
(n = 81 WT female flies). A green LED was illuminated beneath one arm of the Y-Track,
creating a stable, mildly attractive, landmark (Figure 4A). In the presence of this
landmark, neither drowsy (9 of 81) nor restless (66 of 81) flies significantly increase turn
preference in the rewarded direction (One-sample bootstrap test; p = 0.500 and p =
0.695, respectively; Figure 4B-D). This result indicates that the presence of a world
orientation cue does not enhance, and may inhibit, learned change in turn preference.
Together, the results of the thigmotaxis, alternation, and orientation-cue experiments
indicate that neither sensory inputs nor motor patterns explain the change in turn
preference in the direction of reward observed in drowsy flies. We conclude that the
learning produced in this paradigm is navigational and operant in character.
422
Sucrose Promotes Rest and is the Learning-Relevant US in the Y-Track
Operant conditioning is a learned association between behavior and a US, and
the strength of the US influences the strength of the learned association (Rickard et al.,
2009). Rest is strongly regulated by feeding and nutritional state (Murphy et al., 2016),
and the consumption and hedonic value of sugar is dependent upon the state of the fly
(Krashes et al., 2009; Li et al., 2020). While we have shown that there is a correlation
between early rest and learning, it is not clear how they are connected mechanistically.
One possibility is that there may be a between-flies difference in the consumption
of the sucrose US that explains enhanced learning in drowsy flies. In order to test the
hypothesis that drowsy flies eat more sucrose and therefore receive a larger reward, we
analyzed the relationship between time spent adjacent to sucrose and learning in the
WT fly data reported above (Figures 1 and 3). We found that there is a significant main
effect of early rest on time adjacent to sucrose where drowsy flies spend more time next
to the sucrose (Two-way ANOVA; both $F_{(1,1)}$ > 4.5; both p < 0.035; Figure 5A).
Importantly, both drowsy and restless flies spend enough time adjacent to the sucrose
US (>5 seconds on average) to consume a sugar meal (approximately 2 seconds; Ro et
al., 2014). Given this observed difference, we hypothesized that longer interactions with
the US would result in greater learning. However, we found no correlation between time
spent adjacent to sucrose and learning in the square-wall or curved-floor experiments

(Pearson correlation; Square Wall R = 0.12, Curved Floor R = -0.01; Figure 5B). This

analysis indicates that increased sucrose consumption is associated with additional rest

but is not associated with increased learning.

To validate this *post-hoc* analysis demonstrating that sucrose consumption

promotes rest, we trained flies in a Y-Track with no sucrose available (n = 78 WT female

flies). Remarkably, only 2 flies in this cohort had early rest, a 3-fold reduction in the

fraction of drowsy flies compared to training with sucrose available (Figure 5D) and

consistent with the sleep-suppressing effect of food deprivation on flies (Keene et al.,

2010). The restless flies in the no-sucrose experiment did not have a significant

increase in turn preference following training (One-sample bootstrap test; p = 0.119;

Figure 5E-F). Together, increased time adjacent to sucrose in drowsy flies and the

reduction in the number of drowsy flies when sucrose is removed from the Y-Track,

indicate that sugar consumption promotes rest. The lack of residual learning of a

secondary reinforcer in the Y-Track when sucrose is removed also shows that sucrose

is the learning-relevant US.

Optogenetically-Induced Sleep is Not Sufficient to Enhance Learning

Activation of several genetically-targetable cell types in flies is sufficient to induce

sleep (Donlea et al., 2011; Liu et al., 2016). Activation of these cells has the same effect

on sleep-dependent learning as spontaneous or pharmacologically-induced sleep

(Donlea et al., 2011; Dissel et al., 2015). We therefore tested the hypothesis that sleep

is sufficient to enhance learning of turn direction in the Y-Track by optogenetically

activating sleep-promoting neurons. Flies do not synthesize the cofactor of Channel

Rhodopsin (ChR), All-Trans Retinal (ATR), so ATR needs to be added to the food to

functionalize the channels (Zhang et al., 2006). We tested optogenetic sleep induction

using a dorsal Fan-Shaped Body driver (dFSB; 104y-Gal4; n = 39 ATR, 45 Vehicle), or

an Ellipsoid Body driver (EB; VT058968-Gal4; n = 56 ATR, 54 Vehicle) driving

expression of ChR2.XXL (Dawydow et al., 2014), and WT control flies (n = 43 ATR, 36

Vehicle) (Figure 6A). Sleep was induced by turning on blue LEDs located under the Y-

Track for 5 minutes at the mid-point of Training Block 2 (Figure 6B). Lights were

symmetrically located in all arms of the Y-Track, so they did not provide a world

orientation cue. Blue light did not increase rest in WT flies fed ATR or vehicle, but

dramatically increased rest in the dFSB and EB driver flies (Figure 6C,D). The drivers

differed in both potency and on/off kinetics: the dFSB driver rapidly induces sleep in all

ATR fed flies, and flies woke from sleep shortly after the light was turned off, consistent

with prior reports (Tainton-Heap et al., 2020). In contrast, the EB driver induced a lower

level of sleep that persisted even after the blue light was removed (Figure 6B). Both

drivers dramatically increased the % flies with early rest (Figure 6D). Turn preference

for reward was increased in drowsy WT flies (9 of 74), but not restless WT flies (62 of

74) (One-sample bootstrap test; $p = 0.041$ and $p = 0.494$, respectively; Figure 6E).

Significant learning in the drowsy WT flies indicates that the changes in food and blue

light illumination in this experiment did not prevent rest-dependent learning. However,

turn preference for reward was not increased in either drowsy (43 of 84) or restless (36

of 84) flies in the dFSB induction experiment (One-sample bootstrap test; $p = 0.775$ and

$p = 0.964$, respectively; Figure 6F). Nor was turn preference for reward increased in

either drowsy (41 of 104) or restless (57 of 104) flies in the EB induction experiment

(One-sample bootstrap test; $p = 0.098$ and $p = 0.335$, respectively; Figure 6G). The

failure of drowsy flies to learn in the rest induction experiments implies that

optogenetically-induced rest does not enhance learning. And because rest is induced in

a large fraction of the flies without also enhancing their learning, the group of

spontaneously drowsy flies that have learned is diluted with non-learning flies,

effectively erasing the difference between the restless and drowsy groups. We conclude

that spontaneous early rest is associated with increased learning of the navigational

task, but that rest itself is not sufficient to induce learning.

Y-Track Operant Learning is Dependent on cAMP Regulation

The Y-Track task we have developed is an operant, navigational, sucrose-

reinforced learning paradigm, in which learning is revealed by changes in turn

preference from baseline (Figure 1,3). Many neurotransmitters, receptors, and second

messengers necessary for classical conditioning in flies have been identified using

genetic knockouts (Margulies et al., 2005). While cyclic adenosine monophosphate

(cAMP) is important for the formation of classical conditioning (Zars et al., 2000), a

previous study found that activity-regulated cAMP synthesis is not necessary for the formation of aversive operant conditioning (Brembs and Plendl, 2008). In order to determine if regulation of cAMP is necessary for formation of appetitive operant conditioning, we tested the performance of flies mutant for the *dunce* phosphodiesterase (*dnc¹*). *dnc¹* flies fail to learn in classical paradigms and fail to modulate cAMP in response to learning stimuli (Gervasi et al., 2010). In an independent cohort (*n* = 209 WT female flies), we found that drowsy flies (19 of 209) have significantly greater learning than restless flies (183 of 209) (Two-sample bootstrap test; *p* = 0.035; Figure 7A-C). In contrast, flies carrying the *dnc¹* mutation failed to show learning (*n* = 95 female flies): neither drowsy (11 of 95) nor restless (79 of 95) *dnc¹* files significantly increased their likelihood of turning toward reward (One-sample bootstrap test; *p* = 0.181 and *p* = 0.591, respectively; Figure 7D-F).

DISCUSSION

The formal study of associative learning has been remarkably successful: experimentally-induced associative memory has been demonstrated across the animal kingdom and dozens of genes, neurotransmitters, second messengers, and neural structures have been implicated in its formation (Mayford et al., 2012). Within the context of fly learning, animals have been trained to texture (Platt et al., 1980), sound (Menda et al., 2011), color (Schnaitmann et al., 2010), location (Wustmann et al., 1996), and odor (Quinn et al., 1974), among other cues. The fly learning literature developed rapidly, with first reports of training paradigms for classical conditioning to an aversive US, operant conditioning to an aversive US, and classical conditioning to a rewarding US occurring within a decade of one another (Quinn et al., 1974; Booker and Quinn, 1981; Tempel et al., 1983). In this report, we present the first paradigm for operant conditioning of flies to a sucrose US: Y-Track conditioning. The magnitude of the learning observed in WT flies (mean ΔPI = 0.2-0.3; Figure 1,3,7) is modest compared with immediate memory in highly-optimized olfactory learning paradigms (Flyer-Adams et al., 2020), but is similar to the magnitude of learning observed in early reports of olfactory and visual classical conditioning paradigms (Tempel et al., 1983; Vogt et al., 2014), and is also similar in strength to consolidated olfactory memories (Krashes et al.,

2007). Like consolidated memory (Berry et al., 2015; Chouhan et al., 2020), the learning

we observe is dependent upon rest (Figure 1,3). However, rest itself is not sufficient for

learning as we see no learning enhancement in optogenetically induced rest (Figure 6),

indicating that rest does not indiscriminately promote learning in the Y-Track. Y-Track

learning does not depend on sensory cues (Figure 2,4) and requires cAMP as a second

messenger (Figure 7). The properties of Y-Track learning have implications for

navigation, learning and memory, and the connection between sleep and learning in

Drosophila.

Learning, Memory and Rest in Drosophila

Disorders of sleep and associative learning co-occur in several categories of

neurological disease. Primary sleep disorders and sleep deprivation result in decreased

cognitive and memory performance (Kessler et al., 2011; Shekleton et al., 2014;

Zamarian et al., 2015). Conversely, plasticity disorders such as neurodevelopmental

disability and post-traumatic stress disorder have co-morbid sleep abnormalities

(Angriman et al., 2015; Gilbert et al., 2015). Finally, neurodegenerative disorders,

including dementia, Huntington's disease, and Parkinson's disease, frequently disrupt

both sleep and memory (Chaudhuri and Naidu, 2008; Morton, 2013; Robbins and

Cools, 2014; Porter et al., 2015). The widespread connections between learning and

sleep in human neurological disease indicate that there are neuronal circuits linking, or

shared by, sleep and learning in humans. Similar to the sleep/learning connection in

human disease, sleep and learning regulate one another in flies: sleep deprivation

decreases learning (Seugnet et al., 2008; Chouhan et al., 2020) and learning increases

time spent asleep (Ganguly-Fitzgerald et al., 2006). Remarkably, increased sleep is

sufficient to rescue memory formation in learning mutant flies, aged flies, and in a fly

model of Alzheimer's disease (Donlea et al., 2014; Dissel et al., 2015), demonstrating

that the shared circuit can be therapeutically useful.

Learning in the Y-Track conditioning task we have developed depends upon rest

during training (Figure 1,3). We have described prolonged locomotor pauses as "rest"

rather than "sleep" because sleep has a precise, three-fold definition (quiescence,

increased arousal threshold, homeostasis) and it is not possible to properly evaluate

this definition on the Y-Track (Hendricks et al., 2000; Shaw et al., 2000). We

hypothesize that the locomotor pauses that we classify as rest are "sleep-like." In light of

the extensive links between sleep and learning in flies, we proposed two alternative

hypotheses that could account for the correlation between rest and learning in the Y-

Track: First, we hypothesized that learning could drive increased rest. This hypothesis is

inconsistent with the similar amounts of rest observed in populations of flies that learn

and those that do not (*i.e.* yoked controls and *dunce* mutants; Figure 3,7). Second, we

hypothesized that rest drives increased learning. Sleep is known to promote

consolidation of memory both in mammals and flies (Buhry et al., 2011; Donlea et al.,

2011). However, we do not find that optogenetically-induced rest promotes learning

(Figure 6). This failure to promote learning could be due to optogenetically-induced rest

not promoting the learning-associated sleep state (Liu et al., 2019; Wiggin et al., 2020),

as the brain-state of flies with dFSB-induced sleep is significantly different from the

brain-state of spontaneous sleep (Tainton-Heap et al., 2020). It is also possible that the

precise timing of rest is important to its learning-associated function and the timing of

optogenetically-induced rest is suboptimal. Based on the failure of optogenetically-

induced sleep to promote learning, we refine our hypothesis to propose that rest acts as

a gate to learning. In this model, a coincidence between behavior and US must be

detected, presumably by a cAMP-dependent mechanism (Figure 7). Following this

coincidence detection event, sleep is required within a tight temporal window in order to

consolidate the memory and prevent locomotion-related forgetting (Berry et al., 2015).

Neural Circuits of Navigation as a Locus for Operant Plasticity

Identifying plastic neuronal circuits that are responsible for learning is a subject of

intense interest in the effort to understand learning, memory, and cognition. In flies, the

mushroom bodies are the best studied locus of learning-related plasticity. Learning of

sensory cues in flies, including odors and visual cues, is mediated by mushroom body

circuits (Aso et al., 2014; Vogt et al., 2014). The identification of the mushroom body as

an important learning center proceeded primarily from neuroanatomy, including their

connections to sensory projection neurons (Davis, 1993). Because operant conditioning

is not primarily a sensory-driven behavior, a sensory-first search for neural circuits is

unlikely to uncover the locus of plasticity that underlies operant learning. In fact, operant

conditioning in the fly is mushroom body independent (Wolf et al., 1998; Brembs, 2009),

while behavioral output circuits in the ventral nerve cord, such as motor neurons, have

been implicated instead (Booker and Quinn, 1981; Colomb and Brembs, 2016).

However, motor neurons themselves are unlikely to be the location of behavior/US

coincidence detection (Talay et al., 2017). Motor planning circuits in the fly central

complex, such as those responsible for navigation, are therefore an interesting potential

neuronal locus for operant plasticity.

Control of turn direction on the Y-Track is determined by a mix of innate motor

preferences and goal-directed search strategies. Innate handedness is strongly

influenced by the activity of PB-FB-No neurons (PFN) (Buchanan et al., 2015). While

the synaptic partners of PFN neurons implicated in innate handedness are not yet

mapped, the brain structures innervated are all heavily involved in orientation and

navigation in the fly (Giraldo et al., 2018; Shiozaki et al., 2020). Fly orientation circuits

show rapid plasticity and features of short-term memory (Fisher et al., 2019) and are

strongly responsive to visual stimuli (Seelig and Jayaraman, 2015). If fly orientation

circuits are necessary for Y-Track operant conditioning, the mix of plasticity and visual

responses would account for both behavioral plasticity and our finding that a strong

visual stimulus inhibits learning rather than promoting it (Figure 4).

In addition to innate preferences, flies display strong learned place preference

and goal directed search behaviors (Ofstad et al., 2011; Kim and Dickinson, 2017).

Development of a location preference is capable of overriding innate preferences

(Baggett et al., 2018) and foraging flies modify their innate locomotor preferences to

repeatedly visit remembered sites of food and search for nearby food sources (Kim and

Dickinson, 2017). The formation of spatial memories has been previously linked to

cAMP as a coincidence detector (Zars et al., 2000), consistent with our finding that

cAMP regulation is necessary for Y-Track conditioning (Figure 7). Our behavioral results

are congruent with either plasticity happening directly in orientation/innate preference

circuits, or in a foraging/place preference circuit. Further characterization of the neural

components of spatial memory in flies is necessary for these possibilities to be

distinguished.

ACKNOWLEDGEMENTS

The authors would like to thank Hazal Uzunkaya and the Brandeis MakerLab for 3D

printing assistance and would like to thank Junwei Yu for recommending the rest-

promoting Ellipsoid Body driver (VT058968).

FUNDING

This work was supported by National Institutes of Health Grants R01 MH067284 (to

LCG), F32 NS098624 (to TDW), R90 DA033463 (to JBL), the Radcliffe Institute for

Advanced Studies Fellowship (to RH), the Brandeis Provost Research Grant (to TDW),

and the M. R. Bauer Foundation Summer Undergraduate Research Fellowship (to

YYH).

REFERENCES

Angriman M, Caravale B, Novelli L, Ferri R, Bruni O (2015) Sleep in children with
neurodevelopmental disabilities. Neuropediatrics 46:199–210.

Aso Y et al. (2014) Mushroom body output neurons encode valence and guide memory-
based action selection in Drosophila. Elife 3:e04580.

Baggett V, Mishra A, Kehrer AL, Robinson AO, Shaw P, Zars T (2018) Place learning
overrides innate behaviors in Drosophila. Learn Mem 25:122–128.

Berry JA, Cervantes-Sandoval I, Chakraborty M, Davis RL (2015) Sleep Facilitates
Memory by Blocking Dopamine Neuron-Mediated Forgetting. Cell 161:1656–1667.

Booker R, Quinn WG (1981) Conditioning of leg position in normal and mutant
Drosophila. Proc Natl Acad Sci U S A 78:3940–3944.

Brembs B (2009) Mushroom Bodies Regulate Habit Formation in Drosophila. Curr Biol
19:1351–1355.

Brembs B, Plendl W (2008) Double Dissociation of PKC and AC Manipulations on
Operant and Classical Learning in Drosophila. Curr Biol 18:1168–1171.

Buchanan SM, Kain JS, de Bivort BL (2015) Neuronal control of locomotor handedness
in *Drosophila*. Proc Natl Acad Sci 112:6700–6705.

Buhry L, Azizi AH, Cheng S (2011) Reactivation, replay, and preplay: how it might all fit

together. Neural Plast 2011:203462.

Chaudhuri KR, Naidu Y (2008) Early Parkinson's disease and non-motor issues. J

Neurol 255 Suppl:33–38.

Chóliz M (2010) Experimental Analysis of the Game in Pathological Gamblers: Effect of

the Immediacy of the Reward in Slot Machines. J Gambl Stud 26:249–256.

Chouhan NS, Griffith LC, Haynes P, Sehgal A (2020) Availability of food determines the

need for sleep in memory consolidation. Nature.

Colomb J, Brembs B (2016) PKC in motorneurons underlies selflearning, a form of

motor learning in Drosophila. PeerJ 2016.

Davis RL (1993) Mushroom bodies and drosophila learning. Neuron 11:1–14.

Dawydow A, Gueta R, Ljaschenko D, Ullrich S, Hermann M, Ehmann N, Gao S, Fiala A,

Langenhan T, Nagel G, Kittel RJ (2014) Channelrhodopsin-2-XXL, a powerful

optogenetic tool for low-light applications. Proc Natl Acad Sci U S A 111:13972–

13977.

DeJianne D, McGuire TR, Pruzan-Hotchkiss A (1985) Conditioned suppression of

proboscis extension in Drosophila melanogaster. J Comp Psychol 99:74–80.

Dissel S, Angadi V, Kirszenblat L, Suzuki Y, Donlea J, Klose M, Koch Z, English D,

Winsky-Sommerer R, van Swinderen B, Shaw PJ (2015) Sleep restores behavioral

plasticity to Drosophila mutants. Curr Biol 25:1270–1281.

Donelson NC, Donelson N, Kim EZ, Slawson JB, Vecsey CG, Huber R, Griffith LC

(2012) High-resolution positional tracking for long-term analysis of Drosophila sleep

and locomotion using the "tracker" program. PLoS One 7:e37250.

Donlea JM, Ramanan N, Silverman N, Shaw PJ (2014) Genetic rescue of functional

senescence in synaptic and behavioral plasticity. Sleep 37:1427–1437.

Donlea JM, Thimgan MS, Suzuki Y, Gottschalk L, Shaw PJ (2011) Inducing sleep by

remote control facilitates memory consolidation in Drosophila. Science 332:1571–

1576.

Fanselow MS, Wassum KM (2015) The Origins and Organization of Vertebrate

Pavlovian Conditioning. Cold Spring Harb Perspect Biol 8:a021717.

Fisher YE, Lu J, D'Alessandro I, Wilson RI (2019) Sensorimotor experience remaps

visual input to a heading-direction network. Nature 576:121–125.

Flyer-Adams JG, Rivera-Rodriguez EJ, Yu J, Mardovin JD, Reed ML, Griffith LC (2020)
Regulation of olfactory associative memory by the circadian clock output signal
Pigment-dispersing factor (PDF). J Neurosci 40:JN-RM-0782-20.

Ganguly-Fitzgerald I, Donlea J, Shaw PJ, Tononi G, Shaw PJ (2006) Waking
Experience Affects Sleep Need in Drosophila. Science 313:1775–1781.

Gatzounis R, Schrooten MGS, Crombez G, Vlaeyen JWS (2012) Operant Learning
Theory in Pain and Chronic Pain Rehabilitation. Curr Pain Headache Rep 16:117–
126.

Gervasi N, Tchénio P, Preat T (2010) PKA Dynamics in a Drosophila Learning Center:
Coincidence Detection by Rutabaga Adenylyl Cyclase and Spatial Regulation by
Dunce Phosphodiesterase. Neuron 65:516–529.

Gilbert KS, Kark SM, Gehrman P, Bogdanova Y (2015) Sleep Disturbances, TBI and
PTSD: Implications for Treatment and Recovery. Clin Psychol Rev 40:195–212.

Giraldo YM, Leitch KJ, Ros IG, Warren TL, Weir PT, Dickinson MH (2018) Sun
Navigation Requires Compass Neurons in Drosophila. Curr Biol 28:2845-2852.e4.

Greenspan RJ, Dierick HA (2004) "Am not I a fly like thee?" From genes in fruit flies to
behavior in humans. Hum Mol Genet 13 Spec No:R267-73.

Hendricks JC, Finn SM, Panckeri KA, Chavkin J, Williams JA, Sehgal A, Pack AI (2000)
Rest in Drosophila is a sleep-like state. Neuron 25:129–138.

Hughson BN, Anreiter I, Jackson Chornenki NL, Murphy KR, Ja WW, Huber R,
Sokolowski MB (2018) The adult foraging assay (AFA) detects strain and food-
deprivation effects in feeding-related traits of Drosophila melanogaster. J Insect
Physiol 106:20–29.

Jang T, Lee KP (2015) The genetic basis for mating-induced sex differences in
starvation resistance in Drosophila melanogaster. J Insect Physiol 82:56–65.

Kamyshev NG, Kamysheva EA, Ivanova GO (1997) Autosomal Mutations In Drosophila
Which Reduce Operant Learning Ability. Neurosci Behav Physiol 27:254–257.

Keene AC, Duboué ER, McDonald DM, Dus M, Suh GSB, Waddell S, Blau J (2010)
Clock and cycle limit starvation-induced sleep loss in drosophila. Curr Biol
20:1209–1215.

Kessler RC, Berglund PA, Coulouvrat C, Hajak G, Roth T, Shahly V, Shillington AC,

Stephenson JJ, Walsh JK (2011) Insomnia and the performance of US workers:

results from the America insomnia survey. Sleep 34:1161–1171.

Kim IS, Dickinson MH (2017) Idiothetic Path Integration in the Fruit Fly Drosophila

melanogaster. Curr Biol 27:2227-2238.e3.

Kimble GA, Mann LI, Dufort RH (1955) Classical and instrumental eyelid conditioning. J

Exp Psychol 49:407–417.

Korff S, Harvey BH (2006) Animal Models of Obsessive-Compulsive Disorder: Rationale

to Understanding Psychobiology and Pharmacology. Psychiatr Clin North Am

29:371–390.

Krashes MJ, DasGupta S, Vreede A, White B, Armstrong JD, Waddell S (2009) A

neural circuit mechanism integrating motivational state with memory expression in

Drosophila. Cell 139:416–427.

Krashes MJ, Keene AC, Leung B, Armstrong JD, Waddell S (2007) Sequential use of

mushroom body neuron subsets during drosophila odor memory processing.

Neuron 53:103–115.

Levey DF, Gelernter J, Polimanti R, Zhou H, Cheng Z, Aslan M, Quaden R, Concato J,

Radhakrishnan K, Bryois J, Sullivan PF, Stein MB (2020) Reproducible Genetic

Risk Loci for Anxiety: Results From ~200,000 Participants in the Million Veteran

Program. Am J Psychiatry 177:223–232.

Lewis SA, Negelspach DC, Kaladchibachi S, Cowen SL, Fernandez F (2017)

Spontaneous alternation: A potential gateway to spatial working memory in

Drosophila. Neurobiol Learn Mem 142:230–235.

Li Q, DeBeaubien NA, Sokabe T, Montell C (2020) Temperature and Sweet Taste

Integration in Drosophila. Curr Biol 30:2051-2067.e5.

Liu C, Meng Z, Wiggin TD, Yu J, Reed ML, Guo F, Zhang Y, Rosbash M, Griffith LC

(2019) A Serotonin-Modulated Circuit Controls Sleep Architecture to Regulate

Cognitive Function Independent of Total Sleep in Drosophila. Curr Biol 29:3635-

3646.e5.

Liu C, Plaçais P-Y, Yamagata N, Pfeiffer BD, Aso Y, Friedrich AB, Siwanowicz I, Rubin

GM, Preat T, Tanimoto H (2012) A subset of dopamine neurons signals reward for

odour memory in Drosophila. Nature 488:512–516.

Liu S, Liu Q, Tabuchi M, Wu MN (2016) Sleep Drive Is Encoded by Neural Plastic
Changes in a Dedicated Circuit. Cell 165:1347–1360.

Maquet P (2001) The Role of Sleep in Learning and Memory. Science (80-) 294:1048–
1052.

Margulies C, Tully T, Dubnau J (2005) Deconstructing memory in Drosophila. Curr Biol
15:R700-13.

Martin JR, Ernst R, Heisenberg M (1999) Temporal pattern of locomotor activity in
Drosophila melanogaster. J Comp Physiol - A Sensory, Neural, Behav Physiol
184:73–84.

Mayford M, Siegelbaum SA, Kandel ER (2012) Synapses and memory storage. Cold
Spring Harb Perspect Biol 4:1–18.

Meissner GW, Grimm JB, Johnston RM, Sutcliffe B, Ng J, Jefferis GSXE, Cachero S,
Lavis LD, Malkesman O (2018) Optimization of fluorophores for chemical tagging
and immunohistochemistry of Drosophila neurons. PLoS One 13.

Menda G, Bar HY, Arthur BJ, Rivlin PK, Wyttenbach RA, Strawderman RL, Hoy RR
(2011) Classical conditioning through auditory stimuli in Drosophila: Methods and
models. J Exp Biol 214:2864–2870.

Mendes CS, Bartos I, Akay T, Márka S, Mann RS (2013) Quantification of gait
parameters in freely walking wild type and sensory deprived Drosophila
melanogaster. Elife 2013:231.

Morton AJ (2013) Circadian and sleep disorder in Huntington's disease. Exp Neurol
243:34–44.

Murphey RM (1967) Instrumental conditioning of the fruit fly, Drosophila melanogaster.
Anim Behav 15:153–161.

Murphy KR, Deshpande SA, Yurgel ME, Quinn JP, Weissbach JL, Keene AC, Dawson-
Scully K, Huber R, Tomchik SM, Ja WW (2016) Postprandial sleep mechanics in
Drosophila. Elife 5.

Nargeot R, Simmers J (2011) Neural mechanisms of operant conditioning and learning-
induced behavioral plasticity in Aplysia. Cell Mol Life Sci 68:803–816.

Nuwal N, Stock P, Hiemeyer J, Schmid B, Fiala A, Buchner E (2012) Avoidance of Heat
and Attraction to Optogenetically Induced Sugar Sensation as Operant Behavior in

Adult *Drosophila*. J Neurogenet 26:298–305.

Ofstad TA, Zuker CS, Reiser MB (2011) Visual place learning in Drosophila

melanogaster. Nature 474:204–209.

Ollendick TH, Vasey MW, King NJ (2001) Operant Conditioning Influences in Childhood

Anxiety. In: The Developmental Psychopathology of Anxiety, pp 231–252. Oxford

University Press.

Pfeiffer BD, Ngo TTB, Hibbard KL, Murphy C, Jenett A, Truman JW, Rubin GM (2010)

Refinement of tools for targeted gene expression in Drosophila. Genetics 186:735–

755.

Platt SA, Holliday M, Drudge OW (1980) Discrimination learning of an instrumental

response in individual Drosophila melanogaster. J Exp Psychol Anim Behav

Process 6:301–311.

Porter VR, Buxton WG, Avidan AY (2015) Sleep, Cognition and Dementia. Curr

Psychiatry Rep 17:97.

Quinn WG, Harris WA, Benzer S (1974) Conditioned behavior in Drosophila

melanogaster. Proc Natl Acad Sci U S A 71:708–712.

Rickard JF, Body S, Zhang Z, Bradshaw CM, Szabadi E (2009) Effect Of Reinforcer

Magnitude On Performance Maintained By Progressive-Ratio Schedules. J Exp

Anal Behav 91:75–87.

Ro J, Harvanek ZM, Pletcher SD (2014) FLIC: High-Throughput, Continuous Analysis of

Feeding Behaviors in Drosophila Roman G, ed. PLoS One 9:e101107.

Robbins TW, Cools R (2014) Cognitive deficits in Parkinson's disease: a cognitive

neuroscience perspective. Mov Disord 29:597–607.

Schnaitmann C, Vogt K, Triphan T, Tanimoto H (2010) Appetitive and aversive visual

learning in freely moving Drosophila. Front Behav Neurosci 4:10.

Seelig JD, Jayaraman V (2015) Neural dynamics for landmark orientation and angular

path integration. Nature 521:186–191.

Seugnet L, Suzuki Y, Vine L, Gottschalk L, Shaw PJ (2008) D1 receptor activation in the

mushroom bodies rescues sleep-loss-induced learning impairments in Drosophila.

Curr Biol 18:1110–1117.

Shaw PJ, Cirelli C, Greenspan RJ, Tononi G (2000) Correlates of sleep and waking in

Drosophila melanogaster. Science 287:1834–1837.

Shekleton JA, Flynn-Evans EE, Miller B, Epstein LJ, Kirsch D, Brogna LA, Burke LM,

Bremer E, Murray JM, Gehrman P, Lockley SW, Rajaratnam SMW (2014)

Neurobehavioral performance impairment in insomnia: relationships with self-

reported sleep and daytime functioning. Sleep 37:107–116.

Shiozaki HM, Ohta K, Kazama H (2020) A Multi-regional Network Encoding Heading

and Steering Maneuvers in Drosophila. Neuron 106:126-141.e5.

Simon JC, Dickinson MH (2010) A New Chamber for Studying the Behavior of

Drosophila Hashimoto K, ed. PLoS One 5:e8793.

Skinner BF (1948) Superstition in the pigeon. J Exp Psychol 38:168–172.

Smit DJA, Cath D, Zilhão NR, Ip HF, Denys D, den Braber A, de Geus EJC, Verweij

KJH, Hottenga JJ, Boomsma DI (2020) Genetic meta-analysis of obsessive–

compulsive disorder and self-report compulsive symptoms. Am J Med Genet Part B

Neuropsychiatr Genet 183:208–216.

Smith DJ et al. (2016) Genome-wide analysis of over 106 000 individuals identifies 9

neuroticism-associated loci. Mol Psychiatry 21:749–757.

Stowers JR, Hofbauer M, Bastien R, Griessner J, Higgins P, Farooqui S, Fischer RM,

Nowikovsky K, Haubensak W, Couzin ID, Tessmar-Raible K, Straw AD (2017)

Virtual reality for freely moving animals. Nat Methods 14:995–1002.

Susswein AJ, Schwarz M, Feldman E (1986) Learned changes of feeding behavior in

Aplysia in response to edible and inedible foods. J Neurosci 6:1513–1527.

Tainton-Heap LAL, Kirszenblat LC, Notaras ET, Grabowska MJ, Jeans R, Feng K,

Shaw PJ, van Swinderen B (2020) A Paradoxical Kind of Sleep in Drosophila

melanogaster. Curr Biol.

Talay M, Richman EB, Snell NJ, Hartmann GG, Fisher JD, Sorkaç A, Santoyo JF,

Chou-Freed C, Nair N, Johnson M, Szymanski JR, Barnea G (2017) Transsynaptic

Mapping of Second-Order Taste Neurons in Flies by trans-Tango. Neuron 96:783-

795.e4.

Tempel BL, Bonini N, Dawson DR, Quinn WG (1983) Reward learning in normal and

mutant Drosophila. Proc Natl Acad Sci U S A 80:1482–1486.

Vogt K, Schnaitmann C, Dylla K V, Knapek S, Aso Y, Rubin GM, Tanimoto H (2014)

Shared mushroom body circuits underlie visual and olfactory memories in

Drosophila. Elife 3:e02395.

Wagh DA, Rasse TM, Asan E, Hofbauer A, Schwenkert I, Dürrbeck H, Buchner S,

Dabauvalle MC, Schmidt M, Qin G, Wichmann C, Kittel R, Sigrist SJ, Buchner E

(2006) Bruchpilot, a protein with homology to ELKS/CAST, is required for structural

integrity and function of synaptic active zones in Drosophila. Neuron 49:833–844.

Warren TL, Giraldo YM, Dickinson MH (2019) Celestial navigation in Drosophila. J Exp

Biol 222.

Wiggin TD, Goodwin PR, Donelson NC, Liu C, Trinh K, Sanyal S, Griffith LC (2020)

Covert sleep-related biological processes are revealed by probabilistic analysis in

Drosophila. Proc Natl Acad Sci U S A 117:10024–10034.

Wolf R, Heisenberg M (1991) Basic organization of operant behavior as revealed in

Drosophila flight orientation. J Comp Physiol A 169:699–705.

Wolf R, Wittig T, Liu L, Wustmann G, Eyding D, Heisenberg M (1998) Drosophila

mushroom bodies are dispensable for visual, tactile, and motor learning. Learn

Mem 5:166–178.

Wustmann G, Rein K, Wolf R, Heisenberg M (1996) A new paradigm for operant

conditioning of Drosophila melanogaster. J Comp Physiol A 179:429–436.

Xu CS et al. (2020) A Connectome of the Adult *Drosophila* Central Brain.

bioRxiv:2020.01.21.911859.

Zamarian L, Högl B, Delazer M, Hingerl K, Gabelia D, Mitterling T, Brandauer E,

Frauscher B (2015) Subjective deficits of attention, cognition and depression in

patients with narcolepsy. Sleep Med 16:45–51.

Zars T, Wolf R, Davis R, Heisenberg M (2000) Tissue-specific expression of a type I

adenylyl cyclase rescues the rutabaga mutant memory defect: In search of the

engram. Learn Mem 7:18–31.

Zhang F, Wang LP, Boyden ES, Deisseroth K (2006) Channelrhodopsin-2 and optical

control of excitable cells. Nat Methods 3:785–792.

Figure 1 – Flies That Rest Learn the Operant Contingency in a Novel Sucrose Seeking Task (A) Diagram of operant conditioning paradigm. Sucrose is presented at the end of the track in the reinforced direction. A trial is completed after the fly crosses the choice threshold leaving the center of the track. Each training block is 20 trials, reward presentation begins in training block #1. Sucrose is made available for 10 seconds following the fly crossing the choice threshold. (B) Rendering of the 3D model of the novel apparatus used for 3D printing (*Left*) and a photo of a fully assembled apparatus (*Right*). (C) Field of view of overhead camera in the 3D render (*Left*) and a video frame of a fly navigating the Y-Track (*Right*). Red arrowhead indicates the position of the fly. Scale bar: 5 mm. (D) Probability Distribution Function (P.D.F.) histogram of movement (in pixels) per frame (approx. 50ms) for live flies (*Left*) and dead flies (*Right*). One pixel of movement is approximately 90 μm. (E) Histogram of locomotor pause durations from 12 hour recordings of locomotor behavior ($n = 2$ flies, 4127

episodes). Dashed line indicates the 1 minute threshold to distinguish short pauses from

rest episodes. (F) Schematic of hallway geometry with representative fly for scale (*Top*),

Fraction of flies with each rest phenotype (*Bottom*). (G) Mean turn direction Preference

Index (PI) for each training block. Points plotted in color are flies with early rest

(drowsy), points plotted in grey are flies with low rest (restless). (H) Probability

Distribution Function (P.D.F.) histograms of change in turn preference index (ΔPI)

between training block #1 and #7. Drowsy flies are plotted in red, restless flies are

plotted in grey. (I) Mean ΔPI of low rest and early rest flies. Error bars are standard error

of the mean, *n.s.*: groups are not significantly greater than zero, * groups are

significantly greater than zero.

Figure 2 – Y-Track Geometry Significantly Affects Thigmotaxis and Spontaneous Alternation (A-B) Position heatmaps of flies in the square wall (A) and curved floor (B) Y-Mazes. Insets (*right*) show the heatmaps of trajectories through the center of the maze for flies walking from the left zone to the upper zone vs. the lower zone. Scale bar: 5 mm. Color shows relative occupancy of each pixel from low to high. (C) Heatmap of left turn Preference Index (PI) from a population of *in silico* flies. Each heatmap position is the mean of 1000 simulated trials of a fly with a range of left-turn bias and alternation bias. Dashed lines indicate zero bias on an axis. Solid lines are smoothed contours of constant PI. (D,E) Mean thigmotaxis (D) and spontaneous alternation (E) for WT flies in square wall and curved wall experiments. (D) Thigmotaxis is defined for each fly as the median distance from the wall of walking locations. The dashed line indicates the median distance from the wall if walking locations were distributed uniformly across the hallway. (E) Spontaneous alternation is measured in the final training block. The dashed line indicates the random rate of alternation. Error bars are standard error of the mean. Groups with the same letter are not significantly different from one another.

Figure 3 – Learning to Turn Toward Sucrose is Independent of Track Geometry

but Dependent upon an Informative Operant Contingency (A) Schematic of hallway

geometry with representative fly for scale (*Top*), Fraction of flies with each rest

phenotype (*Bottom*). (B, F) Mean turn direction Preference Index (PI) for each training

block for trained (B) and yoked control (F) flies in the curved floor Y-Maze. Points

plotted in color are flies with early rest (drowsy), points plotted in grey are flies with low

rest (restless). (C,G) Probability Distribution Function (P.D.F.) histograms of change in

turn preference index (ΔPI) between training block #1 and #7 for trained (C) and yoked

control (G) flies. Drowsy flies are plotted in color (*cyan*: trained, *green*: yoked), restless

flies are plotted in grey. (D,H) Mean ΔPI of low rest (restless) and early rest (drowsy)

flies trained in curved floor Y-Maze (D), and yoked control flies (H). Error bars are

standard error of the mean, *n.s.*: groups are not significantly greater than zero, * groups

are significantly greater than zero.

Figure 4 – A World Orientation Cue does not Facilitate Learning the Operant Task.

(A) Position heatmaps of flies trained with an LED spatial orientation cue. Scale bar is 5 mm. Color shows relative occupancy of each pixel from low to high. Small inset indicates the position of the green light cue. (B) Percentage flies with each rest phenotype. (C) PI by training block for early rest (magenta) and low rest (grey) flies. (D) Mean ΔPI of early rest and low rest flies. Error bars are standard error of the mean, *n.s.*: groups are not significantly greater than zero, * groups are significantly greater than zero.

Figure 5 – Sucrose Promotes Rest and is the Learning-Relevant US in the Y-Track. (A) Mean time per trial WT flies spend adjacent to sucrose in the square wall and curved floor experiments. Time per trial is capped at 10 seconds, because sucrose is automatically removed 10 seconds after it is made available. Groups with the same letter are not significantly different from one another. (B) Scatter plot of change in turn preference index (ΔPI) between training block #1 and #7 by time adjacent to sucrose in training block #1 for flies in the square wall experiment (red dots) and curved floor experiment (cyan dots). (C-F) Results of Y-Maze training with no sucrose available. (C) Position heatmaps of trained flies. Scale bar is 5 mm. Color shows relative occupancy of each pixel from low to high. (D) Percentage flies with each rest phenotype. PI by training block (E) and mean ΔPI (F) for low rest flies. Only 2 flies had early rest, which is insufficient to calculate a standard error, so these flies are not plotted. Error bars are standard error of the mean, *n.s.*: groups are not significantly greater than zero.

Figure 6 – Optogenetically-Induced Rest is Not Sufficient to Enhance Learning.

(A) Confocal maximum intensity projections of fly brains expressing 104y-Gal4;UAS-GFP (*left*) and VT058968-Gal4;UAS-GFP (*right*) stained for GFP (*green*) and Bruchpilot (nc82, *magenta*). Scale bar is 20 µm. White arrowheads indicate the sleep-promoting neuropil for each Gal4 driver: the dorsal Fan Shaped Body for 104y-Gal4 and the Ellipsoid Body for VT058968-Gal4. (B) Block diagram of operant conditioning paradigm, with the time of optogenetic stimulus indicated in cyan. The track diagram indicates the locations of each blue LED in the track, these positions are equidistant from the center of the track and illuminate the entire Y-track. (C) Peri-stimulus rest plots for WT (*left*), 104y (*center*), and VT058968 (*right*). Blue shaded area shows when the blue LED is on. The plots show 20 minutes of behavior following light onset - this time is not linked to trial times, which vary between animals, and does not encompass the entire duration of the experiment. (D) Percent of each rest phenotype present in each experiment group.

(E-G) Results of training WT (E), 104y (F), and VT058968 (G) flies. Early and low rest

flies for each genotype were pooled from the ATR and vehicle groups. Change in PI

(ΔPI) is plotted for each genotype; colorful points are flies with early rest (*purple:* WT,

magenta: 104y, *green*: VT058968). Error bars are standard error of the mean, *n.s.*:

groups are not significantly greater than zero, * groups are significantly greater than

zero.

Figure 7 – Y-Track Operant Learning is Dependent on cAMP Regulation. (A,D)
Position heatmaps of trained WT (A) and *dnc¹* (D) flies. Scale bar is 5 mm. Color shows
relative occupancy of each pixel from low to high. (B,E) Fraction of WT (B) and *dnc¹* (E)
flies with each rest phenotype. (C,F) Mean ΔPI of low rest (restless) and of early rest
(drowsy) WT (C) and *dnc¹* (F) flies. The mean ΔPI of drowsy and restless WT files are
both greater than zero (One-sample bootstrap test; *p* = 0.005 and *p* = 0.019,
respectively). The mean ΔPI of drowsy WT flies is greater than the mean ΔPI of restless
WT files (Two-sample bootstrap test; p = 0.035). Error bars are standard error of the
mean, *n.s.*: groups are not significantly greater than zero, * groups are significantly
greater than zero or different from one another.