Different combinations of insect Na,K-ATPase α- and β-subunit paralogs enable fine tuning of toxin resistance and enzyme kinetics

Marlena Herbertz[1#], Safaa Dalla[1,2#], Vera Wagschal[1], Rohin Turjalei[1], Marlies Heiser[1], Susanne Dobler[1*]

[1]Institute of Zoology, Molecular Evolutionary Biology, Universität Hamburg, Hamburg, Germany

[2] present address: Tishreen University, Latakia, Syria

[#] Co-first author

[*]Corresponding author: Susanne Dobler

E-mail: susanne.dobler@uni-hamburg.de

Running title: Kinetics and toxin resistance of Na,K-ATPase paralogs

Keywords: Na+/K+-ATPase, cardiac glycosides, toxicity, resistance, molecular evolution, baculovirus expression system

Abstract

Background

Cardiac glycosides are known to fatally inhibit the Na,K-ATPase throughout the animal kingdom. Several animals, however, evolved target-site insensitivity by substitutions in the otherwise highly conserved cardiac glycoside binding pocket located on the Na,K-ATPase α-subunit. The minimal functional enzyme consist of an α- and a β-subunit, the latter considered mainly as a chaperone responsible for correct folding and membrane integration. We here analyze resistance to cardiac glycosides and kinetic properties of different Na,K-ATPase α/β-combinations of the large milkweed bug, *Oncopeltus fasciatus*. These insects have adapted to high concentrations of cardiac glycosides in their food plants via several rounds of Na,K-ATPase gene duplications followed by differential resistance conferring substitutions and subfunctionalization of the enzymes.

Results

To investigate their characteristics we expressed nine combinations of *O. fasciatus* Na,K-ATPase α/β-sunbunits (three each) in *Sf*9 cells and tested them with two structurally distinct cardiac glycosides, calotropin, a host plant compound, and ouabain, a commonly used toxin. Differences in the number and identity of amino acid substitutions in the cardiac glycoside binding site resulted in large differences in activity and toxin resistance of the three α-subunits. The enzymes' kinetics were also influenced by the β-subunits leading to increased activities (αCβ3) or altered resistances. The host plant toxin calotropin proved to be a much more potent inhibitor than ouabain for the

45 phylogenetically oldest αC based enzymes. This effect was compensated for in the αB and

46 αA based enzymes with αAβ1 having higher resistance against calotropin than against

47 ouabain.

Conclusion

49 The originally higher inhibitory potency of the host compound calotropin supports a

50 coevolutionary escalation of plant defenses and herbivore tolerance mechanisms. For the

51 bugs the possession of multiple paralogs improved adaptation to plant toxins in a stepwise

52 manner and mitigates pleiotropic effects by a compromise between ion pumping activity

53 and resistance.

Background

The transmembrane Na,K-ATPase, is ubiquitously expressed in animal cells and is essential

for maintaining membrane potentials, controlling cell volume and the overall functionality

of cells (1). The functional enzyme consists of two subunits, the catalytic α-subunit and an

associated β-subunit serving as chaperone for correct folding and membrane integration

(2). In vertebrates several paralogs are present for both genes and the enzyme's

characteristics differ depending on the specific combination of α and β-subunits (3,4). An

additional modulatory and tissue specific γ-subunit is present in vertebrate tissues (5), but

has not been detected in invertebrates. Much less is known about the importance of

different α/β-subunit combinations in invertebrates (6). Only in *Drosophila* the tissue

distribution and functional roles of some of these combinations have been explored to

some extent (7–9). We here explore the functional role of different Na,K-ATPase

complexes in the large milkweed bug, *Oncopeltus fasciatus*. In this species, the Na,K-

ATPase α1 subunit has undergone three rounds of duplications (10,11) and displays high

insensitivity against cardiac glycosides (12,13), which are universal inhibitors of the Na,K-

ATPase (1).

Inhibition of the Na,K-ATPase by cardiac glycosides, such as the cardenolides ouabain and

calotropin (Fig. 1), depends on a specific binding site located on the extracellular side of

the catalytic α-subunit. Upon binding of cardiac glycosides the enzyme is arrested in the

phosphorylated state and ion pumping is disrupted (1). Inhibition of the Na,K-ATPase is

harmful and can be lethal to almost all animals. There are, however, a few notable

exceptions to this: various mammals, frogs, and reptiles prey unharmed on cardiac

glycoside-containing toads and a number of phytophagous insects feed - often exclusively

- on cardiac glycoside-containing host plants (reviewed in 14,15). Most of these species

avoid the toxic effect of cardiac glycosides and exhibit target site insensitivity, by

drastically lowering binding affinity of cardiac glycosides to their Na,K-ATPase α-subunit

(16,17). Strikingly, identical amino acid substitutions at homologous positions

convergently evolved in various, distantly related insect groups and even resemble the

substitutions observed in cardiac glycoside-resistant vertebrates (11,17–19).

While insect species usually express a single gene copy of the Na,K-ATPase α-subunit

(ATPα1) throughout their body, this gene underwent one to three rounds of gene

duplications in several cardiac glycoside-exposed species (10,11,16,20). These duplicated

ATPα1 copies differ in the number and identity of presumably resistance-causing amino

acid substitutions and show tissue specific expression patterns (10,11,20,21).

The large milkweed bug *Oncopeltus fasciatus* (Hemiptera, Lygaeidae) possesses four

ATPα1 paralogs (A, B, C, D) that differ in the number and identity of amino acid

substitutions (10,11). We previously introduced substitutions observed in these paralogs

at four usually highly conserved amino acid sites (Q111, N122, F786, and T797) known to

be important for high affinity to cardiac glycosides (22–25) into the sensitive ATPα1 gene

of *Drosophila melanogaster* (26,27). The N122H (as observed in ATPα1D of *O. fasciatus*),

N122H+T797S (as observed in ATPα1C), Q111T+N122H+F786N (as in ATPα1B) or

Q111T+N122H+F786N+T797A (as in ATPα1A) substitutions yielded proteins with stepwise

increasing resistance to the cardenolide ouabain. At the same time, enzyme activity

decreased with increasing numbers of substitutions (26).

Additional untested substitutions are present in the *O. fasciatus* ATPα1 paralogs and these

might influence the characteristics of the enzymes, potentially further increasing

resistance but possibly also serving as compensatory mutations mitigating deleterious

intra-molecular epistatic effects of previously characterized mutations (28–30). In

addition, it has been shown that the β-subunit of the Na,K-ATPase not only acts as a

chaperone, guiding the heterodimeric enzyme to the cell membrane, but also influences

the behavior of the Na,K-ATPase including its sensitivity to ouabain (2,4,27).

Although ouabain is a commonly used cardiac glycoside in studies of the Na,K-ATPase, it

does not occur in host plants of *O. fasciatus*. The *Asclepias* species (Apocynaceae) that the

bugs feed on typically contain different types of cardenolides characterized by a *trans-*

trans-cis conformation of the steroidal skeleton and a single sugar moiety attached by a

cyclic bridge as in the abundant calotropin (15) (Fig. 1). The most biologically relevant test

of the bugs' resistance against cardiac glycosides should thus involve a plant-derived

compound like calotropin. To address these issues, we identified and heterologously

expressed nine possible *O. fasciatus* Na,K-ATPase α/β combinations and characterized

their activities and sensitivities to ouabain and calotropin.

Since activity and resistance of the Na,K-ATPase is determined by the number and identity

of amino acid substitutions in the cardenolide binding site located in the α-subunit, (1) we

hypothesized that α-subunit paralogs – associated with different β-subunits – differ

significantly in their activities and insensitivities to the two cardenolides we tested. (2) We

further hypothesized that the different β-subunits modulate the Na,K-ATPase α-subunit

paralog behavior influencing their response to the cardenolides. Additionally, although

ouabain and calotropin both seem to be strong inhibitors of the Na,K-ATPase, (3) we

expected the two cardenolides to differ significantly in their inhibitory potency against the

Na,K-ATPase of *O. fasciatus* due to differences in structure and steric conformation. We

hypothesized that coevolutionary interaction should lead to higher resistance against the

host plant compound calotropin.

Results

Bioinformatic analysis

Four potential β-subunits were identified from the *O. fasciatus* transcriptome and

according to their similarity to published sequences named β1, β2, β3, and βx (Genbank

AccNo xxx-xxx). While β1-3 conform to the standard length of β-subunits of around 900 bp

and are recovered in Blast searches as homologues of similarly numbered β-subunits of

other insects, βx is much shorter and missing most of the conserved N-terminal and half of

the transmembrane region and does not have obvious homologues in other insects (Fig.

S1A). This β-subunit was therefore not further considered here. The 5' end of α1B and α1C

could successfully be recovered by RACE protocols (Genbank AccNo xxx, xxx). The only

minimally expressed α1D (10) was not further investigated here.

Expression of α/β-subunit constructs in Sf9 cells

All nine α/β-subunit constructs (αAβ1, αAβ2, αAβ3, αBβ1, αBβ2, αAβ3, αCβ1, αCβ2, and

αCβ3) were successfully expressed in *Sf*9 cells and the corresponding proteins harvested.

The western blot shows that similar concentrations were loaded onto the gel (12G10 anti-

α-tubulin as positive control), all α-subunits were expressed at similar levels and the

deglycosylated β-subunits can be easily discriminated by their specific patterns (Fig. 2).

Na,K-ATPase activity of different α/β-complexes

The activities of different α/β-subunit combinations of the Na,K-ATPase were tested *in*

vitro to determine whether the amino acid substitutions in the cardenolide binding site or

the associated β-subunit influence the catalytic activity of the enzyme. All enzymes

showed catalytic activity, but the activity of the α/β-complexes differed significantly from

one another (Fig. 3). The α-subunits A and B with their associated β-subunits (1-3) showed

significantly lower activity than αC combined with β1, 2 or 3. The activities of αA and αB

did not differ significantly from each other regardless of the β-subunit, except for αAβ2.

This combination was significantly less active than αB associated with the three β-subunits

(Fig. 3). By far the highest activity was measured for αC in association with β3 that had an

average turnover of 32.7 nmol Pi per mg Protein per minute and differed significantly

from all other enzymes.

Influence of α/β-subunit combination on ouabain sensitivity

In vitro tests with the cardenolide ouabain revealed large differences in sensitivity of the

Na,K-ATPases depending on the number of amino acid substitutions in the cardenolide

binding site and was modulated by the identity of the β-subunit (Fig. 4). Yet, when

comparing the IC_{50} values only the one of αC associated with β1 was significantly lower

than those for the other two αC complexes (ANOVA, $F_{2,8}$=8.035, p=0.02, Tukey's HSD,

p=0.035 and p=0.027, for αC β1 vs β2 and β1 vs β3, respectively; Fig. 4c, Table 1) whereas

the IC_{50} values of the three αA or αB complexes did not differ significantly.

Although analyzing differences in IC_{50} is a standard approach, the IC_{50} values only reflect

one point of the dose response curve. We therefore aimed to get a better insight into the

overall responses of the Na,K-ATPases to increasing ouabain concentrations by analyzing

the differences between the full dose response curves of enzymes with the same α-

subunit as well as enzymes with the same β-subunit (Table 2).

We found that α-subunits associated with different β-subunits behaved differently when

subjected to ouabain. The enzyme αAβ3 was significantly more sensitive to the

cardenolide than the other two αA complexes (Table 2, Fig. 4a), and αBβ1 was inhibited

more strongly by ouabain than αBβ2 (Table 2, Fig. 4b). Nevertheless, the reaction of

enzymes with identical α-subunits was more uniform than the one of Na,K-ATPases with

identical β but different α-subunits. Most of the latter enzymes differed significantly in

their reactions, reflecting their varying insensitivity levels against ouabain (Fig. 4 a-c, Table

2).

Influence of α/β-subunit association on calotropin sensitivity

Testing the inhibitory effect of the host plant cardenolide calotropin on the recombinant

Na,K-ATPases resulted in total inhibition of the most calotropin-sensitive αC/β-

combinations at the highest calotropin concentration. Furthermore, the three αB based

enzymes turned out to be most resistant against calotropin. As for the inhibition by

ouabain only the IC_{50} values of αC associated with the three different β-subunits differed

significantly from each other, αC associated with β3 was significantly more insensitive to

calotropin than αC associated with β1 and β2 (ANOVA, $F_{2,8}$=18.554, p=0.003, Tukey's HSD,

p=0.002 and p=0.027, for αCβ3 versus αCβ1 and αCβ3 versus αCβ2, respectively; Fig. 4c).

When comparing the full dose response curves of the constructs (Table 2) neither the

three αA combinations differed significantly from one another nor the αB or αC based

enzymes, indicating a similar response to calotropin exposition independent of the

associated β-subunit (Fig. 4 a-c). In contrast, the inhibition by calotropin of the three α-

subunits combined with the same β-subunits was significantly different (from high to low

inhibition: αC> αA> αB; Fig. 4a-c). The curves of αAβ1 and αBβ1 did not differ significantly

from one another, since αAβ1 tends to be the most resistant combination of the three αA

enzymes and αBβ1 the most sensitive of the three αB enzymes (Fig. 4a & b).

Comparison of both cardenolides

Calotropin inhibited especially the three αC enzymes to a much higher extent than

ouabain as judged by a comparison of IC_{50} values (ANOVA, $F_{5,17}$=57.25882, p<0.001,

Tukey's HSD, p<0.001 for all αC based enzymes; Fig. 4c, Table 1). αCβ1 was inhibited 18-

fold, αCβ2 20-fold, and αCβ3 14-fold more strongly by calotropin than by ouabain. When

only comparing IC_{50} values αB associated with β3 was also significantly more strongly

inhibited by calotropin than ouabain, resulting in an approximately 24-fold higher IC_{50}

(ANOVA, $F_{5,17}$=7.54487, p=0.00205, Tukey's HSD, p=0.00357; Fig. 4b, Table 1) while the

IC_{50} values of the two cardenolides did not differ significantly for αBβ1 and αBβ2 (Tukey's

HSD, p=0.70595 and p=0.51146, respectively; Fig. 4b, Table 1). Comparison of the two

cardenolides resulted in no significant differences between IC_{50} values of αA associated

with any of the three β-subunits (ANOVA, $F_{5,16}$=1.85583, p=0.18241; Fig. 4b, Table 1).

However, when comparing the full dose response curves of each enzyme inhibited by

either of the cardenolides, we found that αAβ1 was significantly more insensitive against

calotropin than against ouabain, reflected by a flat descending curve at the top (Fig. 4a,

Table 3). For the αC-based enzymes the stronger inhibition by calotropin remained

evident, while for enzymes with αB no difference in inhibitory potency between the two

cardenolides was found (Table 3).

Table 1: Mean IC_{50} values and Hill slopes of different Na,K-ATPase α/β-complexes (3 biological replicates with

standard deviation (± s.d.)) obtained from dose response curves upon inhibition with ouabain and calotropin

(Origin-Pro 9.1G).

α/β-combination	IC_{50} values [M] Ouabain (± s.d.)	Hill slope (± s.d.)	IC_{50} values [M] Calotropin (± s.d.)	Hill slope (± s.d.)
αAβ1	4.148E-04 (± 6.541E-04)	-0.221 (± 0.047)	1.211E-04 (± 4.443E-05)	-0.271 (± 0.03)
αAβ2	3.471E-04 (± 3.598E-04)	-0.234 (± 0.013)	1.449E-05 (± 5.994E-06)	-0.269 (± 0.076)
αAβ3	2.607E-05 (± 2.767E-05)	-0.260 (± 0.044)	6.959E-05 (± 7.54E-05)	-0.27 (± 0.036)
αBβ1	2.089E-03 (± 2.515E-03)	-0.213 (± 0.039)	1.108E-04 (± 3.423E-05)	-0.293 (± 0.044)
αBβ2	2.777E-03 (± 9.676E-04)	-0.462 (± 0.055)	3.281E-04 (± 2.119E-04)	-0.398 (± 0.066)

αBβ3	7.09E-03 (± 3.119E-03)	-0.261 (± 0.045)	2.944E-04 (± 8.195E-05)	-0.254 (± 0.071)
αCβ1	4.274E-04 (± 8.104E-05)	-0.869 (± 0.071)	2.402E-05 (± 4.636E-06)	-0.88 (± 0.029)
αCβ2	7.369E-04 (± 1.001E-04)	-0.869 (± 0.093)	3.699E-05 (± 5.517E-06)	-1.289 (± 0.198)
αCβ3	7.579E-04 (± 1.476E-04)	-0.988 (± 0.35)	5.594E-05 (± 8.55E-06)	-0.912 (± 0.098)

221   Table 2: Comparison between the dose response curves of different α/β-complexes inhibited by ouabain

222   and calotropin analyzed with Origin-Pro 9.1G. Significant differences are highlighted in bold.

Data comparison α/β-combinations	F-statistic Ouabain	F-statistic Calotropin
αAβ1 − αAβ2	$F_{2,8}$= 0.074, p= 0.929	$F_{2,8}$= 1.319, p= 0.320
αAβ1 − αAβ3	**$F_{2,8}$= 5.218, p= 0.035**	$F_{2,8}$= 0.728, p= 0.512
αAβ2 − αAβ3	**$F_{2,8}$= 6.092, p= 0.025**	$F_{2,8}$= 0.079, p= 0.925
αBβ1 − αBβ2	**$F_{2,8}$= 10.981, p= 0.005**	$F_{2,8}$= 2.607, p= 0.134
αBβ1 − αBβ3	$F_{2,8}$= 3.781, p= 0.070	$F_{2,8}$= 0.724, p= 0.514
αBβ2 − αBβ3	$F_{2,8}$= 0.455, p= 0.650	$F_{2,8}$= 0.655, p= 0.545
αCβ1 − αCβ2	$F_{2,8}$= 0.745, p= 0.505	$F_{2,8}$= 1.887, p= 0.213

$\alpha C\beta 1 - \alpha C\beta 3$	$F_{2,8}$= 1.535, p= 0.273	$F_{2,8}$= 3.402, p= 0.085
$\alpha C\beta 2 - \alpha C\beta 3$	$F_{2,8}$= 1.080, p= 0.384	$F_{2,8}$= 2.217, p= 0.171
$\alpha A\beta 1 - \alpha B\beta 1$	$F_{2,8}$= 1.039, p= 0.397	$F_{2,8}$= 0.554, p= 0.595
$\alpha A\beta 1 - \alpha C\beta 1$	**$F_{2,8}$= 22.729, p< 0.001**	**$F_{2,8}$= 8.724, p= 0.010**
$\alpha B\beta 1 - \alpha C\beta 1$	**$F_{2,8}$= 11.975, p= 0.004**	**$F_{2,8}$= 8.795, p= 0.010**
$\alpha A\beta 2 - \alpha B\beta 2$	**$F_{2,8}$= 57.311, p< 0.001**	**$F_{2,8}$= 9.763, p= 0.007**
$\alpha A\beta 2 - \alpha C\beta 2$	**$F_{2,8}$= 4.835, p= 0.042**	**$F_{2,8}$= 6.422, p= 0.022**
$\alpha B\beta 2 - \alpha C\beta 2$	$F_{2,8}$= 2.353, p= 0.157	**$F_{2,8}$= 13.589, p= 0.003**
$\alpha A\beta 3 - \alpha B\beta 3$	**$F_{2,8}$= 32.590, p< 0.001**	**$F_{2,8}$= 37.553, p< 0.001**
$\alpha A\beta 3 - \alpha C\beta 3$	**$F_{2,8}$= 33.295, p< 0.001**	**$F_{2,8}$= 19.271, p< 0.001**
$\alpha B\beta 3 - \alpha C\beta 3$	$F_{2,8}$= 1.614, p= 0.258	**$F_{2,8}$= 8.239, p= 0.011**

224   Table 3: Comparison between dose response curves for individual α/β-complexes inhibited by ouabain

225   versus calotropin analyzed with Origin-Pro 9.1G. Significant differences are highlighted in bold.

Data comparison α/β-combinations	F-statistic calotropin vs. ouabain	more potent inhibitor
$\alpha A\beta 1$	**$F_{2,8}$= 11.752, p= 0.004**	ouabain
$\alpha A\beta 2$	$F_{2,8}$= 2.421, p= 0.151	equal
$\alpha A\beta 3$	$F_{2,8}$= 0.589, p= 0.577	equal
$\alpha B\beta 1$	$F_{2,8}$= 0.343, p= 0.720	equal
$\alpha B\beta 2$	$F_{2,8}$= 4.257, p= 0.055	equal

αBβ3	$F_{2,8}$= 3.054, p= 0.103	equal
αCβ1	**$F_{2,8}$= 35.226, p< 0.001**	calotropin
αCβ2	**$F_{2,8}$= 100.356, p< 0.001**	calotropin
αCβ3	**$F_{2,8}$= 74.665, p< 0.001**	calotropin

Discussion

Here we have identified the full length coding sequences of three Na,K-ATPase α- and

three full length β-subunits of *O. fasciatus* and functionally characterized all nine possible

recombinant α/β enzyme complexes. These nine functional enzymes strongly differed in

their activity levels. We found that enzyme complexes with αC had a significantly higher

activity compared to the other constructs. This perfectly fits with the fact that αC is mainly

expressed in the nervous tissue of *O. fasciatus* (11,21) where the Na,K-ATPase plays a

crucial role in maintaining electric potentials that guarantee the excitability of neurons.

Consequently, a high ionic pumping activity is indispensable. Two of our nine α/β-

complexes clearly stand out amongst the other enzyme complexes: αAβ2 and αCβ3. The

activity of αAβ2 is reduced in comparison to the other αA and αB constructs that all show

similar activity levels. Therefore, we suppose that αAβ2 is not a predominant enzyme *in*

vivo. In contrast, the activity of αC combined with β3 is significantly higher and would

therefore fit well with a natural occurrence of this complex in the nervous tissue. Data of

an ongoing proteomic study support this assumption and show that the subunits αC and

β3 are most abundant in the nervous tissue of *O. fasciatus* (M. Herbertz et al.,

unpublished data).

Regardless of the strikingly high overall resistance to inhibition by cardenolides that Na,K-

ATPase preparations of *O. fasciatus* display (12,13,26) we found that the individual α/β-

combinations strongly differed in their resistance against the two cardenolides ouabain

and calotropin. These two cardenolides differ in their structure, steric conformation and

by their ability to permeate membranes, as reflected by their lipophilicity. Ouabain is

produced in plants of the genus *Acokanthera* and in seeds of *Strophanthus gratus* (both

Apocynaceae), it is commercially available, very polar and routinely used in studies on

Na,K-ATPase resistance to cardiac glycosides (1,12,23,25,27). Calotropin, on the other

hand, is a common compound in plants of the tribe Asclepiadeae (Apocynaceae) to which

the host plants of *O. fasciatus* belong. The two cardenolides differ in the conformation of

the steroid rings, which is supposed to determine inhibitory potency. The A/B and C/D

rings of ouabain, and many other cardenolides, are *cis*-fused and the B/C ring is *trans*-

fused (*cis-trans-cis*) (15,31). In contrast calotropin and other cardenolides of *Asclepias*

species, have a *trans-trans-cis* conformation (15) that is predicted to lower the toxicity

(32) and the binding affinity of the cardenolide (31). This may, however, be compensated

by the doubly-linked sugar moiety, typical of *Asclepias* cardenolides (33, 34). A recent

study testing various cardiac glycosides against the Na,K-ATPase of the monarch butterfly,

Danaus plexippus, which is also specialized on *Asclepias* species, actually found calotropin

and its isomer calactin to be the strongest tested inhibitors, with calotropin inhibiting the

monarch's Na,K-ATPase 60-fold more strongly than ouabain (35).

In contrast to the monarch butterfly, there is evidence that the coevolutionary association

between the milkweed bugs and their host plants led to lowered sensitivity towards

calotropin. While the phylogenetically older αC-subunit (11, 26) was roughly 20-times

more strongly inhibited by calotropin than by ouabain not only the overall sensitivity to

both cardenolides but also the relative inhibition by calotropin compared to ouabain

drastically decreased in the phylogenetically more derived αA- and αB-subunits. Dose-

response curves of αB-based enzymes did not significantly differ with respect to inhibition

by either cardenolide whereas the αAβ1 complex was even more resistant to calotropin

than to ouabain. This argues in favor of a specific adaptation to the host plant compounds

over the course of the evolutionary interaction between the milkweed bugs and their host

plants.

The very pronounced insensitivity of the αA- and αB-subunits of *O. fasciatus* to both

cardenolides fits well with their abundant expression in the Malpighian tubules (21) where

the enzymes are directly exposed to toxins in the hemolymph and a very efficient

reabsorption of cardenolides from the primary urine takes place (36). In contrast, the least

sensitive but most active αC is the dominant Na,K-ATPase α-subunit in the nervous tissue

where it is protected from intruding cardenolides by the perineurium (blood-brain-barrier)

which controls xenobiotica access to the nervous system by a tight network of septate

junctions and active efflux carriers (37).

A comparison of the present results with our previous studies (26,27) on *O. fasciatus* Na,K-

ATPase mimics reveals good concordance regarding our conclusions about enzyme

activities. We previously introduced one, two, three or four of the potentially resistance-

conferring substitutions into the α-subunit of *D. melanogaster* to mimic the known α1-

gene copies of *O. fasciatus* (in this order: αD, αC, αB, αA-mimic; c.f. 10,11) and combined

them with two *D. melanogaster* β-subunits, Nrv2.2 and Nrv3 (26,27). The overall behavior

of these mimics strongly resembled the present data for the original *O. fasciatus* Na,K-

ATPase complexes. Most strikingly, the activity levels of the αA and αB mimics were

significantly lower than of the Drosophila wildtype enzyme, the αC or the αD (N122H)

mimics. The two Drosophila β-subunits also modulated the activity: the association with

Nrv3 strongly reduced the activity of all mimics and the Drosophila wildtype enzyme when

compared to those bearing Nrv2.2. In the present study, such a large influence of the β-

subunit on ion pumping activity was observed only for the αC-subunit while the effect was

less pronounced for the αA and αB-subunits.

The β-subunits also strongly modulated the resistance of the enzymes against ouabain,

but their effect was specific to the α-subunits they were combined with. A combination

with Nrv3 instead of Nrv2.2 previously lead to lowered resistance in the αA- and αB-mimic

and increased resistance in the αC-mimic (27). In the present study, resistance changed in

the same direction with β1, β2, or β3 for αA (decreasing) and αC (increasing IC_{50}) but also

lead to increased IC_{50}-values for αB.

The most striking differences, however, appear when comparing the overall ouabain-

sensitivity of the original enzyme combinations with the previously constructed mimics

(27). While the mimics of the *O. fasciatus* αA-copy (featuring the substitutions

Q111T,N122H,F786N,T797A) were by far the most insensitive constructs followed by the

αB-mimics (with Q111T,N122H,F786N), and the more sensitive αC-mimics (N122H,F786S),

the original αA-constructs tested here are at least five times less sensitive than αB-

constructs. On the other hand, the original αC-constructs are 10-times more resistant than

predicted from the experiments with the αC-mimics.

These differences in sensitivity between mimics and original enzymes strongly point to the

importance of additional substitutions in the *O. fasciatus* Na,K-ATPase α1-paralogs. When

constructing the mimics, we focused on the four most conspicuous amino acid exchanges

in the binding pocket of the Na,K-ATPase that were previously shown functionally to be

essential for cardiac glycoside binding (25,38,39). However, there are additional amino

acid substitutions in the cardiac glycoside binding pocket of the Na,K-ATPase α1-genes of

O. fasciatus that may further increase their resistance (11,19). These and further

additional amino acid exchanges that at first appear inconsequential may be of

importance in modulating the effects of the resistance conferring substitutions or

ameliorating their pleiotropic effects (28-30).

321 Conclusion

Resistance to specific cardiac glycosides, enzyme activity and the emerging knowledge on

tissue distribution of the nine α/β-complexes of the large milkweed bug investigated here

further support a scenario of stepwise adaptation to the bugs' toxic host plants: The

duplications leading to three paralogs of the α1-subunit gene that can be combined with

three functional β-subunits enabled a functional diversification of the resulting Na,K-

ATPases allowing for tissue specific adaptations. The high activity of the αC/β-complexes is

suited for efficient ion transport necessary for excitability of nervous tissue but trades off

against lower resistance to high concentrations of cardiac glycosides. These

phylogenetically older complexes were relatively more susceptible to the host plant toxin

calotropin while the younger αA- and αB-subunits were overall more resistant and

relatively less sensitive to calotropin and thus suited for expression in toxin exposed

tissues like the Malpighian tubules. These physiological details provide new fascinating

insights into the continued escalation of defense and counter-defenses between insects

and their poisonous host plants.

Material and methods

Identification of sequences

Two of the three previously published gene copies of the Na,K-ATPase α1 gene (11) (α1B

Acc.No. JQ771519, α1C Acc.No. JQ771518) were incomplete at the 3' end. Both missing

ends could be recovered by RACE PCR (SMARTer RACE kit, Takara Bio, Mountain View, CA,

USA).

Previously published transcriptomic data of *O. fasciatus* (11) were used to identify the full-

length coding sequences of the Na,K-ATPase β-subunits. The raw data were trimmed and

assembled with Trinity (40) as described in Meinzer et al. (41). *O. fasciatus* β-subunit

sequences were recovered using published orthologous sequences from *Locusta*

migratoria (Orthoptera; Acc.No. KF813098-KF813100) and *Blatta germanica* (Blattodea;

Acc.No. HE819405) in tblastn searches against the transcriptome. *De novo* assembly of the

raw reads and quality control was performed with the program CLC-Genomics Workbench

(Qiagen, Hilden, Germany). The resulting *contigs* were screened for open reading frames

(ORFs) with a length of roughly 900 bp (the consensus length of the conserved Na,K-

ATPase β-subunit) using the program VectorNTI (Thermo Fisher, Darmstadt, Germany).

Homology of the *contigs* were assigned by a further round of tblastx searches against

published sequences in genbank.

356 Construction of recombinant baculoviruses

The coding sequences of the Na,K-ATPase α- and β-subunits genes (α1A, α1B, α1C, β1, β2

and β3) were amplified from cDNA transcribed with Superscript III (Thermo Fisher) from

RNA extracted from nervous tissue, gut, testes or muscles with the RNAmagic-kit (Bio-

Budget Technologies GmbH, Krefeld, Germany). For downstream cloning all subunits were

amplified with the Pfu-Phusion Polymerase (Thermo Scientific) and primers including the

full coding sequence and incorporating suitable restriction sites at the 5' end.

PCR products were restricted with the chosen enzymes (Thermo Fisher), gel extracted

(QIAquick gel extraction kit, Qiagen) and ligated into pFastBac-Dual vectors (Thermo

Fisher). The final plasmids had an α-subunit under the control of the polyhedrin promoter

(PH) and a β-subunit under the p10 promoter. Recombinant baculoviruses were generated

as previously described (27) and transfected into *Sf*9 cells to obtain a first (P1) and second

(P2) generation of recombinant virus stock. Uninfected *Sf*9 cells have no or negligible

Na,K-ATPase abundances so that contaminating endogenous Na,K-ATPase activities can be

excluded (42).

Production of recombinant proteins

Production of different recombinant Na,K-ATPase α-β-subunit constructs started from infection of Sf9 cells with P2 virus stock and followed our previously published protocol (27). The cells were harvested by centrifugation, the pellet resuspended and sonicated and cell debris removed by centrifugation. Cell membranes were recovered by ultracentrifugation of the supernatant and redissolved in pure water (ROTIPURAN® p.a., ACS water; Roth). The concentrations of the resuspended protein pellets were determined by Bradford assay (Sigma-Aldrich, Taufkirchen, Germany) using bovine serum albumin (BSA) as a standard.

SDS-PAGE/ Western blotting

Specific binding of customized β-specific anitbodies (Davids Biotechnology, Regensburg, Germany) was additionally validated in our lab by applying the antibodies on two experimental setups. First, the specificity of the antibodies was tested with heterologously expressed *O. fasciatus* Na,K-ATPase protein and second, with four different tissue extracts (muscles, Malpighian tubules, ovaries and nervous tissue). The α1/β1, α1/β2, α1/β3, and α1/βx cell protein construct extracts (4x 30 µg) and muscles, Malpighian tubules, ovaries, and nervous tissue extract (4x 35 µg for each tissue) were run on a SDS-PAGE followed by western blotting as described below. All four Na,K-ATPase constructs and tissues were immunostained with each of the specific β-antibodies. These stainings verified that no unspecific binding occurs (Fig. S2 & S3), but that the β-specific antibodies solely detect their target-subunit in all tested samples.

Thirty µg of the heterologously expressed proteins were deglycosylated with

endoglycosidase H following the manufacturer's protocol (Promega GmbH, Wiesloch,

Baden Württemberg, Germany) but instead of boiling proteins for 5 minutes at 95°C we

performed the denaturation step at 50°C for 30 minutes to maintain the integrity of the

protein. SDS-polyacrylamide gel electrophoresis and western blotting was performed as

previously reported (27) with few modifications. The membrane proteins were blotted on

a nitrocellulose membrane for 90 minutes at 100 V with an additional ice block at 4°C. To

mask unspecific binding sites the membrane was blocked with BlueBlock PF (Serva,

Heidelberg, Germany). The split membranes were incubated with the primary monoclonal

antibodies α5, 12G10 anti-α-tubulin (DSHB Hybridoma Bank, University of Iowa; deposited

by D.M. Fambrough (43) and by J. Frankel and E.M. Nelsen), and customized specific

polyclonal antibodies for the subunits β1, β2, and β3 (Davids Biotechnology, Regensburg,

Germany). For detection of the primary antibodies the membrane was incubated for one

hour with goat anti-mouse (α5, 12G10 anti-α-tubulin), anti-rabbit (β2), and anti-chicken

(β1, β3) secondary antibodies conjugated with horseradish peroxidase (Dianova,

Hamburg, Germany) and developed with 4-chloro-1-naphtol as previously described (27).

Extraction and preparation of cardenolides

Calotropin was extracted according to (35) from approximately 8 g of dried larvae of the

monarch butterfly *Danaus plexippus* reared on *Asclepias curassavica*. The cardenolides

were extracted with methanol, lipids extracted with hexane, and the cardenolides

retrieved by chloroform extraction. The concentrated methanolic extract was subjected to

preparative HPLC (Agilent 1100) equipped with a photodiode array detector and a semi-

preparative RP-Nucleodur C18 Htec column (5 μm, 250 x 10 mm, Macherey-Nagel, Düren,

Germany) using a water-acetonitrile gradient. Calotropin was identified by its retention

time and checked for molecular weight and LS-MS spectrum by LC-ESI-TOF spectrometry

by Dr. Maria Riedner (Department of Chemistry, Universität Hamburg) and the structure

confirmed by NMR spectroscopy. The resulting calotropin preparation was 80% pure.

Calotropin and the commercially available ouabain (O3125, Sigma-Aldrich) were first

dissolved in DMSO and prepared for Na,K-ATPase activity assays as dilutions in sterile H_2O

reaching final concentrations from 10^{-8} to 10^{-3} M with 2% DMSO.

Na,K-ATPase activity and inhibition assay

Activities of the nine different Na,K-ATPase α/β-subunit combinations of *O. fasciatus* were

determined by photometric measurements of inorganic phosphate released from

enzymatic ATP hydrolysis while subtracting background ATPase activity following

previously described methods (27,44). Based on previous results (27) the buffer regime

used was 50 mM NaCl, 4 mM $MgCl_2$, 50 mM imidazol, 20 mM KCl, pH 7.4. The inhibition of

the different Na,K-ATPases was tested under increasing ouabain and calotropin

concentrations and expressed as percentage activity of a non-inhibited control. For each

α/β-subunit construct technical duplicates of three biological replicates of recombinant

proteins isolated from independently transfected *Sf*9 cells were tested.

Statistic analysis of data

After checking for normal distribution (Shapiro Wilk's test) and variance homogeneity

(Levene's test on squared deviations), differences in total Na,K-ATPase activities were

tested and compared between each α/β-subunit construct by one-way ANOVA followed

by Tukey's HSD posthoc test. The data of each Na,K-ATPase α/β-subunit construct was

subjected to nonlinear curve fitting with dose response function and Levenberg

Marquardt iteration algorithm in Origin-Pro 9.1G (OriginLab Corporation, Northampton,

USA) using a four parameter logistic function, top asymptote was set to 100, bottom

asymptote set to 0 and a variable Hill slope. For each construct IC_{50} values were calculated

and differences were analyzed separately for constructs affected by ouabain and

calotropin by one-way ANOVA followed by Tukey's HSD posthoc tests. A pairwise, F-test

based comparison of similarity between the fitted curves was performed using the

command "compare datasets". All calculations were performed with Origin-Pro 9.1G.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

457 The datasets generated during the current study are available in the Dryad repository at

458 https://doi.org/10.5061/dryad.j0zpc86dg, sequence data will be deposited in the European

459 Nucleotide Archive (ENA) upon acceptance.

Competing interests

461 The authors declare that they have no competing interests.

Funding

463 MW was supported by a stipend of the Nachwuchsförderung Universität Hamburg, and

464 the whole study by grants of the Deutsche Forschungsgemeinschaft (DO527/5-3 and 10-1)

465 and the Templeton Foundation (grant ID #41855) to SDo. These funding bodies played no

466 role in the design of the study and collection, analysis, and interpretation of data and in

467 writing the manuscript.

Author's contribution

470 SDa and SDo conceptualization; SDa, MW, VW, RT, MH cloning and protein expression;

471 SDa and MW enzyme assays, data analysis and visualization; MW and SDo writing-original

472 draft; SDa writing-review and editing.

Acknowledgments

We acknowledge the help of Sebastian Gatemann who purified calotropin as part of his BSc thesis. Tammy Tischer supervised by Jennifer Lohr complemented the 5' end of the αC copy. Shab Mohammadi improved the manuscript by proofreading the English and valuable discussions. We thank Daniel Tal for generating the structures in Fig. 1.

References

1. Horisberger J-D. Recent Insights into the structure and mechanism of the sodium pump. Physiology. 2004;19:377–87.

2. Geering K. The functional role of β subunits in oligomeric P-type ATPases. J Bioenerg Biomembr. 2001;33:425–38.

3. Blanco G, Mercer RW. Isozymes of the Na-K-ATPase: heterogeneity in structure, diversity in function. Am J Physiol Ren Physiol. 1998;275:F633–50.

4. Blanco G. Na,K-ATPase subunit heterogeneity as a mechanism for tissue-specific ion regulation. Semin Nephrol. 2005;25:292–303.

5. Geering K. Function of FXYD proteins, regulators of Na,K-ATPase. J Bioenerg Biomembr. 2005;37:387–92.

6. Okamura H, Yasuhara JC, Fambrough DM, Takeyasu K. P-type ATPases in *Caenorhabditis* and *Drosophila*: implications for evolution of the P-type ATPase subunit families with special reference to the Na,K-ATPase and H,K-ATPase subgroup. J Membr Biol. 2002;191:13–24.

7. Baumann O, Salvaterra PM, Takeyasu K. Developmental changes in β-subunit composition of Na,K-ATPase in the *Drosophila* eye. Cell Tissue Res. 2010;340:215–

497   28.

498   8.   Yasuhara JC, Baumann O, Takeyasu K. Localization of Na/K-ATPase in developing
499   and adult *Drosophila melanogaster* photoreceptors. Cell Tissue Res. 2000;300:239–
500   49.

501   9.   Paul SM, Palladino MJ, Beitel GJ. A pump-independent function of the Na,K-ATPase
502   is required for epithelial junction function and tracheal tube-size control.
503   Development. 2007;134:147–55.

504   10.   Yang L, Ravikanthachari N, Mariño-Pérez R, Deshmukh R, Wu M, Rosenstein A, et al.
505   Predictability in the evolution of Orthopteran cardenolide insensitivity. Philos Trans
506   R Soc Lond B Biol Sci. 2019;374:20180246.

507   11.   Zhen Y, Aardema ML, Medina EM, Schumer M, Andolfatto P. Parallel molecular
508   evolution in an herbivore community. Science. 2012;337:1634–7.

509   12.   Moore L V., Scudder GGE. Ouabain-resistant Na,K-ATPases and cardenolide
510   tolerance in the large milkweed bug, *Oncopeltus fasciatus*. J Insect Physiol.
511   1986;32:27–33.

512   13.   Bramer C, Dobler S, Deckert J, Petschenka G. Na+/K+-ATPase resistance and
513   cardenolide sequestration: basal adaptations to host plant toxins in the milkweed
514   bugs (Hemiptera: Lygaeidae: Lygaeinae). Proc R Soc B Biol Sci. 2015;282:20142346.

515   14.   Agrawal AA, Petschenka G, Bingham RA, Weber MG, Rasmann S. Toxic
516   cardenolides: chemical ecology and coevolution of specialized plant-herbivore
517   interactions. New Phytol. 2012;194:28–45.

518   15.   Malcolm SB. Cardenolide-Mediated Interactions between Plants and Herbivores. In:
519   Rosenthal GA, Berenbaum MR, editors. Herbivores: their Interactions with
520   Secondary Plant Metabolites. San Diego: Academic Press; 1991. p. 251–296.

521   16.   Petschenka G, Wagschal V, Von Tschirnhaus M, Donath A, Dobler S. Convergently
522   evolved toxic secondary metabolites in plants drive the parallel molecular evolution
523   of insect resistance. Am Nat. 2017;190:S29–43.

524   17.   Ujvari B, Casewell NR, Sunagar K, Arbuckle K, Wüster W, Lo N, et al. Widespread
525   convergence in toxin resistance by predictable molecular evolution. Proc Natl Acad
526   Sci. 2015;112:11911–6.

527   18.   Mohammadi S, Gompert Z, Gonzalez J, Takeuchi H, Mori A, Savitzky AH. Toxin-
528   resistant isoforms of Na+/K+- ATPase in snakes do not closely track dietary
529   specialization on toads. Proc R Soc B. 2016;283:20162111.

530   19.   Dobler S, Dalla S, Wagschal V, Agrawal AA. Community-wide convergent evolution
531   in insect adaptation to toxic cardenolides by substitutions in the Na,K-ATPase. Proc
532   Natl Acad Sci. 2012;109:13040–5.

533   20.   Dobler S, Wagschal V, Pietsch N, Dahdouli N, Meinzer F, Romey-Glüsing R, et al.
534   New ways to acquire resistance: imperfect convergence in insect adaptations to a
535   potent plant toxin. Proc R Soc B. 2019;286:20190883.

536   21.   Lohr JN, Meinzer F, Dalla S, Romey-Glüsing R, Dobler S. The function and
537   evolutionary significance of a triplicated Na,K-ATPase gene in a toxin-specialized
538   insect. BMC Evol Biol. 2017;17:256.

539   22.   Price EM, Lingrel JB. Structure - function relationships in the Na,K-ATPase α subunit:
540   site-directed mutagenesis of glutamine-111 to arginine and asparagine-122 to
541   aspartic acid generates a ouabain-resistant enzyme. Biochemistry. 1988;27:8400–8.

542   23.   Laursen M, Gregersen JL, Yatime L, Nissen P, Fedosova NU. Structures and
543   characterization of digoxin- and bufalin-bound Na^+,K^+-ATPase compared with the
544   ouabain-bound complex. Proc Natl Acad Sci. 2015;112:1755–60.

545   24.   Palasis M, Kuntzweiler TA, Argüello JM, Lingrel JB. Ouabain interactions with the
546   H5-H6 hairpin of the Na,K-ATPase reveal a possible inhibition mechanism via the
547   cation binding domain. J Biol Chem. 1996;271:14176–82.

548   25.   Croyle ML, Woo AL, Lingrel JB. Extensive random mutagenesis analysis of the Na + /
549   K + - ATPase a subunit identifies known and previously unidentified amino acid
550   residues that alter ouabain sensitivity implications for ouabain binding. Eur J

Biochem. 1997;248:488–95.

26.   Dalla S, Dobler S. Gene duplications circumvent trade-offs in enzyme function:
insect adaptation to toxic host plants. Evolution (N Y). 2016;70:2767–77.

27.   Dalla S, Baum M, Dobler S. Substitutions in the cardenolide binding site and
interaction of subunits affect kinetics besides cardenolide sensitivity of insect Na,K-
ATPase. Insect Biochem Mol Biol. 2017;89:43–50.

28.   Karageorgi M, Groen SC, Sumbul F, Pelaez JN, Verster KI, Aguilar JM, et al. Genome
editing retraces the evolution of toxin resistance in the monarch butterfly. Nature.
2019;574:409–12.

29.   Taverner AM, Yang L, Barile ZJ, Lin B, Peng J, Pinharanda AP, et al. Adaptive
substitutions underlying cardiac glycoside insensitivity in insects exhibit epistasis in
vivo. Elife. 2019;8:48224.

30.   Mohammadi S, Yang L, Harpak A, Herrera-Álvarez S, Rodriguez-Ordonez M del P,
Peng J, et al. Concerted evolution reveals co-adapted amino acid substitutions in
frogs that prey on toxic toads. Curr Biol. 2021; accepted

31.   Katz A, Lifshitz Y, Bab-Dinitz E, Kapri-Pardes E, Goldshleger R, Tal DM, et al.
Selectivity of digitalis glycosides for isoforms of human Na,K-ATPase. J Biol Chem.
2010;285:19582–92.

32.   Hoch JH. A survey of cardiac glycosides and genins. Columbia, SC, USA: University of
South Carolina Press; 1961.

33.   Venkateswara Rao R, Vaidyanathan CS. Chemistry and biochemical pharmacology of
cardiac glycosides - A review. J Indian Inst Sci. 1991;71:329–64.

34.   Piacente S, Masullo M, De Nève N, Dewelle J, Hamed A, Kiss R, et al. Cardenolides
from *Pergularia tomentosa* display cytotoxic activity resulting from their potent
inhibition of Na+/K+-ATPase. J Nat Prod. 2009;72:1087–91.

35.   Petschenka G, Fei CS, Araya JJ, Schröder S, Timmermann BN, Agrawal AA. Relative

577   selectivity of plant xardenolides for Na+/K+-ATPases from the monarch butterfly

578   and non-resistant insects. Front Plant Sci. 2018;9:1424.

579 36. Meredith J, Moore L, Scudder GG. Excretion of ouabain by Malpighian tubules of

580   *Oncopeltus fasciatus*. Am J Physiol - Regulatory Integrative Comp Physiol

581   1984;246:705-715.

582 37. Dermauw W, Van Leeuwen T. The ABC gene family in arthropods: comparative

583   genomics and role ininsecticide transport and resistance. Insect Biochem Mol Biol.

584   2014;45:89–110.

585 38. Qiu LY, Krieger E, Schaftenaar G, Swarts HGP, Willems PHGM, De Pont JJHHM, et al.

586   Reconstruction of the complete ouabain-binding pocket of Na,K-ATPase in gastric

587   H,K-ATPase by substitution of only seven amino acids. J Biol Chem.

588   2005;280:32349–55.

589 39. Holzinger F, Frick C, Wink M. Molecular basis for the insensitivity of the Monarch

590   (*Danaus plexippus*) to cardiac glycosides. FEBS Lett. 1992;314:477–80.

591 40. Grabherr MG., Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. Trinity:

592   reconstructing a full-length transcriptome without a genome from RNA-Seq data.

593   Nat Biotechnol. 2011;29:644–52.

594 41. Meinzer F, Dobler S, Donath A, Lohr JN. Robust reference gene design and

595   validation for expression studies in the large milkweed bug, *Oncopeltus fasciatus*,

596   upon cardiac glycoside stress. Gene. 2019;710:66–75.

597 42. Blanco G. The Na/K-ATPase and its isozymes: What we have learned using the

598   baculovirus expression system. Front Biosci. 2005;10(SUPPL. 1):2397–411.

599 43. Lebovitz RM, Takeyasu K, Fambrough DM. Molecular characterization and

600   expression of the (Na+ + K+)-ATPase alpha-subunit in *Drosophila melanogaster*.

601   EMBO J. 1989;8:193–202.

602 44. Petschenka G, Fandrich S, Sander N, Wagschal V, Boppré M, Dobler S. Stepwise

603   evolution of resistance to toxic cardenolides via genetic substitutions in the Na+/K+-

604   ATPase of milkweed butterflies (Lepidoptera: Danaini). Evolution (N Y).

605   2013;67:2753–61.

a)

608
b)

Figure 1: Structure of the cardenolides ouabain (a) and calotropin (b). Both are

characterized by a monosaccharide attached to a steroid skeleton at C3 (4 steroid rings: A-

D) and a five-membered unsaturated lactone ring at C-17. a) In ouabain the

monosaccharide is attached by a single glycosidic link to the steroid rings that are *cis-*

trans-cis fused. b) Calotropin is characterized by a doubly linked monosaccharide as well

as a *trans-trans-cis* conformation of the steroid rings (figure generated by Daniel Tal, The

Weizmann Institute of Science, 2020).

Figure 2: Western blot of nine α/β-complexes of *O. fasciatus* expressed heterologously in cell culture. 30 µg of protein were separated by SDS-PAGE and blotted on a nitrocellulose membrane. The upper membrane slice shows at a size of 110 kDa αA, αB and αC in alternating order, the proteins were stained with the primary antibody α5 (DSHB hybridoma bank). At 55 kDa α-tubulin is shown as a positive control (middle membrane), the protein is stained with the primary antibody 12G10 anti-α-tubulin (DSHB hybridoma bank). Primary antibodies of both proteins were detected with the secondary HRP-conjugated antibody goat-anti mouse (Dianova). The associated β-subunits 1, 2, and 3 were deglycosylated in advance. The lower membrane is split into three slices, the first one shows β1, the second β2, and the third β3. The subunits were stained with β-subunit specific customized primary antibodies (produced by Davids Biotechnology) and HRP-conjugated secondary antibodies goat-anti rabbit (for β2, Dianonva) and goat-anti chicken (for β1 and 3, Dianova). The different β-subunits show bands at weights ranging between 35 and 40 kDa. The bands were visualized with 4-CN. Please note: Signal strength is not comparable between the different β-subunits, due to different binding affinities of the antibodies. Due to the grey scale some bands of the multicolor protein ladder were marked by lines.

Figure 3: Na,K-ATPase activity [nmol Pi/mg protein*min] of nine α/β-complexes of *O.*

fasciatus. Each bar depicts the mean and standard deviation of three biological replicates.

Different letters above the bars indicate significant differences between enzymes (ANOVA,

$F_{8,18}$=77.97, p<0.001). Black bars: αA with three β-subunits and overall activities ranging

from **5.5 to 9.3 nmol Pi/mg protein*min**. Bars with a grey dotted outline: data for the αA

mimic (THNA) combined with Nrv2.2 and Nrv3 of a previous study (Dalla et al., 2017). Dark

gray bars: αB/β-complexes ranging from **10.3 to 13 nmol Pi/mg protein*min**. Light gray

bars: αC/β-complexes with activity levels ranging from **19.7 to 32.7 nmol Pi/mg**

protein*min. The activity levels of the three αC/β-enzymes are significantly higher than

those of the other enzymes, with the highest activity reached by αC combined with β3.

Figure 4: *In vitro* inhibition by calotropin and ouabain of nine α/β-complexes of *O. fasciatus* expressed heterologously in cell culture. Dose response relationships between increasing cardenolide concentrations and residual enzyme activities were expressed in % of the non-inhibited control (calculated with Origin Pro 9.1G). Each data point represents the mean of three biological replicates and standard deviation. Effect of calotropin (blue curves and round symbols) and ouabain (black curves and square symbols) a) on αA, b) on αB, and c) on αC combined each with β1, 2, and 3. Please note the different scales.

Supplemental

```
Of_ß1 MKGKTGNH-KEAEYSWGYKKPEDDNRTLKEKIRIAFYNPSTGTVLGRTPLDWGRLLLFYILLYLALALLFAICMKGL
Of_ß2 MEANTQEKKKEVGF---YSAPK--QLTGWAYIQNFLYDPVEGKVLGRTGPSWAKIGVFYLIFYSVLASLFGIMLWIF
Of_ß3 MADKAKEE-----F---YTRPA--KVGNWDAFKLFLWNSETSQFLGRTFSSWAKILLFYLIFYACLMGFFSALLALF
Of_ßx ----------------------------------------------------------------MKHLLAVLFCFL

Of_ß1 LLTIDLKHPKWILEESIIGTNPGMGYRPMPKMAEDGKSLVWYQANNKSNVAMWEESINEFLKPYMNEH---EFTNRM
Of_ß2 YHTLDPRVPRWQLRESLIGQDPGLGFRPMPNDTDSRSTLLWYQGKSRESCNLWTKTLENFLEVYRKPGLTPGRGQNI
Of_ß3 FQTLDFKQPKWQQASSLIGSNPGLGFRPMPPESHVESTLIWFKV-SESNYATWTNKLDEFLKPYREPD--PKYENVM
Of_ßx QHSL--------------GDQITLKVHPLSEDS--KSTLIWF---NSSTYDRWAQQLDNFLEVYRNPTLTDGREENL

Of_ß1 F-CDYDMPKPSEDKVCEVYINETSWGPCTKRNHYSFPSSSPCVFLKLNRIYGWTPEYYEDLSSLPPEMPSNLRSFI
Of_ß2 YNCDYDRP-PGKGQVCDVDVKS--WYPCTLENNFNYHLSSPCVFIKLNKIYGWQPEYYNDSRDFPETMPEDLKNHI
Of_ß3 QSCDFDTP-DTPDRVCKVDIKS--WSPCVKENNYAYHNGAPCIFLKLNKIYGWMPEFYNDTETLPKDMPEDLKSHI
Of_ßx VTCNYNKY-PSTDSVCNIRPEE--MKPCTASNFFGYKSGTPCVFIELEKVLGWKPKYFSSGKSLPKTMPKNLKEKI

Of_ß1 ENEKAIDPRRMNTIWVSCNPETPFDNETMGEVEYYPRQGFPGYYYPYKNQDGYLSPLIAVRFKRPPVGILVNVECR
Of_ß2 LTQKAINPELIKTIWVSCEGESPHDRENIGEIQYIPRQGFPGYFYPYVNTEGYLSPLVAVNFKRPKTGVIINVECR
Of_ß3 NEEKKHNPKSLNMVWVSCEGENPADVENIGSIQYIPRRGFPGFYFPYKNQEGYLSPVIAIWFEKPVNGILINIECK
Of_ßx LETTEVNLKMWETIWLSCDGETPADKEMIGPIQILPQPGFPGYFFPYIGQPGYLNPVVAVHFERPLGGLVINIVCK

Of_ß1 AWAKNIIYQRPTGAKDRQGSVHFEILVDP*
Of_ß2 AWAKNLKVDR----KERIGIVHFELLID*
Of_ß3 AWARNIHHDR----AERRGSVHFELMID*
Of_ßx AWAPGIIHDM----NKPLGVARFELLID*
```

Figure S1: Alignment of deduced *O. fasciatus* Na,K-ATPase α and β-subunit protein sequences. Full length coding sequence of all four β-subunits, tyrosine rich motif typical for β-subunits in orange, cysteine residues forming three conserved intramolecular disulfide bridges in yellow, potential glycosylation sites in blue, and peptides used to raise β-subunit specific antibodies are marked in grey. The N-terminal truncated βx was not further investigated in this study.

Figure S2: Antibody specificity test with 30µg of heterologously expressed α/β enzymes via SDS-PAGE and western blot. Each set of α1 combined with β1, β2, β3, and βx respectively was stained with α5 primary antibody (upper membrane with α1-subunits at 110 kDa) and one of the four β-specific antibodies: a) β1, b) β2, c) β3, and d) βx-specific antibody, lower membrane with dissociated β-subunits). The middle membrane of the last set (d) was additionally stained with 12G10-anti-α-tubulin. Both, the α-tubulin and α1-staining function as a positive control. The primary antibodies were detected with HRP-conjugated secondary antibody (α5 and 12G10-anti-α-tubulin: goat-anti mouse, β1 and β3: goat-anti chicken, β2 and βx: goat-anti rabbit). The western blots show that the β-specific antibodies solely stain their target subunits (bold letters mark the specific stained β-subunits).

Figure S3: Antibody specificity test with 35µg four different tissue extracts (muscle

(mus), Malpighian tubules (malp), ovaries (ov), nervous tissue (ns)) via SDS-PAGE and

western blot. On each membrane set, consisting of an upper and lower slice, one α1/β-

complex is shown. Each set of four different tissue extracts was stained with α5 primary

antibody (α1: 110 kDa) and 12G10-anti-α-tubulin (α-tubulin: 55 kDa) on the upper slices

and one of the four β-specific antibodies (β: 30-45 kDa; from upper row left to bottom

row right: β1-, β2-, β3- and βx-specific antibody) on the lower slices. The 12G10-anti-α-

tubulin staining functions as a positive control. The primary antibodies were detected with

HRP-conjugated secondary antibody (α5 and 12G10-anti-α-tubulin: goat-anti mouse, β1

and β3: goat-anti chicken, β2 and βx: goat-anti rabbit). The immunostained western blots

show differences between abundances and glycosylation pattern in each tissue unique for

each β-subunit.