1

Different combinations of insect Na,K-ATPase α- and β-subunit paralogs enable fine tuning of toxin resistance and enzyme kinetics

Marlena Winter[1#], Safaa Dalla[1,2#], Vera Wagschal[1], Rohin Turjalei[1], Marlies Heiser[1], Susanne Dobler[1*]

[1]Institute of Zoology, Molecular Evolutionary Biology, Universität Hamburg, Hamburg, Germany

[2] present address: Tishreen University, Latakia, Syria

[#] Co-first author

[*]Corresponding author: Susanne Dobler

E-mail: susanne.dobler@uni-hamburg.de

Running title: Kinetics and toxin resistance of Na,K-ATPase paralogs

Keywords: Na+/K+-ATPase, cardiac glycosides, toxicity, resistance, molecular evolution, baculovirus expression system

Financial support - Marlena Winter was supported by a stipend of the Nachwuchsförderung Universität Hamburg, and the whole study by grants of the Deutsche Forschungsgemeinschaft (DO527/5-3 and 10-1) and the Templeton Foundation (grant ID #41855) to Susanne Dobler.

Summary statement

Activity and cardiac glycoside resistance of the *Oncopeltus fasciatus'* Na,K-ATPase is determined by amino acid substitutions in the α-subunit binding pocket, the α/β-subunit combination and the cardiac glycoside identity.

Abstract

Cardiac glycosides are known to fatally inhibit the Na,K-ATPase throughout the animal kingdom. Several animals, however, evolved target-site insensitivity by substitution in the otherwise highly conserved cardiac glycoside binding pocket located on the Na,K-ATPase α-subunit. The minimal functional enzyme consists of an α- and a β-subunit, considered mainly as a chaperone responsible for correct folding and membrane integration. We here analyze resistance to cardiac glycosides and kinetic properties of different Na,K-ATPase α-β-combinations of the large milkweed bug, *Oncopeltus fasciatus.* These insects have adapted to high concentrations of cardiac glycosides in their food plants via several rounds of ATPase α1 gene duplications followed by subfunctionalization of the enzymes.

To investigate their characteristics we expressed nine combinations of Na,K-ATPase α/β-sunbunits (three each) in *Sf*9 cells and tested them with two structurally distinct cardiac glycosides, calotropin, a host plant compound, and ouabain, a commonly used toxin. Differences in the number and identity of amino acid substitutions in the cardiac glycoside binding site resulted in large differences in activity and toxin resistance of the three α-

subunits. The enzymes' kinetics were also influenced by the β-subunits leading to

increased activities (αCβ3) or altered resistances. Calotropin proved to be a much more

potent inhibitor than ouabain. Our results show that the possession of multiple paralogs

enables adaptation to plant toxins and mitigates pleiotropic effects by a compromise

between ion pumping activity and resistance.

Introduction

The transmembrane Na,K-ATPase, is ubiquitously expressed in animal cells and is essential

for maintaining membrane potentials, controlling cell volume and the overall functionality

of cells (Horisberger, 2004). The functional enzyme consists of two subunits, the catalytic

α-subunit and an associated β-subunit serving as chaperone for correct folding and

membrane integration (Geering, 2001). In vertebrates several paralogs are present for

both genes and the enzyme's characteristics differ depending on the specific combination

of α- and β-subunits (Blanco, 2005; Blanco and Mercer, 1998). An additional modulatory

and tissue specific γ-subunit is present in vertebrate tissues (Geering, 2005), but has not

been detected in invertebrates. Much less is known about the importance of different α-

β-subunit combinations in invertebrates (Okamura et al., 2002). Only in *Drosophila* the

tissue distribution and functional roles of some of these combinations have been explored

to some extent (Baumann et al., 2010; Paul et al., 2007; Yasuhara et al., 2000). We here

explore the functional role of different Na,K-ATPase complexes in the large milkweed bug,

Oncopeltus fasciatus, that have undergone three rounds of duplications of the Na,K-

ATPase (Yang et al., 2019; Zhen et al., 2012) and display high insensitivity against cardiac

glycosides (Bramer et al., 2015; Moore and Scudder, 1986), which are universal inhibitors

of the Na,K-ATPase (Horisberger, 2004).

Inhibition of the Na,K-ATPase by cardiac glycosides, such as the cardenolides ouabain and

calotropin (Fig. 1), depends on a specific binding site located on the extracellular side of

the catalytic α-subunit. Upon binding of cardiac glycosides the enzyme is arrested in the

phosphorylated state and ion pumping is disrupted (Horisberger, 2004). Inhibition of the

Na,K-ATPase is harmful and can be lethal to almost all animals. There are, however, a few

notable exceptions to this: various mammals, frogs and reptiles prey unharmed on cardiac

glycoside-containing toads and a number of phytophagous insects feed - often exclusively

- on cardiac glycoside-containing host plants (reviewed in Agrawal et al., 2012; Malcolm,

1991). Most of these species avoid the toxic effect of cardiac glycosides and exhibit target

site insensitivity, by drastically lowering binding affinity of cardiac glycosides to their Na,K-

ATPase α-subunit (reviewed in Petschenka et al., 2017; Ujvari et al., 2015). Strikingly,

identical amino acid substitutions at homologous positions convergently evolved in

various, distantly related insect groups and even resemble the substitutions observed in

cardiac glycoside-resistant vertebrates (Dobler et al., 2012; Mohammadi et al., 2016;

Ujvari et al., 2015; Zhen et al., 2012).

While insect species usually express a single gene copy of the Na,K-ATPase α-subunit

(ATPα1) throughout their body, this gene underwent one to three rounds of gene

duplications in several cardiac glycoside-exposed species (Dobler et al., 2019; Petschenka

et al., 2017; Yang et al., 2019; Zhen et al., 2012). These duplicated ATPα1 copies differ in

the number and identity of presumably resistance-causing amino acid substitutions and

show tissue specific expression patterns (Dobler et al., 2019; Lohr et al., 2017; Yang et al.,

2019; Zhen et al., 2012).

The large milkweed bug *Oncopeltus fasciatus* (Hemiptera, Lygaeidae) possesses four

ATPα1 paralogs (A, B, C, D) that differ in the number and identity of amino acid

substitutions (Yang et al., 2019; Zhen et al., 2012). We previously introduced substitutions

observed in these paralogs at four usually highly conserved amino acid sites (Q111, N122,

F786 and T797) known to be important for high affinity to cardiac glycosides (Croyle et al.,

1997; Laursen et al., 2015; Palasis et al., 1996; Price and Lingrel, 1988) into the sensitive

ATPα1 gene of *Drosophila melanogaster* (Dalla and Dobler, 2016; Dalla et al., 2017). The

N122H (as observed in ATPα1D of *O. fasciatus*), N122H+T797S (as observed in ATPα1C),

Q111T+N122H+F786N (as in ATPα1B) or Q111T+N122H+F786N+T797A (as in ATPα1A)

substitutions yielded proteins with stepwise increasing resistance to the cardenolide

ouabain. At the same time, enzyme activity decreased with increasing numbers of

substitutions (Dalla and Dobler, 2016).

Additional untested substitutions are present in the *O. fasciatus* ATPα1 paralogs and these

might influence the characteristics of the enzymes, potentially further increasing

resistance but possibly also serving as compensatory mutations mitigating deleterious

intra-molecular epistatic effects of previously characterized mutations (Karageorgi et al.,

2019; Taverner et al., 2019). In addition, it has been shown that the β-subunit of the Na,K-

ATPase not only acts as a chaperone, guiding the heterodimeric enzyme to the cell

membrane, but also influences the behavior of the Na,K-ATPase including its sensitivity to

ouabain (Blanco, 2005; Dalla et al., 2017; Geering, 2001).

Although ouabain is a commonly used cardiac glycoside in studies of the Na,K-ATPase, it

does not occur in host plants of *O. fasciatus*. The *Asclepias* species (Apocynaceae) that the

bugs feed on typically contain different types of cardenolides characterized by a *trans-*

trans-cis conformation of the steroidal skeleton and a single sugar moiety attached by a

cyclic bridge as in the abundant calotropin (Malcolm, 1991) (Fig. 1). The most biologically

relevant test of the bugs' resistance against cardiac glycosides should thus involve a plant-

derived compound like calotropin. To address these issues, we identified and

heterologously expressed nine possible *O. fasciatus* Na,K-ATPase α/β combinations and

characterized their activities and sensitivities to ouabain and calotropin.

Since activity and resistance of the Na,K-ATPase is determined by the number and identity

of amino acid substitutions in the cardenolide binding site located in the α-subunit, (1) we

hypothesized that α-subunit paralogs – associated with different β-subunits – differ

significantly in their activities and insensitivities to the two cardenolides we tested. (2) We

further hypothesized that the different β-subunits modulate the Na,K-ATPase α-subunit

paralog behavior influencing their response to the cardenolides. Additionally, although

ouabain and calotropin both seem to be strong inhibitors of the Na,K-ATPase, (3) we

expected the two cardenolides to differ significantly in their inhibitory potency against the

Na,K-ATPase of *O. fasciatus* due to differences in structure, steric conformation and

lipophilicity.

Material and methods

Identification of sequences

Two of the three previously published gene copies of the Na,K-ATPase α1 gene (Zhen et

al., 2012) (α1B Acc.No. JQ771519, α1C Acc.No. JQ771518) were incomplete at the 3' end.

Both missing ends could be recovered by RACE PCR (SMARTer RACE kit, Takara Bio,

Mountain View, CA, USA).

Previously published transcriptomic data of *O. fasciatus* (Zhen et al., 2012) were used to

identify the full-length coding sequences of the Na,K-ATPase β-subunits. The raw data

were trimmed and assembled with Trinity (Grabherr et al., 2011) as described in Meinzer

et al. (2019). *O. fasciatus* β-subunit sequences were recovered using published

orthologous sequences from *Locusta migratoria* (Orthoptera; Acc.No. KF813098-

KF813100) and *Blatta germanica* (Blattodea; Acc.No. HE819405) in tblastn searches

against the transcriptome. *De novo* assembly of the raw reads and quality control was

performed with the program CLC-Genomics Workbench (Qiagen, Hilden, Germany). The

resulting *contigs* were screened for open reading frames (ORFs) with a length of roughly

900 bp (the consensus length of the highly conserved Na,K-ATPase β-subunit) using the

program VectorNTI (Thermo Fisher, Darmstadt, Germany). Homology of the *contigs* were

assigned by a further round of tblastx searches against published sequences in genbank.

Construction of recombinant baculoviruses

The coding sequences of the Na,K-ATPase α- and β-subunits genes (α1A, α1B, α1C, β1, β2

and β3) were amplified from cDNA transcribed with Superscript III (Thermo Fisher) from

RNA extracted from nervous tissue, gut, testes or muscles with the RNAmagic-kit (Bio-

Budget Technologies GmbH, Krefeld, Germany). For downstream cloning all subunits were

amplified with the Pfu-Phusion Polymerase (Thermo Scientific) and primers including the

full coding sequence and incorporating suitable restriction sites at the 5' end.

PCR products were restricted with the chosen enzymes (Thermo Fisher), gel extracted

(QIAquick gel extraction kit, Qiagen) and ligated into pFastBac-Dual vectors (Thermo

Fisher). The final plasmids had an α-subunit under the control of the polyhedrin promoter

(PH) and a β-subunit under the p10 promoter. Recombinant baculoviruses were generated

as previously described (Dalla et al., 2017) and transfected into *Sf*9 cells to obtain a first

(P1) and second (P2) generation of recombinant virus stock.

Production of recombinant proteins

Production of different recombinant Na,K-ATPase α-β-subunit constructs started from

infection of Sf9 cells with P2 virus stock and followed our previously published protocol

(Dalla et al., 2017). The cells were harvested by centrifugation, the pellet resuspended and

sonicated and cell debris removed by centrifugation. Cell membranes were recovered by

ultracentrifugation of the supernatant and redissolved in pure water (ROTIPURAN® p.a.,

ACS water; Roth). The concentrations of the resuspended protein pellets were determined

by Bradford assay (Sigma-Aldrich, Taufkirchen, Germany) using bovine serum albumin

(BSA) as a standard.

SDS-PAGE/ Western blotting

Thirty μg of the heterologously expressed proteins were deglycosylated with

endoglycosidase H following the manufacturer's protocol (Promega GmbH, Wiesloch,

Germany) but instead of boiling proteins for 5 minutes at 95°C we performed the

denaturation step at 50°C for 30 minutes to maintain the integrity of the protein. SDS-

polyacrylamide gel electrophoresis and western blotting was performed as previously

reported (Dalla et al., 2017) with few modifications. The membrane proteins were blotted

on a nitrocellulose membrane for 90 minutes at 100 V with an additional ice block at 4°C.

To mask unspecific binding sites the membrane was blocked with BlueBlock PF (Serva,

Heidelberg, Germany). The split membranes were incubated with the primary monoclonal

antibodies α5, α-tubulin (DSHB Hybridoma Bank, University of Iowa; deposited by D.M.

Fambrough (Lebovitz et al., 1989), and customized specific polyclonal antibodies for the

subunits β1, β2 and β3 (Davids Biotechnology, Regensburg, Germany). For detection of

the primary antibodies the membrane was incubated for one hour with goat anti-mouse

(α5, α-tubulin), anti-rabbit (β2) and anti-chicken (β1, β3) secondary antibodies conjugated

with horseradish peroxidase (Dianova, Hamburg, Germany) and developed with 4-chloro-

1-naphtol as previously described (Dalla et al., 2017).

Extraction and preparation of cardenolides

Calotropin was extracted according to (Petschenka et al., 2018) from approximately 8 g of

dried larvae of the monarch butterfly *Danaus plexippus* reared on *Asclepias curassavica*.

The cardenolides were extracted with methanol, lipids extracted with hexane and the

cardenolides retrieved by chloroform extraction. The concentrated methanolic extract

was subjected to preparative HPLC (Agilent 1100) equipped with a photodiode array

detector and a semi-preparative RP-Nucleodur C18 Htec column (5 µm, 250 x 10 mm,

Macherey-Nagel, Düren, Germany) using a water-acetonitrile gradient. Calotropin was

identified by its retention time and checked for molecular weight and LS-MS spectrum by

LC-ESI-TOF spectrometry by Dr. Maria Riedner (Department of Chemistry, Universität

Hamburg) and the structure confirmed by NMR spectroscopy. The resulting calotropin

preparation was 80% pure.

Calotropin and the commercially available ouabain (Sigma-Aldrich) were first dissolved in

DMSO and prepared for Na,K-ATPase activity assays as dilutions in sterile H_2O reaching

final concentrations from 10^{-8} to 10^{-3} M with 2% DMSO. Ouabain could in addition be

prepared as a 5×10^{-3} M solution.

Na,K-ATPase activity and inhibition assay

Activities of the nine different Na,K-ATPase α/β-subunit combinations of *O. fasciatus* were

determined by photometric measurements of inorganic phosphate released from

enzymatic ATP hydrolysis while subtracting background ATPase activity following

previously described methods (Dalla et al., 2017; Petschenka et al., 2013). Based on

previous results (Dalla et al., 2017) the buffer regime used was 50 mM NaCl, 4 mM $MgCl_2$,

50 mM imidazol, 20 mM KCl, pH 7.4. The inhibition of the different Na,K-ATPases was

tested under increasing ouabain and calotropin concentrations and expressed as

percentage activity of a non-inhibited control. For each α/β-subunit construct duplicates

of at least three biological replicates of recombinant proteins isolated from independently

transfected *Sf*9 cells were tested.

Analysis of data/ Statistics

After checking for normal distribution (Shapiro Wilk's test) and variance homogeneity

(Levene's test on squared deviations), differences in total Na,K-ATPase activities were

tested and compared between each α/β-subunit construct by one-way ANOVA followed

by Tukey's HSD posthoc test. The data of each Na,K-ATPase α/β-subunit construct was

subjected to nonlinear curve fitting with dose response function and Levenberg

Marquardt iteration algorithm in Origin-Pro 9.1G (OriginLab Corporation, Northampton,

USA) using a four parameter logistic function, top asymptote was set to 100, bottom

asymptote set to 0 and a variable Hill slope. For each construct IC_{50} values were calculated

and differences were analyzed separately for constructs affected by ouabain and

calotropin by one-way ANOVA followed by Tukey's HSD posthoc tests. A pairwise, F-test

based comparison of similarity between the fitted curves was performed using the

command "compare datasets". All calculations were performed with Origin-Pro 9.1G.

Results

Bioinformatic analysis

Four potential β-subunits were identified from the *O. fasciatus* transcriptome and

according to their similarity to published sequences named β1, β2, β3 and βx (Genbank

AccNo xxx-xxx). While β1-3 conform to the standard length of β-subunits of around 900

bp, βx is much shorter and missing most of the conserved N-terminal and half of the

transmembrane region (Fig. S1A). This β-subunit was therefore not further considered

here. The 5' end of α1B and α1C could successfully be recovered by RACE protocols (Fig.

S1B). The only minimally expressed α1D (Yang et al., 2019) was not further investigated

here.

Expression of α/β-subunit constructs in Sf9 cells

All nine α/β-subunit constructs (αAβ1, αAβ2, αAβ3, αBβ1, αBβ2, αAβ3, αCβ1, αCβ2, and

αCβ3) were successfully expressed in *Sf*9 cells and harvested. The western blot shows that

identical concentrations were loaded onto the gel (α-tubulin as positive control), all α-

subunits were expressed at similar levels and the deglycosylated β-subunits can be easily

discriminated by their specific patterns (Fig. 2).

Na,K-ATPase activity of different α/β-subunit constructs

The activities of different α/β-subunit combinations of the Na,K-ATPase were tested *in*

vitro to determine whether the amino acid substitutions in the cardenolide binding site or

the associated β-subunit influence the catalytic activity of the enzyme. All enzymes

showed catalytic activity, but the activity of the α/β-subunit constructs differed

significantly from one another (Fig. 3). The α-subunits A and B with their associated β-

subunits (1-3) showed significantly lower activity than αC combined with β1, 2 or 3. The

activities of αA and αB did not differ significantly from each other regardless of the β-

subunit, except for αAβ2. This combination was significantly less active than αB associated

with the three β-subunits (Fig. 3). By far the highest activity was measured for αC in

association with β3 that had an average turnover of 36.5 nmol Pi mg^{-1} Protein * minute

and differed significantly from all other constructs.

Influence of α/β-subunit association on ouabain sensitivity

In vitro tests with the cardenolide ouabain revealed large differences in sensitivity of the

Na,K-ATPases depending on the number of amino acid substitutions in the cardenolide

binding site and was modulated by the identity of the β-subunit (Fig. 4). Yet, when

comparing the IC$_{50}$ values only the one of αC associated with β1 was significantly lower

than those for the other two αC constructs (ANOVA, $F_{2,8}$=8.035, p=0.02, Tukey's HSD,

p=0.035 and p=0.027, for αC β1 vs β2 and β1 vs β3, respectively; Fig. 4c, Table 1) whereas

the IC$_{50}$ values of the three αA or αB constructs did not differ significantly.

Although analyzing differences in IC$_{50}$ is a standard approach, the IC$_{50}$ values only reflect

one point of the dose response curve. We therefore aimed to get a better insight into the

overall responses of the Na,K-ATPases to increasing ouabain concentrations by analyzing

the differences between the full dose response curves of constructs with the same α-

subunit as well as constructs with the same β-subunit (Table 2).

We found that α-subunits associated with different β-subunits behaved differently when

subjected to ouabain. The construct αAβ3 was significantly more sensitive to the

cardenolide than the other two αA complexes (Table 2, Fig. 4a), and αBβ1 was inhibited

more strongly by ouabain than αBβ2 (Table 2, Fig. 4b). Nevertheless, the reaction of

constructs with identical α-subunits was more uniform than the one of constructs with

identical β- but different α-subunits. Most of the latter constructs differed significantly in

their reactions, reflecting their varying insensitivity levels against ouabain (Fig. 4 a-c, Table

2).

Influence of α/β-subunit association on calotropin sensitivity

Testing the inhibitory effect of the host plant cardenolide calotropin on the recombinant

Na,K-ATPases resulted in total inhibition of the most calotropin-sensitive αC/β-subunit

constructs at the highest calotropin concentration. Furthermore, the three αB constructs

turned out to be most resistant against calotropin. As for the inhibition by ouabain only

the IC_{50} values of αC associated with the three different β-subunits differed significantly

from each other, αC associated with β3 was significantly more insensitive than αC

associated with β1 and β2 (ANOVA, $F_{2,8}$=18.554, p=0.003, Tukey's HSD, p=0.002 and

p=0.027, for αCβ3 versus αCβ1 and αCβ3 versus αCβ2, respectively; Fig. 4c).

When comparing the full dose response curves of the constructs (Table 2) neither the

three αA constructs differed significantly from one another nor the αB or αC constructs,

indicating a similar response to calotropin exposition independent of the associated β-

subunit (Fig. 4 a-c). In contrast, the inhibition by calotropin of the three α-subunits

combined with the same β-subunits was significantly different (from high to low

inhibition: αC> αA> αB; Fig. 4a-c). The curves of αAβ1 and αBβ1 did not differ significantly

from one another, since αAβ1 tends to be the most resistant combination of the three αA

constructs and αBβ1 the most sensitive of the three αB constructs (Fig. 4a & b).

Comparison of both cardenolides

Calotropin had a higher inhibitory potency and inhibited especially the three αC-

constructs to a much higher extent than ouabain (ANOVA, $F_{5,17}$=57.25882, p<0.001,

Tukey's HSD, p<0.001 for αC β1, αC β2 and αC β3, respectively; Fig. 4c, Table 1).

Calotropin inhibited αCβ1 18-fold, αCβ2 20-fold and αCβ3 14-fold more strongly than

ouabain did. Similarly, αB associated with β3 was significantly more inhibited by calotropin

than by ouabain, resulting in an approximately 24-fold higher IC_{50} (ANOVA, $F_{5,17}$=7.54487,

p=0.00205, Tukey's HSD, p=0.00357; Fig. 4b, Table 1) while the IC_{50} values of the two

cardenolides did not differ significantly for αBβ1 and αBβ2 (Tukey's HSD, p=0.70595 and

p=0.51146, respectively; Fig. 4b, Table 1).

Additionally, we compared the dose response curves of same constructs inhibited by

calotropin and ouabain. Besides similar results for the αC constructs, we found that the

construct αAβ1 was significantly more insensitive against calotropin than against ouabain,

reflected by a flat descending curve at the top (Fig. 4c & 4a, Table 3).

Table 1: Mean IC_{50} values and Hill slopes of different Na,K-ATPase α-β-constructs (3 biological replicates with

standard deviation (± s.d.)) obtained from dose response curves upon inhibition with ouabain and calotropin

(Origin-Pro 9.1G).

α/β-combination	IC_{50} values [M] Ouabain (± s.d.)	Hill slope (± s.d.)	IC_{50} values [M] Calotropin (± s.d.)	Hill slope (± s.d.)

αAβ1	4.148E-04 (± 6.541E-04)	-0.221 (± 0.047)	1.211E-04 (± 4.443E-05)	-0.271 (± 0.03)
αAβ2	3.471E-04 (± 3.598E-04)	-0.234 (± 0.013)	1.449E-05 (± 5.994E-06)	-0.269 (± 0.076)
αAβ3	2.607E-05 (± 2.767E-05)	-0.260 (± 0.044)	6.959E-05 (± 7.54E-05)	-0.27 (± 0.036)
αBβ1	2.089E-03 (± 2.515E-03)	-0.213 (± 0.039)	1.108E-04 (± 3.423E-05)	-0.293 (± 0.044)
αBβ2	2.777E-03 (± 9.676E-04)	-0.462 (± 0.055)	3.281E-04 (± 2.119E-04)	-0.398 (± 0.066)
αBβ3	7.09E-03 (± 3.119E-03)	-0.261 (± 0.045)	2.944E-04 (± 8.195E-05)	-0.254 (± 0.071)
αCβ1	4.274E-04 (± 8.104E-05)	-0.869 (± 0.071)	2.402E-05 (± 4.636E-06)	-0.88 (± 0.029)
αCβ2	7.369E-04 (± 1.001E-04)	-0.869 (± 0.093)	3.699E-05 (± 5.517E-06)	-1.289 (± 0.198)
αCβ3	7.579E-04 (± 1.476E-04)	-0.988 (± 0.35)	5.594E-05 (± 8.55E-06)	-0.912 (± 0.098)

319   Table 2: Comparison between the dose response curves of different α/β constructs inhibited by ouabain and

320   calotropin analyzed with Origin-Pro 9.1G. Significant differences are highlighted in bold.

Data comparison α/β-combinations	F-statistic Ouabain	F-statistic Calotropin
αAβ1 – αAβ2	$F_{2,8}$= 0.074, p= 0.929	$F_{2,8}$= 1.319, p= 0.320
αAβ1 – αAβ3	**$F_{2,8}$= 5.218, p= 0.035**	$F_{2,8}$= 0.728, p= 0.512
αAβ2 – αAβ3	**$F_{2,8}$= 6.092, p= 0.025**	$F_{2,8}$= 0.079, p= 0.925
αBβ1 – αBβ2	**$F_{2,8}$= 10.981, p= 0.005**	$F_{2,8}$= 2.607, p= 0.134
αBβ1 – αBβ3	$F_{2,8}$= 3.781, p= 0.070	$F_{2,8}$= 0.724, p= 0.514
αBβ2 – αBβ3	$F_{2,8}$= 0.455, p= 0.650	$F_{2,8}$= 0.655, p= 0.545
αCβ1 – αCβ2	$F_{2,8}$= 0.745, p= 0.505	$F_{2,8}$= 1.887, p= 0.213
αCβ1 – αCβ3	$F_{2,8}$= 1.535, p= 0.273	$F_{2,8}$= 3.402, p= 0.085
αCβ2 – αCβ3	$F_{2,8}$= 1.080, p= 0.384	$F_{2,8}$= 2.217, p= 0.171
αAβ1 – αBβ1	$F_{2,8}$= 1.039, p= 0.397	$F_{2,8}$= 0.554, p= 0.595
αAβ1 – αCβ1	**$F_{2,8}$= 22.729, p< 0.001**	**$F_{2,8}$= 8.724, p= 0.010**
αBβ1 – αCβ1	**$F_{2,8}$= 11.975, p= 0.004**	**$F_{2,8}$= 8.795, p= 0.010**
αAβ2 – αBβ2	**$F_{2,8}$= 57.311, p< 0.001**	**$F_{2,8}$= 9.763, p= 0.007**
αAβ2 – αCβ2	**$F_{2,8}$= 4.835, p= 0.042**	**$F_{2,8}$= 6.422, p= 0.022**
αBβ2 – αCβ2	$F_{2,8}$= 2.353, p= 0.157	**$F_{2,8}$= 13.589, p= 0.003**
αAβ3 – αBβ3	**$F_{2,8}$= 32.590, p< 0.001**	**$F_{2,8}$= 37.553, p< 0.001**
αAβ3 – αCβ3	**$F_{2,8}$= 33.295, p< 0.001**	**$F_{2,8}$= 19.271, p< 0.001**

αBβ3 − αCβ3	$F_{2,8}$= 1.614, p= 0.258	**$F_{2,8}$= 8.239, p= 0.011**

322 Table 3: Comparison between dose response curves of same α/β constructs inhibited by ouabain and

323 calotropin analyzed with Origin-Pro 9.1G. Significant results are highlighted in bold.

Data comparison α/β-combinations	F-statistic calotropin vs. ouabain
αAβ1	**$F_{2,8}$= 11.752, p= 0.004**
αAβ2	$F_{2,8}$= 2.421, p= 0.151
αAβ3	$F_{2,8}$= 0.589, p= 0.577
αBβ1	$F_{2,8}$= 0.343, p= 0.720
αBβ2	$F_{2,8}$= 4.257, p= 0.055
αBβ3	$F_{2,8}$= 3.054, p= 0.103
αCβ1	**$F_{2,8}$= 35.226, p< 0.001**
αCβ2	**$F_{2,8}$= 100.356, p< 0.001**
αCβ3	**$F_{2,8}$= 74.665, p< 0.001**

Discussion

Here we have identified the full length coding sequences of three Na,K-ATPase α- and

three full length β-subunits of *O. fasciatus* and functionally characterized all nine possible

recombinant α/β enzyme complexes. These nine functional enzymes strongly differed in

their activity levels. We found that enzyme complexes with αC had a significantly higher

activity compared to the other constructs. This perfectly fits with the fact that αC is mainly

expressed in the nervous tissue of *O. fasciatus* (Lohr et al., 2017; Zhen et al., 2012) where

the Na,K-ATPase plays a crucial role in maintaining electric potentials that guarantee the

excitability of neurons. Consequently, a high ionic pumping activity is indispensable. Two

of our nine α/β constructs clearly stand out amongst the other enzyme complexes: αAβ2

and αCβ3. The activity of αAβ2 is reduced in comparison to the other αA and αB

constructs that all show similar activity levels. Therefore, we suppose that αAβ2 is not a

functional enzyme *in vivo*. In contrast, the activity of αC combined with β3 is significantly

higher and would therefore well fit with a natural occurrence of this complex in the

nervous tissue. Data of an ongoing proteomic study support this assumption and show

that the subunits αC and β3 are most abundant in the nervous tissue of *O. fasciatus* (M.

Winter et al., unpublished data).

Regardless of the strikingly high overall resistance to inhibition by cardenolides that Na,K-

ATPase preparations of *O. fasciatus* display (Bramer et al., 2015; Dalla and Dobler, 2016;

Moore and Scudder, 1986) we found that the individual α/β-combinations strongly

differed in their resistance against the two cardenolides ouabain and calotropin. Overall

calotropin proved to be the much more potent inhibitor. This is despite the fact that *O.*

fasciatus is naturally exposed to calotropin in its host plants (Malcolm, 1991) and thus

could be specifically adapted to this toxin. Yet, our data demonstrate that in particular the

most active αC-subunit was roughly 20-times more strongly inhibited by calotropin than

by ouabain. A differential inhibition can also be observed for the three αB constructs while

the difference in inhibition between ouabain and calotropin is least pronounced for the

αA constructs. Overall, the αA- and αB-subunits are more insensitive to both cardenolides

than αC, which fits well with their abundant expression in the Malpighian tubules where

the enzymes are directly exposed to toxins in the hemolymph (Lohr et al., 2017). In

contrast, the least sensitive but most active αC is the dominant Na,K-ATPase α-subunit in

the nervous tissue where it is protected from intruding cardenolides by the perineurium

(blood-brain-barrier) which controls xenobiotica access to the nervous system by a tight

network of septate junctions and active efflux carriers (Dermauw and Van Leeuwen,

2014).

The difference in inhibitory potencies of the two cardenolides results from differences in

their structure, steric conformation and - *in vivo* - by their ability to permeate membranes,

as reflected by their lipophilicity. Ouabain is produced in plants of the genus *Acokanthera*

and in seeds of *Strophanthus gratus* (both Apocynaceae)*,* whereas calotropin is

synthesized in plants of the tribe Asclepiadeae (Apocynaceae). Cardiotonic activity is

determined by the steroidal aglycone, specifically by the conformation of the steroid rings.

The A/B and C/D rings of ouabain, and many other cardenolides, are *cis*-fused and the B/C

ring is *trans*-fused (*cis-trans-cis*) (Katz et al., 2010; Malcolm, 1991). In contrast calotropin

and other cardenolides produced in the genus *Asclepias,* have a *trans-trans-cis*

conformation (Malcolm, 1991) that is predicted to lower the toxicity (Hoch, 1961) and the

binding affinity of the cardenolide (Katz et al., 2010). However, a doubly-linked sugar

moiety, often found in cardenolides produced in plants of the tribe Asclepiadeae,

enhances the binding to the Na,K-ATPase as well as the toxicity (Piacente et al., 2009;

Venkateswara Rao and Vaidyanathan, 1991). Despite their differences in structure and

steric conformation, molecular docking analyses suggest that both cardenolides fit well

into the cardenolide binding site due to formation of strong H-bonds between the

interaction partners (Zhang et al., 2014). Similar to our results a study testing various

cardiac glycosides against the Na,K-ATPase of the monarch butterfly, *Danaus plexippus*,

(Petschenka et al., 2018) found calotropin and its isomer calactin to be the strongest

tested inhibitors, with calotropin inhibiting 60-fold more strongly than ouabain.

A comparison of the present results with our previous studies (Dalla and Dobler, 2016;

Dalla et al., 2017) on *O. fasciatus* Na,K-ATPase mimics reveals good concordance regarding

our conclusions about enzyme activities. We previously introduced one, two, three or four

of the potentially resistance-conferring substitutions into the α-subunit of *D.*

melanogaster to mimic the known α1-gene copies of *O. fasciatus* (in this order: αD, αC,

αB, αA-mimic; c.f. Yang et al., 2019; Zhen et al., 2012) and combined them with two *D.*

melanogaster β-subunits, Nrv2.2 and Nrv3 (Dalla and Dobler, 2016; Dalla et al., 2017). The

overall behavior of these mimics strongly resembled the present data for the original *O.*

fasciatus Na,K-ATPase complexes. Most strikingly, the activity levels of the αA and αB

mimics were significantly lower than of the Drosophila wildtype enzyme, the αC or the αD

(N122H) mimics. The two Drosophila β-subunits also modulated the activity: the

association with Nrv3 strongly reduced the activity of all mimics and the Drosophila

wildtype enzyme when compared to those bearing Nrv2.2. In the present study, such a

large influence of the β-subunit on ion pumping activity was observed only in the αC-

subunit while the effect was less pronounced for the αA and αB-subunits.

The β-subunits also strongly modulated the resistance of the enzymes against ouabain,

but their effect was specific to the α-subunits they were combined with. Previous work

has shown that a combination with Nrv3 instead of Nrv2.2 lead to lowered resistance in

the αA- and αB-mimic and increased resistance in the αC-mimic (Dalla et al., 2017). In the

present study, resistance changed in the same direction with β1, β2, or β3 for αA

(decreasing) and αC (increasing IC_{50}) but also lead to increased IC_{50}-values for αB.

The most striking differences, however, appear when comparing the overall ouabain-

sensitivity of the original enzyme combinations with the previously constructed mimics

(Dalla et al., 2017). While the mimics of the *O. fasciatus* αA-copy (featuring the

substitutions Q111T,N122H,F786N,T797A) were by far the most insensitive constructs

followed by the αB-mimics (with Q111T,N122H,F786N) and the more sensitive αC-mimics

(N122H,F786S), the original αA-constructs tested here are at least five times less sensitive

than αB-constructs. On the other hand, the original αC-constructs are 10-times more

resistant than predicted from the experiments with the αC-mimics.

These differences in sensitivity between mimics and original enzymes strongly point to the

importance of additional substitutions in the *O. fasciatus* Na,K-ATPase α1-paralogs. When

constructing the mimics, we focused on the four most conspicuous amino acid exchanges

in the binding pocket of the Na,K-ATPase that were previously shown functionally to be

essential for cardiac glycoside binding (Croyle et al., 1997; Holzinger et al., 1992; Qiu et al.,

2005) and confirmed to interact with cardiac glycosides by crystal structure analysis

(Laursen et al., 2015). However, there are additional amino acid substitutions in the

cardiac glycoside binding pocket of the Na,K-ATPase α1-genes of *O. fasciatus* that may

further increase their resistance (Dobler et al., 2012; Zhen et al., 2012). These and further

additional amino acid exchanges that at first appear inconsequential may be of

importance in modulating the effects of the resistance conferring substitutions or

ameliorating their pleiotropic effects (Karageorgi et al., 2019; Taverner et al., 2019;

Mohammadi et al., submitted).

Author contribution - S. Da. and S. Do. conceptualization; S. Da., M. W., V. W., R. T., M. H.

cloning and protein expression; S. Da. and M. W. enzyme assays, data analysis and

visualization; M. W. and S. Do. writing-original draft; S. Da. writing-review and editing.

The authors declare that they have no conflicts of interest with the contents of this article.

Acknowledgments - We acknowledge the help of Sebastian Gatemann who purified the

calotropin as part of his BSc thesis. Tammy Tischer supervised by Jennifer Lohr

complemented the 5' end of the αC copy. Shab Mohammadi improved the manuscript by

proofreading the English and valuable discussions. We thank Daniel Tal for generating the

structures in figure 1.

References

Agrawal, A. A., Petschenka, G., Bingham, R. A., Weber, M. G. and Rasmann, S. (2012).

Toxic cardenolides: chemical ecology and coevolution of specialized plant-herbivore

interactions. *New Phytol.* **194**, 28–45.

Baumann, O., Salvaterra, P. M. and Takeyasu, K. (2010). Developmental changes in β-

subunit composition of Na,K-ATPase in the *Drosophila* eye. *Cell Tissue Res.* 215–228.

Blanco, G. (2005). Na,K-ATPase subunit heterogeneity as a mechanism for tissue-specific

ion regulation. *Semin. Nephrol.* **25**, 292–303.

Blanco, G. and Mercer, R. W. (1998). Isozymes of the Na-K-ATPase: heterogeneity in

structure, diversity in function. *Am. J. Physiol. Ren. Physiol.* F633–F650.

Bramer, C., Dobler, S., Deckert, J. and Petschenka, G. (2015). Na+/K+-ATPase resistance

and cardenolide sequestration: basal adaptations to host plant toxins in the

milkweed bugs (Hemiptera: Lygaeidae: Lygaeinae). *Proc. R. Soc. B Biol. Sci.* 20142346.

Croyle, M. L., Woo, A. L. and Lingrel, J. B. (1997). Extensive random mutagenesis analysis

of the Na + / K + - ATPase a subunit identifies known and previously unidentified

amino acid residues that alter ouabain sensitivity implications for ouabain binding.

Eur. J. Biochem. **248**, 488–495.

Dalla, S. and Dobler, S. (2016). Gene duplications circumvent trade-offs in enzyme

function: insect adaptation to toxic host plants. *Evolution (N. Y).* **70**, 2767–2777.

Dalla, S., Baum, M. and Dobler, S. (2017). Substitutions in the cardenolide binding site

and interaction of subunits affect kinetics besides cardenolide sensitivity of insect

Na,K-ATPase. *Insect Biochem. Mol. Biol.* **89**, 43–50.

Dermauw, W. and Van Leeuwen, T. (2014). The ABC gene family in arthropods:

comparative genomics and role ininsecticide transport and resistance. *Insect*

Biochem. Mol. Biol. **45**, 89–110.

Dobler, S., Dalla, S., Wagschal, V. and Agrawal, A. A. (2012). Community-wide convergent

evolution in insect adaptation to toxic cardenolides by substitutions in the Na,K-

ATPase. *Proc. Natl. Acad. Sci.* **109**, 13040–13045.

Dobler, S., Wagschal, V., Pietsch, N., Dahdouli, N., Meinzer, F., Romey-Glüsing, R. and

Schütte, K. (2019). New ways to acquire resistance: imperfect convergence in insect

adaptations to a potent plant toxin. *Proc. R. Soc. B* **286**, 20190883.

Geering, K. (2001). The functional role of β subunits in oligomeric P-type ATPases. *J.*

Bioenerg. Biomembr. **33**, 425–438.

Geering, K. (2005). Function of FXYD proteins, regulators of Na,K-ATPase. *J. Bioenerg.*

Biomembr. **37**, 387–392.

Grabherr, M. G. ., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I.,

Adiconis, X., Fan, L., Raychowdhury, R., Zeng, Q., et al. (2011). Trinity: reconstructing

a full-length transcriptome without a genome from RNA-Seq data. *Nat. Biotechnol.*

29, 644–652.

Hoch, J. H. (1961). *A survey of cardiac glycosides and genins*. Columbia, SC, USA:

University of South Carolina Press.

Holzinger, F., Frick, C. and Wink, M. (1992). Molecular basis for the insensitivity of the

Monarch (*Danaus plexippus*) to cardiac glycosides. *FEBS Lett.* **314**, 477–480.

Horisberger, J.-D. (2004). Recent Insights into the structure and mechanism of the sodium

pump. *Physiology* **19**, 377–387.

Karageorgi, M., Groen, S. C., Sumbul, F., Pelaez, J. N., Verster, K. I., Aguilar, J. M.,

Hastings, A. P., Bernstein, S. L., Matsunaga, T., Astourian, M., et al. (2019). Genome

editing retraces the evolution of toxin resistance in the monarch butterfly. *Nature*

574, 409–412.

Katz, A., Lifshitz, Y., Bab-Dinitz, E., Kapri-Pardes, E., Goldshleger, R., Tal, D. M. and

Karlish, S. J. D. (2010). Selectivity of digitalis glycosides for isoforms of human Na,K-

ATPase. *J. Biol. Chem.* **285**, 19582–19592.

Laursen, M., Gregersen, J. L., Yatime, L., Nissen, P. and Fedosova, N. U. (2015). Structures

and characterization of digoxin- and bufalin-bound Na^+,K^+-ATPase compared with

the ouabain-bound complex. *Proc. Natl. Acad. Sci.* **112**, 1755–1760.

Lebovitz, R. M., Takeyasu, K. and Fambrough, D. M. (1989). Molecular characterization
and expression of the (Na+ + K+)-ATPase alpha-subunit in *Drosophila melanogaster*.
EMBO J. **8**, 193–202.

Lohr, J. N., Meinzer, F., Dalla, S., Romey-Glüsing, R. and Dobler, S. (2017). The function
and evolutionary significance of a triplicated Na,K-ATPase gene in a toxin-specialized
insect. *BMC Evol. Biol.* **17**, 256.

Malcolm, S. B. (1991). Cardenolide-mediated interactions between plants and herbivores.
In *Herbivores: their Interactions with Secondary Plant Metabolites* (ed. Rosenthal, G.
A.) and Berenbaum, M. R.), pp. 251–296. San Diego, CA: Academic Press.

Meinzer, F., Dobler, S., Donath, A. and Lohr, J. N. (2019). Robust reference gene design
and validation for expression studies in the large milkweed bug, *Oncopeltus fasciatus*,
upon cardiac glycoside stress. *Gene* **710**, 66–75.

Mohammadi, S., Gompert, Z., Gonzalez, J., Takeuchi, H., Mori, A. and Savitzky, A. H.
(2016). Toxin-resistant isoforms of Na+/K+-ATPase in snakes do not closely track
dietary specialization on toads. *Proc. R. Soc. B* **283**, 20162111.

Mohammadi, S., Yang, L., Harpak, A., Herrera-Alvarez, S., Rodriguez-Ordonez, M.d.P.,
Peng, J., Zhang, K., Storz, J.F., Dobler, S., Crawford, A. J., Andolfatto, P. Concerted
evolution reveals co-adapted amino acid substitutions in frogs that prey on toxic
toads. submitted.

Moore, L. V. and Scudder, G. G. E. (1986). Ouabain-resistant Na,K-ATPases and
cardenolide tolerance in the large milkweed bug, *Oncopeltus fasciatus*. *J. Insect*
Physiol. **32**, 27–33.

Okamura, H., Yasuhara, J. C., Fambrough, D. M. and Takeyasu, K. (2002). P-type ATPases
in *Caenorhabditis* and *Drosophila*: implications for evolution of the P-type ATPase
subunit families with special reference to the Na,K-ATPase and H,K-ATPase subgroup.
J. Membr. Biol. **191**, 13–24.

Palasis, M., Kuntzweiler, T. A., Argüello, J. M. and Lingrel, J. B. (1996). Ouabain

Interactions with the H5-H6 Hairpin of the Na,K-ATPase Reveal a Possible Inhibition

Mechanism via the Cation Binding Domain. *J. Biol. Chem.* **271**, 14176–14182.

Paul, S. M., Palladino, M. J. and Beitel, G. J. (2007). A pump-independent function of the

Na,K-ATPase is required for epithelial junction function and tracheal tube-size

control. *Development* **134**, 147–55.

Petschenka, G., Fandrich, S., Sander, N., Wagschal, V., Boppré, M. and Dobler, S. (2013).

Stepwise evolution of resistance to toxic cardenolides via genetic substitutions in the

Na+/K+-ATPase of milkweed butterflies (lepidoptera: Danaini). *Evolution (N. Y).* **67**,

2753–2761.

Petschenka, G., Wagscha, V., Von Tschirnhaus, M., Donath, A. and Dobler, S. (2017).

Convergently evolved toxic secondary metabolites in plants drive the parallel

molecular evolution of insect resistance. *Am. Nat.* **190**, S29–S43.

Petschenka, G., Fei, C. S., Araya, J. J., Schröder, S., Timmermann, B. N. and Agrawal, A. A.

(2018). Relative selectivity of plant xardenolides for Na+/K+-ATPases from the

monarch butterfly and non-resistant insects. *Front. Plant Sci.* **9**, 1424.

Piacente, S., Masullo, M., De Nève, N., Dewelle, J., Hamed, A., Kiss, R. and Mijatovic, T.

(2009). Cardenolides from *Pergularia tomentosa* display cytotoxic activity resulting

from their potent inhibition of Na+/K+-ATPase. *J. Nat. Prod.* **72**, 1087–1091.

Price, E. M. and Lingrel, J. B. (1988). Structure - function relationships in the Na,K-ATPase

α subunit: site-directed mutagenesis of glutamine-111 to arginine and asparagine-

122 to aspartic acid generates a ouabain-resistant enzyme. *Biochemistry* **27**, 8400–

8408.

Qiu, L. Y., Krieger, E., Schaftenaar, G., Swarts, H. G. P., Willems, P. H. G. M., De Pont, J. J.

H. H. M. and Koenderink, J. B. (2005). Reconstruction of the complete ouabain-

binding pocket of Na,K-ATPase in gastric H,K-ATPase by substitution of only seven

amino acids. *J. Biol. Chem.* **280**, 32349–32355.

Taverner, A. M., Yang, L., Barile, Z. J., Lin, B., Peng, J., Pinharanda, A. P., Rao, A. S.,

Roland, B. P., Talsma, A. D., Wei, D., et al. (2019). Adaptive substitutions underlying
cardiac glycoside insensitivity in insects exhibit epistasis in vivo. *Elife* 48224.

Ujvari, B., Casewell, N. R., Sunagar, K., Arbuckle, K., Wüster, W., Lo, N., O'Meally, D.,
Beckmann, C., King, G. F., Deplazes, E., et al. (2015). Widespread convergence in
toxin resistance by predictable molecular evolution. *Proc. Natl. Acad. Sci.* **112**,
11911–11916.

Venkateswara Rao, R. and Vaidyanathan, C. S. (1991). Chemistry and biochemical
pharmacology of cardiac glycosides - A review. *J. Indian Inst. Sci.* **71**, 329–364.

Yang, L., Ravikanthachari, N., Mariño-Pérez, R., Deshmukh, R., Wu, M., Rosenstein, A.,
Kunte, K., Song, H. and Andolfatto, P. (2019). Predictability in the evolution of
Orthopteran cardenolide insensitivity. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.*
20180246.

Yasuhara, J. C., Baumann, O. and Takeyasu, K. (2000). Localization of Na/K-ATPase in
developing and adult *Drosophila melanogaster* photoreceptors. *Cell Tissue Res.* 239–
249.

Zhang, R.-R., Tian, H.-Y., Tan, Y.-F., Chung, T.-Y., Sun, X.-H., Xia, X., Ye, W.-C., Middleton,
D. A., Fedosova, N., Esmann, M., et al. (2014). Structures, chemotaxonomic
significance, cytotoxic and Na + ,K + -ATPase inhibitory activities of new cardenolides
from *Asclepias curassavica*. *Org. Biomol. Chem.* **12**, 8919–8929.

Zhen, Y., Aardema, M. L., Medina, E. M., Schumer, M. and Andolfatto, P. (2012). Parallel
molecular evolution in an herbivore community. *Science (80-.).* **337**, 1634–1637.

Figure 1: Structure of the cardenolides ouabain (a) and calotropin (b). Both are

characterized by a monosaccharide attached to a steroid skeleton at C3 (4 steroid rings: A-

D) and a five-membered unsaturated lactone ring at C-17. a) In ouabain the

monosaccharide is attached by a single glycosidic link to the steroid rings that are *cis-*

trans-cis fused. b) Calotropin is characterized by a doubly-linked monosaccharide as well

as a *trans-trans-cis* conformation of the steroid rings (structures generated by Daniel Tal,

The Weizmann Institute of Science, 2020).

Figure 2: Western blot of nine α/β constructs of *O. fasciatus* expressed heterologously in cell culture. 30 µg of protein were separated by SDS-PAGE and blotted on a nitrocellulose membrane. The upper membrane slice shows at a size of 110 kDa αA, αB and αC in alternating order. At 55 kDa α-tubulin is shown as a positive control. Both proteins were stained with the primary antibody α5 (DSHB hybridoma bank) and the secondary HRP-conjugated antibody goat-anti mouse (Dianova). The associated β-subunits 1, 2 and 3 were deglycosylated in advance. They were stained with β-subunit specific customized primary antibodies (produced by Davids Biotechnology) and HRP-conjugated secondary antibodies goat-anti rabbit (for β2, Dianova) and goat-anti chicken (for β1 and 3, Dianova). The different β-subunits show bands at weights ranging between 35 and 40 kDa. The bands were visualized with 4-CN. Due to the grey scale some bands of the multicolor protein ladder were marked by lines.

Figure 3: Na,K-ATPase activity [nmol Pi/mg protein*min] of nine α/β-constructs of *O.*

fasciatus. Each bar depicts the mean and standard deviation of three biological replicates.

Different letters above the bars indicate significant differences between constructs

(ANOVA, $F_{8,28}$=63.932, p<0.001). Black bars: αA with three β-subunits and overall activities

ranging from **7 to 12 nmol Pi/mg protein*min**. Bars with a grey dotted outline: data for

the αA mimic (THNA) combined with Nrv2.2 and Nrv3 of a previous study (24). Dark gray

bars: αB/β constructs ranging from **13 to 14.3 nmol Pi/mg protein*min**. Light gray bars:

αC/β constructs with activity levels ranging from **22 to 36.5 nmol Pi/mg protein*min**. The

activity levels of the three αC/β constructs are significantly higher than those of the other

constructs, with the highest activity reached by αC combined with β3.

Figure 4: ***In vitro*** **inhibition by calotropin and ouabain of nine α/β constructs of** ***O.*** ***fasciatus*** **expressed heterologously in cell culture. Dose response relationships between** **increasing cardenolide concentrations and residual enzyme activities were expressed in** **% of the non-inhibited control (calculated with Origin Pro 9.1G).** Each data point represent the mean of three biological replicates and standard deviation. a) effect of calotropin (blue curves and round symbols) and ouabain (black curves and square symbols) on αA, b) on αB, and c) on αC combined each with β1, 2 and 3. Please note the different scales.