Identification of Dysregulated Pathways and key genes in Human Retinal Angiogenesis using Microarray Metadata

Umadevi Subramanian [1] and Bharanidharan Devarajan [1,*]

[1] Department of Bioinformatics, Aravind Medical Research Foundation, Madurai, India.

*Corresponding authorE-mail: bharanid@gmail.com

Abstract

Retinal angiogenesis is a common neovascularization mechanism that causes severe irreversible vision loss in the number of retinal diseases worldwide. Patients often do not respond to the current anti-angiogenic therapies and have a vision loss. Understanding the various angiogenic pathways and factors involved in the pathogenic mechanism is vital for disease management. In this study, to identify dysregulated angiogenic pathways and specific angiogenic factors involved in vision-threatening diseases namely proliferative diabetic retinopathy (PDR), retinopathy of prematurity (ROP) and neovascular age-related macular degeneration (nAMD), we downloaded microarray metadata of samples and obtained the differentially expressed genes (DEGs) in all the disease and each disease samples compared to controls. Subsequently, we performed Gene Set Enrichment (GESA) analysis for pathways, a protein-protein interaction (PPI), and angiome network analysis using R and Cytoscape software. We identified highly enriched dysregulated pathways that were neuroactive ligand receptor interaction and cytokine-cytokine receptor interaction. The angiogenic–associated DEGs were predominately related to the cytokine-cytokine receptor interaction pathway, which we further confirmed with RNA-seq data of PDR samples. Together, our analysis of these data elucidated the molecular mechanisms of retinal angiogenesis and provided potential angiogenic targets for therapeutics.

Keywords: Retinal angiogenesis, Microarray-Meta data, Dysregulated angiogenic pathways, Cytokine-cytokine receptor interaction pathway

1. Introduction

Retinal angiogenesis is the common dysregulated pathway of numerous blinding disorders, namely, proliferative diabetic retinopathy (PDR), retinal artery (or vein) occlusion, retinopathy of prematurity (ROP), sickle cell retinopathy[1], and neovascular age-related macular degeneration (nAMD)[2]. Although there are subtle differences in the mechanism of angiogenesis, they are characterized as common neovascularization event. Angiogenesis is a physiological process through which new blood vessels formed from pre-existing vessels under hypoxic conditions[3].Hypoxia-inducible-1α drives the expression of substantial pro-angiogenic growth factors and induces endothelial cells into a tip cell phenotype. The tip cell proliferates slowly and activates matrix metalloproteinases (MMPs), plasmin, uPA, and tPA, and cleaving numerous extracellular matrix (ECM) proteins. The activated MMPs and other proteins stimulate the Notch ligand in tip cells and proliferate further to elongate a new vessel[4,5]. As the newly formed vessels mainly serve a role in a wound healing response, they usually do not restore tissue integrity, but instead, cause visual impairment[6,7].

Vascular endothelial growth factor (VEGF) is the major growth factor, induced in hypoxic conditions. VEGF family members VEGF-A, VEGF-B, VEGF-C, VEGF-D and placental growth factor (PIGF), and their receptors are the validated regulators of angiogenic signaling pathway[8]. VEGF-A is the potent growth factor and mediator of intraocular neovascularization. VEGF-A stimulates not only the development of new blood vessels, but also, microaneurysm formation, capillary occlusion with ischemia, and promoting increased vascular permeability[9]. VEGF-B can regulate the function of endothelial cells. VEGF-C and VEGF-D increase vascular permeability as well as induce angiogenesis. PIGF plays a role in pathogenic angiogenesis by increasing the activity of VEGF-A[10].

The current anti-angiogenic agents, approved by the Food and Drug Administration, inhibit the VEGF pathway[8]. The anti-VEGF drugs are the molecules that target VEGF isoforms, or inhibiting VEGF receptors, or inhibiting VEGF downstream signaling[11]. About 30% of PDR patients fail to respond to initial treatment, and the majority of the responders would require multiple rounds of intravitreal injections[12]. Therefore, anti-VEGF treatment is potentially required for years, and chronic VEGF inhibition may cause side effects and drug resistance. The potential consequence of VEGF blockade is hypertension, proteinuria, impairment of wound healing/collateral vessel development, and inhibition of bone growth, infertility, inhibition of skeletal muscle, regeneration, cardiac remodeling and neuroprotective effect in the ischaemic retina[13]. Also, a dose-dependent decrease in ganglion cells was reported in rats[14].

Studies show that VEGF independent factors may stimulate angiogenesis, particularly in

tumors[15]. Zhang et al. suggested identifying the angiogenic factors other than VEGF for a

better therapeutic approach[16]. Gene expression studies have been widely performed to

identify angiogenic factors. However, such studies will provide numerous key genes that are

differentially expressed, and thus, it is a challenging task to detect reliable targets. On the

molecular level, a combination of the differential gene expression (which may vary in each

individual) is assumed to dysregulate the common cellular pathway[17]. Therefore,

identification of the dysregulated pathways is essential for understanding disease

mechanisms and the identification of targets for future application of custom therapeutic

decisions[18]. The accumulation of large amounts of gene expression data in public databases

provides an opportunity for identifying commonly dysregulated pathways and the gene

targets that are differentially expressed[19].

In the present study, we aim to identify the dysregulated angiogenic pathways in addition

to VEGF dependent angiogenic pathway in retinal diseases. To achieve our aim, we designed

this study with several steps that are (i) to combine the microarray data from different reliable

experiments and adjusting batch effects, (ii) to identify robust differentially expressed genes

(DEGs), (iii) to conduct the pathway enrichment analysis, (iv) to identify key dysregulated

angiogenic pathways and angiogenic factors through network analysis and (v) to compare

and confirm the identified dysregulated angiogenic pathways and angiogenic factors with

available RNA-Seq data

2. Materials and Methods

2.1. Gene expression data and preprocessing

All the microarray samples used in this study were systematically searched and

downloaded from NCBI-GEO[20] after the manual curation of the sample details. Probe IDs

were mapped to official gene symbols in the DAVID database[21], and non-protein coding

genes, as defined by an assignment of HUGO gene symbol, were excluded from the study. In

case of genes with multiple probes, gene expression level was aggregated from probeset to

gene-level data by taking the maximum value across all the samples[22].

2.2. Missing value imputation and normalization of data

All the samples selected for the study were combined by their gene symbol using R

script. The missing gene expression values were imputed by Bayesian Principal Component

Analysis (BPCA) method[23], which performed well even with low entropy. Further, the whole

data were quantile normalized[24] to make the distribution of values in all the samples into the

same. The outlier samples were removed based on the principal component analysis (PCA)

plot. Further it was confirmed with Euclidean distance-based clustering using R script.

2.3. Identification of differentially expressed genes (DEGs) and pathway enrichment

DEGs were selected from the list by filtering, only if the moderated t-test FDR was ≤

0.05 and the fold change (FC - a difference of the means between two groups when all the

values were log2 transformed) ≥ 2 (up-regulated) or ≤ -2 (down-regulated). For the pathway

analysis, Gene Set Enrichment Analysis (GSEA) proposed by Subramanian et al.

implemented with GenePattern tools was performed[25]. The DEGs and FC values were

regarded as a pre-ranked list, and c2.cp.kegg.v6.1 was interrogated as pathway database. We

used default parameters and set the minimum gene set size 2. The output enrichment score

reflects the degree to which a gene set in the pathway is over-represented at the extreme (low

or high) of the entire ranked list of FC. For the pathway enrichment, normalized enrichment

score (NES) was considered as a metric of influence of DEGs with p-value ≤ 0.05.

Analysis was carried out with different perspectives, differential gene expression and

pathway regulation between retinal angiogenesis (case) samples and controls, case and tissue

(retina, and RPE/C) specific controls, cases and disease (dry AMD, non-Proliferative

Diabetic Retinopathy - NPDR, and Diabetic Mellitus - DM) specific control samples, disease

specificity (nAMD, PDR, ROP) and control samples as given in figure 1. Among the DEGs,

the angiogenesis-related genes were identified by GO term in GSEA and the list is given by

Chu et al.[26]. Further, the STRING database was used to retrieve the predicted interactions for

the identified DEGs and graphically represented using Cytoscape 3.2.0.

2.4. RNA-Seq analysis

The available RNA-Seq samples related to this study in NCBI-SRA are PDR (Accession

No.: SRP097696) and whole retinal (SRP119766) control samples. They were downloaded

by SRA Toolkit, aligned to human reference genome hg38 using CLC Genomics Workbench

12.0, quantile normalized in R, and analyzed using EdgeR.

3. Results

3.1. Standardization of gene expression data

The analysis was primarily done with 243 control samples, 95 normal retinal control

(GSE60436 (3 samples), GSE28133 (19), GSE29801 (55), GSE32614 (12), GSE41019 (3),

GSE57864 (3)), 138 normal RPE/C control (GSE18811 (31), GSE29801 (96), GSE43257

(2), GSE50195 (7), GSE5741 (2)), and 32 disease cases includes, 16 (8 retina, and 8 RPE/C)

nAMD (Accession No.: GSE29801), 6 fibrovascular membrane of PDR (GSE60436), and 10

retinal microvascular endothelial cells of ROP (GSE5946) samples. The studies with disease

specific control samples include, 64 dry AMD retina (GSE29801 (61), GSE17938 (3)), 9 dry

AMD RPE/C (GSE50195), 6 neural retinal NPDR (GSE53257) and 6 neural retinal DM

(GSE53257) samples. After preprocessing and merging of data, 15.75% of missing data were

imputed. All the samples which contain 18621 genes were quantile normalized (Figure S1).

The 318 control samples (95 retina, 138 RPE/C) used in the primary study were obtained

from different technical platforms. Eventhough they were quantile normalized, to reduce the

bias the outlier samples have to be removed. In order to this, all the disease samples and

control samples were subjected to PCA (Figure S2). The disease samples were retained, the

control samples which were scatterd/ outgrouped were removed, and clustering tree was

construted with remaining samples. This process was iteratively done once observed the

clear clustering of case and control samples (Figure S3). This results to 31 retina and 34

RPE/C control samples.

3.2. Dysregulated pathways of all the cases with all normal control samples

One thousand three hundred eighty-three genes displayed differential expression with

two-fold change (FDR ≤ 0.05) in all the cases compared to normal control samples (Figure

S4). Of these, 852 were downregulated and 531 upregulated. Gene set enrichment analysis

(GSEA) with respect to pathways showed that 237 DEGs were associated with 22 different

dysregulated pathways (significant with p ≤ 0.05, NES of pathways are given in Table 1). We

then analyzed DEGs that are associated with angiogenesis. We identified 127

angiogenic-associated enriched DEGs, of which 31 were based on gene ontology (GO)

(NES: 1.74 with p = 0.01) and 96 DEGs were based on the literature. The highly enriched

pathways with their significantly altered DEGs and their association with angiogenesis were

graphically represented in Figure 2.

The neuroactive ligand receptor interaction pathway was the highly dysregulated

pathway with the normalized enrichment score (NES) of 3.15, and 15 DEGs were identified

in this pathway. Of these, 11 DEGs were unique to this pathway and while, 4 DEGs GRIN1,

CHRM2, GRIN2C and GRIN2D were connecting other pathways. Only UTS2R was found

to be angiogenic-associated (Figure 2).

3.2.2. Cytokine-cytokine receptor interaction pathway

Cytokine-cytokine receptor interaction pathway, next to the neuroactive ligand receptor

interaction pathway, had enrichment score 2.55 (Table 1). We identified 25 DEGs (Figure 2

and S5) in this pathway. Among these, down-regulated VEGFA, CCL3, EGFR, PDGFC and

up-regulated genes VEGFB, VEGFC, CX3CL1, ACVRL1, CCL23, CCL24, CCL26, OSM,

and TNFRSF25 genes were found to be angiogenic-associated. Of interest, we identified that

three of the angiogenic-associated genes PDGFC, CXCL1, and EGFR were also involved in

the other pathways, as shown in Figures 2 and S5.

3.2.3. Other pathways with DEGs of angiogenic association

In addition to highly enriched cytokine-cytokine receptor interaction and Neuroactive

ligand receptor interaction pathways, we showed 13 other pathways with

angiogenic-associated DEGs. Regulation of the actin cytoskeleton pathway has ten

angiogenic DEGs; Adherens junction and Epithelial cell signaling in Helicobacter pylori

infection have seven angiogenic DEGs; tight junction has six angiogenic DEGs. Three

angiogenic DEGs were involved in Alzheimer's disease and arrhythmogenic right ventricular

cardiomyopathy ARVC pathways and 2 DEGs were in endocytosis and pathogenic

Escherichia coli infection pathways. Five pathways had one DEG, namely, aminoacyl tRNA

biosynthesis (YARS), Huntington's disease (BAX), p53 signaling pathway (BAX), ribosome

(RPL29) and Vibrio cholerae infection (GNAS).

The angiogenic-associated DEGs that were involved in two enriched pathways include

ADAM10, BAX, CXCL1, ITGAV, JAM3, NRAS, and PDGFC. CTNNA1, MAPK1, and

RAC1 were involved in three enriched pathways; RHOA in four pathways; EGFR and

CDC42 were involved in more than five pathways. Despite the above, 92

angiogenic-associated DEGs were not involved in any of the enriched pathways. For

instance, we identified the SEMA3B gene with high fold change of 5.154 and not involved in

any of the enriched pathways.

3.2.4. Angiogenic interactome of DEGs through protein-protein interaction (PPI)

In order to identify the interactome of angiogenic-associated DEGs, their protein-protein

interaction (PPI) partners and pathways associated with them are connected as network and

graphically illustrated in Figure 3. PPP2CA showed the highest connectivity, which is

associated with the tight junction pathway, and it mostly connected to the ribosome pathway.

Followed by PPP2CA, RHOA and EGFR showed more interacting partners and pathways.

3.2.5 Pathway specificity analysis

To examine the specificity of enriched pathways (Table 1), we performed a similar

analysis using two sets of control samples, one from the retina and one from RPE/Choroid

samples (Figure 1). Compare to enriched pathways obtained using all control samples; we

identified that almost all the enriched pathways were enriched either using retina or

RPE/Choroid or both except Cysteine and methionine metabolism and Vibrio cholerae

infection. However, there are other pathways were enriched in the two control data sets that

were not enriched in all control data as shown in Supplementary Table S1. Nevertheless, the

enriched pathways, as shown in Table 1 is not affected by the two different retina and
RPE/Choroid control data sets, not random, although they influence the pathway enrichment
analysis.

3.2.6 Disease-specific analysis

Furthermore, we performed disease-specific pathway enrichment analysis whether a
single disease influences the angiogenic-associated pathways and genes. The heatmap of the
dysregulated pathways with respect to disease specificity is illustrated in Figure 4. The highly
enriched pathways Neuroactive ligand receptor interaction pathway and cytokine-cytokine
receptor interaction pathway was not affected by the specific disease. The cytokine-cytokine
receptor interaction pathway with highly angiogenic-associated DEGs was graphically
illustrated as a heat map in Figure 5. Many angiogenic-associated genes (mentioned
previously) showed similar patterns of fold change in all three retinal diseases. For example,
VEGFA gene was downregulated in three all retinal diseases individually and all three
together. We also performed the disease-specific comparative analysis using different
control data set (Supplementary Table S2). When PDR samples compared with NPDR and
DM samples, we observed the almost same pattern of dysregulated pathways. However, we
did not observe similar pattern when we compared dry AMD with AMD samples.

3.3. Pathway analysis using RNA-seq data

We further extended our analysis with the RNA-sequencing data set in order to show
whether the pathway enrichment analysis is biased with microarray data. We compared nine
PDR samples with 13 normal whole retina samples, and we identified 5816 upregulated and
2031 downregulated genes. The supplementary table 3 shows the dysregulated pathways
with the NES score. The graft versus host disease pathway was enriched with the highest
NES (4.15); however, it did not include the angiogenic DEGs. As expected, the
Cytokine-cytokine receptor interaction pathway with NES of 3.35 is the most dysregulated in
the retinal angiogenesis included with seventeen angiogenic-associated DEG. Among the
DEGs, ACVRL1, CCL19, CCL23, CXCL1, IL10RA, IL18RAP, IL1R2, IL3RA, IL4R,
OSM, TNFRSF25 and TNFRSF4 were observed with similar expression as that of the results
from microarray data analysis. Notably, of the VEGF family, only VEGFC was observed
with significant upregulation with 1.3 fold change.

4. Discussion

Neovascularization by dysregulation of angiogenic pathways is the major cause of vision
loss in many retinal diseases. The purpose of this study is to identify dysregulated angiogenic
pathways in addition to VEGF dependent angiogenic pathways and specific angiogenic

factors involved in vision-threatening retinal diseases, include PDR, ROP and nAMD. We

thought that the most feasible approach to this end would be to combine all the expression

data from retinal diseases with angiogenesis, to identify common dysregulated angiogenic

pathways. We showed that VEGF independent pathways are highly enriched by the

differentially expressed genes (DEGs) of the retinal angiogenesis samples. The two highly

enriched pathways are neuroactive ligand receptor interaction and cytokine-cytokine

receptor interaction pathways (Figure 4). Among the several DEGs involved in neuroactive

ligand receptor interaction pathway, though our angiome-network showed no

angiogenic-associated genes, ADRA1B [27], CHRM2[28], DRD4[29], GALR3[30], GLP2R[31],

GRIN2C[32], GRIN2D[33], HTR5A[34], P2RY2[35], UTS2R[36] DEGs have been reported to be

involved in angiogenesis. The neuroactive ligand receptor interaction pathway has been

reported to be the most significant pathway for cancers and angiogenesis and is also closely

connected with the poor prognosis of cancer[37–41].

Based on the angiome-network with enriched DEGs, we report that the

cytokine-cytokine receptor interaction pathway is the highly dysregulated angiogenic

pathway enriched with fifteen angiogenic-associated DEGs (Supplementary Figure S5)

followed by regulation of actin cytoskeleton, adherens junction and epithelial cell signaling

in *Helicobacter pylori* infection. Jennifer et al. [42] experimentally identified novel

VEGF-independent cytokines and their association with PDR, suggesting the association of

cytokines in angiogenesis. In this study, we have shown that ten angiogenic-associated

cytokines were significantly up-regulated in the dysregulated cytokine-cytokine receptor

interaction pathway. Interestingly, VEGFA along with other angiogenic factors CCL3,

EGFR and PDGFC are significantly down-regulated, suggesting that VEGF independent

cytokines play a vital role in retinal angiogenesis. Apart from the ten up-regulated

angiogenic-associated genes, with widespread literature survey, we account that CCL19[43],

IL18RAP[44], IL1R2[45], LTB[46], IL10RA[47], CNTF[48] up-regulated genes may contribute to the

dysregulation of angiogenesis. Therefore, we report that cytokine-cytokine receptor

interaction pathway is the common dysregulated pathway in retinal angiogenesis as

previously stated in the animal models of retinal angiogenesis[49].

Of interest, to support that cytokine-cytokine receptor interaction pathway as common

dysregulated angiogenic pathways in retinal diseases, we constructed a regulatory pathway

(figure 6) with observed DEGs in this pathway and literature support. First, we exhibit that

the VEGF pathway is downregulated by showing the downregulation of VEGFA gene

expression (Figures 5 and 6) and upregulation of CCL19 with CCR7 in retinal angiogenesis.

The CCL19 and CCR7 have been reported to regulate the VEGF pathway negatively[50–52].

Additionally, the DEGs CCL3[53] and EGFR[54] are reported to induce VEGFA expression,

which is downregulated in this study. We observe the downregulation of other DEGs c-Met,

RAS, ERK, Elk1 and STAT3 that could affect the VEGFA dependent angiogenesis[52]. The

upregulated DEGs ITFRSF25, TNFRSF4, CCL23, CCL26, and CX3CL1 of

cytokine-cytokine interaction pathway is reportedly activating PI3K/AKT pathway[55–59],

which in turn induces the angiogenesis through activating mTOR[54] and NFKB pathway[60].

The upregulated DEGs IL18RAP and CXCL1 has shown to modulates the NFKB and thus

induces the angiogenesis. Furthermore, the upregulated DEG ACVRL1 is reported to

promote endothelial cell proliferation by activating SMAD1[61] and coexpression of IL12 with

CCL19 reported to regulates the JAK/STAT signaling[62], may help to mediate the

angiogenesis indirectly[61–63]. We also observe that the other dysregulated pathways (Table 1)

include aminoacyl tRNA biosynthesis, pathogenic Escherichia coli infection signaling,

epithelial cell signaling in Helicobacter pylori infection and cysteine and methionine

metabolism pathway, reported to trigger VEGF dependent angiogenic pathway[64–66], are

negatively enriched. We also support the fact that VEGF-independent angiogenic pathway in

retinal angiogenesis using the PPI network. Most of the PPI partners of VEGFA namely

EGF, FLT1, HIF1A, IGF1, KDR, PIK3CA, TGFB1, THBS1 are downregulated in this study.

However, we have observed that the upregulation of VEGFB and VEGFC. However,

comparing the expression VEGFB and VEGFC in each disease and results of RNA-seq

analysis, we have identified that VEGFC expression is consistent, individually significant

upregulation was observed in PDR (more than two FC).

Further, to identify the specific angiogenic factors that are related to the

cytokine-cytokine receptor interaction pathway, we have examined the DEGs in the

disease-specific data. The fold change pattern of DEGs related to cytokine-cytokine receptor

interaction pathway is following the same pattern. However, among DEGs, CCL19, CCL26,

CCL24, CXCL1, ACVRL1, OSM, IL4R, CCL23and IL18RAP are significantly upregulated

genes; while, VEGFA, EGFR, and IFNGR2 are downregulated genes in all three retinal

diseases. The DEGs CCL14, CNTF, CX3CL1, IL10RA, IL15RA, IL1R2, IL3RA, LTB,

PDGFC, TNFRSF25 and TNFRSF4 are upregulated genes in any of two retinal diseases.

Further, when we compared the results with DEGs of cytokine-cytokine receptor interaction

pathway enriched using RNA-Seq data, we have observed more than twelve DEGs that have

similar expression viz ACVRL1, CCL19, CCL23, CXCL1, IL10RA, IL18RAP, IL1R2,

IL3RA, IL4R, OSM, TNFRSF25 and TNFRSF4 , suggesting their role in angiogenesis

invariably. Furthermore, we have observed that CXCL1 is the major hub of upregulated
DEGs based on the protein-protein interactions with the angiome-pathway network (figure
3), and thus it would be a potential target. In addition to CXCL1, based their angiogenic role
(Figure 6), the PPI network (figure 3) and the RNA-seq data results, we highlight that
VEGFC, ACVRL1, CCL23, CX3CL1 and TNFRSF25 are the notable targets.

In conclusion, the cytokine-cytokine receptor interaction pathway is highlighted as the
common contributor for the retinal angiogenesis in PDR, ROP and nAMD through
microarray metadata analysis. Further, several DEGs in the cytokine-cytokine receptor
interaction pathway are identified as potential angiogenic factors that may cause the retinal
angiogenesis. However, as only a few studies have been published related to the highlighted
VEGF-independent retinal angiogenic pathways and genes, we hope that this work will lead
future investigations into mechanisms of retinal angiogenesis, as well as identification of
potential drug target.

References

1.   Aiello, L. M. Perspectives on diabetic retinopathy. *Am. J. Ophthalmol.* (2003).
doi:10.1016/S0002-9394(03)00219-8

2.   Das, A. & McGuire, P. G. Retinal and choroidal angiogenesis: Pathophysiology and
strategies for inhibition. *Progress in Retinal and Eye Research* (2003).
doi:10.1016/j.preteyeres.2003.08.001

3.   Folkman, J., Watson, K., Ingber, D. & Hanahan, D. Induction of angiogenesis during
the transition from hyperplasia to neoplasia. *Nature* (1989). doi:10.1038/339058a0

4.   Dulak, J., Józkowicz, A. & Łoboda, A. *Angiogenesis and vascularisation: Cellular*
and molecular mechanisms in health and diseases. Angiogenesis and Vascularisation:
Cellular and Molecular Mechanisms in Health and Diseases (2013).
doi:10.1007/978-3-7091-1428-5

5.   Neve, A., Cantatore, F. P., Maruotti, N., Corrado, A. & Ribatti, D. Extracellular matrix
modulates angiogenesis in physiological and pathological conditions. *BioMed*
Research International (2014). doi:10.1155/2014/756078

6.   Andreoli, C. M. & Miller, J. W. Anti-vascular endothelial growth factor therapy for
ocular neovascular disease. *Current Opinion in Ophthalmology* (2007).

334   doi:10.1097/ICU.0b013e3282f0ca54

7.   Schlingemann, R. O. & Witmer, A. N. Treatment of retinal diseases with VEGF

antagonists. *Progress in Brain Research* (2009).

doi:10.1016/S0079-6123(09)17517-9

8.   Ferrara, N., Hillan, K. J., Gerber, H. P. & Novotny, W. Discovery and development of

bevacizumab, an anti-VEGF antibody for treating cancer. *Nature Reviews Drug*

Discovery (2004). doi:10.1038/nrd1381

9.   Hoeben, A. *et al.* Vascular endothelial growth factor and angiogenesis.

Pharmacological Reviews (2004). doi:10.1124/pr.56.4.3

10.   Usui, Y. *et al.* Angiogenesis and Eye Disease. *Annu. Rev. Vis. Sci.* (2015).

doi:10.1146/annurev-vision-082114-035439

11.   Giet, M., Henkel, C., Schuchardt, M. & Tolle, M. Anti-VEGF Drugs in Eye Diseases:

Local Therapy with Potential Systemic Effects. *Curr. Pharm. Des.* (2015).

doi:10.2174/1381612821666150225120314

12.   Mirshahi, A. *et al.* Bevacizumab-augmented retinal laser photocoagulation in

proliferative diabetic retinopathy: A randomized double-masked clinical trial. *Eur. J.*

Ophthalmol. (2008).

13.   Simó, R. & Hernández, C. Intravitreous anti-VEGF for diabetic retinopathy: Hopes

and fears for a new therapeutic strategy. *Diabetologia* (2008).

doi:10.1007/s00125-008-0989-9

14.   K., N. *et al.* Vascular endothelial growth factor-A is a survival factor for retinal

neurons and a critical neuroprotectant during the adaptive response to ischemic injury.

American Journal of Pathology (2007).

15.   Bałan, B. J. & Słotwiński, R. VEGF and tumor angiogenesis. *Central-European*

Journal of Immunology (2008).

16.   Zhang, F. *et al.* VEGF-B is dispensable for blood vessel growth but critical for their

survival, and VEGF-B targeting inhibits pathological angiogenesis. *Proc. Natl. Acad.*

Sci. U. S. A. (2009). doi:10.1073/pnas.0813061106

17.   Kim, Y. A., Wuchty, S. & Przytycka, T. M. Identifying causal genes and dysregulated

pathways in complex diseases. *PLoS Comput. Biol.* (2011).
doi:10.1371/journal.pcbi.1001095

18.   Ahn, T., Lee, E., Huh, N. & Park, T. Personalized identification of altered pathways in
cancer using accumulated normal tissue data. in *Bioinformatics* (2014).
doi:10.1093/bioinformatics/btu449

19.   Liu, K. Q., Liu, Z. P., Hao, J. K., Chen, L. & Zhao, X. M. Identifying dysregulated
pathways in cancers from pathway interaction networks. *BMC Bioinformatics* (2012).
doi:10.1186/1471-2105-13-126

20.   Barrett, T. *et al.* NCBI GEO: Archive for functional genomics data sets - Update.
Nucleic Acids Res. (2013). doi:10.1093/nar/gks1193

21.   Huang, D. W., Sherman, B. T. & Lempicki, R. A. Systematic and integrative analysis
of large gene lists using DAVID bioinformatics resources. *Nat. Protoc.* (2009).
doi:10.1038/nprot.2008.211

22.   Galavotti, S. *et al.* The autophagy-associated factors DRAM1 and p62 regulate cell
migration and invasion in glioblastoma stem cells. *Oncogene* (2013).
doi:10.1038/onc.2012.111

23.   Oba, S. *et al.* A Bayesian missing value estimation method for gene expression profile
data. *Bioinformatics* (2003). doi:10.1093/bioinformatics/btg287

24.   Bolstad, B. M., Irizarry, R. A., Åstrand, M. & Speed, T. P. A comparison of
normalization methods for high density oligonucleotide array data based on variance
and bias. *Bioinformatics* (2003). doi:10.1093/bioinformatics/19.2.185

25.   Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based approach
for interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. U. S. A.*
(2005). doi:10.1073/pnas.0506580102

26.   Chu, L. H., Rivera, C. G., Popel, A. S. & Bader, J. S. Constructing the angiome: A
global angiogenesis protein interaction network. *Physiol. Genomics* (2012).
doi:10.1152/physiolgenomics.00181.2011

27.   Tressel, S. L. *et al.* Pharmacology, biodistribution, and efficacy of GPCR-based
pepducins in disease models. *Methods in molecular biology (Clifton, N.J.)* (2011).

392   doi:10.1007/978-1-60761-919-2_19

393   28.   Pacini, L. *et al.* M2muscarinic receptors inhibit cell proliferation and migration in
394   urothelial bladder cancer cells. *Cancer Biol. Ther.* (2014).
395   doi:10.4161/15384047.2014.955740

396   29.   Campa, D. *et al.* Polymorphisms of dopamine receptor/transporter genes and risk of
397   non-small cell lung cancer. *Lung Cancer* (2007). doi:10.1016/j.lungcan.2006.11.007

398   30.   Mansouri, S. *et al.* GalR3 activation promotes adult neural stem cell survival in
399   response to a diabetic milieu. *J. Neurochem.* (2013). doi:10.1111/jnc.12396

400   31.   Romero, G., Von Zastrow, M. & Friedman, P. A. Role of PDZ Proteins in Regulating
401   Trafficking, Signaling, and Function of GPCRs. Means, Motif, and Opportunity. in
402   *Advances in Pharmacology* (2011). doi:10.1016/B978-0-12-385952-5.00003-8

403   32.   Shang, Y., Zhang, J. & Huang, E. J. HIPK2-mediated transcriptional control of
404   NMDA receptor subunit expression regulates neuronal survival and cell death. *J.*
405   *Neurosci.* (2018). doi:10.1523/JNEUROSCI.3577-17.2018

406   33.   Ferguson, H. J. M. *et al.* Glutamate dependent NMDA receptor 2D is a novel
407   angiogenic tumour endothelial marker in colorectal cancer. *Oncotarget* (2016).
408   doi:10.18632/oncotarget.7812

409   34.   Zamani, A. & Qu, Z. Serotonin activates angiogenic phosphorylation signaling in
410   human endothelial cells. *FEBS Lett.* (2012). doi:10.1016/j.febslet.2012.05.047

411   35.   Ding, M. *et al.* Enhancer RNA – P2RY2e induced by estrogen promotes malignant
412   behaviors of bladder cancer. *Int. J. Biol. Sci.* (2018). doi:10.7150/ijbs.27151

413   36.   Chu, L. H., Lee, E., Bader, J. S. & Popel, A. S. Angiogenesis interactome and time
414   course microarray data reveal the distinct activation patterns in endothelial cells. *PLoS*
415   *One* (2014). doi:10.1371/journal.pone.0110871

416   37.   Huan, J. *et al.* Insights into significant pathways and gene interaction networks
417   underlying breast cancer cell line MCF-7 treated with 17β-Estradiol (E2). *Gene*
418   (2014). doi:10.1016/j.gene.2013.08.027

419   38.   Wei, P., Tang, H. & Li, D. Insights into Pancreatic Cancer Etiology from Pathway
420   Analysis of Genome-Wide Association Study Data. *PLoS One* (2012).

doi:10.1371/journal.pone.0046887

39. Myers, J. S., von Lersner, A. K. & Sang, Q. X. A. Proteomic upregulation of fatty acid

synthase and fatty acid binding protein 5 and identification of cancer- and

race-specific pathway associations in human prostate cancer tissues. *J. Cancer* (2016).

doi:10.7150/jca.15860

40. Shi, Z. *et al.* Exploring the key genes and pathways of osteosarcoma with pulmonary

metastasis using a gene expression microarray. *Mol. Med. Rep.* (2017).

doi:10.3892/mmr.2017.7577

41. Wei, L., He, F., Zhang, W., Chen, W. & Yu, B. Bioinformatics analysis of microarray

data to reveal the pathogenesis of diffuse intrinsic pontine glioma. *Biol. Res.* (2018).

doi:10.1186/s40659-018-0175-6

42. Bromberg-White, J. L. *et al.* Identification of VEGF-independent cytokines in

proliferative diabetic retinopathy vitreous. *Investig. Ophthalmol. Vis. Sci.* (2013).

doi:10.1167/iovs.13-12518

43. Yan, X. J. *et al.* Identification of outcome-correlated cytokine clusters in chronic

lymphocytic leukemia. *Blood* (2011). doi:10.1182/blood-2011-03-342436

44. Benevides, L. *et al.* IL17 promotes mammary tumor progression by changing the

behavior of tumor cells and eliciting tumorigenic neutrophils recruitment. *Cancer Res.*

(2015). doi:10.1158/0008-5472.CAN-15-0054

45. Mar, A. C. *et al.* Interleukin-1 receptor type 2 acts with c-Fos to enhance the

expression of interleukin-6 and vascular endothelial growth factor A in colon cancer

cells and induce angiogenesis. *J. Biol. Chem.* (2015). doi:10.1074/jbc.M115.644823

46. Wang, D. & DuBois, R. N. Measurement of Eicosanoids in Cancer Tissues. *Methods*

in Enzymology (2007). doi:10.1016/S0076-6879(07)33002-4

47. Zadka, Ł. *et al.* The expression of IL10RA in colorectal cancer and its correlation with

the proliferation index and the clinical stage of the disease. *Cytokine* (2018).

doi:10.1016/j.cyto.2018.04.030

48. Li, R., Wen, R., Banzon, T., Maminishkis, A. & Miller, S. S. CNTF mediates

neurotrophic factor secretion and fluid absorption in human retinal pigment

450   epithelium. *PLoS One* (2011). doi:10.1371/journal.pone.0023148

451 49. Recchia, F. M., Xu, L., Penn, J. S., Boone, B. & Dexheimer, P. J. Identification of

452   genes and pathways involved in retinal neovascularization by microarray analysis of

453   two animal models of retinal angiogenesis. *Investig. Ophthalmol. Vis. Sci.* (2010).

454   doi:10.1167/iovs.09-4006

455 50. Matsumura, A. *et al.* HGF regulates VEGF expression via the c-Met receptor

456   downstream pathways, PI3K/Akt, MAPK and STAT3, in CT26 murine cells. *Int. J.*

457   *Oncol.* (2013). doi:10.3892/ijo.2012.1728

458 51. Ueda, M. & Saji, H. Radiolabeled probes targeting hypoxia-inducible factor-1-active

459   tumor microenvironments. *Scientific World Journal* (2014).

460   doi:10.1155/2014/165461

461 52. Xu, Z. *et al.* CCL19 suppresses angiogenesis through promoting miR-206 and

462   inhibiting Met/ERK/Elk-1/HIF-1α/VEGF-A pathway in colorectal cancer. *Cell Death*

463   *Dis.* (2018). doi:10.1038/s41419-018-1010-2

464 53. Liao, Y. Y. *et al.* CCL3 promotes angiogenesis by dysregulation of miR-374b/

465   VEGF-A axis in human osteosarcoma cells. *Oncotarget* (2016).

466   doi:10.18632/oncotarget.6708

467 54. Karar, J. & Maity, A. PI3K/AKT/mTOR Pathway in Angiogenesis. *Front. Mol.*

468   *Neurosci.* (2011). doi:10.3389/fnmol.2011.00051

469 55. Schreiber, T. H. & Podack, E. R. Immunobiology of TNFSF15 and TNFRSF25.

470   *Immunol. Res.* (2013). doi:10.1007/s12026-013-8465-0

471 56. Song, J., So, T. & Croft, M. Activation of NF-κB1 by OX40 Contributes to

472   Antigen-Driven T Cell Expansion and Survival. *J. Immunol.* (2008).

473   doi:10.4049/jimmunol.180.11.7240

474 57. Jeong, W., Bae, H., Lim, W., Bazer, F. W. & Song, G. Differential expression and

475   functional roles of chemokine (C-C motif) ligand 23 and its receptor chemokine (C-C

476   motif) receptor type 1 in the uterine endometrium during early pregnancy in pigs. *Dev.*

477   *Comp. Immunol.* (2017). doi:10.1016/j.dci.2017.07.007

478 58. Nakayama, T. *et al.* Eotaxin-3/CC Chemokine Ligand 26 Is a Functional Ligand for

CX3CR1. *J. Immunol.* (2010). doi:10.4049/jimmunol.0904126

59.   Gregg, B., Lumeng, C. N. & Bernal-Mizrachi, E. Fractalkine signaling in regulation of

insulin secretion. *Islets* (2014). doi:10.4161/isl.27861

60.   Jo, S. Y. *et al.* CT perfusion can predict overexpression of CXCL8 (Interleukin-8) in

head   and   neck   squamous   cell   carcinoma. *Am.   J.   Neuroradiol.* (2013).

doi:10.3174/ajnr.A3610

61.   Kritharis, A., Al-Samkari, H. & Kuter, D. J. Hereditary hemorrhagic telangiectasia:

Diagnosis and management from the hematologist's perspective. *Haematologica*

(2018). doi:10.3324/haematol.2018.193003

62.   Kochumon, S. *et al.* Adipose tissue expression of CCL19 chemokine is positively

associated   with   insulin   resistance. *Diabetes.   Metab.   Res.   Rev.* (2019).

doi:10.1002/dmrr.3087

63.   Wang, H., Lafdil, F., Kong, X. & Gao, B. Signal transducer and activator of

transcription 3 in liver diseases: A novel therapeutic target. *International Journal of*

Biological Sciences (2011). doi:10.7150/ijbs.7.536

64.   Mirando, A. C., Francklyn, C. S. & Lounsbury, K. M. Regulation of angiogenesis by

aminoacyl-trna synthetases. *International Journal of Molecular Sciences* (2014).

doi:10.3390/ijms151223725

65.   Cane, G. *et al.* Up-regulation of intestinal vascular endothelial growth factor by

Afa/Dr   diffusely   adhering   Escherichia   coli. *PLoS   One* (2007).

doi:10.1371/journal.pone.0001359

66.   Liu, N. *et al.* Helicobacter pylori promotes angiogenesis depending on

Wnt/beta-catenin-mediated   vascular   endothelial   growth   factor   via   the

cyclooxygenase-2   pathway   in   gastric   cancer. *BMC   Cancer* (2016).

doi:10.1186/s12885-016-2351-9

Acknowledgments: The authors are thankful to the Science and Engineering Research

Board, Govt. of India (PDF/2016/001448), for financial support. The authors thank Prof. VR.

Muthukkaruppan for the helpful discussion.

Author Contributions: Conceptualization, U.S., methodology, U.S., writing—original

draft preparation, U.S., Investigation, B.D., writing—review and editing, B.D., supervision,

B.D.

Disclosure: The authors declare no conflict of interest.

Legends:

Table 1. List of dysregulated pathways enriched in retinal angiogenic disease samples

compared to normal control samples.

Figure 1. Schema of the workflow - samples, analysis groups used, and the methods

followed

Figure 2. Dysregulated pathway interaction network for the deferentially expressed genes

(DEGs). The green node represents the dysregulated pathways, pink represents DEGs with

fold change between -3 and -2, red represents FC ≤ -3, pale blue represents FC between 2 and

3, dark blue represents FC ≥ 3. The round edged square shape of node represents pathways,

circle represents DEGs, and yellow circle represents DEGs associated with angiogenesis GO.

Figure 3. PPI network of core angiogenesis GO related differentially expressed genes and

their PPI partners with the interaction of dysregulated pathways. The green node represents

the dysregulated pathways, pink represents DEGs with fold change between -3 and -2, red

represents FC ≤ -3, pale blue represents FC between 2 and 3, dark blue represents FC ≥ 3. The

square shape of node represents pathways, circle represents DEGs, and yellow circle

represents DEGs associated with angiogenesis GO. Edges red in colour represents the

interaction between the pathway and the PPI partners of the DEGs which associated with

angiogenesis. The size of the node corresponds to the degree i.e. number of edges linked to

that node.

Figure 4. Clustering annotation with heat map of dysregulated pathways in retinal

angiogenesis related diseases

Figure 5. Heat map of DEGs in cytokine-cytokine receptor interaction pathway in retinal

angiogenic related diseases

Figure 6. Regulatory role of DEGs in cytokine-cytokine receptor interaction pathway. Dark

boxes indicate the DEGs identified in the retinal angiogenesis samples used in this study.

Table(s)

Table 1. List of dysregulated pathways enriched in retinal angiogenic disease samples

compared to normal control samples.

Dysregulated pathway	Normalized enrichment score	p value
Neuroactive ligand receptor interaction	3.101	0.000
Cytokine cytokine receptor interaction	2.553	0.000
Arachidonic acid metabolism	1.931	0.007
Aminoacyl tRNA biosynthesis	-1.572	0.042
Tight junction	-1.631	0.042
Pathogenic Escherichia coli infection	-1.646	0.032
Cysteine and methionine metabolism	-1.667	0.035
Vibrio cholerae infection	-1.753	0.018
Endocytosis	-1.768	0.014
Regulation of actin cytoskeleton	-1.872	0.012
Epithelial cell signaling in helicobacter pylori infection	-1.873	0.016
Arrhythmogenic right ventricular cardiomyopathy	-1.886	0.011
p53 signaling pathway	-1.892	0.008
Protein export	-1.909	0.004
Lysosome	-1.925	0.004
Adherens junction	-2.040	0.002
Spliceosome	-2.094	0.004
Alzheimers disease	-2.108	0.000

Parkinsons disease	-2.324	0.002
Huntingtons disease	-2.494	0.000
Oxidative phosphorylation	-2.579	0.000
Ribosome	-4.047	0.000

542 **Figures:**

544 **Figure 1.** Schema of the workflow - samples, analysis groups used, and the methods

545 followed

546
547

Figure 2. Dysregulated pathway interaction network for the deferentially expressed genes (DEGs). The green node represents the dysregulated pathways, pink represents DEGs with fold change between -3 and -2, red represents FC ≤ -3, pale blue represents FC between 2 and 3, dark blue represents FC ≥ 3. The round edged square shape of node represents pathways, circle represents DEGs, and yellow circle represents DEGs associated with angiogenesis GO.

Figure 3. PPI network of core angiogenesis GO related differentially expressed genes and their PPI partners with the interaction of dysregulated pathways. The green node represents the dysregulated pathways, pink represents DEGs with fold change between -3 and -2, red represents FC ≤ -3, pale blue represents FC between 2 and 3, dark blue represents FC ≥ 3. The square shape of node represents pathways, circle represents DEGs, and yellow circle represents DEGs associated with angiogenesis GO. Edges red in colour represents the interaction between the pathway and the PPI partners of the DEGs which associated with angiogenesis. The size of the node corresponds to the degree i.e. number of edges linked to that node.

Figure 4. Clustering annotation with heat map of dysregulated pathways in retinal
angiogenesis related diseases

Figure 5. Heat map of DEGs in cytokine-cytokine receptor interaction pathway in retinal

angiogenic related diseases

570 **Figure 6.** Regulatory role of DEGs in cytokine-cytokine receptor interaction pathway. Dark

571 boxes indicate the DEGs identified in the retinal angiogenesis samples used in this study.