The subcellular architecture of the *xyl* gene expression flow of the TOL catabolic plasmid of *Pseudomonas putida* mt-2

by

Juhyun Kim[1m], Angel Goñi-Moreno[2,3] and Víctor de Lorenzo[1]

[1]*Systems Biology Department, Centro Nacional de Biotecnología-CSIC, Campus de Cantoblanco, Madrid 28049, Spain,* [2]*School of Computing, Newcastle University, Newcastle Upon Tyne, NE4 5TG, UK,* [3]*Centro de Biotecnología y Genómica de Plantas (CBGP, UPM-INIA), Universidad Politécnica de Madrid (UPM), Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria (INIA), Campus de Montegancedo-UPM, 28223 Pozuelo de Alarcón, Madrid, Spain*

Running Title: Subcellular localization of TOL transcripts in *P. putida*

Keywords: *Pseudomonas putida, xylUW, xylX,* mRNA localization, specific DNA locus

[m]Current address: Department of Chemical Engineering, Imperial College London, London, SW7 2AZ, UK

____

* Correspondence to: Víctor de Lorenzo

     Centro Nacional de Biotecnología-CSIC

     Campus de Cantoblanco, Madrid 28049, Spain

     Tel.: 34- 91 585 45 36; Fax: 34- 91 585 45 06

     E-mail: vdlorenzo@cnb.csic.es

ABSTRACT. Despite intensive research on the biochemical and regulatory features of the archetypal catabolic TOL system borne by pWW0 of *Pseudomonas putida* mt-2, the physical arrangement and tridimensional logic of the *xyl* gene expression flow remains unknown. In this work, the spatial distribution of specific *xyl* mRNAs with respect to the host nucleoid, the TOL plasmid and the ribosomal pool has been investigated. *In situ* hybridization of target transcripts with fluorescent oligonucleotide probes revealed that *xyl* mRNAs cluster in discrete foci, adjacent but clearly separated from the TOL plasmid and the cell nucleoid. Also, they co-localize with ribosome-rich domains of the intracellular milieu. This arrangement was kept even when the *xyl* genes were artificially relocated at different chromosomal locations. The same happened when genes were expressed through a heterologous T7 polymerase-based system, which originated mRNA foci outside the DNA. In contrast, rifampicin treatment, known to ease crowding, blurred the confinement of *xyl* transcripts. This suggested that *xyl* mRNAs intrinsically run away from their initiation sites to ribosome-rich points for translation—rather than being translated coupled to transcription. Moreover, the results suggest that the distinct subcellular motion of *xyl* mRNAs results both from innate properties of the sequence at stake and the physical forces that keep the ribosomal pool away from the nucleoid in *P. putida.* This scenario is discussed on the background of current knowledge on the 3D organization of the gene expression flow in other bacteria and the environmental lifestyle of this soil microorganism.

IMPORTANCE. The transfer of information between DNA, RNA and proteins in a bacterium is often compared to the decoding of a piece of software in a computer. However, the tridimensional layout and the relational logic of the cognate biological hardware i.e. the nucleoid, the RNA polymerase and the ribosomes, are habitually taken for granted. In this work we inspected the localization and fate of the transcripts that stem from the archetypal biodegradative plasmid pWW0 of soil bacterium *Pseudomonas putida* KT2440 through the non-homogenous milieu of the bacterial cytoplasm. The results expose that—similarly to computers also—the material components that enable the expression flow are well separated physically and they decipher the sequences through a distinct tridimensional arrangement with no indication of transcription/translation coupling. We argue that the resulting subcellular architecture enters an extra regulatory layer that obeys a species-specific positional code that accompanies the environmental lifestyle of this bacterium.

INTRODUCTION

The TOL system encoded by plasmid pWW0 of *Pseudomonas putida* mt-2 is to this date the most

thoroughly studied example of biodegradative system in soil microorganisms. The primary function of this

catabolic device is enabling carrier bacteria to grow on toluene, *m*-xylene, *p*-xylene and other related

aromatics through a set of enzymes encoded by *upper* and *lower* plasmid-borne operons (Fig. 1A; 1-3).

While catabolic traits of this sort are not uncommon in many other environmental isolates, what makes

the TOL system special is the extraordinary regulatory intricacy that controls expression of the *xyl* genes

and their high-level interplay with the host's physiological regulons. The many mechanisms unveiled over

the years in this respect seem to include nearly every device known in the prokaryotic word for ruling the

gene expression flow (4-6). This state of affairs has made the TOL plasmid and its bearer (*P. putida*

KT2440) a beneficiary of the suite of conceptual and material tools of contemporary Systems and

Synthetic Biology (7, 8). In particular, the wealth of experimental data on expression of the *xyl* genes has

enabled the understanding of the cognate regulatory network as a complex device that processes inputs

into outputs following a layer of logic gates implemented with promoters, transcriptional factors and

sRNAs (9, 10). It could thus be argued that we know at this point much about the genetically-encoded

software of the system and the relational logic that rules its performance. In contrast, we know virtually

nothing of the physical arrangement of the *hardware* that sustains the same process. In particular, the

gross spatial disposition of the molecular actors that execute the transfer of information from the *xyl* genes

to production of catabolic enzymes is unknown.

The same sort of questions has been tackled in other prokaryotic systems. There seem to be at least two

different patterns of transcript localization. In one scenario, the mRNA remains close to the site of

transcription. This is the case of the mRNAs of *groESL* and *creS* of *Caulobacter crescentus* and *lacZ* of

E. coli, which appear to remain in the vicinity of the corresponding genomic DNA loci where they are

transcribed (11). This suggests that for mRNAs to be translated they need to be associated to the

ribosomes while being produced (as the prevailing view of generalized transcription-translation coupling

would imply) or short after their creation by RNAP (11, 12). An alternative scenario involves migration of

mRNAs to specific spots of the cytoplasm for translation close to the site(s) where the gene products are

needed. While this setting argues against generalized transcription/coupling translation, there is solid

evidence of its occurrence e.g. the *bglGFB* operon, whose products were located in sub-cellular

localizations where cognate proteins were expected to work (13). This example may not be anecdotal as

transcriptome-wide scale studies of mRNA localization in *E. coli* revealed that a large share of transcripts

were found in cell domains (e.g. membranes, cytoplasm, poles) that coincided with the function of the

cognate proteome (14, 15). Furthermore, the mRNA of archetypal membrane proteins LacY and TetA

are distinctively close to the cell envelope (16, 17). In other instances, a signal-recognition particle (SRP)

is involved in the movement of mRNAs to the membrane, as translation of some mRNAs encoding inner-

membrane proteins produces a signal peptide that recruits such SRP and the complex leads the transcript

to its intracellular address (15, 16, 18). Finally, Rho-dependent transcription termination in *Bacillus subtilis*

is somewhat weak, and *runaway*, untranslated mRNAs are abundant (19). In sum, the fate of each

transcript seems to be both gene (i.e. sequence)-dependent and species-dependent.

In this work we have inspected the localization of mRNAs initiated in the catabolic promoters of the TOL

plasmid pWW0 of *P. putida* mt-2. The starting point for tackling the issue is the earlier observation that

RNA polymerase (RNAP) of *P. putida* fully co-localizes with the chromosomal DNA of the nucleoid while

being entirely apart from the bulk of the ribosomal pool (20). This observation suggested that rather than

being coupled to translation *ab initio,* many (if not most) of the mRNAs initiated at chromosomal promoters

need to move to ribosome-rich, nucleoid-free domains of the intracellular space for translation. Note that

in the case of the *xyl* promoters, transcription initiates in an extrachromosomal element, and therefore

the 3D itinerary of the corresponding mRNAs could be different. As shown below, by merging genetic

analyses with *in situ* RNA-FISH and DNA-FISH technology we could faithfully locate the relative positions

of the nucleoid DNA, the pWW0 plasmid, the *xyl* transcripts and the ribosomes and predict the motion of

cognate mRNAs through the cell interior. The results exposed an unexpected degree of physical partition

among the material actuators of the gene expression flow that may help *P. putida* to deal with its typical

environmental settings.

RESULTS AND DISCUSSION

mRNAs of catabolic genes expressed from the TOL plasmid are spatially organized. To visualize

specific mRNAs of *P. putida* mt-2 stemming from the TOL operons of plasmid pWW0 we adopted an

RNA FISH approach (Fig. 1B). To this end, the strain was cultured in M9 minimal medium with succinate

as the sole carbon source until the cells reached exponential phase. At this point the cultures were

exposed (or not) to saturating vapors of *m*-xylene to induce transcription of the catabolic *xyl* genes. After 2 h, samples were collected and fixed with formaldehyde for hybridization with specific probes as described in the Materials and Methods section. For this, two sets of 48 CAL Fluor Red 610-tagged fluorescent oligonucleotides (20 nt long; Supplementary Table S2) were synthesized that covered, respectively, the leading 1418 bps of the *upper* TOL operon transcript spanning the whole of *xylU* and part of *xylW* (encoding benzyl alcohol dehydrogenase) and the front segment of the TOL *lower* operon, encompassing 1203 pb of the *xylX* gene (alpha subunit of toluate 1,2-dioxygenase). After hybridization with these oligo sets, the samples were washed and red signals inspected with fluorescence microscopy.

As shown in Fig. 1C, distinct, discrete fluorescent foci were clearly noticed under the microscope (1-2 per cell) from the cultures subject to *m*-xylene exposure upon in situ hybridization with *upper*-pathway or *lower*-pathway specific oligonucleotides. In contrast, no signals were detected in bacteria grown in M9-succinate without aromatic effector (Supplementary Fig. S1A). These observations accredited that the designed probe sets and the methodology proper were working as expected, as there were no signals that could be attributed to hybridization with DNA. Therefore, the foci appeared to represent *upper* or *lower xyl* transcripts. To further benchmark the experimental approach, cells were treated with either toluene (an alternative TOL substrate) or *o*-xylene (a gratuitous inducer of the *upper* pathway and downstream activation of the *lower* pathway (21, 22; Fig. 1A). Under these conditions, fluorescent foci for both the *xylUW* and the *xylX* transcripts were expectedly observed. In contrast, cells treated with benzoate or 3-methyl benzoate (substrates of the *lower* pathway) produced signals for *xylX* mRNA only (Supplementary Fig. S1B). Appearance of the fluorescent signals matched the known regulatory network that rules the interplay between the substrates, regulators and catabolic operons of the TOL system (Fig. 1A). We could thus safely consider that the foci inside cells shown in Fig. 1C were the result of the *bona fide* hybridization of the fluorescent probes to specific mRNA of *xylUW* or *xylX*. Closer inspection of the images revealed that no red output ever appeared dispersed throughout the cell, but always as discrete foci (Fig. 1B). Yet, their position in respect to the cell shape varied among cells and signals were located near the center, the poles, the contour or the septum of the cells (Fig. 1C). We thus set out to characterize this asymmetrical localization of the *xyl* transcripts in respect to the nucleoid occupation and to the TOL plasmid, as explained below.

***xyl* mRNAs occupy subcellular nucleoid-free regions.** In order to identify the relative localization of

the *xyl* transcripts in respect to the nucleoid, following hybridization with the fluorescent probes described

above the bacterial genome was stained with 4′, 6-diamidino-2-phenylindole (DAPI) in cells exposed to

m-xylene. The results of this procedure with the *upper* and *lower* pathway TOL probes (*xylUW* and *xylX*)

respectively are shown in Fig. 2. For interpreting these images, note that [i] previous work showed that

the DNA of the nucleoid overlaps spatially with *P. putida*'s RNAP (20) and [ii] DAPI binds both

chromosomal and plasmid DNA. Inspection of the cells under the microscope (Fig. 2A) revealed that

signals stemming from *xylUW* or *xylX* transcripts were located in the nucleoid-free region of the bacterial

inside. In order to quantity the phenomenon, >100 pictures of individual DAPI-stained and fluorescent

oligo-hybridized cells were separately recorded with the blue and red channels of the fluorescence

microscope and they were automatically inspected with the CellShape image analysis tool (23; Fig. 2B).

The outcome of this analysis is shown in Fig. 2C. Very few red foci (< 1%) overlapped with the densest

DAPI signals and as little as 15% of fluorescent spots from either *xylUW* or *xylX* RNA were detected at

the peripheral regions of genomic DNA. Instead, the vast majority of the remaining foci were placed away

from the blue signal (Fig. 2C). It thus looked like the bulk of TOL transcripts were located within subcellular

regions with no or little overlap with the DNA signal and therefore virtually devoid of RNAP (20). This

suggests that once formed, *xyl* mRNAs could migrate for translation to a site different from the place

where transcription is initiated. Note that the pathway substrate (*m*-xylene) dissolves in the cell membrane

(24). Moreover, the *xylM* product (hydroxylase component of the leading pathway enzyme xylene

monooxygenase) is located in the membrane (25). It is not uncommon that genomic sites encoding

envelope-associated proteins are transcribed coupled and co-translationally inserted to the membrane

through a so-called *transertion* mechanism (16). In the case of the TOL transcripts it looked instead that

movement away from the DNA begins after transcription has terminated and the mRNA has disengaged

from the nucleoid. To clarify this, we examined the relative position of the *xyl* mRNA in respect to their

actual origin inside the cells (i.e. the TOL plasmid) as explained below.

***xyl* transcripts map near but do not overlain the TOL plasmid.** Subcellular localization of the TOL

plasmid within the whole of the DAPI-stained DNA of the *P. putida*'s nucleoid required first tagging pWW0

with a different fluorescent label. A 9679 pb DNA sequence including an array of tandemly repeated *tetO*

operators for the TetR repressor (26) was inserted in a site of the plasmid between and close to the *upper*

and *lower* operons i.e. by ORF105 of the current annotation (27); Supplementary Fig. S2A). The

corresponding sequences could then be exposed through FISH with a special type of 6-

Carboxyfluorescein (6-FAM)-labeled 18 nt oligonucleotides. These encoded the *tetO* operator and bore

a distinctive chemical configuration (so called locked nucleic acid LNA structure) that increases specificity

for target DNA (11; Supplementary Fig. 2A). Samples were thus first hybridized with *xyl*-specific (red

fluorescence) and pWW0-specific (green fluorescence) probes and then stained with DAPI. As a result

of this procedure, both red, green and blue fluorescent signals could be mapped in cells exposed to *m*-

xylene, corresponding to *xylUW* or *xylX* mRNA, plasmid, and nucleoid, respectively (Fig. 3B-D and Fig.

4B-D). In contrast, cells grown without the aromatic compound lacked the red signal altogether

(Supplementary Fig. S3A).

The control images (panels 1 and 2 of Figs. 3B-D and 4B-D; Supplementary Fig. S3) show individual

bacteria hosting discrete green spots in cells bearing the TOL plasmid variant tagged with *tetO* array. In

contrast, no defined green foci were observed in cells with the intact TOL plasmid devoid of the same

array (Supplementary Fig. S2B). This verified that the *tetO* probe hybridized specifically to the tagged

plasmid without significantly interact with any other DNA of the cells. Further inspection of the pictures

indicated that the green signals appeared only once or twice in every cell (in the last case in a symmetrical

fashion; Supplementary Fig. S3). Such a subcellular distribution of the plasmid has been observed before

(28) and plausibly reflects the plasmid partition system (29-31). Once the position of the TOL plasmid

was determined, by filtering the images with the adequate color channels we could pinpoint the relative

localization of pWW0, the nucleoid and the TOL transcripts.

Fig. 3B-D and Fig. 4B-D show a few examples of thereby processed cells to locate the *xylUW* (*upper*

pathway) and the *xylX* mRNAs (*lower* pathway), respectively. Each of these images was not only visually

inspected but also the signals quantified with the CellShape software (23; Supplementary Fig. S4). In

general, the transcripts from either the *upper* or the *lower* TOL pathway behaved similarly. The first piece

of information resulting from a detailed image analysis (panels 2/6 of Fig. 3B-D and Fig. 4B-D) is that

TOL plasmid could be found both in DAPI-intense regions and in the peripheral space of the cytoplasm

with little or insignificant DAPI signal. This agrees with the pWW0 predicted segregation system: the

presence of plasmid-encoded *parA* and *parB* indicates pWW0 to bear a Type I partition mechanism (27)

though which the plasmid can either detach or remain connected to the chromosome through a ParA-

ATP-nucleoid complex (30).

Inspection of colored spots merged within the contour of single cells exposed two predominant spatial

arrangements of the different signals. In one case, red and green spots virtually co-localized (e.g. Fig.

3B), while in most others, the two signals were close to each other but clearly separated (Fig. 3C-D and

Fig. 4B-D). It is likely that these two scenarios reflect different stages of transcription of *xyl* genes. In an

early stage, mRNA is necessarily tethered to its DNA template in the plasmid and therefore the gene and

its transcript occupy the same spot. But at a later stage the *xyl* mRNA could move away from the plasmid

towards nucleoid-free ribosome-rich domains of the cell inside (20). Note that the sizes of the *upper-* and

lower TOL operons are ~ 8 Kb and 11 Kb respectively and, if linearly stretched, the 5' ends of their full-

length mRNAs could be found well away from the plasmid while still tethered to the template. However,

it is known that the mRNA from the *meta* pathway is quite short lived and starts being degraded before it

is fully synthesized (32). Note also that mRNA tends to form secondary structures that shorten the

distances between the 5' and 3' ends (33). On this basis, we argue that separation of red and green spots

in the images shown in Fig. 3, Fig. 4 and Supplementary Fig. S4 reflect an authentic migration of the TOL

transcripts away from their DNA template. The next obvious question was whether this was result of TOL

genes being encoded in a type of plasmid that largely stays in the periphery of the nucleoid (see above).

Alternatively, the unexpected localization of *xyl* mRNAs could stem from intrinsic properties of the

transcripts themselves. In order to sample these possibilities we entered various types of perturbations

in the system as described below.

xyl transcripts move away from the nucleoid regardless of their replicon. Although the native

physical location of the TOL genes is in plasmid pWW0, the same growth phenotype on aromatic

compounds can be brought about when the *upper* and the *lower* pathways are placed in the chromosome.

This may happen either naturally (34) or by engineering the corresponding DNA segments in the genome

of a heterologous host (35). We took advantage of this for exploring whether the position of the *xyl* DNA

template affects the spatial distribution of the thereby transcribed RNAs. To this end, we used *P. putida*

PaW140 (36), which carries both the *upper* and *lower* operons and their cognate regulators in its

chromosome (Fig. 5A). The strain was grown and induced with *m*-xylene under identical conditions as

before and cells fixed and hybridized with RNA probes to expose *xylUW* and *xylX* transcripts in respect

to the bacterial nucleoid. The results of Fig. 5B showed the number of foci per cell to increase by ~ 30%

in *P. putida* Paw140 as compared to the lower numbers of the pWW0-bearing *P. putida* mt-2 strain (Fig.

2C), surely due to a higher transcriptional activity. This is not surprising as the pedigree of strain *P. putida*

Paw140 involved random chromosomal insertion of TOL genes and selection for best growers on *m*-

xylene, what plausibly favored implantation in regions of high transcriptional activity (37-39). Yet, when

RNA-red signals were compared to those of the DAPI-stained nucleoid, most of them were observed

away of the denser DNA regions. Specifically, < 3% of the red spots of either *xyl* mRNAs overlapped the

more compacted chromosome, while 74% of *xylUW* and 87% of *xylX* signals were enriched in the

peripheral space of the cell (Fig. 5B and 5C). Preferential localization of *xyl* mRNAs was thus maintained

regardless of the replicon (plasmid of chromosome) that bears the cognate genes. Moreover, the higher

incidence of TOL mRNAs in the nucleoid-free subcellular regions corroborates that *xyl* transcripts move

away from their transcription site.

The drive of xyl mRNAs to escape the nucleoid is detached of the transcriptional machinery. A

second type of perturbation entered in the architecture of the *xyl* gene expression flow involved complete

replacement of the native σ^{54}-dependent *Pu* promoter of the *upper* operon of the pWW0 plasmid (40, 41;

Fig. 1A) by a heterologous expression T7 system. The replacement was entered in the pWW0 plasmid

as explained in the Material and Methods section and the resulting construct (pTOL-PuxT7) then passed

to strain *P. putida* KT2440•T7 (42), which expresses the T7 RNA polymerase through a genomic insertion

of a *lacI^q/P$_{lac}$-T7pol* cassette (Fig. 6A). As shown in Supplementary Fig. S5A, the modified plasmid did

not support *m*-xylene metabolism in cells without T7 RNAP because it did not transcribe the *upper*

pathway. Due to the leakiness of the *P$_{lac}$-T7pol* born by in the *P. putida* chromosome, some red signals

indicative of *xylUW* expression were detected as well in samples without IPTG (Supplemental Fig. S5C).

Note also that there were RNA dots stemming from the *lower* pathway in the wild-type *P. putida* mt-2 host

bearing the modified plasmid pTOL-PuxT7 (Supplementary Fig. S5B). This is because *m*-xylene-

activated XylR caused overexpression of XylS, which in turn suffices to activate the *Pm* promoter even

in the absence of any effector (5, 43, 44). As a consequence, when pTOL-PuxT7 was placed in *P. putida*

KT2440•T7, [i] *P$_{T7}$* activation elicited formation of both *xylUW* transcripts (Fig. 6, panels 1-4) and *xylX*

transcripts (Fig. 6, panels 5-8) in the complete absence of the aromatic inducers used before and [ii] cells

could grow on minimal medium with *m*-xylene as sole carbon source (Supplemental Fig. S5A) due to the

performance of both the *upper* and the lower TOL pathways. This made sense within the current model

of regulation of the *xyl* operons discussed above (Fig. 1A). But in contrast to the effect of the native

effectors in the naturally-occurring system, the surrogate control of the *upper* route by T7 polymerase

originated a higher number of distinct foci per cell. While this reflected the strength of the P_{T7} promoter

as compared to *Pu* (which propagated into a more potent activity of the *Pm* promoter as well), it is

noteworthy that the red signals always appeared focused (i.e. constrained within a subcellular domain)

rather that diffused through the cytoplasm. Furthermore, they again materialized mostly in the nucleoid

free regions. Automated quantification of signals detected with the red and blue channels in individual

cells with pixel precision revealed that 91% of *xylUW* and 98% of the *xylX* were excluded from the DAPI-

stained field (Fig. 6C). Given that such transcripts originate in single promoters per bacterium and that

nucleoid-free domains of *P. putida* cells are filled with ribosomes (20) it is possible that the images of Fig.

6 reflect the detachment of the transcripts from their promoters (whether P_{T7} for *xylUW* and *Pm* for *xylX*)

and their resetting elsewhere in the cytoplasm for translation.

In addition, one important evidence embodied in this set of experiments is that formation of mRNA foci

and apparent motion towards ribosome-rich subcellular sectors occurs regardless of whether the cognate

RNAP can (host polymerase) or cannot (T7pol) couple transcription to translation. This rules out that the

distinct foci of *xylUW* and *xylX* mRNAs shown throughout this work reflect the action of

transcription/translation complexes (the so-called expressome; 45, 46) on the nucleoid surface. Instead,

it seems that the untranslated TOL transcripts move away from their promoters, plausibly towards other

subcellular locations for translation. But, if that were the case, what physical forces prevent their diffusion

through the cytoplasm?

Disruption of intracellular crowding enables xyl transcripts to diffuse throughout the cytoplasm

The data above accredit that *xyl* mRNAs leave the proximity of the nucleoid towards the ribosome-

enriched subcellular domains. Although a role for specific RNA binding proteins cannot be ruled out as

drivers of the process, a simpler explanation is that the corresponding sequences endow the *xyl*

transcripts with physical properties that make them to be quickly discharged from the vicinity of the

nucleoid, especially if the transcribed sequences are large. In fact, it seems that smaller RNAs use to

appear uniformly distributed throughout the bacterial cell, while longer molecules typically display more

limited dispersion (47). Large mRNAs can hardly co-localize with densely packed DNA regions due to

straight physical forces e.g. excluded volume effects (48, 49). As a consequence, the more crowded the

bacterial cytoplasm is the less diffusible cellular components and molecules are, logically, in a size-

dependent fashion (50, 51).

In order to test whether such a mutual exclusion between different intracellular domains accounted for

the unusual behavior of the *xyl* mRNAs we treated *P. putida* KT2440•T7 (pTOL-PuxT7) cells with

rifampicin. In one hand, since T7 RNAP is not sensitive to this antibiotic, the *upper* TOL transcript (*xylUW*)

can still be produced (the *xylX* gene of the *lower* pathway predictably cannot, though). On the other hand,

it is known that diffusion rate of ribosomal proteins is faster (11, 12) and the cDNA of the nucleoid expands

after treatment of the drug (52, 53). Finally, the dearth of ribosomes available for interacting with RNAs

due to inhibition of 16s rRNA synthesis with (54) may also ease diffusion of otherwise tethered transcripts.

As a consequence, rifampicin treatment elicits a major change in the partition of the different components

of the gene expression flow and causes a less compact intracellular milieu.

After treatment of growing *P. putida* KT2440•T7 (pTOL-PuxT7) for 2 h with the drug and *m*-xylene vapors,

the mRNAs were visualized with FISH as before. Expectedly, we could hardly detect any *xylX* mRNA

signals (Supplementary Fig. S6). In contrast, when cells were hybridized with the *xylUW* probe under the

same conditions, red signals did appear (Fig. 7 and Supplementary Fig. S5). Yet, inspection of individual

bacteria revealed a considerable variability in the intensity of the signals and their intracellular distribution.

Some cells had their whole contour non-uniformly filled with red color, while other showed a number of

discrete red signals with lower intensity (Fig. 7 and Supplementary Fig. S7). This outcome is not

altogether unexpected, as the strong T7 promoter in the plasmid (55) drains cellular resources (56, 57)

for the sake of transcription of the *upper* pathway, thereby originating noise. This effect can be

exacerbated in a plasmid, as the loss of gyrase production with the drug and the ensuing accumulation

of local supercoiling can lead to stochastic transcriptional bursts (58-60). In any case, we argue that the

lack of distinct localization patterns in rifampicin-treated cells reflects the diffusion of the TOL transcripts

under the conditions. Moreover, some images showed enrichment of *xylUW* mRNAs in the cells' internal

periphery in bacteria where the chromosome was otherwise highly compacted (Supplementary Fig. S7,

marked). In these cases, it looked like mRNA was unable penetrate the denser DNA regions but could

freely diffuse in the rest of the cytoplasmic space. Taken together, analysis of images shown in Fig. 7

and Supplementary Fig. S7 indicated that the *xylUW* transcript of rifampicin-treated cells lost its restraint

in discrete foci and could then freely circulate as the cellular crowding decreases upon antibiotic addition.

In sum, the data suggests that under native conditions *xyl* mRNAs become localized away from the

nucleoid because of their entrapment with the translational machinery and the physical forces that

determine phase separation between the different components of the gene expression flow. As shown

above, if such a separation is perturbed upon rifampicin addition, *xyl* transcripts can then circulate through

the whole cytoplasmic interior.

Conclusion. This study shows that *xyl* mRNAs are constrained within subcellular regions with limited

mobility rather than being freely diffusible inside *P. putida* mt-2 cells. Most *xyl* transcripts were detected

in the space peripheral to the nucleoid, i.e. in a subcellular domain with virtually no DNA but enriched in

ribosomes (20). That TOL mRNAs were detected away from the genetic loci where they originated

suggests that they can migrate to translational sites from the 3D spot of the cell where they were

synthesized. Our data show also that physical forces and translation is crucial for bringing about this

scenario, which disappeared when in-house transcription was halted to reduce cellular crowding and

ribosome engagement. This picture departs from the standard view of transcription-translation coupling,

where RNAP and ribosomes get together in the so-called expressome complex (45, 46) which, obviously,

must operate in close proximity to the nucleoid's DNA encoding the gene(s) at stake.

While transcript entrapment is a known mechanism of RNA control in eukaryotes (61-63) the

phenomenon as such has not been documented in prokaryotes. However, should transcript diffusion

away from the promoter and capture by ribosomes in a different cell location occur, confinement in given

spots for translation could become apparent. Some mRNAs can indeed travel throughout the *E. coli*'s

interior (64) and translation inhibition makes RNA to remain closer to the nucleoid (47). This suggests

that at least in some cases ribosomes in fact pull mRNA from the nucleoid towards the rest of the

cytoplasm. Reality is that the interplay between the various components of the gene expression flow vary

dramatically among bacterial types. In particular, the nucleocytoplasmic (NC) ratio changes from one

greatly from one species to the other, what originates cytoplasms with different biophysical properties

(65) that affect ribosome mobility and localization. For instance the high NC ratio of *Caulobacter*

crescentus forces ribosomes to mRNAs and their translating ribosomes to remain close to their gene loci

in the chromosomal DNA (11, 65, 66). In contrast the low NC ratio of *E. coli* allows segregation of

ribosomes and mRNA away from their transcriptional sites. *Pseudomonas sp.* has also a low NC ratio

(65) and the ribosomes are spatially segregated from the nucleoid (20). It is thus conceivable that TOL

transcripts are not coupled to translation as they are produced but they instead move after complete

transcription towards ribosome-rich cytoplasmic spots.

What could be the advantage of this setting of the *xyl* gene expression flow in *P. putida* as compared to

an alternative subcellular architecture? This species has bona fide *nusA* and *nusG* homologues encoded

in its genome. This indicates that transcription/translation coupling can indeed occur (45, 46), although

its global incidence as compared to *E. coli* or *Bacillus* (19) is unknown. It may well happen that having

movable *xyl* mRNAs eases assembly of the biodegradative complex encoded by the TOL operons for

degrading aromatic compounds. This is because the cognate catabolic enzymes, in particular those

which are membrane-bound (see above) could be synthesized at an optimal location site close to the site

of action. Moreover, thereby synthesized enzymes could be produced close to each other and enable

metabolic channeling (67). Finally, while Rho and transcription/translation coupling helps avoiding RNAP

collisions and R-loops—with the downside of polarity—some bacterial types could have solved the traffic

jam problem by evolving a different type of termination (19). In fact, interruption of non-translated

transcripts may be detrimental for emergence of new catabolic operons: nosense mutations in lead genes

could prevent expression of the whole cluster and curb evolution of downstream ORFs. These are all of

course hypothetical scenarios that deserve further studies.

MATERIALS AND METHODS

Culture conditions Unless otherwise indicated, *E. coli* and *P. putida* were routinely grown at 37°C and

30°C, respectively, in Luria–Bertani (LB) or M9 minimal medium (6 g l^{-1} Na_2HPO_4, 3 g l^{-1} KH_2PO_4, 1.4 g

l^{-1} $(NH_4)_2SO_4$, 0.5 g l^{-1} NaCl, 0.2 g l^{-1} $MgSO_4·7H_2O$) with 10 mM succinate. Bacteria were cultured in 100-

ml Erlenmeyer flasks with shaking at 170 rpm with 20 ml of the medium specified for the corresponding

experiment. Whenever necessary kanamycin (Km, 50 μg ml^{-1}), ampicillin (Ap, 150 μg ml^{-1}), gentamycin

(Gm, 10 μg ml^{-1}) or chloramphenicol (Cm, 30 μg ml^{-1}) was added to cultures of bacterial cells for ensuring

plasmid retention and maintenance of manipulated genotype. For the induction of TOL catabolic genes

during RNA-FISH experiment, *P. putida* strains, carrying the *xyl* genes either in the pWW0 plasmid or in

the chromosome, were overnight cultured in succinate amended M9 medium and then the bacterial

culture were 100-fold diluted in the same medium and grown until exponential phase (OD_{600} = 0.3- 0.5).

Samples were then either cultivated further without additional substrate or exposed to vaporous *m*-xylene

(1/2 dilution in dibutylphthalate, which is a non-effector for TOL genes) in a flask for 2h. To validate

whether the output of the FISH approach matches the known TOL catabolic system, we cultured the mt-

strain with other vaporous effectors such as toluene, *o*-xylene or soluble substrates such as benzoate

(5 mM) and 3MBz (5 mM). The transcription of the *xyl* genes by bacterial RNA polymerase was halted

by adding 200 µg ml^{-1} rifampicin (Rif) to the grown medium of the cognate sensitive strains.

Fluorescent *in situ* hybridization To visualize *xyl* mRNAs, RNA-FISH was performed as described

previously (11) with some modifications. To this end, cells growing in culture medium with or without

effector for *xyl* genes were fixed in formaldehyde solution (4% formaldehyde and 30 mM NaHPO$_3$ pH

7.5) by incubating for 15 min at room temperature, followed by 30 min further on ice. Samples were

collected by using centrifuge for 3 min at 4,500 g and removed supernatant. The cell pellets were washed

with diethyl pyrocarbonate (DEPC)-treated phosphate-buffered saline (PBS) and resuspended the cells

in 100 ul of GTE buffer (50 mM glucose, 20 mM Tris-HCl pH 7.5, 10 mM EDTA pH 8). Then, 12 µl of the

mixture was transferred to 4 µl of the lysozyme solution containing lysozyme and vanadyl ribonucleoside

complex (VRC), resulting in final concentration as 2.5 µg ml^{-1} of lysozyme and 2 mM VRC respectively.

Next, 3 µl of the mixed solution was applied onto poly-L-lysine coated cover slip, and stored at room

temperature to be dried on the cover slip. After putting the cover slip into a rack, this assembly was

immersed sequentially in methanol for 10 min and acetone for 30s at -20 °C. Once the coverslip was dry,

it was kept at 37 °C in 10% formamide solution (10% formamide, 2X saline-sodium citrate buffer (SSC)

in DEPC treated water, 2 mM VRC). After 60 min later, the solution was removed and 50 µl of the

hybridization solution (10 % formamide, DEPC treated 2X SSC, 10 % dextransulphate, 2 mM VRC, 40 U

RNase inhibitor, and 250 nM CAL Fluor Red 610 labeled FISH probes; Stellaris™, Biosearch

Technologies, Supplementary Tables S3) was spotted onto the cover slip and the sample was incubated

in a dark and humid chamber at 42 °C for the hybridization process. After overnight, the sample was then

washed twice with the solution (10% formamide and DEPC treated 2X SSC) for 15 min at 37 °C and

DAPI (2.5 ug ml^{-1}) staining was performed in second washing step. After brief rinse with PBS, the

coverslip was assembled with slide glass including antifade reagent Prolong (Invitrogen) and sealed by

clear nail polish. The specimen was visualized by fluorescent microscope. To identify cellular position of

the TOL plasmid carrying the tandem copies of the *tet* operators, DNA-FISH was exploited with the 6-

carboxyfluorescein (FAM) labeled locked nucleic acid (LNA) *tetO* probe (5' 6-FAM-

CTCTATCACTGATAGGGA; Bionova). It is the same strategy as RNA-FISH, but the hybridization was

carried out at two different temperatures: at 95 °C to denature DNA for 2 min and followed by at 42 °C

overnight in a dark and humid chamber. The fluorescent signals representing *xyl* mRNAs and the *tetO*-

tagged DNA were achieved by employing sequentially combined RNA/ DNA-FISH. We followed the same

procedure the RNA-FISH with *xyl* probe sets prior to performing the DNA-FISH with the *tetO* probe.

Construction of bacterial strains. The bacterial strains, plasmids, and primers used in this study are

described in Supplementary Tables S1 and S2. General methods for DNA manipulation were followed

standard protocols (68). Plasmid DNA was isolated from bacterial cells using commercial Wizard *Plus*

SV Minipreps DNA Purification kit (Promega) or the QIAprep Spin Miniprep Kit (Qiagen). To replace the

Pu promoter of the *upper* operon of the pWW0 plasmid by the orthogonal transcription T7 system, we

used the seamless allelic replacement method of (69). The delivery plasmid for executing the promoter

replacement pTOL-*PuxT7* was built as follows. First, the upstream (TS1^{Pu}, ~0.5 kb) and downstream

(TS2 ^{Pu}, ~0.5 kb) regions of the pWW0 plasmid around the *Pu* promoter were amplified with primer pairs

PuxT7-TS1F/R and PuXT7-TS2F/R respectively (Supplementary Table S2). SOEing PCR then joined

the amplicon TS1^{Pu} and TS2^{Pu}, including 3'- and 5' theT7 promoter sequence overhang complementary,

by using primer pairs PuXT7-TS1F and PuXT7-TS2R (Supplementary Table S2). The resulting PCR

product was cloned into the pEMG by using restriction enzymes such as *Eco*RI and *Bam*HI and ligation

within the same site of the vector yielded the plasmid pEMG-PuxT7, which was kept into the *E. coli* DH5α

λ*pir* strain. This plasmid was transferred to *P. putida* mt-2 by triparental mating using the *E. coli* HB101

(pRK600) as helper strain (70, 71). Next, the pSW plasmid that expresses I-SceI endonuclease under

the *Pm* promoter (72) was introduced by electroporation into the pEMG-PuxT7 cointegrated strain and

thus had resistance for both Km and Ap. The clones were grown in LB medium with Ap (500 µg ml^{-1}) and

3MBz (15 mM) to activate the *Pm* promoter, allowing I-SceI expression. The cells were plated on LB agar

and we confirmed the promoter replacement in the TOL plasmid, by testing the loss of the pEMG-PuxT7,

encoded Km resistance gene. Km-sensitive clones were selected and PCR further confirmed the

emergent colonies with T7F/ PuXT7-TS2R primer pairs (Supplementary Table S2).Then, the manipulated

plasmid pTOL-*PuxT7* was either maintained in the mt-2 strain or transferred into the KT2440•T7 strain

by conjugation.

To label the pWW0 plasmid with the tandemly repeated *tet* operators, we inserted the arrays into the

orf105 locus of the plasmid. To this end, the 570-bp of PCR product was obtained using primer pairs

105F/ 105R (Supplementary Table S2) corresponding to the 5' portion of the *orf105* gene of the pWW0

plasmid. Then, the amplified fragment was cloned into the *Hind*III and *Not*I cloning site of the pP30D-

FRT-tetO vector (26), generating pP30D-FRT-*tetO*-orf105. The construct was either kept into the CC118

strain and subsequently transferred into *P. putida* mt-2 strain, in turn the backbone of the vector, including

the *tetO* arrays, was integrated into the pWW0 plasmid, generating the strain mt-2 (pTOL-*tetO*).

Microscopy and image analysis. Microscopy was performed using an Olympus BX61 apparatus

equipped with ×100 phase contrast objective and a DP70 camera of the same brand. Signals for red-

RNA, DAPI, and green-DNA were obtained using wide field excitation with following filters; MWIY2, U-

MNU2, and U-MNIBA2. Process of multi-channel images obtained from the microscopy and

measurement of relative fluorescent intensities were carried out by using the software Fiji. To analyze

fluorescent foci representing mRNA molecules in terms of their relative cellular positions in respect of the

DAPI stained-nucleoid, we used the Cellshape (23) . To analyze overlap between the red and the blue

signals, the contour lines of both channels were calculated over the RGB TIF formatted images. Firstly,

pixel values (red/blue channels) were extracted. Secondly, the resulting values were normalised within

the interval [0,1] in order to allow for direct channel comparison regardless any potential technical

inconsistencies in measuring any one particular signal. Finally, contour lines were calculated with sub-

pixel accuracy. The overlap value refers to the overlap between the contours for the red signal and the

contours for the blue signal

SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found on line at XXX

Supplementary Tables S1, S3 and S3

Supplementary Figures S1, S2, S3, S4, S5, S6 and S7

ACKNOWLEDGEMENTS

V.D.L. and J.K. designed the project and the experimental layout. J.K. did all the experiments with the

assistance of A.G-M, who developed and implemented the software for image analysis used throughout

this work. J.K. and V.D.L. wrote the manuscript with further contributions from A.G-M. to data analysis

and interpretation of the results. We thank Christine Jacobs-Wagner for sharing the protocol for FISH

approach. Isabelle Vallet-Gely and Karl-Erich Jaeger are gratefully acknowledged also for generous

sharing of valuable materials, as is Belén Calles for insightful discussions. This work was funded by the HELIOS Project of the Spanish Ministry of Science BIO 2015-66960-C3-2-R (MINECO/FEDER), the ARISYS (ERC-2012-ADG-322797), EmPowerPutida (EU-H2020-BIOTEC-2014-2015-6335536) and MADONNA (H2020-FET-OPEN-RIA-2017-1-766975) Contracts of the European Union, and InGEMICS-CM (B2017/BMD-3691) contract of the Comunidad de Madrid (FSE, FECER). Authors declare no conflict of interest.

REFERENCES

1. Assinder SJ, Williams PA. 1990. The TOL plasmids: determinants of the catabolism of toluene and the xylenes. Adv Microb Physiol 31:1-69.

2. Marques S, Manzanera M, Gonzalez-Perez MM, Gallegos MT, Ramos JL. 1999. The XylS-dependent Pm promoter is transcribed in vivo by RNA polymerase with sigma 32 or sigma 38 depending on the growth phase. Mol Microbiol 31:1105-13.

3. Worsey MJ, Williams PA. 1975. Metabolism of toluene and xylenes by Pseudomonas (putida (arvilla) mt-2: evidence for a new function of the TOL plasmid. J Bacteriol 124:7-13.

4. Domínguez-Cuevas P, Marqués S. Transcriptional Control of the TOL Plasmid Pathways, Handbook of Hydrocarbon and Lipid Microbiology.

5. Ruíz R, Aranda-Olmedo MI, Domínguez-Cuevas P, Ramos-González MI, Marqués S. 2004. Transcriptional regulation of the toluene catabolic pathways, p 509-537, Virulence and Gene Regulation. Springer.

6. Rojo F, Dinamarca MA. 2004. Catabolite repression and physiological control, p 365-387, Virulence and Gene Regulation. Springer.

7. Koutinas M, Lam MC, Kiparissides A, Silva-Rocha R, Godinho M, Livingston AG, Pistikopoulos EN, De Lorenzo V, Dos Santos VAM, Mantalaris A. 2010. The regulatory logic of m-xylene biodegradation by Pseudomonas putida mt-2 exposed by dynamic modelling of the principal node Ps/Pr of the TOL plasmid. Environmental microbiology 12:1705-1718.

8. Koutinas M, Kiparissides A, Silva-Rocha R, Lam M-C, dos Santos VAM, de Lorenzo V, Pistikopoulos EN, Mantalaris A. 2011. Linking genes to microbial growth kinetics—An integrated biochemical systems engineering approach. Metabolic Engineering 13:401-413.

9. Silva-Rocha R, de Jong H, Tamames J, de Lorenzo V. 2011. The logic layout of the TOL network of Pseudomonas putida pWW0 plasmid stems from a metabolic amplifier motif (MAM) that optimizes biodegradation of m-xylene. BMC Systems Biology 5:191.

10. Silva-Rocha R, Tamames J, dos Santos VM, de Lorenzo V. 2011. The logicome of environmental bacteria: merging catabolic and regulatory events with Boolean formalisms. Environmental Microbiology 13:2389-2402.

11. Montero Llopis P, Jackson AF, Sliusarenko O, Surovtsev I, Heinritz J, Emonet T, Jacobs-Wagner C. 2010. Spatial organization of the flow of genetic information in bacteria. Nature 466:77-81.

12. Sanamrad A, Persson F, Lundius EG, Fange D, Gynnå AH, Elf J. 2014. Single-particle tracking reveals that free ribosomal subunits are not excluded from the Escherichia coli nucleoid. Proc Natl Acad Sci U S A 111:11413-8.

13. Nevo-Dinur K, Nussbaum-Shochat A, Ben-Yehuda S, Amster-Choder O. 2011. Translation-independent localization of mRNA in E. coli. Science 331:1081-4.

14. Kannaiah S, Livny J, Amster-Choder O. 2019. Spatiotemporal Organization of the E. coli Transcriptome: Translation Independence and Engagement in Regulation. Mol Cell doi:10.1016/j.molcel.2019.08.013.

15. Moffitt JR, Pandey S, Boettiger AN, Wang S, Zhuang X. 2016. Spatial organization shapes the turnover of a bacterial transcriptome. Elife 5.

16. Libby EA, Roggiani M, Goulian M. 2012. Membrane protein expression triggers chromosomal locus repositioning in bacteria. Proc Natl Acad Sci U S A 109:7445-50.

17. Yang S, Kim S, Kim DK, Jeon An H, Bae Son J, Hedén Gynnå A, Ki Lee N. 2019. Transcription and translation contribute to gene locus relocation to the nucleoid periphery in E. coli. Nat Commun 10:5131.

18. Driessen AJ, Nouwen N. 2008. Protein translocation across the bacterial cytoplasmic membrane. Annu Rev Biochem 77:643-67.

19. Johnson GE, Lalanne J-B, Peters ML, Li G-W. 2020. Functionally uncoupled transcription–translation in Bacillus subtilis. Nature doi:10.1038/s41586-020-2638-5.

20. Kim J, Goni-Moreno A, Calles B, de Lorenzo V. 2019. Spatial organization of the gene expression hardware in Pseudomonas putida. Environ Microbiol 21:1645-1658.

21. Kim J, Perez-Pantoja D, Silva-Rocha R, Oliveros JC, de Lorenzo V. 2016. High-resolution analysis of the m-xylene/toluene biodegradation subtranscriptome of Pseudomonas putida mt-2. Environ Microbiol 18:3327-3341.

22. Abril MA, Michan C, Timmis KN, Ramos JL. 1989. Regulator and enzyme specificities of the TOL plasmid-encoded upper pathway for degradation of aromatic hydrocarbons and expansion of the substrate range of the pathway. J Bacteriol 171:6782-90.

23. Goni-Moreno A, Kim J, de Lorenzo V. 2017. CellShape: A user-friendly image analysis tool for quantitative visualization of bacterial cell factories inside. Biotechnol J 12.

24. Ramos JL, Duque E, Godoy P, Segura A. 1998. Efflux pumps involved in toluene tolerance in Pseudomonas putida DOT-T1E. Journal of bacteriology 180:3323-3329.

25. Bühler B, Witholt B, Hauer B, Schmid A. 2002. Characterization and Application of Xylene Monooxygenase for Multistep Biocatalysis. Applied and Environmental Microbiology 68:560.

26. Vallet-Gely I, Boccard F. 2013. Chromosomal organization and segregation in Pseudomonas aeruginosa. PLoS Genet 9:e1003492.

27. Greated A, Lambertsen L, Williams PA, Thomas CM. 2002. Complete sequence of the IncP-9 TOL plasmid pWW0 from Pseudomonas putida. Environ Microbiol 4:856-71.

28. Goni-Moreno A, Benedetti I, Kim J, de Lorenzo V. 2017. Deconvolution of Gene Expression Noise into Spatial Dynamics of Transcription Factor-Promoter Interplay. ACS Synth Biol 6:1359-1369.

29. Schumacher MA. 2012. Bacterial plasmid partition machinery: a minimalist approach to survival. Curr Opin Struct Biol 22:72-9.

30. Gerdes K, Howard M, Szardenings F. 2010. Pushing and pulling in prokaryotic DNA segregation. Cell 141:927-42.

31. Guynet C, de la Cruz F. 2011. Plasmid segregation without partition. Mob Genet Elements 1:236-241.

32. Marques S, Ramos JL, Timmis KN. 1993. Analysis of the mRNA structure of the Pseudomonas putida TOL meta fission pathway operon around the transcription initiation point, the xylTE and the xylFJ regions. Biochim Biophys Acta 1216:227-36.

33. Leija-Martinez N, Casas-Flores S, Cadena-Nava RD, Roca JA, Mendez-Cabanas JA, Gomez E, Ruiz-Garcia J. 2014. The separation between the 5'-3' ends in long RNA molecules is short and nearly constant. Nucleic Acids Res 42:13963-8.

34. Sinclair MI, Maxwell PC, Lyon BR, Holloway BW. 1986. Chromosomal location of TOL plasmid DNA in Pseudomonas putida. Journal of bacteriology 168:1302-1308.

35. Panke S, Sánchez-Romero JM, de Lorenzo V. 1998. Engineering of quasi-natural Pseudomonas putida strains for toluene metabolism through an ortho-cleavage degradation pathway. Applied and environmental microbiology 64:748-751.

36. Williams PA, Shaw LM, Pitt CW, Vrecl M. 1997. XylUW, two genes at the start of the upper pathway operon of TOL plasmid pWW0, appear to play no essential part in determining its catabolic phenotype. Microbiology 143 (Pt 1):101-7.

37. Bryant JA, Sellars LE, Busby SJ, Lee DJ. 2014. Chromosome position effects on gene expression in Escherichia coli K-12. Nucleic Acids Res 42:11383-92.

38. Chaves JE, Wilton R, Gao Y, Munoz NM, Burnet MC, Schmitz Z, Rowan J, Burdick LH, Elmore J, Guss A, Close D, Magnuson JK, Burnum-Johnson KE, Michener JK. 2020. Evaluation of

chromosomal insertion loci in the Pseudomonas putida KT2440 genome for predictable biosystems design. Metab Eng Commun 11:e00139.

39. Domröse A, Hage-Hülsmann J, Thies S, Weihmann R, Kruse L, Otto M, Wierckx N, Jaeger KE, Drepper T, Loeschcke A. 2019. Pseudomonas putida rDNA is a favored site for the expression of biosynthetic genes. Sci Rep 9:7028.

40. Holtel A, Timmis KN, Ramos JL. 1992. Upstream binding sequences of the XylR activator protein and integration host factor in the xylS gene promoter region of the Pseudomonas TOL plasmid. Nucleic Acids Res 20:1755-62.

41. Perez-Martin J, De Lorenzo V. 1995. Integration host factor suppresses promiscuous activation of the sigma 54-dependent promoter Pu of Pseudomonas putida. Proc Natl Acad Sci U S A 92:7277-81.

42. Troeschel SC, Thies S, Link O, Real CI, Knops K, Wilhelm S, Rosenau F, Jaeger KE. 2012. Novel broad host range shuttle vectors for expression in Escherichia coli, Bacillus subtilis and Pseudomonas putida. J Biotechnol 161:71-9.

43. Marques S, Gallegos MT, Manzanera M, Holtel A, Timmis KN, Ramos JL. 1998. Activation and repression of transcription at the double tandem divergent promoters for the xylR and xylS genes of the TOL plasmid of Pseudomonas putida. J Bacteriol 180:2889-94.

44. Gallegos MT, Marques S, Ramos JL. 1996. Expression of the TOL plasmid xylS gene in Pseudomonas putida occurs from a alpha 70-dependent promoter or from alpha 70- and alpha 54-dependent tandem promoters according to the compound used for growth. J Bacteriol 178:2356-61.

45. Kohler R, Mooney RA, Mills DJ, Landick R, Cramer P. 2017. Architecture of a transcribing-translating expressome. Science 356:194-197.

46. O'Reilly FJ, Xue L, Graziadei A, Sinn L, Lenz S, Tegunov D, Blötz C, Singh N, Hagen WJH, Cramer P, Stülke J, Mahamid J, Rappsilber J. 2020. In-cell architecture of an actively transcribing-translating expressome. Science 369:554-557.

47. Sheng H, Stauffer WT, Hussein R, Lin C, Lim HN. 2017. Nucleoid and cytoplasmic localization of small RNAs in Escherichia coli. Nucleic Acids Res 45:2919-2934.

48. Minton AP. 2001. The influence of macromolecular crowding and macromolecular confinement on biochemical reactions in physiological media. J Biol Chem 276:10577-80.

49. Benny P, Raghunath M. 2017. Making microenvironments: A look into incorporating macromolecular crowding into in vitro experiments, to generate biomimetic microenvironments which are capable of directing cell function for tissue engineering applications. J Tissue Eng 8:2041731417730467.

50. Trevors JT, van Elsas JD, Bej AK. 2013. The Molecularly Crowded Cytoplasm of Bacterial Cells: Dividing Cells Contrasted with Viable but Non-culturable (VBNC) Bacterial Cells. Curr Issues Mol Biol 15:1-6.

51. Parry BR, Surovtsev IV, Cabeen MT, O'Hern CS, Dufresne ER, Jacobs-Wagner C. 2013. The Bacterial Cytoplasm Has Glass-like Properties and Is Fluidized by Metabolic Activity. Cell doi:10.1016/j.cell.2013.11.028.

52. Bakshi S, Siryaporn A, Goulian M, Weisshaar JC. 2012. Superresolution imaging of ribosomes and RNA polymerase in live Escherichia coli cells. Mol Microbiol 85:21-38.

53. Chai Q, Singh B, Peisker K, Metzendorf N, Ge X, Dasgupta S, Sanyal S. 2014. Organization of ribosomes and nucleoids in Escherichia coli cells during growth and in quiescence. J Biol Chem 289:11342-52.

54. Bremer H, Berry L. 1971. Co-transcription of 16S and 23S ribosomal RNA in Escherichia coli. Nat New Biol 234:81-3.

55. Marras SA, Gold B, Kramer FR, Smith I, Tyagi S. 2004. Real-time measurement of in vitro transcription. Nucleic Acids Res 32:e72.

56. Venturelli OS, Tei M, Bauer S, Chan LJG, Petzold CJ, Arkin AP. 2017. Programming mRNA decay to modulate synthetic circuit resource allocation. Nat Commun 8:15128.

57. Kuderova A, Nanak E, Truksa M, Brzobohaty B. 1999. Use of rifampicin in T7 RNA polymerase-driven expression of a plant enzyme: rifampicin improves yield and assembly. Protein Expr Purif 16:405-9.

58. Chong S, Chen C, Ge H, Xie XS. 2014. Mechanism of transcriptional bursting in bacteria. Cell 158:314-326.

59. Yeung E, Dy AJ, Martin KB, Ng AH, Del Vecchio D, Beck JL, Collins JJ, Murray RM. 2017. Biophysical Constraints Arising from Compositional Context in Synthetic Gene Networks. Cell Syst 5:11-24 e12.

60. Kim S, Beltran B, Irnov I, Jacobs-Wagner C. 2019. Long-Distance Cooperative and Antagonistic RNA Polymerase Dynamics via DNA Supercoiling. Cell 179:106-119 e16.

61. Kloc M, Zearfoss NR, Etkin LD. 2002. Mechanisms of subcellular mRNA localization. Cell 108:533-44.

62. Forrest KM, Gavis ER. 2003. Live imaging of endogenous RNA reveals a diffusion and entrapment mechanism for nanos mRNA localization in Drosophila. Curr Biol 13:1159-68.

63. Shav-Tal Y, Darzacq X, Shenoy SM, Fusco D, Janicki SM, Spector DL, Singer RH. 2004. Dynamics of single mRNPs in nuclei of living cells. Science 304:1797-800.

64. Golding I, Cox EC. 2004. RNA dynamics in live Escherichia coli cells. Proc Natl Acad Sci U S A 101:11310-5.

65. Gray WT, Govers SK, Xiang Y, Parry BR, Campos M, Kim S, Jacobs-Wagner C. 2019. Nucleoid Size Scaling and Intracellular Organization of Translation across Bacteria. Cell 177:1632-1648 e20.

66. Campos M, Jacobs-Wagner C. 2013. Cellular organization of the transfer of genetic information. Curr Opin Microbiol 16:171-6.

67. Castellana M, Wilson MZ, Xu Y, Joshi P, Cristea IM, Rabinowitz JD, Gitai Z, Wingreen NS. 2014. Enzyme clustering accelerates processing of intermediates through metabolic channeling. Nat Biotechnol 32:1011-8.

68. Sambrook J, Russell D. 2001. Molecular Cloning: A Laboratory Manual doi:citeulike-article-id:144016. Cold Spring Harbor Laboratory Press.

69. Martinez-Garcia E, de Lorenzo V. 2011. Engineering multiple genomic deletions in Gram-negative bacteria: analysis of the multi-resistant antibiotic profile of Pseudomonas putida KT2440. Environ Microbiol 13:2702-16.

70. de Lorenzo V, Timmis KN. 1994. Analysis and construction of stable phenotypes in gram-negative bacteria with Tn5- and Tn10-derived minitransposons. Methods Enzymol 235:386-405.

71. Kessler B, de Lorenzo V, Timmis KN. 1992. A general system to integrate lacZ fusions into the chromosomes of gram-negative eubacteria: regulation of the Pm promoter of the TOL plasmid studied with all controlling elements in monocopy. Mol Gen Genet 233:293-301.

72. Wong SM, Mekalanos JJ. 2000. Genetic footprinting with mariner-based transposition in Pseudomonas aeruginosa. Proc Natl Acad Sci U S A 97:10191-6.

FIGURES

FIG 1. Interplay between metabolic and regulatory components of the TOL network and visualization of
the transcripts from either the *upper* or the *lower* of the system.

(A) In the presence of *m*-xylene, XylR (expressed from P_R) activates both promoters *Pu* (which
transcribes the *upper* pathway operon) and *Ps,* stimulating expression of *xylS* gene. Also, XylR
expression is negatively auto-regulated. In turn, the XylS protein induces the lower operon by activating
Pm promoter with its effector 3MBz as the metabolite of *m*-xylene. (B) Scheme showing the fluorescent
in situ hybridization (FISH) experiment to visualize mRNAs from the TOL operons of plasmid pWW0. For
this experiment, *P. putida* mt-2 strain was grown in succinate-supplemented minimal medium to the
exponential phase, then further cultured for 2 h with saturating vapors of *m*-xylene to activate the *xyl*
genes. After the fixation of the sample with formaldehyde, the FISH experiment was performed with either
the *upper* (*xylUW*) or the *lower* pathway (*xylX*) probe set, involving the red fluorophore-labeled 48 oligos
complementary to each mRNA sequence. (C) Microscope images of the FISH approach responding to
the *xylUW* mRNA and the *xylX* mRNA, respectively. Phase-contrast (panel 1 and 4), mRNA-red signals
(panel 2 and 5), and the composite images (panel 3 and 6) are shown. Scale bar, 2.5 μm.

1 **FIG 2.** Spatial distribution of the TOL catabolic transcripts with respect to the nucleoid occupation within
2 cells.
3

4
5

(A) The mRNA signals, against the *xylUW* or the *xylX,* were obtained by the FISH assay (panel 2 and 6),
and the nucleoid was stained with DAPI (panel 3 and 7) in the procedure of the experiment to examine
distribution of DNA. The mRNA-red and the DNA-blue channels were merged (panel 4 and 8) to identify
relative position of the transcript with respect to the nucleoid. Phase-contrast images (panel 1 and 5)
corresponding to the red and the blue channels are also shown. Scale bar, 2.5 µm. (B) Using the image
analysis tool as CellShape, fluorescent signals such as mRNA red spots and the nucleoid contour were
identified from exemplary cells appeared in (A). The densest regions of the DAPI-stained regions are
within black lines, while relatively less concentrated DNA regions are depicted with yellow lines. (C) The
abundance of *xyl* mRNAs in the nucleoid-free regions. The *xylUW* (214 spots from 152 cells) or the *xylX*
signals (446 spots from 389 cells) were analyzed concerning the distribution of the nucleoid.
16

FIG 3. Co-visualization of the *xylUW* transcript and the pWW0 plasmid

(A) Sequentially combined RNA-DNA FISH was carried out to visualize both the *xylUW* mRNA and the pWW0 plasmid in the cell. In order to achieve DNA signals with the approach, the tandem copies of the *tet operators* were inserted into the *orf105* locus between the *upper* and *lower* operons and the green-fluorophore tagged *tetO* probe was used for the hybridization in DNA-FISH procedure. (B-D) The mt-2 (pTOL-tetO) strain, exposed to *m*-xylene for 2 h, was fixed for the FISH approach. The resulting images from the experiment were shown with representative cells demonstrating dynamic localization patterns of the mRNA relative to the DNA locus. The microscopy displays phase-contrast (panel 1), the plasmid DNA (green; panel 2), the *xylUW* mRNA (red; panel 3) and the nucleoid (blue; panel 5), respectively. The composite images such as the mRNA/ the plasmid DNA (panel 4), the plasmid DNA/ the nucleoid (panel 6), and the mRNA/ the nucleoid (panel 7) are also shown. Relative intensity was measured along with the mRNA-red and the plasmid-green signals: y-axis refers to relative fluorescent intensity; x-axis indicates pixel distance representing the arrow drawn on the composite image (panel 4). Scale bar, 1 µm.

1   **FIG 4.** Dual labeling of the *xylX* mRNA and the pWW0 plasmid
2
3

4
5
6   (A) The same approach as described in Figure 3A, but red fluorophore-tagged *xylX* probes were applied
7   for the combined RNA-DNA FISH to visualize both the *lower pathway* transcripts and the TOL plasmid.
8   (B-D) Cellular position of the plasmid DNA and the *xylX* mRNA was detected in the example cells by
9   employing the FISH approach. All panels are organized as the same statement of Figure 3.

1 **FIG 5.** Effect of the replicon type on the localization of *xyl* mRNA
2
3

4
5
6
(A) The Paw140 strain, carrying the entire TOL system in the chromosome, was used to identify the
distribution of *xyl* mRNAs by following the same procedure applied for the mt-2 strain. (B) RNA-FISH
microscopy enabled to visualize mRNA signals corresponding to the *xylUW* (panel 2) and *xylX* mRNA
(panel 6), respectively. Their counterpart images such as phase contrast (panel 1 and 5), the nucleoid
(blue; panel 3 and 6), and composite images (panel 4 and 8) are shown. Scale bar, 2.5 µm. (C)
Quantification of mRNA-red foci with respect to the nucleoid occupation. Note that the number of red stop
per cell increased by ~ 30% in *the* Paw140 strain compared to that in the mt-2 strain (Fig. 2C).
14
15
16

1   **FIG 6**. Localization of the *xyl* mRNA transcribed with an orthogonal RNAP
2
3

4
5
6
(A) Sketch of RNA-FISH experiments with the modified plasmid pTOL-PuxT7 in the KT2440•T7 strain,
expressing T7 RNAP by the chromosomal *lacI^q^/Plac-T7pol* cassette. After exposure of the cells for 2 h
with *m*-xylene, *xyl* mRNAs were visualized with the FISH. (B) Images obtained from the FISH experiment
by using either the *xylUW* or the *xylX* probe set are shown: mRNAs (red; panel 2 and 6) , DAPI-stained
DNA (blue; 3 and 7) and merged signals are presented (panel 4 and 8) in the fixed cells (panel 1 and 5).
Scale bar, 2.5 µm. (C) Quantification of overlapping regions between mRNA-red signals and DAPI-blue
signals with sub-pixel precision. Over 100 exemplary cells were considered for the colocalization
analysis.
15
16

1 **FIG 7.** Effect of modulated macromolecular crowding by inhibiting bacterial transcription on the spatial
2 distribution of the *xylUW* mRNA
3

4
5
The *P. putida* KT2440•T7 (pTOL-PuxT7) strain was tested to visualize the *xylUW* mRNA under
transcription inhibited condition. After treatment of the cells for 2 h with *m*-xylene and rifampicin (200 µg
ml^{-1}), RNA-FISH experiment was proceeded. The resulting images were obtained as following: (A)
Phase-contrast. (B) Red signals indicate fluorescently labeled oligos that hybridize to the *xylUW* mRNA,
which are dispersed throughout the cell. (C) DAPI-staining. (D) The merged picture between mRNA-red
and DAPI-blue channels. (E) The subset image of panel (C) was also taken with relatively shorter
exposure time to rule out the blurred signals are not due to the light diffraction. Scale bar, 2.5 µm.
13
14
15
16