Protein Phosphatase 1 in association with Bud14 inhibits mitotic exit in *Saccharomyces*

cerevisiae

Dilara Kocakaplan[1, ‡], Hüseyin Karabürk[1, ‡], Cansu Dilege[1], Idil Kirdok[1], Şeyma Nur

Bektaş[1], Ayse Koca Caydasi[1, *]

[1]Department of Molecular Biology and Genetics, Koç University, Istanbul, Turkey

[‡]These authors contributed equally to this work

*Correspondence to: aykoca@ku.edu.tr

Abstract

Mitotic exit in budding yeast is dependent on correct orientation of the mitotic spindle along

the cell polarity axis. When accurate positioning of the spindle fails, a surveillance mechanism

named the Spindle Position Checkpoint (SPOC) prevents cells from exiting mitosis. Mutants

with a defective SPOC become multinucleated and lose their genomic integrity. Yet, a

comprehensive understanding of the SPOC mechanism is missing. In this study, we identified

the type 1 protein phosphatase, Glc7, in association with its regulatory protein Bud14 as a novel

checkpoint component. We further showed that Glc7-Bud14 promotes dephosphorylation of

the SPOC effector protein Bfa1. Our results suggest a model in which two mechanisms act in

parallel for a robust checkpoint response: first, the SPOC kinase Kin4 isolates Bfa1 away from

the inhibitory kinase Cdc5 and second, Glc7-Bud14 dephosphorylates Bfa1 to fully activate

the checkpoint effector.

Introduction

Budding yeast undergoes asymmetric cell division in every cell cycle and is an intrinsically polarized cell. In every cell cycle, a new cell buds on the mother cell and pinches of from this location at the end of the cell division, giving rise to a young daughter cell and an old mother cell. In order for the daughter cell to receive one copy of the duplicated genetic material, budding yeast has to segregate its chromosomes along its mother-to-bud polarity axis, which requires positioning of the mitotic spindle apparatus along this direction. The correct orientation of the mitotic spindle is achieved by two redundant pathways, namely the "early" and "late" pathways, which are dependent on the Adenomatous Polyposis Coli (APC) homolog Kar9 and the microtubule motor protein Dyn1 respectively (Adames & Cooper, 2000; Beach *et al*, 2000; Li *et al*, 1993; Siller & Doe, 2009).

Correct orientation of the mitotic spindle is monitored by a mitotic checkpoint named the Spindle Position Checkpoint (SPOC) in budding yeast. SPOC prevents cell cycle progression of cells that fail to orient their spindle in the mother-to-bud direction, and hence provides time for cells to correct their spindle orientation before mitotic exit (Adames *et al*, 2001; Bardin *et al*, 2000; Bloecher *et al*, 2000; Pereira *et al*, 2000; Yeh *et al*, 1995). Yeast cells with defects in the spindle positioning pathways or cells with impaired microtubule function rely on the SPOC to maintain their ploidy. Absence of SPOC causes multinucleation, enucleation and aneuploidy in budding yeast. A SPOC like checkpoint also exists in Drosophila and may be present in higher eukaryotes (Cheng *et al*, 2008; O'Connell & Wang, 2000; Pereira & Yamashita, 2011).

In response to spindle mispositioning, SPOC inhibits cell cycle progression by inhibiting the Mitotic Exit Network (MEN) (Baro *et al*, 2017; Caydasi *et al*, 2010a; Caydasi & Pereira, 2012; Weiss, 2012). MEN is a GTPase driven signaling pathway that is essential for mitotic exit in budding yeast (Jaspersen *et al*, 1998; Shou *et al*, 1999). A single conserved phosphatase, named

Cdc14 triggers mitotic exit in budding yeast through inactivation of the mitotic Cyclin

Dependent Kinase (CDK) and dephosphorylation of CDK targets (Manzano-Lopez & Monje-

Casas, 2020). At the anaphase onset, a transient wave of Cdc14 activation is triggered by the

cdcFourteen Early Anaphase Release (FEAR) network; whereas later in anaphase the MEN

promotes fully activation of Cdc14 (Jaspersen *et al.*, 1998; Lee *et al*, 2001a; Rock & Amon,

2009; Shou *et al.*, 1999). A Ras-like small GTPase named Tem1 is the main switch located

near the top of the MEN (Lee *et al.*, 2001a; Morishita *et al*, 1995; Scarfone & Piatti, 2015;

Shirayama *et al*, 1994b). When bound to GTP, Tem1 initiates the MEN signaling by recruiting

the Cdc15 kinase to the spindle pole bodies (centrosome equivalent in yeast, SPB) which

therein activates Dbf2 kinase in association with its regulatory subunit Mob1 (Asakawa *et al*,

2001; Cenamor *et al*, 1999; Visintin & Amon, 2001).

SPOC inhibits the MEN through inactivation of Tem1 by a bipartite GTPase Activating

Complex (GAP) composed of Bfa1 and Bub2 proteins (Geymonat *et al*, 2002). Activity of the

Bfa1-Bub2 GAP complex is tightly regulated at the level of Bfa1 phosphorylation. In cells with

a correctly positioned anaphase spindle, the polo like kinase Cdc5 phosphorylates Bfa1 to

prevent Bfa1-Bub2 GAP activity towards Tem1 (Hu *et al*, 2001). CDK also phosphorylates

Bfa1 in an activating manner until anaphase when this phosphorylation is removed in a FEAR

related manner (Caydasi *et al*, 2017). In the presence of spindle misalignment, however, Kin4

kinase phosphorylates Bfa1 (Maekawa *et al*, 2007). This phosphorylation prevents inhibitory

phosphorylation of Bfa1 by Cdc5 kinase (Pereira & Schiebel, 2005). Consequently, Bfa1-Bub2

GAP complex activity is promoted, Tem1 activity is prevented, and exit from mitosis is

inhibited.

SPBs, serving as a scaffold, play a key role in mitotic exit and its timely regulation by the

spindle position (Campbell *et al*, 2020; Gruneberg *et al*, 2000; Pereira & Schiebel, 2001;

Valerio-Santiago & Monje-Casas, 2011). When the spindle is properly oriented, Bfa1-Bub2

and Tem1 reside in a preferential, asymmetrical manner at the SPB that moves to the daughter

cell (dSPB) (Molk *et al*, 2004; Pereira *et al.*, 2000). On contrary, Kin4 localizes exclusively to

the SPB that stays in the mother cell (mSPB) during an unperturbed anaphase (D'Aquino *et al*,

2005; Pereira & Schiebel, 2005). Spindle misalignment alters SPB localization of SPOC

components. Upon mispositioning of the mitotic spindle, Kin4 localizes to both SPBs (Gislene

Pereira & Schiebel, 2005). Phosphorylation of Bfa1 by Kin4 promotes binding of the 14-3-3

family protein Bmh1 to Bfa1 which causes dissociation of Bfa1-Bub2 from SPBs (Caydasi *et*

al, 2012; Caydasi *et al*, 2014; Caydasi & Pereira, 2009; Monje-Casas & Amon, 2009). Hence,

upon spindle mispositioning, the amount of Bfa1-Bub2 at both SPBs decreases, while their

cytoplasmic levels increase. The release of Bfa1-Bub2 from SPBs is thought to keep Bfa1-

Bub2 away from the inhibitory kinase Cdc5 and thus prevent inactivation of Bfa1-Bub2 by

Cdc5. However, what dephosphorylates the Cdc5 phosphorylated Bfa1 remains elusive.

Here, we identified Bud14 as a novel SPOC protein. Deletion of *BUD14* rescued cold

sensitivity of *lte1Δ* cells, lethality of mitotic exit deficient *lte1Δ spo12Δ* cells, and growth

defects of MEN temperature sensitive mutants *cdc15-1*, *dbf2-2*, *mob1-67*. Our data showed

that *bud14Δ* cells accumulated multinucleated phenotypes when spindle positioning was

impaired in cells via deletion of *KAR9* or *DYN1*. Fluorescence time-lapse microscopy revealed

that *bud14Δ* cells failed to delay mitotic exit in response to spindle mispositioning. We

observed an additive decrease in anaphase duration of *kin4Δ bud14Δ* cells with misaligned

spindles compared to the cells bearing single gene deletions of *KIN4* or *BUD14* suggesting that

Kin4 and Bud14 work in parallel in SPOC. We further found that the mitotic exit inhibitory

function Bud14 required its association with the type 1 protein phosphatase, Glc7. A

temperature sensitive version of Glc7 (*glc7-12*) and a version of Bud14 that cannot interact

with Glc7 (*bud14-F379A*) caused SPOC deficiency and rescued growth of mitotic exit mutants.

Yeast two hybrid data indicated an interaction between Bfa1 and Bud14 that required the

presence of Bub2, which suggests that Bud14 may recognize Bfa1-Bub2. Intriguingly, *bud14Δ*

and *bud14-F379A* cells had more Bfa1-Bub2 localized at the dSPBs, suggestive of a role for

Bud14-Glc7 in limiting SPB-bound levels of Bfa1-Bub2. Similarly to the wild type cells, upon

spindle mispositioning, levels of Bfa1 at the dSPB was able to decrease in *bud14Δ* cells, unlike

kin4Δ cells, which also supported separate functions of Bud14 and Kin4 in SPOC. We further

observed that *lte1Δ* cells lacking Bud14 accumulated hyperphosphorylated Bfa1 forms during

anaphase. Furthermore, overexpression of *BUD14* but not *bud14-F309A* caused a reduction in

this hyperphosphorylated forms of Bfa1. Finally, through in vitro phosphatase assays we

showed that Glc7-Bud14 promoted dephosphorylation of hyperphosphorylated Bfa1. Our data

altogether indicates that Glc7-Bud14 is a novel mitotic exit inhibitor in budding yeast that

works upon spindle mispositioning. We propose a checkpoint model in which two independent

mechanisms activate the checkpoint effector: first, the SPOC kinase Kin4 prevents further

Cdc5 phosphorylation of Bfa1 by rapidly removing Bfa1 away from the Cdc5 kinase and

second, Glc7-Bud14 dephosphorylates Bfa1 to activate the Bfa1-Bub2 GAP complex. Both

mechanisms are crucial for rapid activation of the checkpoint effector Bfa1-Bub2 upon

mispositioning of the anaphase spindle.

Results

Bud14 is a Novel Mitotic Exit Inhibitor

Lte1 is a mitotic exit activator that becomes essential for mitotic exit at cold temperatures (<16 °C) (Shirayama *et al*, 1994a). Although the exact function of Lte1 in mitotic exit is not fully understood, one of the ways that Lte1 promotes mitotic exit is by preventing binding of the SPOC kinase Kin4 to the SPB that has migrated into the bud and by inhibiting its activity therein (Bertazzi *et al*, 2011; Falk *et al*, 2011). At physiological temperatures (i.e. 30 °C) *lte1Δ* cells rely on the presence of the FEAR network to exit mitosis. Deletion of FEAR network components such as *SPO12* or *SLK19* in *lte1Δ* cells causes lethality due to failure of mitotic exit (Stegmeier *et al*, 2002). Lethality of the *lte1Δ spo12Δ* double mutants can be rescued by deletion of the known mitotic exit inhibitors *BFA1*, *BUB2* and *KIN4* (D'Aquino *et al.*, 2005; Stegmeier *et al.*, 2002). In order to identify novel mitotic exit inhibitors, we designed a genetic screen that looked for single gene deletions rescuing *lte1Δ spo12Δ* lethality. In addition to the known mitotic exit inhibitors *BFA1*, *BUB2* and *KIN4*, we identified *BUD14* as a novel gene that contributes to the mitotic arrest of *lte1Δ spo12Δ* cells (data not shown). Deletion of *BUD14* rescued lethality of *lte1Δ spo12Δ* cells (Figure 1A). Deletion of *BUD14* also rescued the cold sensitivity of *lte1Δ* cells (Figure 1B). This data indicates that Bud14 is a novel inhibitor of mitotic exit.

We next asked whether deletion of *BUD14* rescues growth lethality of mitotic exit network (MEN) mutants. For this, we deleted *BUD14* in several MEN temperature sensitive (ts) mutants (Jaspersen *et al.*, 1998) and compared their growth at different temperatures. To allow for better comparison, we also deleted *BFA1*, a known mitotic exit inhibitor, in the same MEN-ts mutants (Caydasi *et al.*, 2017; Hu *et al.*, 2001; Pereira *et al.*, 2000; Scarfone *et al*, 2015; Wang *et al*, 2000). In line with its mitotic exit inhibitory role, deletion of *BFA1* rescued the lethality of

mob1-67, dbf2-2, cdc15-1 and cdc5-10 ts mutants at 33 °C, 37 °C, 35 °C and 37 °C respectively

(Figure 1C). Deletion of *BUD14* also promoted the growth of *mob1-67*, *dbf2-2*, and *cdc15-1*

mutants at 33 °C, 35 °C and 35 °C respectively (Figure 1C). In all cases, growth rescue by

BUD14 deletion was milder than that of *BFA1* deletion. Furthermore, *BUD14* deletion did not

rescue the growth of the *cdc5-10* while *BFA1* deletion did (Figure 1C), suggesting that the

mitotic exit inhibitory function of Bud14 does not contribute to *cdc5-10* lethality. Deletion of

neither *BFA1* nor *BUD14* rescued the growth of the *cdc14-2* mutant (Figure 1C). Cdc14 is the

essential phosphatase that is activated at the very downstream of the MEN pathway. The fact

that *bud14Δ* and *bfa1Δ* does not rescue *cdc14-2* supports their function upstream of the MEN

pathway.

BUD14 is essential for SPOC functionality

The known mitotic exit inhibitors Bfa1, Bub2 and Kin4 are key components of the Spindle

Position Checkpoint (SPOC), which is a surveillance mechanism that inhibits the MEN when

anaphase spindle fails to orient in the mother-to-daughter direction. We thus asked whether the

novel mitotic exit inhibitor Bud14 has a role in SPOC. To address this, we first employed a

population-based approach. Cells with a functional SPOC, arrest in anaphase with

mispositioned nuclei upon spindle mispositioning, whereas SPOC deficient cells such as *bfa1Δ*

or *kin4Δ* continue cell cycle progression resulting in formation of multinucleated and

enucleated cells. We assayed SPOC deficiency in cell populations as a measure of frequency

of multinucleation, relative to that of the spindle mispositioning (SPOC deficiency index).

Kar9- and Dynein-dependent spindle positioning pathways act redundantly to align the spindle

along mother-to-daughter polarity axis in budding yeast (Eshel *et al*, 1993; Li *et al.*, 1993;

Miller & Rose, 1998). To induce spindle mispositioning, we first made use of *kar9Δ* cells. As

expected, deletion of known SPOC components *KIN4* or *BFA1* in the *kar9Δ* background

yielded a remarkably high degree of SPOC deficiency index (Figure 2A). Notably, deletion of

BUD14 in the *kar9Δ* background also caused a significantly high SPOC deficiency index

(Figure 2A). Similar results were obtained using a *dyn1Δ* background (Figure 2B). These data

suggests that *bud14Δ* cells are defective in SPOC.

We then employed a single cell analysis approach to assay SPOC functionality. For this, we

analyzed the timing of mitotic exit in *bud14Δ* cells during spindle misalignment and normal

alignment in a *GFP-TUB1 kar9Δ* cell background. *GFP-TUB1* served as a spindle marker.

kar9Δ population contains cells with misaligned spindles as well as cells with normally aligned

spindles, and thus allows analysis of anaphase duration in both cases. Cells were imaged

through fluorescence time-lapse microscopy for 90 minutes with 1 min time-resolution at 30

ºC. Anaphase duration was calculated as the time elapsed between the start of fast spindle

elongation and the spindle breakage, which are indications of anaphase onset and mitotic exit

respectively. In cells with a normally positioned spindle, mean anaphase duration of all

analyzed cell types were similar - ranging from 18 to 19 minutes with a standard deviation of

2 minutes (Figure 2C, Source data Figure 2C). The majority of *kar9Δ* cells with a misaligned

spindle stayed arrested with an intact spindle during the time-lapse movie (Figure 2C, *kar9Δ*

red triangles), whereas *kin4Δ kar9Δ* cells with misaligned spindles broke their anaphase spindle

in average 27±7 minutes (mean± standard deviation) after the onset of anaphase (Figure 2C,

kar9Δ kin4Δ black squares). *bud14Δ kar9Δ* cells with misaligned spindles also broke their

spindle and with a mean anaphase duration of 23 ± 4 minutes (mean± standard deviation)

(Figure 2C, *kar9Δ bud14Δ*, black squares). These data altogether suggest that Bud14 is needed

as a part of the SPOC mechanism to delay mitotic exit upon spindle mispositioning.

Function of Bud14 in SPOC is parallel to Kin4

Bfa1-Bub2 is the most downstream component of SPOC and Kin4 is the key SPOC kinase that resides upstream of Bfa1 to promote Bfa1-Bub2 GAP activity upon spindle misalignment. We next asked whether Bud14 work parallel to Kin4 in SPOC. We reasoned that if Bud14 and Kin4 work in parallel, concomitant loss of Bud14 and Kin4 would cause a greater SPOC deficiency index than individual loss of Bud14 or Kin4. SPOC deficiency index of *kin4Δ bud14Δ kar9Δ* was significantly higher than that of *kin4Δ kar9Δ* or *bud14Δ kar9Δ* (Figure 2A). Likewise, SPOC deficiency phenotype was more prominent in *kin4Δ bud14Δ dyn1Δ* cells than in *kin4Δ dyn1Δ* or *bud14Δ dyn1Δ* cells (Figure 2B). Thus, *KIN4* and *BUD14* deletion showed an additive effect with respect to SPOC deficiency. Of note, deletion of *BUD14* in *bfa1Δ kar9Δ* cells did not cause any increase in SPOC deficiency index compared to *bfa1Δ kar9Δ* cells (Figure 2A). These data are suggestive of a role for Bud14 parallel to Kin4 and upstream of Bfa1 and in SPOC.

Additive effect of *KIN4* and *BUD14* deletion was also evident in anaphase duration comparisons. Even though *BUD14* deletion caused mitotic exit of cells with misaligned spindles, duration of anaphase in *bud14Δ kar9Δ* cells was significantly longer during spindle misalignment than during spindle normal alignment (Figure 2C, *bud14Δ*, squares vs circles). In addition, one out of 18 cells was able to hold the SPOC arrest longer than an hour. Similar phenomenon was also observed in *kar9Δ kin4Δ* cells (Figure 2C, *kin4Δ*, squares vs circles) and was also reported before (Caydasi *et al.*, 2017; Falk *et al*, 2016b). This indicates that in the absence of Bud14 or Kin4, cells are still able to delay mitotic exit in response to spindle positioning defects, albeit for short time. On contrary, when both *KIN4* and *BUD14* were deleted, cells exited mitosis with the same timing regardless of the spindle position (Figure 2C, *kar9Δ bud14Δ kin4Δ* black squares vs circles). Thus, concomitant deletion of *KIN4* and *BUD14*

results in a more severe loss of SPOC functionality than single deletion of either *KIN4* or

BUD14. This data is in line with our conclusion that Kin4 and Bud14 act in parallel to promote

SPOC arrest.

To rule out the possibility that Bud14 may also be acting upon Kin4, we next analyzed the

effect of Bud14 on Kin4 functionality. For this, we first analyzed cells overexpressing Kin4.

Kin4 overexpression blocks exit from mitosis through constitutive Bfa1-Bub2 activation and

thus results in lethality (D'Aquino *et al.*, 2005). This lethality can be rescued by deletion of

BFA1 or *BUB2,* which are the most downstream SPOC components, and by deletion of *RTS1*

or *ELM1*, which are Kin4 regulators, and by deletion of *BMH1*, which works downstream of

KIN4 (Caydasi *et al*, 2010b; Caydasi *et al.*, 2014). Unlike aforementioned gene deletions,

deletion of *BUD14* did not rescue the lethality of Kin4 overexpression (Figure 2 – Figure

Supplement 1A). Thus, overproduced Kin4 is still able to activate Bfa1 constitutively in the

absence Bud14. Next, we analyzed the effect of Bud14 on Kin4 localization. Kin4 localized to

SPBs in response to microtubule depolymerization independently of the presence of Bud14

(Figure 2 – Figure Supplement 1B). Likewise, Kin4 localized to SPBs during anaphase in *lte1Δ*

cells both in the presence and absence of Bud14 (Figure 2 – Figure Supplement 1C). Kin4

mobility on SDS-PAGE was not altered in cells lacking Bud14 either (Figure 2 – Figure

Supplement 1D). These data indicate that Bud14 does not work upstream of Kin4. As we will

describe later, analysis of Bfa1 localization during spindle misalignment also supported that

Kin4 regulation of Bfa1 takes place independently of Bud14 (Figure 6D, Figure 6- Supplement

3). Taken altogether, our data support a model where Bud14 acts in a pathway parallel to Kin4

in SPOC.

Bud14 function in SPOC is independently of its role in actin regulation

Bud14 regulates actin cable formation through its formin-regulatory motif (amino acids 135-150) (Chesarone *et al*, 2009; Eskin *et al*, 2016). Bud14's function in controlling formin activity also requires its interaction with KELCH domain proteins Kel1 and Kel2 (Gould *et al*, 2014). We addressed the contribution of actin regulatory role of Bud14 to its mitotic exit inhibitory function through two independent approaches. First, we made use of the *bud14-5A* mutant whose formin-regulatory motif is disrupted by five alanine amino acid substitutions at Bud14 residues 135 and 137–140 (Eskin *et al.*, 2016). Ectopic expression of *bud14-5A* and *BUD14* on a centromeric yeast plasmid (pRS416) under *BUD14* endogenous/native promoter rescued the SPOC deficient phenotype of *bud14Δ kar9Δ* cells (Figure 3A) (Figure 3 – Figure supplement 1A). Similarly ectopic expression of *bud14-5A* in *lte1Δ bud14Δ* cells complemented *BUD14* deletion by restoring cold sensitivity of *lte1Δ* (Figure 3B). Thus, *bud14-5A* mutant, which is deficient in formin binding is functional in terms of mitotic exit inhibition. Of note, expression of *bud14-5A* caused a significantly greater extend of reduction in SPOC deficiency index of *bud14Δ kar9Δ* cells than expression of *BUD14* (Figure 3A). Furthermore, Bud14-5A interacted with Glc7 more readily in yeast two hybrid assays (Figure 3 – Figure supplement 1B), which may indicate a link between Glc7-Bud14 interaction and SPOC robustness.

Second, we made used of *kel1Δ* and *kel2Δ* cells. Kel1, Kel2 and Bud14 form a stable complex that regulate the formin Bnr1 and any missing protein from this complex causes actin cable defects to the same extend (Gould *et al.*, 2014). We reasoned that if Kel1-Kel2-Bud14 complex, that is crucial for actin regulation, is also important for SPOC functionality, deletion of *KEL1* or *KEL2* to disturb the complex would also cause SPOC deficiency similar to the *bud14Δ* cells. However, this was not the case. Unlike *bud14Δ kar9Δ* cells, *kel1Δ kar9Δ* or *kel2Δ kar9Δ* cells

did not display SPOC deficiency (Figure 3C). These data altogether suggest that Bud14

function in SPOC is independent of its role in actin cable regulation.

Interestingly, deletion of *KEL1* also rescues cold sensitivity of *lte1Δ* similar to the deletion of

BUD14 (Hofken & Schiebel, 2002; Seshan *et al*, 2002) (Figure 3D). Synthetic lethality of

lte1Δ spo12Δ on the other hand, can only be slightly rescued by the deletion of *KEL1*, whereas

deletion of *BUD14* in *lte1Δ spo12Δ* cells causes a more pronounced growth rescue phenotype

(Figure 3E). Deletion of *KEL2* however does not rescue *lte1Δ spo12Δ* lethality, while it only

has a minor impact on cold sensitivity of *lte1Δ* cells (Figure 3D-E). These data agree with

previous reports that Kel1 and Kel2 have a mitotic exit inhibitory function (Geymonat *et al*,

2010; Hofken & Schiebel, 2002; Seshan *et al.*, 2002) and suggest that this function of Kel1 and

Kel2 in mitotic exit inhibition differs from that of Bud14 as Kel1/Kel2 are not influencing the

SPOC.

Bud14 inhibits mitotic exit through its interaction with Glc7

Bud14 is a regulatory subunit of the type I Protein phosphatase (PP1) Glc7 (Cullen & Sprague,

2002; Knaus *et al*, 2005; Lenssen *et al*, 2005). We next asked whether Bud14 exerts its mitotic

exit inhibitory role through Glc7. To address this question, we used mutants with impaired

Glc7-Bud14 function. First, we used a mutant version of Bud14 that cannot bind Gcl7 (*bud14-*

F379A) (Knaus *et al.*, 2005) (Figure 3- Figure Supplement 1). Introduction of the wild type

BUD14 but not the *bud14-F379A* mutant on a centromeric plasmid restored cold sensitivity of

the *lte1Δ bud14Δ* (Figure 4A). Likewise, expression of *BUD14* but not *bud14-F379A* on a

centromeric plasmid under *BUD14* endogenous/native promoter rescued the SPOC deficient

phenotype of *bud14Δ kar9Δ* cells (Figure 4B). These data suggests that Glc7-Bud14 interaction

is essential for Bud14 function in SPOC.

Next, we used the temperature sensitive *glc7-12* mutant (Andrews & Stark, 2000; Cheng &

Chen, 2010) to further confirm the contribution of Glc7 to mitotic exit inhibition. *glc7-12*

exhibits a growth defect at 33-35 °C and is lethal at 37 °C, as an indication of Glc7 partial and

full inactivation at these temperatures respectively (Figure 4 - Supplement 1A, YPD plates)

(Andrews & Stark, 2000; Cheng & Chen, 2010). Partial inactivation of *glc7-12* at 33-35 °C is

also evident when examining other phenotypes attributed to Glc7 deficiency (Andrews &

Stark, 2000; Cheng & Chen, 2010), such as sensitivity to hydroxyurea (Figure 4 – Figure

Supplement 1A, HU plates) and accumulation of large-budded cells with single nucleus (Figure

4 – Figure Supplement 1B). At 18 °C, which is a permissive temperature for *glc7-12, glc7-12*

lte1Δ spo12Δ cells failed to form colonies (Figure 4C) replicating the lethality of *lte1Δ spo12Δ*

(Figure 1A). However, at the semi-permissive temperatures of *glc7-12* (33 °C-35 °C), *glc7-12*

lte1Δ spo12Δ cells were able to form colonies (Figure 4C). Thus, similar to the deletion of

BUD14, partial inactivation of Glc7 rescues lethality of *lte1Δ spo12Δ*. Furthermore, *glc7-12*

kar9Δ cells were SPOC deficient at 33 °C (Figure 4D). Thus, we conclude that Glc7-Bud14

activity is crucial for SPOC function.

Bud14 interacts with Bfa1 in the yeast two hybrid system

We sought to understand the target of Bud14 among SPOC proteins. For this, we performed a

yeast two hybrid assay. We used Bfa1, Bub2 and Kin4 as bait proteins and Kel1 as a positive

control in this assay. Formation of a blue color indicates activation of the ß-galactosidase gene

and hence physical interaction of LexA DNA binding domain fused with Bud14 (PMM5-

Bud14) and Gal4-Activation Domain fused with the bait protein (pMM6-Bait protein). We

observed blue color formation in cells bearing pMM5-Bud14 and pMM6-Kel1 as expected due

to their known physical interaction (Gould *et al.*, 2014; Krogan *et al*, 2006; Neller *et al*, 2015).

We observed a similar interaction between Bud14 and Bfa1 but not Bub2 or Kin4 (Figure 5A).

We could, however, not detect an interaction between Glc7 and Bfa1, Bub2 or Kin4 in our

yeast two hybrid system (Figure 5 – Figure Supplement 1). Intriguingly, deletion of *BUB2* but

not *KIN4* ceased blue color formation in cells with pMM5-Bud14 and pMM6-Bfa1, indicating

that Bud14-Bfa1 interaction is dependent on the presence of Bub2 (Figure 5B).

We next analyzed the interaction of Bfa1 with mutant forms of Bud14. Bud14-F379A that

cannot bind Glc7 (Figure 3 – Figure Supplement 1B) (Knaus *et al.*, 2005), interacted with Bfa1

similarly to the wild-type Bud14 in our yeast two hybrid setup (Figure 5C). Bud14 formin-

binding mutant (Bud14-5A) also interacted with Bud14 equally well (Figure 5C). Bud14

contains an SH3 domain at its amino terminal (amino acids 262-318), deletion of which

diminishes Bud14 bud cortex localization and also interferes with Bud14-Glc7 interaction

(Knaus *et al.*, 2005) (Figure 3 – Figure Supplement 1A). Notably, deletion of Bud14 SH3

domain also caused a decrease in Bud14-Bfa1 interaction (Figure 5C). Consequently, *bud14-*

SH3Δ was SPOC deficient (Figure 3A) and rescued *lte1Δ* cold sensitivity (Figure 3B). Thus,

SH3 domain of Bud14 is important not only for Bud14 cortex localization but also for Bud14-

Glc7 and Bud14-Bfa1 interaction and hence for its function in SPOC.

Bud14 limits the amount of Bfa1 at dSPBs but does not affect Bfa1 symmetry upon

spindle mispositioning

So far, we showed that Bud14 has a role in SPOC through its interaction with Glc7. We also

showed that it interacts with Bfa1. We next asked whether Bud14-Glc7 affects Bfa1 function.

SPOC functionality relies on a radical change in SPB localization of Bfa1-Bub2. When the

spindle is properly oriented, Bfa1-Bub2 preferentially localizes to the SPB that moves to the

daughter cell (dSPB). Upon spindle misalignment Bfa1-Bub2 is released from dSPB, hence a

low level of Bfa1-Bub2 is attained at both SPBs. Thus, we analyzed the effect of Bud14 on

SPB localization of Bfa1-Bub2. First, we compared the SPB-bound levels of Bfa1-GFP at

SPBs in wild type and *bud14Δ* cells with normally aligned anaphase spindles (spindle length

>3μm) (Figure 6A). To our surprise, more Bfa1-GFP was localized at dSPBs in *bud14Δ* cells

compared to wild type cells. Importantly, increased Bfa1-GFP signal at the dSPB of *bud14Δ*

cells returned to wild type levels when *BUD14* was introduced in these cells on a plasmid

(Figure 6B). Unlike *BUD14*, Introduction of the *bud14*-F379A in *bud14Δ* cells, did not cause

a significant reduction in the levels of Bfa1 at the dSPB (Figure 6C). These data indicate that

presence of Glc7-Bud14 reduces the dSPB bound Bfa1 levels.

Bub2 and Tem1 localization at SPBs also depend on Bfa1 (Pereira *et al.*, 2000). Accordingly,

Bub2-GFP and Tem1-GFP localization at dSPB was also remarkably higher in *bud14Δ* cells

than wild type cells (Figure 6 – Figure Supplement 1A-B). Such increase in SPB-bound levels

of Bfa1-Bub2 and Tem1 was not restricted to anaphase but was also present before anaphase

onset (spindle length = 1,5-2 μm) (Figure 6 – Figure Supplement 1C).

We next asked, whether the increased Bfa1-Bub2 and Tem1 recruitment to the dSPBs observed

in *bud14Δ* cells may cause premature activation of the MEN in these cells. Mitotic exit is

preceded by SPB localization of Cdc15 and Dbf2-Mob1 (Frenz *et al*, 2000; Luca *et al*, 2001;

Pereira *et al*, 2002; Visintin & Amon, 2001; Yoshida & Toh-e, 2001). We assessed MEN

activity using the SPB bound levels of Mob1-GFP owing to its brighter fluorescence signal

(Caydasi *et al.*, 2012). Mob1-GFP did not prematurely localize to SPBs in *bud14Δ* cells (Figure

6 – Figure Supplement 2A-C). The levels of Mob1-GFP at the SPBs of *bud14Δ* cells were not

altered in anaphase either (Figure 6- Figure Supplement 2D-E). Thus, we conclude that despite

the elevated levels of Bfa1-Bub2 and Tem1 at the dSPB, MEN activity is not enhanced by

BUD14 deletion in cells with normally aligned spindles.

We next analyzed SPB-bound Bfa1 levels in cells with misaligned anaphase spindles. Similar

to the situation in cells with correctly aligned spindles, *bud14Δ kar9Δ* cells with misoriented

spindles had more Bfa1-GFP at their dSPBs than *kar9Δ* cells (Figure 6 – Figure Supplement

1D). Thus, more Bfa1 localizes at the dSPB in the absence of Bud14 regardless of the spindle

position and the mitotic stage.

When the anaphase spindle elongates in the wrong direction, phosphorylation of Bfa1 by Kin4

causes dissociation of Bfa1-Bub2 from dSPB. Consequently, upon spindle mispositioning,

SPB localization of Bfa1-GFP changes from asymmetric (more at the dSPB than mSPB) to

more symmetric (similar at both SPBs). We thus asked whether *bud14Δ* cells were able to

change their Bfa1 localization from asymmetric to symmetric in response to spindle

misalignment. As a measure of signal asymmetry at SPBs, we calculated Bfa1-GFP asymmetry

index as the ratio of Bfa1-GFP intensity at SPBs of *kar9Δ* cells with correctly aligned and

misaligned anaphase spindles (Figure 6D). In response to spindle mispositioning, Bfa1-GFP

asymmetry was drastically reduced in *kar9Δ* cells, whereas it did not significantly change in

kin4Δ kar9Δ cells (Figure 6D). Similar to *kar9Δ* cells, Bfa1-GFP asymmetry was also greatly

decreased in *bud14Δ kar9Δ* cells when the spindle was mispositioned (Figure 6D). We also

analyzed the behavior of Bfa1-GFP through time-lapse experiments. Consistent with previous

reports, SPB bound levels of Bfa1-GFP in *kar9Δ* cells drastically decreased within few minutes

after elongation of the anaphase spindle in the wrong direction (Caydasi & Pereira, 2009)

(Figure 6 – Figure Supplement 3A). Bfa1-GFP also rapidly decreased at SPBs of *bud14Δ kar9Δ*

cells (Figure 6 – Figure Supplement 3B), which was not the case for *kar9Δ kin4Δ* cells (Figure

6 – Figure Supplement 3C). Thus, unlike Kin4, Bud14 does not promote a change in SPB

localization of Bfa1 in response to spindle mispositioning, but rather limits the levels of SPB-

bound Bfa1 regardless of the spindle position. These observations altogether suggest a role for

Bud14-Glc7 in controlling Bfa1 sequestration at the SPB in a way differently than Kin4 and

further support the conclusion that Kin4 and Bud14-Glc7 work in different branches of SPOC

activation.

Glc7-Bud14 promotes Bfa1 dephosphorylation

Bfa1 phosphorylation is a key regulator of mitotic exit. Cdc5 phosphorylation of Bfa1 inhibits Bfa1-Bub2 GAP activity towards Tem1 to promote mitotic exit; whereas Kin4 phosphorylation of Bfa1 prevents Cdc5 phosphorylation of Bfa1 to block mitotic exit (Geymonat *et al.*, 2002; Hu *et al.*, 2001; Maekawa *et al.*, 2007). We asked whether Bud14 impinges on Bfa1 phosphorylation. To investigate Bfa1 phosphorylation, we made use of *lte1Δ* Gal1-UPL-*TEM1* cells which arrest in late anaphase upon depletion of Tem1 with Kin4 localized at both SPBs (Bertazzi *et al.*, 2011; Hu *et al.*, 2001). This arrest mimics a SPOC activated state where Cdc5 phosphorylation of Bfa1 is prevented by Kin4 phosphorylation of Bfa1 and consequently Bfa1 hyperphosphorylation is restored upon deletion of *KIN4* in these cells (Bertazzi *et al.*, 2011) (Figure 7- Figure Supplement 1). Markedly, deletion of *BUD14* also resulted in appearance of slow migrating Bfa1 forms in *lte1Δ* Gal1-UPL-*TEM1* cells (Figure 7A-B, Figure 7- Figure Supplement 1B) although Kin4 was localized at both SPBs (Figure 2 – Figure Supplement 1D). Such slow migrating forms of Bfa1 in *lte1Δ bud14Δ* cells were dependent on Cdc5 as they disappeared in *cdc5-10 lte1Δ bud14Δ* cells arrested in anaphase (Figure 7C). These data indicate that Bud14 prevents Cdc5-dependent hyperphosphorylation of Bfa1 in *lte1Δ* cells.

We next analyzed the effect of *BUD14* overexpression on Bfa1 phosphorylation. Bud14 overexpression causes cells to arrest in a pre-anaphase state and, hence, it is lethal (Chesarone *et al.*, 2009; Cullen & Sprague, 2002; Knaus *et al.*, 2005; Pinsky *et al*, 2006). Of importance, this lethality is not dependent on Bfa1 or Kin4 (Figure 7 – Figure Supplement 2A). To obtain cells arrested in anaphase in the presence of high levels of Bud14, we first arrested *cdc15-as* (L99G) (Bishop *et al*, 2000; D'Aquino *et al.*, 2005) bearing cells in anaphase through 1NM-PP1 treatment and then induced Gal1-*BUD14* overexpression by addition of Galactose into the growth medium. Overexpression of *BUD14* (Figure 7D) but not *bud14-F379A* caused a

reduction in hyperphosphorylated forms of Bfa1 (Figure 7E-F). This data suggests that Glc7-

Bud14 causes dephosphorylation of hyperphosphorylated Bfa1 in anaphase arrested cells.

To further understand whether Glc7-Bud14 dephosphorylates Bfa1, we performed an *in vitro*

dephosphorylation assay. For this, we enriched Bfa1-3HA from Gal1-UPL-*TEM1 kin4Δ* cells

arrested in telophase through depletion of Tem1 and incubated it with or without

immunoprecipitated Glc7-TAP in the presence or absence of 1.5 μM okadaic acid to inhibit

Glc7 (Bialojan & Takai, 1988; Cohen *et al*, 1989a; Cohen *et al*, 1989b) (Figure 7G). Addition

of Glc7-TAP in the phosphatase reaction caused a down shift of Bfa1-3HA, but not when

okadaic acid was added (Figure 7G). Same behavior was observed when Bud14-TAP was used

in the phosphatase reaction instead of Glc7-TAP (Figure 7 – Figure Supplement 2B). These

data altogether show that Glc7-Bud14 promotes dephosphorylation of hyperphosphorylated

Bfa1 in anaphase.

Discussion

SPOC is a surveillance mechanism that inhibits mitotic exit in response to spindle

mispositioning. Protein phosphorylation and localization play a key role in SPOC. Especially

the activity of Bfa1, the most downstream effector of the SPOC is tightly controlled with

respect to the spindle orientation through phosphorylation by the polo like kinase Cdc5 and the

mother specific kinase Kin4 in an opposing manner. Phosphorylation of Bfa1 in turn affects

activity and localization of Bfa1-Bub2 and hence the activity and the localization of the GTPase

Tem1 at SPBs. Despite the knowledge on Bfa1 phosphorylating kinases, knowledge on

phosphatases counteracting these kinases has been limited. In this study we established Glc7-

Bud14 as a novel SPOC component that promote dephosphorylation of Bfa1 when Bfa1 is

hyperphosphorylated in anaphase in a Cdc5-dependent manner. Our work supports a model

where Bfa1-Bub2 is activated by Glc7-Bud14 in parallel to Kin4. Upon spindle misalignment

in this model, the SPOC kinase Kin4 phosphorylates Bfa1 to remove Bfa1 from the SPBs,

away from the inhibitory kinase Cdc5, and Glc7-Bud14 dephosphorylates Bfa1 to remove its

inhibitory phosphorylation.

Glc7-Bud14 is a SPOC component

Our study identified Glc7, the budding yeast protein phosphatase 1, in association with its

regulatory subunit Bud14 as a novel SPOC component that inhibits mitotic exit upon spindle

mispositioning. Several lines of evidence led us to this conclusion. First, similar to other SPOC

proteins, *BUD14* deletion rescued growth of several mutants impaired in mitotic exit, including

the cold sensitive *lte1Δ* cells, the temperature sensitive MEN mutants (*cdc15-1*, *dbf2-2* and

mob1-67), and the synthetic lethal *lte1Δ spo12Δ* cells. Partial impairment of Glc7 function via

the *glc7-12* mutant also rescued growth lethality of *lte1Δ spo12Δ* cells. In addition, a Bud14

mutant that cannot bind Glc7 (*bud14-F379A*) (Knaus *et al.*, 2005) also rescued cold sensitivity

of *lte1Δ* cells. Second, similar to other SPOC proteins, *BUD14* deletion caused accumulation

of multinucleated cells when spindle misalignment was induced in cells by deletion of *KAR9*

or *DYN1*, implying that *bud14Δ* cells are deficient in arresting upon mispositioning of the

anaphase spindle. Of importance, similar to *bud14Δ*, *glc7-12* and the *bud14-F379A* mutants

were also SPOC deficient. Bud14 was implicated in the Dynein pathway of spindle positioning

(Knaus *et al.*, 2005). However, deletion of *BUD14* caused accumulation of multinucleated

phenotypes not only in *kar9Δ* but also in *dyn1Δ* backgrounds indicating that the function of

Bud14 in Dynein activation is not contributing to the observed multinucleation phenotype. In

line with this notion, time-lapse analysis revealed that *bud14Δ kar9Δ* cells exited mitosis

regardless of the spindle position. Furthermore, similar to other SPOC components, absence of

BUD14 did not affect the timing of mitotic exit when the spindle was correctly positioned but

was crucial for delaying exit from mitosis when the spindle is misoriented. Data supporting

this included the anaphase duration of *bud14Δ kar9Δ* cells with normally positioned spindles,

which was not different than that of *kar9Δ* cells and also Mob1-GFP localization in *bud14Δ*

cells, which was not different than in wild type cells.

Glc7-Bud14 works in a pathway parallel to Kin4

Our data showed that Bud14 does not work in the Kin4 branch of SPOC. First, Kin4

localization and phosphorylation did not depend on Bud14. Second, Kin4 overexpression still

caused toxicity in *BUD14* deleted cells indicating that Kin4, when produced at high levels, is

able to keep Bfa1 active. Furthermore, time-lapse analysis of Bfa1 localization during spindle

misalignment showed that Bfa1 is being removed from SPBs of *bud14Δ* cells with misaligned

spindles, while removal of Bfa1 from SPBs is largely a Kin4 dependent process (Caydasi &

Pereira, 2009; Monje-Casas & Amon, 2009). Thus, Kin4 is able to exert its effect on Bfa1 in

the absence of *BUD14*. These data altogether exclude the possibility that Bud14 works through

Kin4 branch of the SPOC and suggests a model in which Glc7-Bud14 prevents mitotic exit of

cells with misaligned spindles in a way parallel to Kin4. In support of this model, we observed

an additive decrease in anaphase duration of *kin4Δ bud14Δ* cells with misaligned spindles

compared to the cells bearing single gene deletions of *KIN4* or *BUD14*. Importantly, *kin4Δ*

bud14Δ cells completely failed to delay mitotic exit while *kin4Δ* or *bud14Δ* cells were able to

slightly delay mitotic exit in response to spindle misorientation (Figure 2C). Accordingly,

SPOC activity in *kin4Δ bud14Δ* cells was as disrupted as in *bfa1Δ* cells (Figure 2A). Thus, we

propose that Kin4 and Bud14 work in parallel and presence of both is critical for a robust SPOC

arrest.

Interestingly, neither *KIN4* nor *BUD14* is required for spindle assembly checkpoint dependent

metaphase arrest of cells, whereas Bfa1-Bub2 is (data not shown) (Pereira & Schiebel, 2005).

This suggest that the Kin4 and Bud14 become important to halt mitotic exit after APC/Cdc20

activation at the metaphase-anaphase transition. Anaphase onset also brings about the partial

activation of Cdc14 by the FEAR pathway which primes MEN activation in several ways and

Kin4 was shown to be dispensable for SPOC in cells lacking FEAR activity (Caydasi *et al.*,

2017; Falk *et al*, 2016a; Scarfone *et al.*, 2015). Consistently, Bud14 was also dispensable for

SPOC in *spo12Δ* cells that lack FEAR (data not shown). These data support that Kin4 and

Bud14 work in parallel to keep Bfa1-Bub2 active in anaphase, to provide an arrest of anaphase

cells prior to mitotic exit.

Glc7-Bud14 promotes dephosphorylation of Bfa1

The SPOC component Bfa1 is a highly phosphorylated protein in a cell cycle dependent

manner. So far, Bfa1 has been shown to be the target of kinases Cdk1, Cdc5 and Kin4 (Caydasi

et al., 2017; Geymonat *et al*, 2003; Hu *et al.*, 2001; Pereira & Schiebel, 2005). Among these

kinases Cdc5 plays an inhibitory role, whereas Cdk1 and Kin4 plays and activating role. It is

possible that other kinases may also phosphorylate Bfa1. There has been an ongoing curiosity

for phosphatases that act on Bfa1. So far, Cdc14 was shown to counteract Cdk1

phosphorylation of Bfa1 to partially inhibit Bfa1-Bub2 activity after anaphase onset (Caydasi

et al., 2017) and PP2A-Cdc55 was shown to counteract Cdc5 dependent phosphorylation of

Bfa1 before anaphase onset, and hence downregulation of PP2A-Cdc55 activity after anaphase

onset facilitates Bfa1 inhibition (Baro *et al*, 2013). Our data that Bfa1 appears

hyperphosphorylated in *lte1Δ bud14Δ* cells, in a Cdc5 dependent manner, during anaphase

indicate that Glc7-Bud14 promotes dephosphorylation of hyperphosphorylated Bfa1 after

anaphase onset. Consistently, overexpression of *BUD14* but not the *bud14-F379A*, which

cannot bind Glc7, promoted dephosphorylation of Bfa1 in anaphase arrested cells.

Furthermore, through an in vitro dephosphorylation assay we showed that Glc7-Bud14 is able

to dephosphorylate hyperphosphorylated Bfa1 purified from anaphase arrested cells.

We further showed that Bfa1 and Bud14 interacted in yeast two hybrid assays. Curiously, Bfa1-

Bud14 interaction observed in the yeast two hybrid assay required presence of Bub2. Thus,

Bud14 may recognize Bfa1-Bub2 as a complex, or the interaction between Bud14 and Bfa1

may require *in vivo* modifications of Bfa1 that occur only when bound to Bub2. It has been

known that hyperphosphorylated forms of Bfa1 abolishes in the absence of Bub2 (Lee *et al*,

2001b). In addition, Bfa1 fails to localize to SPBs in the absence of Bub2 (Pereira *et al.*, 2000).

Furthermore, Bfa1 phosphorylation by Cdc5 takes place at SPBs (Maekawa *et al.*, 2007;

Pereira *et al.*, 2000).

These findings, altogether, are suggestive of a model in which Glc7-Bud14 may recognize

specifically a phosphorylated form of Bfa1. We favor a speculative model where Glc7-Bud14

dephosphorylates Cdc5-phosphorylated Bfa1 specifically upon spindle misalignment. This

model is also in concordance with our previous conclusion that Glc7-Bud14 works in a parallel

pathway to Kin4. Removal of Cdc5 dependent phosphorylation of Bfa1 by Glc7-Bud14 and

isolation of Bfa1 away from the inhibitory Cdc5 kinase activity by Kin4 would concertedly

reduce the levels of Cdc5 phosphorylated Bfa1 and thus work together to rapidly increase the

levels of active Bfa1-Bub2.

Kin4 phosphorylation of Bfa1 prevents Cdc5 phosphorylation of Bfa1 both during SPOC

activation (i.e. anaphase arrest of *lte1Δ* cells that mimics SPOC arrest) and during the

metaphase arrest upon nocodazole treatment (Bertazzi *et al.*, 2011; D'Aquino *et al.*, 2005;

Pereira & Schiebel, 2005). However, unlike *kin4Δ* cells, *bud14Δ* cells arrested in metaphase

through nocodazole treatment did not have hyperphosphorylated Bfa1 (data not shown). This

discrepancy between Kin4 and Glc7-Bud14 may indicate alternative scenarios: First, Glc7-

Bud14 may dephosphorylate sites other than those phosphorylated by Cdc5. Second, there may

be redundant mechanisms of Bfa1 dephosphorylation at Cdc5 phosphorylation sites that

especially dominate in metaphase. PP2A-Cdc55 was shown to counteract Bfa1

phosphorylation before anaphase onset (Baro *et al.*, 2013). Therefore, one possibility is that

PP2A-Cdc55 and Glc7-Bud14 may work redundantly to dephosphorylate Cdc5 phosphorylated

Bfa1 at different stages of the cell cycle, metaphase and anaphase respectively.

In addition, a truncated version of Bud14 that lacks the SH3 domain was also unable to inhibit

mitotic exit. This truncated Bud14 is unable to localize at the bud cortex is also unable to bind

Glc7 (Knaus et al., 2005) (Figure 3-Figure Supplement 1B). Our yeast two hybrid analysis also

revealed that Bud14-SH3Δ also have a reduced binding to Bfa1 (Figure 5C). Thus, the loss in

SPOC activity of the Bud14-SH3Δ can be attributed to its reduced association with Glc7 and

Bfa1.

Intriguingly, another version of Bud14, which lacks the formin regulatory motif (Bud14-5A),

was able to yield a stronger interaction with Glc7 in our yeast two hybrid assays (Figure 3-

Figure Supplement 1B). This increased association may explain why Bud14-5A provided a

stronger SPOC activity than the Bud14 wildtype based on SPOC functionality measurements

(Figure 3A).

Bud14 deletion causes an increase in SPB-bound levels of Bfa1-Bub2 and Tem1

We observed that cells lacking *BUD14* had more Bfa1, Bub2, and Tem1 localized at dSPB,

making their localization more asymmetric. The asymmetry of Bfa1 at SPBs was shown to be

dependent on Cdc5, as in the absence of Cdc5 activity Bfa1 localized more symmetrically to

SPBs (Kim *et al*, 2012). Thus, presence of more Cdc5 phosphorylated Bfa1 in *bud14Δ* cells

may account for increased levels of Bfa1 at the dSPB, and also Bub2 and Tem1 which bind

SPBs in association with Bfa1. Alternatively, Glc7-Bud14 may have other targets at the SPB

that might normally decrease Bfa1-Bub2 docking sites at the SPBs.

Studies that used versions of Bfa1 forced to bind SPBs stably (Scarfone *et al.*, 2015; Valerio-

Santiago & Monje-Casas, 2011), suggested that increased Bfa1-Bub2 localization at SPBs can

lead to increased MEN activity, as it also causes increased Tem1 association with SPBs. Does

altered localization of Bfa1-Bub2 and Tem1 observed in *bud14Δ* contribute to mitotic exit?

Analysis of Mob1 localization as a marker for MEN activation suggests that *bud14Δ* cells does

not activate MEN prematurely. This suggests that Bud14-Glc7 may be inhibiting mitotic exit

by activation of Bfa1-Bub2 GAP complex through counteracting an inactivating

phosphorylation of Bfa1 rather than through modulating SPB-bound levels of Bfa1-Bub2. We

favor that the change in SPB localization is a consequence of the change in the phosphorylation

status of Bfa1 or another target at the SPB.

Role of PP1 through the cell cycle

PP1 has numerous functions in the control of the cell cycle (Holder *et al*, 2019; Kim *et al*,

2016; Moura & Conde, 2019; Nasa & Kettenbach, 2018). PP1, together with PP2A, is required

for mitotic exit in animal cells by counteracting CDK activity (Wu *et al*, 2009). In human cells,

PP1 also regulates centrosome splitting by antagonizing the Nek2 kinase (Meraldi & Nigg,

2001). In addition, PP1 facilitates abscission by counteracting Plk1 mediated phosphorylation

of the centrosomal protein CEP55 during late mitosis and thus promoting its midbody

localization (Gao *et al*, 2018). In meiosis of Xenopus oocytes, PP1 regulates G2/M transition

by partially activating Cdc25 (Perdiguero & Nebreda, 2004). From yeast to mammals, PP1 is

essential for silencing of the spindle assembly checkpoint through stabilization of kinetochore

microtubule interactions by counteracting Aurora B and by potentially regulating recruitment

of MCC to kinetochores (Kelly & Funabiki, 2009; Kotwaliwale & Biggins, 2006; Musacchio

& Salmon, 2007; Ruchaud *et al*, 2007). In this study, we show that the budding yeast PP1,

Glc7, in association with its regulatory subunit Bud14, inhibits mitotic exit upon anaphase

spindle mispositioning. This function of PP1 seems to be opposite to the mitotic exit promoting

role of PP1 reported in animal cells. The fact that mitotic exit is achieved by the Cdc14

phosphatase in budding yeast, instead of the PP1 and PP2 phosphatases (Manzano-Lopez &

Monje-Casas, 2020) is likely the reason for Glc7 gaining a different function after the anaphase

onset during the narrow time window until exit from mitosis.

Glc7-Bud14 was implicated in microtubule cortex interactions. Data based on Kin4 activity on

Bfa1 indicates that SPOC mechanism may sense microtubule-cortex interactions (Caydasi &

Pereira, 2009; Maekawa *et al.*, 2007; Monje-Casas & Amon, 2009; Pereira & Schiebel, 2005).

How spindle alignment is sensed, and this information activates SPOC still remain unknown.

Due to their known function in regulating microtubule-cortex interactions (Knaus *et al.*, 2005),

Glc7-Bud14 constitutes a great candidate for sensing of such interactions. Considering that

Glc7 (PP1) is a conserved phosphatase, similar mechanisms may be employed by higher

eukaryotes. Future work will show whether Glc7-Bud14 may act as a sensor that translates the

unattached microtubule-cortex interactions to the downstream SPOC machinery.

Materials Methods

Yeast methods, strains and growth

All yeast strains used are isogenic with S288C and are listed in Table 1. Basic yeast methods

and growth media were as described (Sherman, 1991). Carbon source is glucose unless

otherwise stated. When necessary 2% D-Raffinose, 3% D(+)-Galactose was used instead of

glucose. Yeast strains were grown at 30 °C and in rich medium unless otherwise stated. Plasmid

containing strains were grown in synthetic complete (SC) media lacking the appropriate

auxotrophic nutrients. The temperature sensitive mutants and *kar9Δ* cells were maintained at

23 °C. Cassette PCR-based gene editing method was used for chromosomal gene deletion and

C-terminal or N-terminal tagging (Janke *et al*, 2004; Knop *et al*, 1999). To determine genetic

interactions based on growth, mutant yeast strains bearing the wild type gene on a *URA3*-based

plasmid (pRS316) were spotted on 5-Fluoroorotic acid (5FOA) plates. 5FOA plates negatively

selects for the *URA3*-based plasmid and allows for observation of the growth phenotype in the

absence of the corresponding plasmid. Most *kar9Δ* cells were maintained complemented with

KAR9 on *URA3-based* centromeric plasmid (pGW399), and stroke on 5FOA plates to induce

plasmid lost shortly before experiments on SPOC analysis. Similarly, MEN-ts mutants were

maintained complemented with plasmids carrying the wildtype version of the corresponding

MEN gene. The plasmids used in this study are listed in Table 2.

Cell growth comparison on agar plates

Yeast cultures were grown in appropriate media and growth conditions for 24 hours. The OD_{600}

of the cultures were adjusted to 1 and 10-fold serial dilutions were made using sterile PBS. 10

μl of serial dilutions were spotted on appropriate agar plates and grown at appropriate

temperatures for 1-3 days.

Cell cycle synchronizations

Cells were synchronized in G1 phase by treating log-phase cultures with 10 µg/ml α-factor (Sigma #T6901) dissolved in DMSO and incubating for one and a half doubling time. For synchronization in metaphase without spindles, 15 µg/ml nocodazole (Sigma #M1404) dissolved in DMSO were added to log-phase cultures and incubated for ~2h. The arrest was confirmed by microscopy after fixing the cells with 70% Ethanol and re-suspending in PBS containing 1 µg/ml 4',6-diamino-2-phenylindole (DAPI, Sigma). For nocodazole arrest from G1 synchronized yeast cultures, cells were first treated with α-factor and released in nocodazole containing YPAD. For telophase arrest of Gal1-UPL-Tem1 bearing cells, after cultures were grown to log-phase in Raffinose-Galactose containing medium, cells were transferred into YPAD medium and grown for ~3 h until cells (>%90) were arrested with large buds and separated nuclei. For telophase arrest from G1 arrested cells: Log-phase cultures of Gal1-UPL-Tem1 bearing cells grown in Raffinose/Galactose were treated with α-factor. 30 min before G1 synchronization was completed, glucose was added to the culture to give 2% final concentration. Then cells were released from α-factor in YPAD. The arrest was confirmed by microscopy after ethanol fixation of the cells and staining the nucleus with DAPI. As for anaphase arrest using *cdc15-as*, log phase cell cultures grown in YP-Raffinose medium, were treated with 1NM-PP1 (2.5 µM) and incubated for 3 h. Synchronization was judged based on microscopy after DAPI staining. Anaphase synchronization of *cdc5-10* cells were achieved by arresting the cells in G1 at 23 °C first and then releasing them in pre-warmed alpha factor free medium at 30 °C and incubating them at 30 °C for 1.5 h.

Microscopy Methods

All fluorescence microscopy methods were performed using Carl ZEISS Axio Observer 7 motorized inverted epifluorescence microscope equipped with Colibri 7 LED light source,

Axiocam 702 Monochrome camera, 100X and 63X Plan Apochromat immersion oil objective

lenses, Filter sets 95 (GFP/Cherry), 20 (Rhodamin), FITC and DAPI, and an XL incubation

and climate chamber. For Kin4-GFP, GFP-Tubulin and Bud14-GFP analysis 63X objective

was used. For analysis of Bfa1-GFP, Bub2-GFP, Tem1-GFP and Mob1-GFP 100X objective

was used.

For end-point analysis of samples, wet mounts of samples were prepared by sandwiching 2 µl

of culture between the coverslip and slide. Cells were either fixed using 70% Ethanol and

stained with DAPI (for SPOC assays and mitotic index analysis) or analyzed live (For GFP

signal quantification). For time-lapse experiments, cells were prepared on glass bottom petri

dishes (WVR 10810-054 Matsunami) as described (Caydasi & Pereira, 2017). Briefly, the

glass center of the petri dish was covered with 6% Concanavalin A (*Canavalia ensiformis* Jack

Bean, type IV Sigma C2010-G). Excess Concanavalin A was washed out and 150 µl of culture

was pipetted on the dish followed by a 30 min incubation at 30 °C. Then, cells were aspirated

gently and cells that were not attached on the dish were washed out with pre-warmed fresh

media. After the final wash, the dish was filled with media and taken to the microscope stage

in the preheated chamber, equilibrated there for at least 1 hour before the start of the time-

lapse.

SPOC functionality assays

For end point analysis of SPOC functionality, log-phase *kar9Δ* cells cultured at 23°C were

incubated at 30°C for 3-5 h or log-phase *dyn1Δ* cells cultured at 30 °C were incubated at 18 °C

overnight. Cells were fixed using 70% Ethanol, re-suspended in PBS containing 1 µg/ml DAPI

and analyzed by microscopy. Cells bearing normal and misaligned nuclei, and cells with

SPOC-deficient phenotypes including multinucleated cells were counted. SPOC deficiency

index was calculated as:

$$= \% \textit{ Cells with multinucleation } \div \% \textit{ cells with mispositioned nuclei } \times 10$$

For time lapse analysis of SPOC functionality, GFP-Tubulin bearing *kar9Δ* cells were imaged every minute for 90 min at 30 °C. 10 z-stacks of 0.4 μm thickness were taken at each time point. Anaphase duration of cells with correct and misaligned spindles was calculated as the time from the start of fast spindle elongation (metaphase-to-anaphase transition) until spindle breakdown (Caydasi *et al.*, 2010b; Straight *et al*, 1997). Dot plots were prepared in GraphPad Prism 8.0.1 (GraphPad, La Jolla, CA).

Localization based fluorescence image quantifications

For signal intensity quantification of Bfa1-GFP, Bub2-GFP, Tem1-GFP and Mob1-GFP, we took still images of live cells bearing *SPC42*-eqFP and mCherry-*TUB1* as a mitotic marker. Each still image consisted of 13 z-stacks of 0.3 μm thickness. Z-stacks of the images were sum-projected, and mean fluorescence intensities were measured using Image J (NIH) software and corrected as described (Caydasi *et al.*, 2012). Briefly, a 0.577 μm^2 (42 pixels) box was selected around the spindle pole body under inspection (ROI, region of interest) and the mean fluorescence intensity of the GFP signal was measured. Background fluorescence intensity was determined by measuring an area inside the cytoplasm of respective cell, near to the region of interest. The correction was done by subtracting the mean fluorescence intensity of the background from the mean fluorescence intensity of the ROI.

To detect the behavior of asymmetric localization of Bfa1 at SPBs, ratio of the corrected signal intensities (as described above) at the SPBs were calculated. For cells with normally aligned spindles, the signal intensity at the dSPB was divided to the signal intensity at the mSPB. For cells with misaligned spindles, the SPB with the greater signal was considered as dSPB.

Statistical results and graphs were computed and plotted by using Prism 8.0.1 (GraphPad, La

Jolla, CA). Two-tailed t-test were applied to compare the statistical significance of the signal

intensity measurements in a pair-wise manner. Chi-square test were applied for comparison of

Bud14-GFP cortex localization.

Yeast Two Hybrid Assay

Yeast Two Hybrid analysis was performed as described (Meitinger *et al*, 2014). Briefly,

indicated genes cloned into pMM5 or pEG202 (LexA-myc fusion, *HIS* autotrophy marker) and

pMM6 (Gal4-HA fusion, *LEU2* autotrophy marker) were transformed into the yeast strain

SGY37-VIII,3 that contains a chromosomally integrated LexA(op)-LacZ reporter for

activation of the ß-galactosidase gene. Transformants were selected on SC-His-Leu agar plates,

grown in SC-His-Leu media until stationary phase, diluted to 1 OD_{600} and 10 µl of each were

spotted on SC-His-Leu agar plates. Cells were grown for~ 48 hours and plates were overlaid

with 0.4% soft-agar solution containing 0.4 mg/ml X-Gal, 0.1% SDS, 10 mM KCl and 1 mM

$MgCl_2$. Blue color formation was monitored. Plates were scanned.

Glc7 Functionality Assay

For the analysis of *GLC7* functionality, *GLC7* and *glc7-12* cells were grown at 23 °C until log-

phase. Log-phase cultures were diluted to a OD600 value of 0.1 and shifted to 30 °C, 33 °C,

35 °C and 37 °C. After incubation at these temperature for 4 h cells were fixed using 70%

Ethanol, and then resuspended in with 1 µg/ml DAPI in 1X PBS prior to microscopic

visualization. Then, the cells were analyzed under fluorescence microscopy. All the cells were

counted and categorized according to their bud size and nucleus number. For analyzing the

functionality of *GLC7* by growth assay, the cultures were grown at 23 °C for 24-48 hours.

OD600 value of all the yeast cultures, then, was adjusted to 1.0 using 1X sterile PBS. The

cultures were serially diluted 10-fold using sterile 1X PBS. 7.5 µl of each serial dilutions were

spotted on the appropriate agar plates (YPD and YPD containing 0.01M HU). The plates were

incubated at the appropriate temperatures for 2-3 days.

Protein Methods

Total yeast protein precipitation, sample preparation, and Western blotting following Laemmli

SDS-PAGE were performed as previously described (Meitinger *et al*, 2016). Total cell native

protein extracts were prepared by lysis of cells using acid washed glass beads (Sigma) in a

cooling bead beater homogenizer (Analytik Jena, SpeedMill Plus). Primary antibodies were

Mouse-anti-HA (Gift from Gislene Pereira), Rabbit anti-Clb2 (Gift from Gislene Pereira),

Rabbit-anti-Tubulin (Abcam EPR13799), Rabbit anti-TAP (Thermofisher scientific

CAB1001) and Rabbit anti-GFP (Abcam ab290). Secondary antibodies were Goat-anti-rabbit

(Advansta #R-05062-500) or Goat-anti-mouse (Advansta #R-05071-500) horse radish

peroxidase (HRP) conjugated secondary antibodies.

Phosphatase Assay

Glc7-TAP and Bud14-TAP were enriched from yeast lysates on IgG Sepharose beads (GE

Healthcare BioScience) according to manufacturer's recommendations. In addition, ESM356-

1 (a strain without TAP tag) lysates were incubated with IgG Sepharose beads as a no-tag

control. Briefly, cell pellets of logarithmically growing cultures were lysed using acid washed

glass beads (Sigma) in a lysis buffer containing 50mM Hepes pH:7.5, 150mM NaCl, 1mM

EDTA , %1 SDS, 1mM DTT, 2mM PMSF and complete Protease Inhibitor Coctail (Roche).

The lysate was cleared by centrifugation at 10000 rpm at + 4 $^{\circ}$C and the supernatant was

incubated with IgG beads for 2 h at 4 $^{\circ}$C. Beads were washed with ice cold wash buffer

containing 50mM Hepes pH: 7.5, 150mM NaCl, 0.1mM EDTA, %0.1 SDS, %0.025 Tween

20, 1mM DTT and were immediately used for the phosphatase reaction as the phosphatase

source.

Bfa1-3HA was purified from *BFA1-3HA kin4Δ Gal1 -UPL-TEM1* cells arrested in anaphase.

For this, log-phase culture grown in Raffinose/Galactose containing rich medium were

transferred into YPAD medium. After >90% of the cell arrest was achieved, cell pellets were

lysed using acid washed glass beads in bead beater. Lysis buffer composition was: 50mM

Hepes pH:7.5, 150mM NaCl, 5% Glycerol, 1mM EDTA, 1% NP-40, 160mM β-

Glycerophosphate, 2mM $NaVO_3$, 100mM NaF, 2mM PMSF, 4mM Benzamidine and complete

Protease Inhibitor Coctail (Roche). After clearing the cell lysate by centrifugation at 10000

rpm at + 4 °C, the cell extract was incubated with anti-HA magnetic beads (Thermofisher,

Pierce™) for 2 h at 4 °C. After the incubation the beads were washed in 50mM Hepes pH: 7.5,

150mM NaCl, %1 NP-40, %5 Glycerol, 1mM EDTA and eluted from beads under basic elution

conditions according to manufacturer's protocol. Briefly, beads were incubated with 50mM

NaOH for 5 minutes and the eluate was immediately neutralized by 300mM Tris pH:8.5. Next,

the eluate was applied on PD MiniTrap G-25 Sephadex (GE Healthcare) columns equilibrated

with a buffer containing 50mM Hepes pH: 7.5, 100mM NaCl. Buffer exchange was performed

according to manufacturer's spin protocol.

In vitro phosphatase reaction was performed using the IgG-beads that contained Bud14-TAP

or Glc7-TAP or the IgG-beads that was incubated with the cell extract of the ESM356-1

control. Bfa1-3HA that was prepared as explained above was added as the substrate.

Phosphatase reaction buffer composition was 100mM NaCl, 50mM Hepes pH: 7.5, 2mM DTT,

%0.025 Tween, 2mM $MnCl_2$. Okadaic acid (Abcam) was also added to the indicated reactions

to 1.5 μM final concentration. After reaction was incubated at 30 °C for 1h, 5x sample buffer

was added immediately into the reaction tubes, samples were boiled at 95 °C for 5 min and

analyzed by western blotting.

Quantification of hyperphosphorylation of Bfa1

Bfa1-3HA containing protein samples were run 8% Laemmli SDS-PAGE gels, for about 3-3.5

hours. The gel tank was placed in an ice-filled bucked during running. Gels were run at 50V

for about 30 min (to allow passage of proteins from stacking to separating gel), and at 100V

for the remaining time. Semi-dry blotting was performed for 2h at 0.1 A per gel and signals

were detected on x-ray films using ECL. X-ray films were scanned, and the images were

analyzed in Image J. Mean grey value of the top half and bottom half of the Bfa1-3HA bands

at each line was quantified using Image J measure tool. The area sizes were kept constant for

each blot. Background mean intensity of the film was also quantified and subtracted from that

of the measured Bfa1-3HA band intensities for correction. Corrected intensity of the top half

was divided to the corrected intensity of the bottom half of the Bfa1-3HA bands and the values

were plotted.

Acknowledgements

This work was supported by TÜBITAK Grant No. 117C041. Work of AKC was supported by EMBO installation Grant No. 3918 and ICGEB installation Grant No. CRP/TUR17-04_EC. CD was funded by Koç University, HK and IK were funded by TÜBITAK Grant No. 118Z979 and SNE was funded by TÜBITAK Grant No. 117Z232 granted to AKC. AKC was funded by MSCA Individual Fellowship No. 796599-COHEMEX. We are grateful to Gislene Pereira (COS, Heidelberg), Elmar Schiebel (ZMBH, Heidelberg), Claudio De Virgilio (University of Fribourg, Fribourg), Matthias Peter (ETH, Zürich) and Michael Knop (ZMBH, Heidelberg) for sharing reagents, strains and plasmids.

Author Contributions

HK, DK, CD, IK and SNE designed and performed the experiments. AKC acquired funding, designed experiments, supervised HK, DK, CD, IK, SNE, and wrote the manuscript.

Conflict of Interest

The authors declare that they have no conflict of interest.

 Figure Legends

Figure 1. Bud14 deletion rescues growth of cells with impaired mitotic exit. A. *bud14Δ*

cells rescue the synthetic lethality of *lte1Δ spo12Δ* cells. **B.** *bud14Δ* cells rescue the lethality

of *lte1Δ* cells at 18 °C. **C.** Comparison of growth rescue of MEN-ts mutants upon deletion of

BUD14 and *BFA1*. Serial dilutions of indicated strains were spotted on indicated plates and

grown at given temperatures. 5-FOA plates negatively selects for the *URA3*-based plasmids

(pRS316 containing the *LTE1* in A and B, pRS316 containing the wild type gene copy of the

corresponding MEN mutant in C. Thus, only cells that have lost these plasmids can grow on

5-FOA plates where genetic interactions can be observed.

Figure 2. *bud14Δ* cells are SPOC deficient. A-B. End-point analysis of SPOC integrity of

indicated yeast strains in *kar9Δ* (A) and *dyn1Δ* (B) background. Cells were fixed with ethanol

and stained with DAPI. Cells with normally aligned nuclei, mispositioned nuclei and multi

nucleated cells that failed to correctly position their spindle before mitotic exit were counted

by microscopy and their SPOC deficiency index percentages were plotted, where SPOC

deficiency index = % cells with multinucleation ÷ % cells with mispositioned nuclei x 10.

Graphs are average of three independent experiments. A minimum of 100 cells were counted

from each strain in each experiment. Error bars show standard deviation. Two-tailed Student's

t-test was applied. ** $p < 0.01$, *** $p < 0.001$. **C.** Single cell analysis of SPOC integrity in

indicated strains. Duration of anaphase in cells with misaligned and normally aligned spindles

were calculated as explained in the text and plotted as dot-plots. Data points in red indicate the

cells in which the spindle did not break down during the time lapse movie. In this case, plotted

values are the time duration during which these cells were observed in anaphase through the

time lapse movie. Consequently, the actual anaphase duration is greater than the value plotted

in red. This fact is emphasized in the red part of the y-axis, indicated above the red dashed line,

with addition of the ">" symbol before the y-axis values. Green dash line indicates the mean

value of the anaphase duration in cells with normally aligned spindles. One-way Anova with

uncorrected Fisher's LSD was applied for statistical analysis. *** $p < 0.001$, **** $p < 0.0001$

n: sample size. All pairwise comparisons and descriptive statistics are shown in the

corresponding source data file, whereas only comparisons of normal and misaligned spindles

are shown in the figure.

Figure 3. Function of Bud14 in actin regulation is dispensable for SPOC. A. End-point

analysis of SPOC deficiency index in *bud14Δ kar9Δ* cells carrying *URA3-based* empty plasmid

(empty) or *BUD14*-containing *URA3*-based plasmids (*BUD14, bud14-5A* and *bud14-ΔSH3*).

Graphs are average of three independent experiments. A minimum of 100 cells were counted

from each strain in each experiment. Error bars show standard deviation. *p<0.05 according to

Student's t-test. **B.** Serial dilutions of indicated strains bearing *URA3-based* empty plasmid

(not indicated on figure) or *BUD14*-containing *URA3*-based plasmids (*BUD14, bud14-5A* and

bud14-ΔSH3) were spotted on SC-URA plate and grown at indicated temperatures. **C.** End-

point analysis of SPOC deficiency index in indicated cell types. Graphs are average of three

independent experiments. A minimum of 100 cells were counted from each strain in each

experiment. Error bars show standard deviation. ***p<0.001 according to Student's t-test. **D-**

E. Serial dilutions of indicated strains bearing *LTE1* on *URA3*-based pRS316 plasmid were

spotted on 5-FOA and SC plates and grown at indicated temperatures. 5-FOA negatively

selects for *URA3* containing plasmids, thus cells lose their pRS316-*LTE1* plasmids on 5-FOA

plates and genetic interactions can be observed on this plate.

Figure 4. Glc7-Bud14 interaction is required for SPOC regulatory function of Bud14. A.

Serial dilutions of indicated strains bearing *URA3*-based empty plasmid (not indicated on

figure) or *BUD14*-containing *URA3*-based plasmids (*BUD14* and *bud14-F379A*) were spotted

on SC-URA plate and grown at indicated temperatures. **B.** SPOC deficiency indexes of

indicated strains carrying *URA3*-based empty plasmid (empty) or *BUD14*-containing *URA3*-

based plasmids (*BUD14* and *bud14-F379A*). Graphs are average of three independent

experiments. A minimum of 100 cells were counted from each strain in each experiment. Error

bars show standard deviation. * p<0.05 according to student's t-test. **C.** Serial dilutions of

indicated strains were spotted on 5-FOA plate and grown at indicated temperatures. Cells that

contain *LTE1* on a *URA3* based plasmid are indicated with *pRS316-LTE1*. **D.** SPOC deficiency

indexes of indicated strains. *p< 0.05 according to student's t-test.

Figure 5. Yeast two hybrid analysis of Bud14 and Bud14 mutants with SPOC proteins.

A. Bud14 interacts with Bfa1 but not with Bub2 or Kin4. **B.** Bud14-Bfa1 interaction is

dependent on Bub2. **C.** Bfa1-Bud14 interaction is reduced in Bud14ΔSH3 mutant. SGY37 was

co-transformed with indicated plasmids. Empty plasmids served as a control for any self-

activation. Kel1 served as a positive control. Cells were grown for two days on selective agar

plates before overlay. Blue color formation was monitored as an indication of protein-protein

interaction.

Figure 6. More Bfa1 localizes to dSPB in the absence of *BUD14*. A. Bfa1-GFP signal

intensity quantifications at the SPBs of *BFA1-GFP mCherry-TUB1 or BFA1-GFP mCherry-*

TUB1 bud14Δ cells with normal aligned anaphase spindles were plotted on the right. Black

and grey lines in the dot plots are the mean value and standard deviation respectively.

Representative still images of cells were shown on the left. **B.** Bfa1-GFP signal intensities at

the dSPB were plotted in *BUD14* and *bud14Δ* cells bearing empty plasmid (blue and red), as

well as *bud14Δ* cells bearing a ADH-*BUD14* containing plasmid (green). **C.** Bfa1-GFP signal

intensities at the dSPB were plotted in *bud14Δ cells* bearing empty plasmid (red), ADH1-

BUD14 containing plasmid (green) and ADH1-*bud14-F379A* containing plasmid (yellow) **D.**

Analysis of Bfa1-GFP asymmetry at the SPBs of *kar9Δ, kar9Δ kin4Δ* and *kar9Δ bud14Δ* cells

with correctly aligned and misaligned anaphase spindles. Box and Whisker plot shows the ratio

of Bfa1-GFP signal intensities at the SPB1 and SPB2, where SPB1 always corresponds to SPB

with the greater Bfa1-GFP signal. The box represents first and third quartile while the whiskers

show 10-90 percentile of the data. The horizontal line in the box indicates the median of the

data. Only comparisons of normal and misaligned spindles are shown in the figure.

Representative still images of cells are shown on the right. n: sample size. Scale bar: 2 μm.

One-way Anova with uncorrected Fisher's LSD was applied for all statistical analysis. **

$p<0.01$, *** $p < 0.001$, **** $p < 0.0001$. All pairwise comparisons are shown in the

corresponding source data file.

Figure 7. Glc7-Bud14 promotes Bfa1 dephosphorylation in anaphase. A. Bfa1-3HA Gal1-

UPL-Tem1 containing *lte1Δ* and *lte1Δbud14Δ* cells grown in galactose containing medium

were released from alpha factor imposed G1 arrest (t=0) into an alpha factor free medium

supplemented with glucose to achieve Tem1 depletion and samples were collected at indicated

time points. Bfa1-3HA mobility shift was analyzed via western blotting using anti-HA

antibodies. Red arrow indicates hyperphosphorylated forms of Bfa1-3HA whereas blue arrow

indicates hypophosphorylated forms of Bfa1-3HA. Percentage of cells with single nucleus (1

DAPI) and two separate nuclei (2 DAPI) were plotted as a marker for cell cycle progression.

B. Indicated time points of each cell type from the experiment shown in A were loaded side-

by-side for better comparison of Bfa1-3HA mobility. **C.** Bfa1-3HA mobility in Gal1-UPL-

Tem1 or *cdc5-10* bearing *LTE1 BUD14* or *lte1Δbud14Δ* cells. Percentage of cells with two

separate nuclei (% 2DAPI) were indicated as a measure of cells in anaphase. Cells were first

arrested in G1, then released from the G1 arrest and cultured for 90 min before sample

collection. Gal1-UPL-Tem1 cells were treated as in A to achieve the anaphase arrest, whereas

anaphase arrest of *cdc5-10* cells was achieved through growth at 37 °C. **D.** *cdc15-as* bearing

868 cells bearing *BUD14* or Gal1-*BUD14* at the Bud14 endogenous locus were grown to log-phase

869 in raffinose containing medium, treated with 1NM-PP1 for 2,5h followed by galactose addition

870 (t0) to the medium. Samples were collected at 0h, 2h and 3h after galactose addition and Bfa1

871 mobility was analyzed. Percentage of cells with two separate nuclei (% 2DAPI) were indicated

872 as a measure of cells in anaphase. **E.** *cdc15-as bud14Δ* cells with *BUD14*-9myc, Gal1-*BUD14*-

873 9myc, or Gal1-*bud14-F379A*-9myc integrated at the chromosomal *leu2* locus were grown to

874 log-phase in raffinose containing medium, treated with 1NM-PP1 for 3h, followed by galactose

875 addition (t0) to the medium. Samples were collected at 0h, 1h, 2h and 3h after galactose

876 addition and Bfa1 mobility was analyzed. Percentage of cells with two separate nuclei (%

877 2DAPI) were indicated as a measure of cells in anaphase. **F**. Quantification of relative levels

878 of hypersphosphorylated Bfa1 from the experiment shown in E. Band intensity ratio of slow

879 migrating forms to fast migrating forms of Bfa1 was plotted. **G.** In vitro phosphatase assay of

880 immunoprecipitated Glc7-TAP on IgG beads were incubated with Bfa1-3HA purified from

881 *BFA1-3HA* Gal1-UPL-TEM1 *kin4Δ* cells in the presence or absence of 1.5 μM Okadaic acid.

882 As a no Glc7-TAP control, IgG beads incubated with cell lysates of ESM356-1 were used.

883 Glc7-TAP levels were detected using anti-TAP antibodies. Bfa1-3HA was detected using anti-

884 HA antibodies. Red and blue arrows indicate slow migrating and fast migrating forms of Bfa1-

885 3HA respectively. Quantification of relative levels of hypersphosphorylated Bfa1 is shown on

886 the right. Each color represents a different independent experiment. One-way Anova with

887 uncorrected Fisher's LSD was applied for statistical analysis. ** $p<0.01$, **** $p < 0.0001$.

 Supplementary Figure Legends

Figure 2 – Figure Supplement 1. Bud14 does not influence Kin4 function. A. Serial

dilutions of indicated strains were spotted on Glucose containing (YPD) and Galactose

containing (YP Raf/Gal) agar plates. Gal1-Kin4 overexpression is induced on Galactose

containing plates. **B.** Logarithmic growing cultures bearing Kin4-GFP and Spc42-eqFP were

treated with nocodazole for 2 hours and Kin4 SPB and cortex localization was analyzed by

microscopy. *rts1Δ* cells were used as a control in which Kin4 fails to localize to SPBs and

cortex. Graph is an average of three independent experiments. Error bars are standard

deviation. At least 100 cells were counted from each sample in each experiment. Scale bar: 2

μm. **C.** SPB localization of Kin4-GFP was analyzed in logarithmic growing *lte1Δ* and *lte1Δ*

bud14Δ cultures. Spc42-eqFP and mCherry-Tub1 served as SPB and spindle markers. Only

cells in anaphase were considered. Graphs were plotted as described in (B). **D.** Logarithmic

growing cultures bearing Kin4-6HA and the indicated gene deletions were treated with

nocodazole for 2.5 h. Kin4 mobility shift was analyzed by western blotting using anti-HA

antibodies. *rts1Δ* served as a control for hyperphosphorylated Kin4.

Figure 3 - Figure Supplement 1. Expression and Glc7 binding of Bud14 mutants. A.

Analysis of *BUD14* expression in cells bearing pMK60 (*pRS416-endogenous-BUD14-*

promoter-GFP-BUD14), pDKY001 (*pRS416-endogenous-BUD14-promoter-GFP-*bud14⁵ᴬ),

pHK002 (*pRS416-endogenous-BUD14-promoter-GFP-bud14-F379A),* pDKY003 (*pRS416-*

endogenous-BUD14-promoter-GFP-bud14-ΔSH3) Bud14 was detected using anti-GFP

antibodies. **B.** Interaction of Glc7 with Bud14 and Bud14 mutants in yeast two hybrid system.

Figure 4 – Figure Supplement 1. Assessment of *glc7-12* inactivation at different

temperatures. A. Serial dilutions of indicated strains were spotted on YPD and YPD

containing 0.01M HU and grown at indicated temperatures. The plates then were incubated at

given temperatures for 2-3 days. **B.** Analysis of cell morphology of *GLC7* and *glc7-12* cells at indicated temperatures. Liquid cultures were grown to log-phase at 23 °C, shifted to indicated temperatures and further grown for 4 h. Cells were fixed with ethanol and stained with DAPI. Cells were counted by microscopy and percentages of cells with large buds and single DAPI were plotted. A minimum of 100 cells were counted from each strain.

Figure 5 - Figure Supplement 1. Analysis of Glc7 interaction with SPOC components. A. Yeast two hybrid assays of Glc7 with Bfa1, Bub2 and Kin4. Bud14 was included as a positive control.

Figure 6 - Figure Supplement 1. Bfa1, Bub2 and Tem1 localization in *bud14Δ* cells during anaphase and metaphase. A-B. Bub2-GFP and Tem1-GFP representative images (A) and signal intensity quantifications at the SPBs (B) of *BFA1-GFP mCherry-TUB1* or *BFA1-GFP mCherry-TUB1 bud14Δ* cells with normal aligned anaphase spindles. Black and grey lines in the dot plots are the mean value and standard deviation respectively. Representative still images of cells were shown in A. **C.** Bfa1-GFP, Bub2-GFP and Tem1-GFP signal intensities at the SPBs of cells with metaphase spindles (spindle length = 1,5-2 μm). Graph is plotted as described in B. **D.** Analysis of Bfa1-GFP signal intensities at the SPBs of *kar9Δ*, *kar9Δ kin4Δ* and *kar9Δ bud14Δ* cells with correctly aligned and misaligned anaphase spindles. The graph was plotted as in B. Pairwise comparisons were performed using One-way Anova with uncorrected Fisher's LSD *** $p < 0.001$, **** $p < 0.0001$, n: sample size. Scale bar: 2 μm.

Figure 6- Figure Supplement 2. Analysis of Mob1 SPB localization. A. Percentage of metaphase cells (spindle length = 1,5-2 μm) with Mob1-GFP signal at the SPBs of the indicated cells. Graphs are average of three independent experiments. A minimum of 100 cells were counted from each strain in each experiment. Error bars show standard deviation. **B.**

Representative still images showing Mob1-GFP localization in metaphase cells **C-D.** Dot plot

of Mob1-GFP signal intensity at the SPBs of indicated strains carrying *MOB1-GFP mCherry-*

TUB1 in metaphase (C) and anaphase (D). Black lines represent the mean value, and grey lines

show standard deviation. **E.** Representative still images of anaphase cells. Scale bar: 2 μm.

One-way Anova with uncorrected Fisher's LSD was used for pairwise comparisons. **** $p < $

0.0001.

Figure 6 – Figure Supplement 3. Bfa1-GFP localization during spindle misalignment.

Selected frames from time-lapse series of *BFA1-GFP SPC42-eqFP mCherry-TUB1* bearing

kar9Δ (A), *kar9Δ bud14Δ* (B) and *kar9Δ kin4Δ* (C) gene deletions. Cell boundaries are outlined

with a dashed line. Blue arrows indicate the SPBs. Time points from the beginning of the time

lapse is indicated on the frames. Scale bar: 2 μm.

Figure 7 – Figure Supplement 1. Analysis of Bfa1 mobility in *lte1Δ* and *lte1Δ kin4Δ* cells.

A. Bfa1-3HA Gal1-UPL-Tem1 containing *lte1Δ* and *lte1Δkin4Δ* cells grown in galactose

containing medium were released from G1 arrest (t=0) into an alpha factor free medium

supplemented with glucose. Samples were collected at indicated time points. Bfa1-3HA

mobility shift was analyzed via western blotting. **B.** Indicated time points of each cell type

from the experiment shown in A were loaded side-by-side for better comparison of Bfa1-3HA

mobility. **C.** Percentage of cells with single nucleus (1 DAPI) and two separate nuclei (2 DAPI)

as well as budding index of the experiment shown in A were plotted as a marker for cell cycle

progression.

Figure 7 – Figure Supplement 2. Kin4 overexpression toxicity in *bud14Δ* cells and

phosphatase assay using Bud14-TAP. A. Serial dilutions of indicated strains were spotted

on Glucose containing (YPD) and Galactose containing (YP Raf/Gal) agar plates. Gal1-

Bud14 overexpression is induced on Galactose containing plates. **B.** In vitro phosphatase

assay of immunoprecipitated Bud14-TAP. Bud14-TAP was pulled on IgG Sepharose beads

from the lysates of *BUD14-TAP GLC7-9myc* cells. Beads were incubated with Bfa1-3HA

purified from *BFA1-3HA Gal1-UPL-TEM1 kin4Δ* cells in the presence or absence of Okadaic

acid. Beads incubated with ESM356-1 cell lysates were used as a negative control. Bud14-

TAP levels were detected using anti-TAP, Glc7-9myc levels were detected using anti-myc,

and Bfa1-3HA was detected using anti-HA antibodies. Quantification of relative levels of

hypersphosphorylated Bfa1 is shown on the right. Each color represents a different

independent experiment. One-way Anova with uncorrected Fisher's LSD was applied for

statistical analysis. ** $p < 0.01$.

Tables

Table 1. Yeast strains used in this study.

Strain Name	Description	Reference
AKY038	*ESM356 Bfa1-GFP-kanMX6 Spc42-eqFP-hphNT1 mCherry-Tub1-URA3*	This study
AKY043	*ESM356 BFA1-GFP-kanMX6 Spc42-eqFP-hphNT1 kar9Δ::klTRP1 pRS316-KAR9*	(Caydasi & Pereira, 2009)
AKY1533	*ESM356 leu2::LEU2-pGal1-KIN4*	This study
AKY1574	*ESM356 leu2::LEU2-pGal1-KIN4 bub2Δ::hphNT1*	This study
AKY2526	*FAY145 lte1Δ::kanMX6 pRS316-LTE1 spo12Δ::natNT2*	(Caydasi *et al.*, 2017)
AKY260	*ESM356 kar9Δ::HIS3MX6 pRS316-KAR9*	(Caydasi *et al.*, 2017)
AKY2916	*FAY145 lte1Δ::kanMX6 pRS316-LTE1 spo12Δ::natNT2 bud14Δ::HIS3MX6*	This study
AKY2917	*YPH499 GFP-Tub1-URA3 dyn1Δ::klTRP1 bud14Δ::HIS3MX6*	This study
AKY2918	*YPH499 GFP-Tub1-URA3 dyn1Δ::klTRP1 kin4Δ::hphNT1 bud14Δ::HIS3MX6*	This study
AKY315	*ESM356 kar9Δ::HIS3MX6 pRS316-KAR9 bfa1Δ::klTRP1*	(Caydasi *et al.*, 2017)
AKY321	*ESM356 kar9Δ::HIS3MX6 pRS316-KAR9 kin4Δ::klTRP1*	(Caydasi *et al.*, 2017)
AKY346	*YPH499 kar9Δ::klTRP1 pRS316-KAR9 GFP-Tub1-ADE2*	(Caydasi *et al.*, 2010b)
AKY351	*YPH499 kar9Δ::klTRP1 pRS316-KAR9 kin4Δ::HIS3MX6 GFP-Tub1-ADE2*	(Caydasi *et al.*, 2017)
AKY4001	*ESM356 leu2::LEU2-pGal1-KIN4 bud14Δ::klTRP1*	This study

AKY4006	*ESM356 KIN4-GFP-HIS3MX6 Spc42-eqFP-natNT2 bud14Δ::klTRP1 lte1Δ::hphNT1 mCherry-Tub1-URA3*	This study
AKY4007	*ESM356 Bub2-GFP-kanMX6 Spc42-eqFP-hphNT1 mCherry-Tub1-URA3*	This study
AKY4008	*ESM356 Tem1-GFP-kanMX6 Spc42-eqFP-hphNT1 bud14Δ::HIS3MX6 mCherry-Tub1-URA3*	This study
AKY4009	*ESM356 Bub2-GFP-kanMX6 Spc42-eqFP-hphNT1 bud14Δ::klTRP1 mCherry-Tub1-URA3*	This study
AKY4011	*ESM356 Bfa1-GFP-kanMX6 Spc42-eqFP-hphNT1 bud14Δ::klTRP1 mCherry-Tub1-URA3*	This study
AKY4012	*ESM356 Bfa1-GFP-kanMX6 Spc42-eqFP-hphNT1 mCherry-Tub1-URA3*	This study
AKY4013	*ESM356 Tem1-GFP-kanMX6 Spc42-eqFP-hphNT1 mCherry-Tub1-URA3*	This study
AKY4016	*ESM356 KIN4-GFP-HIS3MX6 Spc42-eqFP-natNT2 lte1Δ::URA3 mCherry-Tub1-kITRP1*	This study
AKY4028	*FAY145 lte1Δ::kanMX6 bud14Δ::HIS3MX6*	This study
AKY4036	*FAY145 lte1Δ::kanMX6 bud14Δ::HIS3MX6 pRS416-endogenous BUD14-promoter- GFP-BUD145A*	This study
AKY4038	*FAY145 lte1Δ::kanMX6 bud14Δ::HIS3MX6 pRS416-endogenous BUD14-promoter- GFP-BUD14ΔSH3*	This study
AKY404	*ESM356 KIN4-GFP-HIS3MX6 Spc42-eqFP-natNT2 rts1Δ::klTRP1*	(Caydasi *et al.*, 2010b)
AKY4040	*ESM356 Spc42-eqFP-hphNT1 BUD14-GFP-his3MX6 mCherry-Tub1-URA3*	This study

AKY4068	*YPH499 kar9Δ::klTRP1 pRS316-KAR9 GFP-Tub1-ADE2 bud14Δ::HIS3MX6 kin4Δ::hphNT1*	This study
AKY4078	*PAY704 MATa ade2-1 his3-11,15 leu2-3,112 ura3-1 can1-100 ssd1-d2 glc7::LEU2 trp1-1::GLC7::TRP1 kar9Δ::hphNT1*	This study
AKY4079	*PAY701 MATa ade2-1 his3-11,15 leu2-3,112 ura3-1 can1-100 ssd1-d2 glc7::LEU2 trp1-1::glc7-12::TRP1 kar9Δ::hphNT1*	This study
AKY4087	*ESM356 cdc15::CDC15-as1(L99G)-URA3 Bfa1-3HA- kITRP1 pGal1-Bud14-natNT2*	This study
AKY4088	*ESM356 Mob1-GFP-kanMX6 Spc42-eqFP-hphNT1 kin4Δ::natNT2 bud14Δ::HIS3MX6*	This study
AKY4091	*YPH499 cdc5-10 Bfa1-3HA-kITRP1 lte1Δ::natNT2 bud14Δ::HIS3MX6*	This study
AKY4094	*YPH499 cdc14-2 pRS316-CDC14 bfa1Δ::klTRP3*	This study
AKY4095	*YPH499 dbf2-2 pRS316-DBF2 bfa1Δ::klTRP3*	This study
AKY4102	*ESM356 Mob1-GFP-kanMX6 Spc42-eqFP-hphNT1 kin4Δ::natNT2 bud14Δ::HIS3MX6 mCherry-Tub1-LEU2*	This study
AKY415	*ESM356 KIN4-6HA-hphNT1 rts1Δ::klTRP1*	(Caydasi *et al.*, 2010b)
BKY032	*ESM356 BFA1-GFP-kanMX6 Spc42-eqFP-hphNT1 kar9Δ::klTRP1 pRS316-KAR9 kin4Δ::HIS3MX6*	(Caydasi & Pereira, 2009)
DGY001	*ESM356 KIN4-6HA-hphNT1 bud14Δ::klTRP1*	This study
DGY004	*ESM356 KIN4-GFP-HIS3MX6 Spc42-eqFP-natNT2 bud14Δ::klTRP1*	This study
DKY056	*YPH499 bud14Δ::kITRP3*	This study
DKY057	*YPH499 cdc14-2 pRS316-CDC14 bud14Δ::klTRP3*	This study

DKY058	*YPH499 mob1-67 pRS316-MOB1 bud14Δ::klTRP3*	This study
DKY060	*YPH499 dbf2-2 pRS316-DBF2 bud14Δ::klTRP3*	This study
DKY061	*YPH499 cdc15-1 pRS316-CDC15 bud14Δ::klTRP3*	This study
DKY063	*ESM356 KIN4-GFP-HIS3MX6 Spc42-eqFP611-natNT2 bud14Δ::klTRP1*	This study
DKY069	*FAY145 lte1Δ::kanMX6 pRS316-LTE1 bud14Δ::HIS3MX6*	This study
DKY070	*PAY704 MATa ade2-1 his3-11,15 leu2-3,112 ura3-1 can1-100 ssd1-d2 glc7::LEU2 trp1-1::GLC7::TRP1*	This study
DKY071	*PAY701 MATa ade2-1 his3-11,15 leu2-3,112 ura3-1 can1-100 ssd1-d2 glc7::LEU2 trp1-1::glc7-12::TRP1*	This study
DKY074	*PAY701 glc7::LEU2 trp1-1::glc7-12::TRP1 lte1Δ::HIS3MX6*	This study
DKY075	*PAY704 glc7::LEU2 trp1-1::GLC7::TRP1 lte1Δ::HIS3MX6*	This study
DKY078	*PAY704 glc7::LEU2 trp1-1::GLC7::TRP1 lte1Δ::HIS3MX6 pRS316-LTE1*	This study
DKY079	*PAY701 glc7::LEU2 trp1-1::glc7-12::TRP1 lte1Δ::HIS3MX6 pRS316-LTE1*	This study
DKY080	*PAY701 glc7::LEU2 trp1-1::glc7-12::TRP1 lte1Δ::HIS3MX6 pRS316-LTE1 spo12Δ::hphNT1*	This study
DKY081	*ESM356 BFA1-GFP-kanMX6 Spc42-eqFP hphNT1 kar9Δ::klTRP1 pRS316-KAR9 bud14Δ::HIS3MX6*	This study
DKY101	*ESM356 GAL1-UPL-TEM1::kITRP1 Bfa1-3HA-hphNT1*	This study
DKY113	*SGY37-VIII,3 bub2Δ::hphNT1*	This study

DKY115	*ESM356 Bfa1-3HA-hphNT1 kin4Δ::natNT2 pWS103 (pGal1-UPL-Tem1-kITRP1)*	This study
DKY118	*ESM356 GAL1-UPL-TEM1::kITRP1 Bfa1-3HA-hphNT1 lte1Δ::natNT2*	This study
DKY123	*YPH499 cdc5-10 Bfa1-3HA-hphNT1*	This study
DKY125	*ESM356 GAL1-UPL-TEM1::kITRP1 Bfa1-3HA-hphNT1 kin4Δ::natNT2 bud14Δ::HIS3MX6*	This study
DKY126	*ESM356 GAL1-UPL-TEM1::kITRP1 Bfa1-3HA-hphNT1 lte1Δ::natNT2 bud14Δ::HIS3MX6*	This study
DKY131	*FAY145 lte1Δ::kanMX6 pRS316-LTE1 bud14Δ::HIS3MX6 GFP BUD14 CEN URA*	This study
DKY132	*ESM356 bud14Δ::klTRP GFP BUD14 CEN URA*	This study
DKY135	*FAY145 lte1Δ::kanMX6 bud14Δ::HIS3MX6 pRS416-ADH-BUD14*	This study
DKY137	*FAY145 lte1Δ::kanMX6 bud14Δ::HIS3MX6 pRS416-ADH-BUD14-F379A*	This study
DKY145	*ESM356 kar9Δ::HIS3MX6 bud14Δ::klTRP1 pRS316*	This study
DKY147	*ESM356 kar9Δ::HIS3MX6 bud14Δ::klTRP1 pMK125*	This study
DKY149	*ESM356 kar9Δ::HIS3MX6 bud14Δ::klTRP1 pMK131*	This study
DKY167	*ESM356 KIN4-GFP-HIS3MX6 Spc42-eqFP-natNT2 bud14Δ::klTRP1 lte1Δ::hphNT1*	This study
DKY179	*YPH499 cdc5-10 pRS316-CDC5 bud14Δ::klTRP3*	This study
DKY188	*ESM356 KIN4-GFP-HIS3MX6 Spc42-eqFP-natNT2 lte1Δ::URA3*	This study
ESM2156	*YPH499 GFP-Tub1-URA3 dyn1Δ::klTRP1 kin4Δ::hphNT1*	(Pereira & Schiebel, 2005)

ESM2282	*YPH499 cdc15-1 pRS316-CDC15*	(Caydasi *et al.*, 2017)
ESM2283	*YPH499 dbf2-2 pRS316-DBF2*	(Caydasi *et al.*, 2017)
ESM2285	*YPH499 mob1-67 pRS316-MOB1*	(Caydasi *et al.*, 2017)
ESM2326	*ESM356 KIN4-6HA-hphNT1*	(Caydasi *et al.*, 2010b)
ESM356	*MATa ura3-52 leu2Δ1 his3Δ200 trp1Δ63*	(Pereira & Schiebel, 2001)
FAY145	*MATa ura3-52 his3Δ200 leu2Δ1 lte1Δ::kanMX6 pRS316-LTE1*	(Bertazzi *et al.*, 2011)
GPY1033	*ESM356-1 KIN4-GFP-his3MX6 Spc42-eqFP-natNT2*	(Pereira & Schiebel, 2005)
GPY1054	*ESM356 Mob1-GFP-kanMX6 Spc42-eqFP-hphNT1*	This study
GPY491	*YPH499 cdc14-2 pRS316- CDC14*	(Caydasi *et al.*, 2017)
GYBY005	*ESM356 Glc7-TAP-kanMX6 (CBP-Tev-4ProtA-kanMX6)*	This study
HKY099	*ESM356 BFA1-GFP-kanMX6 Spc42-eqFP-hphNT1 kar9Δ::klTRP1 pRS316-KAR9 mCherry-Tub1-LEU2*	This study
HKY100	*ESM356 BFA1-GFP-kanMX6 Spc42-eqFP-hphNT1 kar9Δ::klTRP1 pRS316-KAR9 kin4Δ::HIS3MX6 mCherry-Tub1-LEU2*	This study
HKY101	*ESM356 BFA1-GFP-kanMX6 Spc42-eqFP-hphNT1 kar9Δ::klTRP1 pRS316-KAR9 bud14Δ::HIS3MX6 mCherry-Tub1-LEU2*	This study
HKY114	*ESM356 Bfa1-GFP-kanMX6 Spc42-eqFP-hphNT1 mCherry-Tub1-LEU2 pRS416*	This study

HKY115	*ESM356 Bfa1-GFP-kanMX6 Spc42-eqFP-hphNT1 bud14Δ::klTRP1 mCherry-Tub1-LEU2 pRS416*	This study
HKY116	*ESM356 Bfa1-GFP-kanMX6 Spc42-eqFP-hphNT1 bud14Δ::klTRP1 mCherry-Tub1-LEU2 pRS416-BUD14*	This study
HKY133	*FAY145 lte1Δ::kanMX6 pRS316-LTE1 kel1Δ::hphNT1*	This study
HKY134	*FAY145 lte1Δ::kanMX6 pRS316-LTE1 kel2Δ::hphNT1*	This study
HKY135	*FAY145 lte1Δ::kanMX6 pRS316-LTE1 spo12Δ::natNT2 kel1Δ::hphNT1*	This study
HKY136	*FAY145 lte1Δ::kanMX6 pRS316-LTE1 spo12Δ::natNT2 kel2Δ::hphNT1*	This study
HKY139	*ESM356 kar9Δ::HIS3MX6 pRS316-KAR9 kel1Δ::hphNT1*	This study
HKY140	*ESM356 kar9Δ::HIS3MX6 pRS316-KAR9 kel2Δ::hphNT1*	This study
HKY155	*ESM356 BUD14-TAP-kanMX6*	This study
HKY164	*ESM356 Bud14-TAP-kanMX6 Glc7-9myc-HIS3MX6*	This study
HKY171	*ESM356 Mob1-GFP-kanMX6 Spc42-eqFP-hphNT1 mCherry-Tub1-LEU2*	This study
HKY172	*ESM356 Mob1-GFP-kanMX6 Spc42-eqFP-hphNT1 mCherry-Tub1-LEU2 bud14Δ::HIS3MX6*	This study
HKY173	*ESM356 Mob1-GFP-kanMX6 Spc42-eqFP-hphNT1 mCherry-Tub1-LEU2 kin4Δ::natNT2*	This study
HKY174	*ESM356 Mob1-GFP-kanMX6 Spc42-eqFP-hphNT1 mCherry-Tub1-LEU2 bfa1Δ::natNT2*	This study
HKY175	*YPH499 kar9Δ::klTRP1 pRS316-KAR9 GFP-Tub1-ADE2 bud14Δ::HIS3MX6 bfa1Δ::natNT2*	This study
HKY177	*YPH499 bfa1Δ::klTRP1*	This study

HKY180	*YPH499 mob1-67 pRS316-MOB1 bfa1Δ::klTRP3*	This study
HKY182	*YPH499 cdc15-1 pRS316-CDC15 bfa1Δ::klTRP3*	This study
HKY183	*YPH499 cdc5-10 pRS316-CDC5 bfa1Δ::klTRP3*	This study
HKY184	*ESM356 cdc15::CDC15-as1(L99G)-URA3 Bfa1-3HA- klTRP1 bud14Δ::HIS3MX6*	This study
HKY185	*ESM356 pRS416*	This study
HKY186	*ESM356 pMK60 (Bud14-GFP)*	This study
HKY187	*ESM356 pHK002 (Bud14-F379A-GFP)*	This study
HKY188	*ESM356 pDK003 (Bud14-ΔSH3-GFP)*	This study
HKY189	*ESM356 pDK001 (Bud14-5A-GFP)*	This study
HKY190	*ESM356 BFA1-GFP-kanMX6 Spc42-eqFP-hphNT1 bud14Δ::klTRP1 mCherry-TUB1-Leu2 pMK131*	This study
IKY075	*SGY37-VIII,3 kin4Δ::klTRP1*	This study
DKY133	*SGY37-VIII,3 bub2Δ::hphT1*	
IKY192	*ESM356 kar9Δ::HIS3MX6 bud14Δ::klTRP1 pRS416-endogenous BUD14-promoter- GFP-BUD14-5A*	This study
IKY193	*ESM356 kar9Δ::HIS3MX6 bud14Δ::klTRP1 pRS416-endogenous BUD14-promoter- GFP-BUD14-ΔSH3*	This study
JOY79	*YPH499 cdc5-10 pRS316-CDC5*	(Caydasi *et al.*, 2017)
PAY701	*W303 MATa ade2-1 his3-11,15 leu2-3,112 ura3-1 can1-100 ssd1-d2 glc7::LEU2 trp1-1::glc7-12::TRP1*	(Andrews & Stark, 2000)
PAY704	*W303 MATa ade2-1 his3-11,15 leu2-3,112 ura3-1 can1-100 ssd1-d2 glc7::LEU2 trp1-1::GLC7::TRP1*	(Andrews & Stark, 2000)

SEY032	*YPH499 kar9Δ::klTRP3 pRS316-KAR9 GFP-Tub1-ADE2 bud14Δ::HIS3MX6*	This study 975
SEY206	*ESM356 cdc15::CDC15-as1(L99G)-URA3 Bfa1-3HA- kITRP1*	This study 976
SEY212	*ESM356 cdc15::CDC15-as1(L99G)-URA3 Bfa1-3HA- kITRP1 Bud14-9myc-HIS3MX6*	This study
SEY218	*ESM356 cdc15::CDC15-as1(L99G)-URA3 Bfa1-3HA- kITRP1 bud14Δ::HIS3MX6 leu2::LEU2- pGal1-Bud14-9myc-hphNT1*	This study
SEY219	*ESM356 cdc15::CDC15-as1(L99G)-URA3 Bfa1-3HA- kITRP1 bud14Δ::HIS3MX6 leu2::LEU2- pGal1-Bud14- F379A-9myc-hphNT1*	This study
SGY37-VIII,3	*MATa leu2 his3 trp1 ADE2 ura3-52::URA3-lexA-op-LacZ*	(Geissler *et al*, 1996)
SHM562	*YPH499 GFP-Tub1-URA3 dyn1Δ::klTRP1*	(Maekawa *et al.*, 2007)
YDA101	*ESM356 kar9Δ::HIS3MX6 bud14Δ::klTRP1*	(Caydasi *et al.*, 2017)
YPH499	*MATa ura3-52 lys2-801 ade2-101 trp1Δ63 his3Δ200 leu2Δ1*	(Sikorski & Hieter, 1989)

 Table 2. Plasmids used in this study

Plasmid Name	Description	Reference
pRS316	*URA3-dependent CEN-based yeast-E. coli* shuffle plasmid. *AmpR.*	(Sikorski & Hieter, 1989)
pRS416	*URA3-dependent CEN-based yeast-E. coli* shuffle plasmid. *AmpR.*	(Geissler *et al.*, 1996)
pAFS125	*GFP-TUB1-URA3* containing integration plasmid.	(Straight *et al.*, 1997)
pSM1027	*GFP-TUB1-ADE2 containing integration plasmid.*	(Caydasi *et al.*, 2010b)
pAK010	*mCherry-TUB1-kıTRP1* containing integration plasmid.	(Khmelinskii *et al*, 2007)
pAK011	*mCherry-TUB1-URA3* containing integration plasmid.	(Khmelinskii *et al.*, 2007)
pBK067	*mCherry-TUB1-LEU2* containing integration plasmid.	(Caydasi *et al.*, 2014)
pWS103-1	*pRS304-Gal1-UPL-TEM1*	(Shou *et al.*, 1999)
pMM5	p423-Gal1-lexA-Myc	(Geissler *et al.*, 1996)
pMM6	p425-Gal1-Gal4-HA.	(Geissler *et al.*, 1996)
pCDV471	*pEG202 ADH1-LexA-DBD-GLC7*	(Lenssen *et al.*, 2005)
pEG202	*2μm ADH1-LexA-DBD-HIS3*	(Pereira *et al*, 1999)
pCL2-1	*pMM6-BUB2*	(Hofken & Schiebel, 2004)
pCL4a-3	*pMM6-BFA1*	(Hofken & Schiebel, 2004)
pHA132-2	*pMM6-KIN4* (1-750 aa)	Gift from G. Pereira
pHA69-1	*pMM5-BUD14*	Gift from G. Pereira
pHA70-1	*pMM6-BUD14*	Gift from G. Pereira
pTH221	*pMM6-KEL*	(Hofken & Schiebel, 2004)
IKY006	*pMM5-BUD14-5A*	This Study
IKY007	*pMM5-BUD14-F379A*	This Study
IKY008	*pMM5-BUD14-ΔSH3*	This Study
IKY010	*pMM6-BUD14-5A*	This Study

IKY011	*pMM6-BUD14-F379A*	This Study
IKY012	*pMM6-BUD14-ΔSH3*	This Study
pMK125	*pRS416 ADH-BUD14*	(Knaus *et al.*, 2005)
pMK131	*pRS416 ADH-BUD14-F379A*	(Knaus *et al.*, 2005)
pMK60	*pRS416-endogenous BUD14-promoter-GFP-BUD14*	(Knaus *et al.*, 2005)
pDKY001	*pRS416-endogenous BUD14-promoter-GFP-BUD14*$^{5A (135A\ 137A\ 138A\ 139A\ 140A)}$	This Study
pDKY003	*pRS416-endogenous BUD14-promoter-GFP-BUD14*$^{ΔSH3(\ Δaa\ 262–318)}$	This Study
pHK002	*pRS416-endogenous BUD14-promoter-GFP-Bud14-F379A*	This study
pGW399-1	*pRS316-KAR9*	(Caydasi *et al.*, 2010b)
pSM805	*pRS305-pGal1*	(Konig *et al*, 2010)
pHK003	*pSM805 pGal1-Bud14-Leu2*	This study
pHK004	*pSM805 pGal1-Bud14-F379A-Leu2*	This study
pIK007	*pMM5-Bud14-F379A*	This study
pSM903-4	*pRS316-LTE1*	(Hofken & Schiebel, 2002)
pSM908	*pRS316-DBF2*	(Caydasi *et al.*, 2017)
pSM926	*pRS316-MOB1*	(Caydasi *et al.*, 2017)
pUG120	*pRS316-CDC14*	(Gruneberg *et al.*, 2000)
pBS9	*pRS316-CDC15*	(Caydasi *et al.*, 2017)
pCL33	*pRS316-CDC5*	(Hofken & Schiebel, 2004)

References

Adames NR, Cooper JA (2000) Microtubule interactions with the cell cortex causing
nuclear movements in Saccharomyces cerevisiae. *J Cell Biol* 149: 863-874

Adames NR, Oberle JR, Cooper JA (2001) The surveillance mechanism of the spindle
position checkpoint in yeast. *J Cell Biol* 153: 159-168

Andrews PD, Stark MJ (2000) Type 1 protein phosphatase is required for maintenance
of cell wall integrity, morphogenesis and cell cycle progression in Saccharomyces cerevisiae.
Journal of cell science 113 (Pt 3): 507-520

Asakawa K, Yoshida S, Otake F, Toh-e A (2001) A novel functional domain of Cdc15
kinase is required for its interaction with Tem1 GTPase in Saccharomyces cerevisiae. *Genetics*
157: 1437-1450

Bardin AJ, Visintin R, Amon A (2000) A mechanism for coupling exit from mitosis to
partitioning of the nucleus. *Cell* 102: 21-31

Baro B, Queralt E, Monje-Casas F (2017) Regulation of Mitotic Exit in Saccharomyces
cerevisiae. *Methods in molecular biology* 1505: 3-17

Baro B, Rodriguez-Rodriguez JA, Calabria I, Hernaez ML, Gil C, Queralt E (2013) Dual
Regulation of the mitotic exit network (MEN) by PP2A-Cdc55 phosphatase. *PLoS genetics* 9:
e1003966

Beach DL, Thibodeaux J, Maddox P, Yeh E, Bloom K (2000) The role of the proteins
Kar9 and Myo2 in orienting the mitotic spindle of budding yeast. *Curr Biol* 10: 1497-1506

Bertazzi DT, Kurtulmus B, Pereira G (2011) The cortical protein Lte1 promotes mitotic
exit by inhibiting the spindle position checkpoint kinase Kin4. *J Cell Biol* 193: 1033-1048

Bialojan C, Takai A (1988) Inhibitory effect of a marine-sponge toxin, okadaic acid, on
protein phosphatases. Specificity and kinetics. *Biochem J* 256: 283-290

Bishop AC, Ubersax JA, Petsch DT, Matheos DP, Gray NS, Blethrow J, Shimizu E,
Tsien JZ, Schultz PG, Rose MD *et al* (2000) A chemical switch for inhibitor-sensitive alleles
of any protein kinase. *Nature* 407: 395-401

Bloecher A, Venturi GM, Tatchell K (2000) Anaphase spindle position is monitored by
the BUB2 checkpoint. *Nat Cell Biol* 2: 556-558

Campbell IW, Zhou X, Amon A (2020) Spindle pole bodies function as signal amplifiers
in the Mitotic Exit Network. *Molecular biology of the cell* 31: 906-916

Caydasi AK, Ibrahim B, Pereira G (2010a) Monitoring spindle orientation: Spindle
position checkpoint in charge. *Cell division* 5: 28

Caydasi AK, Khmelinskii A, Duenas-Sanchez R, Kurtulmus B, Knop M, Pereira G
(2017) Temporal and compartment-specific signals coordinate mitotic exit with spindle
position. *Nature communications* 8: 14129

Caydasi AK, Kurtulmus B, Orrico MI, Hofmann A, Ibrahim B, Pereira G (2010b) Elm1
kinase activates the spindle position checkpoint kinase Kin4. *J Cell Biol* 190: 975-989

Caydasi AK, Lohel M, Grunert G, Dittrich P, Pereira G, Ibrahim B (2012) A dynamical
model of the spindle position checkpoint. *Molecular systems biology* 8: 582

Caydasi AK, Micoogullari Y, Kurtulmus B, Palani S, Pereira G (2014) The 14-3-3
protein Bmh1 functions in the spindle position checkpoint by breaking Bfa1 asymmetry at
yeast centrosomes. *Molecular biology of the cell* 25: 2143-2151

Caydasi AK, Pereira G (2009) Spindle alignment regulates the dynamic association of
checkpoint proteins with yeast spindle pole bodies. *Dev Cell* 16: 146-156

Caydasi AK, Pereira G (2012) SPOC alert--when chromosomes get the wrong direction.
Experimental cell research 318: 1421-1427

Caydasi AK, Pereira G (2017) Evaluation of the Dynamicity of Mitotic Exit Network
and Spindle Position Checkpoint Components on Spindle Pole Bodies by Fluorescence
Recovery After Photobleaching (FRAP). *Methods in molecular biology* 1505: 167-182

Cenamor R, Jimenez J, Cid VJ, Nombela C, Sanchez M (1999) The budding yeast Cdc15
localizes to the spindle pole body in a cell-cycle-dependent manner. *Mol Cell Biol Res Commun*
2: 178-184

Cheng J, Turkel N, Hemati N, Fuller MT, Hunt AJ, Yamashita YM (2008) Centrosome
misorientation reduces stem cell division during ageing. *Nature* 456: 599-604

Cheng YL, Chen RH (2010) The AAA-ATPase Cdc48 and cofactor Shp1 promote
chromosome bi-orientation by balancing Aurora B activity. *Journal of cell science* 123: 2025-
2034

Chesarone M, Gould CJ, Moseley JB, Goode BL (2009) Displacement of formins from
growing barbed ends by bud14 is critical for actin cable architecture and function. *Dev Cell* 16:
292-302

Cohen P, Klumpp S, Schelling DL (1989a) An improved procedure for identifying and
quantitating protein phosphatases in mammalian tissues. *FEBS Lett* 250: 596-600

Cohen P, Schelling DL, Stark MJ (1989b) Remarkable similarities between yeast and
mammalian protein phosphatases. *FEBS Lett* 250: 601-606

Cullen PJ, Sprague GF, Jr. (2002) The Glc7p-interacting protein Bud14p attenuates
polarized growth, pheromone response, and filamentous growth in Saccharomyces cerevisiae.
Eukaryot Cell 1: 884-894

D'Aquino KE, Monje-Casas F, Paulson J, Reiser V, Charles GM, Lai L, Shokat KM,
Amon A (2005) The protein kinase Kin4 inhibits exit from mitosis in response to spindle
position defects. *Mol Cell* 19: 223-234

Eshel D, Urrestarazu LA, Vissers S, Jauniaux JC, van Vliet-Reedijk JC, Planta RJ,
Gibbons IR (1993) Cytoplasmic dynein is required for normal nuclear segregation in yeast.
Proc Natl Acad Sci U S A 90: 11172-11176

Eskin JA, Rankova A, Johnston AB, Alioto SL, Goode BL (2016) Common formin-
regulating sequences in Smy1 and Bud14 are required for the control of actin cable assembly
in vivo. *Molecular biology of the cell* 27: 828-837

Falk JE, Campbell IW, Joyce K, Whalen J, Seshan A, Amon A (2016a) LTE1 promotes
exit from mitosis by multiple mechanisms. *Molecular biology of the cell* 27: 3991-4001

Falk JE, Chan LY, Amon A (2011) Lte1 promotes mitotic exit by controlling the
localization of the spindle position checkpoint kinase Kin4. *Proc Natl Acad Sci U S A* 108:
12584-12590

Falk JE, Tsuchiya D, Verdaasdonk J, Lacefield S, Bloom K, Amon A (2016b) Spatial
signals link exit from mitosis to spindle position. *eLife* 5

Frenz LM, Lee SE, Fesquet D, Johnston LH (2000) The budding yeast Dbf2 protein
kinase localises to the centrosome and moves to the bud neck in late mitosis. *Journal of cell
science* 113 Pt 19: 3399-3408

Gao K, Zhang Y, Shi Q, Zhang J, Zhang L, Sun H, Jiao D, Zhao X, Tao H, Wei Y *et al*
(2018) iASPP-PP1 complex is required for cytokinetic abscission by controlling CEP55
dephosphorylation. *Cell Death Dis* 9: 528

Geissler S, Pereira G, Spang A, Knop M, Soues S, Kilmartin J, Schiebel E (1996) The
spindle pole body component Spc98p interacts with the gamma-tubulin-like Tub4p of
Saccharomyces cerevisiae at the sites of microtubule attachment. *The EMBO journal* 15: 3899-
3911

Geymonat M, Spanos A, Jensen S, Sedgwick SG (2010) Phosphorylation of Lte1 by Cdk
prevents polarized growth during mitotic arrest in S. cerevisiae. *J Cell Biol* 191: 1097-1112

Geymonat M, Spanos A, Smith SJ, Wheatley E, Rittinger K, Johnston LH, Sedgwick SG
(2002) Control of mitotic exit in budding yeast. In vitro regulation of Tem1 GTPase by Bub2
and Bfa1. *J Biol Chem* 277: 28439-28445

Geymonat M, Spanos A, Walker PA, Johnston LH, Sedgwick SG (2003) In vitro
regulation of budding yeast Bfa1/Bub2 GAP activity by Cdc5. *J Biol Chem* 278: 14591-14594

Gould CJ, Chesarone-Cataldo M, Alioto SL, Salin B, Sagot I, Goode BL (2014)
Saccharomyces cerevisiae Kelch proteins and Bud14 protein form a stable 520-kDa formin
regulatory complex that controls actin cable assembly and cell morphogenesis. *J Biol Chem*
289: 18290-18301

Gruneberg U, Campbell K, Simpson C, Grindlay J, Schiebel E (2000) Nud1p links astral
microtubule organization and the control of exit from mitosis. *The EMBO journal* 19: 6475-
6488

Hofken T, Schiebel E (2002) A role for cell polarity proteins in mitotic exit. *The EMBO
journal* 21: 4851-4862

Hofken T, Schiebel E (2004) Novel regulation of mitotic exit by the Cdc42 effectors Gic1
and Gic2. *J Cell Biol* 164: 219-231

Holder J, Poser E, Barr FA (2019) Getting out of mitosis: spatial and temporal control of
mitotic exit and cytokinesis by PP1 and PP2A. *FEBS Lett* 593: 2908-2924

Hu F, Wang Y, Liu D, Li Y, Qin J, Elledge SJ (2001) Regulation of the Bub2/Bfa1 GAP
complex by Cdc5 and cell cycle checkpoints. *Cell* 107: 655-665

Janke C, Magiera MM, Rathfelder N, Taxis C, Reber S, Maekawa H, Moreno-Borchart
A, Doenges G, Schwob E, Schiebel E *et al* (2004) A versatile toolbox for PCR-based tagging
of yeast genes: new fluorescent proteins, more markers and promoter substitution cassettes.
Yeast 21: 947-962

Jaspersen SL, Charles JF, Tinker-Kulberg RL, Morgan DO (1998) A late mitotic
regulatory network controlling cyclin destruction in Saccharomyces cerevisiae. *Molecular
biology of the cell* 9: 2803-2817

Kelly AE, Funabiki H (2009) Correcting aberrant kinetochore microtubule attachments:
an Aurora B-centric view. *Curr Opin Cell Biol* 21: 51-58

Khmelinskii A, Lawrence C, Roostalu J, Schiebel E (2007) Cdc14-regulated midzone
assembly controls anaphase B. *J Cell Biol* 177: 981-993

Kim HS, Fernandes G, Lee CW (2016) Protein Phosphatases Involved in Regulating
Mitosis: Facts and Hypotheses. *Mol Cells* 39: 654-662

Kim J, Luo G, Bahk YY, Song K (2012) Cdc5-dependent asymmetric localization of
bfa1 fine-tunes timely mitotic exit. *PLoS genetics* 8: e1002450

Knaus M, Cameroni E, Pedruzzi I, Tatchell K, De Virgilio C, Peter M (2005) The
Bud14p-Glc7p complex functions as a cortical regulator of dynein in budding yeast. *The*
EMBO journal 24: 3000-3011

Knop M, Siegers K, Pereira G, Zachariae W, Winsor B, Nasmyth K, Schiebel E (1999)
Epitope tagging of yeast genes using a PCR-based strategy: more tags and improved practical
routines. *Yeast* 15: 963-972

Konig C, Maekawa H, Schiebel E (2010) Mutual regulation of cyclin-dependent kinase
and the mitotic exit network. *J Cell Biol* 188: 351-368

Kotwaliwale C, Biggins S (2006) Microtubule capture: a concerted effort. *Cell* 127:
1105-1108

Krogan NJ, Cagney G, Yu H, Zhong G, Guo X, Ignatchenko A, Li J, Pu S, Datta N,
Tikuisis AP *et al* (2006) Global landscape of protein complexes in the yeast Saccharomyces
cerevisiae. *Nature* 440: 637-643

Lee SE, Frenz LM, Wells NJ, Johnson AL, Johnston LH (2001a) Order of function of
the budding-yeast mitotic exit-network proteins Tem1, Cdc15, Mob1, Dbf2, and Cdc5. *Curr*
Biol 11: 784-788

Lee SE, Jensen S, Frenz LM, Johnson AL, Fesquet D, Johnston LH (2001b) The Bub2-
dependent mitotic pathway in yeast acts every cell cycle and regulates cytokinesis. *Journal of*
cell science 114: 2345-2354

Lenssen E, James N, Pedruzzi I, Dubouloz F, Cameroni E, Bisig R, Maillet L, Werner
M, Roosen J, Petrovic K *et al* (2005) The Ccr4-Not complex independently controls both
Msn2-dependent transcriptional activation--via a newly identified Glc7/Bud14 type I protein
phosphatase module--and TFIID promoter distribution. *Mol Cell Biol* 25: 488-498

Li YY, Yeh E, Hays T, Bloom K (1993) Disruption of mitotic spindle orientation in a
yeast dynein mutant. *Proc Natl Acad Sci U S A* 90: 10096-10100

Luca FC, Mody M, Kurischko C, Roof DM, Giddings TH, Winey M (2001)
Saccharomyces cerevisiae Mob1p is required for cytokinesis and mitotic exit. *Mol Cell Biol*
21: 6972-6983

Maekawa H, Priest C, Lechner J, Pereira G, Schiebel E (2007) The yeast centrosome
translates the positional information of the anaphase spindle into a cell cycle signal. *J Cell Biol*
179: 423-436

Manzano-Lopez J, Monje-Casas F (2020) The Multiple Roles of the Cdc14 Phosphatase
in Cell Cycle Control. *Int J Mol Sci* 21

Meitinger F, Khmelinskii A, Morlot S, Kurtulmus B, Palani S, Andres-Pons A, Hub B,
Knop M, Charvin G, Pereira G (2014) A memory system of negative polarity cues prevents
replicative aging. *Cell* 159: 1056-1069

Meitinger F, Palani S, Pereira G (2016) Detection of Phosphorylation Status of
Cytokinetic Components. *Methods in molecular biology* 1369: 219-237

Meraldi P, Nigg EA (2001) Centrosome cohesion is regulated by a balance of kinase and
phosphatase activities. *Journal of cell science* 114: 3749-3757

Miller RK, Rose MD (1998) Kar9p is a novel cortical protein required for cytoplasmic
microtubule orientation in yeast. *J Cell Biol* 140: 377-390

Molk JN, Schuyler SC, Liu JY, Evans JG, Salmon ED, Pellman D, Bloom K (2004) The
differential roles of budding yeast Tem1p, Cdc15p, and Bub2p protein dynamics in mitotic
exit. *Molecular biology of the cell* 15: 1519-1532

Monje-Casas F, Amon A (2009) Cell polarity determinants establish asymmetry in MEN
signaling. *Dev Cell* 16: 132-145

Morishita T, Mitsuzawa H, Nakafuku M, Nakamura S, Hattori S, Anraku Y (1995)
Requirement of Saccharomyces cerevisiae Ras for completion of mitosis. *Science* 270: 1213-
1215

Moura M, Conde C (2019) Phosphatases in Mitosis: Roles and Regulation. *Biomolecules*
9

Musacchio A, Salmon ED (2007) The spindle-assembly checkpoint in space and time.
Nature reviews Molecular cell biology 8: 379-393

Nasa I, Kettenbach AN (2018) Coordination of Protein Kinase and Phosphoprotein
Phosphatase Activities in Mitosis. *Front Cell Dev Biol* 6: 30

Neller J, Dunkler A, Rosler R, Johnsson N (2015) A protein complex containing Epo1p
anchors the cortical endoplasmic reticulum to the yeast bud tip. *J Cell Biol* 208: 71-87

O'Connell CB, Wang YL (2000) Mammalian spindle orientation and position respond to
changes in cell shape in a dynein-dependent fashion. *Molecular biology of the cell* 11: 1765-
1774

Perdiguero E, Nebreda AR (2004) Regulation of Cdc25C activity during the meiotic
G2/M transition. *Cell Cycle* 3: 733-737

Pereira G, Grueneberg U, Knop M, Schiebel E (1999) Interaction of the yeast gamma-
tubulin complex-binding protein Spc72p with Kar1p is essential for microtubule function
during karyogamy. *The EMBO journal* 18: 4180-4195

Pereira G, Hofken T, Grindlay J, Manson C, Schiebel E (2000) The Bub2p spindle
checkpoint links nuclear migration with mitotic exit. *Mol Cell* 6: 1-10

Pereira G, Manson C, Grindlay J, Schiebel E (2002) Regulation of the Bfa1p-Bub2p
complex at spindle pole bodies by the cell cycle phosphatase Cdc14p. *J Cell Biol* 157: 367-379

Pereira G, Schiebel E (2001) The role of the yeast spindle pole body and the mammalian
centrosome in regulating late mitotic events. *Curr Opin Cell Biol* 13: 762-769

Pereira G, Schiebel E (2005) Kin4 kinase delays mitotic exit in response to spindle
alignment defects. *Mol Cell* 19: 209-221

Pereira G, Yamashita YM (2011) Fly meets yeast: checking the correct orientation of cell
division. *Trends Cell Biol*

Pinsky BA, Kotwaliwale CV, Tatsutani SY, Breed CA, Biggins S (2006) Glc7/protein
phosphatase 1 regulatory subunits can oppose the Ipl1/aurora protein kinase by redistributing
Glc7. *Mol Cell Biol* 26: 2648-2660

Rock JM, Amon A (2009) The FEAR network. *Curr Biol* 19: R1063-1068

Ruchaud S, Carmena M, Earnshaw WC (2007) Chromosomal passengers: conducting
cell division. *Nature reviews Molecular cell biology* 8: 798-812

Scarfone I, Piatti S (2015) Coupling spindle position with mitotic exit in budding yeast:
The multifaceted role of the small GTPase Tem1. *Small GTPases* 6: 196-201

Scarfone I, Venturetti M, Hotz M, Lengefeld J, Barral Y, Piatti S (2015) Asymmetry of
the budding yeast Tem1 GTPase at spindle poles is required for spindle positioning but not for
mitotic exit. *PLoS genetics* 11: e1004938

Seshan A, Bardin AJ, Amon A (2002) Control of Lte1 localization by cell polarity
determinants and Cdc14. *Curr Biol* 12: 2098-2110

Sherman F (1991) Getting started with yeast. *Methods Enzymol* 194: 3-21

Shirayama M, Matsui Y, Tanaka K, Toh-e A (1994a) Isolation of a CDC25 family gene,
MSI2/LTE1, as a multicopy suppressor of ira1. *Yeast* 10: 451-461

Shirayama M, Matsui Y, Toh EA (1994b) The yeast TEM1 gene, which encodes a GTP-
binding protein, is involved in termination of M phase. *Mol Cell Biol* 14: 7476-7482

Shou W, Seol JH, Shevchenko A, Baskerville C, Moazed D, Chen ZW, Jang J,
Charbonneau H, Deshaies RJ (1999) Exit from mitosis is triggered by Tem1-dependent release
of the protein phosphatase Cdc14 from nucleolar RENT complex. *Cell* 97: 233-244

Sikorski RS, Hieter P (1989) A system of shuttle vectors and yeast host strains designed
for efficient manipulation of DNA in Saccharomyces cerevisiae. *Genetics* 122: 19-27

Siller KH, Doe CQ (2009) Spindle orientation during asymmetric cell division. *Nat Cell*
Biol 11: 365-374

Stegmeier F, Visintin R, Amon A (2002) Separase, polo kinase, the kinetochore protein
Slk19, and Spo12 function in a network that controls Cdc14 localization during early anaphase.
Cell 108: 207-220

Straight AF, Marshall WF, Sedat JW, Murray AW (1997) Mitosis in living budding
yeast: anaphase A but no metaphase plate. *Science* 277: 574-578

Valerio-Santiago M, Monje-Casas F (2011) Tem1 localization to the spindle pole bodies
is essential for mitotic exit and impairs spindle checkpoint function. *J Cell Biol* 192: 599-614

Visintin R, Amon A (2001) Regulation of the mitotic exit protein kinases Cdc15 and
Dbf2. *Molecular biology of the cell* 12: 2961-2974

Wang Y, Hu F, Elledge SJ (2000) The Bfa1/Bub2 GAP complex comprises a universal
checkpoint required to prevent mitotic exit. *Curr Biol* 10: 1379-1382

Weiss EL (2012) Mitotic Exit and Separation of Mother and Daughter Cells. *Genetics*
192: 1165-1202

Wu JQ, Guo JY, Tang W, Yang CS, Freel CD, Chen C, Nairn AC, Kornbluth S (2009)
PP1-mediated dephosphorylation of phosphoproteins at mitotic exit is controlled by inhibitor-
1 and PP1 phosphorylation. *Nat Cell Biol* 11: 644-651

Yeh E, Skibbens RV, Cheng JW, Salmon ED, Bloom K (1995) Spindle dynamics and
cell cycle regulation of dynein in the budding yeast, Saccharomyces cerevisiae. *J Cell Biol* 130:
687-700

Yoshida S, Toh-e A (2001) Regulation of the localization of Dbf2 and mob1 during cell
division of saccharomyces cerevisiae. *Genes Genet Syst* 76: 141-147
1232

Figure 1

Figure 2

Figure 3

Figure 4

A

16 °C   23 °C   30 °C

lteΔ bud14Δ
lte1Δ
lteΔ bud14Δ BUD14
lteΔ bud14Δ bud14-F379A

SC-URA

B

SPOC deficiency index

*

BUD14   empty   bud14-F379A

kar9Δ bud14Δ

C

18 °C   33 °C   35 °C   37 °C

GLC7
glc7-12
lte1Δ glc7-12
lte1Δspo12Δ glc7-12

pRS316-LTE1

5FOA

D

SPOC deficiency index

*

kar9Δ GLC7   kar9Δ glc7-12

Figure 5

Figure 6

Figure 7

Figure 2 - Figure Supplement 1

A

Gal1-*KIN4*
Gal1-*KIN4 bub2*Δ
Gal1-*KIN4 bud14*Δ

Galactose   Glucose

B

	Kin4-GFP	Spc42-eqFP	MERGE	MERGE + Phase
WT				
*bud14*Δ				
*rts1*Δ				

*****

% Cells

WT   *bud14*Δ   *rts1*Δ

■ Kin4-GFP at the cortex   ■ Kin4-GFP at the SPB

C

	Kin4-GFP	Spc42-eqFP mCherry-Tubulin	MERGE
*lte1*Δ			
*bud14*Δ *lte1*Δ			

% Anaphase Cells

n.s.   n.s.   n.s.

both SPBs   mSPB   No SPB

■ *lte1*Δ   ■ *bud14*Δ *lte1*Δ

D

KIN4-6HA

*bud14*Δ   *rts1*Δ   -

KIN4-6HA

Clb2

Tubulin

1DAPI %   88   90   85

Figure 3 - Figure Supplement 1

Figure 4 - Supplement 1

Figure 5 - Figure Supplement 1

Figure 6 - Figure Supplement 1

Figure 6 - Figure Supplement 3

Figure 7 - Figure Supplement 1

A

lte1Δ

0 15 30 45 60 75 90 105 120

lte1Δ kin4Δ

0 15 30 45 60 75 90 105 120

Bfa1-3HA

Clb2

Tubulin

Bfa1-3HA Gal1-UPL-Tem1

B

lte1Δ
lte1Δ bud14Δ
lte1Δ kin4Δ

75 min

90 min

105 min

Bfa1-3HA

C

lte1Δ

lte1Δ kin4Δ

— no bud
— small bud
— large bud

— 1 DAPI
— 2 DAPI

Figure 7 - Figure Supplement 2

A

WT
Gal1-*BUD14*
Gal1-*BUD14 kin4Δ*
Gal1-*BUD14 bfa1Δ*

Glucose   Raffinose/Galactose

B

Bfa1-3HA   +   +   +
Bud14-TAP   –   +   +
Okadaic acid –   –   +

Bfa1-3HA

Glc7-9myc

Bud14-TAP

1   2   3