Time-course analysis of *Streptococcus sanguinis* after manganese depletion reveals

changes in glycolytic, nucleotide, and redox metabolites

Tanya Puccio[1], Biswapriya B. Misra[2], Todd Kitten[1*]

[1]Philips Institute for Oral Health Research, Virginia Commonwealth University School of Dentistry,

Richmond 23298, VA USA.

[2]Department of Internal Medicine, Section on Molecular Medicine, Wake Forest School of Medicine,

Medical Center Boulevard, Winston-Salem 27157, NC USA.

*Corresponding author:

Todd Kitten

tkitten@vcu.edu

804-628-7010

Keywords: metabolomics, manganese, endocarditis, multivariate, time-course

Short title: Time-course metabolomics of Mn-deplete *S. sanguinis*

Abstract

Introduction

Manganese is important for the endocarditis pathogen, *Streptococcus sanguinis*. Little is known
about why manganese is required for virulence or how it impacts the metabolome of streptococci.

Objectives

We applied untargeted metabolomics to cells and the Brain Heart Infusion media they were growing
in to understand temporal changes resulting from manganese depletion.

Methods

EDTA was added to a *S. sanguinis* manganese-transporter mutant in aerobic fermentor conditions.
Cell and media samples were collected pre- and post-EDTA treatment. Metabolomics data were
generated using positive and negative modes of data acquisition on an LC-MS/MS system. Data were
subjected to statistical processing using MetaboAnalyst and time-course analysis using Short Time
series Expression Miner (STEM).

Results

We observed quantitative changes in 534 and 422 metabolites in cells and media, respectively. The
173 cellular metabolites identified as significantly different indicated enrichment of purine and
pyrimidine metabolism. Further multivariate analysis revealed that the top 15 cellular metabolites
belonged to lipids, cCMP, cUMP, and redox metabolites. The STEM analysis revealed global
changes in cells and media in comparable metabolic pathways. Products of glycolysis such as
pyruvate and fructose-1,6-bisphosphate increased after EDTA addition, which indicates catabolic
enzymes may require a manganese cofactor. Nucleosides accumulated after depletion, possibly due
to a blockage in the conversion to nucleobases. Accumulation of *ortho*-tyrosine suggests the cells
were oxidized but unable to regulate utilization of redox metabolites such as glutathione.

Conclusion

Differential analysis of metabolites revealed the activation of a number of metabolic pathways in
response to manganese depletion, many of which may be connected to carbon catabolite repression.

1. Introduction

Streptococcus sanguinis is a gram-positive bacterium known for its duplicity. As an early and abundant colonizer of teeth, *S. sanguinis* is associated with oral health (Kreth *et al.*, 2017; Kreth *et al.*, 2005). However, when it enters the bloodstream, whether through dental procedures or activities as routine as eating, it is known to colonize the heart valves or other endocardial surfaces of persons with particular pre-existing cardiac conditions, leading to infective endocarditis (IE) (Moreillon *et al.*, 2002; Widmer *et al.*, 2006). IE has a global mortality rate of 12-40% (Bor *et al.*, 2013; Cahill *et al.*, 2017). Historically, prevention has relied upon administration of prophylactic broad-spectrum antibiotics to high-risk patients prior to dental visits (Wilson *et al.*, 2007). With rising antibiotic resistance (Dodds, 2017), as well as controversial efficacy (Quan *et al.*, 2020; Thornhill *et al.*, 2018), novel drug targets that are required for endocarditis causation but not beneficial colonization are under investigation.

One such putative drug target in *S. sanguinis* is the lipoprotein SsaB, a component of the ATP-binding cassette transporter SsaACB. This transporter and orthologs in related species have been shown to be important for manganese (Mn) transport and virulence (Colomer-Winter *et al.*, 2018; Crump *et al.*, 2014; Dintilhac *et al.*, 1997; Kehl-Fie *et al.*, 2013). Previous studies utilizing a Δ*ssaACB* strain of *S. sanguinis* revealed that this mutant is significantly deficient in cellular Mn levels (Murgas *et al.*, 2020) and virulence in our rabbit model of IE (Baker *et al.*, 2019). These studies also suggested that the reduced virulence of Mn-deficient cells is due to growth arrest in the aerobic, low-Mn environment characteristic of an aortic valve infection, implying the existence of one or more Mn-dependent metabolic pathways that are essential for aerobic growth. The metabolic pathway(s) and individual metabolites involved have not been defined.

Metabolomics is the comprehensive study of small molecules in the molecular weight range of 50-2000 Da in biological systems. Diverse mass spectrometry platforms such as LC-MS/MS, GC-MS and CE-MS with and without chromatography, and spectroscopy technologies such as NMR have enabled high-throughput discovery metabolomics in various biological systems, including bacteria, plants, and humans (Misra and Olivier, 2020). Recent studies have described the metabolomes of certain streptococci using various mass spectrometry methods: *Streptococcus intermedius* under various oxygen conditions (Fei *et al.*, 2016); *Streptococcus pneumoniae* in chemically defined

medium (Leonard *et al.*, 2018); and *Streptococcus thermophilus* in pH-controlled batch fermentation

(Liu *et al.*, 2020; Qiao *et al.*, 2019). To our knowledge, the metabolome of *S. sanguinis* has yet to be

investigated. Here we report the first untargeted metabolomic analysis of *S. sanguinis* or, indeed, of

any *Streptococcus*, under Mn replete vs. deplete conditions.

2. Materials and Methods

2.1 Bacterial strains and growth conditions

S. sanguinis strain SK36 was isolated from human dental plaque (Kilian *et al.*, 1989; Xu *et al.*,

2007). The Δ*ssaACB* strain (JFP169) was generated from SK36 previously by replacement of the

ssaACB genes with the *aphA-3* gene encoding kanamycin resistance (Puccio *et al.*, 2020). Overnight

pre-cultures were created by inoculation of Brain Heart Infusion (BHI) broth (Becton, Dickinson and

Company, Franklin Lakes, NJ) with single-use aliquots of cryopreserved cells by 1000-fold dilution.

Kanamycin (Sigma-Aldrich, St. Louis, MO) was added to 500 µg mL^{-1} for Δ*ssaACB* pre-cultures.

Pre-cultures were incubated at 37°C for 18 h in 6% O_2 (6% O_2, 7% H_2, 7% CO_2 and 80% N_2) using

an Anoxomat (Advanced Instruments, Norwood, MA) jar.

2.2 Fermentor growth conditions and sample collection

Aerobic fermentor growth of Δ*ssaACB* cell culture was achieved using a BIOSTAT® B bioreactor

(Sartorius Stedim, Göttingen, Germany) and samples were collected as described in Puccio and

Kitten (2020). The pre-EDTA sample was collected at T_{-20} (min), where EDTA was added at T_0 to a

final concentration of 100 µM. Post-EDTA samples were collected at T_{25} and T_{50}. All samples were

stored at -80°C until shipped on dry ice to Metabolon, Inc. (Durham, North Carolina) for further

analysis.

2.3 Sample preparation, UPLC-MS/MS, data extraction, compound identification, and

curation

Metabolomics sample processing was completed by Metabolon, Inc. as described in the

Supplementary Methods and in previous publications (Dehaven *et al.*, 2010; Evans *et al.*, 2009).

2.4 Statistical analysis of metabolomics and transcriptomics datasets

Statistical analysis of the metabolomics data sets was performed using statistical software R (Version 3.5.2)(Team, 2018). Normalized, transformed, imputed, outlier-removed, and scaled peak areas representative of relative metabolite amounts obtained using DeviumWeb (Grapov, 2014) are presented. Hierarchical clustering analysis (HCA) was performed on Pearson distances using MetaboAnalyst 4.0 (www.metaboanalyst.ca) (Xia *et al.*, 2015), with the data normalized using z-scores of the relative abundance of the metabolites for heat map display. Correlations reported are Spearman rank correlations. Principal component analysis (PCA) and partial least squared discriminant analyses (PLS-DA) were performed using MetaboAnalyst, with the output displayed as score plots for visualization of sample groups. One-way analysis of variance (ANOVA) followed by post-hoc analysis using Fisher's least significant difference (LSD) test was used for analysis of statistical significance using MetaboAnalyst.

2.5 Time-course analysis of cellular and media metabolomes

For our 70 min time course, we used the Short Time series Expression Miner (STEM) tool, originally used for short microarray time series experiments (3–8 time points for $>$ ~80% of the datasets). The following parameters were used for our analysis: no normalization of data; 0 added as the starting point; number of model profiles = 20; maximum unit change in model profiles between time points = 3. To explain the model profiles, we used an expression of -1 if levels of a metabolite decreased, 0 if levels were unchanged, and 1 if levels increased. For instance, a model profile with an expression of -1, -1, 0, represents decreased, decreased, and unchanged, levels of a given set of metabolites for the 3 time points.

2.6 Metabolic pathway and enrichment analysis

Pathway enrichment analysis was performed using MetaboAnalyst 4.0 and reported pathways are KEGG-based (Kanehisa and Goto, 2000). The Chemical Translation Service (CTS: http://cts.fiehnlab.ucdavis.edu/conversion/batch) was used to convert the common chemical names into their KEGG, HMDB, Metlin, PubChem CID, and ChEBI identifiers.

3. Results

3.1 EDTA treatment of Δ*ssaACB* cells leads to Mn depletion and slowed growth

As described in Puccio *et al.* (2020), EDTA treatment of Δ*ssaACB* aerobic fermentor-grown cells results in the depletion of Mn but no other biologically relevant metals, such as Fe or Zn, as determined by inductively coupled plasma optical emission spectroscopy (ICP-OES) **(Figure 1)**. Beginning ~38 min post-EDTA addition, cell growth slowed, resulting in a steady drop in OD **(Figure 1)**.

3.2 Global metabolomics of *S. sanguinis* cells and BHI media

Our goal was to understand the metabolic consequences of Mn depletion during growth of a *S. sanguinis* Mn-transporter mutant in a rich medium (BHI), as well as to survey changes in the conditioned media during the growth and treatment periods. Extensive global untargeted metabolomics analysis revealed 534 metabolites in cells and 422 metabolites in conditioned media. The raw metabolite abundance values alongside the identified metabolite IDs, super pathways and sub-pathway names, average mass, and identifiers such as Chemical Abstracts Service (CAS), PubChem, Kyoto Encyclopedia of Genes and Genomes (KEGG), and Human Metabolome Database (HMDB) IDs are provided for both cellular and media metabolites (**Tables S1-S2**). Further, these datasets were refined after normalization, transformation, and scaling, followed by imputation (**Tables S3-S4).** The 534 metabolites belong to 57 different KEGG metabolic pathways (**Table S5**). The 422 metabolites quantified in the conditioned BHI media belonged to 50 different metabolic pathways (**Table S6**), all of which overlap with the metabolic pathways found in the cells.

BHI has as its chief constituents bovine and porcine brain and heart extracts. Based on comparison with the pre-inoculation media samples, we identified several metabolites such as sucrose, caprylate (8:0), 5-methyluridine (ribothymidine), 2'-deoxyuridine, 5-aminoimidazole-4-carboxamide that appear to originate from BHI, and were excluded from further statistical processing as they were unique to the growth media alone **(Table S7)**. Any metabolite that occurred in less than 75% of the samples was also excluded from the analysis, which resulted in the exclusion of 9 of the 534 metabolites detected in cells **(Table S7)**.

3.3 Differential accumulation patterns of metabolites over time course and EDTA treatment

We used a false discovery rate (FDR)-corrected ANOVA to determine metabolites that were significantly different in abundance between the different time-points. ANOVA revealed 173 and 13 metabolites that were significantly different in cells and media, respectively **(Tables S8-9)**. There were no metabolites that ranked within the top 10 from both sources. (**Table 1**). To investigate whether these differential metabolites would map to metabolic pathways, we mapped the set of metabolites using the *Streptococcus pyogenes* M1 476 KEGG database within MetaboAnalyst by implementing overrepresentation analysis with Fisher's exact test and pathway topology analysis using relative-betweenness centrality (Jewison *et al.*, 2014). Pathway enrichment analysis of the 173 cellular metabolites that were differential along the time course of EDTA treatment identified only purine and pyrimidine metabolism (nominal P-value < 0.05) (**Figure S1a; Table S10)**. Surprisingly pathway enrichment analysis of the 13 media metabolites that were differential along the time course belonged to purine and pyrimidine metabolism as above, but also glyoxylate and dicarboxylate metabolism, and alanine, aspartate, and glutamate metabolism (nominal P-value < 0.05) (**Figure S1b; Table S11**). When metabolite abundances were compared for the two post-EDTA time points vs T_{-20}, it was revealed that one, five, 13, and 30 metabolites were increased in T_{25} and T_{50} in media and T_{25} and T_{50} in cells, respectively **(Figure S1c)**. Of these, only 2'-deoxyadenosine increased in both cells and media at T_{50} (**Tables S12-13**). The 30 metabolites increased in T_{50} in cells were mostly lipids, energy metabolites, nucleotide phosphates, and dinucleotides (**Table S12**). When significantly decreased metabolites were compared, it was revealed that 1, 1, 13, and 30 metabolites were decreased in T_{25} and T_{50} in media and T_{25} and T_{50} in cells, respectively (**Figure S1d**). Only glutamine levels decreased in both media samples (**Table S13**). The 5 metabolites that decreased in cells at T_{25} included cCMP and cUMP, while the 18 metabolites that decreased at T_{50} in the cells included IMP and XMP (**Table S12**).

3.4 Multivariate and hierarchical clustering analysis

To define the metabolomic changes caused by Mn depletion, we used multivariate analysis and HCA. Using an unsupervised multivariate analysis, PCA, we observed that metabolite abundances alone were able to discriminate between the samples and explain 58.8% of the variation in the dataset by virtue of the first 2 PCs (PC1, PC2) in cells (**Figure 2a)** and 67.5% in media **(Figure 2b).**

Using supervised multivariate analysis, PLS-DA, we observed that metabolite abundances alone
were able to discriminate between the samples and explain 57.1 % of the variation in the dataset by
virtue of the first 2 PCs (Component 1 and 2) in cells (**Figure S2a)** and 57.7% in media **(Figure**
S2b). Additionally, PLS-DA and PCA performed on spent media samples explained 93.4% and
93.5% of the variation, respectively, by virtue of the first 2 PCs **(Figure S2c-d)**.

To identify the metabolites responsible for the discrimination among the metabolomic profiles, the
variable importance in projection (VIP) score was used to select features with the most significant
contribution in a PLS-DA model. VIP scores are a weighted sum of PLS weights for each variable
and measure the contribution of each predictor variable to the model. Further, the VIP statistic
summarizes the importance of the metabolites in differentiating the sample time points in
multivariate space. Metabolites exhibiting high VIP scores ($\geq$1.5) are the more influential variables.
Our VIP analysis revealed that the top 15 metabolites for cells belonged to lipids, cCMP, cUMP, and
redox metabolites **(Figure 2c).** The VIP analysis revealed that the top 15 metabolites for spent media
belonged to amino acids and organic acids **(Figure 2d).** Of these VIP metabolites (cut off $\geq$1.5), 7
metabolites (glutamine, adenosine, adenine, glycerate, forminoglutamate, citrulline, and orotate)
were shared between cells and media across all the time points, indicating their importance.

We performed an HCA using the z-score-normalized metabolite abundances of the cellular and
media metabolites, separately **(Figure S3)**. Results indicated a clear clustering for the three time
points as shown for the top 25 metabolites obtained from the ANOVA for individual sample groups.
In cells, two distinct clusters were formed based on the metabolite abundances, where the upper
cluster (decreased in T_{50}) was represented by acetylated metabolites, purines and pyrimidines, and
glutamyl dipeptides, and the bottom cluster (increased in T_{50}) contained several amino acids and
lipids, and cCMP, cUMP, and UTP **(Figure S3a).** In media, two distinct clusters were formed based
on the metabolite abundances, with the upper cluster (increased in T_{50}) represented by several
important metabolites such as uracil, ribose, pyruvate, nicotinamide, inosine, adenosine, guanosine,
and the bottom cluster (decreased in T_{50}) containing glutamine, adenine, and 3'AMP **(S3b).**

3.5 Time-course analysis of cellular and media metabolites

To understand the time course-dependent changes in metabolite accumulation patterns across the
three time points in this complex study design, we started with a clustering analysis. Using STEM

analysis, we interrogated the time course changes of the metabolites in the cells and media. The metabolite abundances for all quantified metabolites across the three time points were put into 20 model clusters, which revealed differential accumulation of metabolites for media and cells as a function of time. For the cells, the top two significant models were #19 (pattern 0, 1, 1, -1, P-value 5e-115) and #18 (pattern 0, 1, -1, 0, P-value 4e-12) representing 193 and 80 metabolites, respectively **(Figure S4a; Table S14)**. Metabolites following the pattern in model #19 were enriched for amino acid metabolic pathways: valine, leucine and isoleucine biosynthesis and degradation, alanine, aspartate and glutamate metabolism, and glycine, serine and threonine metabolism (P-value, < 0.1). Model #18 metabolites were enriched for arginine biosynthesis, arginine and proline metabolism, histidine metabolism, glyoxylate and dicarboxylate metabolism, and pyrimidine metabolism (P-value < 0.1). For the media, the top three models were #18 (0, 1, -1, 0, P-val- 3e-59), #19 (pattern 0, 1, 1, -1, P-val- 3e-23) and #14 (pattern 1, 1, 1, 1, P-val-6e-24) representing 132, 81, and 4 metabolites, respectively **(Figure S4b and Table S15)**. Metabolites following the pattern in model #18 were enriched for alanine, aspartate and glutamate metabolism, amino acid metabolism, and arginine and proline metabolism. Those in model #19 were enriched for arginine biosynthesis, valine, leucine and isoleucine biosynthesis and degradation, glyoxylate and dicarboxylate metabolism, pyrimidine metabolism, alanine, aspartate and glutamate metabolism, and glycine, serine and threonine metabolism. The metabolites in model #14 included 2-deoxyadenosine, N6-methyladenosine, inosine, and nicotinamide.

4.   Discussion

4.1 Metabolomic analysis of BHI spent media reveals metabolic interactions of *S. sanguinis* with the extracellular environment

Our purpose in conducting this study was to examine the role of Mn in *S. sanguinis* metabolism, particularly in relation to IE. While the perfect medium for such a study would have been serum or plasma, this would not have been feasible, and so we instead used another complex yet commercially accessible medium—BHI. As with plasma, BHI has glucose as its most abundant sugar (0.2% w/v in BHI and ~0.1% w/v in plasma). Although serum and plasma have been the subject of many metabolomic studies, we are not aware of any previous metabolomic analysis of BHI. Thus, the analysis of the pre-inoculated BHI (**Table S2**) may be of interest to the many investigators who use this medium. Likewise, the comparison of the pre-inoculated and T_{-20} media samples tells us much

concerning the metabolic and transport capabilities of *S. sanguinis* under Mn-replete conditions (**Table S13**).

As expected, we observed a significant decrease of glucose in spent media (**Figure 3a**), indicating its utilization as carbon source. Levels of fructose and mannose significantly decreased as well **(Figure 3a),** indicating that they are catabolized by cells. *S. sanguinis* encodes a number of putative sugar transport systems (Ajdic and Pham, 2007; Xu *et al.*, 2007). Lactate and pyruvate levels increased significantly in the media after cell growth (**Figure 3b**), indicating that these products of glycolysis have been secreted from cells. Pyruvate has been shown to be secreted by *S. sanguinis*, presumably to protect the cells from H_2O_2 stress by acting as an antioxidant (Redanz *et al.*, 2020).

Also of interest, all nucleosides were significantly decreased after *S. sanguinis* growth (**Figures 4 and S5a-b**). The opposite trend was observed with nucleobases, where most were significantly increased after cell growth (**Figures 4 and S5c-d**). Nucleoside transport for salvage has been characterized in many bacteria, including the related species *Lactococcus lactis* (Martinussen *et al.*, 2010) and *Streptococcus mutans* (Webb and Hosie, 2006).

4.2 Carbohydrate metabolism and glycolytic regulation in *S. sanguinis* cells show Mn dependence

The levels of glycolytic byproducts in *S. sanguinis* cells and spent media were impacted by Mn depletion. Glucose, fructose, and lactate levels remained constant in cells at all three time points while pyruvate levels increased after Mn depletion **(Figure 3b)**. Mannose and sucrose were not detected in cells at any time point, indicating rapid catabolism by cells (**Figure 3b**). Lactate is known to be produced in high levels by streptococci and other lactic acid bacteria (Jakubovics *et al.*, 2014), which explains the observed increase of lactate in the media after cellular growth. Pyruvate is produced through metabolism of sugars or amino acids. The observed increase in pyruvate levels in cells after Mn depletion **(Figure 4b)** is not due to increased sugar levels, as the flow of media remained constant throughout the experiment. Most amino acid levels remained unchanged or decreased in cells after Mn depletion (**Table S11**). One potential explanation for the increase in pyruvate levels is that fewer pyruvate molecules were oxidized by pyruvate oxidase (SpxB) into H_2O_2 and acetyl phosphate, consistent with our finding of a significant decrease in H_2O_2 levels after Mn depletion (**Figure 1**) (Puccio *et al.*, 2020).

There was a significant accumulation of hexose diphosphates in cells at T_{50} and a slight increase in
spent media as well (**Figure 3**). Since levels of other glycolytic intermediates such as glucose-6-
phosphate, glycerone, and glyceraldehyde-3-phosphate could not be measured using our platform
(Tables S1-2), we are unable to assess the impact on this pathway using metabolomics alone. We
hypothesize that the hexose diphosphate is primarily fructose-1,6-bisphosphate and its accumulation
results from the reduced activity of two potentially Mn-cofactored fructose-1,6-bisphosphate-
consuming enzymes in the glycolytic pathway: fructose-1,6-bisphosphatase (Fbp; SSA_1056) and
fructose-bisphosphate aldolase (Fba; SSA_1992) (Puccio *et al.*, 2020). We further hypothesize that
fructose-1,6-bisphosphate accumulation is at least partly responsible for the glucose-independent
CcpA repression observed in the transcriptome of *S. sanguinis* after Mn depletion (Puccio *et al.*,
2020).

Previous studies with other bacteria support a role for Mn in carbon metabolism. Mn deprivation was
previously found to induce flux to the pentose phosphate pathway in *S. pneumoniae* (Ogunniyi *et al.*,
2010). *Staphylococcus aureus* was found to be more susceptible to calprotectin-mediated Mn
starvation when glucose was the sole carbon source than when amino acids were also present (Radin
et al., 2016). A recent study found that excess Mn modulated glycolysis in *Escherichia coli* biofilms
by decreasing levels of glucose-6-phosphate and glyceraldehyde-3-phosphate (Guo and Lu, 2020).
Here we provide further evidence that Mn levels impact central carbon metabolism.

4.3 Purine and pyrimidine metabolism in Mn-deplete *S. sanguinis* reveal nucleoside utilization
from media and nucleobase accumulation in cells

Mn is known to impact nucleotide metabolism through its role as cofactor for the aerobic
ribonucleotide reductase NrdF (Makhlynets *et al.*, 2014; Rhodes *et al.*, 2014). Here, we observed
further impacts of Mn on nucleotide metabolism. Mean levels of guanosine, inosine, and adenosine
increased in both cells and media at T_{50} (**Figures 4 and S5a & e**). In cells, guanine levels decreased
while hypoxanthine and adenine levels were unchanged at T_{50} (**Figures 4 and S5g**). This indicates
that there may be blockages in the conversion of purine nucleobases into nucleosides. There are three
enzymes encoded by *S. sanguinis* that can catalyze this reaction: PunA (SSA_1258), DeoD
(SSA_1259), and SSA_2046. None of these enzymes have been found to use Mn according to
BRENDA (https://www.brenda-enzymes.org/) (Jeske *et al.*, 2019). In our recent transcriptomics
study, expression of *punA* and *deoD* were significantly decreased after Mn depletion while

SSA_2046 expression was unchanged (Puccio *et al.*, 2020). The operon encoding *deoD* and *punA*

has a carbon responsive element (*cre*) upstream of the first gene, *rpiA* (Bai *et al.*, 2019), which is the

recognition sequence for the carbon catabolite repression (CCR) regulator CcpA (Warner and

Lolkema, 2003). As observed in Puccio *et al.* (2020), Mn depletion results in many changes in the

CcpA regulon, which may explain the repression of this operon at T_{50}. Thus, this may be but one

example of a non-carbon metabolism pathway impacted by Mn depletion through its effect on CCR.

Similar to the purines, the pyrimidine nucleosides appear to be taken up from the media and the

nucleobases were likely generated by cells **(Figures 4 and S5)**. Mean uridine levels in cells

decreased slightly in cells after Mn depletion, whereas UMP (**Figure 4**) and uracil (**Figures 4 and**

S5h) levels dropped significantly. Uracil levels in cells likely decreased due to lower UMP

production but oddly, mean levels of uracil increased in media after Mn depletion. Interestingly,

orotidine levels increased in cells (**Figure 4**), indicating a potential blockage in the conversion to

UMP, although the explanation for this remains elusive as no *pyrF* enzyme listed in BRENDA has

been shown to utilize a Mn cofactor.

Levels of thymine decreased in cells after Mn depletion (**Figures 4 and S5d & h**). which

corresponds to a decrease in expression of *pdp* (pyrimidine nucleoside phosphorylase; SSA_1035;

thymidine to thymine conversion) (Puccio *et al.*, 2020). Oddly, the thymidine levels decreased as

well, although this may be explained by the fact that dTDP-rhamnose levels increased at T_{50} (**Table**

S10), indicating that thymidine may have been shuttled to sugar metabolism after Mn depletion.

Mean cytosine and cytidine levels increased slightly in cells after Mn depletion (**Figures 4 and S5f**

& h), which is the opposite trend from the other pyrimidines. Levels of downstream products 3'-

CMP and 2', 3'-cyclic CMP levels increased as well (**Figure 4**). The discrepancy may be explained

by decreased conversion to uridine as its levels dropped after Mn depletion (**Figures 4 and S5f**).

This is supported by a decrease in expression of *cdd* (cytidine deaminase; SSA_1037; cytidine to

uridine conversion) after Mn depletion (Puccio *et al.*, 2020) and Cdd may be Mn-cofactored (Hosono

and Kuno, 1973).

4.4 Oxidized and reduced glutathione levels in Mn-depleted *S. sanguinis* cells

Glutathione (γ-glutamyl-cysteinylglycine) is a nonprotein thiol produced by cells to prevent damage

caused by reactive oxygen species (ROS) (Carmel-Harel and Storz, 2000; Sies, 1999). The SK36

genome (Xu *et al.*, 2007) encodes a bifunctional γ-glutamate-cysteine ligase/glutathione synthetase (GshF; SSA_2168) (Janowiak and Griffith, 2005). Mean levels of the glutathione precursors cysteine, glutamine, and γ-glutamylcysteine all decreased slightly in cells after Mn depletion, consistent with active synthesis, although glycine levels did not change **(Figure 5a).** Interestingly, levels of reduced glutathione (GSH) increased in cells after Mn depletion, whereas levels of the oxidized form (GSSG) remained constant **(Figure 5b)**. Since the air flow was kept constant throughout the experiment, we expected that GSH would have been utilized by redox enzymes for ROS remediation. While ROS levels were not measured directly by the metabolomics analysis, levels of *ortho*-tyrosine increased (**Figure 5c**), which is an indicator of high ROS states (Ipson and Fisher, 2016; Matayatsuk *et al.*, 2007). Thus, the accumulation of GSH is probably due to Mn depletion, either because of a blockage of GSH utilization by redox enzymes or due to a reduction of ROS.

Due to the presumed decrease in activity of the Mn-cofactored superoxide dismutase, SodA (Crump *et al.*, 2014), it is unlikely that all ROS levels would have decreased after Mn depletion. The notable exception is H_2O_2, which was found to decrease after Mn depletion due to reduced expression of *spxB* (Puccio *et al.*, 2020). This likely led to a decrease in the direct detoxification of H_2O_2 by GSH, although the extent to which this occurs in cells is controversial (Berndt *et al.*, 2014). Additionally, *S. sanguinis* does not encode any known glutaredoxins and the only enzyme thought to utilize GSH in *S. sanguinis* is glutathione peroxidase (GpoA*;* SSA_1523), which uses GSH to detoxify H_2O_2 (**Figure 5d**) (Carmel-Harel and Storz, 2000). This enzyme has been found to contribute to oxidative stress tolerance in *S. pneumoniae* (Potter *et al.*, 2012) and virulence in *S. pyogenes* (Brenot *et al.*, 2004). Additionally, the enzyme that converts GSSG to GSH, glutathione reductase (Gor; SSA_1533), is likely metal-cofactored, which could explain why GSSG levels remained constant instead of decreasing as GSH levels increased. Thus, Mn depletion could explain the accumulation of both reduced and oxidized glutathione.

5.   Conclusions

In this study, we showed system-wide metabolomic changes induced in *S. sanguinis* Mn-transporter mutant cells and spent media in response to EDTA treatment over time. This study captured the Mn-responsive metabolic processes, such as dysregulations in carbohydrate, nucleotide, and redox

metabolism, many of which may contribute to the reduction in bacterial growth rate and virulence.
The decrease in available Mn led to the accumulation of fructose-1,6-bisphosphate, which likely
resulted in induction of carbon catabolite repression. This has widespread consequences, such as the
blockage of nucleobases conversion into nucleosides and accumulation of reduced glutathione. In
addition, we provide insights into the metabolic composition of BHI and the components streptococci
may utilize from this undefined medium.

6. Declarations:

Funding
This work was supported by the National Institute of Allergy and Infectious Diseases of the National
Institutes of Health under award no. R01 AI114926 to TK. TP was supported by a predoctoral
fellowship from the National Institute of Dental and Craniofacial Research of the National Institutes
of Health under award no. F31 DE028468. The content is solely the responsibility of the authors and
does not necessarily represent the official views of the National Institutes of Health.

Conflicts of interest
TP and TK do not have any conflicts of interest. BBM currently works as a Computational Biologist
with Enveda Therapeutics; however, he has no conflict of interest with this study.

Ethics approval
This article does not contain any studies with human participants or animals performed by any of the
authors.

Consent to participate
Not applicable

Consent for publication
All authors have read, approved and have provided consent for this publication.

Availability of data and material
The datasets generated and analyzed during the current study are available as Supplementary Tables
S1 and S2 as provided by Metabolon, Inc.

Code availability
Not applicable

Author's contributions

TP and TK designed the experiments. TP performed the experiments. BBM performed the data analysis. All authors analyzed the results and wrote the manuscript.

Acknowledgements

We thank Karina Kunka, Dr. Shannon Green, Dr. Seon-Sook An, and Brittany Spivey for discussions and assistance with experiments. We also thank Dr. Danny Alexander (Metabolon, Inc.) for his initial analysis of the data.

7. References

Ajdic, D. and Pham, V.T. (2007) Global transcriptional analysis of *Streptococcus mutans* sugar transporters using microarrays. *J Bacteriol* **189,** 5049-59.

Bai, Y., Shang, M., Xu, M., Wu, A., Sun, L. and Zheng, L. (2019) Transcriptome, phenotypic, and virulence analysis of *Streptococcus sanguinis* SK36 wild type and its CcpA-null derivative (ΔCcpA). *Front Cell Infect Microbiol* **9,** 411.

Baker, S.P., Nulton, T.J. and Kitten, T. (2019) Genomic, phenotypic, and virulence analysis of *Streptococcus sanguinis* oral and infective-endocarditis isolates. *Infect Immun* **87,** e00703-00718.

Berndt, C., Lillig, C.H. and Flohe, L. (2014) Redox regulation by glutathione needs enzymes. *Front Pharmacol* **5,** 168.

Bor, D.H., Woolhandler, S., Nardin, R., Brusch, J. and Himmelstein, D.U. (2013) Infective endocarditis in the U.S., 1998-2009: a nationwide study. *PLoS One* **8,** e60033.

Brenot, A., King, K.Y., Janowiak, B., Griffith, O. and Caparon, M.G. (2004) Contribution of glutathione peroxidase to the virulence of *Streptococcus pyogenes*. *Infect Immun* **72,** 408-13.

Cahill, T.J., Baddour, L.M., Habib, G., Hoen, B., Salaun, E., Pettersson, G.B., Schafers, H.J. and Prendergast, B.D. (2017) Challenges in infective endocarditis. *J Am Coll Cardiol* **69,** 325-344.

Carmel-Harel, O. and Storz, G. (2000) Roles of the glutathione- and thioredoxin-dependent reduction systems in the *Escherichia coli* and *Saccharomyces cerevisiae* responses to oxidative stress. *Annual Review of Microbiology* **54,** 439–61.

Colomer-Winter, C., Flores-Mireles, A.L., Baker, S.P., Frank, K.L., Lynch, A.J.L., Hultgren, S.J.,
Kitten, T. and Lemos, J.A. (2018) Manganese acquisition is essential for virulence of *Enterococcus*
faecalis. *PLoS Pathog* **14,** e1007102.

Crump, K.E., Bainbridge, B., Brusko, S., Turner, L.S., Ge, X., Stone, V., Xu, P. and Kitten, T.
(2014) The relationship of the lipoprotein SsaB, manganese and superoxide dismutase in
Streptococcus sanguinis virulence for endocarditis. *Mol Microbiol* **92,** 1243-59.

Dehaven, C.D., Evans, A.M., Dai, H. and Lawton, K.A. (2010) Organization of GC/MS and LC/MS
metabolomics data into chemical libraries. *J Cheminform* **2,** 9.

Dintilhac, A., Alloing, G., Granadel, C. and Claverys, J.-P. (1997) Competence and virulence of
Streptococcus pneumoniae: Adc and PsaA mutants exhibit a requirement for Zn and Mn resulting
from inactivation of putative ABC metal permeases. *Molecular Microbiology* **25,** 727-739.

Dodds, D.R. (2017) Antibiotic resistance: A current epilogue. *Biochem Pharmacol* **134,** 139-146.

Evans, A.M., DeHaven, C.D., Barrett, T., Mitchell, M. and Milgram, E. (2009) Integrated,
nontargeted ultrahigh performance liquid chromatography/electrospray ionization tandem mass
spectrometry platform for the identification and relative quantification of the small-molecule
complement of biological systems. *Analytical Chemistry (Washington)* **81,** 6656–6667.

Fei, F., Mendonca, M.L., McCarry, B.E., Bowdish, D.M.E. and Surette, M.G. (2016) Metabolic and
transcriptomic profiling of *Streptococcus intermedius* during aerobic and anaerobic growth.
Metabolomics **12,** 46.

Grapov, D. (2014) DeviumWeb: Dynamic multivariate data analysis and visualization platform.,
DOI: 10.5281/zenodo.592909.

Guo, R. and Lu, H. (2020) Targeted metabolomics revealed the regulatory role of manganese on
small-molecule metabolism of biofilm formation in *Escherichia coli*. *J Anal Test* **4,** 226-237.

Hosono, H. and Kuno, S. (1973) The purification and properties of cytidine deaminase from
Escherichia coli. *J Biochem* **74,** 797-803.

Ipson, B.R. and Fisher, A.L. (2016) Roles of the tyrosine isomers meta-tyrosine and ortho-tyrosine in
oxidative stress. *Ageing Res Rev* **27,** 93-107.

Jakubovics, N.S., Yassin, S.A. and Rickard, A.H. (2014) Community interactions of oral
streptococci. *Adv Appl Microbiol* **87,** 43-110.

Janowiak, B.E. and Griffith, O.W. (2005) Glutathione synthesis in *Streptococcus agalactiae*. *J Biol*
Chem **280,** 11829-39.

Jeske, L., Placzek, S., Schomburg, I., Chang, A. and Schomburg, D. (2019) BRENDA in 2019: a
European ELIXIR core data resource. *Nucleic Acids Res* **47,** D542-D549.

Jewison, T., Su, Y., Disfany, F.M., Liang, Y., Knox, C., Maciejewski, A., Poelzer, J., Huynh, J.,
Zhou, Y., Arndt, D., Djoumbou, Y., Liu, Y., Deng, L., Guo, A.C., Han, B., Pon, A., Wilson, M.,
Rafatnia, S., Liu, P. and Wishart, D.S. (2014) SMPDB 2.0: big improvements to the Small Molecule
Pathway Database. *Nucleic Acids Res* **42,** D478-84.

Kanehisa, M. and Goto, S. (2000) KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic*
Acids Research **28,** 27-30.

Kehl-Fie, T.E., Zhang, Y., Moore, J.L., Farrand, A.J., Hood, M.I., Rathi, S., Chazin, W.J., Caprioli,
R.M. and Skaar, E.P. (2013) MntABC and MntH contribute to systemic *Staphylococcus aureus*
infection by competing with calprotectin for nutrient manganese. *Infect Immun* **81,** 3395-405.

Kilian, M., Mikkelsen, L. and Henrichsen, J. (1989) Taxonomic study of viridans streptococci:
Description of *Streptococcus gordonii* sp. nov. and emended descriptions of *Streptococcus sanguis*
(White and Niven 1946), *Streptococcus oralis* (Bridge and Sneath 1982), and *Streptococcus mitis*
(Andrewes and Horder 1906). *Int J of Syst Bacteriol* **39**.

Kreth, J., Giacaman, R.A., Raghavan, R. and Merritt, J. (2017) The road less traveled - defining
molecular commensalism with *Streptococcus sanguinis*. *Molecular Oral Microbiology* **32,** 181-196.

Kreth, J., Merritt, J., Shi, W. and Qi, F. (2005) Competition and coexistence between *Streptococcus*
mutans and *Streptococcus sanguinis* in the dental biofilm. *J Bacteriol* **187,** 7193-203.

Leonard, A., Gierok, P., Methling, K., Gomez-Mejia, A., Hammerschmidt, S. and Lalk, M. (2018)
Metabolic inventory of *Streptococcus pneumoniae* growing in a chemical defined environment. *Int J*
Med Microbiol **308,** 705-712.

Liu, G., Qiao, Y., Zhang, Y., Leng, C., Chen, H., Sun, J., Fan, X., Li, A. and Feng, Z. (2020)
Metabolic profiles of carbohydrates in *Streptococcus thermophilus* during pH-controlled batch
fermentation. *Frontiers in Microbiology* **11,** 1131.

Makhlynets, O., Boal, A.K., Rhodes, D.V., Kitten, T., Rosenzweig, A.C. and Stubbe, J. (2014)
Streptococcus sanguinis class Ib ribonucleotide reductase: high activity with both iron and
manganese cofactors and structural insights. *J Biol Chem* **289,** 6259-72.

500
501 Martinussen, J., Sorensen, C., Jendresen, C.B. and Kilstrup, M. (2010) Two nucleoside transporters
502 in *Lactococcus lactis* with different substrate specificities. *Microbiology* **156,** 3148-3157.

503
504 Matayatsuk, C., Poljak, A., Bustamante, S., Smythe, G.A., Kalpravidh, R.W., Sirankapracha, P.,
505 Fucharoen, S. and Wilairat, P. (2007) Quantitative determination of ortho- and meta-tyrosine as
506 biomarkers of protein oxidative damage in beta-thalassemia. *Redox Rep* **12,** 219-28.

507
508 Misra, B.B. and Olivier, M. (2020) High resolution GC-Orbitrap-MS metabolomics using both
509 electron ionization and chemical ionization for analysis of human plasma. *J Proteome Res*.

510
511 Moreillon, P., Que, Y.A. and Bayer, A.S. (2002) Pathogenesis of streptococcal and staphylococcal
512 endocarditis. *Infect Dis Clin North Am* **16,** 297-318.

513
514 Murgas, C.J., Green, S.P., Forney, A.K., Korba, R.M., An, S.S., Kitten, T. and Lucas, H.R. (2020)
515 Intracellular metal speciation in *Streptococcus sanguinis* establishes SsaACB as critical for redox
516 maintenance. *ACS Infect Dis* **6,** 1906-21.

517
518 Ogunniyi, A.D., Mahdi, L.K., Jennings, M.P., McEwan, A.G., McDevitt, C.A., Van der Hoek, M.B.,
519 Bagley, C.J., Hoffmann, P., Gould, K.A. and Paton, J.C. (2010) Central role of manganese in
520 regulation of stress responses, physiology, and metabolism in *Streptococcus pneumoniae*. *J Bacteriol*
521 **192,** 4489-97.

522
523 Potter, A.J., Trappetti, C. and Paton, J.C. (2012) *Streptococcus pneumoniae* uses glutathione to
524 defend against oxidative stress and metal ion toxicity. *J Bacteriol* **194,** 6248-54.

525
526 Puccio, T. and Kitten, T. (2020) Fermentor Growth of *Streptococcus sanguinis*. *protocols.io*.

527
528 Puccio, T., Kunka, K.S., Zhu, B., Xu, P. and Kitten, T. (2020) Manganese depletion leads to
529 multisystem changes in the transcriptome of the opportunistic pathogen *Streptococcus sanguinis*.
530 *bioRxiv*.

531
532 Qiao, Y., Liu, G., Lv, X., Fan, X., Zhang, Y., Meng, L., Ai, M. and Feng, Z. (2019) Metabolic
533 pathway profiling in intracellular and extracellular environments of *Streptococcus thermophilus*
534 during pH-controlled batch fermentations. *Front Microbiol* **10,** 3144.

535
Quan, T.P., Muller-Pebody, B., Fawcett, N., Young, B.C., Minaji, M., Sandoe, J., Hopkins, S.,
Crook, D., Peto, T., Johnson, A.P. and Walker, A.S. (2020) Investigation of the impact of the NICE
guidelines regarding antibiotic prophylaxis during invasive dental procedures on the incidence of
infective endocarditis in England: an electronic health records study. *BMC Med* **18,** 84.

Radin, J.N., Kelliher, J.L., Parraga Solorzano, P.K. and Kehl-Fie, T.E. (2016) The two-component
system ArlRS and alterations in metabolism enable *Staphylococcus aureus* to resist calprotectin-
induced manganese starvation. *PLoS Pathog* **12,** e1006040.

Redanz, S., Treerat, P., Mu, R., Redanz, U., Zou, Z., Koley, D., Merritt, J. and Kreth, J. (2020)
Pyruvate secretion by oral streptococci modulates hydrogen peroxide dependent antagonism. *ISME J*
14, 1074-1088.

Rhodes, D.V., Crump, K.E., Makhlynets, O., Snyder, M., Ge, X., Xu, P., Stubbe, J. and Kitten, T.
(2014) Genetic characterization and role in virulence of the ribonucleotide reductases of
Streptococcus sanguinis. *J Biol Chem* **289,** 6273-87.

Sies, H. (1999) Glutathione and its role in cellular functions. *Free Radical Biology & Medicine* **27,**
916-921.

Team, R.C. (2018) R: A language and environment for statistical computing.

Thornhill, M.H., Gibson, T.B., Cutler, E., Dayer, M.J., Chu, V.H., Lockhart, P.B., O'Gara, P.T. and
Baddour, L.M. (2018) Antibiotic prophylaxis and incidence of endocarditis before and after the 2007
AHA recommendations. *J Am Coll Cardiol* **72,** 2443-2454.

Warner, J.B. and Lolkema, J.S. (2003) CcpA-dependent carbon catabolite repression in bacteria.
Microbiology and Molecular Biology Reviews **67,** 475-490.

Webb, A.J. and Hosie, A.H. (2006) A member of the second carbohydrate uptake subfamily of ATP-
binding cassette transporters is responsible for ribonucleoside uptake in *Streptococcus mutans*. *J*
Bacteriol **188,** 8005-12.

Widmer, E., Que, Y.-A., Entenza, J.M. and Moreillon, P. (2006) New concepts in the
pathophysiology of infective endocarditis. *Current Infectious Disease Reports* **8,** 271–279.

Wilson, W., Taubert, K.A., Gewitz, M., Lockhart, P.B., Baddour, L.M., Levison, M., Bolger, A.,
Cabell, C.H., Takahashi, M., Baltimore, R.S., Newburger, J.W., Strom, B.L., Tani, L.Y., Gerber, M.,
Bonow, R.O., Pallasch, T., Shulman, S.T., Rowley, A.H., Burns, J.C., Ferrieri, P., Gardner, T., Goff,
D. and Durack, D.T. (2007) Prevention of infective endocarditis: Guidelines from the American
Heart Association. *Circulation* **116,** 1736-54.

Xia, J., Sinelnikov, I.V., Han, B. and Wishart, D.S. (2015) MetaboAnalyst 3.0--making
metabolomics more meaningful. *Nucleic Acids Res* **43,** W251-7.

580
581 Xu, P., Alves, J.M., Kitten, T., Brown, A., Chen, Z., Ozaki, L.S., Manque, P., Ge, X., Serrano, M.G.,
582 Puiu, D., Hendricks, S., Wang, Y., Chaplin, M.D., Akan, D., Paik, S., Peterson, D.L., Macrina, F.L.
583 and Buck, G.A. (2007) Genome of the opportunistic pathogen *Streptococcus sanguinis*. *J Bacteriol*
584 **189,** 3166-75.

Table 1. Top 10 significantly differential (ANOVA) metabolites in cells and media.

Metabolites	P-value	FDR	Fisher's LSD
Cells			
N2-methylguanosine	7.23E-15	3.80E-12	T_{25} v T_{50}; T_{-20} v T_{50}
pseudouridine	1.72E-14	4.51E-12	T_{25} v T_{50}; T_{-20} v T_{25}; T_{-20} v T_{50}
N-acetylglucosamine 6-phosphate	7.09E-09	1.24E-06	T_{-20} v T_{25}; T_{-20} v T_{50}
N-acetylmuramyl-alanyl-isoglutamine	1.23E-08	1.36E-06	T_{25} v T_{50}; T_{-20} v T_{25}; T_{-20} v T_{50}
1-stearoyl-GPA (18:0)	1.29E-08	1.36E-06	T_{50} v T_{25}; T_{25} v T_{-20}; T_{50} v T_{-20}
2'-O-methyluridine	1.97E-08	1.73E-06	T_{25} v T_{50}; T_{-20} v T_{50}
gamma-glutamylglutamate	3.01E-08	2.25E-06	T_{25} v T_{50}; T_{-20} v T_{25}; T_{-20} v T_{50}
orotidine	4.93E-08	3.23E-06	T_{50} v T_{25}; T_{50} v T_{-20}
eicosenoate (20:1n9 or 1n11)	5.73E-08	3.25E-06	T_{50} v T_{25}; T_{50} v T_{-20}
1-stearoyl-GPG (18:0)	6.19E-08	3.25E-06	T_{50} v T_{25}; T_{25} v T_{-20}; T_{50} v T_{-20}
Media			
inosine	1.58E-07	6.53E-05	T_{50} v T_{25}; T_{-20} v T_{25}; T_{50} v T_{-20}
2'-deoxyadenosine	1.25E-06	0.00026	T_{50} v T_{25}; T_{50} v T_{-20}
N6-methyladenosine	2.16E-06	0.000299	T_{50} v T_{25}; T_{25} v T_{-20}; T_{50} v T_{-20}
pyruvate	3.18E-05	0.003148	T_{50} v T_{25}; T_{25} v T_{-20}; T_{50} v T_{-20}
thymidine	3.80E-05	0.003148	T_{50} v T_{25}; T_{50} v T_{-20}
nicotinamide	0.000186	0.012825	T_{50} v T_{25}; T_{25} v T_{-20}; T_{50} v T_{-20}
adenine	0.000243	0.014401	T_{25} v T_{50}; T_{-20} v T_{50}
adenosine	0.000328	0.016368	T_{50} v T_{25}; T_{-20} v T_{25}; T_{50} v T_{-20}
2,3-dihydroxyisovalerate	0.000356	0.016368	T_{25} v T_{-20}; T_{50} v T_{-20}
ophthalmate	0.000459	0.019002	T_{50} v T_{25}; T_{25} v T_{-20}; T_{50} v T_{-20}

Fig. 1 Schematic diagram displaying the experimental design, platform and software tools used for the analysis of metabolomic changes in cells and media subjected to EDTA treatment.

Fermentor sample collection, metal, and hydrogen peroxide analysis charts were adapted from
Puccio, et al. (2020). Extraction, derivatization, and annotation were completed by Metabolon, Inc.
ICP-OES, inductively coupled plasma optical emission spectroscopy; UPLC-MS/MS, ultra
performance liquid chromatography with tandem mass spectrometry.

Fig. 2 Multivariate, VIP, and time course analysis of the metabolomic changes in cells and
media.

Score plots of PCA displaying the separation of time-points in cells (a) and spent media (b). Cell
samples n = 6; media samples n = 3. Top 15 metabolites (variables) based on VIP scores from PLS-
DA analysis of cells (c) and spent media (d).

Fig. 3 Relative abundance of carbohydrates and glycolytic intermediates in media and cells.

Levels of sugars in media (a) and cells (c) are depicted. Products of glycolysis in media (b) and cells
(d). Whiskers indicate the range; horizontal bars represent the mean. A two-tailed t-test was used to
compare the pre-inoculum (Pre-Inoc) media samples to post-inoculum (T_{-20}). Red asterisks indicate
P-value < 0.05. Spent media and cell metabolite levels were compared using one-way ANOVA with
a Fisher's least significant difference test to compare the post-EDTA samples to pre-EDTA. Black
asterisks indicate P-value < 0.05.

Fig. 4 Quantitative changes in nucleotide metabolism for cells and media after Mn depletion.

The direction of change in metabolite concentration is depicted in shades of red or blue, for
increasing or decreasing concentration, respectively. Significance was determined by a t-test using
the comparisons shown in the key. Metabolites that do not have a set of boxes were not detected in
any sample. Diamonds indicate nucleobases and stars indicate nucleosides. Figure was made using
Biorender.com.

Fig. 5 Glutathione abundance in cells and model of Mn depletion.

Levels of glutathione precursors (a), glutathione (b), and the oxidative stress indicator *ortho*-tyrosine
(c) are shown. Metabolite levels were compared using one-way ANOVA with a Fisher's least
significant difference test to compare the post-EDTA samples to pre-EDTA. Black asterisks indicate
P-value < 0.05. (d) Model of glutathione utilization by glutathione peroxidase (GpoA) and reduction

by glutathione reductase (Gor) under normal and low-Mn conditions. Figure (d) was made using
Biorender.com.

Supplementary Table Captions

Supplementary Table S1. Raw metabolite abundance data for cellular metabolites captured using
combined LC-MS/MS (positive and negative modes) analysis by Metabolon, Inc. Retention indices
(RIs), quantifier mass, CAS IDs, KEGG IDs, HMDB IDs, PubChem IDs, SMILES, Super Pathway
and Sub Pathway information, biochemical names for the metabolites, and their raw abundances are
also included.

Supplementary Table S2. Raw metabolite abundance data for media metabolites captured using
combined LC-MS/MS (positive and negative mode) analysis by Metabolon, Inc. Retention indices
(RIs), quantifier mass, CAS IDs, KEGG IDs, HMDB IDs, PubChem IDs, SMILES, Super Pathway
and Sub Pathway information, biochemical names for the metabolites, and their raw abundances are
also included.

Supplementary Table S3. Transformed, scaled and normalized metabolite abundance data for
cellular metabolites.

Supplementary Table S4. Transformed, scaled and normalized metabolite abundance data for
media metabolites.

Supplementary Table S5. Pathway enrichment analysis for the 534 quantified cellular metabolites.

Supplementary Table S6. Pathway enrichment analysis for the 424 quantified metabolites in media.

Supplementary Table S7. Unique metabolites found in some but not all samples.

Supplementary Table S8. One-way ANOVA statistical analysis results for metabolites of cells.
Fold changes cut off > 1.2 and < 0.8; P-value < 0.05.

Supplementary Table S9. One-way ANOVA statistical analysis results for metabolites of media.
Fold changes cut off > 1.2 and < 0.8; P-value < 0.05).

Supplementary Table S10. Pathway enrichment analysis for significantly differential (ANOVA)
cellular metabolites.

Supplementary Table S11. Pathway enrichment analysis for significantly differential (ANOVA)
media metabolites.

Supplementary Table S12. Average cell metabolite levels, fold changes, and P-values as
determined by t-tests comparing post-EDTA samples to the pre-EDTA sample.

Supplementary Table S13. Average media metabolite levels, fold changes, and P-values as
determined by t-tests comparing spent media (T_{-20}) to pre-inoculation media as well as post-EDTA
samples to the pre-EDTA sample.

Supplementary Table S14. STEM analysis of cellular metabolites displaying top 2 significant
profiles.

Supplementary Table S15. STEM analysis of media metabolites displaying top 3 significant
profiles.

Sample Collection

ΔssaACB

T₋₂₀, T₀, T₂₅, T₅₀

EDTA

Metal Analysis (ICP-OES)

Mn | Fe | Zn

μg Metal/mg Protein

EDTA

**

***

EDTA

Time (min)

-20 25 50

Hydrogen Peroxide Analysis

[H₂O₂] μM

EDTA

**

***

Time (min)

-20 25 50

Extraction & Derivatization
UPLC-MS/MS

Annotation
LIPID MAPS
Chemical Abstracts Service
The Human Metabolome Database
Kyoto Encyclopedia of Genes and Genome

Analysis
DeviumWeb
MetaboAnalyst
R