Confinement Discerns Swarmers from Planktonic Bacteria

Weijie Chen[1,2], Neha Mani[1,†], Hamid Karani[1], Hao Li[2], Sridhar Mani[2,*], Jay X. Tang[1,*]

[1] Department of Physics, Brown University, 182 Hope St., Providence, RI 02912, USA

[2] Department of Medicine, Albert Einstein College of Medicine, 1300 Morris Park Ave, Bronx, NY10461, USA

* Corresponding Authors

Email: sridhar.mani@einsteinmed.org; jay_tang@brown.edu

[†] Present address: Hunter College High School, 71 E 94th St, New York, NY 10128, USA

Author Contribution

W.C., N.M., J.X.T., and S.M. conceived and designed the work. W.C. and N.M. performed the experiments and analyzed the data. H.K. performed the computational simulations. H.L. isolated the bacteria strains and prepared the mouse tissues. W.C., H.K., S.M., and J.X.T. wrote the paper.

Conflict of interest

Weijie Chen, Neha Mani, Jay X. Tang and Sridhar Mani filed a U.S. patent application (Application No. 63033369). Other authors declare no conflict of interest.

Acknowledgment

We thank Daniel B. Kearns from Indiana University at Bloomington for providing the *Bacillus subtilis* 3610 bacteria strain, Cori Bargmann at Rockefeller University for gifting us the bacteria strain *Serratia marcescens* Db10. We thank Hui Ma for his assistance in the clean room and discussion of the work. NIH Grant 1R01CA222469-01 supported this work.

Abstract

Powered by flagella, many species of bacteria exhibit collective motion on a solid surface commonly known as swarming. Physical changes like cell elongation and hyper flagellation have been shown to accompany swarming phenotype. Less noticeable, however, are the contrasts of collective motion between the swarming cells and the planktonic cells of comparable cell density. Here, we show that when confined by microwells of specific sizes mounted on a soft agar surface, novel bacteria *Enterobacter* sp. SM3 under swarming condition exhibited "single-swirl" motion pattern distinct from "multi-swirl" motion pattern formed by its concentrated planktonic counterpart. We hypothesize that "rafting behavior" of the swarming bacteria upon dilution might account for the motion pattern difference. This was further validated through computational simulation where swarming cells are modeled by lower repulsion and stronger alignment among them than planktonic cells. Our new technical approach also enabled us to observe swarming on a non-agar tissue surface.

Introduction

Motility is an essential characteristic for bacteria. Although energy-consuming, it provides high returns, enabling cells to uptake nutrients efficiently and escape from noxious environments[1]. In a host environment, bacterial motility is also an essential phenotype that intimately relates to virulence through complex regulatory networks[2]. Swimming and swarming are two common motility phenotypes mediated by flagella. Whereas planktonic phenotype defines the motility of individual bacteria, a collective movement powered by rotating flagella[3] on a partially solidified surface defines swarming[4]. In swarming, bacteria utilize their flagella to navigate, two-dimensionally, through a medium and acquire necessary molecules for the maintenance of homeostasis and overall survival[5]. Morphological changes like cell elongation may or may not occur in all swarming bacteria[6]. Thus, concentrated swimming bacteria are often called "a swarm of bacteria" without requiring precise identification of swarming motility, per se. Nevertheless, microbiologists believe that swarming and swimming are fundamentally different motility types. For instance, studies found that compared with swimming cells, the requirement for flagella torque is higher for swarming *B. subtilis*[7]; swarming *E. coli* remodel their chemotaxis pathway[8]; and in swarming *P. aeruginosa*, the production of virulence factors and antibiotic resistance increases[9]. A recent study has demonstrated a medically relevant distinction between swarming and swimming: a particular strain of swarming endobacteria protect against mice intestinal inflammation while their swimming counterparts could not[10]. The evidence to date that shows swarming is different from swimming comes mostly from biological data. However, precise biophysical visualization and quantitation of these differences are lacking. In this report, using *Enterobacter* sp. SM3, a novel bacteria strain that possesses both swimming and swarming motilities, we show distinct biophysical characteristics between these two types of motility under confined, circular geometry of a particular range of sizes.

Studies have shown that geometric constraints have profound influence on patterns of microswimmers' collective motion. For example, these constraints may create mesoscopic or macroscopic coherent structures such as swirls and jets[11-13]. Circular confinement, in particular, could stabilize a suspension of motile bacteria into a spiral vortex[14-16]. Here, we compare the behaviors of bacteria in swarming and planktonic states under quasi-2D circular confinement. Many species of bacteria show distinctive motion patterns while confined. This characteristic may

lead to future diagnostic applications since there are growing associations between bacterial
swarming and virulence pathologies.

Results

Swarming *Enterobacter* Sp. SM3 forms large single swirls up to 100 μm diameter

A novel bacterial strain *Enterobacter* Sp. SM3 (NCBI BioProject PRJNA558971), isolated in 2014
from DSS induced clitic mice, has been previously studied for motility[17] and infectivity[10]. SM3 is
a strong swarmer, expanding rapidly on 0.5% agar with collective motion of multilayers of cells
at the edge. We mounted a PDMS chip containing circular microwells on the agar so that bacteria
in confinement could rotate for more than 3 hours (details with illustration in Methods). Under
confinement in circular wells in the diameter range of 31-90 μm, swarming SM3 shows single
swirls. In contrast, SM3 planktonic cells concentrated from the liquid medium form mesoscale
vortices (multiple swirls) in the same size range, with the exception of the smallest well diameter
of 31 μm. A clear difference is shown at the well diameter of 74 μm (Fig. 1A-D, Movie S1 & S2).
This striking difference persists in several well depths, except that the concentrated cells yields
small but non-zero vortex order parameter (VOP, defined as illustrated in Fig. 1E) in deeper wells,
as opposed to nearly zero VOP in shallow wells (Fig. 1F).

The confinement well diameter has strong influence on the motion pattern in the wells. In smaller
wells like 31 μm, even concentrated planktonic SM3 forms a single vortex (Fig. 2A) whereas in
larger wells, such as ones of 112 μm diameter, swarming SM3 also breaks into mesoscale vortices
(Fig. 2B). The phase diagram shows a single swirl in small confinement for both types. The
patterns diverge as confinement size increases, but they converge towards multiple swirls as
confinement size reaches 144 μm and larger (Fig. 2C). To further compare the dynamics of the
confined swarming and swimming SM3, spatial correlation of the velocity field was calculated for
$d = 90$ μm where the motion patterns differ for swarming and swimming SM3 and for $d = 500$ μm
where both motilities show mesoscale vortices (see method). We computed the correlation
function for the inscribed square within a well, which shows the extent to which the velocity at an
arbitrary location correlated with the velocity at a distance of Δr away from that location. In 90
μm wells, swarming SM3 velocity correlates positively or negatively throughout the whole well
(negative values have resulted from the opposite sides of a single swirl). In contrast, swimming

velocity of planktonic cells of comparable concentration does not correlate anymore as Δr went up

beyond 25 μm (Fig. 2D). However, in a large open space where both swarming and swimming

SM3 break into small vortices, the correlation functions look similar. The characteristic length as

the curve first crosses $C_r(\Delta r) = 0$, which also represents the size of the mesoscale vortices, of

planktonic SM3 is 27 μm and 33 μm for swarming SM3 (Fig. 2E).

We also tested other bacteria such as *Enterobacter* Sp. SM1, *Serratia marcescens* (including one

lab strain Db10 and another strain H3 isolated from a human patient), *Citrobacter koseri* (H6), and

Bacillus subtilis 3610. All the tested strains showed similar motion pattern divergence between

confined planktonic cells and swarming cells like SM3 with the exception of *B. subtilis* (Fig. S1A,

see discussion).

The large single swirl behavior is indicative of cohesive cell-cell interaction

We performed several experiments to explore the cause for the divergence of motion patterns in

confinement. First, we rule out cell density difference as the reason for the difference in the

confined motion patterns by concentrating planktonic cells to comparable density to that of a

naturally expanding swarm on agar (see methods) before mounting the PDMS chip. Second, we

noticed that SM3 tends to get elongated when they swarm. We hypothesize that elongated bacteria

may enhance the local alignment of the rod-shaped cells and increase the size of vortices in

mesoscale turbulence[18]. Thus, we treated SM3 planktonic cells with Cephalexin (CEP) which has

been shown to elongate *E. coli* [19] . This treatment indeed caused the cell length to reach that of

swarming cells on average (Fig. 3A). However, we found no significant change following

centrifugation and CEP treatment of the planktonic SM3 (Fig. 3B). Although CEP treated

planktonic SM3 has similar cell length, cell density, and cell speed as that of swarming SM3, we

could not restore the single swirl pattern in 74 μm confinement wells (Fig. 3C). Third, noticing a

surfactant rim on the swarming SM3 colony edge, we conjectured that surfactants secreted by

swarming SM3 might help align the swarmers in confinement. Surfactin was added in several

concentrations to planktonic SM3 in order to test whether it could promote a single swirl pattern.

However, it did not establish a stable single swirl. Last but not least, we found that adding

lyophilized swarming supernatant to swimming SM3 also did not help increase the VOP (Fig. 3C).

Unable to make the concentrated planktonic SM3 form a single swirl in the 74 μm well, we tackled

the problem from another angle, by altering the conditions of swarming SM3 in order to break the

single swirls. Initially, we tried to physically "damage" the swarming colony by rubbing the

swarming colony gently with a piece of PDMS offcut. This operation did not break the single swirl

pattern in the wells (Fig. 3D). Then, 0.2% D-mannose was added to the swarming colony to de-

cluster bacteria bundles due to cells' sticking to each other. However, this treatment could not alter

the single swirl pattern, either (Fig. 3D). Finally, we diluted the swarming cells in LB by 20-fold.

After re-concentrating the cells by centrifugation and removing extra LB to recover the initial cell

density, these "swarming" SM3 cells were pipetted back on the agar plate. After this treatment,

the previous single swirl turned to multiple swirls under the confinement (Fig. 3D), suggesting that

these cells now behave much like planktonic cells. We conclude that the single swirl pattern

depends on cohesive cell-cell interaction mediated by biochemical factor/s removable by matrix

dilution.

Diluted swarming SM3 show unique dynamic clustering patterns

We suspected that specific interactions between the neighboring swarming cells were weakened

or even diminished upon dilution with the Luria broth (LB) medium. A fifty (50) μL water droplet

was applied to the swarming and the concentrated planktonic SM3 colony edges to investigate the

potential alignment among the cells at a microscopic scale within the bacterial colony. In the

diluted swarming colony, groups of cells formed bacterial rafts, a characteristic feature previously

associated with gliding motility[3,20]. Those cells within a polar cluster moving in the same direction

in a cohesive pack at the same speed (Movie S3). In contrast, upon dilution of the concentrated

planktonic SM3, the cells disperse uniformly, and their moving directions appear random (Movie

S4). Swarming SM3 cells tend to move together near the agar surface, while planktonic SM3 cells

swim freely in the bulk fluid (Fig. 4A-B). We used the MATLAB PIV toolkit to track the moving

bacteria in the image sequences of diluted swarming and planktonic SM3 for comparison. We

found that swarming SM3 formed clusters with more than 20 cells on average, while we did not

see such clusters of planktonic SM3 cells (Fig. 4C-D). The lingering clusters of cells in the

swarming phase upon dilution point to stronger cell-cell cohesive interaction than between

planktonic cells.

Numerical simulation reveals cell-cell interaction to be the key player

To further verify that rafting in swarming is a crucially relevant factor to the motion pattern

To further verify that rafting in swarming is a crucially relevant factor to the motion pattern

discrepancy, we performed computer simulations using the Zonal Model where the interactions

among the moving particles (short-range repulsion, velocity alignment, and anti-alignment) are

considered, all as functions of the particle-particle distance[21,22]. The speed of the particles is fixed

for simplicity, but the initial particle positions and initial moving directions are randomized. In the

simulations, we interpret the rafting as a lower repulsion force and stronger alignment among the

swarmers (see methods and SI methods). We simulated the situation of confined swarmers and

planktonic cells in different sizes of circular confinement, as in the experiments. The simulation

results mirror the experiment results well: both swarmers and planktonic cells start with single

swirl pattern; as the circle size is increased, the planktonic cells break into multi-swirl motion

pattern earlier than the swarmers and finally both converge to multi-swirl region (Fig. 5A,

compared with Fig. 2C, also see Fig. S2 and Movie S7). We then performed the "dilution"

simulation for both states, finding that swarming cells form dynamic clusters when the cell density

is above $\rho = 235$ whereas the planktonic cells form a "gas" phase without clustering at all densities

(Fig. 5B, Movie S8). This result echoes the experimental results in Fig. 4A. Thus, by encoding

stronger cell-cell interaction among the swarming cells, we recovered the experimental results in

both confinement and dilution experiments.

Identifying SM3 motility type on mice mucosal surface

The difference in confined motion patterns enables us to detect bacterial swarming on surfaces

other than agar, including physiological environments such as on a mucosal surface. We are

unaware of any previous studies or examples regarding bacterial swarming on non-agar surfaces,

likely due to technical challenges in dealing with uneven or non-controlled surfaces. The mouse

intestinal tissue, for instance, is more than 1 mm thick and non-transparent. Since light cannot

penetrate the tissue, observing bacteria directly on the tissue surface is not feasible. Staining or

fluorescence labeling may alter the bacterial swarming motility (e.g. SM3 becomes non-swarming

once GFP labeled. Unpublished observation). If labeled biochemically, the fluorescence signal

weakens when the cells reproduce (e.g., there will be progressively less cell wall labeling of SM3

with Alexa Fluor 488 when cells divide). Using fluorescent beads coated PDMS chips mounted

on SM3 inoculated C57BL6 mouse intestine tissue, we were able to detect swarming motility

based on the swirling motion of the beads. This experiment on the mouse intestine tissue confirms

that bacterial swarming indeed occurs on a non-agar, physiologically relevant surface (Fig. S3;

Movie S5&6, also see SI method).

Discussion

Mounting PDMS chip on a soft agar plate, we have shown the motion pattern differences between

confined planktonic and swarming *Enterobacter* sp. SM3 in the size range of 40 μm ≤ d ≤ 90 μm.

Compared with previous work, our experimental setup has the advantage of ensuring stable and

sustainable patterns. First, PDMS material does not harm living bacteria cells and is permeable to

oxygen[23], which prevents suffocation. Second, we mounted the microchip on a soft agar containing

over 97% water, which, via permeability and capillary flow, automatically fills the wells. Finally,

the LB agar also provides the necessary nutrients to fuel the bacterial movement in the wells.

Therefore, bacterial cells confined in the microwells remain motile for hours, much longer than in

droplets surrounded by mineral oil[14,19] or in microfluidic chambers with glass surfaces[11,16], where

bacteria movement typically lasted no more than 10 minutes.

Prior studies have proposed different models to explain the circularly confined motion of rod-

shaped swimmers[15,19,24]. However, these theories cannot explain the motion pattern difference we

observed for confined swarming and planktonic SM3. Noticing that swarming SM3 washed in LB

lost the single swirl pattern, we hypothesize that other than cell length or cell speed, the strong

cell-cell interaction may be a key factor responsible for the persistence of single swirls in the wells.

The mechanism of the rafting phenomenon of swarming cells has not been fully deciphered yet[3].

It might be due to cohesive interaction among neighboring cells together with hydrodynamic

effects among 2D-confined peritrichously flagellated bacteria[25]. The cell-cell interaction may

further result from biochemical change of cell envelope during swarming (e.g., more long

sidechain lipopolysaccharides) or secretions[26]. Once these surrounding matrix or polymers are

washed away by ~ 100-fold dilution, the cohesive interactions are diminished, resulting in no

dynamic clusters in the dilution experiment and multi-swirl motion pattern under confinement.

Reproducing the experimental results via computer simulation, we confirm that the lower repulsion

and higher alignment are the key factors, which differentiate swarmers and planktonic cells. Future

work is called upon to explore further the swarmer rafting phenomenon and to investigate on the

molecular level which category of substances are mainly responsible for the cell-cell cohesive

interaction among the swarming cells.

A spectrum of swarming bacteria has the same characteristic as SM3 (Fig. S1A). These bacteria

tested, including SM1, H6, H3, and Db10, all behave like SM3. They all show clustering or

cohesive cell-cell interaction when the swarming colony was diluted and uniformly dispersed

when the concentrated planktonic cells were diluted. One notable exception is *Bacillus subtilis*.

Swarming and concentrated planktonic *Bacillus subtilis* 3610 show the same motion pattern across

different confinement sizes. For well diameter $d \leq 90$ μm, both swarming and swimming *B. subtilis*

form single swirls while for well diameter $d \geq 112$ μm, they both break into mesoscale vortices. *B.*

subtilis is Gram-positive bacteria different from SM3, SM1, H6, H3, and Db10 and we speculate

that swarming *B. subtilis* does not have as strong cell-cell interaction as SM3. The interaction is

not so different between the swarming and planktonic cells since we found the diluted swarming

B. subtilis 3610 to disperse uniformly, much like diluted planktonic *B. subtilis* 3610, with no

clustering behavior. The swarming colony thickness might also be a key factor for *B. subtilis* to be

an outlier here. It is known that swarming *B. subtilis* produces abundant surfactant, resulting in a

wide-spread, monolayer, non-compact colony[20,27]. In contrast, swarming SM3 and the other tested

bacteria are multilayer colonies that can be as thick as 20 - 40 μm, which may enhance the spatial

cell-cell alignment that is much stronger than that among the planktonic cells (Fig. S1B).

Our observation on SM3 confirms the prediction made by Beppu *et al.* that single vortex occurs

when the confinement diameter d is smaller than a critical length $l*$[16]. Here, the critical length for

swarming SM3 is ~ 49 μm, whereas, for concentrated planktonic SM3, it is ~ 17 μm. Interestingly,

the same bacteria strain in different motility states has two distinct critical lengths. Thus, we were

able to use this property to identify the motility types on mouse mucosal surfaces. The beads

motion is not a perfect swirl in every well on the colitic tissue because the mucosal surface is not

as smooth as the agar surface. There are sags and crests on the inflamed mucosal surface due to

the disrupted mucin layer. We conjectured that this unevenness would hinder the swirl formation

to a certain extent and intact swirl patterns can be spotted only on limited locations where the

mucosal surface is relatively flat. Nevertheless, capturing only a few wells where beads show

single swirl motion suffices to show that swarming can happen on the mucosal surface.

Evidence of genetic up/down regulations[28-31] and cell morphology changes (e.g., cell elongation

and hyper-flagellation) indicates that swarming is a different phenotype from swimming. Lacking

comparison under the same conditions of experiments, one might suspect that bacterial swarming

might just be a dense group of cells swimming on a surface[3]. Here, through geometry confinement,

we show *Enterobacter* sp. SM3 as an example that swarming manifests different biophysical

characteristics from swimming. The key experimental method used in this study differentiates

swarming motility from swimming motility and provides a straightforward assay to detect

swarming behavior on a given surface visually. The findings of this study provide the rationale for

developing applications such as isolating bacterial swarmers from a polymicrobial environment

and developing diagnostics for the presence of *in vivo* swarming. A quantitative ranking system

for different swarmers could potentially be established based on characteristic well size that

stabilizes the confined motion pattern into a single swirl. Such a ranking system will be significant

for future investigations on the implications of swarming bacteria on host health and diseases.

Methods

PDMS confinement sheet fabrication. Polydimethylsiloxane (PDMS) microwell confinement

sheets with different combinations of well sizes and depths were fabricated using the technique of

soft photolithography. Patterns of the confinement were first designed using the software "L-Edit"

and then uploaded into a maskless aligner (MLA 150, Heidelberg). On a 3.5-inch silicon wafer

(University Wafer Inc.), photoresist gel SQ25 (KemLab, Inc.) was spin-coated at 2,000 rpm (spin

speed varies according to the desired coating thickness). After baking, UV exposure, and chemical

development, the microwells' designed pattern was shown on the wafer (moulding). Then, PDMS

(Dow Corning Sylgard 184) base elastomer was mixed with the curing agent at the ratio of 10:1 in

weight. The mixture was cast onto the patterned silicon wafer. Two grams of the mixture ended

up with a PDMS sheet about 0.5 mm thick. The PDMS solidified at room temperature within 48

hours and it was cut into pieces and peeled off from the silicon wafer before use (demoulding).

Bacterial growth and confinement (Fig. 6A). *Enterobacter* sp. SM3 is a novel swarming

bacterial strain isolated from inflammatory mice[10]. SM3 was transferred from - 80℃ glycerol stock

to fresh LB (Lysogeny Broth: water solution with 10 g/L tryptone, 5 g/L yeast, and 5 g/L NaCl)

and shaken overnight (~ 16 h) in a 37℃ incubator at 200 rpm. For swarming under confinement

assay (Fig. 6A, red arrows), 2 μL overnight bacterial culture was inoculated on the center of a LB

agar plate (10 g/L tryptone, 5 g/L yeast, 5 g/L NaCl, and 5 g/L Agar; volume = 20 mL/plate) and

kept in a 37°C incubator. After 2.5 h of swarming, a PDMS chip (~ 1 cm^2) was mounted upon the

edge of the swarming colony and the Petri dish was transferred onto the microscope stage for

observation. For swimming under confinement assay (Fig. 6A, blue arrows), overnight bacterial

culture was resuspended in fresh LB (1:100 in volume) and shaken in the 37°C incubator at 200

rpm for 2.5 h. The freshly grown culture was centrifuged at 1,500 g for 10 min and $\sim 98.6\%$ of the

supernatant was removed so that the resultant cell density is about 70 times the fresh grown culture.

Ten (10) μL concentrated bacteria culture was inoculated on the LB agar plate, and PDMS chip

was mounted immediately. The plate was then transferred onto the microscope stage for

observation. For other strains of bacteria, including *Bacillus Subtilis* 3610, the procedure was the

same as that of SM3. On one PDMS chip, there are thousands of wells and when mounted on a

bacteria spot or colony edge, hundreds of them are occupied by bacteria. The PDMS chip was first

brought to contact with the bacteria and then gently mounted onto the agar. In this case, there was

a cell density gradient across an array of wells, with the wells closer to the bacteria spot or colony

center having relatively higher cell density. In the experiment, we focused on the area where the

confined bacteria showed collective motion, i.e. the cell density was not too high to jam the well

or too low so that each cell was moving independently.

Bacterial cell density measurement (Fig. 6B). 2.5 h freshly grown SM3 was subjected to

different factors of dilution in LB, such as 10^2, 10^3, until 10^8. 50 μL of each diluted culture was

inoculated and spread on 1.5 % LB agar plate (10 g/L tryptone, 5 g/L yeast, 5 g/L NaCl, and 15

g/L Agar; volume = 20 mL/plate) and was incubated at 37°C for 16 h. Bacterial colonies appeared

on the agar plates and the number of colonies was counted for the dilution that resulted in the

colony's number on the order of 100. The colony forming unit per microliter (CFU/mL) was

calculated by dividing the colony number by the sampled volume. For swarming SM3, the cell

density was measured in a similar way. On the edge of the swarming colony, a chunk of swarming

SM3 (~ 1 mm wide) was picked by an eight (8) mm-wide square spatulate containing a small piece

of agar on the bottom to ensure all the cells in that region were sampled. The 1 mm x 8 mm chunk

of swarming SM3 was then mixed into 1 mL LB for CFU determination. The colony thickness

was assumed to be uniform across the sample and was measured by microscopy focusing on the

top of the colony and the top of the agar surface (i.e., at the bottom of the colony), keeping track

of the readings on the fine adjustment knob. Particles of baby powder (~ several micrometers in
diameter) were spread on the surfaces of the swarm colony and the agar to aid in the microscope
focus. The thickness of the swarming colony was calculated based on the calibration of the knob
turning tick readings. Then the cell density was estimated by CFU/mL. CFU was calculated for
both swarming and swimming SM3 to make sure the cell densities of these two cases were
comparable inside the wells. We consider colony forming unit counting a better way to control the
live cell number than simply using the volume fraction because: 1, Dead cells that count in the
volume fraction will not contribute to the motion in the well, but they will be excluded in CFU
calculation; 2, It is technically difficult to measure the volume of dense bacterial suspension using
pipetting method due to high viscosity.

Bacterial cell length and motility. For swimming SM3, 2.5 h freshly grown culture was diluted
100 times in LB and 50 μL of which was transferred on a glass slide and covered with a coverslip.
The sample slide was placed under the microscope (Olympus CKX41, 20X) and image sequences
were captured. Cell lengths were measured using ImageJ (v1.59e) freehand label tool. Cell speed
was calculated by traveling trajectory length divided by the traveling duration (~ 1s). For swarming
SM3, a chunk of swarming bacteria was collected from the edge of the swarming colony and mixed
with 1 mL LB. A droplet of 50 μL mixed culture was sandwiched by a glass slide and a cover slip,
and the rest of the procedure was the same as that for the swimming SM3.

Swimming SM3 with different treatments. i), Cephalexin treatment. Overnight SM3 culture
was diluted 100 times in fresh LB and incubated in a 37℃ shaker at 200 rpm for 1.5 h. Cephalexin
(CEP) was added to the culture so that the resultant concentration of CEP was 60 μg/mL. The
culture was kept in the shaker for another two (2) h before use. **ii), Surfactin additions.** After 2.5
h regrown culture was centrifuged, more supernatant was removed than usual, and surfactin was
added so that the resulting concentrations of surfactin were 10, 50, 100, 500 μM while the cell
density remained comparable to that of swarming SM3. **iii), Addition of swarming supernatant.**
Before swarming SM3 covered the plate, the colony was scratched carefully using a piece of
PDMS (~ 0.5 cm^2) and transferred into 1 mL deionized water. The mixture was sucked into a
syringe and filtered with a 0.2 μm filter. Then the solution was lyophilized to powder and then
dissolved into the concentrated swimming SM3 of roughly the same volume as the collected

swarm fluid. Thus, the concentration of the swarming supernatant was kept the same after being

transferred to swimming SM3.

Swarming SM3 with different treatments. i), Rubbed with PDMS. After SM3 swarmed on the

agar plate for 2.5 h, a piece of PDMS (~ 0.5 cm^2) was used to rub gently on the edge of the

swarming colony so that the swarming cells were disturbed. A PDMS confinement chip was then

mounted on the disturbed region for observation. **ii), Spun down in LB.** After swarming for 2.5

h, SM3 cells were collected from the edge of the colony using the blotting method[32]. The cells

were blotted by a piece of spare PDMS and transferred to 1 mL LB. The swarming cells were

centrifuged at 1,500g, and LB was removed in order to restore the initially high cell density. Ten

(10) μL of the swarming cells thus treated were inoculated on a new swarm agar and a PDMS

confinement chip was mounted for observation. **iii), Added D-mannose.** A droplet of 50 μL 0.2%

(w/v) D-mannose was pipetted on a swarming SM3 colony edge. After 1-2 minutes, when the cell

density became uniform again, a piece of PDMS confinement chip was applied to the D-mannose

treated region for observation under the microscope.

VOP measurement and spatial autocorrelation function. Image sequences of swarming or

swimming SM3 under confinement were taken by a microscope camera (ThorLabs, Kiralux

CS505MU) and then processed using a particle image velocimetry (PIV) package in MATLAB.

The velocity field was marked for the confined bacteria and the VOP was calculated using the

equation in Fig. 1E. Using the velocity field information, the spatial autocorrelation function was

calculated through the equation $C_r(\Delta r) = < \frac{v(r_0) \cdot v(r_0 + \Delta r)}{|v(r_0)^2|} >$, where r$_0$ is the local position vector

and Δr is the displacement vector[33]. A Python script was written to calculate all the C$_r$ values in

the region of interest (ROI) with a label of Δr values. These C$_r$ values were then plotted as a

function of Δr.

Clustering analysis. On the swarming SM3 colony edge or concentrated swimming SM3

inoculation, a droplet of 50 μL deionized water was added via a pipette. Once the fluid flow

stabilizes, image sequences were captured at the locations of the diluted swarming or swimming

SM3 samples. In a region of 130 μm x 130 μm, using the PIV toolkit, the velocity field was

calculated and the vectors with magnitude below four (4) μm/s were removed. The purpose of the

vector validation was to exclude non-motile bacteria. Once the moving cells were identified, a

Python script was implemented to perform the clustering analysis using the function of DBSCAN[34]

where the parameter ε was set to 50, which specifies how close points should be to each other to

be considered a part of a cluster, and the minimum number of points to form a cluster was set to

20.

Numerical Simulations. The numerical simulation consists of a 2D system of N particles. The

position **r** of each particle is modeled via the following overdamped Langevin equation:

$$\partial_t \boldsymbol{r_i} = v_0 \widehat{p}_i - \sum_{j \neq i} G_\theta(d_{ex}, r_{ji}) + \sqrt{2D_T}\xi_i \qquad (1)$$

It is assumed that particles are cruising at a constant speed of v_0 in the direction of $\widehat{p}_i =$

$[\cos(\theta_i), \sin(\theta_i)]$. The second term includes the exclusion forcing term from all neighboring

particles residing at a distance r_{ji} closer than the exclusion range d_{ex}. The last term is the thermal

fluctuation term with the translational diffusivity of D_T and a zero-mean and delta-correlated noise

term ξ. The direction of motion θ_i of each particle is updated by the interaction terms F_θ which

includes alignment, anti-alignment and repulsion effects with all neighboring particles and also the

rotational diffusion term with diffusivity of D_r and noise term ζ:

$$\partial_t \theta_i = \sum_{j \neq i} F_\theta(\boldsymbol{r_{ji}}, \widehat{p}_i, \widehat{p}_j) + \sqrt{2D_r}\zeta_i \qquad (2)$$

The details of the binary interaction terms G_θ and F_θ are provided in the Supplementary Material.

The simulation starts with random initial position and orientations, followed by numerical

integration of equations (1) and (2) using a first-order Euler method. The integration time step Δt

is chosen small enough to ensure numerical stability and also independence of long-term dynamics

from the time step increment. The interaction of particles with circular bounded domain is modeled

through a reflective boundary condition, where the particles are reflected off the boundary with an

angle equal to their incident angle. In all diluted cases, reflecting solid boundary is replaced with

a periodic boundary condition to ensure that boundary scattering is not affecting the dynamics in

the bulk.

Figure 1

Figure 1 | Swirls of *Enterobacter* sp. SM3 under circular confinement. (A-B) Motion pattern of
concentrated planktonic (A) and swarming (B) SM3 in the PDMS microwells of 74 μm in diameter. Circular
arrows indicate the direction of bacterial collective motion. (C-D) Velocity field of concentrated planktonic
(C) and swarming (D) SM3 in a single microwell. (E) Illustration of how vortex order parameter (VOP) is
defined. |·| denotes the absolute value while ‖·‖ denotes the Euclidean norm. (F) VOP of swarming and
swimming SM3 in 74 μm microwells of different depths. The sample size n = 5 for each group and data are
represented as mean and standard deviation (+SD).

Figure 2

Figure 2 | The effect of well diameter on confined *Enterobacter* Sp. SM3 motility patterns. (A-B)
Motion pattern of concentrated planktonic SM3 confined in 31 μm (A) and swarming SM3 confined in 112
μm (B) diameter microwells. (C) VOP of swarming and concentrated planktonic SM3 as a function of well
diameter. The error bars represent the standard deviations (± SD) for each data point, and the sample size
is n = 5. (D-E) Spatial autocorrelations of the bacterial velocity field in the well diameter of 90 μm (D) and
500 μm (E). Unless otherwise noted, the depth of the wells is 22 μm.

Figure 3

Figure 3 | Factors that possibly influence the bacterial motion pattern in the well. (A) Bacterial cell

length of planktonic, swarming, and Cephalexin (CEP) treated planktonic SM3, n = 500 for each group.

Data are represented as median and interquartile range. **** indicates $P < 0.0001$. ns indicates not

significant (Kruskal-Wallis test). (B) Bacterial cell speed of swimming, swarming, centrifuged, and CEP

treated swimming SM3, n = 10 for each group. ns, not significant, one-way ANOVA followed by Tukey's

post hoc test. (C) VOP of swimming SM3 under 74 μm diameter confinement with different treatments, n

= 5 for each group. (D) VOP of swarming SM3 under 74 μm diameter confinement with different treatments,

n = 5 for each group. B-D, Data are represented as mean and standard deviation (+SD).

Figure 4

Figure 4 | Spatial distribution of swarming and swimming SM3 cells. (A-B) Snapshots showing diluted

swarming SM3 (A) and swimming SM3 (B) on a soft agar surface, respectively. (C-D) DBSCAN clustering

analysis of diluted swarming SM3 (C) and swimming SM3 (D). Black dots represent moving bacterial cells

and colored markers show cells in clusters, as determined by the program. The axis represents the dimension

of the image in pixels.

Figure 5

Figure 5 | Numerical simulations of planktonic and swarimng SM3 in confinement and open space.

(A) VOP of swarming and concentrated planktonic SM3 as a function of well diameter. The error bars

represent the standard deviations (± SD) for each data point, and the sample size is n = 5. The circles on the

upper right corner and the lower left corner show representative motion patterns of swarmers and

concentrated planktonic cells in the confinement size between 0.38 and 0.5. (B) Planktonic cells (left) and

diluted swarming cells (right) with same cell density in a space of periodic boundary condition.

Figure 6

Figure 6 | Illustration of experimental procedure. (A) Schematic of sample preparation procedure. Red
arrows represent the assay procedure for swarming bacteria. Blue arrows represent the assay procedure for
swimming planktonic bacteria. (B) Cell density measured by colony forming unit (CFU/mL) of swarming
SM3 and swimming SM3. Swarming SM3 cell density is measured after SM3 swarming on an agar surface
for 2.5 h while swimming SM3 cell density is measured for overnight SM3 culture being regrown in fresh
Lysogeny Broth (LB) for 2.5 h. Since cell density of swarming SM3 was higher than that of planktonic
SM3, the latter was concentrated before being applied on the agar plate to acquire comparable cell density.

References

1. Webre, D. J., Wolanin, P. M. & Stock, J. B. Bacterial chemotaxis. *Curr Biol* **13**, R47-R49, doi:Doi 10.1016/S0960-9822(02)01424-0 (2003).

2. Josenhans, C. & Suerbaum, S. The role of motility as a virulence factor in bacteria. *Int J Med Microbiol* **291**, 605-614, doi:Doi 10.1078/1438-4221-00173 (2002).

3. Kearns, D. B. A field guide to bacterial swarming motility. *Nat Rev Microbiol* **8**, 634-644, doi:10.1038/nrmicro2405 (2010).

4. Partridge, J. D. & Harshey, R. M. Swarming: Flexible Roaming Plans. *J Bacteriol* **195**, 909-918, doi:10.1128/Jb.02063-12 (2013).

5. Darnton, N. C., Turner, L., Rojevsky, S. & Berg, H. C. Dynamics of Bacterial Swarming. *Biophys J* **98**, 2082-2090, doi:10.1016/j.bpj.2010.01.053 (2010).

6. Michaels, B. & Tisa, L. S. Swarming motility by Photorhabdus temperata is influenced by environmental conditions and uses the same flagella as that used in swimming motility. *Can J Microbiol* **57**, 196-203, doi:10.1139/W10-119 (2011).

7. Hall, A. N., Subramanian, S., Oshiro, R. T., Canzoneri, A. K. & Kearns, D. B. SwrD (YlzI) Promotes Swarming in Bacillus subtilis by Increasing Power to Flagellar Motors. *J Bacteriol* **200**, doi:UNSP e0052910.1128/JB.00529-17 (2018).

8. Partridge, J. D., Nhu, N. T. Q., Dufour, Y. S. & Harshey, R. M. Escherichia coli Remodels the Chemotaxis Pathway for Swarming. *Mbio* **10**, doi:ARTN e00316-1910.1128/mBio.00316-19 (2019).

9. Overhage, J., Bains, M., Brazas, M. D. & Hancock, R. E. W. Swarming of Pseudomonas aeruginosa is a complex adaptation leading to increased production of virulence factors and antibiotic resistance. *J Bacteriol* **190**, 2671-2679, doi:10.1128/Jb.01659-07 (2008).

10. Chen, W. *et al. Bacterial Swarmers exhibit a Protective Response to Intestinal Stress*, <https://www.biorxiv.org/content/10.1101/759886v3> (2020).

11. Wioland, H., Lushi, E. & Goldstein, R. E. Directed collective motion of bacteria under channel confinement. *New J Phys* **18**, doi:Artn 07500210.1088/1367-2630/18/7/075002 (2016).

12. Wioland, H., Woodhouse, F. G., Dunkel, J. & Goldstein, R. E. Ferromagnetic and antiferromagnetic order in bacterial vortex lattices. *Nat Phys* **12**, 341-U177, doi:10.1038/Nphys3607 (2016).

13. Theillard, M., Alonso-Matilla, R. & Saintillan, D. Geometric control of active collective motion. *Soft Matter* **13**, 363-375, doi:10.1039/c6sm01955b (2017).

14. Wioland, H., Woodhouse, F. G., Dunkel, J., Kessler, J. O. & Goldstein, R. E. Confinement Stabilizes a Bacterial Suspension into a Spiral Vortex. *Phys Rev Lett* **110**, doi:ARTN 26810210.1103/PhysRevLett.110.268102 (2013).

15. Lushi, E., Wioland, H. & Goldstein, R. E. Fluid flows created by swimming bacteria drive self-organization in confined suspensions. *P Natl Acad Sci USA* **111**, 9733-9738, doi:10.1073/pnas.1405698111 (2014).

16. Beppu, K. *et al.* Geometry-driven collective ordering of bacterial vortices. *Soft Matter* **13**, 5038-5043, doi:10.1039/c7sm00999b (2017).

17. Araujo, G., Chen, W. J., Mani, S. & Tang, J. X. Orbiting of Flagellated Bacteria within a Thin Fluid Film around Micrometer-Sized Particles. *Biophys J* **117**, 346-354, doi:10.1016/j.bpj.2019.06.005 (2019).

18   Doostmohammadi, A., Adamer, M. F., Thampi, S. P. & Yeomans, J. M. Stabilization of active
matter by flow-vortex lattices and defect ordering. *Nat Commun* **7**, doi:ARTN
1055710.1038/ncomms10557 (2016).
19   Hamby, A. E., Vig, D. K., Safonova, S. & Wolgemuth, C. W. Swimming bacteria power microspin
cycles. *Sci Adv* **4**, doi:ARTN eaau012510.1126/sciadv.aau0125 (2018).
20   Be'er, A. & Ariel, G. A statistical physics view of swarming bacteria. *Mov Ecol* **7**, doi:ARTN
910.1186/s40462-019-0153-9 (2019).
21   Grossmann, R., Romanczuk, P., Bar, M. & Schimansky-Geier, L. Pattern formation in active
particle systems due to competing alignment interactions. *The European Physical Journal*, 1325-
1347, doi:10.1140/epjst/e2015-02462-3 (2015).
22   Grossmann, R., Romanczuk, P., Bar, M. & Schimansky-Geier, L. Vortex arrays and mesoscale
turbulence of self-propelled particles. *Phys Rev Lett* **113**, 258104,
doi:10.1103/PhysRevLett.113.258104 (2014).
23   Turner, L., Zhang, R. J., Darnton, N. C. & Berg, H. C. Visualization of Flagella during Bacterial
Swarming. *J Bacteriol* **192**, 3259-3267, doi:10.1128/Jb.00083-10 (2010).
24   Tsang, A. C. H. & Kanso, E. Circularly confined microswimmers exhibit multiple global patterns.
Phys Rev E **91**, doi:ARTN 04300810.1103/PhysRevE.91.043008 (2015).
25   Li, Y., Zhai, H., Sanchez, S., Kearns, D. B. & Wu, Y. L. Noncontact Cohesive Swimming of
Bacteria in Two-Dimensional Liquid Films. *Phys Rev Lett* **119**, doi:ARTN
01810110.1103/PhysRevLett.119.018101 (2017).
26   Judith P. Armitage, D. G. S., R. J. Rowbury. Alternation in the cell envolope composition of
Proteus Mirabilis during the development of swarmer cells. *Biochimica et Biophisica Acta* **584**,
389-397 (1979).
27   Jeckel, H. *et al.* Learning the space-time phase diagram of bacterial swarm expansion. *P Natl Acad
Sci USA* **116**, 1489-1494, doi:10.1073/pnas.1811722116 (2019).
28   Tremblay, J. & Deziel, E. Gene expression in Pseudomonas aeruginosa swarming motility. *Bmc
Genomics* **11**, doi:Artn 58710.1186/1471-2164-11-587 (2010).
29   Wang, Q. F., Frye, J. G., McClelland, M. & Harshey, R. M. Gene expression patterns during
swarming in Salmonella typhimurium: genes specific to surface growth and putative new motility
and pathogenicity genes. *Mol Microbiol* **52**, 169-187, doi:10.1111/j.1365-2958.2003.03977.x
(2004).
30   Morgenstein, R. M., Szostek, B. & Rather, P. N. Regulation of gene expression during swarmer
cell differentiation in Proteus mirabilis. *Fems Microbiol Rev* **34**, 753-763, doi:10.1111/j.1574-
6976.2010.00229.x (2010).
31   Daniels, R., Vanderleyden, J. & Michiels, J. Quorum sensing and swarming migration in bacteria.
Fems Microbiol Rev **28**, 261-289, doi:10.1016/j.femsre.2003.09.004 (2004).
32   Darnton, N., Turner, L., Breuer, K. & Berg, H. C. Moving fluid with bacterial carpets. *Biophys J*
86, 1863-1870, doi:Doi 10.1016/S0006-3495(04)74253-8 (2004).
33   Patteson, A. E., Gopinath, A. & Arratia, P. E. The propagation of active-passive interfaces in
bacterial swarms. *Nat Commun* **9**, doi:ARTN 537310.1038/s41467-018-07781-y (2018).
34   Ester, M. K., H. Sander, J. Xu, X. in *Proceedings of the Second International Conference on
Knowledge Discovery and Data Mining.* 226-231 (AAAI Press).
518

Supplementary Information

Figure S1

Figure S1 | Other bacteria under confinement and their swarm front. (A) VOP of concentrated

planktonic and swarming Enterobacter sp. SM1, *Citrobacter koseri* (H6), *Serratia marcescens* (H3),

Serratia marcescens (Db10) and *Bacillus subtilis 3610* confined in the PDMS microwells of 58 μm in

diameter. The bars indicate averages with standard deviation (+SD) over 5 microwells. (B) Swarm front of

the tested bacteria. *B. subtilis* 3610 shows monolayer, loose swarming colony while all the other bacteria

strains show multilayer, compact swarming colony.

Figure S2

Figure S2 | Representative patterns at different sizes of bounded domain. Top row: Swarming; Bottom
row: Planktonic. The corresponding domain sizes and VOP values are marked as filled symbols. The
particle density is kept constant as the area of the simulated region increases.
534

Figure S3

Figure S3 | Fluorescent beads motion in microwells mounted on infected murine tissue. PDMS chips
were coated with 0.5 μm fluorescent beads and mounted on SM3 inoculated colitic (A) or normal (B) mice
intestine tissue surfaces. Average velocity field was calculated by tracing the beads motion using PIV
toolkit. (A) On colitic tissue, wells with VOP > 0.7 were found and marked with yellow squares. We
conclude that, in these wells, the single swirl motion pattern of the beads was powered by the confined
swarming SM3. Since the tissue surface was not as smooth as on agar surface, the motion of the beads in
some wells did not form a complete vortex, yet jets indicating partial vortices can be seen. (B) On a normal
tissue lacking swarming bacteria, the average velocity of the beads in the wells due to random motion is
close to zero, thus the VOP is uniformly small. We could infer that the confined SM3 in these wells were
predominantly swimming rather than swarming.

SI Methods

Detecting bacterial motility on mouse intestine tissue using PDMS chips. Six-week-old female C57BL/6 mice (Jackson Laboratories, Bar Harbor, ME; #000664) were administered 3%(w/v) DSS (Dextran Sulfate Sodium) (MPI; # 160110) in animal facility drinking water daily to induce acute colitis. After 9-12 days, when the mice weight loss reached 20%, mice were euthanized using isoflurane anesthesia and large intestines were harvested. For control, conventional six-week-old female C57BL/6 mice exposed to drinking water not containing DSS treatment were also sacrificed and the intestines were collected. This study was approved by the Institute of Animal Studies at the Albert Einstein College of Medicine, Inc (IACUC # 20160706 & 00001172). Intestine tissues were cut open, cleaned with 35%(v/v) ethanol, and rinsed with PBS twice. Tissues were spread on a 1% agar plate with inner side facing up, and overnight SM3 bacterial culture were inoculated on one end of the tissue. The agar plate was incubated under 37°C for 4.5 hours to allow SM3 bacteria to duplicate and move on the tissue surface. PDMS chips (d = 38 μm) were coated with 0.5 μm fluorescent beads (Dragon green; Bangs Laboratory, IN) and cut into strips to fit the size of the tissue. The PDMS strip was mounted and covered the tissue surface. Bead motion was observed under the fluorescent microscope (Olympus CKX41) with 20X objectives.

Numerical simulations. The dynamics of N interacting active particles have been modeled in a 2-dimensional space using the overdamped Langevin-based equations, assuming that inertia is negligible in a low Reynolds number environment. The position $\boldsymbol{r}$ and orientation θ of particle i are calculated using the following stochastic differential equations:

$$\partial_t \boldsymbol{r_i} = v_0\widehat{p_i} - \sum_{j\neq i} k_{ex} r_{ji} \mathcal{H}(d_{ex} - r_{ji}) + \sqrt{2D_T}\xi_i \tag{1}$$

$$\partial_t \theta_i = \sum_{j\neq i} F_\theta(\boldsymbol{r_{ji}}, \widehat{p_i}, \widehat{p_j}) + \sqrt{2D_r}\zeta_i \tag{2}$$

Based on our experimental observations, bacterial velocity in the suspension is largely independent of the local cell density. Accordingly, the self-propulsion speed of particles is set to be a constant v_0 along the direction $\widehat{p_i} = [\cos(\theta_i), \sin(\theta_i)]$. The second term incorporates the central exclusion force term with a spring constant k_{ex} which acts over the relative distance r_{ji} with all the neighboring particles j. This exclusion force term applies only when r_{ji} gets smaller than the

exclusion range d_{ex} (represented as a Heaviside function H). The last term in Eq. (1) is the

Brownian fluctuation term with the corresponding translation diffusivity D_T and ξ_i is the white

noise with zero mean and correlation $\delta(t)$.

The temporal change in the orientation of each particles is influenced by two terms. The first term

on the right-hand side of Eq. (2) includes all the binary interaction terms. The last term on the

right-hand side of Eq. (2) is the contribution from the angular Brownian fluctuation with the

rotational diffusion D_r and a zero mean and delta-correlated stochastic white noise ς. In the present

study, we employ the pair-wise interaction model introduced previously[21,22], which successfully

reproduces various macroscopic patterns reminiscent of bacterial suspensions. The pair-wise

interaction term is based on a zonal model (Figure S4 below) which captures the alignment, anti-

alignment and repulsion effects, and is formulated in the following form[21,22]:

$$F_\theta\left(\boldsymbol{r_{ji}}, \widehat{p_i}, \widehat{p_j}\right) = k_r\, \mathcal{H}\left(r_r - r_{ji}\right)\sin\left(\theta_i - \theta_{ji}\right) + \mu \sin\left(\theta_j - \theta_i\right) \tag{3}$$

583　k_r is the magnitude of the constant repulsion interaction that applies over distance of r_r around the

584　particle (Fig. 1). The second term in Eq. (3) represents the alignment and anti-alignment effects,

585　which operate over a range of r_a and r_{aa}, respectively. The magnitude of the aligning interaction m

586　is distance-dependent and is defined as [1, 2]:

$$\mu = \begin{cases} \mu^+(1 - (r_{ji}/r_a)^2) & 0 \le r_{ji} \le r_a \\ -\mu^- \dfrac{4(r_{ji} - r_a)(r_{aa} - r_{ji})}{(r_{aa} - r_a)^2} & r_a \le r_{ji} \le r_{aa} \end{cases} \tag{4}$$

where m$^+$ and m$^-$ are the strength of alignment and anti-alignment interactions, respectively.

We numerically integrate Eqs. (1) and (2) using the first order Euler scheme. Initially, the particles

are randomly distributed with random orientations. The integration time step Δt is selected

sufficiently small to ensure both numerical stability and also independence of long-term statistics

from Δt. The simulation time is set long enough to let the system reach a dynamic steady-state.

The interaction of particles with the bounded circular domain is modeled via a reflective boundary

condition.

Figure S4 | Schematic of the zonal pair-wise interaction model showing anti-alignment, alignment, and repulsion zones with the corresponding interaction radii r_{aa}, r_a and r_r.

In order to differentiate the Swarming and Planktonic cases, two different sets of interaction parameters have been used, which are summarized in Table 1. The values are unitless. In both systems, we set the exclusion parameters k_{ex} and d_{ex} to fixed values of 0.02 and 0.035, respectively. It is also assumed that particles only experience a rotational diffusion D_r of 0.75. The simulations for both Swarming and Planktonic cases have been performed at two particle densities $r = N/A_{dom}$, where N is the number of particles and A is the area of the simulation domain. In the high density case, $r = 4300$ and in the diluted case, we set $r = 235$. In the diluted case, in order to further minimize the boundary effects, we replace the bounded domain with a periodic boundary.

		Swarming	Planktonic
repulsion	k_r	2	3
	r_r	0.05	0.08
alignment	m^+	0.5	0.2
	r_a	0.2	0.2
Anti-alignment	m^-	0.5	4.0
	r_{aa}	0.25	0.25

SI Methods Table 1 | Simulation parameters used for the Swarming and Planktonic cases

SI Movies

All videos play in real time, with the exception of Movie S5 & S6 which were taken in 20 fps but compressed in 30 fps.

Movie S1: Confined swarming SM3 showing single swirl motion pattern. Swarming SM3 was confined in 74 µm diameter PDMS wells.

Movie S2: Confined concentrated planktonic SM3 showing turbulent motion pattern. Swimming SM3 was confined in 74 µm diameter PDMS wells.

Movie S3: Diluted swarming SM3 colony. Swarming SM3 colony edge was diluted by adding a 50 µL water droplet. Clusters of bacteria cells formed rafts.

Movie S4: Diluted swimming SM3 suspension. Concentrated planktonic SM3 was diluted by adding a 50 µL water droplet. Bacteria cells were observed to swim independently without clustering.

Movie S5: Fluorescent beads motion on DSS induced colitic mouse intestine tissue. The unidirectional rotation motion in 38 µm diameter wells indicates swarming SM3 on the tissue surface.

Movie S6: Fluorescent beads motion on normal mouse intestine tissue. The random motion in 38 µm diameter wells indicates planktonic SM3 on the normal mice tissue surface.

Movie S7: Numerical simulations of circularly confined SM3. Swarming SM3 (left) and concentrated planktonic SM3 (right) were simulated in the well size of 0.48. The video shows the representative confined motion pattern. Arrows indicate the moving direction of the particles.

Movie S8: Numerical simulations of SM3 cells in open space. Diluted swarming SM3 (left) and planktonic SM3 (right) were simulated without confinement with a periodic boundary condition. Cell density in both cases are $\rho = 235$, and the arrows indicate the moving directions of the particles.