EVOLUTION WITHOUT VARIATION AND SELECTION

LIANE GABORA AND MIKE STEEL

ABSTRACT. A central tenet of standard evolutionary theory is that evolution requires variation, which is subjected to selection. This theory provides a unifying explanation for how species have evolved, but it does not as readily apply to the origin of life, or to cultural evolution. We describe a means of attaining cumulative, adaptive, open-ended change that can occur in the absence of variation and selective exclusion, and in the absence of generations (i.e., no explicit birth or death). This second evolutionary process occurs through the assimilation, restructuring, and extrusion of products into the environment by identical, interacting Reflexively Autocatalytic and Foodset-generated networks. We refer to this more primitive evolutionary process as Self–Other Reorganisation (SOR) because it involves internal self-organising and self-maintaining processes within entities, as well as interaction between entities. SOR cannot replace natural selection; it operates alongside it. Since there is no self-assembly code, it is more haphazard than natural selection, and there is no discarding of acquired traits (a signature characteristic of natural selection). In the extreme, SOR can operate with just one entity, but it differs from learning because in general it operates across groups. In entities undergoing *both* natural selection and SOR, SOR can produce adaptive change across biological generations. We discuss the relationship between SOR and Lamarckism. We suggest that this more primitive process is operative during the initial stage of an evolutionary process, and that it is responsible for the origin and early evolution of both organic life, and human culture. In cultural evolution, SOR could increase homogeneity amongst members of a group and thereby foster group identity and cohesion.

Keywords: autocatalytic network, cultural evolution, evolution, origin of life, social learning

Addresses for Correspondence:

Liane Gabora
Department of Psychology, University of British Columbia, Kelowna BC, Canada
liane.gabora@ubc.ca

Mike Steel
Biomathematics Research Centre, University of Canterbury, Christchurch, New Zealand
mike.steel@canterbury.ac.nz

1. Introduction

A central tenet of evolutionary theory is that evolution requires variation upon which selection can act. This follows from Darwin's theory of natural selection, which holds that evolution is due to differential replication of randomly generated heritable variations in a population over generations, such that some traits become more prevalent than others. As Darwin himself made clear [9], this theory cannot explain how life began[1], but it has provided the foundation of modern biology, and a unifying principle for describing species change well after the origin of life. Darwin's theory arose in response to the paradox of how organisms accumulate adaptive change despite that traits acquired over a lifetime are eliminated at the end of each generation. His solution was to come up with a population-level explanation: although acquired traits are discarded, inherited traits are retained, so evolution can be explained in terms of preferential selection for those inherited traits that confer fitness benefits on their bearers. However, neither origin of life research nor cultural evolution research is plagued by the problem that acquired traits are extinguished at the end of a generation; therefore, in these domains, the paradox that Darwin's theory was designed to solve does not exist [19]. Accordingly, it would seem appropriate to pay particular attention in these domains to the role of epigenetic and non-Darwinian (e.g., Lamarckian) processes. However, it is widely assumed that any kind of evolutionary process requires variation and selection. Even in research on cultural evolution— which is sometimes (though not universally) regarded as Lamarckian—it is generally assumed that variation and selection are what enables evolution to take place (e.g., [3, 6, 61]).

Indeed, although the term 'evolution' is sometimes assumed to be synonymous with variation and selection, it is increasingly accepted that non-Darwinian processes play a role in evolution [17, 48, 49, 80, 87]. This paper goes a step further by providing an existence proof that evolution is possible in the *absence* of variation and selection. Although this has been suggested before

---

[1]Since natural selection requires the existence of multiple competing entities that are selected amongst, it cannot explain how the first living entity arose [17].

[20], here, we formulate the argument in a more rigorous fashion by mathematically describing what kind of structure an entity must have to engage in this kind of evolutionary process. This is important, because although entities of many kinds change due to acquired traits—e.g., a rock tumbling down a stream will acquire rounded edges—this change may not be adaptive, or contribute to the entity's efficiency, survival, or replication. Indeed, in the case of the rock, the process that rounds its edges culminates in its disintegration. In contrast to the rock—and also, in contrast to neutral evolution [51, 52]—in the example we will provide, the change is adaptive.

In origin of life research, there is a sizeable literature on models of the evolvability of primitive metabolic networks prior to DNA (see for example [55, 70, 72, 78, 88], and the references therein). Here, in order to describe evolutionary processes as seemingly disparate as the origin of life and the origin of culture within a common framework, we necessarily take a higher level and more general approach, using a different (though related) mathematical modelling technique from these earlier studies. The paper begins with a list of definitions of terms and a table of the acronyms used, followed by an introduction to the framework that will be used to develop the argument. We then provide a mathematical investigation of a simple model for evolution in the absence of variation and selection with this framework. Next, we discuss what sort of structure is able to exhibit this form of evolution. We conclude with a discussion of the implications.

2. Preliminaries

We begin by alphabetically listing and defining key terms used in this paper. We believe this is necessary in order to maintain clarity, since miscommunication has arisen from inconsistencies in how evolutionary concepts are applied in the sciences and social sciences.

- **Acquired trait:** a trait obtained during the lifetime of its bearer (e.g., a scar, tattoo, or the memory of a song) and transmitted horizontally (i.e., laterally).

- **Culture:** extrasomatic adaptations—including behavior and artifacts—that are socially rather than sexually transmitted.

- **Darwinian (or 'selectionist') process:** an evolutionary process—i.e., a process that exhibits cumulative, adaptive, open-ended change—occurring through natural or artificial selection.

- **Darwinian threshold:** transition from a non-Darwinian to a Darwinian evolutionary process [87, 80].

- **Evolution:** descent with modification, or cumulative, adaptive change over time, giving rise to new species that share a common ancestor [9]. The most well-known theories to explain how evolution works are natural selection and Lamarckism. The concept of evolution has been extended to apply to, not just to cumulative, adaptive change in biological species, but also cumulative, adaptive *cultural* change [3, 6]. Thus, here we use the more general definition of evolution used in cultural evolution research: cumulative, adaptive change over time.

- **Generation:** a single transition period from the internalized to the externalized form of a trait.[2]

- **Horizontal transmission:** non-germ-line transmission of an acquired trait from one entity to another. Thus, social transmission is horizontal because the information is not inherited, i.e., it is not passed from parent to offspring by way of genes.

- **Lamarckian process (or 'soft inheritance'):** widely understood to mean the transmission of characteristics to offspring that were acquired through use or disuse during its lifetime.[3] Some maintain that Lamarckian evolution requires genetic transmission to

---

[2]Note that, with respect to biological evolution, a new generation generally (though not in horizontal gene transfer) begins with the birth of one or more organism(s). With respect to cultural evolution, a new 'generation' may begin with the expression of an idea (which is considered one transmission event). Thus, over the course of a single discussion, an idea (a cultural trait) may undergo multiple generations.

[3]We note that although this is how the term is standardly defined, it has been pointed out that this idea was not unique to Lamarck, and transmission of characteristics was only one component of Lamarck's theory [31].

biological offspring, a view held by early biologists (e.g., [71]), though current scholars (e.g., [61]) sometimes take a more equivocal view.

- **Neutral theory of molecular evolution**: the theory that a significant degree of genetic variation in populations is the result of mutation and genetic drift, not selection [51, 52].

- **Selection:** differential replication of randomly generated heritable variation in a population over generations such that some traits become more prevalent than others. Selection may be natural (due to non-human elements of the environment) or artificial (due to human efforts such as selective breeding), and it can occur at multiple levels, e.g., genes, individuals, or groups [57].

- **Self-assembly code:** a set of coded instructions that is: (i) *actively interpreted* through a developmental process to generate a soma, and (ii) *passively copied without interpretation* through a reproduction process to generate self-copies, i.e., new sets of self-assembly instructions that are in turn used in these two distinct ways.

- **Self–Other Reorganisation (SOR):** a theory of how both culture, and early life, evolve through interacting, self-organizing networks, based on theory and findings that have come out of the post-modern synthesis in biology [17, 74].

- **Vertical transmission:** Germ-line inheritance of a trait from one generation to the next. (In other words, transmission occurs by way of a self-assembly code, such as DNA.)

It is important to emphasize that we are using the term 'selection' in its technical, scientific sense. The word 'selection' also has an 'everyday' sense in which it is synonymous with 'choosing' or 'picking out'. One could say that selection—in the everyday sense of the term—occurs in a competitive marketplace through the winnowing out of inferior products. However, in the discussion that follows. the term 'selection' is used in its scientific sense.

Our definition of 'self-assembly code' is consistent with Von Neumann's work on universal constructors and relf-replicating automata [81], and subsequent insights by Holland [37], and Langton [56]. In a strictly Darwinian or selectionist process, traits acquired through interaction with the soma (the culmination of the *active interpretation* of the code) do not affect traits vertically transmitted through reproduction (the culmination of the *passive uninterpreted* use of the code). As Holland [37] put it, the self-assembly code passed down through the germ-line is *sequestered*, i.e., protected from the influence of acquired traits. Thus, it is the division of labour between these two ways of using the self-assembly code that is responsible for the signature characteristic of a selectionist process: lack of transmission of acquired traits.

Note that vertical and horizontal transmission must be defined with respect to the relevant evolutionary process. A common error is to refer to the transmission of cultural information from parent to offspring as vertical transmission (e.g., [6]). Although the individuals in question are in a parent–child relationship with respect to their status as biologically evolving organisms, this may not be the case with respect to their status as participants in cultural evolution. Indeed, although childbirth entails one mother and one father, there is no limit to the number of 'parental influences' on the 'birth' of an idea.

A related error is to say that in cultural evolution, there is a third form of transmission, namely, oblique transmission, in which traits are transmitted from non-kin members of the parental generation (e.g., [6]), because as far as cultural evolution is concerned it is not strictly relevant whether the information comes from biological kin or non-kin. In a similar vein, although dual inheritance theorists speak of culture as a second form of inheritance [35, 68, 85, 64], the distinguishing feature of an inherited trait is that it is transmitted vertically (e.g., from parent to offspring) by way of a self-assembly code (e.g., DNA), and therefore not obliterated at the end of a generation. This is not the case with respect to cultural traits [19] (nor is it even the case for all biological traits).

The acronyms used in this paper are listed alphabetically in Table 1.

Acronym	Meaning
CRS	Catalytic Reaction System
RAF	Reflexively Autocatalytic and Foodset-generated (F-generated)
OOL	Origin of Life
OOC	Origin of Culture
MR	Mental Representation
CCP	Cognitive Catalytic Process
SOR	Self–Other Reorganisation

TABLE 1. Acronyms used throughout this paper.

3. AUTOCATALYTIC NETWORKS

The type of evolution without variation and selection process that we will describe involves entities that (1) have an organisation that is self-maintaining (i.e., they have a structure, and a mechanism for maintaining that structure), and (2) interact with each other by way of their environment. The theory of autocatalytic networks grew out of studies of the statistical properties of *random graphs* consisting of nodes randomly connected by edges [11]. As the ratio of edges to nodes increases, the size of the largest cluster increases, and the probability of a phase transition resulting in a single giant connected cluster also increases. The recognition that connected graphs exhibit phase transitions led to their application to efforts to develop a formal model of the origin of life (OOL), namely, of how abiogenic catalytic molecules crossed the threshold to the kind of collectively self-sustaining, self-replicating, evolving structure we call 'alive' [48, 47]. Applying graph theory to the OOL, the nodes represent catalytic molecules, and the edges represent reactions. It is exceedingly improbable that any catalytic molecule present in the primordial soup of Earth's early atmosphere catalysed its own formation. However, reactions generate new molecules that catalyse new reactions, and as the variety of molecules

increases, the variety of reactions increases faster. As the ratio of reactions to molecules increases, the probability increases that the system will undergo a phase transition. When, for each molecule, there is a catalytic pathway to its formation, they are said to be collectively *autocatalytic*, and the process by which this state is achieved has been referred to as *autocatalytic closure* [48]. The molecules thereby become a self-sustaining, self-replicating structure, such as a living protocell [43]. Thus, the theory of autocatalytic networks provides a promising avenue for modelling the OOL and thereby understanding how biological evolution began [88]. In this section, we summarize how autocatalytic networks have proven particularly useful in both in OOL and origin of culture (OOC) settings.

3.1. **Reflexively Autocatalytic and Foodset-generated (RAF) networks.** Autocatalytic networks have been developed mathematically in the theory of Reflexively Autocatalytic and Foodset-generated (RAF) networks [44, 76]. The term *reflexively* is used in its mathematical sense, meaning that every element is related to the whole. The term *foodset* refers to the reactants that are initially present, as opposed to those that are the products of catalytic reactions. (Elements of a 'foodset' are not the same thing as 'food'; the foodset is simply the raw building blocks available for catalyzed reactions.)

We now summarise the key concepts of RAF theory. A *catalytic reaction system* (CRS) is a tuple $\mathcal{Q} = (X, \mathcal{R}, C, F)$ consisting of a set X of element types, a set $\mathcal{R}$ of reactions, a catalysis set C indicating which element types catalyse which reactions, and a subset F of X called the foodset. A *Reflexively Autocatalytic and F-generated* set (i.e., a RAF) is a non-empty subset $\mathcal{R}' \subseteq \mathcal{R}$ of reactions that satisfies the following two properties:

(1) *Reflexively autocatalytic (RA)*: each reaction $r \in \mathcal{R}'$ is catalysed by at least one element type that is either produced by $\mathcal{R}'$ or is present in the foodset F; and

(2) *F-generated*: all reactants in $\mathcal{R}'$ can be generated from the foodset F by using a series of reactions only from $\mathcal{R}'$ itself.

A set of reactions that forms a RAF is simultaneously self-sustaining (by the F-generated condition) and (collectively) autocatalytic (by the RA condition) because each of its reactions is catalysed by an element associated with the RAF. Note that in RAF theory, 'catalysis' does not necessarily require that a catalyst to cause more of something to be produced (or at a faster rate); rather, the catalyst is simply the impetus that allows a reaction to proceed.

A CRS need not have a RAF; alternatively, it may contain many RAFs. When a CRS does contain RAFs there is a unique maximal one, referred to as the *maxRAF*. It is the fact that a CRS may contain multiple RAFs that allows RAFs to evolve, as demonstrated both in theory and in simulation studies, through selective proliferation and drift acting on the RAFs that are subsets of the maxRAF [44, 78]. Fig. 1 provides an example that illustrates how a RAF can contain multiple RAFs within it. The maxRAF, and possibly some of the RAFs it contains are *closed*, in the sense that the RAF contains every reaction r in R for which each reactant and at least one catalyst of r is either in the food-set or is a product of a reaction in the RAF.

Table 2 summarises how RAF theory terms apply in biological and cultural/cognitive settings. The fact that RAFs have proven useful in both domains suggests that RAF theory may provide a broad conceptual framework that could turn out to be applicable to the origins and early stages of diverse evolutionary processes. We now examine OOL and OOC applications of RAFs in more detail.

RAF Theory	Origin of Life (OOL)	Origin of Culture (OOC)
node	catalytic molecule	mental representation (MR)
edge	reaction pathway	association
cluster	elements connected via reactions	MRs connected via associations
connected graph	autocatalytic closure [47, 48]	conceptual closure[4] [13]

TABLE 2. Application of graph theoretic concepts to the origin of life and origin of culture.

FIGURE 1. A RAF can contain a vast array of other RAFs within it. The analysis of this 7-reaction system in [41] (which arose in an earlier RNA system constructed in a laboratory-based experimental model of early life) revealed that more than half of all the $2^7 = 128$ subsets of this RAF are themselves RAFs. In this figure, the reactions (r_i) are labelled by square nodes, and round nodes indicate element types (which consist of a single food-set element f, along with derived element types x_i). Catalysis is indicated by dashed arrows. Figure produced using the RAF program *CatlyNet*.

3.2. **Biological Applications of RAFs.** RAFs have proven useful for modelling the OOL and the onset of biological evolution [44, 76, 78, 88]. In the OOL context, a RAF emerges in systems of polymers (molecular strings composed of repeated units called monomers) when the complexity of these polymers (as measured by their maximum length) reaches a certain threshold [48, 63]. The phase transition from no RAF to a RAF incorporating most or all of the elements depends on (1) the probability of any one polymer catalyzing the reaction by which a given other polymer was formed, and (2) the maximum length (number of monomers) of polymers in the system. This transition has been formalised and analysed mathematically, and by using simulations, and RAF theory has been applied to real biochemical systems [39,

---

[4]Conceptual closure is the focus of another paper [22].

40, 42, 44, 63], as well as to ecologies [7]. The theory has proven useful for identifying how phase transitions might occur and at what parameter values. The evolution and reproduction of RAFs in an OOL context has been studied [63].

3.3. **Applications of RAFs to Culture and its Cognition.** Autocatalytic networks have been used to model cognitive transitions associated with the origin of culture and the capacity for cumulative cultural evolution [13, 14, 25, 27, 26], and the onset of behavioral modernity (i.e., the capacity to think and act like modern humans) [26]; for related approaches see [1, 5, 65]. Also, the evolution and reproduction of RAFs has been studied in an OOC context [27]. RAFs have also been used to develop a process model of conceptual change (i.e., the process by which a child becomes an active participant in cultural evolution) [22], using data on conceptual change in childrens' mental models of the shape of the earth [83].

Whereas in the OOL context, a reaction network may be bounded by a semipermeable lipid membrane, in the OOC context, the network is a conceptual network of knowledge and experiences, which supervenes upon a brain within a human body, which thus constitutes a boundary on the cognitive RAF. Whereas the OOL setting involves networks of chemical reactions, in the OOC context, the network is a conceptual networks of knowledge and memories, and the products and reactants are not catalytic molecules but culturally transmittable *mental representations*[5] (MRs) of knowledge, experiences, ideas, schemas, and scripts. Despite these marked differences, the OOL and OOC settings both exhibit an underlying RAF structure. Just as reactions between molecules generate new molecules, interactions between MRs generate new MRs, which in turn enable new interactions. In cognitive RAFs, the foodset (i.e., elements of F) is composed of pre-existing information, i.e., MRs that do not have to be derived from scratch. They could be innate, acquired through social learning processes such as imitation,

---

[5]Although we use the term 'mental representation,' our model is consistent with the view (common amongst ecological psychologists and in cognition and quantum cognition communities) that what we call mental representations do not 'represent' but instead act as contextually elicited bridges between mind and world.

or acquired through individual learning from the natural world or human-made artifacts. The set of foodset-derived elements (i.e., elements of $\neg F$) result from creative processes such as representational redescription or restructuring, resulting in new information.

There will generally be multiple RAFs in the conceptual network of a single individual, which may be subsumed by a larger RAF (the maxRAF) that connects them all; alternatively, they may be disconnected (as when inconsistent views are held by the same individual).

The RAF approach to culture and its cognitive underpinnings offers several advantages over other approaches:

- **Semantic grounding.** A long recognized problem in cognitive science is the circularity of defining mental representations in terms of other MRs [34]. The distinction between foodset items and foodset-derived elements provides a natural means of grounding abstract concepts in direct experiences; foodset-derived elements emerge through 'reactions' that can be traced back to foodset-items.

- **Reactivity of ideas.** In cognitive RAFs, mental representations (MRs) become aligned with needs and desires (e.g., the need to solve a problem or resolve cognitive dissonance), and the emotions they elicit. In so doing, they incite ('catalyze') interactions between other mental representations (modeled as 'reactions'), which can be carried out recursively. As a result, whereas conventional conceptual network models require external input to continue processing, RAFs 'catalyze' conceptual change endogenously (in the absence of new external input), resulting in new conduits by which needs and goals can be met. Thus, a RAF network is self-organizing, and conceptual restructuring can percolate throughout the network and affect its global structure.

- **Track cultural change within and across conceptual networks.** The approach tags mental representations with their source, i.e., whether they were (1) innate, (2) acquired through social learning or individual learning (of pre-existing information), or (3) the result of creative thought (resulting in new information). This demarcation makes it possible to trace innovations back to the individuals that generated them, and precisely describe and track how new ideas and cultural outputs emerge from previous ones.

4. A RAF MODEL OF EVOLUTION WITHOUT VARIATION AND SELECTION

We now demonstrate a primitive non-Darwinian form of evolution using RAF networks. We begin by describing a simple process involving a group $\mathcal{G}$ of indistinguishable entities, as illustrated in Fig. 2. (They may be CRSs such as those associated with very early life [2, 8, 17, 30, 45, 75, 80], culturally evolving human conceptual networks (e.g., [12, 28]), or some other structure we have never encountered.) The entities are described as identical RAFs.

One entity in $\mathcal{G}$, which we call entity i, encounters a stimulus from the environment, which we refer to as s. In the case of a primitive biochemical reaction network, s could be an element that crosses the lipid membrane that encases the reaction network. In the case of culture, s could be a visual or auditory stimulus. What makes s a novel stimulus is that at t_0 it was not part of the reaction network of any entity in $\mathcal{G}$. Stimulus s is nonetheless part of the foodset, because it is available as an initial reactant or catalyst, as opposed to being directly or indirectly *constructed from* the set of initial reactants.[6]

Stimulus s triggers one or more catalytic reaction(s) in i. In a OOL scenario the reactions would be chemical reactions, while in a OOC scenario they would be mental operations such as

---

[6]It is a matter of time before one of the entities encountered s; the more abundant s is, the less time it is expected to take for this chance encounter to occur. It is possible that the encounter with s results from the discovery of a previously uncharted part of the environment that is rich in s, but it could also be that s entered the environment from elsewhere, or that s was always there but by chance had not previously interacted with any of the entities.

FIGURE 2. Identical entities exhibit cumulative adaptive change over time, as indicated by transition from light to dark, as a result of cumulative environmental changes, indicated by p, p' .... The first product p, generated by i due to a stimulus s catalyses a reaction in the RAF of entity i. The structure of i's RAF is illustrated in the blow up on the upper left (with element types $\alpha - \epsilon$ and s, reactions indicated by squares, and catalysis indicated by dashed arrows. The foodset here is $F = \{\alpha, \delta, s\}$, and ϵ gives rise to p). The product p in turn catalyses a reaction sequence in j at time t_2, leading to p', and so on. In the simple scenario outlined here, stimulation of i by s occurs only once, but that is sufficient to result in the generation of p, which is sufficient to set in motion a sequence of adaptive changes.

restructuring or concept combination that modify MRs. (n RAF models, recursive sequences of such 'reactions' have been referred to as *cognitive catalytic processes* (CCPs) [25, 27]).[7] This reaction (or chain of reactions) culminates in the generation of a product p into the environment. Thus, p was not part of the foodset; it is the end result of the series of reactions (or interactions) triggered by s. Note that s can only trigger the formation of p if the necessary sequence of

[7]See [18] for an explanation of why the innovation generation process does not require selection, and is not Darwinian.

reactions is able to take place, and it is the RAF that enables this reaction sequence to take place. Product p confers some adaptive benefit on i. Exposure to p may improve the efficiency of i's reaction network (or conceptual network) by reducing the number of steps required for a particular reaction sequence (or problem-solving process). It may have a protective function (for example, by making the environment less inviting for a predator) or (in a cultural context) it may serve as a source of pleasure that enhances wellbeing (such as a work of art). This triggering event initiates a sequence of one or more reactions within the RAF network of entity i that generates p, and for which the initial reaction in the sequence is catalysed by s. For example, in the OOC context, s catalyses the formation of a transient CCP that is part of individual i's RAF (as illustrated in Fig. 2(ii) of [28]). Having generated p, i returns to its initial state, except that it now experiences the adaptive benefit of p.

Once p is released into into the environment, it can be encountered by other entities in $\mathcal{G}$. (In an OOL context, p may be transmitted because it passes through a lipid membrane or protocell wall. In an OOC context, p may be transmitted via social learning, or by way of an artifact.) A recipient of p could have had the capacity to generate p on its own, but by chance had never done so. (In a OOL context, for example, a particular catalyst may have never come in contact with its reactant. In a cultural context, an individual may possess two concepts but never thought of combining them.) Alternatively, it is possible that the recipient did not previously possess the capacity to generate the product.

Since all the entities are identical to i and exist in the same environment as i, the adaptive benefit of p percolates throughout $\mathcal{G}$.[8] The generation of p makes it possible to generate a

---

[8]The word 'percolates' is used here to refer to the 'communal exchange' (between-entities) component of the process, as opposed to the 'self-organization' (within-entities) component of the process. If one entity encounters p before another, that first entity may temporarily function more efficiently. We note also that the scenario could unfold such that each entity is affected solely by its own products, and thus after the impact of the initial generation of p by i, there could be little interaction, direct or indirect, amongst them. In other words, it is possible that j is affected by the p' it generated itself (as when someone is affected by looking at their own art). However, it is also possible that j is affected by the p' generated by another entity in $\mathcal{G}$ (as in when someone is affected by looking at art made by someone else). Finally, we note that possession of a reactant (which in the cognitive scenario is a MR of a particular product) does not guarantee catalysis of that reactant.

new product, p', which confers even greater adaptive benefit than p. An entity j assimilates p, which triggers a sequence of one or more reactions in j, resulting in p'. Entity j benefits from the enhanced adaptive quality of p' over p. The ability to produce p' and experience its adaptive benefit percolates throughout the group $\mathcal{G}$. Thus, each version of p paves the way for its successor. The entities in $\mathcal{G}$ continue to generate increasingly beneficial versions of the original p (e.g., $p', p'', p''', \ldots$). They collectively come to possess the ability to generate and benefit from this family of products through the restructuring of their RAFs, catalyzed by the products of other entities. Thus we have cumulative adaptive change over time. Notice that in this model neither the birth nor the death of entities occurs; it is the same set of entities at the beginning as at the end, and there is no competitive exclusion and no selection of entities.

4.1. **Modelling this process.** We consider a simple model based on two processes: percolation and generation. Let ρ denotes the rate of percolation of products, or knowledge (i.e., MRs) of these products through the group $\mathcal{G}$, with neighbourhood structure represented by a directed graph $D_{\mathcal{G}}$. (The nodes of this graph are the entities in $\mathcal{G}$, and the arcs indicate directed percolation links). For the generation process, we let λ denote the rate (per entity) at which new products are generated. Provided that the ratio ρ/λ is not too small, the entities evolve via cumulative, adaptive change, with the only variation being between those in $\mathcal{G}$ that have not yet switched to a new product and those that have[9]. Moreover, for large values of ρ/λ, each percolation step will be complete before the next new product is generated, so there will be no variation amongst the entities.

Conversely, as $\rho/\lambda \to 0$, the entities diverge from each other with respect to the products they generate. Their complexity (which may be estimated through analysis of the maximal

---

For example, someone may have a memory of an artwork, but while the initial experience of it yielded little cognitive restructuring, the second experience of it triggers a productive stream of thought.

[9]This might be likely in an cultural setting when the entity generating a new product is equidistant from all the others or (more particularly in a social media context) where everyone sees something at the same time.

number of modifications to any product at any given time for the entities in $\mathcal{G}$) is expected to be lower than in the previous scenario of shared cumulative adaptive change.

To help formalise the model, we adopt the following terminology. Given the products $p, p', p'', \ldots$, we say that the entity that first gives rise to one of these products *generates* it. In the case where this product is produced from an earlier product in the series (rather than being the original product p in response to a stimulus), we say the entity *transforms* the existing product to the new one. We model the increase in the adaptive value of products within $\mathcal{G}$ using two stochastically independent non-deterministic processes.

First, the generation of new products by entity i involves either generating a new product, or transforming the most recent version of any available product; for simplicity, we assume that these are equally probable. For example, if entity i has the products (p, p', q, r, r', r'') currently available, then it can either transform p', or q, or r'', or generate a new product, and each of these four outcomes has the same probability, namely $1/4$. This process across entities is assumed to be described by independent exponential random variables with a fixed rate per entity of λ. (Making λ independent of the entity is consistent with the assumption that all entities are initially identical).

Second, each newly generated product begins to percolate through the group by moving along the arcs of the directed graph $D_{\mathcal{G}}$ according to a continuous-time random walk at rate ρ. For simplicity, we will treat the two processes—generation and percolation—as stochastically independent. Let $N = |\mathcal{G}|$ (the number of entities in $\mathcal{G}$). We will assume that the directed graph $D_{\mathcal{G}}$ that describes the community interactions within $\mathcal{G}$ has the property that from each entity in $\mathcal{G}$, it is possible to reach any other entity in $\mathcal{G}$ by following some directed path in $D_{\mathcal{G}}$ (i.e., $D_{\mathcal{G}}$ is 'strongly connected'); however, no further assumptions are made regarding this graph.

We now introduce some notation to keep track of the different versions of products that arise in the process described in Section 4. Suppose that product p first arises from entity i_1, and

product p is then further modified by entity i_2, and so on, with the last modification (before time t) being made by entity i_k. We denote the sequence of products thus generated within $\mathcal{G}$ up to time t as: $p(i_1), p(i_1, i_2), p(i_1, i_2, i_3), \ldots$. More generally, we denote such a sequence by writing $(p(i_1), \ldots, p(i_1, i_2, \ldots, i_k) : k \geq 1)$ (thereby allowing the possibility that a product is generated but not transformed, in the case where $k = 1$). We refer to the number k of terms in this sequence as the *complexity* of the final product; thus, when an entity transforms a product, it increases its complexity by 1 (in particular, the first product $p(i_1)$ has complexity 1).

Note that under the assumptions of the model, the entities $i_1, \ldots, i_k$ are not necessarily distinct (i.e., an entity may enhance a product more than once, either consecutively, or later in the sequence). There may also be several such sequences generated within $\mathcal{G}$; for example, in addition to the previous sequence, one might also have $p(j_1), p(j_1, j_2), \ldots, p(j_1, j_2, \ldots, j_l)$, along with possibly other sequences generated over the time interval $[0, t]$.

In this section, we let $\mathbb{P}(*)$ denote the probability of event $*$ and $\mathbb{E}[**]$ denote the expectation of random variable $**$. Let $T_\rho(i)$ be the expected time for a product generated by entity i to percolate (within $D_\mathcal{G}$) to every entity in $\mathcal{G}$, and let $T_\rho = \max\{T_\rho(i) : i \in G\}$. For a wide range of standard percolation processes, the following properties then hold: (i) for $\rho > 0$, we have $\mathbb{E}[T_\rho] < \infty$; (ii) for all $\eta > 0$, $\lim_{\rho \to 0} \mathbb{P}(T_\rho(i) > \eta) = 1$, and (iii) $\lim_{\rho \to \infty} \mathbb{E}[T_\rho] = 0$. This last property implies that when ρ is large, items are highly likely to percolate throughout the entire group $\mathcal{G}$ in a short time.

If we start this process at time 0 with no products present, let $\tau_1, \tau_2, \ldots, \tau_k$ be the random variables that describe the time intervals between the generation of products across the collection of entities in $\mathcal{G}$. By the assumptions of the model, the τ_i variables are independent and exponentially distributed, with each variable having an expected value of $1/(N\lambda)$. Thus $\sum_{i=1}^{k} \tau_i$ is the time until k products have been generated (this has a gamma distribution with expected value $k/(N\lambda)$). Let $\mu = N\lambda$. Then, for any $\eta > 0$, $\mathbb{P}\left(\bigcap_{i=1}^{k}\{\tau_i \geq \eta\}\right) = e^{-\mu k \eta}$ and $\mathbb{P}(T_\rho \leq \eta) \geq 1 - \mathbb{E}[T_\rho]/\eta$ (by the Markov inequality). Let $\mathcal{E}_k$ denote the following event: for

each of the first k products generated, each product percolates to each entity in $\mathcal{G}$ before the next new product (in this collection of size k) is generated in $\mathcal{G}$. We then have:

$$(1) \qquad \mathbb{P}(\mathcal{E}_k) \geq e^{-\mu k \eta} \cdot (1 - \mathbb{E}[T_\rho]/\eta)^k = (e^{-\mu \eta}(1 - \mathbb{E}[T_\rho]/\eta))^k.$$

Setting $\eta = \sqrt{\mathbb{E}[T_\rho]}$ in (1) and applying Property (iii) above gives:

$$\lim_{\rho \to \infty} \mathbb{P}(\mathcal{E}_k) = 1.$$

Thus, as ρ becomes large, the entities evolve collectively, and any variation is transient and short-lived. We will refer to this limiting case as the *community-based model*. One can model this process by the following novel type of Pólya Urn model:

> Consider an urn that initially has a single white ball. At each step (the timing of which follows a Poisson process at rate r), a ball is selected from the urn uniformly at random. If the selected ball is white, it is returned to the urn along with a ball of a new colour (not present already in the urn). If the selected ball is coloured, it is removed and replaced by a ball of the same colour but a darker shade.

To connect this urn process to the community-based model described above, note that selecting a white ball corresponds to the generation of a new product (which results in a ball of a new colour being added to the urn), while selecting a coloured ball and exchanging it for a darker one of that colour corresponds to the transformation of an existing product. Thus $r = N\lambda$.

We now compare the community-based model (corresponding to ρ large) to the opposite extreme, where ρ becomes small. In that latter case, the probability that there is percolation between any two entities in $\mathcal{G}$ over the interval $[0, t]$ tends to 0, and so products are only generated within entities but not shared between them. We will refer to this limiting case as

the *individual-based model.* Note that in this individual-based model, entity i_j may possibly generate a new product $p(i_j)$, or generate $p(i_j)$ and then transform it (producing $p(i_j, i_j)$ and so on (or it might not generate any new products at all). Note that, in general, $p(i_j)$ may be different from $p(i_k)$ (for $k \neq j$) (i.e., different entities may either produce or transform different products).

For the individual-based model, we have N independent samples of the above Urn model but with $r = \lambda$. By contrast, with the community-based model, we have a single sample of the above Urn model, but with $r = N\lambda$. Note that both models have the same expected number of generation events, but they have quite different dynamics, as we now describe.

Firstly, in the community-based model, there is only short-lived or transient variation among the entities, whereas in the individual-based model, the individuals diverge from each other in terms of the collections of products available to them. However, a subtler difference is that in the community model, the complexity of items is significantly higher than in the individual model, in a sense that we now make precise.

To analyse this in more detail, let X_t denote the number of steps in this Urn proces (i.e., where a ball is sampled and the urn modified) over the interval $[0, t]$. Then X_t has a Poisson distribution with mean rt. Next, let Y_t denote the number of times a white ball is selected from the urn over the interval $[0, t]$, let Z_t denote the number of times a coloured ball is selected from the urn over the time interval $[0, t]$, and let C_t denote the number of coloured balls in the urn at time t. Notice that the following two identities hold:

$$(2) \qquad\qquad X_t = Y_t + Z_t,$$

$$(3) \qquad Y_t = C_t = \text{the number of balls in the urn at time } t \text{ minus } 1.$$

Let $k_r := \lfloor 2\sqrt{rt} \rfloor$. We claim that as $r \to \infty$:

(4) $$\mathbb{P}\left(X_t \geq \frac{1}{2}rt \right) \to 1 \text{ and } \mathbb{P}(Y_t \leq k_r) \to 1,$$

and so (applying the Bonferroni inequality):

(5) $$\mathbb{P}\left(X_t/Y_t \geq \frac{1}{2}rt/k_r \right) \to 1.$$

The first limit in (4) holds because X_t has a Poisson distribution with mean rt, and so X_t/rt converges in probability to 1. For the second limit in (4), let T_{k_r} denote the time until Y_t first hits k_r, in which case, a well known identity applies:

(6) $$\mathbb{P}(Y_t \leq k_r) = \mathbb{P}(T_{k_r} \geq t).$$

Now, T_{k_r} is a sum of k_r independent exponential random variables, with means $1/r, 2/r, \ldots, k_r/r$ and variances $1^2/r^2, 2^2/r^2, \ldots, k_r^2/r^2$. Thus, T_{k_r} has expected value

$$\mathbb{E}[T_{k_r}] \sim k_r^2/2r \sim 4rt/2r = 2t,$$

and variance

$$\mathbb{V}ar[T_{k_r}] \sim k_r^3/3r^2 = 8r^{3/2}t^{3/2}/3r^2 = \Theta(r^{-1/2}) \to 0 \text{ as } r \to \infty.$$

Applying Eqn. (6) now shows that $\mathbb{P}(Y_t \leq k_r) \to 1$ as r grows, thereby justifying the second part of Eqn. (4).

We now apply Eqns. (2) and (3). These reveal that the condition $X_t/Y_t \geq k_r$ is equivalent to the condition that $Z_t/C_t \geq k_r - 1$. By the well-known 'pigeonhole principle' in combinatorics, this last inequality implies that at least one of the (C_t) coloured balls must be at least $k_r - 2$ shades darker than when that colour first appeared in the urn (because otherwise, each of the C_t coloured balls must been selected at most $k_r - 2$ times, in which case $Z_t \leq (k_r - 2)C_t$).

It now follows from Eqn. (5) (noting that the term in that equation (namely $\frac{1}{2}rt/k_r$) is of order $\sqrt{r}$)), that for r large (for fixed t), there is high probability that at least one coloured ball is present the urn that is of order $\sqrt{r}$ shades darker than it was originally was when it first appeared in the urn. Since the community-based model has $r = N\lambda$, we arrive at the following conclusion regarding the influence of the size of $\mathcal{G}$ on complexity:

Over a given period of time, some products in the community-based model have an expected complexity of order at least $\sqrt{N}$.

By contrast, for the individual-based model, we have $r = \lambda$ for each entity, and so we have N independent and identically distributed samples of a process where the maximum complexity of products across each group $\mathcal{G}$ will exhibit a lower (logarithmic) dependence on N. (Moreover, these complex products are likely to exist only in one or a few entities, rather than being shared across the group). To see this, note that the complexity of any product associated with an entity (up to time t) is bounded above by the number of generation steps for that entity, which has a Poisson distribution with mean λt, and the maximum of N independent and identically distributed Poisson random variables is known to be dominated asymptotically (with N) by a $\log(N)$ term [50].

5. Why evolution without variation and selection is possible

We have shown how entities that possess an abstract structure mathematically described by RAFs can exhibit evolution—i.e., cumulative, adaptive change—and how it is possible for this process to operate in the absence of variation and/or selection. We refer to this kind of evolutionary process as Self–Other Reorganisation (SOR) because it consists of (i) self-organization of RAF networks, and (ii) interactions amongst these networks that alter their potential for future configurations [20, 21, 24, 74, 82]. We now compare SOR to other establised

concepts in evolutionary theory: natural selection, Lamarckian evolution, the neutral theory of molecular evolution, and learning.

5.1. **SOR versus Natural Selection.** SOR involves not *competition and survival of some at the expense of others*, but *transformation of all*.[10] Like natural selection, SOR has mechanisms for preserving continuity and introducing novelty, but unlike natural selection, reproduction is a low-fidelity process, because it is the culmination of haphazard catalyzed interactions, as opposed to the accuracy afforded by copying from a code, as in a Darwinian process. The distinction between Darwinian evolution and SOR is summarised in Table 3 and illustrated in Fig. 3.

Feature	Variation–Selection	Self–Other Reorganisation (SOR)
Unit of self-replication[11]	Organism	RAF network
Preservation of continuity	Reproduction (vertical)	Communal exchange (horizontal)[12]
Generation of novelty	Mutation, recombination	Creativity, catalysis, transmission error
Self-assembly code	DNA or RNA	None
High fidelity	Yes	No
Transmission of acquired traits	No	Yes
Type	Selectionist	Lamarckian (by some standards; see above)
Evolution processes explained	Biological	Early life, horizontal gene transfer, culture

TABLE 3. Comparison between evolution through selection and evolution through Self–Other Reorganisation.

---

[10]SOR is an evolutionary process, and therefore by definition adaptive, but we note that communal interaction in and of itself is not necessarily beneficial to the recipient. Transmission of useful plasmids through horizontal exchange among bacteria or protists may be beneficial, but transmission of viruses may be damaging. Similarly, transmission of useful technologies may be beneficial to the recipient, but transmission of misinformation may be harmful.

[11]We do not use the term 'replicator' here since it is often assumed that replicator evolution necessarily involves inheritance of germ-line material.

[12]Note that communal exchange is not necessarily beneficial to the recipient. Transmission of useful plasmids through horizontal exchange among bacteria or protists may be beneficial, but transmission of viruses may be damaging. Similarly, transmission of useful technologies may be beneficial to the recipient, but transmission of misinformation may be harmful.

FIGURE 3. Comparison of (a) evolution through natural selection and (b) evolution by Self–Other Reorganisation (SOR). An X indicates that the entity was selected against (i.e., did not contribute to the next generation). Individual entities are represented by circles, with variation among them represented by different patterns. (In the example shown in (b), there is variation but there need not be, as illustrated in Fig. 2). In both (a) and (b), the darker the color the better adapted the individual (regardless of pattern). In (a), the darker color of one of the individuals in the third generation is due to mutation. Both (a) and (b) result in cumulative, adaptive change over time. However, (a) is relatively slow because it works due to differential replication of randomly generated heritable variation in a population over generations, such that some traits become more prevalent than others, with acquired changes being discarded at the end of each generation. In contrast, change in (b) is entirely due to acquired change over time.

SOR is distinctly different from a Darwinian or selectionist process. Recall that by 'self-assembly code' we mean a set of coded instructions that is: (i) *actively interpreted* through a developmental process to generate a soma, and (ii) *passively copied without interpretation* through a reproduction process to generate self-copies, i.e., new sets of self-assembly instructions that are in turn used in these two distinct ways. As mentioned above, in Darwinian evolution, interpretation of the self-assembly code in development is separate from the uninterpreted copying of the self-assembly code in reproduction. As a result, while acquired traits (those obtained during an organism's lifetime, such as a tattoo, or knowledge of a recipe) may affect the soma, and potentially affect interpretation of the self-assembly instructions, they do not affect the *content* of the self-assembly instructions. Therefore, though inherited traits (e.g., blood type) get transmitted vertically from parent to offspring by way of genes, acquired traits do not. Thus, it is the division of labour between these two ways of using the self-assembly code that is responsible for the *sine qua non* of a Darwinian process: lack of transmission of acquired traits.

Now let us turn to the earliest structures that could be said to be alive, prior to the evolution of something as complex as a DNA- or RNA-based self-assembly code. Without a self-assembly code, there were no vertically inherited traits; all change was horizontally transmitted (i.e., acquired). Therefore, the evolution of early life was not due to differential replication of heritable variation in response to selection; it was non-Darwinian [17, 80, 87]. The situation is analogous for culture. Human culture does not (currently) involve a self-assembly code that gets used in these two distinct ways, and as such, it does not exhibit this signature trait of a selectionist process. No traits are inherited by way of a self-assembly code; all change is acquired, and acquired change is transmitted. We posit that both early life and culture evolve through SOR.

A Darwinian explanation works in biology to the extent that retention of acquired change is negligible compared with retention of selected change; otherwise, the first (which can operate instantaneously) can overwhelm the second (which takes generations). Moreover, the lengthy

period we associate with biological generations, from birth through to reproductive maturity and parenthood, is in general significantly longer than the stretch of time between when an individual acquires a cultural trait (e.g., an idea) and then expresses (their own version of, or their own take on) that cultural trait. This can make the relative speed of Darwinian evolution slower still.

5.2. **SOR versus Lamarckian Evolution.** As mentioned above, the term 'Lamarckian' is widely understood to refer to the transmission of characteristics to offspring that were acquired through use or disuse during the entity's lifetime. According to interpretations of Lamarckism that specify that that Lamarckian evolution requires genetic transmission to biological offspring (e.g., [61]), SOR would not be classified as a form of Lamarckian evolution, as it is not restricted to biologically evolving entities. According to less restrictive interpretations of Lamarckism that encompass cultural as well as biological change (e.g., [61]), SOR could be considered a form of Lamarckian evolution. However, unlike Lamarckism, SOR is precise about the kind of structure required (specifically, SOR is only possible with RAF networks), and why this mechanism makes cumulative, adaptive change possible (as described above).

5.3. **SOR versus the neutral theory of molecular evolution.** The neutral theory of molecular evolution posits that much genetic variation in populations is the result of mutation and genetic drift, not selection [51, 52]. Random genetic drift in geographically isolated populations can lead to new forms appearing and others disappearing, without selection. However, drift is as likely to be disadvantageous as it is to be advantageous—it is (statistically) neutral. SOR, in contrast to neutral evolution, is not (statistically) neutral. Like natural selection, SOR is a mechanism by which adaptive change can take place.

5.4. **SOR versus Learning.** Since RAFs are central to SOR, and in cognitive RAFs the foodset consists of MRs acquired through social learning and individual learning, and learning

is a key component of SOR. These processes constitute the external component of the process, as opposed to the internal component, which involves catalyzed reactions amongst MRs resulting in new MRs that were *not* available as initial reactants, i.e., foodset-derived MRs. In the extreme, SOR could work with just one entity. In this case, the foodset consists of MRs acquired through individual learning; there are no MRs acquired through social learning. However, so long as individually learnt information can trigger cognitive 'reactions', it would be possible for some degree of cumulative, adaptive change to take place. There is a long history in psychology of viewing the assimilation of new information and the accommodation of existing knowledge in response to this new information as working hand in hand. Assimilation—the learning component—adds foodset itmes, but accommodation—an internal process—adds foodset-derived items. Thus, to the extent that they do work hand-in-hand, learning can be considered a kind of SOR. However, SOR is a broader concept than learning, because in the general case it operates on groups of entities, and can span biological generations. It is unlikely that the converse is true, i.e., that evolution by SOR could work through social learning alone, with no individual learning, because there would be nothing to socially transmit. As demonstrated in a computational model of cultural evolution, turning off social learning resulted in cumulative adaptive change (exclusively within individual agents), albeit much more slowly. However, if social learning was the *only* means of learning available, there was nothing to kickstart the process, so there was no cumulative adaptive change [12].

One might suggest that natural language is a cultural self-assembly code. However, (1) although natural language involves *encoding,* it is not a set of encoded instructions for the self-replication of natural languages, and (2) language evolution does not exhibit the signature characteristic of evolution by way of a self-assembly code: lack of transmission of acquired traits. Though some have argued that humans are biologically predisposed to achieve language, language itself is characterised by horizontal—not vertical—transmission. Therefore, language evolution is not due to the mechanism Darwin proposed: differential replication of heritable

variation in response to selection [15, 20]. Results from computational modelling suggest that to cross the Darwinian threshold from non-selectionist to selectionist evolution requires the emergence of a self-assembly code [80]. There is no evidence that language or any other component of culture has crossed this threshold, though it is not impossible that culture is moving toward a 'cultural Darwinian threshold'; in other words, it may exist in the state biological life was in before the last universal common ancestor [86].

6. Implications

The feasibility of evolution in the absence of variation and selection —i.e., evolution by SOR— and the fact that early life and cultural evolution are both promising candidates for this second form of evolution, imply that we must be cautious about applying concepts and methodologies developed in a Darwinian evolutionary context in these two domains. Since biological acquired traits are usually (though not always) discarded, and since a self-assembly code must stay intact to preserve its self-replication capacity, the joining of bifurcations in biological lineages is infrequent; thus, a phylogenetic tree correctly captures its branching structure. Speciation makes inter-lineage transfer of information relatively rare in biological organisms. By comparison, since cultural acquired traits are not discarded, and there is no cultural self-assembly code, the joining of bifurcations in cultural 'lineages' is commonplace, and thus cultural 'lineages' tend to be network-like rather than tree-like [16, 58, 77]. These distinctions become clear in phylogenetic studies. For multicellular species, phylogenetic trees calculated for different protein or RNA sequences tend to be conserved. However, phylogenetic trees constructed for different protein or RNA sequences across protists, bacteria or viruses reveal much more blending and greater network-like organisation [73]. Gene expression profiles exhibit cross-similarity and greater convergent evolution as well. Since cultural relatedness frequently arises through horizontal (inter-lineage) transmission, there is extensive blending of knowledge from

different sources. Extensive horizontal transmission gives a bushy, reticulated appearance to a phylogenetic tree, which is misleading, because it implies not just chronology but ancestry.

Another (related) problem that arises when methods developed for selectionist evolutionary processes are applied to culture is due to convergent evolution, in which similar forms arise independently because they are alternative solutions within similar design constraints. Examples include (i) the body shape and structure similarity between the Tasmanian tiger and the fox, (ii) wasp-imitating hover flies and (iii), the origin of basic and more complex brain structure across the tree of life [62, 89]. Because biological organisms must solve many problems (reproduction, locomotion, digestion, etc.), the probability that a species will be mis-categorised because of convergent evolution (i.e., on the basis of how it solves any one problem) is low. Cultural artifacts, on the other hand, are generally constructed with a single use in mind (though artifacts developed for use in one context may be used to solve other problems; for example, a screwdriver may be used to open a can of paint.) Thus, for cultural outputs, the probability of mis-categorisation arising through the assumption that similarity reflects homology is significantly higher. Therefore, the cost of proceeding as if a Darwinian framework were applicable to culture when it is not is high. This problem does not arise with SOR because it does not assume that superficially similar artifacts are homologous. Indeed, a SOR take on culture suggests that what is evolving is the structures of the conceptual networks that generate artifacts, not the artifacts themselves. Some have claimed that in practice this does not invalidate a phylogenetic approach to culture [32]. However, such conclusions come from analyses of datasets that involve little horizontal transmission (indeed, the creative blends that are the source of cultural novelty are often treated as 'outliers' and are intentionally discarded from analysis).

Such considerations have led some to develop network-based models of cultural evolution [4, 10, 12, 24, 23, 53, 58, 79]. This body of research suggests that horizontal transmission can significantly alter the pattern of relatedness. For example, a network-based analysis of Baltic

psaltery data that incorporated not just superficial physical attributes but also abstract conceptual attributes (such as markings indicative of sacred symbolic imagery), it was possible to resolve ambiguities arising from a phylogenetic analysis and generate a lineage more consistent with other historical data [79]. Horizontal cultural transmission may involve change in superficial features despite a preservation of deep structure, as occurs in metaphor [54], analogy [29, 38], and cross-domain transfer, in which a source from one domain (e.g., music) inspires or influences a creative work in another (e.g., painting) [67, 69]. This kind of complexity and hierarchical structure cannot be captured without taking a network approach to cultural evolution, which provides further support for the theory that culture evolves through SOR.

Interestingly, similar issues arise with the simplest life forms. Because of phenomena such as mutualism, lineage reticulation (due to horizontal gene transfer and allopolyploidy—the combining the genomes of different parental species), certain traits evolve with astonishing speed, thereby diminishing the continuity and distinctiveness of species [66, 87]. Indeed, the stability of genetic information is so compromised that sequencing three *Escherichia coli* genomes revealed that fewer than 40% of the genes were common to all three [84]. As a result, the boundaries between many prokaryote species are fuzzy, and exhibit reticulate patterns of evolution, thus calling into question the appropriateness of the notion of the "tree of life" [36, 46, 60]. The limitations of Darwinism as an explanation of the forms and dynamics of living things is increasingly recognised, while the role of epigenetic processes has become increasingly appreciated. Nevertheless, because such phenomena are much less present in biological evolution than cultural evolution, natural selection provides a reasonable approximation.

7. CONCLUDING COMMENTS

Using RAF networks, this paper demonstrates by way of example, or 'existence proof' that evolution is possible in the absence of variation, selection, and competition. We refer to this kind of evolution as Self–Other Reorganisation (SOR) because it occurs not through competitive

exclusion such that only the fittest reproduce, but through the assimilation, restructuring, and exchange of components. SOR is a primitive form of evolution that can generate change quickly since it does not require the discarding of acquired traits, and it can take place in the absence of birth or death. Because it does not use precisely coded self-assembly instructions, it is more haphazard than natural or artificial selection, but sufficient for cumulative, adaptive change.

RAFs have proven useful in two areas of research that might appear to be quite distinct, but that both involve networks and evolutionary processes. In the OOL, the underlying reaction system is a biochemical reaction network, and in the OOC it is a conceptual network. Since cultural evolution lacks a self-assembly code, and lacks the signature characteristic of Darwinian evolution—discarding of acquired traits at the end of a generation—it seems reasonable to investigate the extent to which the OOL and the OOC share a common theoretical framework. This is consistent with the proposal that the entities that evolve through culture are not discrete artifacts or units of behaviour such as songs, jokes or memes, but the minds that generate them [15], which have a structure that is self-organising and self-maintaining [33, 59], and therefore lends itself to the application of RAF formalisms [25, 27, 26]. We suggest that SOR may be operative during the early stage of any evolutionary process, that it was responsible for both the origin and early evolution of both organic life and human culture, and that RAFs provide a means of modeling evolution through SOR. The approach offers an established formal framework for integrating biological, cultural, and cognitive proceses, and embedding this synthesis in the study of self-organizing structures and their role in evolutionary processes.

SOR is a broader concept than learning, as it can produce adaptive change in groups of entities, and it can span biological generations. Competition may occur in SOR, but unlike selectionist evolution it is not essential. Since SOR does not require competition or variation, it may increase homogeneity among members of a culturally evolving group of individuals by increasing the amount of shared knowledge and experience among group members. It may

thereby play an important role in fostering group identity, cohesion, and cooperation [82] (see also [1]).

References

[1] C. Andersson and P. Törnberg, *Toward a macroevolutionary theory of human evolution: The social protocell*, Biological Theory, 14 (2019), pp. 86–102, https://doi.org/10.1007/s13752-018-0313-y.

[2] D. Baum, *The origin and early evolution of life in chemical composition space*, Journal of Theoretical Biology, 456 (2018), pp. 295–304.

[3] R. Boyd and P. Richerson, *Culture and the evolutionary process*, University of Chicago Press, Chicago, IL, 1988.

[4] A. Buskell, M. Enquist, and F. Jansson, *A systems approach to cultural evolution*, Palgrave Communications, 5 (2019), pp. 1–15.

[5] K. R. Cabell and J. Valsiner, *The catalyzing mind: Beyond models of causality (Annals of Theoretical Psychology, Volume 11)*, Springer, Berlin, 2013, https://doi.org/10.1007/978-1-4614-8821-7.

[6] L. L. Cavalli-Sforza and M. W. Feldman, *Cultural transmission and evolution: A quantitative approach*, Princeton University Press, Princeton, NJ, 1981.

[7] R. Cazzolla Gatti, B. Fath, W. Hordijk, S. Kauffman, and R. Ulanowicz, *Niche emergence as an autocatalytic process in the evolution of ecosystems*, Journal of Theoretical Biology, 454 (2018), pp. 110–117, https://doi.org/10.1016/j.jtbi.2018.05.038.

[8] A. Cornish-Bowden and M. L. Cárdenas, *Life before LUCA*, Journal of Theoretical Biology, 434 (2017), pp. 68–74.

[9] C. Darwin, *The origin of species*, PF Collier & Son, New York, 1909.

[10] M. Enquist, S. Ghirlanda, and K. Eriksson, *Modelling the evolution and diversity of cumulative culture*, Philosophical Transactions of the Royal Society B: Biological Sciences, 366 (2011), pp. 412–423, https://doi.org/10.1098/rstb.2010.0132.

[11] P. Erdös and A. Rényi, *On the evolution of random graphs*, Publication of the Mathematical Institute of the Hungarian Academy of Sciences, 5 (1960), pp. 17–61.

[12] L. Gabora, *Meme and variations: A computer model of cultural evolution*, in 1993 Lectures in Complex Systems, A. Goel, C. Seifert, and C. Freska, eds., Addison-Wesley, Boston MA, 1995, pp. 471–486.

[13] L. Gabora, *Autocatalytic closure in a cognitive system: A tentative scenario for the origin of culture*, Psycoloquy, 9 (1998), pp. [adap–org/9901002].

[14] L. Gabora, *Conceptual closure: How memories are woven into an interconnected worldview.*, in Closure: Emergent Organizations and their Dynamics, G. Van de Vijver and J. Chandler, eds., no. 901 in Annual Review Series, Annals of the New York Academy of Sciences, 2000, pp. 42–53, `https://doi.org/10.1111/j.1749-6632.2000.tb06264.x.`

[15] L. Gabora, *Ideas are not replicators but minds are*, Biology and Philosophy, 19 (2004), pp. 127–143.

[16] L. Gabora, *The fate of evolutionary archaeology: Survival or extinction?*, World Archaeology, 38 (2006), pp. 690–696.

[17] L. Gabora, *Self-other organization: Why early life did not evolve through natural selection*, Journal of Theoretical Biology, 241 (2006), pp. 443–450.

[18] L. Gabora, *Why the creative process is not darwinian*, Creativity Research Journal, 19 (2007), pp. 361–365.

[19] L. Gabora, *Five clarifications about cultural evolution*, Journal of Cognition and Culture, 11 (2011), pp. 61–83.

[20] L. Gabora, *An evolutionary framework for culture: Selectionism versus communal exchange*, Physics of Life Reviews, 10 (2013), pp. 117–145, `https://doi.org/10.1016/j.plrev.2013.03.006.`

[21] L. Gabora, *From deep learning to deep reflection: Toward an appreciation of the integrated nature of cognition and a viable theoretical framework for cultural evolution*, in Proceedings of the 2019 Annual Meeting of the Cognitive Science Society, L. Nadel and D. D. Stein, eds., Cognitive Science Society, Austin, TX, 2019, pp. 1801–1807.

[22] L. Gabora and N. Beckage, *Modeling cognitive development with reflexively autocatalytic networks*, Topics in Cognitive Science, (in prep).

[23] L. Gabora, S. Leijnen, T. Veloz, and C. Lipo, *A non-phylogenetic conceptual network architecture for organizing classes of material artifacts into cultural lineages*, in Proceedings of the 33rd annual meeting of the Cognitive Science Society, L. Carlson, C. Hőlscher, and T. F. Shipley, eds., Cognitive Science Society, Psychology Press, 2011, pp. 2923–2928.

[24] L. Gabora and C. Smith, *Two cognitive transitions underlying the capacity for cultural evolution*, Journal of Anthropological Science, 96 (2018), pp. 27–52, `https://doi.org/10.4436/jass.96008.`

[25] L. Gabora and M. Steel, *Autocatalytic networks in cognition and the origin of culture*, Journal of Theoretical Biology, 431 (2017), pp. 87–95, `https://doi.org/10.1016/j.jtbi.2017.07.022`.

[26] L. Gabora and M. Steel, *A model of the transition to behavioral and cognitive modernity using reflexively autocatalytic networks*, Proceedings of the Royal Society Interface, 17 (2020), p. 20200545, `https://doi.org/http://doi.org/10.1098/rsif.2020.0545`.

[27] L. Gabora and M. Steel, *Modeling a cognitive transition at the origin of cultural evolution using autocatalytic networks*, Cognitive Science, 44 (2020).

[28] L. Gabora and S. Tseng, *The social benefits of balancing creativity and imitation: Evidence from an agent-based model*, Psychology of Aesthetics, Creativity, and the Arts, 11 (2017), pp. 457–473.

[29] D. Gentner, *Structure-mapping: A theoretical framework for analogy*, Cognitive Science, 7 (1983), pp. 155–170, `https://doi.org/10.3389/fpsyg.2019.01426`.

[30] N. Goldenfeld, T. Biancalani, and F. Jafarpour, *Universal biology and the statistical mechanics of early life*, Philosophical Transactions of the Royal Society A, 375 (2017), p. 20160341, `https://doi.org/2017`.

[31] S. J. Gould, *The structure of evolutionary theory*, Belknap Press of Harvard University Press, Cambridge, MA, 2002.

[32] S. J. Greenhill, T. E. Currie, and R. D. Gray, *Does horizontal transmission invalidate cultural phylogenies?*, Proceedings of the Royal Society B: Biological Sciences, 276 (2009), pp. 2299–2306.

[33] A. G. Greenwald, M. R. Banaji, L. A. Rudman, S. D. Farnham, B. A. Nosek, and D. S. Mellott, *A unified theory of implicit attitudes, stereotypes, self-esteem, and self-concept*, Psychological Review, 109 (2002), pp. 3–25.

[34] S. Harnad, *The symbol grounding problem*, Physica D, 42 (1990), pp. 335—-346.

[35] J. Henrich and R. McElreath, *Dual-inheritance theory: The evolution of human cultural capacities and cultural evolution*, in Oxford Handbook of Evolutionary Psychology, Oxford University Press, 2007, pp. 741–750.

[36] E. Hilario and J. P. Gogarten, *Horizontal transfer of atpasegenes: The tree of life becomes a net of life*, Biosystems, 31 (1993), pp. 111–119.

[37] J. H. Holland, *Adaptation in natural and artificial systems: An introductory analysis with applications to biology, control, and artificial intelligence*, MIT Press, Urbana, 1992.

[38] K. J. HOLYOAK AND P. THAGARD, *Mental leaps: Analogy in creative thought*, MIT Press, Cambridge, MA, 1996.

[39] W. HORDIJK, J. HEIN, AND M. STEEL, *Autocatalytic sets and the origin of life*, Entropy, 12 (2010), pp. 1733–1742, https://doi.org/10.3390/e12071733.

[40] W. HORDIJK, S. A. KAUFFMAN, AND M. STEEL, *Required levels of catalysis for emergence of autocatalytic sets in models of chemical reaction systems*, International Journal of Molecular Science, 12 (2011), pp. 3085–3101, https://doi.org/10.3390/ijms12053085.

[41] W. HORDIJK AND S. M., *A formal model of autocatalytic sets emerging in an rna replicatior system*, Journal of the Systems Chemistry, 4:3 (2013), https://doi.org/10.1186/1759-2208-4-3.

[42] W. HORDIJK AND M. STEEL, *Detecting autocatalytic, self-sustaining sets in chemical reaction systems*, Journal of Theoretical Biology, 227 (2004), pp. 451–461, https://doi.org/10.1016/j.jtbi.2003.11.020.

[43] W. HORDIJK AND M. STEEL, *Autocatalytic sets and boundaries*, J. Syst. Chem., 6:1 (2015).

[44] W. HORDIJK AND M. STEEL, *Chasing the tail: The emergence of autocatalytic networks*, Biosystems, 152 (2016), pp. 1–10, https://doi.org/10.1016/j.biosystems.2016.12.002.

[45] W. HORDIJK, M. STEEL, AND P. DITTRICH, *Autocatalytic sets and chemical organizations: Modeling self-sustaining reaction networks at the origin of life*, New Journal of Physics, 20 (2018), p. 015011.

[46] A. L. HUGHES, *The origin of adaptive phenotypes*, Proceedings of the National Academy of Sciences USA, 105 (2008), pp. 13193–13194.

[47] S. A. KAUFFMAN, *Autocatalytic sets of proteins*, Journal of Theoretical Biology, 119 (1986), pp. 1–24, https://doi.org/10.3390/ijms12053085.

[48] S. A. KAUFFMAN, *The origins of order*, Oxford University Press, 1993.

[49] P. KILLEEN, *The non-Darwinian evolution of behavers and behaviors*, Behavioural Processes, 161 (2019), pp. 45–53.

[50] A. C. KIMBER, *A note on Poisson maxima*, Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete, 63 (1983), pp. 551–552.

[51] M. KIMURA, *Evolutionary rate at the molecular level*, Nature, 217 (1968), pp. 624—-626, https://doi.org/doi:10.1038/217624a0.PMID5637732.

[52] J. L. KING AND T. H. JUKES, *Non-darwinian evolution*, Science, 164 (1969), pp. 788–98, https://doi.org/doi:10.1126/science.164.3881.788.PMID5767777.

[53] S. KIRBY, *Culture and biology in the origins of linguistic structure*, Psychonomic Bulletin & Review, 24 (2017), pp. 118–137.

[54] G. LAKOFF, *The contemporary theory of metaphor*, in Metaphor and thought (2nd ed), A. Ortony, ed., Cambridge University Press, Cambridge, UK, 1993, pp. 202–252.

[55] D. LANCET, R. ZIDOVETZKI, AND O. MARKOVITCH, *Coevolution of compositional protocells and their environment*, Journal of the Royal Society Interface, 15:20180159 (2018).

[56] C. G. LANGTON, *Artificial life*, in Lectures in Complex Systems, Vol. IV, L. Nadel and D. Stein, eds., MIT Press, Reading, MA, 1992, pp. 189–242.

[57] R. LEWONTIN, *The units of selection*, Annual Review of Ecology and Systematics, 61 (1970), pp. 1–18.

[58] C. P. LIPO, *The resolution of cultural phylogenies using graphs*, in Mapping our Ancestors, C. P. Lipo, M. J. O'Brien, M. Collard, and S. J. Shena, eds., vol. 5, Aldine de Gruyter, New York, 2005, pp. 89–107.

[59] H. MATURANA AND F. VARELA, *Autopoiesis and cognition: The realization of the living*, in Boston Studies in the Philosophy of Science, R. S. Cohen and M. W. Wartofsky, eds., vol. 42, Reidel, Dordecht, 1973.

[60] J. O. MCINERNEY, D. PISANI, E. BAPTESTE, AND M. J. O'CONNELL, *The public goods hypothesis for the evolution of life on earth*, Biology Direct, 6 (2011).

[61] A. MESOUDI, A. WHITEN, AND L. K. N., *Is human cultural evolution Darwinian? Evidence reviewed from the perspective of the origin of species*, Evolution, 58 (2004), pp. 1–11.

[62] L. MOROZ AND A. KOHN, *Independent origins of neurons and synapses: insights from ctenophores.*, Philosophical Transactions of the Royal Soceity B (Biol. Sci.), 371(1685) (2016), p. 20150041, `https://doi.org/10.1098/rstb.2015.0041`.

[63] E. MOSSEL AND M. STEEL, *Random biochemical networks and the probability of self-sustaining autocatalysis*, Journal of Theoretical Biology, 233 (2005), pp. 327–336, `https://doi.org/10.1016/j.jtbi.2004.10.011`.

[64] G. B. MÜLLER, *Why an extended evolutionary synthesis is necessary*, Interface Focus, 7 (2017), p. 20170015.

[65] M. MUTHUKRISHNA, M. DOEBELI, M. CHUDEK, AND J. HENRICH, *The cultural brain hypothesis: How culture drives brain expansion, sociality, and life history*, PLoS Computational Biology, 14 (2018), p. e1006504, `https://doi.org/10.1371/journal.pcbi.1006504`.

[66] G. J. P. AND J. P. TOWNSEND, *Horizontal gene transfer, genome innovation and evolution*, Nature Reviews, 3 (2005), pp. 679–687.

[67] A. Ranjan, L. Gabora, and B. O'Connor, *Evidence that cross-domain re-interpretations of creative ideas are recognizable*, in Proceedings of the Association for the Advancement of Artificial Intelligence (AAAI) Spring Symposium (Creativity and Cognitive Development: A Perspective from Artificial Creativity, Developmental Artificial Intelligence, and Robotics), AAAI Press, Menlo Park, USA, 2013.

[68] P. Richerson and R. Boyd, *A dual inheritance model of the human evolutionary process i: Basic postulates and a simple model*, Journal of Social and Biological Structures, 1 (1978), pp. 127–154.

[69] V. Scotney, S. Weissmeyer, N. Carbert, and L. Gabora, *The ubiquity of cross-domain thinking in the early phase of the creative process*, Frontiers in Psychology, 10 (2019), p. 1426, `https://doi.org/10.3389/fpsyg.2019.01426`.

[70] D. Segré, D. Ben-Eli, and D. Lancet, *Compositional genomes: Prebiotic information transfer in mutally catalytic noncovalent assemblies*, Proceedings of the Royal Society B, 97 (200), pp. 4112–4117.

[71] R. Semon, *The mneme*, Allen and Unwin, London, 1921.

[72] B. Shenhav, A. Oz, and D. Lancet, *Coevolution of compositional protocells and their environment*, Philosophical Transactions of the Royal Society B, 362 (2007), pp. 1813–1819.

[73] M. Shi, X. Lin, and J. Tian, *Redefining the invertebrate rna virosphere*, Nature, 540 (2016), pp. 539–543, `https://doi.org/10.1038/nature20167`.

[74] C. Smith, L. Gabora, and W. Gardner-O'Kearny, *The extended evolutionary synthesis facilitates evolutionary models of culture change*, Cliodynamics: The Journal of Quantitative History and Cultural Evolution, 9 (2018), pp. 84–107.

[75] M. Steel, *The emergence of a self-catalyzing structure in abstract origin-of-life models*, Applied Mathematics Letters, 13 (2000), pp. 91–95.

[76] M. Steel, W. Hordijk, and J. C. Xavier, *Autocatalytic networks in biology: Structural theory and algorithms*, Journal of the Royal Society Interface, 16 (2019), p. rsif.2018.0808, `https://doi.org/10.1098/rsif.2018.0808`.

[77] I. Temkin and N. Eldredge, *Phylogenetics and material cultural evolution*, Current Anthropology, 48 (2007), pp. 146–153.

[78] V. Vasas, C. Fernando, M. Santos, S. Kauffman, and E. Szathmáry, *Evolution before genes*, Biology Direct, 7 (2012).

[79] T. VELOZ, I. TEMPKIN, AND L. GABORA, *A conceptual network-based approach to inferring cultural phylogenies*, in Proceedings of the 34th annual meeting of the Cognitive Science Society, N. Miyake, D. Peebles, and R. P. Cooper, eds., Cognitive Science Society, Austin TX, 2012, pp. 2487–2492.

[80] K. VETSIGIAN, C. WOESE, AND N. GOLDENFELD, *Collective evolution and the genetic code*, Proceedings of the National Academy of Sciences USA, 103 (2006), pp. 10696–10701.

[81] J. VON NEUMANN, *Theory of Self-Replicating Automata*, University of Illinois Press, Urbana, 1966.

[82] B. VOORHEES, D. READ, AND L. GABORA, *Identity, kinship, and the evolution of cooperation*, Current Anthropology, 61 (2020), pp. 194–218.

[83] S. VOSNIADOU AND W. F. BREWER, *Mental models of the earth: A study of conceptual change in childhood*, Cognitive Psychology, 24 (1992), pp. 535–585.

[84] R. A. WELCH, *Extensive mosaic structure revealed by the complete genome sequence of uropathogenic Escherichia coli*, Proceedings of the National Academy of Sciences USA, 99 (2002), pp. 17020–17024.

[85] A. WHITEN, *A second inheritance system: the extension of biology through culture*, Interface Focus, 7 (2017), p. 20160142.

[86] C. WOESE, *The universal ancestor*, Proceedings of the National Academy of Sciences, 95 (1998), pp. 6854–6859.

[87] C. R. WOESE, *On the evolution of cells*, Proceedings of the National Academy Science, 99 (2002), pp. 8742–8747.

[88] J. C. XAVIER, W. HORDIJK, S. KAUFFMAN, M. STEEL, AND W. F. MARTIN, *Autocatalytic chemical networks at the origin of metabolism*, Proceedings of the Royal Society of London. Series B: Biological Sciences, 287 (2020), p. 20192377.

[89] M. YOSHIDA, A. OGURA, K. IKEO, S. SHIGENO, T. MORITAKI, G. C. WINTERS, A. B. KOHN, AND L. L. MOROZ, *Molecular evidence for convergence and parallelism in evolution of complex brains of cephalopod molluscs: Insights from visual systems*, Integrative and Comparative Biology, 55(6) (2015), pp. 1070–1083, https://doi.org/10.1093/icb/icv049.