Feedback from retinal ganglion cells to the inner retina

Authors: Anastasiia Vlasiuk[1,2] and Hiroki Asari[*1,3]

Affiliations:

1. European Molecular Biology Laboratory (EMBL), Epigenetics and Neurobiology Unit, Rome, 00015, Italy

2. Candidate for Joint PhD degree from EMBL and Heidelberg University, Faculty of Biosciences, Heidelberg, 69117, Germany

3. Previous affiliations: Department of Molecular and Cellular Biology and Center for Brain Science, Harvard University, Cambridge, Massachusetts, 02138, USA; Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, California, 91125, USA

Corresponding author / Lead contact: Hiroki Asari, E-mail: asari@embl.it

Acknowledgments: We gratefully acknowledge Markus Meister for assistance with experiments. We also thank all the members of the Asari lab at EMBL Rome as well as Cornelius Gross and Santiago Rompani for many useful discussions.

Grants: This work was supported by Gosney Postdoctoral Fellowship from CalTech and research grants from EMBL (H.A.).

Author contributions: A.V. and H.A. designed the study, performed the experiments, analyzed the results, and wrote the manuscript.

Competing interests: The authors declare no competing financial interests.

Summary

Retinal ganglion cells (RGCs) are thought to be strictly postsynaptic within the retina. They carry visual signals from the eye to the brain, but do not make chemical synapses onto other retinal neurons. Nevertheless, they form gap junctions with other RGCs and amacrine cells, providing possibilities for RGC signals to feed back into the inner retina. Here we identified such feedback circuitry in the salamander and mouse retinas. First, using biologically inspired circuit models, we found mutual inhibition among RGCs of the same type. We then experimentally determined that this effect is mediated by gap junctions with amacrine cells. Finally, we found that this negative feedback lowers RGC visual response gain without affecting feature selectivity. The principal neurons of the retina therefore participate in a recurrent circuit much as those in other brain areas, not being a mere collector of retinal signals, but are actively involved in visual computations.

Keywords

Retinal ganglion cell, gap junction, feedback, circuit model, gain control.

Acronyms

GABA, γ-aminobutyric acid; GCM, ganglion cell module; LNFDSCNF, linear-nonlinear-feedback-delayed-sum-coupling-nonlinear-feedback; MFA, meclofenamic acid; PSTH, peri-stimulus time histogram; PTX, picrotoxin; RGC, retinal ganglion cell; STR, strychnine.

Introduction

Many brain circuits involve recurrent connections between principal neurons (Ito, 2000; Maass and Markram, 2002; Wang, 2008). This allows for feedback control that can shape the dynamics of neuronal activity, and is suggested to play critical roles in both local and global circuit functions, ranging from sensory information processing to decision making (Carandini and Heeger, 2012).

In the retina, the principal neurons—i.e., ganglion cells—are often thought to be strictly output elements (Gollisch and Meister, 2010; Masland, 2012). Previous studies have thus focused on characterizing how ganglion cells integrate signals from inner retinal neurons distributed in space and across different cell-types via chemical synapses (Berry et al., 1999; Wässle, 2004; Werblin, 2011). As a result, two major mechanisms have been found that are important to control ganglion cell activity: feedforward signaling from bipolar cells to ganglion cells via amacrine cells (Chen et al., 2010; Roska et al., 2006, 2000), and feedback signaling between bipolar cells and amacrine cells (Dong and Werblin, 1998; Nirenberg and Meister, 1997; Tachibana and Kaneko, 1988).

Ganglion cells are, however, part of extensive gap junction networks in the retina (Bloomfield and Völgyi, 2009; Cook and Becker, 1995; Puller et al., 2020). While retinal neurons are generally coupled among those of the same type, ganglion cells also form electrical synapses with amacrine cells (Nath and Schwartz, 2017; Roy et al., 2017; Völgyi et al., 2009). Therefore, ganglion cells could in principle signal back to the inner retina to modify the synaptic inputs they receive. Importantly, such feedback control of ganglion cell activity can be both excitatory and inhibitory. On the one hand, excitatory effects are expected from couplings through gap junction networks as they allow for electrical and bidirectional interactions among coupled neurons (Bloomfield and Völgyi, 2009). Different architectures of the gap junction networks are indeed responsible for distinct firing patterns among ganglion cells over multiple time-scales. For example, direct couplings between ganglion cells underlie their nearly synchronous activity within a few milliseconds, whereas indirect couplings via amacrine cells are suggested to mediate more broadly correlated activity of ganglion cells over tens to hundreds of milliseconds (Brivanlou et al., 1998; Mastronarde, 1983; Meister et al., 1995; Nath and Schwartz, 2017;

Puller et al., 2020; Roy et al., 2017; Trong and Rieke, 2008; Völgyi et al., 2013). On the other hand, inhibitory effects will be obtained if ganglion cells send feedback signals to amacrine cells through electrical synapses, while the amacrine cells in turn inhibit the ganglion cells or presynaptic bipolar cells through chemical synapses (Greschner et al., 2016; Kenyon and Marshak, 1998; Sakai and Naka, 1988, 1990). The computational power of ganglion cells should then be greater than is commonly thought, allowing for more complex and diverse visual processing. It remains unclear, though, if these feedback pathways are truly functional in the retina, and how they contribute to visual information processing in the retina.

Here we characterized such feedback circuitry in the salamander and mouse retinas by combining computational and experimental approaches. Using a circuit model with biologically interpretable elements (Real et al., 2017), we first show that ganglion cells exchange signals with other ganglion cells in two distinct ways: faster, enhancing signals from proximal cells and slower, suppressing signals from distal cells. To experimentally test this model prediction, we perturbed the activity of ganglion cells in the dark by electrically driving antidromic spikes at the tip of the optic nerve emerging from an isolated retina, while simultaneously monitoring the consequences of the perturbation by multi-electrode array recordings. Consistent with our model's prediction, the optic nerve stimulation produced a short period of enhanced firing (for tens of milliseconds), followed by a suppression of firing on longer time scales (for hundreds of milliseconds). The slower suppression was largely eliminated by pharmacologically blocking electrical or inhibitory synaptic transmission. Because inhibition is mediated by amacrine cells in the inner retina (Masland, 2012), these results support that ganglion cells form negative feedback circuits with amacrine cells via gap junctions (Greschner et al., 2016; Kenyon and Marshak, 1998; Sakai and Naka, 1988, 1990). Finally, by pairing the nerve shock with visual stimulation, we found that the negative feedback modulates the visual response gain but not the stimulus feature selectivity of ganglion cells. Therefore, we conclude that ganglion cells can actively control their own spiking activity and that of other ganglion cells via recurrent connections with amacrine cells, providing additional mechanism for adaptive gain control in the retina.

Results

Circuit models with electrical couplings recapitulate retinal ganglion cell visual responses better than non-coupled models

How do ganglion cells interact with each other in the retina? To address this question, we started with a modeling approach that includes subthreshold effects of electrical couplings between retinal ganglion cells (Bloomfield and Völgyi, 2009; Cook and Becker, 1995). Here we followed the cascade model framework and extended the neural circuit models of Real et al. (2017): specifically, the full "linear-nonlinear-feedback-delayed-sum-nonlinear-feedback" (LNFDSNF) model as well as the reduced "linear-nonlinear-sum-nonlinear" (LNSN) model were employed as a baseline non-coupled model. Briefly, the first LNF stages collectively work as spatial subunits that correspond to bipolar cells and their upstream circuitry; the middle D stage is a delay in lateral propagation introduced by amacrine cells; and the last SNF stages represent the spatial summation by a target ganglion cell (see Methods for details).

Our coupled models (LNFDSCNF, Supplementary Figure 1A; and LNSCN, Figure 1A) were then built by incorporating the following two parameters into the non-coupled models. The first parameter is the net coupling strength that represents the effect of both direct coupling between ganglion cells and indirect pathways via amacrine cells (a_k in Eq.(2) in Methods). A positive value is expected for a directly coupled pair of ganglion cells (Brivanlou et al., 1998), where the coupling strength largely depends on the number of electrical synapses between the cells and their conductance (Bloomfield and Völgyi, 2009). In contrast, a negative value is expected for an indirectly coupled cell pair, where signals from one ganglion cell transmit to an amacrine cell via electrical synapses and the amacrine cell in turn inhibits the other ganglion cell via chemical synapses (Greschner et al., 2016; Kenyon and Marshak, 1998; Sakai and Naka, 1988, 1990). The second parameter is the net latency for signals to arrive from the neighboring ganglion cell to the modelled cell (l_k in Eq.(1) in Methods). A short latency is expected for a directly coupled pair, whereas a long latency is expected for an indirectly coupled pair. Importantly, here we kept the model size minimum to achieve reliable data fitting, with only two extra free parameters for each surrounding ganglion cell to describe the net effect of its action

potentials on a target cell. Nevertheless, our coupled models are flexible enough to account for the overall effect of the interactions between retinal ganglion cells through the entire circuitry of the inner retina.

When fitted, these coupled and non-coupled models were both able to approximate the firing patterns of salamander (Asari and Meister, 2012, 2014; Real et al., 2017) and mouse (Lefebvre et al., 2008) retinal ganglion cells in response to white-noise visual stimuli consisting of randomly flickering bars (e.g., Figure 1B–D; see Methods for details). The non-coupled models, however, occasionally predicted visual responses at times when they should not, while the coupled models often correctly suppressed such false-positive responses by the firing of negatively coupled cells that have the same response polarity as the target cell (Figure 1C,D). This led to a small but statistically significant improvement of the coupled models in recapitulating the ganglion cell visual responses compared to the non-coupled models (R^2=0.314±0.091 versus 0.297±0.087, p<0.001, for salamander cells, Figure 2A; R^2=0.215±0.088 versus 0.208±0.087, p<0.001, for mouse cells, Figure 2C; mean ± standard deviation of the coefficient of determination (Eq.(3) in Methods) for the reduced models, paired t-test; see also Supplementary Figure 1 for the full models). In contrast, surrounding cells of the opposite response polarities contributed much less to the performance of the coupled models (R^2=0.322±0.095 versus 0.309±0.0100, p<0.001, for salamander cells, Figure 2B; R^2=0.209±0.090 versus 0.208±0.087, p>0.5, for mouse cells, Figure 2D). Importantly, to avoid confounding effects of common visual inputs in the coupled model, we considered only those surrounding cells that have low spike correlations—hence virtually no receptive field overlap—with the target cell (Supplementary Figure 2A,B). Furthermore, we confirmed the convergence and robustness of the model fitting by the perturbation analysis where we reoptimized the models after resetting the coupling parameters (Supplementary Figure 2C,D; see Methods for details). Therefore, as suggested by anatomical studies (Völgyi et al., 2009), our circuit model analysis supports the presence and solid contributions of couplings between retinal ganglion cells, especially those of the same response polarity, to shaping their visual response dynamics.

Circuit model analysis predicts both positive and negative couplings between retinal ganglion cells

We next analyzed the model parameters to gain insights into the retinal circuits underlying ganglion cell couplings and to derive experimentally testable predictions. The data sets from ON and OFF cells were combined here as they showed the same trend (Figure 3 and Supplementary Figure 3).

First, we found both positive and negative couplings between ganglion cells of the same response polarities (Figure 3A,B and Supplementary Figure 3A,B for salamander retinas; Figure 3E,F and Supplementary Figure 3E,F for mouse retinas). These couplings were frequently reciprocal, either mutually positive or negative between given cell pairs (Figure 3G,I and Supplementary Figure 3G,I). This indicates that a single ganglion cell can be involved in multiple feedback pathways, imposing distinct effects on different surrounding cells. In contrast, positive couplings dominated between the cell pairs with different response polarities (Figure 3C,D,H and Supplementary Figure 3C,D,H for salamander cells; mouse data not analyzed because the coupled model performance did not improve significantly; Figure 2D and Supplementary Figure 1E). Consistently, such a cross-talk between the ON and OFF channels via gap junctions has been reported recently (Cooler and Schwartz, 2021). Taken together, negative couplings are suggested to be formed exclusively between ganglion cells of the same response polarities.

Second, positive couplings were on average found between cell pairs at a shorter distance and had a shorter latency than negative couplings (Figure 3A–F and Supplementary Figure 3A–F). Given that the diameter of ganglion cell dendritic fields is around 0.3 mm (Völgyi et al., 2009; Zhang and Wu, 2010), the distance of the positive couplings (0.31 ± 0.14 and 0.33 ± 0.25 mm for salamander and mouse cell pairs, respectively, median $\pm$ interquartile range from the reduced models) is consistent with direct interactions between ganglion cells, although this does not exclude possible contributions of indirect electrical couplings via amacrine cells (Roy et al., 2017). In contrast, the distance of the negative couplings (0.40 ± 0.26 and 0.78 ± 0.74 mm, respectively) exceeds what one expects for direct coupling. The circuit model thus predicts that faster enhancing effects between proximal ganglion cells can arise from

direct coupling, whereas slower suppressing effects between distal cells should be mediated by indirect coupling, presumably via inhibitory amacrine cells in the inner retina (Greschner et al., 2016; Sakai and Naka, 1988, 1990). The prediction on the positive couplings is well supported by previous studies (Brivanlou et al., 1998; Cocco et al., 2009; Nath and Schwartz, 2017; Pillow et al., 2008; Puller et al., 2020), but direct experimental evidence is still lacking on the slow negative coupling between ganglion cells over a long distance.

Third, there was no marked difference in the common parameters between coupled and non-coupled models (Supplementary Figure 4). Consistently, the outputs of these two models showed nearly identical dynamics (e.g., Figure 1D), suggesting that the ganglion cell couplings have little effect on the cell's stimulus feature selectivity. Importantly, however, non-desired response peaks were present more frequently in the output of the non-coupled models than that of the coupled models (e.g., Figure 1D). This indicates that the ganglion cell couplings have some transient effects on the response gain of a target cell, suppressing such false-positive responses that were present otherwise.

In summary, our circuit model analysis predicts that, besides bidirectional excitation via gap junction networks (Bloomfield and Völgyi, 2009), retinal ganglion cells of the same response polarity can have reciprocal indirect inhibition between each other for a transient gain control from outside their receptive fields.

Optic nerve stimulation imposes diverse effects on ganglion cell activity

To directly test the model prediction of an indirect inhibition between retinal ganglion cells (Figures 1–3 and Supplementary Figures 1–4), we next experimentally examined how the firing patterns of ganglion cells are affected by their own action potentials and the action potentials of other ganglion cells. For going beyond correlation analysis (Greschner et al., 2016), here we sought to perturb the activity of many ganglion cells simultaneously to strongly drive the recurrent signaling pathways, but in a manner independent of light signaling pathways. Using a suction electrode (Stys et al., 1991), we thus electrically stimulated the tip of the optic nerve (around 5 mm long) emerging from an isolated retina in the dark, and simultaneously recorded the spiking activity of ganglion cells to monitor the outcome of the perturbation (Figure 4A; see

Methods for details). We then focused on those cells with sufficiently high baseline firing rates (>1 Hz) to reliably detect a change in their activity. In particular, a decrease of the firing rates should follow the nerve stimulation if ganglion cells are part of the proposed negative feedback circuitry, whereas no significant change should be observed if the stimulated cells are all purely postsynaptic within the retina.

We found that the optic nerve stimulation had pronounced effects on the spontaneous activity of ganglion cells in both salamander and mouse retinas (Figures 4 and 5 and Supplementary Figure 5). About a third of the recorded cells showed immediate antidromically-evoked spikes after the nerve shock (e.g., Figure 4B,C and Supplementary Figure 5A,B). In both cells with and without the antidromic spikes, we frequently observed a period of enhanced firing over tens of milliseconds (e.g., Figure 4D and Supplementary Figure 5B) and a suppression of firing on a longer time scale over hundreds of milliseconds (e.g., Figure 4C,D and Supplementary Figure 5A,B). The proportion and time course of these indirect enhancing and suppressing effects varied greatly between ganglion cells (Figure 5A and Supplementary Figure 5C). Importantly, these indirect effects of the nerve stimulation were observed regardless of the presence of the antidromic spikes (e.g., Figure 4D). Moreover, applying the electrical stimulation to the bath without holding the optic nerve did not affect the ganglion cell spontaneous activity (*data not shown*). Therefore, the observed changes in the firing after the nerve stimulation were likely mediated by signaling through circuits.

To examine how the intrinsic properties of the cells affect the outcome of the optic nerve stimulation, we repeated the experiments using a high-potassium Ringer's medium (13 mM KCl replacing NaCl; Figure 5B). In this condition, ganglion cells generally showed an increased spontaneous activity (from 0.7 ± 1.6 Hz to 2.0 ± 3.2 Hz, mean $\pm$ standard deviation; $p<0.001$, sign-test; $N=855$ salamander cells). Nevertheless, there was no change in the fraction of the cells with antidromic spikes (control, 39% , Figure 5A; high-potassium, 40%, Figure 5B; $p=0.5$, χ^2-test), validating the reliability of our experimental procedures. The fraction of the cells with a suppression after the nerve shock remained unchanged as well (21% versus 19%; $p=0.8$). This supports little contribution of the cell intrinsic mechanisms to the suppression, such as after-hyperpolarization (Gouras, 1969). In contrast, the fraction of the cells with an enhanced firing

increased significantly (from 31% to 43%; p=0.008). This indicates that the fast enhancement of

the ganglion cell activity after the nerve shock was susceptible to a change in the cell intrinsic

properties, and thus cannot be attributed solely to the network effects as we cannot exclude

possible changes in the ganglion cell properties due to the nerve stimulation.

Taken together, consistent with our model predictions, these data provide direct evidence

that ganglion cells can drive negative feedback signaling in the inner retina besides previ-

ously described excitatory interactions via gap junction networks (Brivanlou et al., 1998;

Mastronarde, 1983; Meister et al., 1995; Nath and Schwartz, 2017; Puller et al., 2020; Roy

et al., 2017; Trong and Rieke, 2008; Völgyi et al., 2013).

247 Recurrent inhibitory circuits from retinal ganglion cells involve gap junc-248 tions and amacrine cells

What are the retinal circuits underlying such recurrent inhibitory signaling from ganglion

cells? Previous anatomical and computational studies suggest that ganglion cells can form a

recurrent circuit with amacrine cells via gap junctions (Bloomfield and Völgyi, 2009; Kenyon

and Marshak, 1998; Völgyi et al., 2009). Here we took a pharmacological approach in the

salamander retina to test this circuit hypothesis. We first blocked gap junctions by applying

100 μM meclofenamic acid (Figure 6A,B). This had little effect on the fast enhancement of

the firing, indicating that the optic nerve stimulation itself can affect the intrinsic excitability

of the cells for a short period of time beyond driving the antidromic spikes (see also Figure 5).

Nevertheless, the drug application generally led to an elimination of the slow suppression after

the nerve shock in the dark (7 out of 8 cells; p=0.036, Fisher's exact test). Electrical synapses

were thus found indispensable for the negative feedback signaling from ganglion cells. To test

the involvement of amacrine cells in this signaling pathway, we next blocked γ-aminobutyric

acid (GABA) and glycine transmission by applying 100 μM picrotoxin and 1.0 μM strychnine

(Figure 6C,D). We found that these inhibitory transmission blockers also abolished the slow

suppression after the nerve stimulation in the dark (7 out of 7 cells; p=0.008, Fisher's exact

test), while the fast enhancement remained intact. Taken together, these results support that the

negative feedback pathway is physiologically functional from ganglion cells to amacrine cells

via gap junctions (Greschner et al., 2016; Kenyon and Marshak, 1998; Sakai and Naka, 1988,
1990).

Negative feedback signaling modulates gain but not selectivity of ganglion cell visual responses

Thus far we have electrophysiologically examined the negative feedback without presenting
visual stimuli. But, how does it affect the visual response properties of retinal ganglion
cells? Our model predicts a role in transiently controlling the response gain (Figure 1D). To
experimentally test this, we next presented white-noise visual stimuli in combination with
the optic nerve stimulation, and exploited the reverse-correlation methods to systematically
examine changes in the response gain and feature selectivity of individual cells due to the nerve
stimulation (Figure 7A and Supplementary Figure 6A; Chichilnisky, 2001; Wu et al., 2006).
Specifically, we fitted a linear-nonlinear (LN) cascade model to the visual responses of each
cell at different time points from the nerve shock, and assessed i) the feature selectivity by an
ON-OFF index (e.g., Figure 7C,D and Supplementary Figure 6C,D), defined as the difference
of the peak and valley values of the linear filter, normalized by the sum of the two (see Eq.(4)
in Methods for details) and the spectral peak frequency of the linter filter (e.g., Figure 7E,F and
Supplementary Figure 6E,F); and ii) the response gain by a sigmoid function fitted to the profile
of the static nonlinearity from the LN model (Eqs.(5)–(6) in Methods; e.g., Figure 7G,H and
Supplementary Figure 6G,H). In the present experiments, due to the presence of the stimulation
pipette, we could not avoid a distortion of the visual stimuli projected from above the retina.
Our analysis is thus limited to the full-field response properties, but not extended to the spatio-
temporal properties.

During a period of suppressed firing after the nerve shock (Figure 7B and Supplementary
Figure 6B), the linear filter profile remained largely unchanged (change in the ON-OFF
index, -0.00 ± 0.09, mean $\pm$ standard deviation, $p=0.5$, sign-test, $N=25$ salamander cells,
Figure 7I; change in the spectral peak frequency, 0.01 ± 0.46 Hz, $p=0.8$, Figure 7J; see also
Supplementary Figure 6I,J for the mouse cells). In contrast, it sometimes showed a marked
change during a period of enhanced firing: e.g., from monophasic (having one positive

or negative phase) to biphasic (having both positive and negative phases; Supplementary Figure 6C). Nevertheless, both measures of the feature selectivity did not show significant changes across the population (0.02 ± 0.14 and -0.00 ± 1.18 Hz for the ON-OFF index and spectral peak frequency, respectively; both with $p>0.3$; $N=95$ salamander cells). These data suggest that the recurrent signals from ganglion cells had little effect on the feature selectivity of their visual responses.

The response gain, in contrast, was affected during both periods of enhancement and suppression after the nerve stimulation (Figure 7G,H,K,L and Supplementary Figure 6G,H,K,L). On the one hand, the static nonlinear gain function was down-regulated during a period of suppression (e.g., Figure 7G,H and Supplementary Figure 6G,H). In particular, the upper bound spike probability showed a significant decrease over the population (-0.03 ± 0.02; $p<0.001$, sign-test, Figure 7K; see also Supplementary Figure 6K) while the lower bound remained the same (-0.00 ± 0.00; $p>0.5$; Figure 7L; see also Supplementary Figure 6L). On the other hand, the gain was generally up-regulated during a period of enhancement (e.g., Supplementary Figure 6G,H) with marginally increased upper bounds (0.03 ± 0.08, $p=0.08$, Figure 7K; see also Supplementary Figure 6K) and significantly increased lower bounds (0.06 ± 0.08, $p<0.001$, Figure 7L; see also Supplementary Figure 6L) over the population. It should be noted, though, that this non-specific increase in the activity resulted not only from the gap junction network effect, but also from a possible brief change in the cell's excitability due to the nerve stimulation (see also Figures 5 and 6).

Taken together, these results indicate that the negative feedback signals primarily contribute to modulating the response gain, while ganglion cells may adaptively change their visual response properties by exploiting the whole recurrent circuitry.

Discussion

Circuit modeling is a powerful approach to integrate brain anatomy and physiology for better understanding the overall function of the system (Herz et al., 2006). By explicitly representing individual neurons and their connections with the model parameters, circuit models can provide

predictions on the structure and function of the target neuronal circuitry, and subsequently these predictions can be tested by experiments (Real et al., 2017). Here we took this theory-driven approach to functionally characterize the ganglion cell feedback circuits in the inner retina. We first extended a retinal cascade model (Real et al., 2017) to incorporate interactions between ganglion cells as suggested by anatomical studies (Bloomfield and Völgyi, 2009; Völgyi et al., 2009), and derived a prediction that retinal ganglion cells of the same response polarity form reciprocal inhibition over a long distance for a transient gain modulation (Figures 1–3 and Supplementary Figures 1–4). We then experimentally validated that ganglion cells can indeed suppress the firing among themselves by propagating signals to amacrine cells via gap junctions (Figures 4–6 and Supplementary Figure 5). Furthermore, as predicted by our circuit model, we showed that this negative feedback lowers the visual response gain without much affecting the stimulus feature selectivity of the cells (Figure 7 and Supplementary Figure 6). The discovery of this new gain control mechanism in the retina not only highlights the importance of gap junctions in visual processing, but also promises that theory-driven approaches will further reveal neural circuit functions in future studies.

Retinal gap junction networks have been well conserved across species over evolution (Völgyi et al., 2013). In both salamander and mouse retinas, our circuit model analysis indeed showed the presence of negative couplings among some distal ganglion cells, whereas positive couplings were found more frequently among proximal cells (Figure 3 and Supplementary Figure 3). This spatial organization is consistent with previous studies in both salamander (Cocco et al., 2009) and primate retinas (Greschner et al., 2016; Pillow et al., 2008). Moreover, ganglion cells in both salamander and mouse retinas showed a period of suppression following the optic nerve stimulation under the *ex vivo* condition (Figures 4 and 5 and Supplementary Figure 5). Such suppression was also reported in the catfish retina (Sakai and Naka, 1988, 1990) as well as in the primate retina (Gouras, 1969), although the time scale in these previous reports was shorter than that in this study. Nevertheless, later modelling (Kenyon and Marshak, 1998) and spike correlation studies (Greschner et al., 2016) support that such suppression arises from the recurrent circuitry involving amacrine cells via gap junctions, rather than a transient after-hyperpolarization due to the antidromic spikes evoked by the optic nerve stimulation. Taken

together, recurrent circuits from retinal ganglion cells are likely present widely across species. To gain full insights into these circuits' function, it will be critical next to unveil and compare the underlying synaptic mechanisms across species (Baden et al., 2020), especially if these ganglion-cell-coupled amacrine cells directly inhibit ganglion cells (postsynaptic inhibition) or instead inhibit bipolar cells that provide excitatory inputs to the ganglion cells (presynaptic inhibition) or both.

In this study, we employed an optic nerve stimulation to perturb ganglion cell activity independently of the light signaling pathway (Figure 4A). This successfully drove antidromic spikes in many ganglion cells in a time-locked manner (e.g., Figure 4B) and the obtained results provided direct experimental evidence supporting our model predictions on ganglion cell coupling properties. There are, however, some caveats, such as possible changes in the intrinsic properties of ganglion cells (Figures 5 and 6) and a potential activation of efferent axons from the brain to the retina (Koves et al., 2016; Repérant et al., 2006). Previous studies suggest that dopaminergic amacrine cells receive such centrifugal inputs under the control of circadian rhythms (Gastinger et al., 2006). Although dopamine can affect gap junction networks at nearly every stage of retinal processing, the kinetics are much slower than the time scales considered in this study ($\sim$100 ms) (Pereda et al., 2013; Roy and Field, 2019; Witkovsky and Dearry, 1991). The effects of the efferent signaling, if any, should thus be negligible on the outcome of the nerve stimulation we observed. These dopaminergic amacrine cells are also suggested to receive direct inputs from intrinsically-photosensitive retinal ganglion cells via axon collaterals (Zhang et al., 2012, 2008). For the same reason, however, the contribution of this dopaminergic pathway should also be minimal in this study.

Another caveat is that the optic nerve stimulation potentially led to an extensive perturbation of the coupling pathways due to a strong synchronous activation of ganglion cell populations. This may explain why the measured effect of the negative feedback lasted long, over hundreds of milliseconds (Figures 4 and 5 and Supplementary Figure 5) as opposed to tens of milliseconds in the model predictions (Figure 3 and Supplementary Figure 3). Single-cell stimulation, however, can be too weak to reveal any indirect negative interaction among ganglion cells (Mastronarde, 1983; Trong and Rieke, 2008; but see Sakai and Naka, 1988, 1990). It is a future challenge

to further investigate the circuit mechanisms under more physiological conditions and test remaining model predictions, such as reciprocal negative couplings between retinal ganglion cells of the same response polarity (Figure 3G–I and Supplementary Figure 3G–I).

Recurrent normalization plays important roles in many brain functions (Carandini and Heeger, 2012). Consistent with the previous study in the primate retina (Greschner et al., 2016), both our computational model and experimental data show that the ganglion cell recurrent network plays a role in controlling the visual response gain in both salamander and mouse retinas (Figure 7 and Supplementary Figure 6). Interestingly, the interactions between ganglion cells extend over a wide spatial range (up to ∼1 mm; Figure 3 and Supplementary Figure 3) well beyond the extent of their dendritic fields (Völgyi et al., 2009; Zhang and Wu, 2010). Moreover, this gain control mechanism works on a slower time scale (∼100 ms), if not the slowest (Wark et al., 2009), than many other local gain control mechanisms in the retina, such as the synaptic adaptation achieved at bipolar cell terminals at the cellular level (Euler et al., 2014; Matthews, 1999), or negative feedback loop among bipolar cells and amacrine cells at the circuit level (Dong and Werblin, 1998; Nirenberg and Meister, 1997; Tachibana and Kaneko, 1988). Therefore, the ganglion cell feedback circuits likely contribute to slowly equalizing the response gain across populations, but not mediating visual functions that need precise spike timing of individual cells (Geffen et al., 2007; Gollisch and Meister, 2008; Van Rullen and Thorpe, 2001; Victor, 1999). To investigate the structure and function of the feedback circuits for each ganglion cell type, future studies will benefit from further elaborations on the circuit models combined with calcium or voltage imaging techniques to monitor the subcellular activity of individual neurons and trace signal flow within the retina (Baden et al., 2016; Franke et al., 2017).

Methods

No statistical method was used to predetermine the sample size. The significance level is 0.05 (with Bonferroni correction where appropriate) in all analyses unless noted otherwise. All experiments were performed in strict accordance with the protocols approved by the

Institutional Animal Care and Use Committee at Harvard University or California Institute of Technology, or under the license 233/2017-PR from Italian Ministry of Health. The data analysis and circuit modeling were done in Matlab (Mathworks) and Python. All data and codes are available upon request.

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Hiroki Asari (asari@embl.it).

Modeling

We first reanalyzed the data sets in Asari and Meister (2012, 2014) and Real et al. (2017) for the salamander retinal ganglion cells and those in Lefebvre et al. (2008) for the mouse ganglion cells. Specifically, we focused on the responses to the random noise stimulus consisting of a 1-dimensional array of adjacent bars 8.3–80 μm in width. The light intensities of these bars were drawn from a binary black-or-white distribution (luminance range 0.5–36 mW/m^2), and changed simultaneously, independently, and randomly with a refresh rate of 60–100 Hz.

The distance between the cells (Figure 3 and Supplementary Figure 3) was calculated from the receptive field centers. The spatiotemporal receptive fields of the cells (10–17 ms bin size; 0.4 s window) were estimated by reverse-correlation methods using randomly flickering checkerboard stimuli (30–83 μm square fields; 60–100 frames/s), and their center location was estimated by the 3-dimensional Gaussian curve fit. Cells with positive receptive field center values were classified as ON cells, whereas those with negative values as OFF cells.

Data selection

The raw data sets contained 479 ganglion cells from 10 isolated salamander retinas, and 35 ganglion cells from a mouse retina. Of those, 185 OFF and 4 ON salamander cells and 10 ON and 12 OFF mouse cells were selected for the subsequent modeling analyses according to the following criteria.

1. Cells should have a high spike sorting quality. To ensure low false positives, cells with >15% of spikes with <1.7 ms inter-spike intervals were discarded. To ensure low

false negatives, cells with nearly identical spatiotemporal receptive field profiles were

eliminated except for the one with the highest spike counts.

2. Cells should respond well to the visual stimulus for robust model fitting, assessed by the

spike counts during the stimulus presentation period and the performance of a linear-

nonlinear (LN) cascade model (Chichilnisky, 2001; Wu et al., 2006). The LN model was

fitted by reverse-correlation methods, and cells with $<$3,000 spikes and $<$10% LN model

prediction (coefficient of determination; see below Eq.(3)) were discarded.

3. Cells should have little response correlation due to common visual inputs for modeling

interactions between cells. Neighboring cells were included in the coupling models (see

below Model formalism) only if Pearson correlation between their spike trains and those

of the target cell was low (between -0.1 and 0.1; Supplementary Figure 2A,B). This

generally resulted in no overlap of the receptive field centers (e.g., Figure 1B).

Model formalism

We employed the cascade model framework as described in Real et al. (2017). As a non-

coupled model (Figures 1 and 2 and Supplementary Figures 1 and 4), we thus used the so-called

"linear-nonlinear-feedback-delayed-sum-nonlinear-feedback" (LNFDSNF) or full model, and

the "linear-nonlinear-sum-nonlinear" (LNSN) or reduced model. In short, the first LNF stages

collectively work as spatial subunits of upstream bipolar cells (Bipolar Cell Modules; Eq.(S3)–

(S7) in Real et al., 2017); the middle D is a delay due to lateral signal propagation via amacrine

cells (Amacrine Cell Modules; Eq.(S8) in Real et al., 2017); and the last SNF stages represent

the spatial summation by a target ganglion cell (Ganglion Cell Modules; GCM; Eq.(S9) in

Real et al., 2017), followed by the cell's output nonlinearity (Eq.(S6') below) and the feedback

(Eq.(S7) in Real et al., 2017). Here we replaced the Eq.(S6) in Real et al. (2017) with the

following formula:

$$y(t) = \begin{cases} 0, & \text{if } z(t) \leq \theta \\ \alpha(z(t) - \theta), & \text{otherwise,} \end{cases} \tag{S6'}$$

where α and θ are the slope and threshold of a half-wave rectification function, respectively. The GCM nonlinearity was thus equipped with two free parameters in this study (Supplementary Figure 4), while it was fixed in Real et al. (2017), equivalent to $\alpha = 1$ and $\theta = 0$ in Eq.(S6').

A full coupling model (LNFDSCNF; Supplementary Figure 1A) and a reduced one (LNSCN; Figure 1A) were then built by introducing the coupling step (C) before the GCM nonlinearity (N) and feedback (F). Specifically, the coupling effects on a target ganglion cell were modelled as a delayed weighted sum of the neighboring ganglion cell activities. For each k-th neighboring cell, two free parameters were hence assigned: l_k for the latency of the signal transmission and a_k for the coupling strength. The latency parameter l_k is non-negative and treated as a delay function that requires interpolation of the measured firing rate $r_k(t)$ of the k-th neighboring cell. The delayed activity $r_k^*(t)$ is thus given as:

$$r_k^*(t) = (1 - \{l_k\})\, r_k(t - \lfloor l_k \rfloor) + \{l_k\}\, r_k(t - \lfloor l_k \rfloor - 1), \tag{1}$$

where $\lfloor l_k \rfloor$ is the largest integer not greater than l_k, and the fractional part $\{l_k\} = l_k - \lfloor l_k \rfloor$. The delayed activity $r_k^*(t)$ is then weighted by a coupling strength a_k and added to the signal $x(t)$ coming from the previous summation (S) step (Eq.(S9) in Real et al., 2017):

$$y(t) = x(t) + \sum_k a_k r_k^*(t). \tag{2}$$

The resulting signal $y(t)$ is in turn used as an input to the following GCM nonlinearity (Eq.(S6')) and feedback steps (Eq.(S7) in Real et al., 2017) to obtain an estimated firing rate of the target cell (e.g., Figure 1).

Model fitting and assessment

We wrote custom codes in Python to fit the models to the ganglion cell firing rates (bin size, 1/100–1/60 s) in response to the randomly flickering bar stimuli. For each cell, we fitted the coupled models in two configurations (Figures 1–3 and Supplementary Figures 1–4): the one including only those surrounding cells of the same response polarity as the target cell (e.g., OFF target cell with OFF surrounding cells), and the other with those of different response polarities

alone (e.g., OFF target cell with ON surrounding cells). The Kolmogorov–Smirnov test was performed to compare the model performance gain over the non-coupling models (see below) in the two configurations (Figure 2 and Supplementary Figure 1B–E).

Model performance was assessed by the coefficient of determination between the measured ganglion cell firing rate $r(t)$ and the model prediction $\hat{r}(t)$:

$$R^2 = 1 - \frac{\sum_t (r(t) - \hat{r}(t))^2}{\sum_t (r(t) - \langle r(t) \rangle)^2}. \tag{3}$$

where $\langle \cdot \rangle$ denotes the mean. It reaches its maximum of 1 in the case of an exact agreement between the two binned sequences, and is around 0 or less in the case of unrelated sequences. Paired t-test was used to compare the model performance with and without couplings (Figure 2 and Supplementary Figure 1B–E). Because the coupled model did not improve its performance for mouse cells with surrounding cells of different response polarities, this data set was excluded from the subsequent model parameter analyses (Figure 3 and Supplementary Figures 2–4).

To avoid over-fitting, the model parameters were optimized using a training data set ($\sim 80\%$ of the data) and the model performance was evaluated on a separate testing data set. The testing data set for the salamander data included 8–12 repeats of the identical flicker sequence, and hence the model's output was compared to the average firing rate over all these trials.

Model analysis

We ran a shuffling analysis to evaluate the noise level of the coupling parameters in the LNSCN and LNFDSCNF models (lower and upper thresholds at 0.5 and 99.5 percentiles, respectively; Figure 3 and Supplementary Figures 2 and 3). Specifically, for each modelled cell, we fitted the model parameters to its true spike trains together with randomly jittered spikes trains from all coupled neighboring cells. This keeps intact the stimulus-dependence of the modelled cell's response, but breaks the correlation to the responses of the other cells. Thus the obtained coupling parameters form a distribution expected from a chance level.

We performed a perturbation analysis to test the convergence and sensitivity of the model parameters. Specifically, after fitting LNSCN and LNFDSCNF models, we reset the coupling parameters to be zero while keeping the other parameters intact, and refitted the models to obtain

a new set of the parameters. We then compared the original and reoptimized coupling strength and delay parameters using Pearson's correlation coefficient (Supplementary Figure 2C,D). The difference of the model performance (R^2 in Eq.(3)) between the original and reoptimized models was used to determine the confidence interval of the performance gain from the non-coupling to the coupling models (lower and upper thresholds at 5 and 95 percentiles, respectively; Figure 2 and Supplementary Figure 1B–E).

For the population analysis of the coupling properties (Figure 3 and Supplementary Figure 3), we first pulled together the parameter values for each cell pair across all coupling models. We then performed the Wilcoxon rank-sum test to examine the distance and the signal delay between the positively and negatively coupled cells (Figure 3A–F and Supplementary Figure 3A–F). To analyze the symmetry of the coupling effects between cells, we examined by a χ^2-test if the polarity of the signal from one cell to another depends on that in the opposite direction (df=4; Figure 3G–I and Supplementary Figure 3G–I). The data sets from ON and OFF cells were combined as they showed the same trend.

We analyzed in three ways how the coupling affects the behavior of the ganglion cell module (GCM) in the circuit models (Supplementary Figure 4). First, we assessed the input dynamics to GCM as a collective measure of the model's upstream circuit properties. Specifically, for each cell, we compared the outputs of the summation (S) stage in the coupled and non-coupled models (Eq.(S9) in Real et al., 2017) using Pearson's correlation coefficient (Supplementary Figure 4A,D). Second, we examined the GCM feedback filters from the parameters of the second feedback (F) stage in the full models (Eq.(S7) in Real et al. (2017); Supplementary Figure 4B,E). Finally, we compared the GCM nonlinearity parameters (threshold θ and slope α for the second nonlinearity (N) stage; Eq.(S6')) using Pearson's correlation coefficient between the coupled and non-coupled models (Supplementary Figure 4C,F).

Electrophysiology

The dark-adapted retina of a larval tiger salamander (*Ambystoma tigrinum*) or an adult wild-type mouse (*Mus musculus*; C57BL/6J strain, 2–6 months old) was isolated with an intact optic nerve attached (∼5 mm long), and placed on a flat array of 61 extracellular electrodes with

the ganglion cell side down. The salamander retina was superfused with oxygenated Ringer's medium (in mM: NaCl, 110; NaHCO$_3$, 22; KCl, 2.5; MgCl$_2$, 1.6; CaCl$_2$, 1; and D-glucose, 10; equilibrated with 95% O$_2$ and 5% CO$_2$ gas) at room temperature, and the mouse retina with oxygenated Ames' medium (Sigma-Aldrich, A1420) at 37 °C. The electrode array recorded the extracellular signals from ganglion cells with each electrode sampled at 10 kHz, while photoreceptors and/or the optic nerve were stimulated visually and/or electrically, respectively. Spike trains from individual ganglion cells were extracted from raw voltage traces by a semi-automated spike-sorting algorithm (Pouzat et al., 2002) written in IGOR Pro (Wave Metrics).

We adapted the methods described in Stys et al. (1991) to electrically stimulate the tip of the optic nerve emerging from an isolated retina using a suction electrode (Figure 4A). Specifically, we used bipolar pulses (10–50 V, 0.02–0.5 ms) from a stimulus isolator (Grass Instrument, SD9) at 2/3–1 Hz (100–200 trials in the dark; 1,300–2,000 trials with visual stimulation) controlled by custom software written in LabView (National Instruments). Antidromically evoked spikes were observed in some ganglion cells at a latency of around 5 ms after the nerve stimulation (e.g., Figure 4B,C and Supplementary Figure 5A,B). Due to the stimulus artifacts, we were not able to detect any spikes within a few milliseconds after the nerve stimulation (e.g., Figure 4B). When the suction electrode was inserted into the bath without holding the nerve ending, the electrical stimulation did not affect the spontaneous activity of ganglion cells (*data not shown*).

In total, recordings were made from 855 ganglion cells in 19 salamander retinas and 368 cells in 7 mouse retinas for the optic nerve stimulation experiments. Of those, all salamander retinas were examined with a high-potassium Ringer's medium (13 mM KCl replacing NaCl; Figure 5B), 349 cells in 11 salamander retinas with a gap junction blocker (100 μM meclofenamic acid; Figure 6A,B), 380 cells in 6 salamander retina with inhibitory synaptic transmission blockers (100 μM picrotoxin and 1.0 μM strychnine; Figure 6C,D), and 167 cells in 6 salamander retinas and 172 cells in 3 mouse retinas with visual stimulation (Figure 7 and Supplementary Figure 6). As we focused on a suppression of ganglion cell activity, only those cells with a sufficiently high baseline firing rate (>1 Hz) were selected for subsequent analyses.

Visual stimulation

Visual stimuli were displayed on a gamma-corrected cathode-ray tube monitor (DELL E773c; frame rate 100 Hz; mean luminance 18 mW/m^2) and projected onto the photoreceptor layer of the retina from above through a custom-made lens system. We presented full-field random flicker stimuli (100 frames per second; light intensities drawn from Gaussian distribution with mean luminance of 18 mW/m^2 and standard deviation of 7 mW/m^2) for examining the effects of the optic nerve stimulation on the ganglion cell visual responses (Figure 7 and Supplementary Figure 6). Here we could not use spatially-structured stimuli because the stimulation pipette created a distortion in an uncontrollable manner.

Data analysis

Optic nerve stimulation in the dark

To measure the effects of the optic nerve stimulation in the dark (Figures 4–6 and Supplementary Figure 5), we first computed the peri-stimulus time histogram (PSTH) for each ganglion cell with increasing bin sizes (0–10, 10–20, 20–40, 40–80, 80–160, 160–320, and 320–640 ms after the nerve stimulation), and identified those time bins that had significantly different firing rates from the baseline activity (320 ms period before the onset of the nerve stimulation) using bootstrap resampling methods over trials (10,000 repeats). Here we chose unequal bin sizes 1) to robustly assess a decrease of the firing rate even for those cells with relatively low baseline firing rates (threshold at 1 Hz); and 2) to avoid the problem of multiple comparisons. Cells with a significantly increased firing rate in the first time bin (0–10 ms) were considered as the ones directly evoked by the nerve stimulation (Figure 5A,B and Supplementary Figure 5C, *top* rows) because antidromic spikes typically had a latency of around 5 ms (e.g., Figure 4B,C and Supplementary Figure 5A,B). Cells with significantly different firing rates in the second time bins and thereafter (i.e., >10 ms latency) were considered as the ones indirectly affected by the nerve stimulation (Figure 5A,B and Supplementary Figure 5C, *bottom* rows). We ran a χ^2-test to compare the proportions of the cells in the normal (Figure 5A) and high-potassium (Figure 5B) conditions that showed antidromic spikes, enhanced firing, or suppressed firing after the nerve stimulation, respectively. For a display purpose, we computed the PSTHs with 20 ms bins in

Figure 4C,D and Supplementary Figure 5A,B, and labeled those significantly below or above the baseline (250 ms period before the onset of the nerve stimulation) in blue and red shades, respectively.

Optic nerve stimulation with visual stimulation

We used stimulus ensemble statistical techniques ("reverse correlation" methods; 400 ms window; 10 ms bin width) to calculate the linear filter and static nonlinear gain function of the recorded cells (Figure 7 and Supplementary Figure 6). Specifically, for each ganglion cell, we first computed the PSTH (100 ms bin width) with respect to the nerve stimulation during the full-field random flicker stimulus presentation, and identified those time bins with significantly higher or lower firing rates than the baseline (500 ms period before the nerve stimulation) using the bootstrap resampling methods over trials (10,000 repeats; e.g., Figure 7B and Supplementary Figure 6B). For each time bin, we then estimated the linear filter by a spike-triggered average stimulus (e.g., Figure 7C and Supplementary Figure 6C). The obtained linear filters indicate the average stimulus features that made the cell fire action potentials. We then characterized the change in their profile from the baseline in two ways. First, we used the ON-OFF index, defined as the difference between the peak and valley values of the linear filter, normalized by the sum of the two:

$$\text{ON-OFF index} = \frac{|\text{peak}| - |\text{valley}|}{|\text{peak}| + |\text{valley}|}. \tag{4}$$

The ON-OFF index value of -1, 1, and 0 indicates the stimulus feature selectivity towards purely OFF stimuli, purely ON stimuli, and both ON and OFF stimuli, respectively (e.g., Figure 7D and Supplementary Figure 6D). Second, we calculated the power spectral density of the linear filter and estimated the peak frequency by fitting a gaussian curve as a measure of the cell's temporal frequency tuning (e.g., Figure 7E,F and Supplementary Figure 6E,F). Sign-test was used to evaluate the change in these properties over the population during the suppressed and enhanced firing periods (Figure 7I,J and Supplementary Figure 6I,J).

For each epoch, we also computed the static nonlinear gain function $P(\text{response}|\text{stimulus})$ as a function of the linear filter output values by the convolution with stimulus fragments:

$$P(\text{response}|\text{stimulus}) = \frac{N(\text{stimulus}|\text{response})}{N(\text{stimulus})}, \tag{5}$$

where $N(\text{stimulus})$ and $N(\text{stimulus}|\text{response})$ are the distributions obtained from all stimuli and spike-triggered stimulus ensembles, respectively (e.g., Figure 7G and Supplementary Figure 6G). For quantification, we fitted the following sigmoid function $f(x)$ to the gain function profiles:

$$f(x) = \frac{u - l}{1 + \exp[-s(x - c)]} + l, \tag{6}$$

and calculated the change in the parameters (u, upper bound; l, lower bound; c, center; s, slope) from the baseline (e.g., Figure 7H and Supplementary Figure 6H). Sign-test was performed to test the statistical significance (Figure 7K,L and Supplementary Figure 6K,L).

Figure legends

Figure 1: Coupling model suppresses false-positive responses in non-coupling model

(A) Schematic diagram of the circuit model with couplings among retinal ganglion cells ("LNSCN" model; see also Supplementary Figure 1). The target ganglion cell (red) integrates inputs from upstream bipolar cells as well as surrounding retinal ganglion cells (blue) with certain coupling strengths and delays. The model without couplings is identical to the "LNSN" model in Real et al. (2017).

(B) Receptive field centers of 20 OFF ganglion cells simultaneously recorded from an isolated salamander retina (red, representative modelled cell in C and D; blue, cells included in the coupling model for the representative cell; gray, cells excluded from the coupling model due to high spike correlations with the modelled cell; see also Supplementary Figure 2A). Each outline

represents a two-dimensional Gaussian fit to the receptive field profile (contour at 1 standard deviation).

(C)   Responses of ganglion cells (red, representative modelled cell from B; blue, cells included in the coupling model) to repetitions of the white noise stimulus (8 repeats in total). Each row in the raster denotes spikes from a single stimulus repeat for each cell. Cyan lines indicate timepoints where false positive responses in the non-coupled LNSN model were suppressed in the coupled LNSCN model (arrows in D).

(D)   Time course of the firing rate of the representative cell (red) and that of the model outputs with (black, LNSCN model) and without (cyan, LNSN model) couplings fitted to this cell. The LNSN model occasionally mis-predicted responses at which the cell did not fire spikes, but the LNSCN model correctly suppressed them in many cases (arrows). The coupling parameters (coupling strength and delay) for each surrounding cell included in the LNSCN model are shown next to the raster graph in C.

Figure 2: Coupling model outperforms non-coupling model

Cumulative probability distributions (black; confidence intervals at 5 and 95 percentiles from perturbation analysis in gray; see Methods for details) of the LNSCN model performance gain over LNSN model for salamander ganglion cells, calculated as a difference of the coefficients of determination (R^2, Eq.(3) in Methods; see also Supplementary Figure 1).

(A,B)   The LNSCN model outperforms the LNSN model more with the neighboring cells of the same response polarity (A; N=185 OFF cells with OFF neighbors; ΔR^2=0.017$\pm$0.016, mean $\pm$ standard deviation; p<0.001, paired t-test) than with those of different polarities (B; N=35 OFF cells with ON neighbors and 4 ON cells with OFF neighbors; ΔR^2=0.013$\pm$0.017; p<0.001): p=0.006 from Kolmogorov–Smirnov test. Here and thereafter, three stars ($\star \star \star$) indicate p<0.001; $\star\star$, p<0.01; and $\star$, p<0.05.

(C,D) Corresponding figure panel for the mouse ganglion cells. The LNSCN model out-

performs the LNSN model when the neighboring cells of the same response polarity are

included (C; N=10 ON cells with ON neighbors and 12 OFF cells with OFF neighbors;

ΔR^2=0.007±0.007; p<0.001), but not with the neighboring cells of different polarities (D;

N=10 ON cells with OFF neighbors and 12 OFF cells with ON neighbors; ΔR^2=0.001±0.010;

p>0.5): p<0.001 from Kolmogorov–Smirnov test.

Figure 3: Coupling model predicts faster enhancing effects among proxi-

mal retinal ganglion cells and slower suppressing effects among distal cells

(A,B) Coupling strength (A; a_k in Eq.(2) in Methods) and delay (B; l_k in Eq.(1) in Methods)

parameter values from the LNSCN model plotted as a function of the distance between

salamander retinal ganglion cells of the same response polarity (1460 OFF cell pairs in total;

yellow circles, representative cases in Figure 1B–D). The noise level was determined by

shuffling analysis (gray, 0.5 and 99.5 percentile; see Methods for details). Positive couplings

(red, N=211) were found at a shorter distance (A; 0.31±0.14 mm versus 0.40±0.26 mm;

median±interquartile range as shown by the box plot; p<0.001, rank-sum test) with a shorter

latency (B; 3.5±3.6 ms versus 4.6±4.0 ms; p<0.001) than negative couplings (blue, N=98).

(C,D) Corresponding figure panels for salamander cells with different response polarities

(47 OFF cells with ON neighbors and 46 ON cells with OFF neighbors in total). Couplings

above the noise level were mostly positive (N=10) with the distance of 0.28±0.08 mm (C;

median±interquartile range) and the latency of 4.2±1.5 ms (D).

(E,F) Corresponding figure panels for mouse retinal ganglion cells of the same response

polarity (76 ON cell pairs and 116 OFF cell pairs in total). Positive couplings (red, 18 ON and

15 OFF cell pairs) were found at a shorter distance (C; 0.33±0.25 mm versus 0.78±0.74 mm;

p=0.002) and with a shorter latency (D; 0.0±0.0 versus 21.9±13.8 ms; p<0.001) than negative

couplings (blue, 7 ON and 9 OFF cell pairs). Data not analyzed for the mouse cells with different

response polarities because the coupled model performance did not show an improvement (Figure 2B).

(G–I) Comparison of the coupling strengths from one cell to another and vice versa (black, couplings above the noise level in both directions; dark grey, above the noise level only in one direction; light gray, both below the noise level). The number of data points in each category is shown in the figure panels. Symmetric, either mutually positive or negative, couplings were found more frequently than expected in both salamander (G, between cells of the same response polarity; H, between cells with different response polarities) and mouse (I, between cells of the same response polarity) retinas ($p<0.001$ in all three cases, χ^2-test with $df=4$)

Figure 4: Optic nerve stimulation produces fast excitation and slow inhibition in retinal ganglion cell firing activity

(A) Schematic diagram of the experiment. The tip of the optic nerve emerging from an isolated retina was electrically stimulated with a suction electrode (inter-stimulus interval, 1–1.5 s), while a population of ganglion cells was simultaneously recorded with a multi-electrode array (see panel B for example). A, amacrine cell; B, bipolar cell; G, ganglion cell; open circle, inhibitory synapse; closed circle, excitatory synapse; resistor symbol, electrical synapse. The other retinal cell types and synapses are omitted for simplification.

(B) Example raw data traces from an isolated salamander retina overlaid across trials of the optic nerve stimulation in the dark. The lightning symbol indicates the stimulation onset, and the gray area indicates the window for which the magnified traces are shown at the bottom panel. The antidromic spikes are indicated by the asterisk.

(C,D) Representative responses of retinal ganglion cells to the optic nerve stimulation (top, raster graphs; bottom, peri-stimulus time histogram (PSTH), magnified at the bottom for the window indicated by the gray area). A period of suppression was observed in cells either with (C) or without (D) antidromically evoked spikes (asterisk). Red- and blue-shaded bins in the

PSTHs indicate those in which the firing rate significantly increased or decreased from the spontaneous activity, respectively.

Figure 5: Intrinsic cellular property affects the fast excitation but not slow inhibition after the optic nerve stimulation

Population data of the salamander retinal ganglion cell responses to the optic nerve stimulation in the normal (A) and high-potassium (B; 13 mM KCl) Ringer's media. Each gray line represents the evoked change in the firing rate of a cell from its baseline (A, N=193 in total; B, N=324; red and blue circles, bins with significant increase and decrease, respectively), and the black line shows the mean over the cells in each group: top row, with antidromic spikes (A, N=75, B, N=131; p=0.5, χ^2-test); bottom row, without antidromic spikes; first column, no indirect effect; second and third columns, enhanced firing after the nerve stimulation (A, N=59; B, N=140; p=0.009); third and fourth columns, suppressed firing after the nerve stimulation (A, N=41; B, N=60; p=0.8). The representative responses in Figure 4C,D are shown in magenta in A.

Figure 6: Negative feedback involves electrical synapses and inhibitory synaptic transmission

(A) Spiking activity of a salamander retinal ganglion cell in the dark (top, raster graphs; bottom, PSTHs) in response to the optic nerve stimulation in the absence (black; control) and presence (brown) of gap junction blockers (100 μM meclofenamic acid; MFA).

(B) Summary of the effects of blocking gap junctions on the ganglion cell firing patterns after the nerve stimulation (top, control; bottom, with MFA), shown in the same format as Figure 5 (magenta, the representative cell in A). The suppression after the nerve stimulation was abolished after blocking electrical synapses (right: 7 out of 8 cells; p=0.036, Fisher's exact test), whereas the enhancement remained (left; 11 out of 11 cells).

(C,D)   A representative example (C; in the same format as A) and the summary (D; in the same format as B) of the effects of blocking inhibitory synaptic transmission (100 μM picrotoxin and 1.0 μM strychnine; PTX+STR, green) on the ganglion cell firing patterns after the nerve stimulation. The suppression after the nerve stimulation was abolished after blocking inhibitory transmission (right: 7 out of 7 cells; p=0.008), whereas the enhancement remained (left; 11 out of 11 cells).

Figure 7: Negative feedback modulates the visual response gain but not the feature selectivity

(A)   Schematic diagram of the experiment and analysis. Full-field white-noise visual stimuli were presented together with the optic nerve stimulation (inter-stimulus intervals, 1–1.5 s) to probe changes in the LN model parameters at different time windows from the nerve stimulation (see B–H for example). The linear filter represents the stimulus feature selectivity of a cell, while the static nonlinearity indicates the visual response gain.

(B)   PSTH of a representative salamander ganglion cell with respect to the optic nerve stimulation during the visual stimulus presentation. Blue-shaded bins (0–100 ms after the nerve shock) indicate those in which the firing rate was significantly lower than the baseline response of the cell (500 ms window before the nerve stimulation).

(C)   Linear filters of the example cell obtained by the reverse-correlation methods (i.e., spike-triggered average stimulus) using the spikes in different time bins from the PSTH (blue, 0–100 ms bin with significantly lower firing rates from B; gray, all the other 100 ms bins without significant firing rate changes; black, baseline).

(D)   Polarity of the linear filters (C) quantified by an ON-OFF index (i.e., the difference between the peak and valley values divided by the sum of the two; Eq.(4) in Methods).

(E)   Normalized power spectral density of the linear filters in C.

(F) Peak frequencies of the linear filters, estimated by fitting a gaussian curve to the power

spectra in E.

(G) Static nonlinearity of the example cell, computed for each corresponding linear filter at

different time bins (Eq.(5) in Methods). A sigmoid function (Eq.(6) in Methods) was fitted to

quantify the nonlinear profile (e.g., light blue for the suppressed period at 0–100 ms after the

nerve shock, with the upper and lower bounds of the cell's firing probability in dotted horizontal

lines).

(H) The upper and lower bound dynamics of the example cell's firing probability with respect

to the time from the nerve shock. The cell's response gain (upper bound) became lower during

the period of suppressed firing, while the stimulus feature selectivity remained unchanged (D

and F).

(I,J) Summary of the changes in the ON-OFF indices (I) and the spectral peak frequencies (J)

between the periods with and without significant firing rate changes after the nerve stimulation

(blue, decrease $N{=}25$; red, increase $N{=}94$; magenta, the representative cells in B–H). The

horizontal and vertical lines represent the mean and standard deviation, respectively. On average

there was no significant difference from the baseline across population ($p{>}0.3$ for both cases in

I and J, sign-test).

(F,G) Summary of the changes in the sigmoid function parameters fitted to the nonlinearity.

The upper bounds (K) significantly decreased during the period of suppression (blue, $p{<}0.001$,

sign-test), but only marginally increased during the period of enhancement (red, $p{=}0.08$). In

contrast, the lower bounds (L) significantly increased during the enhancement ($p{<}0.001$) but

remained the same during the suppression ($p{>}0.5$). The other parameters (center and slope) did

not change significantly (*not shown*).

Supplementary Figure Legends

Supplementary Figure 1: Retinal ganglion cell couplings improve the performance of the full circuit model

(A) Schematic diagram of the full circuit model with couplings ("LNFDSCNF" model). The model without couplings is identical to the "LNFDSNF" model in Real et al. (2017).

(B–E) Cumulative probability distributions (black; confidence intervals in gray) of LNFDSCNF model performance gain over LNFDSNF model for salamander (B, $\Delta R^2 = 0.008 \pm 0.008$ (mean +/- standard deviation), $p < 0.001$ (paired t-test), for coupled cells with the same response polarity; C, $\Delta R^2 = 0.005 \pm 0.007$, $p < 0.001$, for those with different polarities; $p = 0.054$ from Kolmogorov–Smirnov test) and mouse (D, $\Delta R^2 = 0.004 \pm 0.002$, $p < 0.001$; E, $\Delta R^2 = 0.000 \pm 0.001$, $p = 0.4$; $p < 0.001$ from Kolmogorov–Smirnov test) retinal ganglion cells. The data are shown in the same format as Figure 2.

Supplementary Figure 2: Ganglion cell pairs with low spike correlations can have strong couplings in either polarity

(A) Pairwise spike correlations of the representative cell (#25 from Figure 1B–D; auto-correlation) and all the other simultaneously recorded 19 cells (cross-correlations). Those cells with too high correlations (gray; ≥ 0.1 or ≤ -0.1 at the peak) were excluded from our model analysis to minimize the confounding effects of common visual inputs (see Methods for details). The coupling strength is shown on the bottom-right for each cell included in the LNSCN model targeting the cell #25 (black).

(B) Coupling strength from the LNSCN (top) and LNFDSCNF (bottom) models as a function of the Pearson cross-correlation of firing patterns between ganglion cell pairs (left, salamander; right, mouse). Although we selected only those cells with low spike correlations (from -0.1 to 0.1) in the coupling models, we frequently found strong couplings in either polarity. The cells in the representative data set (from A) are highlighted with yellow circles.

(C,D)   Comparison of the model parameters (C, coupling strength; D, delay) between the original and reoptimized models after resetting them to be zero (top, LNSCN; bottom, LNFDSCNF; left, salamander; right, mouse). High Pearson cross-correlation values (R) in the perturbation analysis support a good convergence of the model fitting (see Methods for details).

Supplementary Figure 3: The circuit model predictions hold with the full circuit model

The data are shown in the same format as Figure 3 but for the LNFDSCNF model.

(A,B)   Coupling strength (A; a_k in Eq.(2) in Methods) and delay (B; l_k in Eq.(1) in Methods) parameter values plotted as a function of the distance between salamander retinal ganglion cells of the same response polarity. As is the case with the LNSCN model (Figure 3), positive couplings (red, N=154) were found at a shorter distance (A; 0.31±0.14 mm versus 0.36±0.20 mm; median ± interquartile range (box plot); p<0.001, rank-sum test) and with a shorter latency (B; 2.3±4.7 versus 4.7±5.2 ms; p<0.001) than negative couplings (blue, N=64).

(C,D)   Corresponding figure panels for salamander cells with different response polarities. Couplings above the noise level were all positive (N=11) with the distance of 0.29±0.06 mm (C; median ± interquartile range) and the latency of 3.8±6.5 ms (D).

(E,F)   Corresponding figure panels for mouse retinal ganglion cells of the same response polarity. Likewise, positive couplings (red, N=37) were found at a shorter distance (C; 0.35±0.28 mm versus 0.72±0.65 mm; p=0.02) and with a shorter latency (D; 0.00±0.04 ms versus 19.1±7.7 ms; p<0.001) than negative couplings (blue, N=6). Data not analyzed for the mouse cells with different response polarities because the coupled model performance did not improve significantly (Supplementary Figure 1E).

(G–I)   Comparison of the coupling strengths from one cell to another and vise versa. Ganglion cells have symmetric couplings more frequently than expected in both salamander (G, p<0.001,

χ^2-test with $df{=}4$, between cells of the same response polarity; H, $p{<}0.001$, between cells with different response polarities) and mouse (I; $p{<}0.001$, between cells of the same response polarity) retinas. The number of data points in each category is shown in the figure panels.

Supplementary Figure 4: Ganglion cell module properties do not differ between the coupled and non-coupled models

(A) Distribution of the ganglion cell module (GCM) input correlation between coupled and non-coupled models fitted to salamander cells (black, reduced models: LNSCN versus LNSN, $0.987{\pm}0.020$, median $\pm$ interquartile range; gray, full models: LNFDSCNF versus LNFDSNF, $0.996{\pm}0.005$). The correlation was calculated using the outputs of the summation (S) stage in the models (Eq.(S9) in Real et al., 2017). High correlations indicate that the GCM input dynamics are nearly identical in response to the visual stimuli (arrow, representative cell in Figure 1B–D).

(B) The second feedback (F) stage (GCM feedback filters; Eq.(S7) in Real et al., 2017) was nearly identical between the non-coupled (left, LNFDSNF) and the coupled (right, LNFDSCNF) models (yellow, representative cell in Figure 1B–D; gray, all the other cells; red, mean).

(C) Comparison of the second nonlinear (N) stage (GCM nonlinearity) between the coupled and non-coupled models (top, LNSN versus LNSCN; bottom, LNFDSNF versus LNFDSCNF; yellow, representative cell in Figure 1B–D. Couplings did not affect the slope (left, α in Eq.(S6'); Pearson's $R{=}0.979$ and 0.997 for the reduced and full models, respectively) or the threshold (right, θ in Eq.(S6'); $R{=}0.955$ and 0.997, respectively).

(D–F) Corresponding figure panels for mouse retinal ganglion cells. There was no marked difference between the coupled and non-coupled models in GCM input correlations (D; black, reduced models, $0.994{\pm}0.005$; gray, full models, $0.998{\pm}0.003$), GCM feedback (E), GCM slope (F, left; $R{=}0.967$ and 0.978 for the reduced (top) and full (bottom) models, respectively) or GCM slope (F, right; $R{=}0.991$ and 0.996, respectively).

Supplementary Figure 5: Optic nerve stimulation produces fast excitation and slow inhibition in ganglion cell firing activity in isolated mouse retinas

The data are shown in the same format as Figures 4 and 5 but for the mouse retina.

(A,B) Firing patterns of two representative retinal ganglion cells, showing a period of suppression after the optic nerve stimulation (asterisk, antidromically evoked spikes) in the dark. Note the cell in B showed a period of enhanced firing before the suppression, while the one in A did not.

(C) Population data of the ganglion cell responses to the nerve stimulation (top row, cells with antidromic spikes; bottom row, cells without antidromic spikes): first column, no indirect effect; second and third columns, enhanced firing after the nerve stimulation (red circles, bins with significantly increased firing from the baseline); third and fourth columns, suppressed firing after the nerve stimulation (blue circles, significant decrease).

Supplementary Figure 6: Negative feedback modulates the visual response properties of mouse retinal ganglion cells

The data are shown in the same format as Figure 7 but for the mouse retina.

(A) Schematic diagram of the experiment and analysis.

(B) PSTH of a representative mouse ganglion cell with respect to the optic nerve stimulation during the visual stimulus presentation. Blue- and red-shaded bins indicate those with significantly higher and lower firing rates than the baseline, respectively.

(C,D) Linear filters of the example cell (left) and corresponding ON-OFF indices (D; Eq.(4) in Methods) at different time bins from the nerve stimulation. The filter became more biphasic during which the visual responses were enhanced by the nerve stimulation (red, 0–100 ms bin with significantly higher firing rates), but it went back to normal even when the visual responses

were suppressed (blue, 200–300 ms bin with significantly lower firing rates; gray, all the other 100 ms bins, with the mean in black).

(E,F) Normalized power spectral density (E) and the peak frequency (F; estimated by a Gaussian curve fit) of the linear filters in C.

(G,H) Static nonlinearities of the example cell (G; Eq.(5) in Methods) and the upper and lower bounds of the cell's firing probability from sigmoidal curve fits (H; Eq.(6) in Methods) at different time bins from the nerve stimulation. The upper bound was lower during the suppressed firing period (blue, 200–300 ms bin) while the lower bound was higher during the enhanced firing period (red, 0–100 ms bin) compared to the control period (gray, all the other 100 ms bins; black, mean).

(I,J) Summary of the changes in the ON-OFF indices (I; suppressed, -0.09 ± 0.08; enhanced, 0.12 ± 0.21) and the peak frequencies (J; 0.38 ± 0.83 and 0.26 ± 2.29) between the periods with and without significant firing rate changes after the nerve stimulation (blue, decrease $N=4$; red, increase $N=17$; magenta, example cell in B–H).

(E,F) Summary of the changes in the sigmoid function parameters fitted to the nonlinearity (K, upper bounds, -0.04 ± 0.05 and -0.00 ± 0.03; L, lower bounds, -0.00 ± 0.01 and 0.01 ± 0.02; center and slope, not shown; magenta, example cell in B–H).

References

Asari, H. and Meister, M. (2012). Divergence of visual channels in the inner retina. Nat Neurosci, 15, 1581–1589.

Asari, H. and Meister, M. (2014). The projective field of retinal bipolar cells and its modulation by visual context. Neuron, 81, 641–652.

Baden, T., Berens, P., Franke, K., Román Rosón, M., Bethge, M., and Euler, T. (2016). The functional diversity of retinal ganglion cells in the mouse. Nature, 529, 345–350.

Baden, T., Euler, T., and Berens, P. (2020). Understanding the retinal basis of vision across species. Nat Rev Neurosci, 21, 5–20.

Berry, M. J., Brivanlou, I. H., Jordan, T. A., and Meister, M. (1999). Anticipation of moving stimuli by the retina. Nature, 398, 334–338.

Bloomfield, S. A. and Völgyi, B. (2009). The diverse functional roles and regulation of neuronal gap junctions in the retina. Nat Rev Neurosci, 10, 495–506.

Brivanlou, I. H., Warland, D. K., and Meister, M. (1998). Mechanisms of concerted firing among retinal ganglion cells. Neuron, 20, 527–539.

Carandini, M. and Heeger, D. J. (2012). Normalization as a canonical neural computation. Nat Rev Neurosci, 13, 51–62.

Chen, X., Hsueh, H.-A., Greenberg, K., and Werblin, F. S. (2010). Three forms of spatial temporal feedforward inhibition are common to different ganglion cell types in rabbit retina. J Neurophysiol, 103, 2618–2632.

Chichilnisky, E. J. (2001). A simple white noise analysis of neuronal light responses. Network, 12, 199–213.

Cocco, S., Leibler, S., and Monassond, R. (2009). Neuronal couplings between retinal ganglion cells inferred by efficient inverse statistical physics methods. Proc Natl Acad Sci U S A, 106, 14058–14062.

Cook, J. E. and Becker, D. L. (1995). Gap junctions in the vertebrate retina. Microsc Res Tech, 31, 408–419.

Cooler, S. and Schwartz, G. W. (2021). An offset ON-OFF receptive field is created by gap junctions between distinct types of retinal ganglion cells. Nat Neurosci, 24, 105–115.

Dong, C. J. and Werblin, F. S. (1998). Temporal contrast enhancement via GABA$_c$ feedback at bipolar terminals in the tiger salamander retina. J Neurophysiol, 79, 2171–2180.

Euler, T., Haverkamp, S., Schubert, T., and Baden, T. (2014). Retinal bipolar cells: elementary building blocks of vision. Nat Rev Neurosci, 15, 507–519.

Franke, K., Berens, P., Schubert, T., Bethge, M., Euler, T., and Baden, T. (2017). Inhibition decorrelates visual feature representations in the inner retina. Nature, 542, 439–444.

Gastinger, M. J., Tian, N., Horvath, T., and Marshak, D. W. (2006). Retinopetal axons in mammals: Emphasis on histamine and serotonin. Curr Eye Res, 31, 655–667.

Geffen, M. N., de Vries, S. E. J., and Meister, M. (2007). Retinal ganglion cells can rapidly change polarity from Off to On. PLoS Biol, 5, e65.

Gollisch, T. and Meister, M. (2008). Rapid neural coding in the retina with relative spike latencies. Science, 319, 1108–1111.

Gollisch, T. and Meister, M. (2010). Eye smarter than scientists believed: Neural computations in circuits of the retina. Neuron, 65, 150–164.

Gouras, P. (1969). Antidromic responses of orthodromically identified ganglion cells in monkey retina. J Physiol, 204, 407–419.

Greschner, M., Heitman, A. K., Field, G. D., Li, P. H., Ahn, D., Sher, A., Litke, A. M., and Chichilnisky, E. J. (2016). Identification of a retinal circuit for recurrent suppression using indirect electrical imaging. Curr Biol, 26, 1935–1942.

Herz, A. V. M., Gollisch, T., Machens, C. K., and Jaeger, D. (2006). Modeling single-neuron dynamics and computations: a balance of detail and abstraction. Science, 314, 80–85.

Ito, M. (2000). Mechanisms of motor learning in the cerebellum. Brain Res, 886, 237–245.

Kenyon, G. T. and Marshak, D. W. (1998). Gap junctions with amacrine cells provide a feedback pathway for ganglion cells within the retina. Proc Biol Sci, 265, 919–925.

Koves, K., Csaki, A., and Vereczki, V. (2016). Recent research on the centrifugal visual system in mammalian species. Anat Physiol, 6, 3.

Lefebvre, J. L., Zhang, Y., Meister, M., Wang, X., and Sanes, J. R. (2008). Gamma-protocadherins regulate neuronal survival but are dispensable for circuit formation in retina. Development, 135, 4141–4151.

Maass, W. and Markram, H. (2002). On the computational power of recurrent circuits of spiking neurons. J Comput Syst Sci, 69, 593–616.

Masland, R. H. (2012). The neuronal organization of the retina. Neuron, 76, 266–280.

Mastronarde, D. N. (1983). Interactions between ganglion cells in cat retina. J Neurophysiol, 49, 350–365.

Matthews, G. (1999). Synaptic mechanisms of bipolar cell terminals. Vision Res, 39, 2469–2476.

Meister, M., Lagnado, L., and Baylor, D. A. (1995). Concerted signaling by retinal ganglion cells. Science, 270, 1207–1210.

Nath, A. and Schwartz, G. W. (2017). Electrical synapses convey orientation selectivity in the mouse retina. Nat Commun, 8, 2025.

Nirenberg, S. and Meister, M. (1997). The light response of retinal ganglion cells is truncated by a displaced amacrine circuit. Neuron, 18, 637–650.

Pereda, A. E., Curti, S., Hoge, G., Cachope, R., Flores, C. E., and Rash, J. E. (2013). Gap junction-mediated electrical transmission: regulatory mechanisms and plasticity. Biochim Biophys Acta, 1828, 134–146.

Pillow, J. W., Shlens, J., Paninski, L., Sher, A., Litke, A. M., Chichilnisky, E. J., and Simoncelli, E. P. (2008). Spatio-temporal correlations and visual signalling in a complete neuronal population. Nature, 454, 995–999.

Pouzat, C., Mazor, O., and Laurent, G. (2002). Using noise signature to optimize spike-sorting and to assess neuronal classification quality. J Neurosci Methods, 122, 43–57.

Puller, C., Duda, S., Lotfi, E., Arzhangnia, Y., Block, C. T., Ahlers, M. T., and Greschner, M. (2020). Electrical coupling of heterotypic ganglion cells in the mammalian retina. J Neurosci, 40, 1302–1310.

Real, E., Asari, H., Gollisch, T., and Meister, M. (2017). Neural circuit inference from function to structure. Curr Biol, 27, 189–198.

Repérant, J., Ward, R., Miceli, D., Rio, J. P., Médina, M., Kenigfest, N. B., and Vesselkin, N. P. (2006). The centrifugal visual system of vertebrates: a comparative analysis of its functional anatomical organization. Brain research reviews, 52, 1–57.

Roska, B., Molnar, A., and Werblin, F. S. (2006). Parallel processing in retinal ganglion cells: how integration of space-time patterns of excitation and inhibition form the spiking output. J Neurophysiol, 95, 3810–3822.

Roska, B., Nemeth, E., Orzo, L., and Werblin, F. S. (2000). Three levels of lateral inhibition: A space-time study of the retina of the tiger salamander. J Neurosci, 20, 1941–1951.

Roy, K., Kumar, S., and Bloomfield, S. A. (2017). Gap junctional coupling between retinal amacrine and ganglion cells underlies coherent activity integral to global object perception. Proc Natl Acad Sci U S A, 114, E10484–E10493.

Roy, S. and Field, G. D. (2019). Dopaminergic modulation of retinal processing from starlight to sunlight. J Pharmacol Sci, 140, 86–93.

Sakai, H. M. and Naka, K. I. (1988). Dissection of the neuron network in the catfish inner retina. II. Interactions between ganglion cells. J Neurophysiol, 60, 1568–1583.

Sakai, H. M. and Naka, K. I. (1990). Dissection of the neuron network in the catfish inner retina. IV. bidirectional interactions between amacrine and ganglion cells. J Neurophysiol, 63, 105–119.

Stys, P. K., Ransom, B. R., and Waxman, S. G. (1991). Compound action potential of nerve recorded by suction electrode: a theoretical and experimental analysis. Brain Res, 546, 18–32.

Tachibana, M. and Kaneko, A. (1988). Retinal bipolar cells receive negative feedback input from GABAergic amacrine cells. Vis Neurosci, 1, 297–305.

Trong, P. K. and Rieke, F. (2008). Origin of correlated activity between parasol retinal ganglion cells. Nat Neurosci, 11, 1343–1351.

Van Rullen, R. and Thorpe, S. J. (2001). Rate coding versus temporal order coding: What the retinal ganglion cells tell the visual cortex. Network: Comput Neural Syst, 13, 1255–1283.

Victor, J. D. (1999). Temporal aspects of neural coding in the retina and lateral geniculate. Network: Comput Neural Syst, 10, R1–R66.

Völgyi, B., Chheda, S., and Bloomfield, S. A. (2009). Tracer coupling patterns of the ganglion cell subtypes in the mouse retina. J Comp Neurol, 512, 664–687.

Völgyi, B., Pan, F., Paul, D. L., Wang, J. T., Huberman, A. D., and Bloomfield, S. A. (2013). Gap junctions are essential for generating the correlated spike activity of neighboring retinal ganglion cells. PloS one, 8, e69426.

Wang, X.-J. (2008). Decision making in recurrent neuronal circuits. Neuron, 60, 215–234.

Wark, B., Fairhall, A., and Rieke, F. (2009). Timescales of inference in visual adaptation. Neuron, 61, 750–761.

Wässle, H. (2004). Parallel processing in the mammalian retina. Nat Rev Neurosci, 5, 747–757.

Werblin, F. S. (2011). The retinal hypercircuit: a repeating synaptic interactive motif underlying visual function. J Physiol, 589, 3691–3702.

Witkovsky, P. and Dearry, A. (1991). Functional roles of dopamine in the vertebrate retina. Prog Retinal Res, 11, 247–292.

Wu, M. C.-K., David, S. V., and Gallant, J. L. (2006). Complete functional characterization of sensory neurons by system identification. Annu Rev Neurosci, 29, 477–505.

Zhang, A.-J. and Wu, S. M. (2010). Responses and receptive fields of amacrine cells and ganglion cells in the salamander retina. Vision Res, 50, 614–622.

Zhang, D.-Q., Belenky, M. A., Sollars, P. J., Pickard, G. E., and McMahon, D. G. (2012). Melanopsin mediates retrograde visual signaling in the retina. PloS one, 7, e42647.

Zhang, D.-Q., Wong, K. Y., Sollars, P. J., Berson, D. M., Pickard, G. E., and McMahon, D. G. (2008). Intraretinal signaling by ganglion cell photoreceptors to dopaminergic amacrine neurons. Proceedings of the National Academy of Sciences of the United States of America, 105, 14181–14186.

	Coupling strength (%)	Delay (ms)
cell 33	-4.0	6.1
cell 40	-13.0	6.9
cell 46	21.0	3.6
cell 50	5.0	0.8
cell 56	-7.6	7.6
cell 59	-14.9	6.8

Change in model performance (ΔR²)
LNSCN vs. LNSN

Change in model performance (ΔR²)
LNFDSCNF vs. LNFDSNF