Identification of novel avian and mammalian deltaviruses provides new insights

into deltavirus evolution

Masashi Iwamoto[1,2], Yukino Shibata[3], Junna Kawasaki[4,5], Shohei Kojima[4,6], Yung-Tsung

Li[7], Shingo Iwami[2], Masamichi Muramatsu[1], Hui-Lin Wu[7,8], Kazuhiro Wada[3], Keizo

Tomonaga[4,5,9], Koichi Watashi[1,10], Masayuki Horie[4,11,*]

[1]Department of Virology II, National Institute of Infectious Diseases, Tokyo, Japan

[2]Mathematical Biology Laboratory, Department of Biology, Faculty of Sciences, Kyushu University,

Fukuoka, Japan

[3]Graduate School of Life Science, Hokkaido University, Sapporo, Hokkaido, Japan

[4]Institute for Frontier Life and Medical Science, Kyoto University, Kyoto, Japan

[5]Department of Mammalian Regulatory Network, Graduate School of Biostudies, Kyoto University,

Kyoto, Japan

[6]Genome Immunobiology RIKEN Hakubi Research Team, RIKEN Cluster for Pioneering Research

[7]Hepatitis Research Center, National Taiwan University Hospital, Taipei, Taiwan

[8]Graduate Institute of Clinical Medicine, National Taiwan University College of Medicine

[9]Department of Molecular Virology, Graduate School of Medicine, Kyoto University, Kyoto, Japan

[10]Department of Applied Biological Sciences, Tokyo University of Science, Noda, Japan

[11]Hakubi Center for Advanced Research, Kyoto University

*Corresponding author

Masayuki Horie, DVM, PhD

Hakubi Center for Advanced Research, Kyoto University

53 Kawahara-cho, Shogo-in, Sakyo, Kyoto 606-8507, Japan

Email: horie.masayuki.3m@kyoto-u.ac.jp

Abstract

Hepatitis delta virus (HDV) is a satellite virus that requires hepadnavirus envelope

proteins for its transmission. Although recent studies identified HDV-related deltaviruses

in certain animals, the evolution of deltaviruses, such as the origin of HDV and the

mechanism of its coevolution with its helper viruses, is unknown, mainly because of the

phylogenetic gaps among deltaviruses. Here we identified novel deltaviruses of passerine

birds, woodchucks, and white-tailed deer by extensive database searches and molecular

surveillance. Phylogenetic and molecular epidemiological analyses suggest that HDV

originated from mammalian deltaviruses and the past interspecies transmission of

mammalian and passerine deltaviruses. Further, metaviromic and experimental analyses

suggest that the satellite-helper relationship between HDV and hepadnavirus was

established after the divergence of the HDV lineage from non-HDV mammalian

deltaviruses. Our findings enhance our understanding of deltavirus evolution, diversity,

and transmission, indicating the importance of further surveillance for deltaviruses.

Introduction

Hepatitis delta virus (HDV) is the only member of the genus *Deltavirus* which is not

assigned to a family (1). The HDV genome is an approximately 1.7-kb circular, negative

single-stranded RNA, harboring a single open reading frame (ORF) encoding the small

and large hepatitis delta antigens (S-HDAg, 24 kDa and L-HDAg, 27 kDa) that are

translated from the same transcriptional unit via RNA-editing of the stop codon, which is

catalyzed by the host protein ADAR1 (2-5). The 19 amino acid residue extension of the

C-terminal region of L-HDAg contains a farnesylation site required to interact with

helper virus envelope proteins (6). The genome structure of HDV is unique in that it has

genomic and antigenomic ribozymes, which are essential for its replication (7, 8), and is

highly self-complementarity, generating a rod-like structure (9-11). Although HDV can

autonomously replicate, it requires an envelope protein of other "helper" viruses to

produce infectious virions. Hepatitis B virus (HBV) (family *Hepadnaviridae*) provides

the envelope proteins required for HDV transmission between humans (12).

Approximately 15–20 million people worldwide are estimated to be infected with HDV

among 350 million HBV carriers (13). Compared with mono-infection with HBV,

coinfection of HDV and HBV accelerates the pathogenic effects of HBV, such as severe

or fulminant hepatitis and progression to hepatocellular carcinoma, through unknown

mechanisms (14).

The evolutionary origin of HDV presents an enigma. However, recent discoveries of

deltaviruses of vertebrate and invertebrate species (15-18), significantly changed our

understanding of deltavirus evolution. These non-HDV deltaviruses are distantly related

to HDV but may share the same origin because of their similarly structured circular RNA

genomes (approximately 1.7 kb), which encode DAg-like proteins, possess ribozymes

sequences, and are highly self-complementary (15-18). These findings provide clues to

the mechanism of deltavirus evolution. For example, a recent study hypothesizes that

mammalian deltaviruses codiverged with their host mammalian species (18). However,

the few known deltaviruses are highly divergent (15-18). Therefore, the phylogenetic

gaps between the deltaviruses must be filled through the identification of putative novel

deltaviruses.

The discoveries of non-HDV deltaviruses provides insights into the relationships

between deltaviruses and their helper viruses. Recently identified non-HDV deltaviruses

likely do not coinfect with hepadnaviruses, suggesting the presence of other helper

viruses (15-18). This hypothesis is supported by absence of a large isoform of DAg,

which is required for the interaction of HDV with the HBV envelope proteins, in rodent

deltavirus (18). Further, viral envelope proteins of reptarenavirus and hartmanivirus, but

not HBV, confer infectivity upon the snake deltavirus (19). These findings suggest that

hepadnaviruses do not serve as helper viruses for non-HDV deltaviruses and that the

deltavirus-hepadnavirus relationship is specific to the HDV lineage. However, the large

phylogenetic gap between HDV and the few other deltaviruses makes it difficult to assess

the hypothesis, raising the importance of further research.

In this study, to understand the evolution of deltaviruses, we analyzed publicly

available transcriptome data and found novel mammalian and avian deltaviruses. Our

phylogenetic analysis suggests that HDVs originated from non-HDV mammalian

deltaviruses and does not support the codiversification hypothesis of deltavirus and

mammalian evolution. Moreover, *in silico* and experimental analyses, together with

previous findings, suggest that the satellite-helper relationship between HDV and

hepadnavirus was established after the divergence of the novel non-HDV mammalian

deltaviruses and the HDV lineage. Further, we present evidence for recent interfamily

transmission of deltaviruses among passerine birds. Our findings therefore provide novel

insights into the evolution of deltaviruses.

Results

Identification of deltavirus-related sequences in avian and mammalian transcriptomes

We first assembled 46,359 RNA-seq data of birds and mammals. Using the resultant

contigs as queries, we identified five deltavirus-related contigs in the SRA data of birds

and mammals, including the zebra finch (*Taeniopygia guttata*), common canary (*Serinus*

canaria), Gouldian finch (*Erythrura gouldiae*), Eastern woodchuck (*Marmota monax*),

and white-tailed deer (*Odocoileus virginianus*). We named the deltavirus-like sequences

Taeniopygia guttata deltavirus (tgDeV), Serinus canaria-associated deltavirus (scDeV),

Erythrura gouldiae deltavirus (egDeV), Marmota monax deltavirus (mmDeV), and

Odocoileus virginianus (ovDeV), respectively (Table 1).

The amino acid sequences tlanslated from the contigs are 36.0%–66.7% identical to

those of the DAg proteins of known deltaviruses (Table 1, Supp Tables 1 and 2). The

tgDeV, mmDeV, and ovDeV contigs, which comprise approximately 1,700 nucleotides,

encode one ORF with a sequence similar to those of DAg genes of known deltaviruses

(Fig 1a, Table 1, Supp Tables 1 and 2). In contrast, the contigs scDeV and egDeV are 761

and 596 nucleotides in length, respectively (Fig. 1b, Supp Tables 1 and 2). Note that the

nucleotide sequences of tgDeV and egDeV are 97.7% identical, and we therefore

analyzed tgDeV instead of tgDeV and egDeV.

Genome structures of novel avian and mammalian deltaviruses

The three contigs (tgDeV, mmDeV, and ovDeV) are almost identical in length to the

full-length genomes of known deltaviruses. We therefore checked for potential circularity

of the contigs. Dot-plot analyses revealed that each of both ends of these three contigs is

identical (Supp Fig. 1), suggesting that the contigs were derived from circular RNAs. We

further mapped the original RNA-seq data to the corresponding circularized contigs using

the Geneious mapper, revealing that some of the reads properly spanned the junctions

(data not shown), indicating that these contigs are derived from circular RNAs. Therefore,

we designated the resultant circular contigs of tgDeV, mmDeV, and ovDeV as full-length

novel deltavirus genomes (1,706, 1,712, and 1,690 nucleotides, respectively) (Fig. 1a).

These novel genomes are characterized by high self-complementarity, genomic and

antigenomic ribozymes, and poly(A) signals, which are conserved among known

deltaviral genomes (Fig. 1 and Supp Table 2) (15-18). Further, the predicted secondary

structures of the ribozymes are highly similar to those of HDV as well as those of other

deltaviruses (Supp Fig. 2).

Characterization of DAg proteins encoded by the novel deltaviruses

We next characterized the putative DAg proteins encoded by the novel deltaviruses. Most

of their biochemical features, biologically relevant amino acid residues, and functional

domains (15-18) are conserved among the DAg proteins (Fig. 2a). The isoelectric points

of DAg proteins from the novel deltaviruses range from 10.35 to 10.63 (Supp Table 2),

which are nearly identical to those of known deltaviruses. All the post-translational

modification sites in HDAg are conserved among those of all DAg proteins of the novel

deltaviruses, except for the serine phosphorylation site on scDeV-DAg (Fig. 2a). The

NLS is conserved among the DAg proteins, although location of the predicted NLS of

scDeV DAg protein differs (Fig. 2a).

We next investigated whether the novel deltaviruses utilize A-to-I RNA-editing. To

answer this question, we mapped short reads of the SRA data, which we initially used to

detect the deltaviruses, to identify the nucleotide variations among the stop codons. We

found a potential RNA-editing site within the stop codon of the ovDeV-DAg gene, in

which there was 0.4% nucleotide variation (5 of 1160 reads) at the second nucleotide

position of the stop codon (U<u>A</u>G), all of which were G instead of the consensus

nucleotide A (Fig. 2b). The quality scores of the five G variants ranged from 35 to 41

(Supp Fig. 3), which likely exclude the possibility of a sequencing error. This variation

may be explained by A-to-I editing by ADAR1, as known for HDV (2). However,

possible RNA-editing generates a protein two amino acid residues longer because of a

stop codon immediately downstream (Fig. 2c). Further, the C-terminal farnesylation

motif (CXXQ) required for the interaction with hepadnaviral envelope proteins (6) was

absent from the longer product. These observations suggest that even if RNA-editing

occurs, the resultant gene product does not contribute to the interaction with hepadnaviral

envelope proteins. Further, we were unable to identify nucleotide variations of the

mapped reads at the stop codons in the genomes of tgDeV and mmDeV (data not shown).

The novel deltaviruses potentially replicate in their hosts

To determine whether the novel deltaviruses potentially replicate in their respective,

putative host species, we evaluated the mapping pattern of viral reads described above.

We found that the read depths of the predicted transcribed regions (the DAg coding

regions to poly-A signals) were much greater than those of the other genomic regions (Fig.

3), indicating that most viral reads were derived from viral mRNAs. These findings

suggest that the novel deltaviruses replicate in their hosts.

The mapping pattern on tgDeV differed slightly from the others. Specifically,

although the read depth of the DAg ORF region was higher, the reads represented only

80% of the ORF (Fig. 3a). This trend was apparent in another tgDeV-positive RNA-seq

data (Supp Fig. 4). However, it is not clear whether this is attributed to an artifact or

actually reflects the transcription pattern of tgDeV.

Transmission of tgDeV- and tgDeV-like viruses among passerine birds

We next investigated whether the novel deltaviruses are transmitted among animal

populations. We first analyzed tgDeV infections in birds using RNA-seq data (Table 2

and Supp Table 3). Among 6453 SRA data, tgDeV-derived reads were identified in 34

SRAs, including the SRAs in which tgDeV and egDeV were initially detected. The 34

tgDeV-positive SRA data were obtained from tissues such as blood, kidney, and muscles,

suggesting broad tropism and viremia, or systemic infection, or both, with tgDeV.

Further, tgDeV sequences were detected in several bird species such as the

black-headed bunting (*Emberiza melanocephala*) and yellow-bellied tit (*Pardaliparus*

venustulus). All tgDeV-positive bird species belong to the order Passeriformes. These

tgDeV-positive SRA data are included in the nine BioProjects deposited by independent

researchers, and thus the birds were likely from different sources. Further, the

tgDeV-positive sample in SRR9899549 (BioSample accession: SAMN12493457) is

derived from a black-headed bunting caught in the wild. These data suggest that tgDeV

(or tgDeV-like viruses) circulate among diverse passerine birds, even in the wild.

During the above analysis, we found that SRA data from the yellow-bellied tit

(SRR7244693 and SRR7244695–SRR7244698) contain many reads mapped to the

tgDeV genome. Therefore, we extracted the mapped reads of SRA data and performed *de*

novo assembly. We obtained a 1707-nt circular complete genome sequence, which we

designated pvDeV. The pvDeV nucleotide sequence is 98.2% identical to that of tgDeV,

and the properties of its DAg protein sequence are similar to those of tgDeV DAg (Supp

Fig. 5).

We next employed real-time RT-PCR to further evaluate potential deltavirus

infections of passerine birds. We analyzed 30 and 5 whole-blood samples from zebra and

Bengalese finches (*Lonchura striata* var. *domestica*), respectively, and found that one

Bengalese finch was positive for real-time RT-PCR test. To exclude the possibility of

contamination of a plasmid used as a control for real-time PCR, we performed RT-PCR

using a primer set that distinguishes viral from plasmid amplicons (Figs. 4a and b). We

obtained a band of the expected size only from the cDNA sample (Fig. 4c), revealing that

the bird was truly positive for a tgDeV-like virus. Therefore, we named this virus lsDeV,

and further analysis revealed that its full-length genome nucleotide sequence (1708 nt) is

98.2% and 98.4% identical to those of tgDeV and pvDeV, respectively. Moreover, its

genome features are almost identical to those of tgDeV and pvDeV (Supp Fig. 5).

Evidence for recent interfamily transmission of deltaviruses among passerine birds

We found that the sequence similarities among the passerine deltaviruses (tgDeV, pvDeV,

and ls DeV) (Fig. 4d) were not consistent with evolutionary codivergence. According to

the TimeTree (20), deltavirus-positive passerine birds diverged approximately 44 million

years ago (Fig. 4e and Supp Fig. 6). Considering the rapid evolutionary rates of HDVs

(approximately 10^{-3} substitutions per site per year) (21-23), it is unlikely that these

viruses codiverged with their hosts. Most likely, interfamily transmission occurred

relatively recently among passerine birds.

Transmission of mmDeV among woodchucks

We similarly analyzed mmDeV infections using SRA data for the order Rodentia, other

than mice (*Mus musculus*) and rats (*Rattus norvegicus*). Our analysis of 4776 SRA

datasets detected mmDeV reads in 20 SRA data of seven woodchucks (Table 2 and Supp

Table 3). Although these mmDeV-positive SRA data were contributed by the same

research group, the animals were apparently obtained at different times (24, 25),

suggesting that mmDeV was transmitted among woodchucks. The mmDeV-positive SRA

data are derived from samples of liver or peripheral blood mononuclear cells.

We next used real-time RT-PCR to analyze 81 woodchuck samples (liver, n= 43;

serum, n = 38). However, mmDeV was undetectable (data not shown), which may be

explained by the absence of mmDeV infection or clearance, low level of infection, or

both.

No evidence of transmission of other deltaviruses

We next focused on ovDeV and scDeV sequences of ruminant animals and passerine

birds, respectively (Table 2 and Supp Table 3). We detected ovDeV-derived reads only in

five SRA data. The SRA data were obtained from brain, muscle, testis, pedicles, and

antlers, suggesting systemic infection, viremia, or both. However, we were unable to

determine if these samples were derived from multiple individuals. We detected

scDeV-derived reads only from the SRA data in which the virus was initially detected. We

therefore were unable to provide evidence for the transmission of ovDeV and scDeV in

their host animals.

Phylogenetic relationships among deltaviruses

To decipher the evolutionary relationships among deltaviruses, we conducted a

phylogenetic analysis using known and the novel deltavirus sequences discovered here.

We did not include sequences of recently identified fish, toad, newt, termite, and

duck-associated deltaviruses because they share very low amino acid sequence identities

with the novel deltaviruses as well as with HDVs (Fig. 5a), which may reduce the

accuracy of tree (16). We further excluded scDeV for this reason, and we did not include

tgDeV-like viruses, because their sequences are nearly identical to that of tgDeV. The

reconstructed tree shows that the newly identified tgDeV forms a strongly supported

cluster with snake DeV and rodent DeV, although they are distantly related to each other

(Fig. 5b). Note that mmDeV and ovDeV are more closely related to HDVs than the other

deltaviruses.

Candidate helper viruses

To gain insights into helper viruses of the novel deltaviruses, we first analyzed the

coexisting viruses in the SRA data using BLASTx. Note that we omitted experimental

woodchuck hepatitis virus (WHV) infections associated with the mmDeV-positive

woodchuck-derived SRA data (SRR2136864 to SRR2136999). We also excluded viruses

that infect invertebrates and endogenous retroviruses as well. These analyses reveal that

polyomavirus, bornavirus, and circovirus sequences are present in the deltavirus-positive

SRA data of passerine birds (Table 3 and Supp Table 4). Further, we detected contigs with

98%–100% identities to human viruses (human mastadenovirus or mammalian

rubulavirus 5) (Supp Table 4), although these may represent contamination, index

hopping, or both. Among these viruses, only the genome of bornavirus encodes an

envelope protein. Note that scDeV and bornavirus-positive SRA data was obtained from

pooled samples (SAMN04260514), and we therefore were unable to determine whether

scDeV and canary bornavirus 3 infected the same individual.

We next cross-referenced the mmDeV reads and the metadata, which also provide

insights into the mmDeV helper virus. Among 20 mmDeV-positive SRA data, 18 were

obtained from animals experimentally infected with the hepadnavirus WHV, which was

experimentally shown to serve as a helper virus for HDV (26, 27). However, the other

two SRA data (SRR437934 and SRR437938) were derived from animals negative for

antibodies against WHV as well as WHV DNA (24). These observations suggest that

mmDeV was transmitted to the two animals without WHV and that WHV therefore was

not the helper virus for mmDeV that infected these two individuals.

Replication of tgDeV and mmDeV in human and woodchuck cell lines

To investigate whether the novel deltavirus sequences are replicable or not, we performed

transfection-based assays in cell culture systems. We constructed plasmid expression

vectors harboring the minus-strand genome of the tgDeV or mmDeV dimer sequence

under the transcriptional control of the CMV promoter (see Materials and Methods). We

first determined if the replication initiated by transfecting these plasmids. These plasmids

express the minus-strand genome and therefore DAg protein is expressed if the viral

genome replicates (19). We transfected the plasmid vectors into Huh7 human hepatic

cells and WCH-17 woodchuck hepatic cells and used western blotting (Figs. 6a and b)

and immunofluorescence assay (IFA) (Figs. 6c–f) to detect the expression of DAg

proteins. Western blotting detected the expected bands (approximately 22 kDa) only in

transfected cells (Figs. 6a and b). Note that a single specific band was detected in lysates

prepared from each cell type, suggesting that tgDeV and mmDeV expressed only one

DAg isoform. Consistent with the above results, specific signals were observed only in

the transfected cells in IFA (Figs. 6c and d, red signals). Together, these data suggest that

the tgDeV and mmDeV initiated replication from the constructed plasmids in the cell

culture system.

The DAg proteins predominantly localized to the nucleus 2 days post-transfection

(Figs. 6e and f). Interestingly, large viral speckles were observed in the nucleus, similar to

those detected in cells infected with HDV (28, 29). These results suggest that tgDeV and

mmDeV employ a nuclear replication strategy similar to that used by HDV.

HBV envelope proteins do not contribute to the production of infectious tgDeV and

mmDeV

As described in the above section "*Candidate helper viruses*", there is no evidence of

coinfections of hepadnaviruses with tgDeV or mmDeV. However, this does not

necessarily mean hepadnaviruses do not serve as helper viruses for the novel deltaviruses.

To determine whether tgDeV or mmDeV utilize the HBV envelope proteins (HBs), we

transfected the deltavirus expression plasmids together with an HBs expression vector or

the cognate empty vector into Huh7 cells. The culture supernatants were incubated with

HepG2-NTCP cells, which are susceptible to HBs-dependent HDV infection (30, 31).

HDV served as a control to monitor HBs-dependent virus release and subsequent cell

entry. Viral RNA was undetected in supernatants of cells that did not express HBs (Fig.

7a). In contrast, cotransfection of the HBs plasmid released large amounts of HDV RNA

into the supernatant, consistent with published data (32), whereas tgDeV or mmDeV

RNA was undetectable (Fig. 7a).

We next measured the amounts of viral RNA and detected DAg protein in

HepG2-NTCP cells 7 days after incubation with the supernatants. HDV RNA and DAg

protein were highly expressed in the infected cells (Fig. 7b and c). HDV infection was

inhibited by the preS1 peptide (Myrcludex B), which was shown to inhibit

HBs-dependent HDV infection (33). These indicate that HDV infection is indeed

mediated by the HBs. On the other hand, tgDeV and mmDeV RNA or DAg protein was

undetectable, suggesting that HBs do not contribute to the production of infectious tgDeV

or mmDeV.

Discussion

Important aspects of the evolution of deltaviruses are unknown, such as the origin of

HDV and the coevolution of deltaviruses and their helper viruses, mainly because few

deltaviruses are known, and they are highly genetically divergent (15-18). Therefore, the

resulting large gaps in the deltavirus phylogenetic tree create a formidable obstacle to

understanding deltavirus evolution. Here we identified five complete genomes of novel

deltaviruses from birds and mammals (Fig. 1 and Supp Fig. 5), which partially fill these

phylogenetic gaps (Fig. 5b). Moreover, our present findings reveal that the evolution of

deltaviruses is much more complicated than previously thought. For example, one

hypothesis states that mammalian deltaviruses codiverged with their mammalian hosts

(18). However, our phylogenetic analysis shows that the tree topology of mammalian

deltaviruses is incongruent with their hosts'. For example, ovDeV, which we detected in

deer, is most closely related to human HDV (Fig. 5b). Further, the distantly related

mmDeV and rodent DeV (detected in *Proechimys semispinosus* (18)) were detected in

rodent species. These data suggest that deltaviruses were transmitted among mammalian

species and did not always codiverge with their hosts. Moreover, we discovered recent

interfamily transmission of passerine deltaviruses (tgDeV and its relatives) (Fig. 4e).

Therefore, avian and mammalian deltaviruses may have, at least partially, evolved by

interspecies transmission.

Our present phylogenetic analysis also gives insights into the origin of human HDVs.

As described above, ovDeV and mmDeV are close relatives of human HDVs, suggesting

that HDVs arose from other mammalian deltaviruses. Recent studies on the phylogeny of

bat deltaviruses support our findings and conclusions (34, 35) (Supp Fig. 7). However, we

were unable to exclude the possibility of infection of animal lineages apart from

mammals with unknown deltaviruses phylogenetically located between those viral

lineages. Further investigations are required for a better understanding of the deltavirus

evolution.

There is a paucity of knowledge about helper viruses for non-HDV deltaviruses,

other than the snake deltavirus (19), although evidence indicates that hepadnaviruses may

not serve as helper viruses for the novel deltaviruses discovered here, as suggested for

other non-HDV deltaviruses (15-18). Here we only detected bornavirus, circovirus, and

polyomavirus, but not hepadnavirus sequences in association with deltavirus-positive

SRA data (Table 3). Further, mmDeV was detected in two woodchuck individuals that

were demonstrated to be negative for WHV (Table 2 and Supp Table 3). Moreover, we

found that HBs did not contribute to the formation of infectious tgDeV and mmDeV in

cell culture experiments. These observations suggest that hepadnaviruses do not serve as

helper viruses for the novel non-HDV deltaviruses detected here.   Further, we were

unable to demonstrate that the deltaviruses identified here express proteins similar to the

L-HDAg protein (Figs. 6a and b), which is expressed via RNA-editing and is essential for

HDV to interact with HBs (2, 37). Although RNA-editing may alter the stop codon of

ovDeV DAg, this does not lead to the expression of a large isoform of DAg (L-DAg)

protein containing a farnesylation site (Figs. 2b and c). The lack of L-DAg expression

was also observed in rodent deltaviruses (18). Therefore, L-DAg expression phenotype

may have been acquired after the divergence of ovDeV and the HDV lineage (Fig. 5b).

Among the coexisting viruses, only bornavirus produces an envelope glycoprotein

(G protein), which might be used by non-HDV deltaviruses to produce infectious virions.

Indeed, snake deltavirus utilizes the envelope proteins of reptarenaviruses and

hartmaniviruses to produce infectious particles (19). Furthre, HDV forms infectious

virions using envelope proteins *in vitro* of RNA viruses such as vesiculovirus and

hepacivirus (38). Therefore, the bornavirus G protein might envelop non-HDV

deltaviruses.

In contrast, the coexisting viruses, circoviruses and polyomaviruses, are

nonenveloped. Therefore, it is unlikely that these viruses can serve as helper viruses for

the deltaviruses. However, we cannot exclude the possibility that these viral capsid

proteins might contribute to the transmissibility of deltaviruses through unknown

mechanisms. Additionally, virus-derived sequences in host genomes, called endogenous

viral elements (EVEs), might mediate the formation of infectious particles. Here detected

the expression of retroviral EVEs in certain deltavirus-positive SRA data (data not

shown). Although HDVs do not use retroviral envelope proteins (38), non-HDV

deltaviruses might utilize strategies distinct from those employed by HDVs. Alternatively,

non-HDV deltaviruses may not require helper viruses and utilize extracellular vesicles for

transmission. Further biological experiments, together with molecular surveillance, are

therefore required to understand the satellite-helper relationships of deltaviruses.

Here we show that the sequences of tgDeV and tgDeV-like viruses, such as pvDeV

and lsDeV, are relatively closely related to known vertebrate deltaviruses (Fig. 5b).

Although a previous study found a deltavirus from duck, this duck-associated virus was

detected in oropharyngeal/cloacal swabs and is distantly related to vertebrate deltaviruses,

suggesting the possibility of its dietary origin (15, 18). This may be true for scDeV

studied here. scDeV was detected in skin (Table 1). Although scDeV was excluded from

our phylogenetic analysis, the amino acid identities between the DAg protein of scDeV

with other vertebrate deltaviruses range from 32.7%–39.5% (Fig. 5a). Therefore, scDeV

may be derived from contaminants, which should be addressed in the future. In contrast,

tgDeV and tgDeV-like viruses were detected in tissues such as the spleen and muscles

(Table 2), suggesting that tgDeV and tgDeV-like viruses are authentic avian deltaviruses.

Here we show that certain novel deltaviruses are transmitted among animal

populations (Table 2 and Supp Table 3). Note that few reads were mapped to the virus

genomes in some SRA data, which may be attributed to index hopping (39-43) from SRA

data containing numerous deltavirus-derived reads. Therefore, these data should be

interpreted with caution. Nevertheless, our conclusions are not affected, because they are

supported by robust data (Table 2). For data in which index hopping has possibly

occurred, further analyses are needed to confirm deltavirus infections.

Our present analysis provides virological insights into important characteristics of

deltavirus infections, such as tissue and host tropism. For example, infections with tgDeV

(and tgDeV-like viruses), mmDeV, and ovDeV were not limited to the liver and were

detected in at least two different tissues (Table 2). These observations are consistent with

those of previous studies that non-HDV deltaviruses in multiple organs and blood and

that they replicate in numerous cell types (18, 19). Therefore non-HDV deltaviruses may

infect diverse tissues and cause systemic infection, viremia, or both. Furthre, tgDeV and

mmDeV replicated in human and woodchuck cells (Fig. 6), which is consistent with the

ability of the snake deltavirus to replicate in mammalian cells (19). These observations

suggest that the host range of deltaviruses is broad and that the helper viruses of

non-HDV deltaviruses may be the determinants of host range.

Our analyses further suggest that tgDeV and mmDeV are sensitive to host immune

responses. We cross-referenced tgDeV reads and metadata and made an intriguing

finding that may contribute to the virus-host interaction. BioProject PRJNA297576

contains 12 RNA-seq data for six zebra finches (44). Interestingly, the tgDeV reads were

almost exclusively detected in birds treated with testosterone vs the controls (Supp Fig.

8a and Supp Table 5). Therefore, the immunosuppressive effects of testosterone (45) may

increase the transcription or replication of tgDeV, or both, to enable detection using

RNA-seq. Furthre, when we cross-referenced the mmDeV reads and the metadata, we

found that 18 of 20 mmDeV-positive SRA data derived from five individuals were

acquired through an experiment lasting 27 weeks (PRJNA291589) (25). Among 18 SRA

data, those of one individual (ID 1008) provide insights into mmDeV infection as

follows: At first (–3 weeks), mapped reads were not detected, although the proportion of

mapped reads were highest at one week and then drastically decreased (Supp Fig. 8b and

Supp Table 6). Interestingly, a previous study suggested that the host's immune response

can clear rodent deltaviruses (18). Our present observations together with this previous

finding, suggest that the host immune response suppresses and then clear deltavirus

infections. This may explain the low prevalence of RT-PCR-positive samples of

woodchucks and passerine birds as described in the Results (sections "*Circulation of*

tgDeV and tgDeV-like viruses among passerine birds" and "*Circulation of mmDeV in*

woodchucks"). Note that latent or low levels of persistent deltavirus infections may occur.

Indeed, snake deltavirus establishes a persistent infection in a cell culture system (19).

Therefore, deltaviruses might persistently infect host cells with a low level of virus

replication, and some stimulations, such as immunosuppression, may trigger robust virus

replication. Further studies are therefore required to understand the interactions between

deltaviruses and their hosts.

Together, our present data contribute to a deeper understanding of the evolution of

deltaviruses and suggest the presence of undiscovered deltaviruses that infect diverse

animal species. Further investigations will provide further insights into deltavirus

evolution.

Materials and methods

Detection of deltaviruses from publicly available transcriptome data

Paired-end, RNA-seq data from birds and mammals were downloaded from NCBI SRA
(46). The SRA accession numbers used in this study are listed in Supplementary material.
The downloaded SRA files were dumped using pfastq-dump (DOI:
10.5281/zenodo.2590842; https://github.com/inutano/pfastq-dump), and then
preprocessed using fastp 0.20.0 (47). If genome data of either the same species or the
same genus were available, the preprocessed reads were mapped to the corresponding
genome sequences (the genome information is available upon request) by HISAT2 2.1.0
(48), and then unmapped paired-end reads were extracted using SAMtools 1.9 (49) and
Picard 2.20.4 (http://broadinstitute.github.io/picard/). The extracted unmapped reads
were used for *de novo* assembly. If genome data were unavailable, the preprocessed reads
were directly used for *de novo* assembly. *De novo* assembly was conducted using SPAdes
(50) and/or metaSPAdes (51) 3.13.0 with k-mer of 21, 33, 55, 77, and 99. The resultant
contigs were clustered by cd-hit-est 4.8.1 (52, 53) with a threshold of 0.95. Finally, the
clustered contigs $\geq$ 500 nt were extracted by SeqKit 0.9.0 (54), and they were used for the
downstream analyses.

Two-step sequence similarity searches were performed to identify RNA virus-like
sequences. First, BLASTx searches were performed against a custom database for RNA
viruses using the obtained contigs as queries employing BLAST+ 2.9.0 (55) with the
following options: -word_size 2, -evalue 1e^{-3}, max_target_seqs 1. The custom database of
RNA viruses consisted of clustered sequences (by cd-hit 4.8.1 with a threshold of 0.98)
from viruses of the realm *Riboviria* in the NCBI GenBank (the sequences were
downloaded on June 2, 2019) (46). Next, the query sequences with viral hits were

subjected to second BLASTx analyses, which were performed against the NCBI nr

database. Finally, the second blast hits with the best hit against deltaviruses were regarded

as deltavirus-like agents, and they were used for detailed analyses.

Confirmation of circularities of deltavirus contigs

Self dot-plot analyses of linear deltavirus contigs were conducted using the YASS online

web server (29). Based on the analysis, the contigs were manually circularized using

Geneious 11.1.5 (https://www.geneious.com). Further confirmation of the circularities of

deltavirus contigs was obtained by mapping short reads to circular deltavirus contigs

using Geneious software as follows. The reads used for the *de novo* assembly were first

imported to Geneious, after which they were mapped to the circular contigs using the

Geneious mapper. The mapped reads across the circularized boundaries were confirmed

manually.

Detection of possible RNA-editing sites at stop codons of DAg genes

We used the mapped reads obtained by the analyses described above to detect possible

RNA-editing at the stop codons of DAg genes of deltaviruses. We analyzed the

nucleotide variations (presence of variations, variant nucleotide(s), and variant

frequency) of mapped reads at each of the stop codons of newly identified deltaviruses

using the "Find Variation/SNPs" function in Geneious. We used a custom Python script to

visualize base quality scores of the NGS reads mapped at the second nucleotide of the

stop codon genome. The codes are available at following URL:

https://github.com/shohei-kojima/iwamoto_et_al_2020.

Sequence characterization

DAg ORFs were detected by the "Find ORFs" function in Geneious with a threshold of

500 nucleotides. Poly(A) signals were manually detected. Putative ribozyme sequences

were identified using nucleotide sequence alignment with other deltaviruses. Ribozyme

structures were first inferred using the TT2NE webserver (56), and the obtained data were

then visualized using the PsudoViewer3 web server (57). We used the visualized data as

guides to draw ribozyme structures.

The self-complementarities of deltavirus-like contigs were analyzed using the Mfold

web server (58). Coiled-coil domains and nuclear export signals (NLSs) were predicted

using DeepCoil (59) and NLS mapper

(http://nls-mapper.iab.keio.ac.jp/cgi-bin/NLS_Mapper_form.cgi) web servers,

respectively.

Short read mapping for detection of deltavirus infection

To detect deltavirus-derived reads in publicly available RNA-seq data, short reads were

mapped to deltavirus genomes and then the numbers of mapped reads were counted as

follows. SRA files were downloaded from NCBI, dumped, and preprocessed following

the procedure described above. The preprocessed reads were then mapped to linearized

deltavirus contigs by HISAT2 with the default setting. SAM tools were used to extract the

mapped BAM files from the resultant BAM files, and the mapped read numbers were

counted using BamTools 2.5.1 (60).

Recovery of a deltavirus genome from RNA-seq data of Pardaliparus venustulus

Mapped reads obtained from SRR7244693, SRR7244695, SRR7244696, SRR7244697,

and SRR7244698 in the above analysis (section *Short read mapping for detection of*

deltavirus infection) were extracted by Geneious. All the extracted reads were

co-assembled using Geneious Assembler with the circular contig assembly function. The

obtained circular contigs were characterized as described previously.

Animals and samples

Zebra finches (n = 30) and Bengalese finches (n = 5) were obtained from breeding

colonies at Wada lab, Hokkaido University. The founder birds were originally obtained

from local breeders in Japan. Five to twelve birds were kept together in cages in an aviary

and were exposed to a 13:11 light-dark cycle. Blood samples were collected from the

wing vein using 30 G × 8 mm syringe needles (Becton Dickinson; Franklin Lakes, NJ,

USA). Each blood sample was diluted 1.5 times with PBS, frozen immediately on dry ice

after collection, and maintained at −80°C until further requirement. These experiments

were conducted under the guidelines and with the approval of the Committee on Animal

Experiments of Hokkaido University. These guidelines are based on the national

regulations for animal welfare in Japan (Law for the Humane Treatment and

Management of Animals with partial amendment No.105, 2011).

Woodchucks (*Marmota monax*) were purchased from Northeastern Wildlife

(Harrison, ID, USA) and kept at the Laboratory Animal Center, National Taiwan

University College of Medicine. At three days of age, the animal supplier infected

captive-born woodchucks with WHV from the same infectious pool. Wild-caught

woodchucks were infected naturally and live trapped. Serum samples were collected

from the woodchucks periodically via the femoral vein by means of venipuncture. Liver

tissues of woodchucks were obtained at autopsy, snap-frozen in liquid nitrogen, and

stored at −80°C until RNA extraction. This study used liver tissues from 10 wild-caught

and 33 captive-born woodchucks and serum samples from 33 wild-caught and five

captive-born woodchucks. In this study, all the experimental procedures involving

woodchucks were performed under protocols approved by the Institutional Animal Care

and Use Committee of National Taiwan University College of Medicine.

Real-time and Endpoint RT-PCR detection of deltaviruses from animal specimens

Total RNAs were isolated from the whole blood samples from zebra finches and serum

samples from woodchucks using Quick RNA Viral Kit (Zymo Research; Irvine, CA,

USA). The obtained RNA samples were stored at −80°C until further requirement. Total

RNAs were also extracted from 50 mg of the woodchuck liver tissues using either Trizol

(Thermo Fisher Scientific; Waltham, MA, USA) or ToTALLY RNA kit (Thermo Fisher

Scientific; Waltham, MA, USA) according to the manufacturers' instructions.

The obtained RNA was reverse-transcribed into cDNA using ReverTra Ace qPCR

RT Master Mix (TOYOBO; Osaka, Japan), and these were used as templates for real-time

PCR analyses. Real-time PCR was performed with KOD SYBR qPCR Mix (TOYOBO)

and primers (Supp Table 7) using the CFX Connect Real-Time PCR Detection System

(Bio-Rad; Hercules, CA, USA) according to the manufacturer's instructions. The

real-time PCR systems for mmDeV and tgDeV were validated using

pcDNA3-mmDeV(-) and pcDNA3-tgDeV(-) monomer, respectively, as controls.

End-point RT-PCR was also performed to confirm deltavirus infections. PCR was

performed with Phusion Hot Start II DNA Polymerase (Thermo Fisher Scientific) using

the above-described cDNAs and primers listed in Supp Table 7. The PCR products were

analyzed by agarose gel electrophoresis. The obtained PCR products were purified and

sequenced by Sangar sequencing in FASMAC (Atsugi, Japan).

Determination of a full genome sequence of deltavirus in passerine birds

To determine the full genome sequence of detected deltaviruses, the cDNA obtained in

the section "*Realtime and Endpoint RT-PCR detection of deltaviruses from animal*

specimens" was amplified using illustra GenomiPhi V2 Kit (GE healthcare; Chicago, IL,

USA). The amplified DNA was then purified with innuPREP PCRpure Kit (Analytik

Jena: Jena, Germany). PCR was performed with Phusion Hot Start II DNA Polymerase

using the primers listed in Supp Table 7. The PCR products were analyzed using agarose

gel electrophoreses. When single bands were observed, the amplicon was purified with

innuPREP PCRpure Kit. When several bands were detected, bands of the expected sizes

were extracted and purified using Zymoclean Gel DNA Recovery Kit (Zymo Research).

The purified amplicons were sequenced in FASMAC (Atsugi, Japan).

Phylogenetic analysis

Deduced amino acid sequences of DAg proteins were used to infer the phylogenetic

relationship between deltaviruses. Multiple alignment was performed by MAFFT 7.427

using the E-INS-i algorithms (61), and ambiguously aligned regions were then removed

using trimAl 1.2rev59 with the --strict option (62). The phylogenetic relationship was

inferred by the maximum likelihood method using RAxML Next Generation v. 0.9.0 (63).

The LG+G model, which showed the lowest BIC by prottest3 3.4.2 (64), was used. The

reliability of the tree was assessed by 1,000 bootstrap resampling using the transfer

bootstrap expectation method (65). The alignment file is available in Supporting

materials.

Detection of co-infected viruses

To identify co-infected viruses in deltavirus–positive SRAs, a three-step BLASTx search

was performed. First, BLASTx searches were performed against a custom database,

including RefSeq protein sequences from viruses using the assembled contigs (see the

subsection *Detection of deltaviruses from publicly available transcriptome data*) as

queries. The custom database was prepared as follows. Virus-derived protein sequences

in the RefSeq protein database (46) were downloaded on July 17, 2020, and were

clustered by cd-hit 4.8.1 (threshold = 0.9). Then, sequences of more than 100 amino acid

residues were extracted using SeqKit 0.10.1 and these were used as a BLAST database.

The first BLAST hits were extracted, which were used for the second BLASTx analysis.

The second BLASTx analysis was performed against the NCBI RefSeq protein database.

The BLAST hits with the best hit to viral sequences were extracted and used for the final

BLASTx searches. The final BLASTx searches were performed against the NCBI nr

database. The BLAST hits with the best hit to viral sequences were extracted and

analyzed manually.

Cell culture

HepG2-NTCP cells were cultured with Dulbecco's modified Eagle's medium

(DMEM)/F-12 + GlutaMax (Thermo Fisher Scientific) supplemented with 10 mM

HEPES (Sigma Aldrich; St. Louis, MO, USA), 100 unit/ml penicillin (Meiji; Tokyo,

Japan), 100 mg/ml streptomycin (Meiji), 10% FBS (Sigma Aldrich), 5 μg/ml insulin

(Wako; Tokyo, Japan) and 400 g/ml G418 (Nacalai tesque). Huh7 and WCH-17 cells

were maintained in DMEM (Wako) containing 10% FBS (Sigma Aldrich), 100 unit/ml

penicillin (Meiji), 100 mg/ml streptomycin (Meiji), 100 mM nonessential amino acids

(Thermo Fisher Scientific), 1 mM sodium pyruvate (Sigma Aldrich), and 10 mM HEPES

(Sigma Aldrich).

Antibody production

The peptides corresponding to 65 to 78 aa (DSSSPRKRKRGEGG) of tgDeV DAg and

174 to 187 aa (ESPYSRRGEGLDIR) of mmDeV DAg conjugated with cysteine at N

terminus were synthesized. Each of the peptides was injected into mice, and antisera were

obtained from the mice at 42 days after the peptide injections. Each of the antisera was

affinity-purified using the corresponding peptide. The whole procedure was performed in

SCRUM (Tokyo, Japan).

Rescue of mmDeV and tgDeV

The DNA of negative-strand genomes of mmDeV and tgDeV was synthesized in

GenScript Japan (Tokyo, Japan). The synthesized DNAs were inserted into the KpnI

-XbaI site of the pcDNA3 vector, designated as pcDNA3-mmDeV(-) monomer and

pcDNA-tgDeV (-) monomer. In addition, tandem sequences of mmDeV and tgDeV

genome were inserted into the pcDNA3 vector, which were named pcDNA3-mmDeV(-)

dimer and pcDNA-tgDeV (-) dimer, respectively. To rescue these viruses,

pcDNA3-mmDeV(-) dimer or pcDNA-tgDeV (-) dimer was transfected into Huh7 and

WCH-17 cells using Lipofectamine 3000 and Lipofectamine 2000 (Thermo Fisher

Scientific), respectively, according to the manufacturer's instructions. The transfected

cells were cultured for 48 h and were used for western blotting, IFA to verify DAg protein

expression.

Western blotting

Cells were lysed with SDS sample buffer [100 mM Tris-HCl (pH 6.8) (Sigma Aldrich),

4% SDS (Nippon gene; Tokyo, Japan), 20% glycerol (Nacalai tesque), 10%

2-mercaptoethanol (Wako)]. The cell lysates were subjected to SDS-PAGE and

transferred onto polyvinylidene difluoride membranes (Merck Millipore; Darmstadt,

Germany). After blocking the membranes with 5% skim milk (Morinaga; Tokyo, Japan),

they were reacted with anti-tgDeV DAg, anti-mmDeV DAg, or anti-actin (Sigma

Aldrich) antibodies as primary antibodies, followed by reaction with horseradish

peroxidase (HRP)–conjugated secondary antibodies (Cell Signaling Technology;

Danvers, MA, USA).

Indirect immunofluorescence assay (IFA)

The cells were fixed in 4% paraformaldehyde (Wako) and then permeabilized using 0.3%

Triton X-100 (MP Biomedicals; Santa Ana, CA, USA). After blocking the cells by

incubation in PBS containing 1% bovine serum albumin (BSA) (KAC; Kyoto, Japan),

they were treated with the primary antibodies against HDAg, tgDeV DAg, or mmDeV

DAg and then incubated with Alexa555-conjugated secondary antibody (Thermo Fisher

Scientific), together with DAPI (Nacalai tesque). To detect deltavirus-positive cells, the

fluorescence signal was observed using fluorescence microscopy, BZ-X710 (KEYENCE;

Osaka, Japan). High magnification examination of the subcellular localization was

performed using confocal microscopy, LSM900 (ZEISS; Oberkochen, Germany).

Deltavirus preparation and infection assay

HDV was produced from the culture supernatants of Huh7 cells transfected with HDV

(pSVLD3) and HBs (pT7HB2.7) expressing plasmid, as described previously (32, 66).

tgDeV and mmDeV were also subjected to the same assay. The supernatants of

transfected cells were collected at 6, 9, and 12 days post-transfection, and they were then

filtrated and concentrated using 0.45-μm filters and Amicon Ultra (Merck Millipore),

according to the manufacturer's instructions. The concentrated supernatants were

inoculated into HepG2-NTCP cells with 5% PEG8000 (Sigma Aldrich) for 24 h followed

by washing to remove free viruses. The inoculated cells were cultured for 6 days and used

for the downstream analyses.

Acknowledgments

HDV and HBs expression plasmids were kindly provided by Dr. John Taylor (the Fox

Chase Cancer) and Dr. Camille Sureau (Institute National de la Transfusion Sanguine).

We thank Dr. Keiko Takemoto (Kyoto University) for her kind help with the setting of

computer resources. The super-computing resources were provided by Human Genome

Center, the Institute of Medical Science, the University of Tokyo, and the NIG

supercomputer at ROIS National Institute of Genetics. All the silhouette images except

for woodchuck were downloaded from silhouetteAC (http://www.silhouette-ac.com/).

This study was supported by the Hakubi project at Kyoto University (MH);

Grant-in-Aid for Scientific Research on Innovative Areas from the Ministry of Education,

Culture, Science, Sports, and Technology (MEXT) of Japan, Grant Numbers

JP19H04839 (SI), JP16H06429 (KT), JP16K21723 (KT), JP16H0643 (KT),

JP17H05821 (MH), and JP19H04833 (MH); the Japan Society for the Promotion of

Science KAKENHI, Grant Numbers JP19K16672 (MI), JP20J00868 (MI), and

JP20H03499 (KW); AMED Grant Numbers JP20jm0210068j0002 (KW) and

JP20fk0310114j0004 (KW).

Author contributions

MH conceived the study. MI, KW, and MH designed the study. MI conducted cell culture

experiments. MH, JK, SK performed *in silico* analyses. YS, YTL, HLW, KW, and MH

prepared and analyzed animal specimens. All the authors analyzed and discussed the data.

MI and MH wrote the manuscript.

Figure legends

Figure 1. Genome organization of novel deltaviruses.

Genomes of **(a)** tgDeV, mmDeV, and ovDeV (complete genomes) and **(b)** scDeV and

egDeV (partial genomes). Annotations (ORF, poly-A signal, and ribozymes) are shown

by colored arrow pentagons. The numbers indicate nucleotide positions. **(c)**

Self-complementarities of novel deltaviruses. The predicted RNA structures were

visualized using the Mfold web server (58). Red, blue, and green arcs indicate G-C, A-U,

and G-U pairs, respectively.

Figure 2. Amino acid sequence characterization of putative delta antigens of novel

deltaviruses.

(a) Alignment and functional features of the putative S-HDAg and DAgs of

representative HDVs and novel deltaviruses. (Putative) functional domains are shown by

colored boxes. Me: arginine methylation site, Ac: lysine acetylation site, P: Serine

phosphorylation site. **(b)** ovDeV mRNA (upper panel) and a possible A-to-I RNA-editing

site (lower panel). Consensus ovDeV-DAg mRNA sequence and mapped read sequence

with potential RNA-edited nucleotides (blue boxes). Pink boxes indicate the ORF of

ovDeV DAg. **(c)** Deduced amino acid sequences of ovDeV-DAg proteins translated from

the viral mRNA with or without RNA-editing. The blue letter shows the possible

RNA-editing site.

Figure 3. Mapping coverages of original short reads of each contig.

Mapped read graphs of **(a)** tgDeV, **(b)** mmDeV, and **(c)** ovDeV. Lines, arrow pentagons,

and arrowheads indicate viral genomes, ribozymes, and poly(A) signals, respectively.

710 The numbers above the graphs show nucleotide positions. The light pink box indicates a

711 low read depth region in the putative transcript of tgDeV.

713 **Figure 4. Interfamily transmission of deltaviruses among passerine birds.**

714 **(a–c)** RT-PCR detection of a deltavirus from *Lonchura striata*. **(a)** Plasmid used for the

715 establishment of real-time PCR detection system for tgDeV and **(b)** the tgDeV circular

716 genome. The blue arrows indicate the primers used for endo-point RT-PCR detection. **(c)**

717 Endo-point RT-PCR for detection of the circular deltavirus genome. M, 100-bp ladder

718 marker. **(d)** Pairwise nucleotide identities between deltaviruses detected in passerine

719 birds. **(e)** Phylogenetic tree of passerine birds positive for deltaviruses. Phylogenetic tree

720 of birds and deltavirus infections are indicated. MYA: million years ago.

722 **Figure 5. Phylogenetic analysis of deltaviruses.**

723 **(a)** Heat map of pairwise amino acid sequence identities between deltaviruses. **(b)** The

724 phylogenetic tree was inferred by the maximum likelihood method using an amino acid

725 sequence alignment of representative deltaviruses. Known phenotypes (RNA-editing and

726 expression of the large isoform of DAg protein) and helper virus(es) of each virus are

727 shown on the right. Note that the SDeV phenotypes are shown in gray letters, because

728 there is insufficient information, evidence, or both for the RNA-editing and L-DAg

729 expression. The deltaviruses identified in this study are indicated by the blue circles.

730 Bootstrap values >70 are shown. SDeV: snake deltavirus, RDeV: rodent deltavirus.

732 **Figure 6. Detection of tgDeV DAg and mmDeV DAg in cells ectopically expressing**

733 **the tgDeV or mmDeV dimer genome.**

(**a** and **b**) Western blotting analysis of Huh7 or WCH-17 cells transfected with a tgDeV or

mmDeV dimer-sequence expression plasmid. The numbers on the left side of panels

indicate the size marker of protein (kDa). (**c–f**) Indirect immunofluorescence analysis of

the expression of tgDeV or mmDeV DAg protein. The cells were observed using

fluorescent microscopy (**c** and **d**) or a confocal microscopy (**e** and **f**). Blue; DAPI, Red;

tgDeV or mmDeV DAg. Scale bars = 50 μm (c and d) and 5 μm (e and f).

Figure 7. No infectious particle of tgDeV and mmDeV was produced by

supplementation of HBV envelop proteins.

(**a**) Quantification of deltavirus RNAs in culture supernatants. HDV, tgDeV, or mmDeV

expression plasmid was transfected with or without plasmid expressing HBV envelope

proteins into Huh7 cells. Viral RNA levels in supernatants were quantified using

quantitative RT-PCR (n = 3). (**b** and **c**) HepG2-NTCP cells were incubated with the

culture supernatants of the transfectants for 24 h in the presence or absence of 500 nM

Myrcludex B (MyrB), an inhibitor of HBV envelope-dependent viral entry. The cells

were cultured for an additional 6 days, and viral RNA levels and protein exprssion were

analyzed using quantitative RT-PCR (n = 3) (**b**) and IFA (**c**), respectively. The numbers in

(**c**) correspond to those of (**b**). Blue, DAPI; Red, HDV; tgDeV, or mmDeV DAg. Scale

bar = 50 μm.

Reference

1.   Magnius L, Taylor J, Mason WS, Sureau C, Deny P, Norder H, Ictv Report C.

2018. ICTV Virus Taxonomy Profile: Deltavirus. J Gen Virol 99:1565-1566.

2.   Wong SK, Lazinski DW. 2002. Replicating hepatitis delta virus RNA is edited in

the nucleus by the small form of ADAR1. Proc Natl Acad Sci U S A

99:15118-23.

3.   Bergmann KF, Gerin JL. 1986. Antigens of hepatitis delta virus in the liver and

serum of humans and animals. J Infect Dis 154:702-6.

4.   Bonino F, Heermann KH, Rizzetto M, Gerlich WH. 1986. Hepatitis delta virus:

protein composition of delta antigen and its hepatitis B virus-derived envelope. J

Virol 58:945-50.

5.   Taylor JM. 2020. Infection by Hepatitis Delta Virus. Viruses 12.

6.   O'Malley B, Lazinski DW. 2005. Roles of carboxyl-terminal and farnesylated

residues in the functions of the large hepatitis delta antigen. J Virol 79:1142-53.

7.   Kuo MY, Sharmeen L, Dinter-Gottlieb G, Taylor J. 1988. Characterization of

self-cleaving RNA sequences on the genome and antigenome of human hepatitis

delta virus. J Virol 62:4439-44.

8.   Sharmeen L, Kuo MY, Dinter-Gottlieb G, Taylor J. 1988. Antigenomic RNA of

human hepatitis delta virus can undergo self-cleavage. J Virol 62:2674-9.

9.   Chen PJ, Kalpana G, Goldberg J, Mason W, Werner B, Gerin J, Taylor J. 1986.

Structure and replication of the genome of the hepatitis delta virus. Proc Natl

Acad Sci U S A 83:8774-8.

10. Kos A, Dijkema R, Arnberg AC, van der Meide PH, Schellekens H. 1986. The

hepatitis delta (delta) virus possesses a circular RNA. Nature 323:558-60.

11. Wang KS, Choo QL, Weiner AJ, Ou JH, Najarian RC, Thayer RM, Mullenbach

GT, Denniston KJ, Gerin JL, Houghton M. 1986. Structure, sequence and

expression of the hepatitis delta (delta) viral genome. Nature 323:508-14.

12. Rizzetto M. 2015. Hepatitis D Virus: Introduction and Epidemiology. Cold

Spring Harb Perspect Med 5:a021576.

13. Littlejohn M, Locarnini S, Yuen L. 2016. Origins and Evolution of Hepatitis B

Virus and Hepatitis D Virus. Cold Spring Harb Perspect Med 6:a021360.

14. Rizzetto M. 2016. The adventure of delta. Liver Int 36 Suppl 1:135-40.

15. Wille M, Netter HJ, Littlejohn M, Yuen L, Shi M, Eden JS, Klaassen M, Holmes

787   EC, Hurt AC. 2018. A Divergent Hepatitis D-Like Agent in Birds. Viruses 10.

788   16.   Chang WS, Pettersson JH, Le Lay C, Shi M, Lo N, Wille M, Eden JS, Holmes

789   EC. 2019. Novel hepatitis D-like agents in vertebrates and invertebrates. Virus

790   Evol 5:vez021.

791   17.   Hetzel U, Szirovicza L, Smura T, Prahauser B, Vapalahti O, Kipar A, Hepojoki J.

792   2019. Identification of a Novel Deltavirus in Boa Constrictors. mBio 10.

793   18.   Paraskevopoulou S, Pirzer F, Goldmann N, Schmid J, Corman VM, Gottula LT,

794   Schroeder S, Rasche A, Muth D, Drexler JF, Heni AC, Eibner GJ, Page RA,

795   Jones TC, Muller MA, Sommer S, Glebe D, Drosten C. 2020. Mammalian

796   deltavirus without hepadnavirus coinfection in the neotropical rodent

797   Proechimys semispinosus. Proc Natl Acad Sci U S A

798   doi:10.1073/pnas.2006750117.

799   19.   Szirovicza L, Hetzel U, Kipar A, Martinez-Sobrido L, Vapalahti O, Hepojoki J.

800   2020. Snake Deltavirus Utilizes Envelope Proteins of Different Viruses To

801   Generate Infectious Particles. mBio 11.

802   20.   Hedges SB, Dudley J, Kumar S. 2006. TimeTree: a public knowledge-base of

803   divergence times among organisms. Bioinformatics 22:2971-2.

804   21.   Chao YC, Tang HS, Hsu CT. 1994. Evolution rate of hepatitis delta virus RNA

805   isolated in Taiwan. J Med Virol 43:397-403.

806   22.   Krushkal J, Li WH. 1995. Substitution rates in hepatitis delta virus. J Mol Evol

807   41:721-6.

808   23.   Alvarado-Mora MV, Romano CM, Gomes-Gouvea MS, Gutierrez MF, Carrilho

809   FJ, Pinho JR. 2011. Dynamics of hepatitis D (delta) virus genotype 3 in the

810   Amazon region of South America. Infect Genet Evol 11:1462-8.

811   24.   Fletcher SP, Chin DJ, Ji Y, Iniguez AL, Taillon B, Swinney DC, Ravindran P,

812   Cheng DT, Bitter H, Lopatin U, Ma H, Klumpp K, Menne S. 2012.

813   Transcriptomic analysis of the woodchuck model of chronic hepatitis B.

814   Hepatology 56:820-30.

815   25.   Fletcher SP, Chin DJ, Gruenbaum L, Bitter H, Rasmussen E, Ravindran P,

816   Swinney DC, Birzele F, Schmucki R, Lorenz SH, Kopetzki E, Carter J, Triyatni

817   M, Thampi LM, Yang J, AlDeghaither D, Murreddu MG, Cote P, Menne S. 2015.

818   Intrahepatic Transcriptional Signature Associated with Response to

819   Interferon-alpha Treatment in the Woodchuck Model of Chronic Hepatitis B.

820  PLoS Pathog 11:e1005103.

821 26. Ryu WS, Bayer M, Taylor J. 1992. Assembly of hepatitis delta virus particles. J
822   Virol 66:2310-5.

823 27. Gudima S, He Y, Chai N, Bruss V, Urban S, Mason W, Taylor J. 2008. Primary
824   human hepatocytes are susceptible to infection by hepatitis delta virus
825   assembled with envelope proteins of woodchuck hepatitis virus. J Virol
826   82:7276-83.

827 28. Zuccola HJ, Rozzelle JE, Lemon SM, Erickson BW, Hogle JM. 1998. Structural
828   basis of the oligomerization of hepatitis delta antigen. Structure 6:821-830.

829 29. Noe L, Kucherov G. 2005. YASS: enhancing the sensitivity of DNA similarity
830   search. Nucleic Acids Res 33:W540-3.

831 30. Yan H, Zhong G, Xu G, He W, Jing Z, Gao Z, Huang Y, Qi Y, Peng B, Wang H,
832   Fu L, Song M, Chen P, Gao W, Ren B, Sun Y, Cai T, Feng X, Sui J, Li W. 2012.
833   Sodium taurocholate cotransporting polypeptide is a functional receptor for
834   human hepatitis B and D virus. Elife 1:e00049.

835 31. Iwamoto M, Watashi K, Tsukuda S, Aly HH, Fukasawa M, Fujimoto A, Suzuki
836   R, Aizaki H, Ito T, Koiwai O, Kusuhara H, Wakita T. 2014. Evaluation and
837   identification of hepatitis B virus entry inhibitors using HepG2 cells
838   overexpressing a membrane transporter NTCP. Biochem Biophys Res Commun
839   443:808-13.

840 32. Sureau C, Guerra B, Lee H. 1994. The middle hepatitis B virus envelope protein
841   is not necessary for infectivity of hepatitis delta virus. J Virol 68:4063-6.

842 33. Gripon P, Cannie I, Urban S. 2005. Efficient inhibition of hepatitis B virus
843   infection by acylated peptides derived from the large viral surface protein. J
844   Virol 79:1613-22.

845 34. Bergner LM, Orton RJ, Broos A, Tello C, Becker DJ, Carrera JE, Patel AH, Biek
846   R, Streicker DG. 2020. Diversification of mammalian deltaviruses by host
847   shifting. bioRxiv doi:10.1101/2020.06.17.156745.

848 35. Edgar RC, Taylor J, Altman T, Barbera P, Meleshko D, Lin V, Lohr D,
849   Novakovsky G, Al-Shayeb B, Banfield JF, Korobeynikov A, Chikhi R, Babaian
850   A. 2020. Petabase-scale sequence alignment catalyses viral discovery. bioRxiv
851   doi:10.1101/2020.08.07.241729.

852 36. Anonymous. !!! INVALID CITATION !!! (34, 35).

853   37.   Chang FL, Chen PJ, Tu SJ, Wang CJ, Chen DS. 1991. The large form of

854   hepatitis delta antigen is crucial for assembly of hepatitis delta virus. Proc Natl

855   Acad Sci U S A 88:8490-4.

856   38.   Perez-Vargas J, Amirache F, Boson B, Mialon C, Freitas N, Sureau C, Fusil F,

857   Cosset FL. 2019. Enveloped viruses distinct from HBV induce dissemination of

858   hepatitis D virus in vivo. Nat Commun 10:2098.

859   39.   Kircher M, Sawyer S, Meyer M. 2012. Double indexing overcomes inaccuracies

860   in multiplex sequencing on the Illumina platform. Nucleic Acids Res 40:e3.

861   40.   Nelson MC, Morrison HG, Benjamino J, Grim SL, Graf J. 2014. Analysis,

862   optimization and verification of Illumina-generated 16S rRNA gene amplicon

863   surveys. PLoS One 9:e94249.

864   41.   Renaud G, Stenzel U, Maricic T, Wiebe V, Kelso J. 2015. deML: robust

865   demultiplexing of Illumina sequences using a likelihood-based approach.

866   Bioinformatics 31:770-2.

867   42.   D'Amore R, Ijaz UZ, Schirmer M, Kenny JG, Gregory R, Darby AC, Shakya M,

868   Podar M, Quince C, Hall N. 2016. A comprehensive benchmarking study of

869   protocols and sequencing platforms for 16S rRNA community profiling. BMC

870   Genomics 17:55.

871   43.   Wright ES, Vetsigian KH. 2016. Quality filtering of Illumina index reads

872   mitigates sample cross-talk. BMC Genomics 17:876.

873   44.   Fuxjager MJ, Lee JH, Chan TM, Bahn JH, Chew JG, Xiao X, Schlinger BA.

874   2016. Research Resource: Hormones, Genes, and Athleticism: Effect of

875   Androgens on the Avian Muscular Transcriptome. Mol Endocrinol 30:254-71.

876   45.   Duffy DL, Bentley GE, Drazen DL, Ball GF. 2000. Effects of testosterone on

877   cell-mediated and humoral immunity in non-breeding adult European starlings.

878   Behavioral Ecology 11:654-662.

879   46.   Coordinators NR. 2018. Database resources of the National Center for

880   Biotechnology Information. Nucleic Acids Res 46:D8-D13.

881   47.   Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ

882   preprocessor. Bioinformatics 34:i884-i890.

883   48.   Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome

884   alignment and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol

885   37:907-915.

886   49.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis
887   G, Durbin R, Genome Project Data Processing S. 2009. The Sequence
888   Alignment/Map format and SAMtools. Bioinformatics 25:2078-9.

889   50.   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin
890   VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N,
891   Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly
892   algorithm and its applications to single-cell sequencing. J Comput Biol
893   19:455-77.

894   51.   Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. 2017. metaSPAdes: a new
895   versatile metagenomic assembler. Genome Res 27:824-834.

896   52.   Li W, Godzik A. 2006. Cd-hit: a fast program for clustering and comparing large
897   sets of protein or nucleotide sequences. Bioinformatics 22:1658-9.

898   53.   Fu L, Niu B, Zhu Z, Wu S, Li W. 2012. CD-HIT: accelerated for clustering the
899   next-generation sequencing data. Bioinformatics 28:3150-2.

900   54.   Shen W, Le S, Li Y, Hu F. 2016. SeqKit: A Cross-Platform and Ultrafast Toolkit
901   for FASTA/Q File Manipulation. PLoS One 11:e0163962.

902   55.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden
903   TL. 2009. BLAST+: architecture and applications. BMC Bioinformatics 10:421.

904   56.   Bon M, Orland H. 2011. TT2NE: a novel algorithm to predict RNA secondary
905   structures with pseudoknots. Nucleic Acids Res 39:e93.

906   57.   Byun Y, Han K. 2009. PseudoViewer3: generating planar drawings of
907   large-scale RNA structures with pseudoknots. Bioinformatics 25:1435-7.

908   58.   Zuker M. 2003. Mfold web server for nucleic acid folding and hybridization
909   prediction. Nucleic Acids Res 31:3406-15.

910   59.   Ludwiczak J, Winski A, Szczepaniak K, Alva V, Dunin-Horkawicz S. 2019.
911   DeepCoil-a fast and accurate prediction of coiled-coil domains in protein
912   sequences. Bioinformatics 35:2790-2795.

913   60.   Barnett DW, Garrison EK, Quinlan AR, Stromberg MP, Marth GT. 2011.
914   BamTools: a C++ API and toolkit for analyzing and managing BAM files.
915   Bioinformatics 27:1691-2.

916   61.   Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software
917   version 7: improvements in performance and usability. Mol Biol Evol
918   30:772-80.

919   62.   Capella-Gutierrez S, Silla-Martinez JM, Gabaldon T. 2009. trimAl: a tool for
920   automated alignment trimming in large-scale phylogenetic analyses.
921   Bioinformatics 25:1972-3.
922   63.   Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A. 2019. RAxML-NG: a
923   fast, scalable and user-friendly tool for maximum likelihood phylogenetic
924   inference. Bioinformatics 35:4453-4455.
925   64.   Darriba D, Taboada GL, Doallo R, Posada D. 2011. ProtTest 3: fast selection of
926   best-fit models of protein evolution. Bioinformatics 27:1164-5.
927   65.   Lemoine F, Domelevo Entfellner JB, Wilkinson E, Correia D, Davila Felipe M,
928   De Oliveira T, Gascuel O. 2018. Renewing Felsenstein's phylogenetic bootstrap
929   in the era of big data. Nature 556:452-456.
930   66.   Kuo MY, Chao M, Taylor J. 1989. Initiation of replication of the human hepatitis
931   delta virus genome from cloned DNA: role of delta antigen. J Virol 63:1945-50.
932

Table 1. Summary of novel deltaviruses

Virus name		Host species	Tissue	SRA accession	DDBJ accession	Contig length (nt)	GC content (%)	BLASTx best hit		
								Virus name	Accession	Identity (%)
Taeniopygia guttata DeV	tgDeV	*Taeniopygia guttata*	Scapulohumeralis caudalis	SRR2545946	BR001665	1706	56.6	Rodent deltavirus	QJD13558	63.3
Marmota monax DeV	mmDeV	*Marmota monax*	Liver	SRR2136906	BR001661	1712	53.4	Hepatitis delta virus	AIR77039	60.0
Odocoileus virginianus DeV	ovDeV	*Odocoileus virginianus*	Pedicle	SRR4256033	BR001662	1690	56.4	Hepatitis delta virus	AHB60712	66.7
Erythrura gouldiae DeV	egDeV	*Erythrura gouldiae*	Skin	SRR7504989	BR001660	596	59.4[a]	Rodent deltavirus	QJD13555	63.5
Serinus canaria-associated DeV	scDeV	*Serinus canaria*	Skin	SRR2915371	BR001664	761	54.4[a]	Hepatitis delta virus	AIR77012	36.0
Pardaliparus venustulus DeV	pvDeV	*Pardaliparus venustulus*	Lung, Kidney, Cardiac muscle, Flight muscle, Liver	SRR7244693 SRR7244695 SRR7244696 SRR7244697 SRR7244698	BR001663	1708	55.8	Rodent deltavirus	QJD13562	62.4
Lonchura striata DeV	lsDeV	*Lonchura striata* var. *domestica*	Blood	-	LC575944	1708	56.2	Rodent deltavirus	QJD13555	62.9

a) GC content of the partial genome sequences.

Table 2. Detection of deltavirus-derived reads in RNA-seq data.

Virus	BioProject/ BioStudy	SRA	Host Taxonomy Family	Host Taxonomy Species	RPM [a] (read per million)	Tissue
tgDeV	PRJNA297576	SRR2545943	Estrildidae	*Taeniopygia guttata*	10.28	Pectoralis
		SRR2545944			1.02	Scapulohumeralis caudalis
		SRR2545946			56.73	Scapulohumeralis caudalis
	PRJNA558524	SRR9899549	Emberizidae [b]	*Emberiza melanocephala*	3.11	Blood
	PRJNA470787	SRR7244693	Paridae	*Pardaliparus venustulus*	10.68	Lung
		SRR7244695			2.07	Kidney
		SRR7244696			2.65	Cardiac muscle
		SRR7244697			7.12	Flight muscle
		SRR7244698			1.77	Liver
	PRJNA478907	SRR7504989	Estrildidae	*Erythrura gouldiae*	1.07	Skin
mmDeV	PRJNA291589	SRR2136906		*Marmota monax*	70.86	Liver
		SRR2136907			63.08	Liver
		SRR2136916			1.02	Liver
	SRP011132	SRR437934			46.34	PBMC
		SRR437938			19.83	PBMC
ovDeV	PRJNA317745	SRR4256033		Odocoileus virginianus	180.73	Pedicle
scDeV	PRJNA300534	SRR2915371		*Serinus canaria*	9.79	Skin

The full version of the table is available as Supplementary Table 3.

a) This table only shows the samples with RPM >1.

b) Emberizidae is regarded as the subfamily Emberizinae of the family Fringillidae in TimeTree.

Table 3. Coexisting viruses in deltavirus-positive SRAs

| SRA accession | Host | | Virus name | Envelope | Deltavirus infection |
	Species	Common name			
SRR2545944	*Taeniopygia guttata*	Zebra finch	Serinus canaria polyomavirus	-	tgDeV
SRR5001849	*Taeniopygia guttata*	Zebra finch	Serinus canaria polyomavirus	-	tgDeV
SRR5001850	*Taeniopygia guttata*	Zebra finch	Serinus canaria polyomavirus	-	tgDeV
SRR5001851	*Taeniopygia guttata*	Zebra finch	Serinus canaria polyomavirus	-	tgDeV
SRR2915371	*Serinus canaria*	Common canary	Canary bornavirus 3	+	scDeV
			Canary circovirus	-	
SRR7504989	*Erythrura gouldiae*	Gouldian finch	Erythrura gouldiae polyomavirus 1	-	egDeV

Fig. 1

a

b

c

Fig. 2

a

Conserved post-translational modification sites	(Predicted) NLS	Me Methylation site	P Phosphorylation site
Conserved leucine residues	(Predicted) coiled-coil domain	Ac Acetylation site	

b

c

No editing	...CAG UUU CCA UAG GAC UGA UUU UUG...
	Q F P *
A-to-I editing	...CAG UUU CCA UGG GAC UGA UUU UUG...
	Q F P W D *

Fig. 3

a tgDeV

b mmDeV

c ovDeV

d scDeV

■ Delta antigen ORF	■ Putative genomic ribozyme
▷ Poly(A) signal	■ Putative antigenomic ribozyme

Fig. 4

a

pcDNA3-tgDeV(-)

| CMV pro | tgDeV genome |

b

185 bp

tgDeV
genome

c

d

Nucleotide identity (%)

	pvDeV	tgDeV	lsDeV
pvDeV		98.2	98.4
tgDeV	98.2		98.2
lsDeV	98.4	98.2	

e

Family	Species	Deltavirus reads	
Paridae	*Pardaliparus venustulus*	+	pvDeV
Emberizidae	*Emberiza melanocephala*	+	
Estrildidae	*Taeniopygia guttata*	+	tgDeV
	Lonchura striata	+	lsDeV
	Erythrura gouldiae	+	egDeV

44 MYA

Fig. 5

a

	M55042 HDV1	AY261459 HDV2	L22063 HDV3	AF209859 HDV4	AM183328 HDV5	AM183332 HDV6	AM183333 HDV7	AM183330 HDV8	ovDeV	mmDeV	MK598012 RDeV	tgDeV	MH988742 SDeV	MH824555 duck DeV	scDeV	MN031240 fish DeV	MK962760 toad DeV	MN031239 newt DeV	MK962759 termite DeV
M55042 HDV1		75.0	67.9	74.0	73.0	70.9	72.6	74.0	63.3	57.7	55.1	51.3	48.5	40.2	35.0	22.4	21.5	23.6	21.3
AY261459 HDV2	75.0		66.2	74.9	76.5	73.3	72.4	76.9	62.6	56.4	54.8	51.5	46.2	35.6	37.1	19.0	24.6	25.6	20.4
L22063 HDV3	67.9	66.2		64.9	63.1	62.1	64.2	63.1	62.4	60.3	55.6	54.3	49.7	36.7	35.1	19.0	20.6	20.0	20.4
AF209859 HDV4	74.0	74.9	64.9		72.4	73.3	74.0	76.9	67.0	60.8	56.1	52.8	47.2	39.4	33.2	19.0	23.6	23.3	20.4
AM183328 HDV5	73.0	76.5	63.1	72.4		74.4	74.5	79.5	62.6	53.4	52.3	50.0	45.3	36.8	38.1	20.0	23.7	24.7	22.4
AM183332 HDV6	70.9	73.3	62.1	73.3	74.4		71.9	77.4	66.7	54.4	54.8	52.0	48.7	37.2	37.1	20.6	22.6	23.3	21.5
AM183333 HDV7	72.6	72.4	64.2	74.0	74.5	71.9		79.1	62.7	54.5	53.4	49.6	46.4	37.9	37.7	19.1	24.2	23.4	22.0
AM183330 HDV8	74.0	76.9	63.1	76.9	79.5	77.4	79.1		65.6	57.4	54.3	52.0	46.2	37.2	39.5	19.5	22.6	27.4	20.4
ovDeV	63.3	62.6	62.4	67.0	62.6	66.7	62.7	65.6		67.0	57.7	58.9	51.3	37.2	36.6	24.5	23.1	22.3	21.9
mmDeV	57.7	56.4	60.3	60.8	53.4	54.4	54.5	57.4	67.0		54.6	55.8	49.7	39.9	32.7	22.1	23.7	22.8	21.5
MK598012 RDeV	55.1	54.8	55.6	56.1	52.3	54.8	53.4	54.3	57.7	54.6		64.1	54.7	36.0	33.0	19.9	19.4	20.0	21.8
tgDeV	51.3	51.5	54.3	52.8	50.0	52.0	49.6	52.0	58.9	55.8	64.1		54.2	37.2	33.8	20.2	23.3	21.8	19.9
MH988742 SDeV	48.5	46.2	49.7	47.2	45.3	48.7	46.4	46.2	51.3	49.7	54.7	54.2		35.9	35.1	17.2	21.7	20.8	19.6
MH824555 duck DeV	40.2	35.6	36.7	39.4	36.8	37.2	37.9	37.2	37.2	39.9	36.0	37.2	35.9		29.5	15.3	16.8	22.3	16.2
scDeV	35.0	37.1	35.1	33.2	38.1	37.1	37.7	39.5	36.6	32.7	33.0	33.8	35.1	29.5		17.3	21.8	22.3	17.9
MN031240 fish DeV	22.4	19.0	19.0	19.0	20.0	20.6	19.1	19.5	24.5	22.1	19.9	20.2	17.2	15.3	17.3		15.0	12.5	14.0
MK962760 toad DeV	21.5	24.6	20.6	23.6	23.7	22.6	24.2	22.6	23.1	23.7	19.4	23.3	21.7	16.8	21.8	15.0		15.4	14.0
MN031239 newt DeV	23.6	25.6	20.0	23.3	24.7	23.3	23.4	27.4	22.3	22.8	20.0	21.8	20.8	22.3	22.3	12.5	15.4		15.0
MK962759 termite DeV	21.3	20.4	20.4	20.4	22.4	21.5	22.0	20.4	21.9	21.5	21.8	19.9	19.6	16.2	17.9	14.0	14.0	15.0	

Identity (%)
High
Low

b

	RNA-editing/ L-DAg	Candidate helper virus	HBs usage
● BR001665 tgDeV	-	?	-
MH988742 SDeV	??/ +22 aa??	Reptarenavirus Hartmanivirus	-
MK598012 RDeV	-	?	?
● BR001661 mmDeV	-	?	-
● BR001662 ovDeV	?/ +2 aa?	?	?
L22063 HDV3			
M55042 HDV1			
AF209859 HDV4			
AY261459 HDV2	+/ +19 aa	HBV	+
AM183332 HDV6			
AM183328 HDV5			
AM183330 HDV8			
AM183333 HDV7			

Delta-hepadna relationship?

0.05

Fig. 6

Fig. 7

a

Delta virus production in the medium

Delta virus RNA (copies/µl)

Delta virus :	−	+	+	−	+	+	−	+	+
HBV envelope :	−	−	+	−	−	+	−	−	+
		HDV			tgDeV			mmDeV	

b

Delta virus RNA in the cells

Delta virus RNA (copies/cell)

lane :	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
Delta virus :	−	−	+	+	+	+	−	−	+	+	+	+	−	−	+	+	+	+
HBV envelope :	−	−	−	−	+	+	−	−	−	−	+	+	−	−	−	−	+	+
MyrB :	−	+	−	+	−	+	−	+	−	+	−	+	−	+	−	+	−	+
			HDV						tgDeV						mmDeV			

c

Blue: nucleus   Red: DAg