Title

Activation of the Cpx-envelope stress response system promotes tolerance to antibacterials delivered by arginine-rich peptides and aminoglycosides in *Escherichia coli*

Authors

Jakob Frimodt-Møller[1*], Andreas Koulouktsis[1], Godefroid Charbon[1], Marit Otterlei[3], Peter E. Nielsen[2], Anders Løbner-Olesen[1*]

Affiliations

[1] Department of Biology, Center for Peptide-Based Antibiotics, University of Copenhagen, Copenhagen, Denmark. [2] Department of Cellular and Molecular Medicine, Center for Peptide-Based Antibiotics, The Panum Institute, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark. [3] Department of Clinical and Molecular Medicine, Faculty of Medicine and Health Sciences, NTNU, Norwegian University of Science and Technology, Trondheim, Norway.

* Corresponding authors: Jakob Frimodt-Møller (jakob.frimodtmoller@bio.ku.dk) and Anders Løbner-Olesen (lobner@bio.ku.dk).

Abstract

Cell penetrating peptides (CPP) are increasingly used for cellular drug delivery in both pro- and eukaryotic cells, and oligoarginines have attracted special attention. However; their mechanism of action, particularly for prokaryotes is still unknown. Arginine-rich CPPs (R-CPP) efficiently delivers the antimicrobial peptide nucleic acid (PNA) into bacteria. Here, we show that resistance to an R-CPP PNA conjugate in *Escherichia coli* requires multiple genetic modifications and is specific to R-CPP and not to the PNA-part. An integral part of the resistance was the constitutively activated Cpx-envelope stress response system (*cpx**), which decreased the cytoplasmic membrane potential and thereby indicates an indirectly energy dependent uptake mechanism. Interestingly, *cpx** mutants also showed increased tolerance to aminoglycosides and R-CPP conjugated to a peptide targeting the DNA sliding clamp; i.e., similar uptake in *E. coli* for these antimicrobial compounds. We speculate that the *cpx** phenotype could create an evolutionary opportunity to adapt and evolve in the presence of either compounds.

Author summary

The emergence of multidrug resistant bacteria is raising the need for new classes of antibiotics. Peptide nucleic acids (PNAs) may fill this requirement by their ability to block translation of essential mRNAs and hence inhibit growth. PNA needs conjugation to a delivery peptide (cell penetrating peptide; CPP) to enter the bacteria. Arginine-rich CPPs (CPP$_R$) are receiving a lot of attention for use as delivery vessels. Here, we show, for the first time, CPP$_R$-PNA resistance in *Escherichia coli* directed towards the delivery peptide. Consequently, resistance also applies to other antimicrobial compounds delivered by the same carrier. An

integral part of CPP$_R$ resistance is due to a constitutive active Cpx-response system, which leads to a

decreased electric potential ($\Delta\Psi$) across the inner membrane. The decreased $\Delta\Psi$ is a result of down-

regulation of two aerobic respiratory operons, namely NADH:ubiquinone oxidoreductase complex I and

cytochrome bo$_3$ ubiquinol oxidase. The decreased $\Delta\Psi$ also led to increased tolerance to aminoglycosides.

This shows that a (large) negative $\Delta\Psi$ is important for providing sufficient free energy for membrane

translocation of both CPP$_R$ and that the inner membrane is the main barrier for entry of both arginine-rich

delivery peptides and aminoglycosides.

48
Introduction
Antimicrobial resistance is one of the major challenges of the 21st century. Thus, efforts to bring

novel antimicrobial compounds, including antimicrobial peptides (AMP), into clinical use are

accelerating. Antisense technology, as a gene targeted precision drug modality has recently

produced several new drugs in clinical use (Inotersen, Valonesorsen and Golodirsen), and agents

based on the pseudopeptide DNA mimic, peptide nucleic acid (PNA)(1) and phosphorodiamidate

morpholino oligonucleotide (PMO)(2) have been proposed as a future antibiotic. Naked PNA and

PMO, like many AMPs, are unable to traverse the cell envelope in *Escherichia coli*(2, 3). Hence,

these requires a carrier molecule (i.e., cell-penetrating peptide (CPP)) for efficient translocation

into the cytoplasm. Very little is known about how CPP conjugated to PNA/PMO/AMP enters the

cell and for CPP-PNA only one paper has addressed this topic(4). Here, PNA is designed to target

the translation initiation region of the mRNA of an essential gene causing steric hindrance of

ribosome binding, and thereby inhibition of gene expression and cell death(1). PNA conjugated to

the lysine-rich CPP *L*((KFF)$_3$K) (CPP$_{KFF}$-PNA) enters *E. coli* by the non-essential, inner membrane

transporter SbmA(4); consequently SbmA deficient *E. coli* are highly resistant to this peptide-PNA.

However, PNA conjugated to an arginine-rich CPP, (R-Ahx-R)$_4$-Ahx-(βAla)-PNA (CPP$_{RXR}$-PNA) enters

the cell in an unknown and SbmA-independent manner(4). Cationic arginine-rich CPPs (R-CPP) are

increasingly used as molecular carriers and several CPP-based drugs are currently being pursued

in drug discovery in relation to for example wound healing, cancer, and Duchenne Muscular

Dystrophy(5). How R-CPP enters eukaryotic and prokaryotic cells is poorly understood. In

eukaryotes, evidence supports predominantly an endocytotic (energy-dependent) pathway,

although direct translocation (energy-independent mechanism) may also play a role (5), while it

is unclear how R-CPPs penetrate through membrane lipids in prokaryotes.

The envelope of Gram-negative bacteria consists of three essential structurally and chemically diverse layers: 1) the inner membrane (cytoplasmic membrane); 2) the periplasm containing a thin peptidoglycan layer; and 3) the outer membrane that contains phospholipids in the inner leaflet and lipopolysaccharide (LPS) in the outer leaflet. *E. coli* can actively transport compounds across the cytoplasmic membrane using either ATP directly via membrane transporter proteins or indirectly via the membrane potential (partly created by the chemical proton gradient, and also being part of theproton motive force (PMF), generated by the electron transport chain). The energetics of the cell varies considerably between growth with or without oxygen, with oxidative growth conditions providing the highest energy turnover. For instance, PMF is decreased during anoxic growth conditions. The electron transport chain will under aerobic conditions predominantly contain NADH:ubiquinone oxidoreductase complex I (organized from the *nuoABCEFGHIJKLMN*-operon; NDH-I) and cytochrome bo_3 ubiquinol oxidase (organized from the *cyoABCDE* operon; cytochrome bo_3 oxidase)(6). PMF generated by translocating protons from the cytoplasm into the periplasmic space(6), serves as energy storage, which is used to drive ex. ATP synthase, efflux pumps, and transport of metabolites(7). ATP is generated from ADP and inorganic phosphate by the multi-subunit protein complex F_1F_0-ATPase(8). F_1F_0-ATPase consist of two structural domains; F_1 containing the catalytic core and F_0 containing the membrane proton channel(8). The PMF across the cytoplasmic membrane is a combination of the electric potential ($\Delta\Psi$) and the transmembrane proton gradient (ΔpH: internal pH − external pH). $\Delta\Psi$ is always negative inside growing neutrophils, such as *E. coli*, and $\Delta\Psi$ decreases as ΔpH increases(9); i.e. bacteria may regulate $\Delta\Psi$ and/or ΔpH in order to control PMF.

The Bae, Cpx, Psp, Rcs, and σ^E pathways constitute the *E. coli* signaling systems that detect and respond to alterations of the bacterial envelope(10). These pathways are involved in the biogenesis, maintenance, and repair of the bacterial envelope, and thus contribute to cell surface integrity(10). One of the most well studied extracytoplasmic stress response-systems is the CpxA/R two-component signal transduction system, which among other genes down-regulate respiratory operons when activated(11). Like many other histidine kinases, CpxA is localized to the cytoplasmic membrane through two transmembrane helices and contains both periplasmic and cytoplasmic domains(12). The cytoplasmic response regulator, *cpxR*, is predicted to encode an OmpR-like transcriptional activator(12). CpxR consist of an N-terminal receiver domain (phospho-acceptor domain) and C-terminal DNA-binding domain(13). Under non-stress conditions the

auxiliary regulator CpxP inhibits the Cpx-response by a predicted direct interaction with the

periplasmic domain of CpxA(14). Under Cpx-inducing conditions, CpxP is degraded by the

periplasmic protease and chaperone DegP(15). The CpxP relieved CpxA autophosphorylates on a

conserved histidine residue (His-248), using ATP as its phosphoryl donor(16). Subsequently,

phosphorylated CpxA donates its phosphoryl group to the conserved aspartate residue (Asp-51)

in the N-terminal receiver domain of CpxR. Phosphorylated CpxR binds to DNA and act as a

transcriptional regulator(16). In addition, CpxA dephosphorylates CpxR-P, which ensures that

CpxR remains inactive in the absence of an activating signal(17). The outer membrane lipoprotein

NlpE that sense surface adhesion, acts as a 'sentry protein' against envelope stress and activates

the Cpx-response system through an undefined interaction with CpxA(17).

Here we show that *E. coli* can develop tolerance to antisense PNA conjugated to R-CPPs through

a constitutive Cpx-response that result in a decreased $\Delta\Psi$ across the cytoplasmic membrane.

The decreased $\Delta\Psi$ also confers cross-resistance to aminoglycosides indicating a similar uptake

mechanism for the two classes of compounds.

Results

To understand bacterial uptake and resistance to PNA R-CPP conjugates, we studied the widely used CPP$_{RXR}$(4) (Table 1). Unless stated otherwise the PNA part always targets translation of the *acpP* mRNA, encoding the acyl carrier protein, which is essential for fatty acid synthesis(4). The CPP$_{RXR}$-PNA exhibited a minimum inhibitory concentration (MIC) of 0.5 μM towards *E. coli*, whereas the corresponding CPP$_{RXR}$-PNA containing two mismatches in the PNA sequence (PNA$_{mm}$) did not inhibit bacterial growth up to 16 μM (Table 1); *i.e.*, the CPP$_{RXR}$ part alone did not confer any antimicrobial effect supporting an antisense mechanism of action. As a previous study failed to identify single-gene deletion mutants of *E. coli* resistant to CPP$_{RXR}$-PNA(4), we used adaptive laboratory evolution (ALE; see Methods) to generate CPP$_{RXR}$-PNA resistant mutants. Wild-type cells were grown in successively increasing concentrations of CPP$_{RXR}$-PNA, to select for mutation(s) that accumulate during long-term exposure to CPP$_{RXR}$-PNA. The ALE experiments were done in a biological duplicate (two independent linages) and was terminated when resistant clones appeared after 20 successive passages, corresponding to a 16-fold increase in MIC (from 0.5 μM to 8 μM) for CPP$_{RXR}$-PNA (Table 1).

Five individual clones from each experiment at day 20 (ten clones in total) were isolated and sequenced (Table 2; Table S1; Table S2). From the ten CPP$_{RXR}$-PNA resistant clones we identified three different genotypes, which were annotated as such: Ahx^R-1 to 3. Ahx^R-1 and 2 were isolated from lineage one and Ahx^R-3 from lineage two. None of the mutations identified in the CPP$_{RXR}$-PNA resistant clones resulted from adaptation to the growth medium, as none of them occurred in wild-type cells adapted to the same growth medium without CPP$_{RXR}$-PNA (Table S3).

Linage two from the ALE-experiment appeared to be clonal. This lineage contained genes/operons that were mutated in all of Ahx^R-1 to 3; the *gltF-yhcA* intergenic region, *waaB, waaO,* and *rpsL*, as well as a mutation in *cpxR* specific to only Ahx^R-3 (Table 2). *gltF* belongs to the *gltBDF*-operon, where *gltB* and *gltD* encode the large and small subunit of glutamate synthase(18), respectively. The function of GltF and the hypothetical protein YhcA, is unknown. The *waa*-operon contains the major core-oligosaccharide assembly operons in *E. coli(19)*. RpsL is the only mutated gene that is essential. The product of *rpsL* is the S12 protein; a component of the 30S subunit of the ribosome where it plays a role in translational accuracy(20). Lastly, *cpxAR* encodes the two-component regulatory system CpxA/R(21) (see above).

Resistance to CPP$_{RXR}$-PNA is towards the arginine-rich CPP-part and not to PNA-part

The MIC were determined for Ahx^R-1-3 (collectively CPP$_{RXR}$-PNA^R) towards additional peptide-PNA conjugates (Table 1). R-PNA^R strains were also resistant to the analogous CPP$_{RXR}$ targeting the essential division protein *ftsZ* instead of *acpP*: CPP$_{RXR}$-PNA$_{ftsZ}$ (Table 1). On the contrary, all R-PNA^R strains remained sensitive to an *acpP* targeting PNA conjugated to the lysine-phenylalanine CPP, (KFF)$_3$K (CPP$_{KFF}$-PNA) (Table 1). Together this shows that resistance to CPP$_{RXR}$-PNA is not directed towards the PNA-part but towards the (arginine-rich) delivery peptide, supporting the notion of different uptake mechanism for conjugates with these two peptides.

An activated Cpx-response leads to R-CPP tolerance

The Cpx-system is known to be involved in extracytoplasmic stress response and was mutated in five out of ten sequenced CPP$_{RXR}$-PNA resistance clones. Thus, we proceed to investigate the role of the Cpx-response in tolerance to R-CPP. The highest tolerance level to CPP$_{RXR}$-PNA was 8 µM (i.e. CPP$_{RXR}$-PNA^R level), which we define as CPP$_{RXR}$-PNA resistance. Therefore, everything between wild-type CPP$_{RXR}$-PNA sensitivity (>0.5 µM) and CPP$_{RXR}$-PNA resistance is referred to as CPP$_{RXR}$-PNA tolerance.

Loss of the Cpx-response, by deletion of *cpxA* or *cpxR* in otherwise wild-type cells had no effect on the tolerance to CPP$_{RXR}$-PNA (Fig. 1A). Interestingly, increased levels of CpxR led to a small but reproducible increase in tolerance to CPP$_{RXR}$-PNA (Fig. 1A), which could indicate that Cpx activation promotes tolerance to CPP$_{RXR}$-PNA. We proceeded to introduce the *cpxR$_{L20Q}$* mutation identified in the ALE experiment (Table 2) into wild-type cells and included a *cpxA$_{Δ16-17}$* mutant into this study from our strain collection (isolated in a different project). In CpxR$_{L20Q}$, leucine is replaced by glutamine at position 20, which is part of a highly conserved hydrophobic core in the N-terminal receiver domain shared by OmpR-family proteins[13]. In CpxA$_{Δ16-17}$ threonine at position 16 and leucine at position 17 of the first CpxA transmembrane domain are missing. Both mutations increased the MIC for CPP$_{RXR}$-PNA from 0.5 µM to 4 µM, relative to wild-type cells (Fig. 1A). Thus, although more tolerant to CPP$_{RXR}$-PNA than wild-type cells, the *cpxR$_{L20Q}$* and *cpxA$_{Δ16-17}$* mutants did not reach the CPP$_{RXR}$-PNA resistance level of the evolved mutant Ahx^R-3. Next, the Cpx-response was tested against an oligoarginine conjugate to a peptide targeting the DNA sliding clamp (DnaB; β-clamp) (RWLVK-GILQWRKI-RRRRRRRRRRR ; CPP$_{R11}$-APIM), which has been shown

to inhibit bacterial growth at micromolar concentrations both *in vitro* and *in vivo*(22) (Table 1) .

Loss of the Cpx-response by deletion of *cpxA* in an otherwise wild-type cell had no effect on

tolerance to CPP_{R11}-APIM, however tolerance was reproductively increased from 1 µM to 2 µM in

the *cpxR$_{L20Q}$* and *cpxA$_{Δ16-17}$* mutants, relative to wild-type cells (Fig. 1B). Thus this peptide also

exhibits tolerance to the Cpx-mutants but only at a 2 fold increase compared to 8 fold for the

CPP_{RXR}-PNA.

We used *cpxP* transcription as a readout for activation of the *cpx* response(23). *cpxR$_{L20Q}$*, *cpxA$_{Δ16-}$*

$_{17}$, and Ahx^R-3 all had increased expression of *cpxP* compared to wild-type (Fig. 1C), albeit not to

the level achieved by overproduction of NlpE (full activation of the Cpx-response; Fig. 1C). This

confirmed that the Cpx-response was indeed activated by the *cpxR$_{L20Q}$* and *cpxA$_{Δ16-17}$* mutations

and that these provided CPP_{RXR}-PNA (and CPP_{R11}-APIM tolerance) to the same level as full

activation of the Cpx-response by NlpE overproduction (Fig. 1B, D). Both mutations are therefore

gain-of-function mutations leading to a constitutively active Cpx-response, which previously have

been denoted *cpx**(16, 17).

Both *cpxR$_{L20Q}$* and *cpxA$_{Δ16-17}$* mutations are dependent on a functional Cpx-system for conferring

CPP_{RXR}-PNA tolerance

To assess whether the *cpxA$_{Δ16-17}$* and *cpxR$_{L20Q}$* mutations were dependent on a functional Cpx

system for conferring CPP_{RXR}-PNA tolerance, *cpxA* was deleted in the *cpxR$_{L20Q}$* strain and *cpxR* was

deleted in the *cpxA$_{Δ16-17}$* strain. In either case, CPP_{RXR}-PNA tolerance was reduced to wild-type

level (Fig. 1A). Thus, the substitution to glutamine in position 20 in CpxR does not make it

constitutively active by mimicking phosphorylation of the conserved Asp-51, and CpxR$_{L20Q}$ still

relies on a functional CpxA to be activated. Overproduction of wild-type CpxR or CpxA in the

cpxR$_{L20Q}$ or *cpxA$_{Δ16-17}$* mutants, respectively, also restored CPP_{RXR}-PNA susceptibility to wild-type

level (Fig. 1A). Together, this shows that the specific *cpxR$_{L20Q}$* or *cpxA$_{Δ16-17}$* mutations mediates

CPP_{RXR}-PNA tolerance, but do need a functional Cpx-system.

The Cpx-response is the only extracytoplasmic stress response conferring tolerance to CPP_{RXR}-

PNA

The Cpx-response is only one of five extracytoplasmic stress response systems known to maintain

cell envelope integrity during stress. While the Cpx-response can be activated by overproducing

nlpE, the four other extracytoplasmic stress responses can be individually activated by

overproduction of *baeR* for the Bae-pathway, *rcsB* for the Rcs-pathway, *pspF* for the Psp-pathway,

and *rpoE* for the σ^E-pathway(10). Activation of neither the Bae, Psp, Rcs, nor the σ^E-responses led

to a CPP$_{RXR}$-PNA tolerant phenotype (Fig. 1D), showing CPP$_{RXR}$-PNA tolerance is specific to an

activated Cpx-response. However, wild-type treated with sub-MIC concentrations (0.5 X MIC) of

CPP$_{RXR}$-PNA or CPP$_{RXR}$-PNA$_{mm}$ did not result in an increased *cpxP* expression (Fig. 1D). Hence,

although an activated Cpx-response confers CPP$_{RXR}$-PNA tolerance, low-level CPP$_{RXR}$-PNA

treatment in itself does not trigger extracytoplasmic stress.

219   Downregulation of respiratory operons leads to CPP$_{RXR}$-PNA tolerance

Because Cpx mediates downregulation of aerobic respiratory operons(11), expression of the

nuoABCEFGHIJKLMN and *cyoABCDE* operons were determined by RT-qPCR. As expected,

transcription of *nuoA* (to evaluate expression of the *nuoABCEFGHIJKLMN* operon) and *cyoA* (to

evaluate expression of the *cyoABCDE* operon) were downregulated in wild-type/pCA24n:*nlpE*,

cpxR$_{L20Q}$, *cpxA$_{\Delta16-17}$*, and Ahx^R-3 cells (Fig. 2A). We found that such Cpx activated cells in addition

to being CPP$_{R11}$-APIM (*cpxR$_{L20Q}$* and *cpxA$_{\Delta16-}$*17; Fig. 2B) and CPP$_{RXR}$-PNA tolerant (Fig. 2B) were also

more tolerant to the aminoglycosides gentamicin, amikacin, and kanamycin (Fig. 2C) than the

wild-type. When the entire *nuoABCEFGHIJKLMN* and *cyoABCDE* operon were individually deleted

from wild-type cells, the resultant Δ*nuo* and Δ*cyo* strains were more tolerant to both CPP$_{R11}$-APIM

(only tested with Δ*cyo*), CPP$_{RXR}$-PNA and aminoglycosides relative to the wild-type (Fig. 2B,C). This

confirms that CPP$_{R11}$-APIM, CPP$_{RXR}$-PNA, and aminoglycoside tolerance in *cpxR$_{L20Q}$* and *cpxA$_{\Delta16-17}$*

is related to aerobic oxidative phosphorylation and result from downregulation of NDH-I and

cytochrome *bo$_3$* oxidase.

234   The cytoplasmic membrane potential (ΔΨ) directly affects CPP$_{RXR}$-PNA and CPP$_{R11}$-APIM activity

Downregulation of respiratory operons suggested that the ΔΨ formed across the inner membrane

plays a significant role in CPP$_{R11}$-APIM CPP$_{RXR}$-PNA tolerance. We determined the ΔΨ at the

intracellular pH 7.6 of *E. coli* based on the distribution of the lipophilic tetraphenylphosphonium

ion (TPP$^+$) using a TPP$^+$-selective electrode(24). At this pH value the ΔpH is zero and the PMF is

only dependent on $\Delta\Psi$. Wild-type cells had a $\Delta\Psi$ of approx. -140 mV, in accordance with previous

observations(25) (Fig. 2D). The $\Delta\Psi$ was reduced by the $cpxR_{L20Q}$ as well as the $cpxA_{\Delta16-17}$ mutation

(Fig. 2D). Δcyo and Δnuo cells also had a decreased $\Delta\Psi$ compared to wild-type; with values

comparable to $cpxR_{L20Q}$ and $cpxA_{\Delta16-17}$ (Fig. 2D). Thus, it is conceivable that the decreased $\Delta\Psi$ of

$cpxR_{L20Q}$ and $cpxA_{\Delta16-17}$ cells may result from downregulation of the respiratory operons.

E. coli maintains a stable PMF across the inner membrane, such that $\Delta\Psi$ decreases with an

increase in ΔpH in an acidic medium, vice versa in an alkaline media (9) (Fig. 3). Thus, $\Delta\Psi$ can be

modulated by growing the cells at various pH levels. In acidic media (pH 6.0) where $\Delta\Psi$ is

decreased(9) (Fig. 3A), wild-type cells were tolerant to CPP$_{RXR}$-PNA, CPP$_{R11}$-APIM and gentamicin

(Fig. 3A), while an increased $\Delta\Psi$ due to growth in alkaline media (pH 8.0)(9) resulted in sensitivity

to both CPP$_{RXR}$-PNA, CPP$_{R11}$-APIM and gentamicin (Fig. 3B). The changes in $\Delta\Psi$ in the two different

pH-adjusted settings had no effect on tolerance to CPP$_{KFF}$-PNA (Fig. 3A,B). *E. coli* cells deficient of

the F$_1$F$_0$-ATPase or supplemented with alanine have been shown to have an increase in $\Delta\Psi$(26,

27) and were more sensitive to CPP$_{RXR}$-PNA, gentamicin, and CPP$_{R11}$-APIM (only tested with

alanine), but sensitivity to CPP$_{KFF}$-PNA was similar to wild-type (Fig. 3B). Finally, $\Delta\Psi$ is decreased

under anaerobic growth conditions(26) explaining why wild-type under these conditions are more

tolerant to CPP$_{RXR}$-PNA and gentamicin (Fig. 3A). Most likely the effect of $\Delta\Psi$ on antimicrobial

activity is explained by increased inner membrane passage of the polycationic molecules

energetically driven by the negative membrane potential.

Neither $cpxR_{L20Q}$ nor the $cpxA_{\Delta16-17}$ mutations conferred resistance to cationic antimicrobial

peptides

Cationic antimicrobial peptides represent the biggest class of antimicrobial peptides and the

majority of these are amphipathic. Here, Cap11, Cap18, cecropin P1, apidaecin 1B, indolicidin,

protamine, and the most well-known polypeptide antibiotic colistin were tested against wild-type,

$cpxR_{L20Q}$, $cpxA_{\Delta16-17}$, and Ahx^R-3 (Table S4). Colistin, Cap11, Cap18, protamine, and cecropin P1 are

believed to target the cell envelope (disruption of bilayers), while indolicidin (disruption of bilayers

and DNA synthesis) and apideacin 1B (ribosomes) have intracellular targets. An activated Cpx-

response and a decrease in $\Delta\Psi$ did not confer tolerance to any of the tested AMPs (Table S4).

Interestingly, the evolved strain Ahx^R-3 was more susceptible to Cecropin P1, Cap11, Cap18, and

colistin than wild-type (Table S4). One of the major differences between wild-type and Ahx^R-3 is

the two IS1-insertions in *waaB* and *waaO* in Ahx^R-3 (Table 2). This is expected to result in LPS

deficiency, which is associated with an increased susceptibility to cationic antimicrobial peptides

in *E. coli*(28). Accordingly, when *waaBO* were deleted in the wild-type (Δ*waaBO*) they became

more susceptible to Cecropin P1, Cap11, Cap18, and colistin but not to CPP$_{R11}$-APIM.

High CPP$_{RXR}$-PNA tolerance is associated with fitness cost

The doubling times of *cpxR$_{L20Q}$* and *cpxA$_{Δ16-17}$* mutants were 28 min and 27 min, respectively, when

grown aerobically in MHB I at 37°C and somewhat slower than that of wild-type cells (25 min).

The reduced doubling times are likely to result from repression of the respiratory operons in the

respective mutants. Of interest, the evolved mutant, Ahx^R-3 (58 min), which conferred the highest

resistance to R-PNA (8 µM vs. 4 µM for the *cpxR$_{L20Q}$* and *cpxA$_{Δ16-17}$* mutants), grew more than twice

as slow as both the wild-type and the *cpxR$_{L20Q}$* and *cpxA$_{Δ16-17}$* mutants. This shows that CPP$_{RXR}$-PNA

resistance comes with a high fitness cost.

Discussion

Since the late eighties, natural and synthetic CPPs have been developed and modified for applications in both basic- and applied biology including delivery of therapeutic agents, gene editing and cell imaging. Due to efficient delivery, CPPs as carriers for therapeutic compounds to treat various diseases has gained attraction in recent years(5). Although there are many types of CPPs with different physicochemical properties and applications, R-CPPs, are among the most extensively employed and studied(29). PNA, PMO and AMPs are novel, promising antimicrobial agents, of which PNA and PMO require conjugation to a CPP for delivery to the intracellular targets in bacteria(4). Here, we isolated and characterize resistant strains towards an R-CPP. High-level resistance to the arginine-rich CPP-part was difficult to obtain, came with a high fitness cost, and it conferred cross-resistance to aminoglycosides, which is seen as a clear disadvantage for the use of these carriers in antibacterial therapy.

The cytoplasmic membrane is the main barrier for CPP$_{RXR}$-PNA

The initial interaction with the envelope differs between CPP$_{R11}$-APIM/CPP$_{RXR}$-PNA and cationic AMPs. Thus, loss of LPS had no effect on the activity of CPP$_{RXR}$-PNA or CPP$_{R11}$-APIM while this sensitized wild-type to the cationic AMPs (Table S4). It therefore remains unclear why LPS deficiency were selected in Ahx^R-1, 2, and 3. A reduction of $\Delta\Psi$ in the cytoplasmic membrane was found to diminish/reduce CPP$_{RXR}$-PNA and CPP$_{R11}$-APIM activity, demonstrating that a (large) negative $\Delta\Psi$ is important for activity, and most likely for providing sufficient free energy required for membrane translocation. The more detailed mechanism of membrane passage is not known, but could rely on direct, local membrane interruption or be aided by one or more still unidentified transporter protein. However, for CPP$_{RXR}$-PNA neither our ALE study nor a previous screening of the KEIO collection(4) provides evidence for the existence of a single CPP$_{RXR}$-PNA transport mechanism. However, we cannot exclude that CPP$_{RXR}$-PNA is the substrate of multiple transport systems, in which case the absence of one would be compensated by the remaining other(s).

CPP$_{RXR}$-PNA resistance arise in multiple steps

Several genetically different R-PNA^R mutants arose both within and across the two lineages in the ALE-experiment. This suggest clonal interference within the populations of each ALE lineage,

where mutants will compete not just against the susceptible parent but also against each

other(30). Clonal interference influences evolutionary trajectories, which typically leads to a

population with one dominant clone(31), as suggested here for clone Ahx^R-3. The evolutionary

paths leading to three genotypes, from the two linages, speaks to either multiple ways to become

CPP$_{RXR}$-PNA resistance or that they all, more or less, confer the same universal CPP$_{RXR}$-PNA^R

phenotype. Nonetheless, resistance development to PNA conjugated to an R-CPP is complex,

requiring multiple mutations in independent loci to create the CPP$_{RXR}$-PNA^R phenotype (Table 2).

This is in stark contrast to the straightforward resistance development to PNA conjugated to the

lysine-rich CPP, i.e. CPP$_{KFF}$-PNA, by a loss-of-function mutation in the *E. coli sbmA* gene(4). We

show that the *cpx**-phenotype is a major part of the CPP$_{RXR}$-PNA resistance phenotype in Ahx^R-3,

but that additional mutations are required for high-level resistance. As observed for *waaBO*, not

all mutations from the ALE-experiment on their own resulted in CPP$_{RXR}$-PNA tolerance. Either, the

waaBO mutation requires another mutation to synergistically provide CPP$_{RXR}$-PNA tolerance or it

was selected as a mutation that compensate a fitness cost associated with mutations conferring

CPP$_{RXR}$-PNA resistance. Indeed, CPP$_{RXR}$-PNA^R strains have increased doubling time, highlighting

that CPP$_{RXR}$-PNA resistance comes with a cost.

The cpx* phenotype relies on a functional Cpx-system

Our results show that alterations in the first transmembrane domain of CpxA can produce a *cpx**

phenotype. Several gain-of-function mutations in *cpxA* have been reported(16, 17), but most

cluster to and around the second transmembrane domain of CpxA(17). We suggest that the

deletion in the first transmembrane domain leads to a conformational change in the periplasmic

part of CpxA, which either disallows CpxP regulation or simulates the stress signal that actives the

Cpx-response; i.e. a constitutive active CpxA. To the best of our knowledge, *cpxR$_{L20Q}$* is the first

reported mutation in *cpxR* resulting in a *cpx** phenotype. The activated Cpx-response in the

cpxR$_{L20Q}$ mutant was CpxA-dependent; i.e. the *cpxR$_{L20Q}$* mutation does not result in a constitutive

phosphorylated CpxR. Therefore, it is likely that the *cpxR$_{L20Q}$* mutation inhibits/diminish the ability

of CpxA to dephosphorylate CpxR-P to CpxR, which in turn results in a constitutive active Cpx-

response. CpxA and CpxR-P both function as dimers(32). Thus overproducing the respective wild-

type allele in the two mutants restored sensitivity to CPP$_{RXR}$-PNA by either hetero-dimerization of

a mutated and a wild-type protein, with the wild-type being dominant to the mutant, or by homo-

dimerization of either mutated or wild-type proteins, but with wild-type dimer domination due to
abundance.

Low cytoplasmic membrane potential correlates with tolerance to CPP$_{RXR}$-PNA, CPP$_{R11}$-APIM, and
aminoglycosides

We found that a reduction of the $\Delta\Psi$ component of PMF across the cytoplasmic membrane
increased tolerance to both CPP$_{RXR}$-PNA, CPP$_{R11}$-APIM, and aminoglycosides; the latter in
agreement with previous studies in *Bacillus subtilis* and *E. coli*(33). The reduced activity of both
CPP$_{RXR}$-PNA, CPP$_{R11}$-APIM, and aminoglycosides when $\Delta\Psi$ was decreased indicates that crossing
of the inner (and not the outer) membrane is the rate limiting and energy dependent step shared
by this type of (polycationic) compounds as previously proposed for aminoglycosides (34). Once
inside the bacterial cells, aminoglycosides interacts with the 30S subunit of ribosomes, RXR-PNA
prevent *acpP* translation, and CPP$_{R11}$-APIM prevent β-clamp function (Fig. 4A). In wild-type cells,
CPP$_{RXR}$-PNA (and possibly also aminoglycosides and, CPP$_{R11}$-APIM) do not activate the Cpx-
response. We propose that *cpx** dependent tolerance to CPP$_{RXR}$-PNA, CPP$_{R11}$-APIM, and
aminoglycosides is caused by downregulation of respiratory complexes in the cytoplasmic
membrane (Fig. 4B) thereby decreasing $\Delta\Psi$ across the cytoplasmic membrane at least partly due
to the reduced ATP level and consequently reduced proton chemical potential over the
membrane, making it less energetically favorable for R-CPP to transvers the cytoplasmic
membrane (Fig. 4B). CPP$_{KFF}$-PNA is internalized in a SbmA-dependent manner(4) (for which a
membrane potential driving force apparently is not important) and therefore the CPP$_{RXR}$-PNA^{R}
strains were sensitive to CPP$_{KFF}$-PNA.

The cpx* phenotype could lead to the acquisition of other mutations to promote resistance or
persistence

The *cpx** phenotype provides moderate resistance to both CPP$_{RXR}$-PNA and aminoglycosides – all
without a major loss of fitness. Thus, in a clinical setting, *cpx** mutants could grow in the presence
of moderate levels of either compound, which in turn would create an evolutionary opportunity
to adapt and evolve by subsequent acquisition of mutations to further increase resistance or to
assist in persistent colonization. Several recent studies have shown that adaptation, may have a

significant impact on the selection of highly resistant/persistent clones for *Mycobacterium*

(ribosomal mutations)(35), *Pseudomonas aeruginosa* (efflux pump regulator mutation)(36), and

E. coli (formation of subpopulations with an increased overall mutation rate)(37). Thus, the *cpx**

phenotype may be added to this list.

Materials and Methods
Growth conditions

Cells were grown in Luria–Bertani Broth (LB) medium or Müller-Hinton I Broth (MHBI) at 37°C with aeration unless stated otherwise. When necessary, antibiotics were added to the following concentrations: chloramphenicol, 20 µg/mL; ampicillin, 150 µg/mL; kanamycin, 50 µg/mL; tetracyclin, 10 µg/mL.

Peptide conjugates

PNA-peptide conjugates were synthezised as described previously (1, 4). CPP$_{R11}$-APIM (RWLVK* with the complete sequence Ac-MD-RWLVK-GILQWRKI-RRRRRRRRRRR) is described previously(22).

Bacterial strains and plasmids

A MG1655 strains used in this study are listed in Table S5 and all primer sequences in Table S6. For deleting and overexpressing genes, we used the KEIO collection(38) and the ASKA library(39), respectively. Shortly, the *cpxA, cpxR, degP,* or *cpxP* deletion/replacement mutations were obtained from the KEIO collection(38) and P1 transduced(40) into *E. coli* MG1655 to create ALO6524, ALO6525, ALO7548, and ALO7549, respectively. For overexpression of genes, pCA24n based plasmids from the ASKA collection was transformed into *E. coli* MG1655 selecting for chloramphenicol.

Deletion of the *nuoABCEFGHIJKLMN, cyoABCDE,* and *waaBO* chromosomal regions in wild-type was constructed by the lambda red procedure(41). Briefly, primer sequences were obtained from Baba *et al.* (38) and are listed in Suppl. Table 2. The *nuo*_cat_Fw primer corresponds to the 5' primer used to delete *nuoA* in Baba *et al.* (38), while the *nuo*_cat_Rv primer corresponds to the 3' primer used to delete *nuoN* in Baba *et al.* (38). The *cyo*_cat_Fw primer corresponds to the 5' primer used to delete *cyoA* in Baba *et al.* (38), while the *cyo*_cat_Rv primer corresponds to the 3' primer used to delete *cyoE* in Baba *et al.* (38). The *waaB*_cat_Fw primer corresponds to the 5' primer used to delete *waaB* in Baba *et al.* (38), while the *waaO*_cat_Rv primer corresponds to the 3' primer used to delete *waaO* in Baba *et al.* (38). DNA fragments were PCR amplified using the

primer sets *nuo*_cat_Fw + *nuo*_cat_Rv, *cyo*_cat_Fw + *cyo*_cat_Rv, and *waaB*_cat_Fw +

*waaO*_cat_Rv to delete *nuoABCEFGHIJKLMN, cyoABCDE,* and *waaBO,* respectively. PCR

amplification was done using high-fidelity Phusion DNA polymerase (Thermo Scientific™)

according to the manufacturer's specifications with pKD3 as template. The resultant DNA

fragments were introduced into wild-type cells bearing pKD46(41), selecting for chloramphenicol

resistance. Each deletion was verified by PCR creating Δ*nuoABCEFGHIJKLMN* (Δ*nuo*; ALO7555),

Δ*cyoABCDE* (Δ*cyo*; ALO7556), and Δ*waaBO* (ALO6972).

The *cpxR*$_{L20Q}$ mutations were introduced into the wild-type using the Tet/SacB counter-selection

and recombineering approach(42). Briefly, the *tet-sacB* cassette flanked by appropriate homology

arms was amplified using the primer set CpxR_*tet-sacB*_Fw + CpxR_*tet-sacB*_Rv using pTETSACB

as a template for exchange of *cpxR* in the wild-type. The resultant DNA fragments were introduced

into wild-type bearing pKD46(41), selecting for tetracycline and ampicillin resistance at 32°C (to

keep pKD46 in the cell) creating *cpxR*:*tet-sacB* pKD46. The deletion was verified by PCR. Next, the

CpxR$_{L20Q}$ mutations were amplified from Ahx^R-3 with homology flaking the inserted *tet-sacB*

cassettes, using the primer sets CpxR$_{L20Q}$_Fw + CpxR$_{L20Q}$_Rv. The resultant DNA fragments

containing the *cpxR*$_{L20Q}$ mutation were introduced into *cpxR*:*tet-sacB* selecting for growth on LB

agar containing 6% sucrose. *cpxR* genes from clones that were tetracycline sensitive and able to

grow on sucrose were sequenced to confirm the introduction of the specific mutations from Ahx^R-

3 into wild-type, thus creating the strain *cpxR*$_{L20Q}$ (ALO7180). *cpxA* was deleted in *cpxR*$_{L20Q}$ and

cpxR was deleted in *cpxA*$_{Δ16-17}$ by P1 transduction (as above) creating *cpxR*$_{L20Q}$ Δ*cpxA* (ALO7550)

and *cpxA*$_{Δ16-17}$ Δ*cpxR* (ALO7551), respectively. In addition, *nuoABCEFGHIJKLMN* and *cyoABCDE*

were deleted as above in *cpxR*$_{L20Q}$ using the lambda red procedure(41) created *cpxR*$_{L20Q}$ Δ*nuo*

(ALO7552) and *cpxR*$_{L20Q}$ Δ*cyo* (ALO7553), respectively.

Adaptive Laboratory Evolution (ALE) experiment

Wild-type was evolved for 20 days to CPP$_{RXR}$-PNA, with two lineages evolved in parallel for each

drug. Here, 1 x 10^5 CFU/mL of cells were inoculated in 2-fold peptide-PNA gradient in 8 dilutions

in 100 µl MHBI in a 96-well polystyrene microtitre plate and grown for 18-24 hours at 37°C. The

following day the population grown in the highest concentration of peptide-PNA where re-diluted

to 1 x 10^5 CFU/mL and inoculated in a fresh 2-fold peptide-PNA gradient as above. After 20

successive passages, five peptide-PNA resistant single clones were isolated from each lineage and

whole genome sequenced. As control, *E. coli* MG1655 was evolved to the MHBI in four independent biological replicates without the presence of peptide-PNA. Here, two clones were isolated and whole genome sequenced for each lineage.

Extraction of genomic DNA

Genomic DNA of *E. coli* MG1655 was isolated using phenol/chloroform extraction. Shortly, cells from a 15 ml culture were collected at an OD_{600} of 0.5 by centrifugation at 5.000 ×g for 10 minutes. The pellet was resuspeded in 1 ml 0.9% NaCl, followed by a centrifugation at 10.000 ×g for 5 minutes. The cells were lysed by incubation at 37°C for 1 hour with 600 µl lysis buffer (Tris-EDTA buffer (TE buffer, pH 8.0), 10% (w/v) SDS, 20 mg/ml proteinase K, 1 mg/ml RNase A). Following two phenol- and two chloroform-extraction, DNA was precipitated using NaCl (0.2 M) and 0.6 vol. isopropanol, washed in 80% ethanol and resuspended in TE buffer (pH 8.0).

Whole genome sequencing and data analysis

Reads were mapped to MG1655 NC_000913.3 using BWA-MEM. Variants were extracted using Freebayes (only variants with >50 % frequency were retained). For location of IS insertion, mapped reads with <60 mapQuality were selected and paired reads (CIGAR-Left-clip/Right-Clip, CIGAR-D and CIGAR-I) were aligned to NC_000913.3 using NCBI Blast to confirm deletion /insertion.

Antimicrobial Susceptibility Testing

MIC values were determined by broth microdilution according to the standard protocol(43) with a few modifications. Shorty, an overnight bacterial cell culture was diluted to approximately 5 × 10^5 CFU/mL in MHBI. 100 µL bacterial solution was dispensed into a low-bind 96-well plate (Thermo-Scientific, cat.no. 260895) along with 11 µL of the test compound (2-fold dilutions). The plate was incubated at 37°C for 18-24 h. The MIC was determined as the lowest concentration, which inhibited visible growth in the wells (OD_{595} nm < 0.1). Peptide-PNAs were dissolved in 0,02% acetic acid containing 0,4% bovine serum albumine and dilution were performed in 0,01% acetic acid containing 0,2% BSA, while gentamicin, amikacin, kanamycin, cecropin P1, Cap11 (Genescript), Cap18 (Genescript), apidaecin 1B (Genescript), colistin, indolicidin and protamine

(Sigma-Aldrich) was diluted according to vendors specifications. To determine MIC under acidic

and alkaline conditions, MHBI was buffered to pH 6.0 or pH 8.0 using citrate buffer (pH 3.0) and

carbonate-bicarbonate buffer (pH 10.0), respectively. Anaerobic growth for MIC determinations

was performed in a double sealed bag using anaerobic atmosphere generation bags (Becton,

Dickinson and Company, New Jersey, USA). MHBI was supplemented with 0.1 mM Isopropyl β-D-

1-thiogalactopyranoside (IPTG), when tested strains harbored pCA24n-based plasmids.

475 Total RNA extraction

RNA were isolated from cells by the hot phenol method. Shortly, cells were grown in cation

adjusted MHBI to mid-log phase, an OD_{595} of 0.2 (2.5×10^9 cells/ml), by shaking at 150 rpm and

37°C. To induce protein expression in cells harboring the inducible pCA24n-based plasmid, IPTG

was added to a concentration of 0.1 mM at an optical density OD_{595} 0.005. Bacteria from 2-ml

aliquot were harvested by centrifugation at 4°C and 17.000×g for 5 minutes. The pellets were

resuspended in the same volume of RNAprotect Bacteria Reagent (QIAGEN, Denmark), incubated

at room temperature for 5 minutes, followed by a centrifugation using the same conditions as

above. The supernatants were discarded and the pellets were either stored at -80°C until needed

or immediately used for RNA extraction.

To extract RNA, the pellets were lyzed in 200 µl Solution I [0.3 M sucrose, 0.01 M sodium acetate

(NaOAc) and pH 4.5] at 4°C and mixed thoroughly. Then, 200 µl of Solution II (2% (w/w) Sodium

dodecyl sulfate (SDS), 0.01 M NaOAc and pH 4.5) were added at room temperature and solutions

were gently mixed by inversion of tubes. Next, a volume of 400 µl of acid

phenol:chloroform:isoamyl alcohol 125:24:1 was added and immediately mixed (vortex) to

emulsify. The samples were incubated at 65°C for 10 minutes and mixed (vortex) for 5-10 seconds

every minute. Then, tubes were transferred on ice for 5 minutes and centrifuged at 14.000 rpm

for 5 minutes. The aqueous phase was transferred to a new polypropylene tube and one-fifth

volume of chloroform was added and gently mixed by inversion, followed by centrifugation at

14.000 rpm for 5 minutes. This step was repeated by addition of the same volume of

chloroform:isoamyl alcohol 24:1. To precipitate RNA, the aqueous phase was transferred to a new

tube, mixed with the same volume of isopropanol and incubated at -80°C for a minimum of 1 hour.

RNA was precipitated by spinning down at 4°C and 12.000 ×g for 30 minutes, followed by wash

with 1 ml 70% ethanol and centrifugation at 4°C and 12.000 ×g for 5 minutes. The pellets were resuspended in 30 µl nuclease free water and RNA concentration and purity was determined using a NanoDrop.

Following removal of genomic DNA by DNase I (Qiagen, Venlo, Netherlands), RNA were purified using RNeasy MinElute Cleanup kit (Qiagen) and following the vendors instructions.

qRT-PCR.

For qRT-PCR experiments, after genomic DNA digestion, 1 µg of total RNA was retrotranscribed using QuantiTect Reverse Transcription kit (Qiagen). Primers designed to amplify *cpxP*, *nuoA*, and *cyoA* (Table 8) were targeted to regions of unique sequences within the genes in wild-type *E. coli* MG1655. The qRT-PCR was performed using the TB Green Premix Ex Taq II (Takara Bio Inc., Shiga, Japan) on a QuantStudio 3 Real-Time PCR System (Thermo Fisher Scientific, Massachusetts, United States). All data were normalized to the reference genes *hctA* and *cysG*. These data were transformed to log2 to obtain a change difference (n-fold) between strains.

Determination of membrane potential (Δψ) by measuring Tetraphenyl phosphonium (TTP$^+$) uptake using the TPP$^+$-selective electrode method

The membrane potential was determined by measuring the uptake of the permeating lipophilic cationic probe, Tetraphenyl phosphonium (Merck KGaA, Germany). An overnight culture was diluted (5.000-fold) in MHBI, and cells from 25 ml culture were harvested when OD$_{595}$ reached 0.2. Cells were washed twice in 0.1 M Tris·HCl (pH 8.1) using the conditions mentioned above, and the pellets were gently resuspended in 1 ml of the same buffer. To generate cells permeable to TPP$^+$, after a first incubation for 6 minutes at 36°C with occasional agitation, bacteria were treated with K$^+$-EDTA to a concentration of 10 mM and then incubated at 36°C for 3 more minutes. Cells were diluted 10-fold in ice-cold 0.1 M potassium phosphate buffer (0.1 M K$_2$HPO$_4$; 0.1 M KH$_2$PO$_4$; pH 7.5), to remove EDTA, followed by immediate centrifugation in the cold at 13.000 rpm for 7 minutes. The cells were washed twice in the same ice-cold buffer using the parameters mentioned above, and the pellets were resuspended, to an OD$_{595}$ of 20.0, in 10 mM potassium phosphate

buffer (pH 7.5) containing 200 mM NaCl, 0.14 mM $CaCl_2$, 0.1 mM $MgSO_4$ and 0.01 mM $MnCl_2$. The

cells were kept on ice until use.

Bacteria were added to an OD_{595} of 4.0 in 10 ml of the same buffer supplemented with glucose

and TPP^+ to a concentration of 5 mM and 10 μM, respectively, and TPP^+ uptake was measured

utilizing the TPP^+-selective electrode method described by Hosoi *et al*(44). TPP^+ concentration in

the external medium was determined using a Kwik-Tip^{TM} Ag/AgCl half-cell electrode, which was

constructed according to the instructions provided by the manufacturer (World Precision

Instruments, Sarasota, FL). Both the TPP^+-selective electrode and a Flexible Dri-Ref reference

electrode (World Precision Instruments, Sarasota, FL) were connected to an Jenway 3510 ion

meter (Cole-Parmer, Staffordshire, UK) and LabTrax 4-Channel Data Acquisition system (WPI).

Finally, TPP^+ uptake measurements were carried out in a 15 ml closed tube, at 30°C and pH 7.5 for

10 minutes, and data were recorded using the iWorx LabScribe recording and analysis software

(iWorx Systems Inc., Dover, NH).

539
540
541

References and Notes

1.   Good L, Nielsen PE. Antisense inhibition of gene expression in bacteria by PNA targeted to mRNA. Nat Biotechnol. 1998;16(4):355-8.

2.   Wesolowski D, Tae HS, Gandotra N, Llopis P, Shen N, Altman S. Basic peptide-morpholino oligomer conjugate that is very effective in killing bacteria by gene-specific and nonspecific modes. Proc Natl Acad Sci U S A. 2011;108(40):16582-7.

3.   Good L, Sandberg R, Larsson O, Nielsen PE, Wahlestedt C. Antisense PNA effects in Escherichia coli are limited by the outer-membrane LPS layer. Microbiology. 2000;146 (Pt 10):2665-70.

4.   Ghosal A, Vitali A, Stach JE, Nielsen PE. Role of SbmA in the uptake of peptide nucleic acid (PNA)-peptide conjugates in E. coli. ACS Chem Biol. 2013;8(2):360-7.

5.   Habault J, Poyet JL. Recent Advances in Cell Penetrating Peptide-Based Anticancer Therapies. Molecules. 2019;24(5).

6.   Ingledew WJ, Poole RK. The respiratory chains of Escherichia coli. Microbiol Rev. 1984;48(3):222-71.

7.   Gunner MR, Amin M, Zhu X, Lu J. Molecular mechanisms for generating transmembrane proton gradients. Biochim Biophys Acta. 2013;1827(8-9):892-913.

8.   Vik SB, Antonio BJ. A mechanism of proton translocation by F1F0 ATP synthases suggested by double mutants of the a subunit. J Biol Chem. 1994;269(48):30364-9.

9.   Kashket ER. The proton motive force in bacteria: a critical assessment of methods. Annu Rev Microbiol. 1985;39:219-42.

10.   Bury-Mone S, Nomane Y, Reymond N, Barbet R, Jacquet E, Imbeaud S, et al. Global analysis of extracytoplasmic stress signaling in Escherichia coli. PLoS Genet. 2009;5(9):e1000651.

11.   Raivio TL, Leblanc SK, Price NL. The Escherichia coli Cpx envelope stress response regulates genes of diverse function that impact antibiotic resistance and membrane integrity. J Bacteriol. 2013;195(12):2755-67.

12.   Weber RF, Silverman PM. The cpx proteins of Escherichia coli K12. Structure of the cpxA polypeptide as an inner membrane component. J Mol Biol. 1988;203(2):467-78.

13.   Itou H, Tanaka I. The OmpR-family of proteins: insight into the tertiary structure and functions of two-component regulator proteins. J Biochem. 2001;129(3):343-50.

14.   Raivio TL, Popkin DL, Silhavy TJ. The Cpx envelope stress response is controlled by amplification and feedback inhibition. J Bacteriol. 1999;181(17):5263-72.

15.   Buelow DR, Raivio TL. Cpx signal transduction is influenced by a conserved N-terminal domain in the novel inhibitor CpxP and the periplasmic protease DegP. J Bacteriol. 2005;187(19):6622-30.

16.   Danese PN, Snyder WB, Cosma CL, Davis LJ, Silhavy TJ. The Cpx two-component signal transduction pathway of Escherichia coli regulates transcription of the gene specifying the stress-inducible periplasmic protease, DegP. Genes Dev. 1995;9(4):387-98.

17.   Raivio TL, Silhavy TJ. Transduction of envelope stress in Escherichia coli by the Cpx two-component system. J Bacteriol. 1997;179(24):7724-33.

18.   Castano I, Bastarrachea F, Covarrubias AA. gltBDF operon of Escherichia coli. J Bacteriol. 1988;170(2):821-7.

19.   Raetz CR, Dowhan W. Biosynthesis and function of phospholipids in Escherichia coli. J Biol Chem. 1990;265(3):1235-8.

20.   Alksne LE, Anthony RA, Liebman SW, Warner JR. An accuracy center in the ribosome conserved over 2 billion years. Proc Natl Acad Sci U S A. 1993;90(20):9538-41.

21.   Otto K, Silhavy TJ. Surface sensing and adhesion of Escherichia coli controlled by the Cpx-signaling pathway. Proc Natl Acad Sci U S A. 2002;99(4):2287-92.

22.   Nedal A, Raeder SB, Dalhus B, Helgesen E, Forstrom RJ, Lindland K, et al.
Peptides containing the PCNA interacting motif APIM bind to the beta-clamp and inhibit
bacterial growth and mutagenesis. Nucleic Acids Res. 2020;48(10):5540-54.
23.   Price NL, Raivio TL. Characterization of the Cpx regulon in Escherichia coli strain
MC4100. J Bacteriol. 2009;191(6):1798-815.
24.   Hirota N, Matsuura S, Mochizuki N, Mutoh N, Imae Y. Use of lipophilic cation-
permeable mutants for measurement of transmembrane electrical potential in metabolizing cells
of Escherichia coli. J Bacteriol. 1981;148(2):399-405.
25.   Suzuki T, Murakami T, Iino R, Suzuki J, Ono S, Shirakihara Y, et al. F0F1-
ATPase/synthase is geared to the synthesis mode by conformational rearrangement of epsilon
subunit in response to proton motive force and ADP/ATP balance. J Biol Chem.
2003;278(47):46840-6.
26.   Jensen PR, Michelsen O. Carbon and energy metabolism of atp mutants of
Escherichia coli. J Bacteriol. 1992;174(23):7635-41.
27.   Ye JZ, Su YB, Lin XM, Lai SS, Li WX, Ali F, et al. Alanine Enhances
Aminoglycosides-Induced ROS Production as Revealed by Proteomic Analysis. Front Microbiol.
2018;9:29.
28.   Ebbensgaard A, Mordhorst H, Aarestrup FM, Hansen EB. The Role of Outer
Membrane Proteins and Lipopolysaccharides for the Sensitivity of Escherichia coli to
Antimicrobial Peptides. Front Microbiol. 2018;9:2153.
29.   Futaki S, Nakase I. Cell-Surface Interactions on Arginine-Rich Cell-Penetrating
Peptides Allow for Multiplex Modes of Internalization. Acc Chem Res. 2017;50(10):2449-56.
30.   Hughes D, Andersson DI. Evolutionary Trajectories to Antibiotic Resistance. Annu
Rev Microbiol. 2017;71:579-96.
31.   Gerrish PJ, Lenski RE. The fate of competing beneficial mutations in an asexual
population. Genetica. 1998;102-103(1-6):127-44.
32.   Mechaly AE, Haouz A, Sassoon N, Buschiazzo A, Betton JM, Alzari PM.
Conformational plasticity of the response regulator CpxR, a key player in Gammaproteobacteria
virulence and drug-resistance. J Struct Biol. 2018;204(2):165-71.
33.   Bryan LE, Kwan S. Roles of ribosomal binding, membrane potential, and electron
transport in bacterial uptake of streptomycin and gentamicin. Antimicrob Agents Chemother.
1983;23(6):835-45.
34.   Vanhoof R, Sonck P, Hannecart-Pokorni E. The role of lipopolysaccharide anionic
binding sites in aminoglycoside uptake in Stenotrophomonas (Xanthomonas) maltophilia. J
Antimicrob Chemother. 1995;35(1):167-71.
35.   Gomez JE, Kaufmann-Malaga BB, Wivagg CN, Kim PB, Silvis MR, Renedo N, et
al. Ribosomal mutations promote the evolution of antibiotic resistance in a multidrug
environment. Elife. 2017;6.
36.   Frimodt-Moller J, Rossi E, Haagensen JAJ, Falcone M, Molin S, Johansen HK.
Mutations causing low level antibiotic resistance ensure bacterial survival in antibiotic-treated
hosts. Sci Rep. 2018;8(1):12512.
37.   El Meouche I, Dunlop MJ. Heterogeneity in efflux pump expression predisposes
antibiotic-resistant cells to mutation. Science. 2018;362(6415):686-90.
38.   Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, et al. Construction of
Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio collection. Mol Syst
Biol. 2006;2:2006 0008.
39.   Kitagawa M, Ara T, Arifuzzaman M, Ioka-Nakamichi T, Inamoto E, Toyonaga H,
et al. Complete set of ORF clones of Escherichia coli ASKA library (a complete set of E. coli K-
12 ORF archive): unique resources for biological research. DNA Res. 2005;12(5):291-9.
40.   J M. Experiments in Molecular Genetics1972.

41.	Datsenko KA, Wanner BL. One-step inactivation of chromosomal genes in Escherichia coli K-12 using PCR products. Proc Natl Acad Sci U S A. 2000;97(12):6640-5.

42.	Li XT, Thomason LC, Sawitzke JA, Costantino N, Court DL. Positive and negative selection using the tetA-sacB cassette: recombineering and P1 transduction in Escherichia coli. Nucleic Acids Res. 2013;41(22):e204.

43.	Hancock REW. MIC Determination By Microtitre Broth Dilution Method 1999 [Available from: http://cmdr.ubc.ca/bobh/method/mic-determination-by-microtitre-broth-dilution-method/.

44.	Hosoi S, Mochizuki N, Hayashi S, Kasai M. Control of membrane potential by external H+ concentration in Bacillus subtilis as determined by an ion-selective electrode. Biochim Biophys Acta. 1980;600(3):844-52.

45.	Liu X, Li Y, Guo Y, Zeng Z, Li B, Wood TK, et al. Physiological Function of Rac Prophage During Biofilm Formation and Regulation of Rac Excision in Escherichia coli K-12. Sci Rep. 2015;5:16074.

46.	Keyhani NO, Bacia K, Roseman S. The transport/phosphorylation of N,N'-diacetylchitobiose in Escherichia coli. Characterization of phospho-IIB(Chb) and of a potential transition state analogue in the phosphotransfer reaction between the proteins IIA(Chb) AND IIB(Chb). J Biol Chem. 2000;275(42):33102-9.

47.	Conter A, Menchon C, Gutierrez C. Role of DNA supercoiling and rpoS sigma factor in the osmotic and growth phase-dependent induction of the gene osmE of Escherichia coli K12. J Mol Biol. 1997;273(1):75-83.

48.	Sukhodolets MV, Cabrera JE, Zhi H, Jin DJ. RapA, a bacterial homolog of SWI2/SNF2, stimulates RNA polymerase recycling in transcription. Genes Dev. 2001;15(24):3330-41.

49.	Yethon JA, Heinrichs DE, Monteiro MA, Perry MB, Whitfield C. Involvement of waaY, waaQ, and waaP in the modification of Escherichia coli lipopolysaccharide and their role in the formation of a stable outer membrane. J Biol Chem. 1998;273(41):26310-6.

50.	Ma Q, Yang Z, Pu M, Peti W, Wood TK. Engineering a novel c-di-GMP-binding protein for biofilm dispersal. Environ Microbiol. 2011;13(3):631-42.

51.	Guyer MS, Reed RR, Steitz JA, Low KB. Identification of a sex-factor-affinity site in E. coli as gamma delta. Cold Spring Harb Symp Quant Biol. 1981;45 Pt 1:135-40.

52.	Charbon G, Campion C, Chan SH, Bjorn L, Weimann A, da Silva LC, et al. Re-wiring of energy metabolism promotes viability during hyperreplication stress in E. coli. PLoS Genet. 2017;13(1):e1006590.

Acknowledgments

This paper is dedicated to the memory of Mads Christian Guldbæk (1991-2019). **Funding**: This research was funded by grants from the Danish National Research Foundation (DNRF120), the Novo Challenge Center for Peptide-Based Antibiotics (NNF16OC0021700), A.P. Møller Lægefonden, Augustinus Fonden, Aase og Ejnar Danielsens Fond, and Brødrene Hartmanns Fond, the program NTNU Health at Norwegian University of Science and Technology (NTNU), and Trond Mohn foundation, Norway. **Author contributions**: Conceptualization: JFM, PEN, and ALO; data curation: JFM and AK; formal analysis: JFM, AK, and ALO; investigation: JFM and AK; methodology: JFM, AK, GC, and ALO; project administration: JFM and ALO; resources: JFM, PEN, and ALO; bioinfomatics: GC; supervision: PEN and ALO; manuscript writing: JFM; manuscript

687 editing: JFM, AK, GC, MO, PEN, and ALO. **Competing interests**: The authors declare that they

688 have no competing interests. **Data and materials availability**: All data needed to evaluate the

689 conclusions in the paper are present in the paper and/or the Supplementary Materials.

690 Additional data are available from authors upon request. All figures were created with

691 BioRender.com.

692
693

 Figures and Tables

Fig. 1. *cpxR$_{L20Q}$* and *cpxA$_{Δ16-17}$* activates the Cpx-response.

A schematic model of the Cpx, Rcs, Psp, Bae, and σ^E stress response to R-PNA in the cytoplasmic membrane.

Here, R-PNA does not activate the Cpx-response; however, an activated Cpx-response is the only of the

five-tested extracytoplasmic stress responses that diminish R-PNA activity. The green plus sign indicates a

constitutive active ESR where an activated Cpx-response confers resistance to R-PNA. A and B: MIC

determinations for CPP-PNA/APIM for wild-type and mutants in µM. MIC were determined using broth

dilutions (see Methods) at 37°C (no shaking). Peptide-PNA sequences are listed in Table 1. pCA24n-based

plasmids were induced with 0.1 mM IPTG. C: *cpxP* expression determined by RT-qPCR (see Methods). All

values are indicated as fold change relative to wild-type (dashed line). Experiments were performed in

triplicate. D: MIC determinations for activated extracytoplasmic stress response-systems in µM (see text

for details). The five extracytoplasmic stress responses were individually activated by overproduction of

nlpE for the Cpx-pathway, *baeR* for the Bae-pathway, *rcsB* for the Rcs-pathway, *pspF* for the Psp-pathway,

and *rpoE* for the σ^E-pathway(10). A dashed line indicates the wild-type MIC. Peptide-PNA sequences are

listed in Table 1. pCA24n-based plasmids were induced with 0.1 mM IPTG.

Fig 2. Cpx-mediated downregulation of NDH-I and cytochrome *bo$_3$* oxidase leads to a decreased

ΔΨ across the cytoplasmic membrane that confers resistance to CPP$_{R11}$-APIM, CPP$_{RXR}$-PNA, and

aminoglycosides.

A schematic model of an activated Cpx-response (shown here with a green plus sign) that leads to

downregulation of both *nuoA* (NDH-I) and *cyoA* (cytochrome *bo$_3$* oxidase), which consequently results in a

decreased ΔΨ across the cytoplasmic membrane that confers cross resistance to CPP$_{RXR}$-PNA and

aminoglycosides. A: *nuoA* and *cyoA* expression determined by RT-qPCR (see Methods). All values are

indicated as fold change relative to wild-type (dashed line). Experiments were performed in biological

triplicates. B and C: MIC values for CPP-PNA/APIM (in µM) and aminoglycosides (in µg/mL) for wild-type

and mutants. MIC were determined using broth dilutions (see Methods) at 37°C (no shaking). Peptide-PNA

sequences are listed in Table 1. D: ΔΨ across the cytoplasmic membrane as measured by the TPP$^+$ uptake

for wild-type, *cpxR$_{L20Q}$*, *cpxA$_{Δ16-17}$*, Δ*cyo*, and Δ*nuo*. Experiments were performed as a minimum in biological

triplicates.

Fig. 3. The magnitude of ΔΨ across the cytoplasmic membrane determines CPP$_{RXR}$-PNA and

aminoglycoside tolerance.

A schematic model of PMF across the cytoplasmic membrane. pH_i is pretty constant over the growth pH_o range (5.5 to 8.5), but the contributions of $\Delta\Psi$ and ΔpH vary(9). A: MIC determinations for wild-type tested against CPP_{RXR}-PNA, CPP_{R11}-APIM, CPP_{KFF}-PNA and gentamicin anaerobically or at pH 6.0. CPP-PNA/APIM MICs are given in µM (left y-axes) and gentamicin MICs are given in µg/mL (right y-axes). Wild-type MIC are given for aerobic growth conditions at pH 7.2 identical to the growth conditions for the ALE-experiment. B: MIC determinations for wild-type tested against CPP_{RXR}-PNA, CPP_{R11}-APIM, CPP_{KFF}-PNA and gentamicin at pH 8.0, supplemented w. 100 mM alanine, or with either *atpA* or *atpB* deleted. Peptide-PNA MICs are given in µM (left y-axes) and gentamicin MICs are given in µg/mL (right y-axes). Wild-type MIC are given for aerobic growth conditions at pH 7.2 identical to the growth conditions for the ALE-experiment.

Fig. 4. CPP_{RXR}-PNA, CPP_{R11}-APIM, and aminoglycoside uptake wild-type (A) and *cpx** (B).

A schematic model of CPP_{RXR}-PNA, CPP_{R11}-APIM, and aminoglycoside (AG) internalization across the cytoplasmic membrane and action in wild-type (A) and a *cpx**mutant (B). A: In the wild-type CPP_{RXR}-PNA, CPP_{R11}-APIM, and AG crossing of the cytoplasmic membrane from the periplasmic space and into the cytosol correlates with the magnitude of the $\Delta\Psi$. The respiratory complexes NADH:ubiquinone oxidoreductase complex I (NDH-I) transcribed from the *nuoABCEFGHIJKLMN*-operon (*nuo*) and cytochrome bo_3 ubiquinol oxidase (Cyo) transcribed from the *cyoABCDE* operon (*cyo*) are shown. CPP_{RXR}-PNA, AG, and CPP_{R11}-APIM in the cytosol leads to growth arrest by steric hindrance of ribosome binding to *acpP* mRNA for CPP_{RXR}-PNA (A.a), binding to the 30S subunit of ribosomes impairing translational accuracy (A.b), and binding to the β-clamp, respectively. B: In a *cpx** mutant CPP_{RXR}-PNA, CPP_{R11}-APIM, and AG are not able to transvers the cytoplasmic membrane from the periplasmic space and into the cytosol. The constitutive active Cpx-response downregulates *nuo, cyo*, which results in a reduced $\Delta\Psi$, which correlates with reducing killing for all the compounds. Lack of CPP_{RXR}-PNA and AG in the cytosol leads to translation of *acpP* mRNA (B.a) wild-type like 30S ribosomal function (B.b), and wild-type like β-clamp function, respectively, both leading to survival during CPP_{RXR}-PNA, CPP_{R11}-APIM, and AG treatment.

Table 1. Minimum inhibitory concentration to peptide-PNA (in µM).

Compound[a]	CPP	Target gene /protein	Wild-type	R-PNA[R] Ahx[R]-1	R-PNA[R] Ahx[R]-2	R-PNA[R] Ahx[R]-3
CPP_{RXR}-PNA(4)		$acpP$[d]	0.5	8	8	8
CPP_{RXR}-PNA$_{mm}$(4)	Arginine-rich CPP_{RXR}[a]	$acpP$ mismatch[e]	>16	>16	>16	>16
CPP_{RXR}-PNA$_{ftsZ}$(4)		$ftsZ$[f]	0.5	8	8	8
CPP_{RXR}-PNA$_{ftsZ\_mm}$(4)		$ftsZ$ mismatch[g]	>16	>16	>16	>16
CPP_{R11}-APIM(22)	Arginine-rich CPP_{R11}[b]	β-clamp (DnaN)[h]	1	N/A	N/A	N/A
PP_{KFF}-PNA(4)	Lysine-phenylalanine CPP_{KFF}[c]	$acpP$[d]	0.25	0.25	0.25	0.25
CPP_{KFF}-PNA$_{mm}$(4)		$acpP$ mismatch[f]	>16	>16	>16	>16

a a Arginine-rich CPP_{RXR}: H-(R-Ahx-R)$_4$-Ahx-(βAla). For the CPP-part amino acids are shown in uppercase letters. Ahx: 6-aminohexanoic acid;; [b] Arginine-rich CPP_{R11}: RRRRRRRRRRR; [c] Lysine-rich CPP_KFF: H-(KFF)$_3$K-eg1; [d] $acpP$ target sequence (5'-3'): H-ctcatactct-NH$_2$; [e] $acpP$ mismatch sequence (the two mismatch are underlined, 5'-3'): H-ctcttacact-NH$_2$; [f] $ftsZ$ target sequence (5'-3'): H-ttcaaacatagt-NH$_2$; [g] $ftsZ$ mismatch sequence (the two mismatch are underlined, 5'-3'): [h] H-ttctaacaaagt-NH$_2$; [i] Ac-MD-RWLVK-GILQWRKI (RWLVK*). MIC determinations for peptide-PNA are in µM. MIC were determined using broth dilutions (see Methods) at 37°C (no shaking).

Table 2. CPP$_{RXR}$-PNA resistant genotypes

Mutation type[a] IS	IN	DEL	SNP	Region/gene[b]	Linage[c] Frequency[d] Strain	Evolved towards CPP$_{RXR}$-PNA #1 2/5 Ahx[R]-1	Evolved towards CPP$_{RXR}$-PNA #1 3/5 Ahx[R]-2	Evolved towards CPP$_{RXR}$-PNA #2 5/5 Ahx[R]-3
IS1				IG: $chbB$ - $osmE$		x	x	
IS5				IG: $gltF$ - $yhcA$		x	x	x
IS1				$rapA$		x	x	
IS1				$waaB$		x	x	x
IS1				$waaO$		x	x	x
			x	$yfaP_{V55G}$		x		
			x	$rpsL_{I82D}$		x	x	x
			x	$bdcA_{A53T}$		x		
			x	$cpxR_{L20Q}$				x
		x		Δrac			x	

Mutational profile of CPP$_{RXR}$-PNA[R] clones. [a] Mutation type; IS: IS insertion; IN: insertion; DEL: deletion; SNP: single-nucleotide polymorphism. [b] Mutated genomic region; IG: inter genic region between two genes;

766   Δ*rac*; excerted *rac* prophage(45). [c] Indicates which lineage the evolved clones where isolated from within

767   the ALE-experiment with CPP$_{RXR}$-PNA. [d] Frequency of the evolved clone within the lineage. See Table S1 for

768   mutation details and Table S2 for function of the mutated genes.

 Fig.1

774   **Fig. 2**

776   **Fig. 3.**

Fig. 4.

781   **Supplementary Materials**
782

783   **Table S1. Mutation type.**

Gene	Start (bp)	Mutation type	Mutation	Change
IG: *chbB* - *osmE*	1821764	IS1 insertion		
IG: *gltF* - *yhcA*	3362023	IS5 insertion		
rapA	63248	IS1 insertion		
waaB	3803557	IS1 insertion		
waaO	3802515	IS1 insertion		
yfaP$_{V55G}$	2327980	SNP	A $\rightarrow$ G	Valine-55-Glycine
rpsL$_{I82D}$	3474308	SNP	A $\rightarrow$ T	Isoleucine-82-Asparagine
bdcA$_{A53T}$	4473895	SNP	C $\rightarrow$ T	Alanine-53-Threonine
cpxR$_{L20Q}$	4105612	SNP	A $\rightarrow$ T	Leucine-20-Glutamine
Δrac	1411941	Deletion	23 kb del	Excerted *rac* prophage(45)

786   **Table S2. Function of mutated genes.**

Gene	Function	Reference
chbB	*N,N'*-diacetylchitobiose-specific phosphotransferase system enzyme IIB component	(46)
osmE	Osmotically-inducible lipoprotein, which is induced at the onset of stationary phase	(47)
gltF	Unknown	(18)
yhcA	Hypothetical protein, function unknown	
rapA	RapA enables recycling of stalled RNA polymerases and stimulates RNA synthesis *in vitro*	(48)
waaB	The *waa*-operon consists of *waaQGPSBOJYZU* whose products are involved in the assembly of the core region of the lipopolysaccharide molecule in the outer membrane of the cell envelope.	(49)
waaO		
yfaP	Unknown	
rpsL	The product of *rpsL* is the S12 protein; a component of the 30S subunit of the ribosome that plays a role in translational accuracy.	(20)
bdcA	BdcA is a c-di-GMP-binding biofilm dispersal mediator protein	(50)
cpxA	*cpxAR* encodes the two-component regulatory system CpxA/R, that consists of the transmembrane sensor kinase CpxA in conjunction with the cytoplasmic transcription factor CpxR.	(21)
cpxR		

789   **Table S3. Accumulated fitness gain mutations from the ALE-experiment to the media alone.**

Gene	Start (bp)	Mutation type	Mutation	Change
yiaD	3716565	SNP	C → T	Leucine-7-Phenylalanine
uspC	1979381	IS1 insertion		
uspC	1979385	IS1 insertion		
uspC	1979486	IS1 insertion		
uspC	1979489	IS1 insertion		
uspC	1979507	IS1 insertion		
uspC	1979510	IS1 insertion		
ccmA	2297378	IS1 insertion		
ccmA	2297389	IS1 insertion		
fimE	4542161	Inversion/duplication		
fimE	4542164	Inversion/duplication		
fimE	4542682	Inversion/duplication		
fimE	4542689	Inversion/duplication		
fimE	4542690	Inversion/duplication		
fimE	4542987	Inversion/duplication		
fimE	4542995	Inversion/duplication		

792 **Table S4.** *cpxR_{L20Q} and cpxA_{Δ16-17} does not provide tolerance to cationic antimicrobial peptides.*

Minimum inhibitory concentration to peptide-PNA/APIM (in μM) and AMPs (in μg/mL)					
AMP/strain	**Wild-type**	*cpxA_{Δ16-17}*	*cpxR_{L20Q}*	**ΔwaaBO**	**Ahx^R-3**
CPP$_{RXR}$-PNA	0.5	4	4	0.5	8
CPP$_{R11}$-APIM	1	2	2	1	N/A
Cecropin P1	32	32	32	16	16
Cap11	2	2	2	1	1
Cap18	2	2	2	1	1
Indolicidin	4	4	4	4	4
Apidaecin 1B	8	8	8	8	8
Colistin	0.125	0.125	0.125	0.0625	0.0625
Protamine	16	16	16	16	16

793 Table S4. MIC values for peptide-PNA (in μM) and AMPs (in μg/mL) for wild-type and indicated mutants.
794 MIC were determined using broth dilutions (see Methods) at 37°C (no shaking). Peptide-PNA sequences
795 are listed in Table 1.

798   **Table S5**. Bacterial strains.

Strain	Genotype	Abbreviated form	Reference/source
MG1655	F⁻λ⁻*rph-1*	Wild-type	(51)
ALO6524	Δ*cpxA*:kan[a]	Δ*cpxA*	This work
ALO6525	Δ*cpxR*:kan[a]	Δ*cpxR*	This work
ALO7548	Δ*degP*:kan[a]	Δ*degP*	This work
ALO7549	Δ*cpxP*:kan[a]	Δ*cpxP*	This work
ALO4576	Δ*atpA*:cat[a]	Δ*atpA*	(52)
ALO4577	Δ*atpB*:cat[a]	Δ*atpB*	(52)
ALO7180	$cpxR_{L20Q}$[a]	$cpxR_{L20Q}$	This work
ALO7550	$cpxR_{L20Q}$ Δ*cpxA*:kan[a]	$cpxR_{L20Q}$ Δ*cpxA*	This work
ALO7552	$cpxR_{L20Q}$ Δ*nuoABCEFGHIJKLMN*:cat[a]	$cpxR_{L20Q}$ Δ*nuo*	This work
ALO7553	$cpxR_{L20Q}$ Δ*cyoABCDE*:cat[a]	$cpxR_{L20Q}$ Δ*cyo*	This work
ALO7554	$cpxA_{Δ16-17}$[a]	$cpxA_{Δ16-17}$	This work
ALO7551	$cpxA_{Δ16-17}$ Δ*cpxR*:kan[a]	$cpxA_{Δ16-17}$ Δ*cpxR*	This work
ALO7555	Δ*nuoABCEFGHIJKLMN*:cat[a]	Δ*nuo*	This work
ALO7556	Δ*cyoABCDE*:cat[a]	Δ*cyo*	This work
ALO6972	Δ*waaBO*:cat[a]	Δ*waaBO*	This work

799   [a] Genotype otherwise as *E. coli* MG1655

Table S6. Primer sequences.

Primers for cloning	
Primer	**Sequence**
*nuo*_cat_Fw	ccgccgagcatgtcggcggctttctgacttacaaagtaacagagtgtaggctggagctgcttc
*nuo*_cat_Rv	catctgccgtgaagagcagtgaatctggcgctactttgatgagtaagcacatatgaatatcctccttag
*cyo*_cat_Fw	gctaagaaaaaggtgccgtagcacctttttaatagagagagagtttttggtgtaggctggagctgcttc
*cyo*_cat_Rv	tgcccacacactttaaacgccaccagatcccgtggaattgaggtcgttaacatatgaatatcctccttag
*waaB*_cat_Fw	ggaaatgagtccacaatgctacccttatatcattactttatagtttcccagttgtgtaggctggagctgcttc
*waaO*_cat_Rv	gacaaataaaaccgttaaaacagtgagttttactcaattaatgagagtttctcatatgaatatcctccttag
CpxR_*tet-sacB*_Fw	ggcgaagatgcgcgcggttaagctgcctatcatgaagcagaaaccatcagatcaaagggaaaactgtccatat
CpxR_*tet-sacB*_Rv	ttagcgacgtctgatgacgtaatttctgcctcggaggtatttaaacaatgtcctaattttttgttgacactcta
CpxR$_{L20Q}$_Fw	tgaatcgagcttgggtaaca
CpxR$_{L20Q}$_Rv	ggatggtgtctatggcaagg
Primers for qPCR	
Primer	**Sequence**
*cpxP*_qPCR_Fw	ggcatccgggtgaagaactt
*cpxP*_qPCR_Rv	aacttatgccgtcgaacatatgg
*nuoA*_qPCR_Fw	caccagataaaccagacctgcc
*nuoA*_qPCR_Rv	atctgttcgcatggtcaacctc
*cyoA*_qPCR_Fw	gactgcttcgcttttgcgac
*cyoA*_qPCR_Rv	aacgaacccggcacttatgac
*hctA*_qPCR_Fw	gctttttgccacgtattcgcc
*hctA*_qPCR_Rv	acttctctttgctgtcgccttc
*cysG*_qPCR_Fw	cgtcgattattgaccgctcacc
*cysG*_qPCR_Rv	cgaactgttgtttcactcgccc