2
3  **Steady-state visually evoked potentials and feature-based attention: Pre-**
4  **registered null results and a focused review of methodological considerations.**
5

6  Kirsten C.S. Adam[1,2], Lillian Chang[3], Nicole Rangan[1], & John T. Serences[1,2,4]
7
[1] Department of Psychology, *University of California San Diego*
[2] Institute for Neural Computation*, University of California San Diego*
[3] Department of Neuroscience, *Georgetown University Medical Center*
[4] Neurosciences Graduate Program, *University of California San Diego*

**Word count:** 226 (abstract), 1121 (introduction), 3318 (results), 1094 (discussion)
**Tables:** 0
**Figures:** 6
16

17  **Keywords:** attention; EEG; SSVEP; feature-based attention
18

19  **Contributions:** K.A. & J.S. planned the study. K.A., L.C., and N.R. collected data and
20  performed the literature review. K.A. performed analyses and drafted the manuscript. All
21  authors revised the manuscript.
22

23  **Funding:** Research was supported by National Eye Institute grant R01 EY025872
24  (J.S.). and National Institute of Mental Health grant 5T32-MH020002 (K.A.).

26  **Pre-registration information:** We published a pre-registered research plan on the
27  Open Science Framework prior to data collection (https://osf.io/kfg9h/).
28

29  **Data availability:** Data and analysis code will be made available online on the Open
30  Science Framework at https://osf.io/ew7dv/ upon acceptance for publication.
31

32  **Acknowledgements:** We thank Matteo d'Amico for additional assistance with data
33  collection and Kelvin Lam for help as lab manager. We thank Rosanne Rademaker and
34  Angus Chapman for sharing Psychtoolbox code that was adapted for use here.

36  **Conflicts of interest:** none
37

**Correspondence to:**
Kirsten C. S. Adam
University of California San Diego
9500 Gilman Drive, Mail Code: 0109
La Jolla, CA 92093-010
kadam@ucsd.edu

45                                    **Abstract**

Feature-based attention is the ability to selectively attend to a particular feature (e.g.,

attend to red but not green items while looking for the ketchup bottle in your refrigerator),

and steady-state visually evoked potentials (SSVEPs) measured from the human

electroencephalogram (EEG) signal have been used to track the neural deployment of

feature-based attention. Although many published studies suggest that we can use trial-

by-trial cues to enhance relevant feature information (i.e., greater SSVEP response to the

cued color), there is ongoing debate about whether participants may likewise use trial-by-

trial cues to ignore a particular feature. Here, we report the results of a pre-registered

study in which participants either were cued to attend or to ignore a color. Counter to prior

work, we found no attention-related modulation of the SSVEP response in either cue

condition. However, positive control analyses revealed that participants paid some degree

of attention to the cued color (i.e., we observed a greater P300 component to targets in

the attended versus the unattended color). In light of these unexpected null results, we

conducted a focused review of methodological considerations for studies of feature-based

attention using SSVEPs. In the review, we quantify potentially important stimulus

parameters that have been used in the past (e.g., stimulation frequency; trial counts) and

we discuss the potential importance of these and other task factors (e.g., feature-based

priming) for SSVEP studies.

## Introduction

Attending to a specific feature leads to systematic changes in the firing rates of neurons that encode the relevant feature space. For example, when looking for a ripe tomato, the firing rate of neurons tuned to red will be enhanced and the firing rate of neurons tuned to other colors will be suppressed (e.g. responses to green; Ipata et al., 2006; Martinez-Trujillo & Treue, 2004; Störmer & Alvarez, 2014). Although there is broad agreement that participants may learn to suppress irrelevant distractors with sufficient experience, there is disagreement about whether these behavioral suppression effects may be volitionally implemented on a trial-by-trial basis in response to an abstract cue (i.e., a "volitional account"), or if they instead are solely implemented via implicit or statistical learning mechanisms (i.e., a "priming-based" account). Consistent with a volitional account, some work has found that participants can learn to use a trial-by-trial cue to ignore a particular color (Arita et al., 2012; Conci et al., 2019; Moher & Egeth, 2012; Reeder et al., 2017; for a more nuanced discussion see: Van Moorselaar & Slagter, 2020). However, other work has found that a specific color needs to be repeated over many trials to be suppressed, consistent with the priming account (Cunningham & Egeth, 2016; Failing et al., 2019; Theeuwes, 2013; Vatterott & Vecera, 2012).

Although many studies have examined the effects of feature-based suppression on later selection-related event-related potential (ERP) markers such as the N2pc and Pd (Arita et al., 2012; Carlisle & Nitka, 2019; Donohue et al., 2018; Sawaki & Luck, 2010), a key open question is whether trial-by-trial cues to ignore a color modulate earlier stages of visual processing. Recent work by Reeder and colleagues (Reeder et al., 2017) hypothesized that cues about which feature to ignore (i.e., "negative cues") may down-regulate processing in visual cortex during the pre-stimulus period. Consistent with this hypothesis, they found that overall BOLD activity in early visual cortex was lower when participants were given a negative cue about the target color than when participants were given either a positive or neutral cue. One limitation of this study, however, is that the authors were unable to test whether these univariate effects were actually feature-specific (as opposed to task-general anticipation). However, evidence from ERP studies suggests

that feature-based attention modulates early visual processing in a feature-specific

manner, as indexed by the P1 component (Moher et al., 2014; W. Zhang & Luck, 2009).

Critically, however, these ERP studies used a design where the same target and

distractor colors were repeated over many trial events. Thus, it is not clear whether

feature-based attention can modulate early visual processing on a trial-by-trial basis, or if

modulation of early visual processing may only be achieved via inter-trial priming

(Theeuwes, 2013). To attempt to address this gap, we conducted a pre-registered

experiment in which we measured steady-state visually evoked potentials (SSVEPs)

while giving participants trial-by-trial cues to suppress feature information.

When visual input flickers continuously at a given frequency (e.g., one stimulus at 24 Hz

and another at 30 Hz), the visually evoked potential in the electroencephalogram (EEG)

signal is entrained to these "steady states" and time-frequency analyses can be used to

derive estimates of the strength of neural responses to each stimulus (Adrian & Matthews,

1934; Regan, 1977). The amplitude of the frequency-specific SSVEP response has been

shown to be modulated  by both spatial and feature-based attention (higher amplitude

when attended; Chen et al., 2003; Morgan et al., 1996; Müller et al., 1998, 2006; Pei et

al., 2002). When participants are cued on a trial-by-trial basis to attend to a particular

feature (e.g., color), the SSVEP amplitude is higher for the attended feature (Andersen et

al., 2008; Chen et al., 2003; Müller et al., 2006). Further, the time-course of the SSVEP

response to an attended color reveals an early enhancement followed by a suppressed

response to the irrelevant, non-attended color (Andersen & Müller, 2010; Forschack et

al., 2017).

Our primary manipulation was whether we cued participants to actively attend or to

actively ignore a color on a trial-by-trial basis. We planned to use this method to track

enhancement vs. suppression of the SSVEP response, and to test whether the time-

course of enhancement and suppression varies with cue type. In the "attend cue"

condition, participants were cued about the relevant color to attend. This condition was

expected to replicate prior work examining the time-course of feature-based attention

using SSVEPs (Andersen & Müller, 2010), whereby enhancement of the attended color

is followed by suppression of the ignored color. In the "ignore cue" condition, we instead

cued participants about which color to *ignore.* If participants can use a cue to *directly*

suppress a color on a trial-by-trial basis independent of target enhancement (i.e., a strong

version of a volitional suppression account), we predicted that the time-course of

enhancement vs. suppression of the SSVEP signal would be reduced or reversed (i.e.,

that suppression of the cued, to-be-ignored color may happen even *prior to* enhancement

of the other color). Alternatively, if participants recode the "ignore" cue to serve as an

indirect "attend" cue (e.g., "Since I'm cued to ignore blue, that means I should attend red";

Beck & Hollingworth, 2015; Becker et al., 2015), then we predicted that target

enhancement would always precede distractor suppression regardless of whether

participants were cued to attend or ignore a particular color.

To preview the results, we were unable to fully test our hypotheses about the time-course

of feature-based enhancement and suppression because we did not find evidence for an

overall attention effect with our task procedures. Despite robust SSVEP entrainment (d >

5), we observed no credible evidence that the SSVEP response was higher for an

attended versus unattended color in either cue condition. Positive control analyses

revealed that our lack of SSVEP effect was not due to a complete lack of attention to the

attended color: ERP responses (P3) to the targets were modulated by attention as

expected (Adamian et al., 2019; Andersen et al., 2013; Andersen, Fuchs, et al., 2011). In

light of our inconclusive results, we also performed a focused methodological review of

key potential task differences between our work and prior work that may have resulted in

our failure to detect the effect of feature-based attention on SSVEP amplitude. We

considered whether task factors such as stimulus flicker frequency, sample size, stimulus

duration, and stimulus color might have impacted our ability to observe an attention effect.

No single methodological factor that we considered neatly explains our lack of effect.

Given our results and literature review, we propose that future work is needed to

systematically explore two key factors: (1) variation in feature-based attention effects

across stimulus flicker frequencies and (2) the extent to which feature-based priming

modulates SSVEP attention effects.

<div align="center">

**Methods**

</div>

We published a pre-registered research plan on the Open Science Framework

prior to data collection (https://osf.io/kfg9h/). Our raw data and analysis code will be

made available online on the Open Science Framework at https://osf.io/ew7dv/ upon

acceptance for publication.

**Participants**

Healthy volunteers ($n = 32$; gender = 17 female, 15 male; mean age = 21.5 years

[SD = 3.84, min = 18, max = 39]; handedness not recorded; corrected-to-normal visual

acuity; normal color vision) participated in one 3.5 to 4 hour experimental session at the

University of California San Diego (UCSD) campus, and were compensated \$15/hr.

Procedures were approved by the UCSD Institutional Review Board, and all participants

provided written informed consent. A sample size of 24 was pre-registered, and exclusion

criteria were pre-registered (see section "EEG preprocessing" below for more details).

After running each participant, we checked whether the data were usable (i.e., sufficient

number of artifact-free trials) so that we would know when to stop data collection. To

reach our final sample size (n = 23 participants with usable data), we ran a total of 32

participants. Nine participants' data were not used for the following reason: Subjects with

an error in the task code (n = 3), subjects who stopped the study early due to technical

issues or to participants' preferences (n = 4), subjects with too many artifacts (n = 2).

Note, we were one subject short of our pre-registered target sample size of 24 because

data collection was suspended due to COVID-19. However, as our later power analyses

will show, we do not believe the addition of 1 further subject would have meaningfully

altered our conclusions.

**Stimuli and Procedures**

**Heterochromatic flicker photometry task.** We chose perceptually equiluminant colors

for each participant using a heterochromatic flicker photometry task. Participants viewed

a large circular, flickering stimulus (8° radius) on a black screen (0.08 cd/m2). We

generated 5 circular color spaces in CIE LAB-space with varying luminance (circles

centered on: L = 35-65, a = 0, b = 0; 5 colors equally spaced around circle with radius =

35) for use in the task. Participants matched each of the 5 colors to a medium-gray

reference color (RGB = 105.6 105.6 105.6).

On each trial, the circular background was flickered between two different colors.

One color was always medium-gray, and the other color was the to-be-matched color on

that trial. The colors of circular background were phase reversed at a rate of 24 Hz, giving

the appearance of a fast flicker when the subjective luminance values were not matched.

On top of the flickering circular stimulus small oriented bars were drawn in the medium-

gray reference color (the bars changed locations at a rate of 1Hz). The oriented bars

served no purpose other than subjectively making it easier to discriminate fine-grained

differences in luminance between the flickering colors (i.e., these bars gave secondary

visual cues about equiluminance via the "minimally distinct border" phenomenon, Kaiser,

1988). Participants increased or decreased the luminance of the to-be-matched color

(using up and down arrow keys) until the amount of perceived flicker was minimized – the

point of perceptual equiluminance. The luminance starting value of the to-be-matched

color was chosen at random on each trial. Once satisfied with their response, the

participant pressed spacebar to continue to the next trial. Each to-be-matched color was

repeated 3 times (15 trials total).

**Feature-based attention task.** All stimuli were viewed on a luminance calibrated

CRT monitor (1024 x 768 resolution, 120 Hz refresh rate) from a distance of ~50 cm in a

dimly lit room. Stimuli were generated using Matlab 2016a and the Psychophysics toolbox

(Brainard, 1997; Kleiner et al., 2007; Pelli, 1997). Participants rested their chin on a chin-

rest and fixated a central dot (0.15º radius) throughout the experiment. The stimulus was

a circular aperture (radius = ~9.5º) filled with 120 oriented bars (each bar ~1.1º long and

~.1º wide). Bars were centered on a grid and separated by ~1 bar length such that they

never overlapped with one another. On each individual frame (~8.33 ms) this grid was

randomly phase shifted (0:2π in x and y coordinates) and rotated (1:360º), thus giving the

appearance of random flicker. To achieve Steady State Visually Evoked Potentials

(SSVEP) half of the bars flickered at 24 Hz (3 frames on, 2 frames off) and the other half

flickered at 30 Hz (2 frames on, 2 frames off; Figure S1).

On each trial (Figure 1), the participants viewed the stimulus array of flickering,

randomly oriented bars presented on a black background (0.08 cd/m2). Half of these bars

were shown in one color (randomly chosen from the 5 possible colors) and the other half

were in another randomly chosen color (with the constraint that the two sets of bars must

be two different colors). During an initial baseline (1,333 ms), participants viewed the

flickering dots while they did not yet know which color to attend; during this baseline, the

fixation point was a medium gray color (same as the reference color in the flicker

photometry task). After the baseline, the fixation dot changed color, cuing the participants

about which color to attend or ignore. In the "attend cue" condition, the color of the fixation

point indicated which color should be attended. In the "ignore cue" condition, the color of

the fixation point indicated which color should be ignored. These two conditions were

blocked, and the order was counterbalanced across participants (further details below).

During the stimulus presentation (2,000 ms), participants monitored the relevant color for

a brief "target event" (333 ms). During this brief target event, a percentage of lines in the

relevant color will be coherent (iso-oriented). Critically, the orientation of each coherent

target or distractor event was completely unpredictable (randomly chosen between 1-180

degrees); thus, participants could not attend to a particular orientation in advance in order

to perform the task. A target event occurred on 50% of trials, and participants were

instructed to press the spacebar as quickly as possible if they detected a target event.

Importantly, physically identical events (iso-oriented lines in a random orientation, 333

ms) could also appear in the distractor color (50% of trials). Participants were instructed

that they should only respond to target events; if they erroneously responded to the

distractor event, the trial was scored as incorrect. The target and/or distractor events

could begin as early as cue onset (0 ms) and no later than 1,667 ms after stimulus onset).

Participants could make responses up to 1 second into the inter-trial interval. If both a

target and distractor event were present, their onset times were separated by at least 333

ms.

**Figure 1. Feature-based attention task.** (A) Trial events in an example 'attend cue' trial where there is a target event. Note, this figure is a schematic and the stimuli are not drawn to scale. After a baseline period, participants were cued to attend or ignore one color via a change to the fixation point color. If participants noticed a target event (~75% iso-oriented lines in the to-be-attended color), they pressed the space bar. (B) Five colors were used, and these 5 colors appeared with equal probability. Thus, the target and distractor colors could be either 72 degrees or 144 degrees apart on a color wheel. This figure shows all possible color pairings if red was the target color. (C) Examples of cues, target events, and distractor events in the 2 main conditions. In the 'attend cue' condition, participants made a response when the iso-oriented lines were the same color as the cue (target event) and did not respond if the iso-oriented lines occur on the uncued color (distractor event). In the 'ignore cue' condition, participants made a response when the iso-oriented lines occurred on the uncued color (target event), and they did not respond if the iso-oriented lines occurred on the cued color (distractor event). Note, all lines were of equal size in the real experiment; lines are shown at different widths here for easier visualization of the target and distractor colors. Here, the iso-oriented lines are drawn at vertical in all 4 examples. In the actual task, the iso-oriented lines could be any orientation (1-180).

To ensure that the task was effortful for participants, the coherence of the lines in the target stimulus was adapted at the end of each block if behavior was outside the range of 70 - 85% correct. At the beginning of the session, the target had 50% coherent iso-oriented lines. If accuracy over the block of 80 trials was >85%, coherency decreased by 5%. If block accuracy was <70%, coherency increased by 5%. The maximum allowed coherence was 80% iso-oriented lines (so that participants would not be able to simply

individuate and attend a single position to perform the task) and the minimum allowed

coherence was 5%. The presence and absence of target and distractor events was

balanced within each block yielding a total of 4 sub-conditions within each cue type (25%

each): (1) target event + no distractor event, T1D0 (2) no target event + distractor event,

T0D1 (3) target event + distractor event, T1D1 (4) no target event + no distractor event,

T0D0.

Participants completed both task conditions (attend cue and ignore cue). The two

conditions were blocked and counterbalanced within a session (i.e., half of participants

performed the "attend cue" task for the first half of the session and the "ignore cue" task

for the second half of the session.) Each block of 80 trials took approximately 6 min 50

sec. Participants completed 18 blocks (9 per condition) for a total of 720 trials per cue

condition. Note, we originally planned for 20 blocks (10 per condition) in the pre-

registration, but the block number was reduced to 18 after the first few participants did

not finish all blocks.

**Summary of deviations from the registered procedures.** As described in-line

above, there were some minor deviations from the pre-registration: (1) We made changes

to the pre-registered task code to fix errors that we discovered while running the first 3

subjects (e.g., incorrect cues and behavioral feedback in the 'ignore cue' condition). (2)

We included code for eye-tracking, which allowed us to give participants automated real-

time feedback if they blinked when they were not supposed to, i.e., during the stimulus

period. (3) We reduced the total number of experimental blocks from 20 (10 per cue

condition) to 18 (9 per cue condition) due to time constraints. (4) We had to prematurely

stop data collection at n = 23 out of 24 due to COVID-19. (5) Due to unanticipated failure

to detect an overall attention effect, we performed additional non-pre-registered control

analyses to attempt to rule out possible explanations of this null effect (see section: "Non

pre-registered control analyses" below).

**EEG pre-processing**

Continuous EEG data were collected online from 64 Ag/AgCl active electrodes

mounted in an elastic cap using a BioSemi ActiveTwo amplifier (Cortech Solutions,

Wilmington, NC). Eight external electrodes were placed on the left and right mastoids,

above and below each eye (vertical EOG), and lateral to each eye (horizontal EOG).

Continuous gaze-position data were collected from an SR Eyelink 1000+ eye-tracker

(sampling rate: 1,000 Hz; SR Research, Ottawa, Ontario). We also measured stimulus

timing with a photodiode affixed to the upper left-hand corner of the monitor (a white dot

flickered at the to-be-attended color's frequency; the photodiode and this corner of the

screen were covered with opaque black tape to ensure it was not visible). Data were

saved unreferenced and unfiltered, then re-referenced offline to the algebraic average of

the left and right mastoids, low-pass filtered (<80 Hz) and high-pass filtered (>.01 Hz).

Artifacts were detected using automatic criteria described below, and the data were

visually inspected to confirm that the artifact rejection criteria worked as expected. We

excluded subjects with fewer than 400 trials remaining per cue condition.

**Eye movements and blinks.** We used the eye-tracking data and the

HEOG/VEOG traces to detect blinks and eye movements. Blinks were detected on-line

during the task using the eye tracker. If a blink was detected (i.e., missing gaze position

returned from the eye tracker), the trial was immediately terminated and the participant

was given feedback that they had blinked (i.e., the word "blink" was written in white text

in the center of the screen). If eye-tracking data could not be successfully collected (e.g.,

calibration issues), the VEOG trace was used to detect blinks and/or eye movements

during offline artifact rejection. To do so, we used a split-half sliding window step function

(Luck, 2005; window size = 150 ms, step size = 10 ms, threshold = 30 microvolts.) We

also used a split-half sliding-window step function to check for eye-movements in the

gaze-coordinate data from the eye-tracker (window size = 80 ms, step size = 10 ms,

threshold = 1º) and in the horizontal electrooculogram (HEOG), window size = 150 ms,

step size = 10 ms, threshold = 30 microvolts, and to detect blinks and/or eye movements

in the vertical electrooculogram (VEOG),

**Drift, muscle artifacts, and blocking:** We checked for drift (e.g., skin potentials)

by comparing the absolute change in voltage from the first quarter of the trial to the last

quarter of the trial. If the change in voltage exceeded 200 microvolts, the trial was rejected

for drift. In addition to slow drift, we also checked for sudden, step-like changes in voltage

with a sliding window (window size = 250 ms, step size = 20 ms, threshold = 200

microvolts). We excluded trials for muscle artifacts if any electrode had peak-to-peak

amplitude greater than 200 microvolts within a 15 ms time window (step size = 10 ms).

We excluded trials for blocking if any electrode had ~120 ms during which all values within

1 microvolt of each other (sliding 200 ms window, step size = 50 ms).

**Pre-registered SSVEP analyses.**

**General method for SSVEP quantification.** We planned to quantify SSVEP

entrainment by filtering the data with a Gaussian wavelet function. First, we calculated an

average ERP for each condition at electrodes of interest (O1, Oz, O2). We chose these

3 electrodes based on large SSVEP modulations at these sites in prior work (e.g.,

Itthipuripat et al., 2013; Müller et al., 2006). After calculating an ERP for each condition,

we filtered the data with a Gaussian wavelet function with .1 fractional bandwidth to obtain

frequency-domain coefficients from 20 to 35 Hz in 1-Hz steps. For a similar analytic

approach see: (Canolty et al., 2006; Itthipuripat et al., 2013; Rungratsameetaweemana

et al., 2018). Signal-to-noise ratio for each SSVEP frequency was calculated as the power

at a given frequency divided by the average power of the 2 adjacent frequencies on each

side. For example, SNR of 24 Hz would be calculated as the power at 24 Hz divided by

the average power at 22, 23, 25, and 26 Hz. We also pre-registered an analysis plan for

examining the time-course of SSVEP amplitude. However, because our data failed to

satisfy pre-registered pre-requisite analyses, we do not report these time-course effects

here (for completeness, we show the time-course of SNR in Figure S2).

**Checking that an SSVEP is elicited at the expected frequencies before**

**collecting the full sample.** At n = 5, we planned to confirm that our task procedure

successfully produced reliable SSVEP responses (i.e., check that we observed peaks at

the correct stimulus flicker frequencies). If our task procedures failed to elicit an SSVEP

at the expected frequencies, we had planned to stop data collection and alter the task to

troubleshoot the problem (e.g., optimize timing, choose different flicker frequencies, make

stimuli brighter, etc.). We planned to begin data collection over again if we failed this

trouble-shooting step. Note, at this early stage we only verified if the basic method worked

(SSVEP frequencies were robustly entrained): we did not test whether any hypothesized

attention effects were present, as this could inflate our false discovery rate (Kravitz &

Mitroff, 2017).

**Checking achieved power for the basic attention effect.** Without adequate

power for the basic attention effect, we would not be able to robustly interpret the time-

course of enhancement vs. suppression. At the full sample size, we thus planned to check

whether we had sufficient achieved power for the overall attention effect (>=80% power

for attended vs. ignored color collapsed across the entire stimulus period) as a

prerequisite for interpreting the time-course of enhancement and suppression.

**Checking if a priori electrodes are reasonable.** We chose a priori to analyze the

SSVEP at electrodes O1, Oz, and O2 (Itthipuripat et al., 2013; Rungratsameetaweemana

et al., 2018), but we planned to plot the topography of SSVEP modulation across all

electrodes to check that these a priori electrodes were responsive to the SSVEP

manipulation. If these electrodes were not responsive to the SSVEP, we planned to

perform a cluster-based permutation test to select a new set of electrodes.

**Checking if target and/or distractor presence alters results.** Our core analyses

planned to use all trials for each condition (e.g., target event present or absent, distractor

event present or absent). To confirm that the act of making a response did not

contaminate the SSVEP results, we planned to compare SSVEPs for each of the 4 sub-

conditions (the 4 possible combinations of target present/absent and distractor

present/absent). We predicted that the main SSVEP attention effect (entire stimulus

period) would not be different across these 4 conditions. But, if we found an effect of

target or distractor presence on the main SSVEP attention effect, then we planned to use

only the trials without target or distractor events for the time-course analyses (as has been

done in prior work, e.g., Andersen & Müller, 2010).

**Checking if the distance between the target and distractor color alters**

**results.** We planned to test whether the magnitude and time-course of attentional

selection differs as a function of target-distractor color similarity (~72° vs. ~144°

separation between the attended and ignored color). Prior work found that it was more

difficult to simultaneously attend opposite colors (180° apart) than to simultaneously

attend two moderately-spaced colors (60º apart) (Störmer & Alvarez, 2014). Given this

result, we predicted that it should be easier to suppress a color that is drastically different

from the target color (and behavioral accuracy should likewise be higher for the 144º

condition).

**Additional non pre-registered SSVEP and ERP control analyses.**

405            We did not anticipate our failure to find an overall attention effect with these task

406    procedures and set of pre-registered "sanity check" analyses described above. To further

407    understand the lack of SSVEP attention effect, we performed additional non-pre-

408    registered control analyses.

409            **Positive control: Frequency analysis of the photodiode voltage.** During the

recording, a photodiode was used to ensure that the flicker frequencies were faithfully

presented, and the photodiode recorded voltage fluctuations induced by a small white dot

that flickered at the to-be-attended target frequency on each trial. The electrical activity

from the photodiode was recorded as an additional "electrode" in the data matrix (with the

structure: trials x electrodes x timepoints*).* Thus, we performed a fast Fourier transform

(FFT, Matlab function "fft.m") to ensure that the trial indexing and FFT aspects of our

analysis were correct. If these aspects of the analysis were correct, we should expect a

near-perfect modulation of photodiode FFT amplitude by attention condition as only the

flicker frequency of the attended stimulus was tagged. For 1 subject, the photodiode was

not plugged in (leaving 22 subjects for this analysis).

420            **Analysis control: Using a more similar frequency analysis procedure to prior**

**published work.** Because we were interested in characterizing a time-course effect, we

chose to use a Gaussian wavelet procedure to quantify power for each frequency of

interest. However, given the lack of overall attention effect, we were not able to

meaningfully look at the time-course effects. Thus, we additionally used a fast Fourier

transform (FFT) to assess overall entrainment during the entire stimulus period. This

method is more commonly used in prior published work (e.g., Andersen et al., 2008;

Andersen & Müller, 2010). Following prior work, we used the entire stimulus epoch

starting from 500 ms onward (500 ms – 2000 ms), we detrended the data, and we zero-

padded this time window (2,048 points) to precisely estimate our frequencies.

**Positive control: Analysis of event-related potential (P3) for an attention**

**effect.** Prior work has found that attention-related SSVEP modulations are accompanied

by changes to event-related potentials (ERPs) associated with target selection and

processing. To measure the P3, we calculated event-related potentials time-locked to the

target or distractor onset (baselined to -200 ms to 0 ms relative to target onset). We

included trials where there was only one target or distractor event on that trial, to avoid

the possibility of overlap between the two signals. We calculated P3 voltage at electrodes

Pz and POz during the time window 450-700 ms after target onset, similar to prior work

(Adamian et al., 2019; Andersen et al., 2013; Andersen, Fuchs, et al., 2011).

<div align="center">**Results**</div>

**Behavior**

Participants were overall accurate at the task (percent correct = 65.7%, d-prime =

1.25), and were significantly above chance (percent correct >50%, $p < .001$; d-prime >1,

$p = .006$). There was no significant effect of cue condition (attend cue versus ignore cue)

on performance, $p = .85$. Average percent correct was lower than our pre-specified target

range of 70-85%, meaning that most participants saw targets and distractors that were

maximally coherent (80% iso-oriented lines) for the majority of blocks (mean coherence

of the target/distractor events = 73.7%, SD = 5.08%). Although overall accuracy was

slightly out of the range we had expected when planning the study, participants were well

above chance and they saw targets with coherence values typical of prior work (Andersen

et al., 2008; Andersen, Fuchs, et al., 2011; Andersen & Müller, 2010).

**Pre-registered SSVEP results**

We first confirmed that our SSVEP procedure was effective at eliciting robust,

frequency-specific modulations of the EEG signal. After collecting the first 5 participants,

we checked that overall entrainment to our two target frequencies (24 and 30 Hz) was

robust when collapsed across conditions (Fig 2A) before proceeding with data collection.

We indeed found that entrainment was robust during the stimulus period even with n=5

for both the 24 Hz frequency (mean SNR = 4.45, SD = .14, $p < .001$) and for the 30 Hz

frequency (mean SNR = 2.97, SD = .15, $p < .001$). These values were similar for the full

n=23 sample (Fig 2B). To compute an effect size, we compared SNR values for each

target frequency (24 Hz and 30 Hz) to SNR for a baseline frequency (15 Hz). SNR values

for the target frequencies were significantly higher than baseline, Cohen's d = 5.3 and

6.32, respectively. As planned, we also confirmed that the electrodes we selected *a priori*

(O1, Oz, and O2) were reasonable given the topography of overall SSVEP entrainment

(i.e., they fell approximately centrally within the brightest portion of the heat map; Figure

2C-D).

**Figure 2. Overall SSVEP entrainment at the expected frequencies (collapsed across**

**all experimental conditions).** (A) Power as a function of frequency during the stimulus

period at electrodes O1, O2, and Oz for the first 5 participants. As expected, we observed

robust peaks at the stimulated frequencies (24 and 30 Hz). (B) Power as a function of

frequency for the stimulus period for the full sample (n=23). (C-D) Topography of signal

to noise ratio values for 24 Hz (C) and 30 Hz (D) for all participants collapsed across all

experimental conditions. Color scale indicates SNR. As expected, the *a priori* electrodes

O1, O2, and Oz (magenta circles) showed robust SNR during the stimulus period.

Next, we checked for a basic attention effect, defined as a larger amplitude

response evoked by the attended frequency compared to the ignored frequency). Note,

for the sake of clarity, all conditions are translated into "attend" terminology. That is, if a

participant was cued to "ignore blue" (24 Hz) during the "ignore cue" condition (and the

other color was red and 30 Hz), this will instead be plotted as "attend red" (30 Hz). Figure

3 shows the Gaussian wavelet-derived frequency spectra during the stimulus period (500-

2000 ms) as a function of cue type (attend versus ignore) and attended frequency (attend

24 Hz or attend 30 Hz). We found a main effect of measured frequency, whereby SNR

was overall higher for 24 versus 30 Hz, $F(1,22) = 57.89$, $p < .001$, $\eta^2_p = .73$. However, we

found no main effect of attended frequency ($p = .27$) or cue type ($p = .83$), and we found

no significant interactions ($p >= .18$). Collapsed down to a paired t-test, the effect size for

attended versus unattended SNR values was Cohen's $d = .03$. To detect an effect of this

size with 80% power (1-$\beta$ = .8; $\alpha$ = .05) would require a sample size $n > 7,000^*$.  Given

that we did not find an overall attention effect, we did not analyze or interpret analysis of

the SSVEP time-course. However, for completeness we have shown the time course in

Figure S2.

---

*As we did not pre-register Bayesian analysis choices (e.g., choices about priors, etc.), we did not calculate a Bayes Factor for this pre-registered analysis. However, the post-hoc power analysis gives a sense of the degree to which this is a null effect.

**Figure 3. Overall attention effect in the attend cue and ignore cue conditions.** (A-B) Frequency spectra in the attend cue (A) and ignore cue (B) conditions during the stimulus period. Although we observe expected peaks at 24 Hz and 30 Hz, this SSVEP response is not modulated by the attention manipulation. (C-D). Violin plots of the signal-to-noise ratio at the SSVEP frequencies in the attend cue (C) and ignore cue (D) conditions.

Although we pre-registered that we would analyze all trials (those with and without target/distractor events), most prior studies have included only trials without any target or distractor events in the main SSVEP analysis (e.g., Andersen et al., 2008; Müller et al., 2006). To ensure that our null result was not due to this analysis choice, we also planned in our pre-registration to examine the SSVEP attention effect for trials with and without target and distractor events. When restricting our analysis to only trials without targets or distractors (25% of the 1440 trials, or 360 trials total before artifact rejection), we likewise found no attention effect. As before, we found a main effect of measured frequency (24 > 30 Hz), $p < .001$, but no effect of cue condition ($p = .053$) or attended frequency ($p = .073$), and, most critically, we found no interaction between measured frequency and attended

514    frequency ($p$ = .33). Frequency spectra for all combinations of target and distractor

515    presence are shown in Figures S3 and S4.

516            Finally, we also pre-registered that we would check whether the similarity of the

target and distractor colors (72 versus 144 degrees apart on a circular color wheel; Figure

1B) would modulate the SSVEP attention effect. We likewise found that the similarity of

the distractor colors did not significantly modulate the SSVEP response, and we found no

attention effect (interaction of measured frequency and attended frequency) in either color

distance condition ($p$ >= .26; Figure S5).

**Non-pre-registered control analyses**

524            We conducted additional control analyses to rule out possible sources of failure to

find an attention effect. First, we examined the photodiode recording to rule out any

failures due to trial indexing. The photodiode measured the luminance of a white dot that

flickered at the attended frequency on each trial. As expected, performing an FFT on the

photodiode time-course thus yielded near-perfect tracking of the attended frequency

(Figure 4A-B, $p$ < .001). On the other hand, we again found null results for the main

attention manipulation (Figure 4C-F) when using an FFT analysis that more closely

followed prior work. We ran a repeated measures ANOVA on the signal to noise ratio

values during the stimulus period, including the factors Measured Frequency (24 Hz, 30

Hz), Attended Frequency (24 Hz, 30 Hz), and Cue Type (Attend, Ignore). We found no

main effect of measured frequency ($p$ = .91), attended frequency ($p$ = .45), or cue type ($p$

= .54), and we found no significant interactions ($p$ >= .38). However, the average signal-

to-noise ratio of the entrained frequencies was overall robust (M = 4.45, SD = 1.45,

greater than chance value of 1: $p < 1 \times 10^{-9}$), so our inability to observe the attention effect

was not due to lack of overall SSVEP entrainment.

**Figure 4. FFT analysis of the photodiode and SSVEPs at attended and unattended frequencies during the stimulus period (500-2000 ms).** (A-B) As a positive control for our analysis pipeline, we performed an FFT analysis on the photodiode trace. The photodiode recorded a flickering white dot at the attended frequency on each trial. As expected, this provides a near-perfect tracking of the attended frequency in both the attend cue condition (A) and the ignore cue condition (B). (C-F) To ensure our null effect was not due to using Gaussian wavelets rather than an FFT, we repeated the main analysis with an FFT. Frequency spectra for the attend cue condition (C) and ignore cue condition (D) reveal overall robust entrainment at 24 Hz and 30 Hz, but no modulation by attention. Likewise, violin plots of signal-to-noise ratios again show robust signal but no modulation by attention in either the attend cue condition (E) or the ignore cue condition (F).

**Positive control: Analysis of event-related potential (P3b) for an attention**

**effect.** Consistent with prior work, we found a significantly larger P3 component for target

onsets compared to distractor onsets (Figure 5). A repeated measures ANOVA with

within-subjects factors cue type (attend cue or ignore cue) and event type (target or

distractor onset) revealed a robust main effect of event type (target > distractor), $F(1,22)$

$= 51.64$, $p < 1 \times 10^{-5}$, $\eta^2_p = .70$, and a main event of cue type (attend > ignore), $F(1,22) =$

$4.96$, $p = .037$, $\eta^2_p = .18$, but no interaction between event type and cue type ($p = .65$).

Control analyses confirmed this P3 modulation was not due to differential rates of making

a motor response for targets and distractors (Figure S6). The main effect of event type

(target > distractor) remained when analyzing only trials where participants made a motor

response ($p < .001$). Thus, the P3 was overall larger for target than distractor events,

consistent with prior work that found this ERP attention effect alongside an SSVEP

attention effect.

**Figure 5. P3 amplitude at electrodes Pz and POz.** (A) P3 amplitude in the attend cue
condition, as a function of whether the event onset (iso-oriented lines) was a target that
should be reported or a distractor that should be ignored. (B) P3 amplitude in the ignore
cue condition. Shaded error bars represent standard error of the mean; the gray rectangle
indicates the time period used for the statistical tests.

573
574
575
576

**Focused review of feature-based attention studies using SSVEPs**

Given that our results are inconsistent with prior work, we conducted a focused review to

try to pinpoint critical methodological differences that may have led to our failure to

replicate a basic attention effect on SSVEP amplitude in this specific task. To do so, we

first read review papers to identify an initial set of empirical studies employing a feature-

based attention manipulation and SSVEPs (Andersen, Müller, et al., 2011; Norcia et al.,

2015; Vialatte et al., 2010). From this initial set of papers, we used Google Scholar to

check citations and citing papers for mention of the terms feature-based attention and

SSVEPs. Our inclusion criteria included: (1) published journal article (2) healthy young

adults (3) SSVEPs were measured from either an EEG or MEG signal and (4) a feature-

based attention manipulation was included.

We defined "feature-based attention manipulation" as having the following characteristics:

(1) Participants were cued to attend a feature(s) within a feature dimension (e.g., attend

red, ignore blue) rather than across a feature dimension (e.g. attend contrast, ignore

orientation), (2) The attended and ignored feature were both frequency-tagged in the

same trials (rather than only 1 feature tagged per trial), (3) Each frequency was both

"attended" and "ignored" on different trials, so that the amplitude of a given frequency

could be examined as a function of attention, (4) The task could not be performed by

adopting a strategy of splitting spatial attention to separate spatial locations.

After applying these screening criteria, some of the studies that we initially identified were

excluded (brackets indicate exclusion reason(s)): Appelbaum & Norcia, 2009 [1,4];

Boylan et al., 2019 [3]; Bridwell & Srinivasan, 2012 [2,3]; Clementz et al., 2008 [3]; Garcia

et al., 2013 [1,4]; Hasan et al., 2017 [2]; Itthipuripat et al., 2019 [1]; Talsma et al., 2006

[1,4]; Thigpen et al., 2019 [4]; Verghese et al., 2012 [1,4]).

We identified a total of 34 experiments from 28 unique papers (Tables S1-S4) meeting

our inclusion criteria. From these experiments, we quantified variables such as the

number of subjects, number of trials, frequencies used, and the presence or absence of

an attention effect in the expected direction (attended > ignored). If more than one group

of participants was used (e.g., an older adults group) then we included the study but only

quantified results for the healthy young adult group (Quigley et al., 2010; Quigley & Müller,

2014).

**Task used in each study.**

The tasks used in these studies fell broadly into one of 4 categories: (1) a competing

gratings task, (2) a whole-field flicker task, (3) a hemifield flicker task and (4) a central

task with peripheral flicker.

In the competing gratings task (Table S1), participants viewed a stream of centrally-

presented, superimposed gratings (e.g., a red horizontal grating and a green vertical

grating). Because colored, oriented gratings were typically used, participants could thus

generally choose to attend based on either one or both features (color and/or orientation).

Each grating flickered at its own frequency (e.g. green grating shown at 7.41 Hz, red

grating shown at 8.33 Hz, as in Chen et al., 2003). Because the gratings were

superimposed, on any given frame only one of the two gratings was shown. On frames

where both gratings should be presented according to their flicker frequencies, a hybrid

"plaid" stimulus was shown. Studies using a competing gratings task include: (Allison et

al., 2008; Chen et al., 2003; Keitel & Müller, 2016; J. Wang et al., 2007).

In the whole-field flicker task (Table S2), participants viewed a spatially global stimulus

comprised of small, intermingled dots or lines. Typically, half of the dots or lines were

presented in one feature (e.g., red) and the other half of the lines were presented in

another (e.g., blue); each set of dots flickered at a unique frequency. Although the most

common attended feature was color, some task variants included (1) attending high or

low contrast stimuli (2) attending a particular orientation or (3) attending a particular

conjunction of color and orientation. The whole-field flicker task was the most common

task variant, and it is also most similar to the task performed here. Studies using a whole-

field flicker task include: (Andersen et al., 2008, 2009, 2012, 2015; Andersen & Müller,

2010; Forschack et al., 2017; Martinovic et al., 2018; Martinovic & Andersen, 2018; Müller

et al., 2006; Quigley et al., 2010; Quigley & Müller, 2014; Steinhauser & Andersen, 2019;

D. Zhang et al., 2010)

In the hemifield flicker task, participants viewed a stimulus within each hemifield, and each

stimulus was comprised of small, intermingled dots or lines. Often, these studies included

both a feature-based attention manipulation and a spatial attention manipulation (e.g.,

attend red on the left-hand side). However, the feature-based attention task could not be

achieved with spatial attention alone, as participants needed to attend to a particular color

and ignore a distractor color within the attended hemifield. In addition, the unattended

hemifield could often be used to track the spatially global spread of feature-based

attention. Studies using a hemi-field flicker task include: (Adamian et al., 2019; Andersen

et al., 2013; Andersen, Fuchs, et al., 2011; Müller et al., 2018; Störmer & Alvarez, 2014)

Finally, in the central task with peripheral flicker, participants performed a task near

fixation (e.g., visual search), and feature-based attention was measured indirectly via a

peripheral flickering stimulus (i.e., this design took advantage of the spatially global

spread of feature-based attention). Studies using a central task with peripheral flicker

include: (Chu & D'Zmura, 2019; Jiang et al., 2017; Painter et al., 2014, 2015).

**Sample Size, Trial Counts, and Stimulus Duration**

First, we examined whether insufficient power (e.g., fewer subjects and/or trials relative

to prior work) could have led to our failure to detect an attention effect. The number of

studies employing each task variant is plotted in Figure 6A, the number of subjects per

experiment is plotted in Figure 6B, the number of trials per experiment is plotted in Figure

6C, and stimulus duration is plotted in Figure 6D. Bars are color coded according to

whether each experiment overall found an expected attention effect (attended > ignored),

a reverse attention effect (ignored > attended), mixed results across conditions within the

experiment, or ambiguous results (3 studies: Pei et al., 2002: reported statistics for the

harmonics but not the fundamental frequency; Allison et al., 2008: missing formal

statistical tests; Martinovic et al., 2018: statistical tests measured if attention effect

differed between conditions, but did not formally test that the attention effect was overall

significant). Our study had N = 23, 1,440 total trials per subject (720 trials per cue

condition), and a stimulus duration of 2 seconds. In comparison to the literature, these

factors are unlikely to explain our failure to find an effect. On average, prior studies had

a median sample size of N = 15 (SD = 4.1, min = 9, max = 23) and a median trial count

of 440 (SD = 314.2, min = 8, max =1600) and a median stimulus duration of 4.1 sec (SD

= 37.35, min = 1 sec, max = 120 sec). Likewise, our trial counts were reasonable when

we restricted our analysis to trials without targets or distractors (360 trials total; 180 trials

per condition) relative to this estimated value for the literature (median = 96 trials per

condition, SD = 95.6, min = 4, max = 400). Upon initially plotting the data, we noticed that

studies using the competing gratings task produced inconsistent results, with as many

experiments showing a mixture of effects across conditions, overall reversed effects

(ignored > attended), and null effects (2 studies in each category). We think this

inconsistency is most likely due to the below-average trial counts for these studies

(median of only ~8 trials) (Boudewyn et al., 2018).

**Figure 6. Study characteristics from the literature review of feature-based attention and SSVEPs.** (a) Number of experiments in each of the 4 task variants. Different colors in the stacked bar graphs indicate whether the experiment found an expected attention effect (attended > ignored), reverse effect (ignored > attended), mixed effects across conditions, or a null / ambiguously reported effect. (b) Number of participants. (c) Number of trials in the experiment (before artifact rejection or excluding trials with targets). (d) Stimulus duration. (e) Percentage of trials, on average, where the attended feature was repeated on the next trial. (f) Stimulation frequencies.

**Percentage of trials where an attended feature was repeated.**

Next, we examined the percentage of trials where the attended feature was repeated. The priming-based account of feature-based attention posits that participants cannot use trial-by-trial cues to enhance a particular feature, but rather, feature-based enhancement happens automatically when a particular feature is repeated (Theeuwes, 2013). Thus, if there is a substantial proportion of trials where the repeated color was attended (e.g. with 2 possible colors, both the attended and ignored color will be repeated on 50% of trials), then the observed attentional enhancement effects might be driven primarily by incidental repetitions of attended features. In our study, we used 5 different colors to reduce the potential effect of inter-trial priming on the observed SSVEP attention effects (20% repeats of the attended color, 4% repeats of the attended color and the ignored color).

We quantified the approximate percentage of trials on which an attended feature on one

trial is repeated on the next trial (within a given block of trials). In some studies,

participants were cued to attend more than one feature on a given trial, or they sometimes

attended to a conjunction of features. In these cases, we calculated the expected number

of repeats for either of the 2 attended features based on the total number of conditions

(Andersen et al., 2008, 2013, 2015; Martinovic et al., 2018).

Figure 6E shows a histogram of the percentage of trials that repeated an attended feature.

In two studies, the to-be-attended color was held constant on each block (100% repeats

(Jiang et al., 2017; Painter et al., 2014). In the majority of the remaining studies, only two

unique features were used so the percentage of trials where the attended feature was

repeated was on average quite high (overall median = 50%, SD = 27.4%, min = 0%, max

= 100%). Finally, in three studies, the attention conditions were perfectly alternated (0%

repeat trials; Allison et al., 2008; Chen et al., 2003; J. Wang et al., 2007). Consistent with

a priming account, 81% of the studies with a high percentage of repeats showed a

consistent positive attention effect, whereas none of the studies with 0% repeat trials

showed a consistent positive attention effect. However, we think the inconsistent effects

in the studies with 0% repeats might be equally attributed to their low trial counts (median

= 8 trials per condition; Boudewyn et al., 2018). Only one study had a similar proportion

of repeats as the present study (Störmer & Alvarez, 2014). Störmer and Alvarez found a

significant attention effect while using 5 unique colors (intermixed randomly from trial to

trial). The findings by Störmer and Alvarez provide evidence against the feature-based

priming account, and suggest the task factor "number of colors" cannot definitively explain

our inability to observe an attention effect. However, given the lack of extant work using

unpredictable color cues, we think future, systematic work is needed to determine the

degree to which inter-trial priming effects may modulate the size and reliability of feature-

based attention effects.

**SSVEP Frequencies.**

Finally, we examined frequencies that have been most commonly used in the literature.

In our study, we chose relatively high frequencies (24 and 30 Hz) in order to have

increased temporal resolution for detecting potential time-course effects. In addition,

some have argued that using higher frequencies as advantages for driving a more

localized portion of visual cortex, as opposed to broadly driving visual, parietal and frontal

cortex when using lower frequencies in the theta/alpha bands (i.e., ~7-12Hz; Ding et al.,

2006; Srinivasan et al., 2006). For the purposes of temporal resolution, earlier work

examining the time course of *spatial* attention with SSVEPs used the frequencies 20 and

28 Hz, (Müller et al., 1998). However, upon reviewing the feature-based attention

literature, we found that our chosen frequencies were outside the range that has

previously been used with a feature-based attention task (Figure 6F; median = 10 Hz, SD

= 3.6 Hz, min = 2.4 Hz, max = 19.75 Hz). Thus, it is possible that feature-based attention,

unlike spatial attention, cannot be easily tracked with flicker frequencies above ~20 Hz.

Although some studies have argued that feature-based attention effects do not

qualitatively appear to vary with frequency (Martinovic et al., 2018; Steinhauser &

Andersen, 2019), the vast majority of reviewed studies only reported statistical

significance of an overall attention effect collapsed across frequencies. Only a handful

studies have reported statistical significance of individual frequencies (e.g., Chu &

D'Zmura, 2019; Painter et al., 2014; Quigley & Müller, 2014; Steinhauser & Andersen,

2019). As such, future work is needed to systematically investigate the effect of frequency

choice on feature-based attention effects, particularly for frequencies >20 Hz.

761                                    **Discussion**

In this pre-registered study, we sought to test whether cuing participants to *ignore* a

particular color modulates the time-course of feature-based attention as indexed by

steady-state visually evoked potentials (SSVEPs). As a baseline point of comparison, we

also included a condition in which participants were cued to *attend* a particular color. This

"attend cue" condition was intended as a close replication of much prior work showing

that SSVEP amplitudes are modulated by attention (greater amplitude for the attended

feature; e.g., Andersen et al., 2008; Andersen & Müller, 2010; Müller et al., 2006).

However, we failed to replicate this basic overall attention effect; we found no difference

in SSVEP amplitude as a function of attention in either the attend cue or the ignore cue

condition. Thus, because we found no overall SSVEP attention effect, we were unable to

test our hypotheses about how this attention effect was modulated by being cued to attend

versus cued to ignore. Despite the lack of an SSVEP attention effect, positive control

analyses indicated that that participants did successfully select the cued target color (i.e.,

we observed a significantly larger P3 component for target events in the attended color

than in the ignored color).

Given our failure to observe an effect of attention on SSVEP amplitude with our task

procedures, we performed a focused review of the literature to quantify key

methodological aspects of prior studies using SSVEPs to study feature-based attention.

Based on this review, we concluded that sample size and trial counts likely did not explain

our failure to find an effect; our sample size and trial counts were near the maximum

values found in the surveyed literature. However, two key, intentional design differences

may have hampered our ability to find an effect: (1) the number of colors in our stimulus

set and (2) the frequencies used to generate the SSVEP.

The first key design difference in our study was the number colors in our stimulus set. We

purposefully minimized the influence of inter-trial priming on our estimates of feature-

based attention (Theeuwes, 2013) by using 5 unique colors and randomly chose target

and distractor colors on each trial. According to a priming account of feature-based

attention, a relatively high proportion of trials where the attended color is repeated back-

to-back could inflate or even entirely drive apparent feature-based attention effects. Using

5 colors somewhat protects against this possibility, because it ensures that the attended

color is repeated on 20% of trials, and both the attended/ignored colors are repeated on

only 4% of trials. In the literature, we found that most studies had back-to-back color

repeats on at least 50% of trials. It is thus plausible that inter-trial priming could contribute

to observed attention differences in these studies. Contrary to a priming account,

however, one study found robust feature-based attention effects using a set of 5 unique
colors (Störmer & Alvarez, 2014), suggesting that participants can use a cue to direct
feature-based attention even when the proportion of repeated trials is relatively low. To
date, however, no study has directly manipulated the proportion of repeated trials or the
number of possible stimulus colors in an SSVEP study. Given emerging evidence that
history-driven effects play an important role in shaping both spatial and feature-based
attentional selection (Adam & Serences, in preparation; Awh et al., 2012; Failing et al.,
2019; B. Wang & Theeuwes, 2018a, 2018b), we think that future work is needed to directly
investigate whether and to what degree SSVEP estimates of feature-based attention are
modulated by inter-trial priming.
808
The second key design difference in our study was the chosen set of frequencies. To
ensure adequate temporal resolution to characterize time-course effects, we chose to use
slightly higher frequencies (24 and 30 Hz). We believed these values would be
reasonable, because an initial study of the time-course of spatial attention used SSVEP
frequencies in a similar range (20 and 28 Hz; Müller et al., 1998). In addition, frequencies
>20 Hz have commonly been used in other SSVEP studies of spatial attention (Garcia et
al., 2013; Kashiwase et al., 2012; Müller, Picton, et al., 1998; Müller & Hillyard, 2000;
Toffanin et al., 2009; Won et al., 2016), and SSVEPs are overall robust using a wide array
of frequencies (at least 1 to 50 Hz; Herrmann, 2001). However, the overall degree of
entrainment, estimated spatial extent of entrainment, and the size of spatial attention
effects vary with frequency (Ding et al., 2006; Gulbinaite et al., 2014; Herrmann, 2001).
Given differences in the cortical processing of locations and features (Cohen & Maunsell,
2011; Haxby et al., 1994; Owen et al., 1996), and differences in the spatial extent and
degree  of entrainment with frequency (Ding et al., 2006; Gulbinaite et al., 2014), it is
plausible that feature-based attention can only be tracked with a limited range of
frequencies (e.g., frequencies near the alpha band). Future work will be needed to
systematically investigate the effect of SSVEP frequency on feature-based attention.
826

It is perhaps puzzling that frequencies above 20 Hz have been commonly used in the

spatial attention literature but have not been used in the feature-based attention literature.

The truncation of the frequency distribution in the reviewed literature could be a piece of

the "file drawer" in action. It is possible that other researchers likewise discovered that

they were unable to track feature-based attention using certain frequencies, but that these

null results were never published due to journals' and authors' biases toward publishing

positive results (Ferguson & Heene, 2012; Rosenthal, 1979). Thus, our results highlight

the practical and theoretical importance of regularly publishing null results. On the

practical side, if prior null results had been published, we may have better known which

frequencies to use or avoid, and we would have been able to test our key hypotheses.

On the theoretical side, our results highlight how seemingly unimportant null results can

have implications for theory when viewed in the context of the broader literature. For

example, if certain frequencies track spatial but not feature-based attention, this may

inform our understanding of the brain networks and cognitive processes differentially

modulated by flicker frequency (Ding et al., 2006; Srinivasan et al., 2006).

In short, we found no evidence that SSVEPs track the deployment of feature-based

attention with our procedures, and future methodological work is needed to determine

constraints on generalizability of the SSVEP method for tracking feature-based attention.

We performed a focused review of prior studies using SSVEPs to study feature-based

attention, and from this review we identified two key factors (frequencies used; likelihood

of inter-trial feature priming) that should be systematically investigated in future work.

## References

Adam, K. C. S., & Serences, J. T. (in preparation). *History-driven modulations of population codes in early visual cortex during visual search*.

Adamian, N., Slaustaite, E., & Andersen, S. K. (2019). Top–Down Attention Is Limited Within but Not Between Feature Dimensions. *Journal of Cognitive Neuroscience*, *31*(8), 1173–1183. https://doi.org/10.1162/jocn_a_01383

Adrian, E. D., & Matthews, B. H. C. (1934). The Berger Rhythm: Potential Changes from the Occiptial Lobes in Man. *Brain*, *57*(4), 355–385.

Allison, B. Z., McFarland, D. J., Schalk, G., Zheng, S. D., Jackson, M. M., & Wolpaw, J. R. (2008). Towards an independent brain–computer interface using steady state visual

evoked potentials. *Clinical Neurophysiology*, *119*(2), 399–408.
https://doi.org/10.1016/j.clinph.2007.09.121

Andersen, S. K., Fuchs, S., & Müller, M. M. (2011). Effects of Feature-selective and Spatial
Attention at Different Stages of Visual Processing. *Journal of Cognitive Neuroscience*,
*23*(1), 238–246. https://doi.org/10.1162/jocn.2009.21328

Andersen, S. K., Hillyard, S. A., & Müller, M. M. (2008). Attention Facilitates Multiple Stimulus
Features in Parallel in Human Visual Cortex. *Current Biology*, *18*(13), 1006–1009.
https://doi.org/10.1016/j.cub.2008.06.030

Andersen, S. K., Hillyard, S. A., & Müller, M. M. (2013). Global Facilitation of Attended Features
Is Obligatory and Restricts Divided Attention. *Journal of Neuroscience*, *33*(46), 18200–
18207. https://doi.org/10.1523/JNEUROSCI.1913-13.2013

Andersen, S. K., & Müller, M. M. (2010). Behavioral performance follows the time course of
neural facilitation and suppression during cued shifts of feature-selective attention.
*Proceedings of the National Academy of Sciences*, *107*(31), 13878–13882.
https://doi.org/10.1073/pnas.1002436107

Andersen, S. K., Müller, M. M., & Hillyard, S. A. (2009). Color-selective attention need not be
mediated by spatial attention. *Journal of Vision*, *9*(6), 2–2. https://doi.org/10.1167/9.6.2

Andersen, S. K., Müller, M. M., & Hillyard, S. A. (2011). Tracking the allocation of attention in
visual scenes with steady-state evoked potentials. In M. I. Posner (Ed.), *Cognitive
Neuroscience of Attention* (2nd ed.). The Guilford Press.

Andersen, S. K., Müller, M. M., & Hillyard, S. A. (2015). Attentional Selection of Feature
Conjunctions Is Accomplished by Parallel and Independent Selection of Single Features.
*Journal of Neuroscience*, *35*(27), 9912–9919. https://doi.org/10.1523/JNEUROSCI.5268-
14.2015

Andersen, S. K., Müller, M. M., & Martinovic, J. (2012). Bottom-Up Biases in Feature-Selective
Attention. *Journal of Neuroscience*, *32*(47), 16953–16958.
https://doi.org/10.1523/JNEUROSCI.1767-12.2012

Appelbaum, L. G., & Norcia, A. M. (2009). Attentive and pre-attentive aspects of figural
processing. *Journal of Vision*, *9*(11), 18–18. https://doi.org/10.1167/9.11.18

Arita, J. T., Carlisle, N. B., & Woodman, G. F. (2012). Templates for rejection: Configuring
attention to ignore task-irrelevant features. *Journal of Experimental Psychology: Human
Perception and Performance*, *38*(3), 580–584. https://doi.org/10.1037/a0027885

Awh, E., Belopolsky, A. V., & Theeuwes, J. (2012). Top-down versus bottom-up attentional
control: A failed theoretical dichotomy. *Trends in Cognitive Sciences*, *16*(8), 437–443.
https://doi.org/10.1016/j.tics.2012.06.010

Beck, V. M., & Hollingworth, A. (2015). Evidence for negative feature guidance in visual search
is explained by spatial recoding. *Journal of Experimental Psychology: Human Perception
and Performance*, *41*(5), 1190–1196. https://doi.org/10.1037/xhp0000109

Becker, M. W., Hemsteger, S., & Peltier, C. (2015). No templates for rejection: A failure to
configure attention to ignore task-irrelevant features. *Visual Cognition*, *23*(9–10), 1150–
1167. https://doi.org/10.1080/13506285.2016.1149532

Boudewyn, M. A., Luck, S. J., Farrens, J. L., & Kappenman, E. S. (2018). How many trials does
it take to get a significant ERP effect? It depends. *Psychophysiology*, *55*(6), e13049.
https://doi.org/10.1111/psyp.13049

Boylan, M. R., Kelly, M. N., Thigpen, N. N., & Keil, A. (2019). Attention to a threat-related
feature does not interfere with concurrent attentive feature selection. *Psychophysiology*,
*56*(6), e13332. https://doi.org/10.1111/psyp.13332

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436.
https://doi.org/10.1163/156856897X00357

Bridwell, D. A., & Srinivasan, R. (2012). Distinct Attention Networks for Feature Enhancement and Suppression in Vision. *Psychological Science*, *23*(10), 1151–1158. https://doi.org/10.1177/0956797612440099

Canolty, R. T., Edwards, E., Dalal, S. S., Soltani, M., Nagarajan, S. S., Kirsch, H. E., Berger, M. S., Barbaro, N. M., & Knight, R. T. (2006). High Gamma Power Is Phase-Locked to Theta Oscillations in Human Neocortex. *Science*, *313*(5793), 1626–1628. https://doi.org/10.1126/science.1128115

Carlisle, N. B., & Nitka, A. W. (2019). Location-based explanations do not account for active attentional suppression. *Visual Cognition*, *27*(3–4), 305–316. https://doi.org/10.1080/13506285.2018.1553222

Chen, Y., Seth, A. K., Gally, J. A., & Edelman, G. M. (2003). The power of human brain magnetoencephalographic signals can be modulated up or down by changes in an attentive visual task. *Proceedings of the National Academy of Sciences*, *100*(6), 3501–3506. https://doi.org/10.1073/pnas.0337630100

Chu, V. C., & D'Zmura, M. (2019). Tracking feature-based attention. *Journal of Neural Engineering*, *16*(1), 016022. https://doi.org/10.1088/1741-2552/aaed17

Clementz, B. A., Wang, J., & Keil, A. (2008). Normal Electrocortical Facilitation But Abnormal Target Identification during Visual Sustained Attention in Schizophrenia. *Journal of Neuroscience*, *28*(50), 13411–13418. https://doi.org/10.1523/JNEUROSCI.4095-08.2008

Cohen, M. R., & Maunsell, J. H. R. (2011). Using Neuronal Populations to Study the Mechanisms Underlying Spatial and Feature Attention. *Neuron*, *70*(6), 1192–1204. https://doi.org/10.1016/j.neuron.2011.04.029

Conci, M., Deichsel, C., Müller, H. J., & Töllner, T. (2019). Feature guidance by negative attentional templates depends on search difficulty. *Visual Cognition*, *27*(3–4), 317–326. https://doi.org/10.1080/13506285.2019.1581316

Cunningham, C. A., & Egeth, H. E. (2016). Taming the White Bear: Initial Costs and Eventual Benefits of Distractor Inhibition. *Psychological Science*, *27*(4), 476–485. https://doi.org/10.1177/0956797615626564

Ding, J., Sperling, G., & Srinivasan, R. (2006). Attentional Modulation of SSVEP Power Depends on the Network Tagged by the Flicker Frequency. *Cerebral Cortex*, *16*(7), 1016–1029. https://doi.org/10.1093/cercor/bhj044

Donohue, S. E., Bartsch, M. V., Heinze, H.-J., Schoenfeld, M. A., & Hopf, J.-M. (2018). Cortical Mechanisms of Prioritizing Selection for Rejection in Visual Search. *The Journal of Neuroscience*, *38*(20), 4738–4748. https://doi.org/10.1523/JNEUROSCI.2407-17.2018

Failing, M., Feldmann-Wüstefeld, T., Wang, B., Olivers, C., & Theeuwes, J. (2019). Statistical regularities induce spatial as well as feature-specific suppression. *Journal of Experimental Psychology: Human Perception and Performance*, *45*(10), 1291–1303. https://doi.org/10.1037/xhp0000660

Ferguson, C. J., & Heene, M. (2012). A Vast Graveyard of Undead Theories: Publication Bias and Psychological Science's Aversion to the Null. *Perspectives on Psychological Science*, *7*(6), 555–561. https://doi.org/10.1177/1745691612459059

Forschack, N., Andersen, S. K., & Müller, M. M. (2017). Global Enhancement but Local Suppression in Feature-based Attention. *Journal of Cognitive Neuroscience, 29*(4), 619–627. https://doi.org/10.1162/jocn_a_01075

Garcia, J. O., Srinivasan, R., & Serences, J. T. (2013). Near-Real-Time Feature-Selective Modulations in Human Cortex. *Current Biology*, *23*(6), 515–522. https://doi.org/10.1016/j.cub.2013.02.013

Gulbinaite, R., Johnson, A., de Jong, R., Morey, C. C., & van Rijn, H. (2014). Dissociable mechanisms underlying individual differences in visual working memory capacity. *NeuroImage*, *99*, 197–206. https://doi.org/10.1016/j.neuroimage.2014.05.060

Hasan, R., Srinivasan, R., & Grossman, E. D. (2017). Feature-based attentional tuning during biological motion detection measured with SSVEP. *Journal of Vision*, *17*(9), 22. https://doi.org/10.1167/17.9.22

Haxby, J., Horwitz, B., Ungerleider, L., Maisog, J., Pietrini, P., & Grady, C. (1994). The functional organization of human extrastriate cortex: A PET-rCBF study of selective attention to faces and locations. *The Journal of Neuroscience*, *14*(11), 6336–6353. https://doi.org/10.1523/JNEUROSCI.14-11-06336.1994

Herrmann, C. S. (2001). Human EEG responses to 1?100 Hz flicker: Resonance phenomena in visual cortex and their potential correlation to cognitive phenomena. *Experimental Brain Research*, *137*(3–4), 346–353. https://doi.org/10.1007/s002210100682

Ipata, A. E., Gee, A. L., Gottlieb, J., Bisley, J. W., & Goldberg, M. E. (2006). LIP responses to a popout stimulus are reduced if it is overtly ignored. *Nature Neuroscience*, *9*(8), 1071–1076. https://doi.org/10.1038/nn1734

Itthipuripat, S., Deering, S., & Serences, J. T. (2019). When Conflict Cannot be Avoided: Relative Contributions of Early Selection and Frontal Executive Control in Mitigating Stroop Conflict. *Cerebral Cortex*, *29*(12), 5037–5048. https://doi.org/10.1093/cercor/bhz042

Itthipuripat, S., Garcia, J. O., & Serences, J. T. (2013). Temporal dynamics of divided spatial attention. *Journal of Neurophysiology*, *109*(9), 2364–2373. https://doi.org/10.1152/jn.01051.2012

Jiang, Y., Wu, X., & Gao, X. (2017). A category-specific top-down attentional set can affect the neural responses outside the current focus of attention. *Neuroscience Letters*, *659*, 80–85. https://doi.org/10.1016/j.neulet.2017.07.029

Kaiser, P. K. (1988). Sensation luminance: A new name to distinguish CIE luminance from luminance dependent on an individual's spectral sensitivity. *Vision Research*, *28*(3), 455–456. https://doi.org/10.1016/0042-6989(88)90186-1

Kashiwase, Y., Matsumiya, K., Kuriki, I., & Shioiri, S. (2012). Time Courses of Attentional Modulation in Neural Amplification and Synchronization Measured with Steady-state Visual-evoked Potentials. *Journal of Cognitive Neuroscience*, *24*(8), 1779–1793. https://doi.org/10.1162/jocn_a_00212

Keitel, C., & Müller, M. M. (2016). Audio-visual synchrony and feature-selective attention co-amplify early visual processing. *Experimental Brain Research*, *234*(5), 1221–1231. https://doi.org/10.1007/s00221-015-4392-8

Kleiner, M., Brainard, D., & Pelli, D. (2007). *What's new in Psychtoolbox-3?* European Conference on Visual Perception (ECVP), Arezzo, Italy. https://pdfs.semanticscholar.org/04d4/7572cec08b7a582a9366e5ac61dcfd633f2a.pdf

Kravitz, D., & Mitroff, S. (2017). Estimates of a priori power and false discovery rates induced by post-hoc changes from thousands of independent replications. *Journal of Vision*, *17*(10), 223. https://doi.org/10.1167/17.10.223

Luck, S. J. (2005). *An Introduction to the Event-Related Potential Technique* (1st ed.). MIT Press.

Martinez-Trujillo, J. C., & Treue, S. (2004). Feature-Based Attention Increases the Selectivity of Population Responses in Primate Visual Cortex. *Current Biology*, *14*(9), 744–751. https://doi.org/10.1016/j.cub.2004.04.028

Martinovic, J., & Andersen, S. K. (2018). Cortical summation and attentional modulation of combined chromatic and luminance signals. *NeuroImage*, *176*, 390–403. https://doi.org/10.1016/j.neuroimage.2018.04.066

Martinovic, J., Wuerger, S. M., Hillyard, S. A., Müller, M. M., & Andersen, S. K. (2018). Neural mechanisms of divided feature-selective attention to colour. *NeuroImage*, *181*, 670–682. https://doi.org/10.1016/j.neuroimage.2018.07.033

Moher, J., & Egeth, H. E. (2012). The ignoring paradox: Cueing distractor features leads first to selection, then to inhibition of to-be-ignored items. *Attention, Perception, & Psychophysics*, *74*(8), 1590–1605. https://doi.org/10.3758/s13414-012-0358-0

Moher, J., Lakshmanan, B. M., Egeth, H. E., & Ewen, J. B. (2014). Inhibition Drives Early Feature-Based Attention. *Psychological Science*, *25*(2), 315–324. https://doi.org/10.1177/0956797613511257

Morgan, S. T., Hansen, J. C., & Hillyard, S. A. (1996). Selective attention to stimulus location modulates the steady-state visual evoked potential. *Proceedings of the National Academy of Sciences of the United States of America*, *93*(10), 4770–4774.

Müller, M. M., Andersen, S., Trujillo, N. J., Valdes-Sosa, P., Malinowski, P., & Hillyard, S. A. (2006). Feature-selective attention enhances color signals in early visual areas of the human brain. *Proceedings of the National Academy of Sciences*, *103*(38), 14250–14254. https://doi.org/10.1073/pnas.0606668103

Müller, M. M., Gundlach, C., Forschack, N., & Brummerloh, B. (2018). It takes two to tango: Suppression of task-irrelevant features requires (spatial) competition. *NeuroImage*, *178*, 485–492. https://doi.org/10.1016/j.neuroimage.2018.05.073

Müller, M. M., & Hillyard, S. (2000). Concurrent recording of steady-state and transient event-related potentials as indices of visual-spatial selective attention. *Clinical Neurophysiology*, *111*(9), 1544–1552. https://doi.org/10.1016/S1388-2457(00)00371-0

Müller, M. M., Picton, T. W., Valdes-Sosa, P., Riera, J., Teder-Sälejärvi, W. A., & Hillyard, S. A. (1998). Effects of spatial selective attention on the steady-state visual evoked potential in the 20–28 Hz range. *Cognitive Brain Research*, *6*(4), 249–261. https://doi.org/10.1016/S0926-6410(97)00036-0

Müller, M. M., Teder-Sälejärvi, W., & Hillyard, S. A. (1998). The time course of cortical facilitation during cued shifts of spatial attention. *Nature Neuroscience*, *1*(7), 631–634. https://doi.org/10.1038/2865

Norcia, A. M., Appelbaum, L. G., Ales, J. M., Cottereau, B. R., & Rossion, B. (2015). The steady-state visual evoked potential in vision research: A review. *Journal of Vision*, *15*(6), 4. https://doi.org/10.1167/15.6.4

Owen, A. M., Milner, B., Petrides, M., & Evans, A. C. (1996). Memory for object features versus memory for object location: A positron-emission tomography study of encoding and retrieval processes. *Proceedings of the National Academy of Sciences*, *93*(17), 9212–9217. https://doi.org/10.1073/pnas.93.17.9212

Painter, D. R., Dux, P. E., & Mattingley, J. B. (2015). Causal involvement of visual area MT in global feature-based enhancement but not contingent attentional capture. *NeuroImage*, *118*, 90–102. https://doi.org/10.1016/j.neuroimage.2015.06.019

Painter, D. R., Dux, P. E., Travis, S. L., & Mattingley, J. B. (2014). Neural Responses to Target Features outside a Search Array Are Enhanced during Conjunction but Not Unique-Feature Search. *Journal of Neuroscience*, *34*(9), 3390–3401. https://doi.org/10.1523/JNEUROSCI.3630-13.2014

Pei, F., Pettet, M. W., & Norcia, A. M. (2002). Neural correlates of object-based attention. *Journal of Vision*, *2*(9), 1. https://doi.org/10.1167/2.9.1

Pelli, D. G. (1997). The VideoToolbox software for visual psychophysics: Transforming numbers into movies. *Spatial Vision*, *10*(4), 437–442. https://doi.org/10.1163/156856897X00366

Quigley, C., Andersen, S. K., Schulze, L., Grunwald, M., & Müller, M. M. (2010). Feature-selective attention: Evidence for a decline in old age. *Neuroscience Letters*, *474*(1), 5–8. https://doi.org/10.1016/j.neulet.2010.02.053

Quigley, C., & Müller, M. M. (2014). Feature-Selective Attention in Healthy Old Age: A Selective Decline in Selective Attention? *Journal of Neuroscience*, *34*(7), 2471–2476. https://doi.org/10.1523/JNEUROSCI.2718-13.2014

Reeder, R. R., Olivers, C. N. L., & Pollmann, S. (2017). Cortical evidence for negative search templates. *Visual Cognition*, *25*(1–3), 278–290. https://doi.org/10.1080/13506285.2017.1339755

Regan, D. (1977). Steady-state evoked potentials. *Journal of the Optical Society of America*, *67*(11), 1475. https://doi.org/10.1364/JOSA.67.001475

Rosenthal, R. (1979). The file drawer problem and tolerance for null results. *Psychological Bulletin*, *86*(3), 638–641. https://doi.org/10.1037/0033-2909.86.3.638

Rungratsameetaweemana, N., Itthipuripat, S., Salazar, A., & Serences, J. T. (2018). Expectations Do Not Alter Early Sensory Processing during Perceptual Decision-Making. *The Journal of Neuroscience*, *38*(24), 5632–5648. https://doi.org/10.1523/JNEUROSCI.3638-17.2018

Sawaki, R., & Luck, S. J. (2010). Capture versus suppression of attention by salient singletons: Electrophysiological evidence for an automatic attend-to-me signal. *Attention, Perception, & Psychophysics*, *72*(6), 1455–1470. https://doi.org/10.3758/APP.72.6.1455

Srinivasan, R., Bibi, F. A., & Nunez, P. L. (2006). Steady-State Visual Evoked Potentials: Distributed Local Sources and Wave-Like Dynamics Are Sensitive to Flicker Frequency. *Brain Topography*, *18*(3), 167–187. https://doi.org/10.1007/s10548-006-0267-4

Steinhauser, M., & Andersen, S. K. (2019). Rapid adaptive adjustments of selective attention following errors revealed by the time course of steady-state visual evoked potentials. *NeuroImage*, *186*, 83–92. https://doi.org/10.1016/j.neuroimage.2018.10.059

Störmer, V. S., & Alvarez, G. A. (2014). Feature-Based Attention Elicits Surround Suppression in Feature Space. *Current Biology*, *24*(17), 1985–1988. https://doi.org/10.1016/j.cub.2014.07.030

Talsma, D., Doty, T. J., Strowd, R., & Woldorff, M. G. (2006). Attentional capacity for processing concurrent stimuli is larger across sensory modalities than within a modality. *Psychophysiology*, *43*(6), 541–549. https://doi.org/10.1111/j.1469-8986.2006.00452.x

Theeuwes, J. (2013). Feature-based attention: It is all bottom-up priming. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *368*(1628), 20130055. https://doi.org/10.1098/rstb.2013.0055

Thigpen, N., Petro, N. M., Oschwald, J., Oberauer, K., & Keil, A. (2019). Selection of Visual Objects in Perception and Working Memory One at a Time. *Psychological Science*, *30*(9), 1259–1272. https://doi.org/10.1177/0956797619854067

Toffanin, P., de Jong, R., Johnson, A., & Martens, S. (2009). Using frequency tagging to quantify attentional deployment in a visual divided attention task. *International Journal of Psychophysiology*, *72*(3), 289–298. https://doi.org/10.1016/j.ijpsycho.2009.01.006

Van Moorselaar, D., & Slagter, H. A. (2020). Inhibition in selective attention. *Annals of the New York Academy of Sciences*, *1464*(1), 204–221.

Vatterott, D. B., & Vecera, S. P. (2012). Experience-dependent attentional tuning of distractor rejection. *Psychonomic Bulletin & Review*, *19*(5), 871–878. https://doi.org/10.3758/s13423-012-0280-4

Verghese, P., Kim, Y.-J., & Wade, A. R. (2012). Attention Selects Informative Neural
  Populations in Human V1. *Journal of Neuroscience*, *32*(46), 16379–16390.
  https://doi.org/10.1523/JNEUROSCI.1174-12.2012

Vialatte, F.-B., Maurice, M., Dauwels, J., & Cichocki, A. (2010). Steady-state visually evoked
  potentials: Focus on essential paradigms and future perspectives. *Progress in
  Neurobiology*, *90*(4), 418–438. https://doi.org/10.1016/j.pneurobio.2009.11.005

Wang, B., & Theeuwes, J. (2018a). Statistical regularities modulate attentional capture. *Journal
  of Experimental Psychology: Human Perception and Performance, 44*(1), 13–17.
  https://doi.org/10.1037/xhp0000472

Wang, B., & Theeuwes, J. (2018b). How to inhibit a distractor location? Statistical learning
  versus active, top-down suppression. *Attention, Perception, & Psychophysics*.
  https://doi.org/10.3758/s13414-018-1493-z

Wang, J., Clementz, B. A., & Keil, A. (2007). The neural correlates of feature-based selective
  attention when viewing spatially and temporally overlapping images. *Neuropsychologia*,
  *45*(7), 1393–1399. https://doi.org/10.1016/j.neuropsychologia.2006.10.019

Won, D.-O., Hwang, H.-J., Dähne, S., Müller, K.-R., & Lee, S.-W. (2016). Effect of higher
  frequency on the classification of steady-state visual evoked potentials. *Journal of
  Neural Engineering*, *13*(1), 016014. https://doi.org/10.1088/1741-2560/13/1/016014

Zhang, D., Maye, A., Gao, X., Hong, B., Engel, A. K., & Gao, S. (2010). An independent brain–
  computer interface using covert non-spatial visual selective attention. *Journal of Neural
  Engineering*, *7*(1), 016010. https://doi.org/10.1088/1741-2560/7/1/016010

Zhang, W., & Luck, S. J. (2009). Feature-based attention modulates feedforward visual
  processing. *Nature Neuroscience*, *12*(1), 24–25. https://doi.org/10.1038/nn.2223