2
3   **Steady-state visually evoked potentials and feature-based attention: Pre-**
4   **registered null results and a focused review of methodological considerations.**
5

6   Kirsten C.S. Adam[1,2], Lillian Chang[3], Nicole Rangan[1], & John T. Serences[1,2,4]

7
8   [1] Department of Psychology, *University of California San Diego*
9   [2] Institute for Neural Computation*, University of California San Diego*
10   [3] Department of Neuroscience, *Georgetown University Medical Center*
11   [4] Neurosciences Graduate Program, *University of California San Diego*

12
13   **Tables:** 0
14   **Figures:** 6
15   **Appendices:** 20
16
17   **Keywords:** attention; EEG; SSVEP; feature-based attention
18

19   **Contributions:** K.A. & J.S. planned the study. K.A., L.C., and N.R. collected data and
20   performed the literature review. K.A. performed analyses and drafted the manuscript. All
21   authors revised the manuscript.
22
23   **Funding:** Research was supported by National Eye Institute grant R01 EY025872
24   (J.S.). and National Institute of Mental Health grant 5T32-MH020002 (K.A.).
25
26   **Pre-registration information:** We published a pre-registered research plan on the
27   Open Science Framework prior to data collection (https://osf.io/kfg9h/).
28
29   **Data availability:** Data and analysis code will be made available online on the Open
30   Science Framework at https://osf.io/ew7dv/ upon acceptance for publication.
31
32   **Acknowledgements:** We thank Matteo d'Amico for additional assistance with data
33   collection and Kelvin Lam for help as lab manager. We thank Rosanne Rademaker and
34   Angus Chapman for sharing Psychtoolbox code that was adapted for use here.
35
Conflicts of interest: none

Correspondence to:
Kirsten C. S. Adam
University of California San Diego
9500 Gilman Drive, Mail Code: 0109
La Jolla, CA 92093-010
kadam@ucsd.edu

45   **Abstract**

Feature-based attention is the ability to selectively attend to a particular feature (e.g.,

attend to red but not green items while looking for the ketchup bottle in your

refrigerator), and steady-state visually evoked potentials (SSVEPs) measured from the

human electroencephalogram (EEG) signal have been used to track the neural

deployment of feature-based attention. Although many published studies suggest that

we can use trial-by-trial cues to enhance relevant feature information (i.e., greater

SSVEP response to the cued color), there is ongoing debate about whether participants

may likewise use trial-by-trial cues to voluntarily ignore a particular feature. Here, we

report the results of a pre-registered study in which participants either were cued to

attend or to ignore a color. Counter to prior work, we found no attention-related

modulation of the SSVEP response in either cue condition. However, positive control

analyses revealed that participants paid some degree of attention to the cued color (i.e.,

we observed a greater P300 component to targets in the attended versus the

unattended color). In light of these unexpected null results, we conducted a focused

review of methodological considerations for studies of feature-based attention using

SSVEPs. In the review, we quantify potentially important stimulus parameters that have

been used in the past (e.g., stimulation frequency; trial counts) and we discuss the

potential importance of these and other task factors (e.g., feature-based priming) for

SSVEP studies.

76   **Introduction**

Attending to a specific feature leads to systematic changes in the firing rates of

neurons that encode the relevant feature space. For example, when looking for a ripe

tomato, the firing rate of neurons tuned to red will be enhanced and the firing rate of

neurons tuned to other features will be suppressed (e.g. responses to green; Bartsch et

al., 2017; Ipata et al., 2006; Kiyonaga & Egner, 2016; Martinez-Trujillo & Treue, 2004;

Störmer & Alvarez, 2014; Y. Wang et al., 2015). Although there is broad agreement that

participants may learn to suppress irrelevant distractors with sufficient experience, there

is disagreement about whether these behavioral suppression effects may be volitionally

implemented on a trial-by-trial basis in response to an abstract cue (i.e., a "volitional

account"), or if they instead are solely implemented via implicit or statistical learning

mechanisms (i.e., a "priming-based" account). Consistent with a volitional or proactive

account, some work has found that participants can learn to use a trial-by-trial cue to

ignore a particular color (Arita et al., 2012; Carlisle & Nitka, 2019; Chang & Egeth, 2019;

Conci et al., 2019; Moher & Egeth, 2012; Reeder et al., 2017; Z. Zhang et al., 2020, for

nuanced reviews, see: Geng, 2014; Van Moorselaar & Slagter, 2020). However, other

work has found that a specific color needs to be repeated over many trials to be

suppressed, consistent with a priming or passive account (Cunningham & Egeth, 2016;

Failing et al., 2019; Geng et al., 2019; Lamy et al., 2008; Stilwell & Vecera, 2019;

Theeuwes, 2013; Vatterott & Vecera, 2012; B.-Y. Won & Geng, 2020).

Although many studies have examined the effects of feature-based suppression

on later selection-related event-related potential (ERP) markers such as the N2pc and

Pd (Arita et al., 2012; Carlisle & Nitka, 2019; Donohue et al., 2018; Sawaki & Luck,

2010), a key open question is whether trial-by-trial cues to ignore a color modulate

earlier stages of visual processing. Recent work by Reeder and colleagues (Reeder et

al., 2017) hypothesized that cues about which feature to ignore (i.e., "negative cues")

may down-regulate processing in visual cortex during the pre-stimulus period.

Consistent with this hypothesis, they found that overall BOLD activity in early visual

cortex was lower when participants were given a negative cue about the target color

than when participants were given either a positive or neutral cue. One limitation of this

study, however, is that the authors were unable to test whether these univariate effects

were actually feature-specific (as opposed to task-general anticipation). However,

evidence from ERP studies suggests that feature-based attention modulates early

visual processing in a feature-specific manner, as indexed by the P1 component (Moher

et al., 2014; W. Zhang & Luck, 2009). Critically, however, these ERP studies used a

design where the same target and distractor colors were repeated over many trial

events. Thus, it is not clear whether feature-based attention can modulate early visual

processing on a trial-by-trial basis, or if modulation of early visual processing is

achieved primarily via inter-trial priming (Lamy & Kristjansson, 2013; Theeuwes, 2013).

To attempt to address this gap, we conducted a pre-registered experiment in which we

measured steady-state visually evoked potentials (SSVEPs) while giving participants

trial-by-trial cues to suppress feature information.

When visual input flickers continuously at a given frequency (e.g., one stimulus at

24 Hz and another at 30 Hz), the visually evoked potential in the electroencephalogram

(EEG) signal reflects these "steady states" and time-frequency analyses can be used to

derive estimates of the strength of neural responses to each stimulus (Adrian &

Matthews, 1934; Regan, 1977). The amplitude of the frequency-specific SSVEP

response has been shown to be modulated by both spatial and feature-based attention

(higher amplitude when attended; Chen et al., 2003; Morgan et al., 1996; Müller et al.,

1998, 2006; Pei et al., 2002). When participants are cued on a trial-by-trial basis to

attend to a particular feature (e.g., color), the SSVEP amplitude is higher for the

attended feature (Andersen et al., 2008; Chen et al., 2003; Müller et al., 2006). Further,

the time-course of the SSVEP response to an attended color reveals an early

enhancement followed by a suppressed response to the irrelevant, non-attended color

(Andersen & Müller, 2010; Forschack et al., 2017).

Our primary manipulation was whether we cued participants to actively attend or

to actively ignore a color on a trial-by-trial basis. We planned to use this method to track

enhancement vs. suppression of the SSVEP response, and to test whether the time-

course of enhancement and suppression varies with cue type. In the "attend cue"

condition, participants were cued about the relevant color to attend. This condition was

expected to replicate prior work examining the time-course of feature-based attention

using SSVEPs (Andersen & Müller, 2010), whereby enhancement of the attended color

is followed by suppression of the ignored color. In the "ignore cue" condition, we instead

cued participants about which color to *ignore.* If participants can use a cue to *directly*

suppress a color on a trial-by-trial basis independent of target enhancement (i.e., a

strong version of a volitional suppression account), we predicted that the time-course of

enhancement vs. suppression of the SSVEP signal would be reduced or reversed (i.e.,

that suppression of the cued, to-be-ignored color may happen even *prior to*

enhancement of the other color). Alternatively, if participants recode the "ignore" cue to

serve as an indirect "attend" cue (e.g., "Since I'm cued to ignore blue, that means I

should attend red"; Beck & Hollingworth, 2015; Becker et al., 2015; Williams et al.,

2020), then we predicted that target enhancement would always precede distractor

suppression regardless of whether participants were cued to attend or ignore a

particular color.

To preview the results, we were unable to fully test our hypotheses about the

time-course of feature-based enhancement and suppression because we did not find

evidence for an overall attention effect with our task procedures. Despite robust SSVEP

amplitude (Cohen's $d > 5$), we observed no credible evidence that the SSVEP response

was higher for an attended versus unattended color in either cue condition. Positive

control analyses revealed that our lack of SSVEP effect was not due to a complete lack

of attention to the attended color: ERP responses (P3) to the targets were modulated by

attention as expected (Adamian et al., 2019; Andersen et al., 2013; Andersen, Fuchs, et

al., 2011). In light of our inconclusive results, we also performed a focused

methodological review of key potential task differences between our work and prior work

that may have resulted in our failure to detect the effect of feature-based attention on

SSVEP amplitude. We considered whether task factors such as stimulus flicker

frequency, sample size, stimulus duration, and stimulus color might have impacted our

ability to observe an attention effect. No single methodological factor that we considered

neatly explains our lack of effect. Given our results and literature review, we propose

that future work is needed to systematically explore two key factors: (1) variation in

feature-based attention effects across stimulus flicker frequencies and (2) the extent to

which feature-based priming modulates SSVEP attention effects.

Methods

Pre-registration and data availability

We published a pre-registered research plan on the Open Science Framework

prior to data collection (https://osf.io/kfg9h/). Our raw data and analysis code will be

made available online on the Open Science Framework at https://osf.io/ew7dv/ upon

acceptance for publication.

Participants

Healthy volunteers ($n = 32$; gender = 17 female, 15 male; mean age = 21.5 years

[SD = 3.84, min = 18, max = 39]; handedness not recorded; corrected-to-normal visual

acuity; normal color vision) participated in one 3.5 to 4 hour experimental session at the

University of California San Diego (UCSD) campus, and were compensated $15/hr.

Procedures were approved by the UCSD Institutional Review Board, and all participants

provided written informed consent. Inclusion criteria included normal or corrected-to-

normal visual acuity, normal color vision, age between 18 and 60 years old, and no self-

reported history of major neurological disorders (e.g., epilepsy, stroke). Data were

excluded from analysis if there were fewer than 400 trials in either cue condition (either

due to leaving the study early or after artifact rejection). A sample size of 24 was pre-

registered, and artifact rejection criteria were pre-registered (see section "EEG

preprocessing" below for more details). After running each participant, we checked

whether the data were usable (i.e., sufficient number of artifact-free trials) so that we

would know when to stop data collection. To reach our final sample size (n = 23

participants with usable data), we ran a total of 32 participants. Nine participants' data

were not used for the following reason: Subjects with an error in the task code (n = 3),

subjects who stopped the study early due to technical issues or to participants'

preferences (n = 4), subjects with too many artifacts (n = 2). Note, we were one subject

short of our pre-registered target sample size of 24 because data collection was

suspended due to COVID-19. However, as our later power analyses will show, we do

not believe the addition of 1 further subject would have meaningfully altered our

conclusions.

Stimuli and Procedures

Heterochromatic flicker photometry task. We chose perceptually equiluminant colors

for each participant using a heterochromatic flicker photometry task. Participants viewed

a large circular, flickering stimulus (8° radius) on a black screen (0.08 cd/m2). We

generated 5 circular color spaces in CIELAB-space with varying luminance (circles

centered on: L = 35-65, a = 0, b = 0; 5 colors equally spaced around circle with radius =

35) for use in the task. Participants matched each of the 5 colors to a medium-gray

reference color (RGB = 105.6 105.6 105.6).

On each trial, the circular background was flickered between two different colors.

One color was always medium-gray, and the other color was the to-be-matched color

on that trial. The colors of circular background were phase reversed at a rate of 24 Hz,

giving the appearance of a fast flicker when the subjective luminance values were not

matched. On top of the flickering circular stimulus small oriented bars were drawn in the

medium-gray reference color (the bars changed locations at a rate of 1Hz). The oriented

bars served no purpose other than subjectively making it easier to discriminate fine-

grained differences in luminance between the flickering colors (i.e., these bars gave

secondary visual cues about equiluminance via the "minimally distinct border"

phenomenon, Kaiser, 1988). Participants increased or decreased the luminance of the

to-be-matched color (using up and down arrow keys) until the amount of perceived

flicker was minimized – the point of perceptual equiluminance. The luminance starting

value of the to-be-matched color was chosen at random on each trial. Once satisfied

with their response, the participant pressed spacebar to continue to the next trial. Each

to-be-matched color was repeated 3 times (15 trials total).

Feature-based attention task. All stimuli were viewed on a luminance calibrated

CRT monitor (1024 x 768 resolution, 120 Hz refresh rate) from a distance of ~50 cm in a

dimly lit room. Stimuli were generated using Matlab 2016a and the Psychophysics

toolbox (Brainard, 1997; Kleiner et al., 2007; Pelli, 1997). Participants rested their chin

on a chin-rest and fixated a central dot (0.15° radius) throughout the experiment. The

stimulus was a circular aperture (radius = ~9.5°) filled with 120 oriented bars (each bar

~1.1° long and ~.1° wide). Bars were centered on a grid and separated by ~1 bar length

such that they never overlapped with one another. On each individual frame (~8.33 ms)

this grid was randomly phase shifted (0:2π in x and y coordinates) and rotated (1:360°),

thus giving the appearance of random flicker. To achieve Steady State Visually Evoked

Potentials (SSVEP) half of the bars flickered at 24 Hz (3 frames on, 2 frames off) and

the other half flickered at 30 Hz (2 frames on, 2 frames off). Due to the jittered rotation

of bar positions and to the random assignment of colors to bars on each "on" frame, this

means that the individual pixels that were "on" for each color varied from frame to

frame. The unpredictable nature of each bar's exact position is thus quite similar to

unpredictable stimuli that have been used in past work (e.g., Andersen et al., 2008). For

each "off frame" no bars of that color were shown (e.g., if 24 Hz had an "on" frame and

30 Hz had an "off" frame, then only 60 out of 120 bars would be shown on the black

background). If both the 24 Hz and 30 Hz bars were "off", then a black screen would be

shown on that frame. See Appendix A for an illustration of some example frame-by-

frame screenshots of the stimuli.

On each trial (Figure 1), the participants viewed the stimulus array of flickering,

randomly oriented bars presented on a black background (0.08 cd/m2). Half of these

bars were shown in one color (randomly chosen from the 5 possible colors) and the

other half were in another randomly chosen color (with the constraint that the two sets

of bars must be two different colors). During an initial baseline (1,333 ms), participants

viewed the flickering dots while they did not yet know which color to attend; during this

baseline, the fixation point was a medium gray color (same as the reference color in the

flicker photometry task). After the baseline, the fixation dot changed color, cuing the

participants about which color to attend or ignore. In the "attend cue" condition, the color

of the fixation point indicated which color should be attended. In the "ignore cue"

condition, the color of the fixation point indicated which color should be ignored. These

two conditions were blocked, and the order was counterbalanced across participants

(further details below). During the stimulus presentation (2,000 ms), participants

monitored the relevant color for a brief "target event" (333 ms). During this brief target

event, a percentage of lines in the relevant color will be coherent (iso-oriented).
Critically, the orientation of each coherent target or distractor event was completely
unpredictable (randomly chosen between 1-180 degrees); thus, participants could not
attend to a particular orientation in advance in order to perform the task. A target event
occurred on 50% of trials, and participants were instructed to press the spacebar as
quickly as possible if they detected a target event. Importantly, physically identical
events (iso-oriented lines in a random orientation, 333 ms) could also appear in the
distractor color (50% of trials). Participants were instructed that they should only
respond to target events; if they erroneously responded to the distractor event, the trial
was scored as incorrect. The target and/or distractor events could begin as early as cue
onset (0 ms) and no later than 1,667 ms after stimulus onset). Participants could make
responses up to 1 second into the inter-trial interval. If both a target and distractor event
were present, their onset times were separated by at least 333 ms.

Figure 1. Feature-based attention task. (A) Trial events in an example 'attend cue'
trial where there is a target event. Note, this figure is a schematic and the stimuli are not
drawn to scale. After a baseline period, participants were cued to attend or ignore one
color via a change to the fixation point color. If participants noticed a target event (~75%
iso-oriented lines in the to-be-attended color), they pressed the space bar. (B) Five
colors were used, and these 5 colors appeared with equal probability. Thus, the target
and distractor colors could be either 72 degrees or 144 degrees apart on a color wheel.
This figure shows all possible color pairings if red was the target color. (C) Examples of
cues, target events, and distractor events in the 2 main conditions. In the 'attend cue'

condition, participants made a response when the iso-oriented lines were the same
color as the cue (target event) and did not respond if the iso-oriented lines occur on the
uncued color (distractor event). In the 'ignore cue' condition, participants made a
response when the iso-oriented lines occurred on the uncued color (target event), and
they did not respond if the iso-oriented lines occurred on the cued color (distractor
event). Note, all lines were of equal size in the real experiment; lines are shown at
different widths here for easier visualization of the target and distractor colors. Here, the
iso-oriented lines are drawn at vertical in all 4 examples. In the actual task, the iso-
oriented lines could be any orientation (1-180).
287

To ensure that the task was effortful for participants, the coherence of the lines in

the target stimulus was adapted at the end of each block if behavior was outside the

range of 70 - 85% correct. At the beginning of the session, the target had 50% coherent

iso-oriented lines. If accuracy over the block of 80 trials was >85%, coherency

decreased by 5%. If block accuracy was <70%, coherency increased by 5%. The

maximum allowed coherence was 80% iso-oriented lines (so that participants would not

be able to simply individuate and attend a single position to perform the task) and the

minimum allowed coherence was 5%. The presence and absence of target and

distractor events was balanced within each block yielding a total of 4 sub-conditions

within each cue type (25% each): (1) target event + no distractor event, T1D0 (2) no

target event + distractor event, T0D1 (3) target event + distractor event, T1D1 (4) no

target event + no distractor event, T0D0.

Participants completed both task conditions (attend cue and ignore cue). The two

conditions were blocked and counterbalanced within a session (i.e., half of participants

performed the "attend cue" task for the first half of the session and the "ignore cue" task

for the second half of the session.) Each block of 80 trials took approximately 6 min 50

sec. Participants completed 18 blocks (9 per condition) for a total of 720 trials per cue

condition. Note, we originally planned for 20 blocks (10 per condition) in the pre-

registration, but the block number was reduced to 18 after the first few participants did

not finish all blocks.

Summary of deviations from the registered procedures. As described in-line

above, there were some minor deviations from the pre-registration: (1) We made

changes to the pre-registered task code to fix errors that we discovered while running

the first 3 subjects (e.g., incorrect cues and behavioral feedback in the 'ignore cue'

condition). (2) We included code for eye-tracking, which allowed us to give participants

automated real-time feedback if they blinked when they were not supposed to, i.e.,

during the stimulus period. (3) We reduced the total number of experimental blocks from

20 (10 per cue condition) to 18 (9 per cue condition) due to time constraints. (4) We had

to prematurely stop data collection at n = 23 out of 24 due to COVID-19. (5) We forgot

to specify a specific statistical test for quantifying the robustness of overall SSVEPs in

section "Checking that an SSVEP is elicited at the expected frequencies before

collecting the full sample", so we have described our justification for the statistical tests

we present here. (6) Due to unanticipated failure to detect an overall attention effect, we

performed additional non-pre-registered control analyses to attempt to rule out possible

explanations of this null effect (see section: "Non pre-registered control analyses"

below).

EEG pre-processing

Continuous EEG data were collected online from 64 Ag/AgCl active electrodes

mounted in an elastic cap using a BioSemi ActiveTwo amplifier (Cortech Solutions,

Wilmington, NC). An additional 8 external electrodes were placed on the left and right

mastoids, above and below each eye (vertical EOG), and lateral to each eye (horizontal

EOG). Continuous gaze-position data were collected from an SR Eyelink 1000+ eye-

tracker (sampling rate: 1,000 Hz; SR Research, Ottawa, Ontario). We also measured

stimulus timing with a photodiode affixed to the upper left-hand corner of the monitor (a

white dot flickered at the to-be-attended color's frequency; the photodiode and this

corner of the screen were covered with opaque black tape to ensure it was not visible).

Data were collected with a sampling rate of 1024 Hz and were not downsampled offline.

Data were saved unfiltered and unreferenced (see: Kappenman & Luck, 2010), then

referenced offline to the algebraic average of the left and right mastoids, low-pass

filtered (<80 Hz) and high-pass filtered (>.01 Hz). Artifacts were detected using

automatic criteria described below, and the data were visually inspected to confirm that

the artifact rejection criteria worked as expected. We excluded subjects with fewer than

400 trials remaining per cue condition.

Eye movements and blinks. We used the eye-tracking data and the

HEOG/VEOG traces to detect blinks and eye movements. Blinks were detected on-line

during the task using the eye tracker. If a blink was detected (i.e., missing gaze position

returned from the eye tracker), the trial was immediately terminated and the participant

was given feedback that they had blinked (i.e., the word "blink" was written in white text

in the center of the screen). If eye-tracking data could not be successfully collected

(e.g., calibration issues), the VEOG trace was used to detect blinks and/or eye

movements during offline artifact rejection. To do so, we used a split-half sliding window

step function (Luck, 2005; window size = 150 ms, step size = 10 ms, threshold = 30

microvolts.) We also used a split-half sliding-window step function to check for eye-

movements in the gaze-coordinate data from the eye-tracker (window size = 80 ms,

step size = 10 ms, threshold = 1°) and in the horizontal electrooculogram (HEOG),

window size = 150 ms, step size = 10 ms, threshold = 30 microvolts, and to detect

blinks and/or eye movements in the vertical electrooculogram (VEOG),

Drift, muscle artifacts, and blocking: We checked for drift (e.g., skin potentials)

by comparing the absolute change in voltage from the first quarter of the trial to the last

quarter of the trial. If the change in voltage exceeded 200 microvolts, the trial was

rejected for drift. In addition to slow drift, we also checked for sudden, step-like changes

in voltage with a sliding window (window size = 250 ms, step size = 20 ms, threshold =

200 microvolts). We excluded trials for muscle artifacts if any electrode had peak-to-

peak amplitude greater than 200 microvolts within a 15 ms time window (step size = 10

ms). We excluded trials for blocking if any electrode had ~120 ms during which all

values within 1 microvolt of each other (sliding 200 ms window, step size = 50 ms).

Pre-registered SSVEP analyses.

General method for SSVEP quantification. We planned to quantify the SSVEP

response by filtering the data with a Gaussian wavelet function. First, we calculated an

average ERP for each condition at electrodes of interest (O1, Oz, O2). We chose these

3 electrodes based on large SSVEP modulations at these sites in prior work (e.g.,

Itthipuripat et al., 2013; Müller et al., 2006). After calculating an ERP for each condition,

we filtered the data with a Gaussian wavelet functions (frequency-domain) with .1

fractional bandwidth to obtain frequency-domain coefficients from 20 to 35 Hz in 1-Hz

steps. The frequency-domain Gaussian filters thus had a full-width half maximum

(FWHM) that varied according to frequency, as specified by the 0.1 fractional bandwidth

parameter (e.g., 1 Hz filter = FWHM of .1 Hz, 20 Hz filter = FWHM of 2 Hz, 30 Hz filter

= FWHM of 3 Hz, etc). For a similar analytic approach see: (Canolty et al., 2006;

Itthipuripat et al., 2013; Rungratsameetaweemana et al., 2018). Signal-to-noise ratio for

each SSVEP frequency was calculated as the power at a given frequency divided by

the average power of the 2 adjacent frequencies on each side. For example, SNR of 24

Hz would be calculated as the power at 24 Hz divided by the average power at 22, 23,

25, and 26 Hz. We also pre-registered an analysis plan for examining the time-course of

SSVEP amplitude. However, because our data failed to satisfy pre-registered pre-

requisite analyses, we do not report these time-course effects here (for completeness,

we show the time-course of SNR in Figure S3).

Checking that an SSVEP is elicited at the expected frequencies before

collecting the full sample. At n = 5, we planned to confirm that our task procedure

successfully produced reliable SSVEP responses (i.e., check that we observed peaks at

the correct stimulus flicker frequencies). If our task procedures failed to elicit an SSVEP

at the expected frequencies, we had planned to stop data collection and alter the task to

troubleshoot the problem (e.g., optimize timing, choose different flicker frequencies,

make stimuli brighter, etc.). We planned to begin data collection over again if we failed

this trouble-shooting step. Note, at this early stage we only verified if the basic method

worked (SSVEP frequencies were robust): we did not test whether any hypothesized

attention effects were present, as this could inflate our false discovery rate (Kravitz &

Mitroff, 2017). Note, in the original pre-registration we failed to specify what test we

would run to determine if SSVEP frequencies were robustly represented in the EEG

signal. Theoretical chance for SNR would be 1, so the simplest test would be to

compare the SNR for our stimulation frequencies (24 and 30 Hz) to 1 using a t-test,

which we report. However, it is often is better to compare to an empirical baseline with a

reasonable amount of noise (Combrisson & Jerbi, 2015). As such, we opted to also

compute an effect size comparing the SNR for our stimulation frequencies to all other

frequencies (with the exception that we did not use frequencies +/- 2 Hz of 24 or 30 Hz

as baseline values, since SNR was calculated as the power at frequency F divided by

the power in the 2 adjacent 1-hz bins).

406   **Checking achieved power for the basic attention effect.** Without adequate

407   power for the basic attention effect, we would not be able to robustly interpret the time-

408   course of enhancement vs. suppression. At the full sample size, we thus planned to

409   check whether we had sufficient achieved power for the overall attention effect (>=80%

410   power for attended vs. ignored color collapsed across the entire stimulus period) as a

411   prerequisite for interpreting the time-course of enhancement and suppression.

412   **Checking if a priori electrodes are reasonable.** We chose a priori to analyze

the SSVEP at electrodes O1, Oz, and O2 (Itthipuripat et al., 2013;

Rungratsameetaweemana et al., 2018), but we planned to plot the topography of

SSVEP modulation across all electrodes to check that these a priori electrodes were

responsive to the SSVEP manipulation. If these electrodes were not responsive to the

SSVEP, we planned to perform a cluster-based permutation test to select a new set of

electrodes.

419   **Checking if target and/or distractor presence alters results.** Our core

analyses planned to use all trials for each condition (e.g., target event present or

absent, distractor event present or absent). To confirm that the act of making a

response did not contaminate the SSVEP results, we planned to compare SSVEPs for

each of the 4 sub-conditions (the 4 possible combinations of target present/absent and

distractor present/absent). We predicted that the main SSVEP attention effect (entire

stimulus period) would not be different across these 4 conditions. But, if we found an

effect of target or distractor presence on the main SSVEP attention effect, then we

planned to use only the trials without target or distractor events for the time-course

analyses (as has been done in prior work, e.g., Andersen & Müller, 2010).

Checking if the distance between the target and distractor color alters
results. We planned to test whether the magnitude and time-course of attentional
selection differs as a function of target-distractor color similarity (~72° vs. ~144°
separation between the attended and ignored color). Prior work found that it was more
difficult to simultaneously attend opposite colors (180° apart) than to simultaneously
attend two moderately-spaced colors (60° apart) (Chapman et al., 2019; Geweke et al.,
2018; Störmer & Alvarez, 2014). Given this result, we predicted that it should be easier
to suppress a color that is drastically different from the target color (and behavioral
accuracy should likewise be higher for the 144° condition).

Additional non pre-registered SSVEP and ERP control analyses.

We did not anticipate our failure to find an overall attention effect with these task
procedures and set of pre-registered "sanity check" analyses described above. To
further understand the lack of SSVEP attention effect, we performed additional non-pre-
registered control analyses.

Positive control: Frequency analysis of the photodiode voltage. During the
recording, a photodiode was used to ensure that the flicker frequencies were faithfully
presented, and the photodiode recorded voltage fluctuations induced by a small white
dot that flickered at the to-be-attended target frequency on each trial. The electrical
activity from the photodiode was recorded as an additional "electrode" in the data matrix
(with the structure: trials x electrodes x timepoints). Thus, we performed a fast Fourier
transform (FFT, Matlab function "fft.m") to ensure that the trial indexing and FFT aspects
of our analysis were correct. If these aspects of the analysis were correct, we should
expect a near-perfect modulation of photodiode FFT amplitude by attention condition as
only the flicker frequency of the attended stimulus was tagged. For 1 subject, the
photodiode was not plugged in (leaving 22 subjects for this analysis).

Analysis control: Using a more similar frequency analysis procedure to
prior published work. Because we were interested in characterizing a time-course
effect, we chose to use a Gaussian wavelet procedure to quantify power for each
frequency of interest. However, given the lack of overall attention effect, we were not

able to meaningfully look at the time-course effects. Thus, we additionally used a fast

Fourier transform (FFT) to measure SSVEP amplitude during the entire stimulus period.

This method is more commonly used in prior published work (e.g., Andersen et al.,

2008; Andersen & Müller, 2010). Following prior work, we used the entire stimulus

epoch starting from 500 ms onward (500 ms – 2000 ms), we detrended the data, and

we zero-padded this time window (2,048 points) to precisely estimate our frequencies.

465   **Positive control: Analysis of event-related potential (P3) for an attention**

effect. Prior work has found that attention-related SSVEP modulations are

accompanied by changes to event-related potentials (ERPs) associated with target

selection and processing. To measure the P3, we calculated event-related potentials

time-locked to the target or distractor onset (baselined to -200 ms to 0 ms relative to

target onset). We included trials where there was only one target or distractor event on

that trial, to avoid the possibility of overlap between the two signals. We calculated P3

voltage at electrodes Pz and POz during the time window 450-700 ms after target

onset, similar to prior work (Adamian et al., 2019; Andersen et al., 2013; Andersen,

Fuchs, et al., 2011).

476   **Results**

477   **Behavior**

478   Participants were overall accurate at the task (percent correct = 65.7%, d-prime =

1.25), and were significantly above chance (percent correct >50%, $p < .001$; d-prime >0,

$p < .001$). There was no overall significant effect of cue condition (attend cue versus

ignore cue) on performance, $p = .85$, but analysis of target-present trials suggested that

participants could more quickly use attend cues than ignore cues ($p < .05$ when the

target appeared between 0 ms and 275 ms, but $p > .05$ if the target appeared after 275

ms; Appendix B).

485   Average percent correct was lower than our pre-specified target range of 70-

85%, meaning that most participants saw targets and distractors that were maximally

coherent (80% iso-oriented lines) for the majority of blocks (mean coherence of the

target/distractor events = 73.7%, SD = 5.08%). Although overall accuracy was slightly

out of the range we had expected when planning the study, participants were well

above chance and they saw targets with coherence values typical of prior work

(Andersen et al., 2008; Andersen, Fuchs, et al., 2011; Andersen & Müller, 2010).

Pre-registered SSVEP results

We first confirmed that our SSVEP procedure was effective at eliciting robust,

frequency-specific modulations of the EEG signal. After collecting the first 5 participants,

we checked that overall SSVEP amplitudes for our two target frequencies (24 and 30

Hz) were robust when collapsed across conditions (Fig 2A) before proceeding with data

collection. We indeed found that the SSVEP signal was robust during the stimulus

period even with n=5 for both the 24 Hz frequency (mean SNR = 4.45, SD = .14, SNR >

1: $p < .001$) and for the 30 Hz frequency (mean SNR = 2.97, SD = .15, SNR > 1: $p <$

.001). These values were similar for the full n=23 sample (Fig 2B). To compute an effect

size, we compared SNR values for each target frequency (24 Hz and 30 Hz) to the SNR

values for each baseline frequency (frequencies from 3-33 Hz not within +/-2 Hz of 24 or

30 Hz). SNR values for the target frequencies were significantly higher than baseline,

mean Cohen's d = 5.10 (SD = 1.11) and 5.99 (SD = 2.66), respectively (See Appendix

C). As planned, we also confirmed that the electrodes we selected *a priori* (O1, Oz, and

O2) were reasonable given the topography of overall SSVEP amplitudes (i.e., they fell

approximately centrally within the brightest portion of the heat map; Figure 2C-D).

Figure 2. SSVEP amplitude at the expected frequencies (collapsed across all experimental conditions). (A) Power as a function of frequency during the stimulus period at electrodes O1, O2, and Oz for the first 5 participants. As expected, we observed robust peaks at the stimulated frequencies (24 and 30 Hz). (B) Power as a function of frequency for the stimulus period for the full sample (n=23). (C-D) Topography of signal to noise ratio values for 24 Hz (C) and 30 Hz (D) for all participants collapsed across all experimental conditions. Color scale indicates SNR. As expected, the *a priori* electrodes O1, O2, and Oz (magenta circles) showed robust SNR during the stimulus period.

Next, we checked for a basic attention effect, defined as a larger amplitude response evoked by the attended frequency compared to the ignored frequency). Note, for the sake of clarity, all conditions are translated into "attend" terminology. That is, if a participant was cued to "ignore blue" (24 Hz) during the "ignore cue" condition (and the other color was red and 30 Hz), this will instead be plotted as "attend red" (30 Hz). Figure 3 shows the Gaussian wavelet-derived frequency spectra during the stimulus period (500-2000 ms) as a function of cue type (attend versus ignore) and attended frequency (attend 24 Hz or attend 30 Hz). We found a main effect of measured frequency, whereby SNR was overall higher for 24 versus 30 Hz, $F(1,22) = 57.89$, $p < .001$, $\eta^2_p = .73$. However, we found no main effect of attended frequency ($p = .27$) or

cue type (*p* = .83), and we found no significant interactions (*p* >= .18). Collapsed down

to a paired t-test, the observed effect size for attended versus unattended SNR values

was Cohen's *d* = .03. To detect an effect of this size with 80% power (1-β = .8; α = .05)

would require a sample size *n* > 7,000[*]. Given that we did not find an overall attention

effect, we did not analyze or interpret analysis of the SSVEP time-course. However, for

completeness we have shown the time course in Appendix D.

Figure 3. Overall attention effect in the attend cue and ignore cue conditions. (A-
B) Frequency spectra in the attend cue (A) and ignore cue (B) conditions during the
stimulus period. Although we observe expected peaks at 24 Hz and 30 Hz, this SSVEP
response is not modulated by the attention manipulation. (C-D). Violin plots of the
signal-to-noise ratio at the SSVEP frequencies in the attend cue (C) and ignore cue (D)
conditions.

---

[*]As we did not pre-register Bayesian analysis choices (e.g., choices about priors, etc.), we did not
calculate a Bayes Factor for this pre-registered analysis. However, the post-hoc power analysis gives a
sense of the degree to which this is a null effect.

Although we pre-registered that we would analyze all trials (those with and

without target/distractor events), most prior studies have included only trials without any

target or distractor events in the main SSVEP analysis (e.g., Andersen et al., 2008;

Müller et al., 2006). To ensure that our null result was not due to this analysis choice,

we also planned in our pre-registration to examine the SSVEP attention effect for trials

with and without target and distractor events. When restricting our analysis to only trials

without targets or distractors (25% of the 1440 trials, or 360 trials total before artifact

rejection), we likewise found no attention effect. As before, we found a main effect of

measured frequency (24 > 30 Hz), $p < .001$, but no effect of cue condition ($p = .053$) or

attended frequency ($p = .073$), and, most critically, we found no interaction between

measured frequency and attended frequency ($p = .33$). Frequency spectra for all

combinations of target and distractor presence are shown in Appendices E and F.

Finally, we also pre-registered that we would check whether the similarity of the

target and distractor colors (72 versus 144 degrees apart on a circular color wheel;

Figure 1B) would modulate the SSVEP attention effect. We likewise found that the

similarity of the distractor colors did not significantly modulate the SSVEP response,

and we found no attention effect (interaction of measured frequency and attended

frequency) in either color distance condition ($p >= .26$; Appendix G).

Non-pre-registered control analyses

We conducted additional control analyses to rule out possible sources of our

failure to find an attention effect. First, we examined the photodiode recording to rule out

any failures due to trial indexing. The photodiode measured the luminance of a white

dot that flickered at the attended frequency on each trial. As expected, performing an

FFT on the photodiode time-course thus yielded near-perfect tracking of the attended

frequency (Figure 4A-B, $p < .001$). On the other hand, we again found null results for the

main attention manipulation (Figure 4C-F) when using an FFT analysis that more

closely followed prior work. We ran a repeated measures ANOVA on the signal to noise

ratio values during the stimulus period, including the factors Measured Frequency (24

Hz, 30 Hz), Attended Frequency (24 Hz, 30 Hz), and Cue Type (Attend, Ignore). We

found no main effect of measured frequency (p = .91), attended frequency (p = .45), or

cue type (p = .54), and we found no significant interactions (p >= .38). However, the

average signal-to-noise ratio of the stimulus frequencies was overall robust (M = 4.45,

SD = 1.45, greater than chance value of 1: $p < 1\times10^{-9}$), so our inability to observe the

attention effect was not due to lack of overall SSVEP signal. Likewise, we ran an

additional analysis to ensure that our choice of pre-registered choice of SNR measure

did not explain our null result (Appendix H).

Given that some work has reported significant effects only for the second

harmonic (e.g., Kim et al., 2007; Vissers et al., 2017), we likewise examined SSVEP

amplitude at 48 Hz and 60 Hz, with the caveat that the 60 Hz harmonic is contaminated

by line noise (Appendices I and J). We found no significant attention effects for either

second harmonic frequency. We also re-ran the FFT analysis with other electrode-

selection methods to ensure our *a priori* choice of electrodes did not impede our ability

to observe an effect. We found no evidence that electrode choice led to our null effect,

as exploiting information from all 64 electrodes by implementing rhythmic entrainment

source separation (RESS) likewise yielded null effects (Appendices K and L; M. X.

Cohen & Gulbinaite, 2017). To ensure that inconsistent task performance did not lead to

null effects, we repeated the main FFT analysis on only accurate trials. We likewise

found null attention effects when analyzing only accurate trials (Appendix M).

Finally, we tested whether phase consistency, rather than power, may track

attention in our task (e.g., Nunez et al., 2015; Tallon-Baudry et al., 1996). To do so, we

performed an FFT on single trials rather than on condition-averaged waveforms, and we

extracted single-trial phase values. We calculated a phase-locking index by computing

mean-resultant vector length on each condition's histogram of single-trial phase values.

Mean-resultant vector length ranges from 0 (fully random values) to 1 (perfectly identical

values), for reference, see Zar (2010). We found no effect of attention on this phase-

locking index (Appendix N).

Figure 4. FFT analysis of the photodiode and SSVEPs at attended and unattended frequencies during the stimulus period (500-2000 ms). (A-B) As a positive control for our analysis pipeline, we performed an FFT analysis on the photodiode trace. The photodiode recorded a flickering white dot at the attended frequency on each trial. As expected, this provides a near-perfect tracking of the attended frequency in both the attend cue condition (A) and the ignore cue condition (B). (C-F) To ensure our null effect was not due to using Gaussian wavelets rather than an FFT, we repeated the main analysis with an FFT. Frequency spectra for the attend cue condition (C) and ignore cue condition (D) reveal an overall robust SSVEP signal at 24 Hz and 30 Hz, but no modulation by attention. Likewise, violin plots of signal-to-noise ratios again show robust signal but no modulation by attention in either the attend cue condition (E) or the ignore cue condition (F).

617
Positive control: Analysis of event-related potential (P3b) for an attention
effect. Consistent with prior work, we found a significantly larger P3 component for
target onsets compared to distractor onsets (Figure 5). A repeated measures ANOVA
with within-subjects factors cue type (attend cue or ignore cue) and event type (target or
distractor onset) revealed a robust main effect of event type (target > distractor), $F(1,22)$
$= 51.64$, $p < 1 \times 10^{-5}$, $\eta^2_p = .70$, and a main event of cue type (attend > ignore), $F(1,22) =$
4.96, $p = .037$, $\eta^2_p = .18$, but no interaction between event type and cue type ($p = .65$).
Control analyses confirmed this P3 modulation was not due to differential rates of
making a motor response for targets and distractors (Appendix O). The main effect of
event type (target > distractor) remained when analyzing only trials where participants
made a motor response ($p < .001$). Thus, the P3 was overall larger for target than
distractor events, consistent with prior work that found this ERP attention effect
alongside an SSVEP attention effect.

631
632   **Figure 5. P3 amplitude at electrodes Pz and POz.** (A) P3 amplitude in the attend cue
633   condition, as a function of whether the event onset (iso-oriented lines) was a target that
634   should be reported or a distractor that should be ignored. (B) P3 amplitude in the ignore
635   cue condition. Shaded error bars represent standard error of the mean; the gray
636   rectangle indicates the time period used for the statistical tests.

Focused review of feature-based attention studies using SSVEPs

Given that our results are inconsistent with prior work, we conducted a focused review

to try to pinpoint critical methodological differences that may have led to our failure to

replicate a basic attention effect on SSVEP amplitude in this specific task. To do so, we

first read review papers to identify an initial set of empirical studies employing a feature-

based attention manipulation and SSVEPs (Andersen, Müller, et al., 2011; Norcia et al.,

2015; Vialatte et al., 2010). From this initial set of papers, we used Google Scholar to

check citations and citing papers for mention of the terms feature-based attention and

SSVEPs. Our inclusion criteria included: (1) published journal article (2) healthy young

adults (3) SSVEPs were measured from either an EEG or MEG signal and (4) a feature-

based attention manipulation was included.

We defined "feature-based attention manipulation" as having the following

characteristics: (1) Participants were cued to attend a feature(s) within a feature

dimension (e.g., attend red, ignore blue) rather than across a feature dimension (e.g.

attend contrast, ignore orientation), (2) The attended and ignored feature were both

frequency-tagged in the same trials (rather than only 1 feature tagged per trial), (3)

Each frequency was both "attended" and "ignored" on different trials, so that the

amplitude of a given frequency could be examined as a function of attention, (4) The

task could not be performed by adopting a strategy of splitting spatial attention to

separate spatial locations.

After applying these screening criteria, some of the studies that we initially

identified were excluded (brackets indicate exclusion reason(s)): Appelbaum & Norcia,

2009 [1,4]; Boylan et al., 2019 [3]; Bridwell & Srinivasan, 2012 [2,3]; Clementz et al.,

2008 [3]; Garcia et al., 2013 [1,4]; Hasan et al., 2017 [2]; Itthipuripat et al., 2019 [1];

Talsma et al., 2006 [1,4]; Thigpen et al., 2019 [4]; Verghese et al., 2012 [1,4]).

We identified a total of 34 experiments from 28 unique papers (Appendices P-S)

meeting our inclusion criteria. From these experiments, we quantified variables such as

the number of subjects, number of trials, frequencies used, and the presence or

absence of an attention effect in the expected direction (attended > ignored). If more

than one group of participants was used (e.g., an older adults group) then we included

the study but only quantified results for the healthy young adult group (Quigley et al.,
2010; Quigley & Müller, 2014).

Task used in each study.

The tasks used in these studies fell broadly into one of 4 categories: (1) a
competing gratings task, (2) a whole-field flicker task, (3) a hemifield flicker task and (4)
a central task with peripheral flicker.

In the competing gratings task (Appendix P), participants viewed a stream of
centrally-presented, superimposed gratings (e.g., a red horizontal grating and a green
vertical grating). Because colored, oriented gratings were typically used, participants
could thus generally choose to attend based on either one or both features (color and/or
orientation). Each grating flickered at its own frequency (e.g. green grating shown at
7.41 Hz, red grating shown at 8.33 Hz, as in Chen et al., 2003). Because the gratings
were superimposed, on any given frame only one of the two gratings was shown. On
frames where both gratings should be presented according to their flicker frequencies, a
hybrid "plaid" stimulus was shown. Studies using a competing gratings task include:
(Allison et al., 2008; Chen et al., 2003; Keitel & Müller, 2016; J. Wang et al., 2007).

In the whole-field flicker task (Appendix Q), participants viewed a spatially global
stimulus comprised of small, intermingled dots or lines. Typically, half of the dots or
lines were presented in one feature (e.g., red) and the other half of the lines were
presented in another (e.g., blue); each set of dots flickered at a unique frequency.
Although the most common attended feature was color, some task variants included (1)
attending high or low contrast stimuli (2) attending a particular orientation or (3)
attending a particular conjunction of color and orientation. The whole-field flicker task
was the most common task variant, and it is also most similar to the task performed
here. Studies using a whole-field flicker task include: (Andersen et al., 2008, 2009,
2012, 2015; Andersen & Müller, 2010; Forschack et al., 2017; Martinovic et al., 2018;
Martinovic & Andersen, 2018; Müller et al., 2006; Quigley et al., 2010; Quigley & Müller,
2014; Steinhauser & Andersen, 2019; D. Zhang et al., 2010)

In the hemifield flicker task, participants viewed a stimulus within each hemifield, and

each stimulus was comprised of small, intermingled dots or lines. Often, these studies

included both a feature-based attention manipulation and a spatial attention

manipulation (e.g., attend red on the left-hand side). However, the feature-based

attention task could not be achieved with spatial attention alone, as participants needed

to attend to a particular color and ignore a distractor color within the attended hemifield.

In addition, the unattended hemifield could often be used to track the spatially global

spread of feature-based attention. Studies using a hemi-field flicker task include:

(Adamian et al., 2019; Andersen et al., 2013; Andersen, Fuchs, et al., 2011; Müller et

al., 2018; Störmer & Alvarez, 2014)

Finally, in the central task with peripheral flicker, participants performed a task

near fixation (e.g., visual search), and feature-based attention was measured indirectly

via a peripheral flickering stimulus (i.e., this design took advantage of the spatially

global spread of feature-based attention, Sàenz et al., 2002, 2003; White & Carrasco,

2011). Studies using a central task with peripheral flicker include: (Chu & D'Zmura,

2019; Jiang et al., 2017; Painter et al., 2014, 2015).

Sample Size, Trial Counts, and Stimulus Duration

First, we examined whether insufficient power could have led to our failure to

detect an attention effect. Both sample size and the number of trials per condition are

critical for determining power (Baker et al., 2019; Boudewyn et al., 2018; Button et al.,

2013; Button & Munafò, 2017; Clayson et al., 2019; Thigpen et al., 2017). The number

of studies employing each task variant is plotted in Figure 6A, the number of subjects

per experiment is plotted in Figure 6B, the number of trials per experiment is plotted in

Figure 6C, and stimulus duration is plotted in Figure 6D. Bars are color coded according

to whether each experiment overall found an expected attention effect (attended >

ignored), a reverse attention effect (ignored > attended), mixed results across conditions

within the experiment, or ambiguous results (3 studies: Pei et al., 2002: reported

statistics for the harmonics but not the fundamental frequency; Allison et al., 2008:

missing formal statistical tests; Martinovic et al., 2018: statistical tests measured if

attention effect differed between conditions, but did not formally test that the attention

effect was overall significant). Our study had N = 23, 1,440 total trials per subject (720

trials per cue condition), and a stimulus duration of 2 seconds. In comparison to the

literature, these factors are unlikely to explain our failure to find an effect. On average,

prior studies had a median sample size of N = 15 (SD = 4.1, min = 9, max = 23) and a

median trial count of 440 (SD = 314.2, min = 8, max =1600) and a median stimulus

duration of 4.1 sec (SD = 37.35, min = 1 sec, max = 120 sec). Likewise, our trial counts

were reasonable when we restricted our analysis to trials without targets or distractors

(360 trials total; 180 trials per condition) relative to this estimated value for the literature

(median = 96 trials per condition, SD = 95.6, min = 4, max = 400). Upon initially plotting

the data, we noticed that studies using the competing gratings task produced

inconsistent results, with as many experiments showing a mixture of effects across

conditions, overall reversed effects (ignored > attended), and null effects (2 studies in

each category). We think this inconsistency is most likely due to the below-average trial

counts for these studies (median of only ~8 trials) (Boudewyn et al., 2018).

Figure 6. Study characteristics from the literature review of feature-based
attention and SSVEPs. (a) Number of experiments in each of the 4 task variants.
Different colors in the stacked bar graphs indicate whether the experiment found an

expected attention effect (attended > ignored), reverse effect (ignored > attended),
mixed effects across conditions, or a null / ambiguously reported effect. (b) Number of
participants. (c) Number of trials in the experiment (before artifact rejection or excluding
trials with targets). (d) Stimulus duration. (e) Percentage of trials, on average, where the
attended feature was repeated on the next trial. (f) Stimulation frequencies.
755
Percentage of trials where an attended feature was repeated.

Next, we examined the percentage of trials where the attended feature was

repeated (e.g., if the attended color was red on trial n, what was the chance that red

would also be attended on trial $n+1$?). The priming-based account of feature-based

attention posits that participants cannot use trial-by-trial cues to enhance a particular

feature, but rather, feature-based enhancement happens automatically when a

particular feature is repeated (Theeuwes, 2013). Thus, if there is a substantial

proportion of trials where the repeated color was attended (e.g. with 2 possible colors,

both the attended and ignored color will be repeated on 50% of trials), then the

observed attentional enhancement effects might be driven primarily by incidental

repetitions of attended features. In our study, we used 5 different colors to reduce the

potential effect of inter-trial priming on the observed SSVEP attention effects (20%

repeats of the attended color, 4% repeats of the attended color and the ignored color).

We quantified the approximate percentage of trials on which an attended feature on one

trial is repeated on the next trial (within a given block of trials). In some studies,

participants were cued to attend more than one feature on a given trial, or they

sometimes attended to a conjunction of features. In these cases, we calculated the

expected number of repeats for either of the 2 attended features based on the total

number of conditions (Andersen et al., 2008, 2013, 2015; Martinovic et al., 2018).

Figure 6E shows a histogram of the percentage of trials that repeated an

attended feature. In two studies, the to-be-attended color was held constant on each

block (100% repeats (Jiang et al., 2017; Painter et al., 2014). In the majority of the

remaining studies, only two unique features were used so the percentage of trials where

the attended feature was repeated was on average quite high (overall median = 50%,

SD = 27.4%, min = 0%, max = 100%). Finally, in three studies, the attention conditions

were perfectly alternated (0% repeat trials; Allison et al., 2008; Chen et al., 2003; J.

Wang et al., 2007). Consistent with a priming account, 81% of the studies with a high

percentage of repeats showed a consistent positive attention effect, whereas none of

the studies with 0% repeat trials showed a consistent positive attention effect. However,

we think the inconsistent effects in the studies with 0% repeats might be equally

attributed to their low trial counts (median = 8 trials per condition; Boudewyn et al.,

2018). Only one study had a similar proportion of repeats as the present study (Störmer

& Alvarez, 2014). Störmer and Alvarez found a significant attention effect while using 5

unique colors (intermixed randomly from trial to trial). The findings by Störmer and

Alvarez provide evidence against the feature-based priming account, and suggest the

task factor "number of colors" cannot definitively explain our inability to observe an

attention effect. However, given the lack of extant work using unpredictable color cues,

we think future, systematic work is needed to determine the degree to which inter-trial

priming effects may modulate the size and reliability of feature-based attention effects.

SSVEP Frequencies.

We examined frequencies that have been most commonly used in the literature.

In our study, we chose relatively high frequencies (24 and 30 Hz) in order to have

increased temporal resolution for detecting potential time-course effects. In addition,

some have argued that using higher frequencies as advantages for driving a more

localized portion of visual cortex, as opposed to broadly driving visual, parietal and

frontal cortex when using lower frequencies in the theta/alpha bands (i.e., ~7-12Hz;

Ding et al., 2006; Srinivasan et al., 2006). For the purposes of temporal resolution,

earlier work examining the time course of *spatial* attention with SSVEPs used the

frequencies 20 and 28 Hz, (Müller et al., 1998). However, upon reviewing the feature-

based attention literature, we found that our chosen frequencies were outside the range

that has previously been used with a feature-based attention task (Figure 6F; median =

10 Hz, SD = 3.6 Hz, min = 2.4 Hz, max = 19.75 Hz). Thus, it is possible that feature-

based attention, unlike spatial attention, cannot be easily tracked with flicker

frequencies above ~20 Hz. Although some studies have argued that feature-based

attention effects do not qualitatively appear to vary with frequency (Martinovic et al.,

2018; Steinhauser & Andersen, 2019), the vast majority of reviewed studies only

reported statistical significance of an overall attention effect collapsed across

frequencies. Only a handful studies have reported statistical significance of individual

frequencies (e.g., Chu & D'Zmura, 2019; Painter et al., 2014; Quigley & Müller, 2014;

Steinhauser & Andersen, 2019). As such, future work is needed to systematically

investigate the effect of frequency choice on feature-based attention effects, particularly

for frequencies >20 Hz.

820   **Task Type and Task Difficulty**

Finally, we examined whether the type of task and task difficulty may have

influenced our ability to detect an attention effect. In particular, the specific targets that

we used may differ slightly from prior work. In our experiment, participants detected a

brief period (333 ms) of an on average ~75% coherent orientation (the coherent line

orientation was a random, unpredictable direction, from 1-180 degrees). In this task,

participants performed well above chance, but the task was still fairly challenging overall

(d' = 1.25). This raises the possibility that, compared to prior SSVEP studies, subjects

were giving up on some percentage of the trials and that this contributed to the lack of

attention effects.

For the reviewed papers in which participants detected a target within the

flickering stimulus ("whole-field flicker task" and "hemifield flicker task"), we compiled

information about participants' accuracy, the duration of the target, the type of target,

and the percentage of dots/lines that comprised the target (Table S5). We found that

our particular task (detect a coherent orientation in the cued color) was slightly different

from the other tasks that have been used. Two other prior studies did not use a

behavioral task at all: participants were simply instructed to monitor a particular feature

without making any overt response (Pei et al., 2002; D. Zhang et al., 2010). In three

papers, participants attended to a brief (200 ms) luminance decrement in 20% of the

attended dots (Adamian et al., 2019; Andersen et al., 2009, 2013). In the remaining

papers, participants monitored for a brief coherent motion event (230 ms – 500 ms) in

50-85% of the attended dots/lines (Andersen et al., 2008, 2012; Andersen & Müller,

2010; Forschack et al., 2017; Martinovic et al., 2018; Martinovic & Andersen, 2018;

Müller et al., 2006, 2018; Quigley et al., 2010; Quigley & Müller, 2014; Steinhauser &

Andersen, 2019; Störmer & Alvarez, 2014).

Although the particulars of the luminance and motion tasks subtly differ from our

orientation task, it is not clear why SSVEPs would track attention when the target is a

coherent luminance value or motion direction, but not when the target is a coherent

orientation. For example, just like in the coherent motion direction tasks used by others,

the angle of the coherent orientation in our task was completely unpredictable. Thus,

participants in our task and in other tasks could not form a template of an orientation or

motion direction they should attend in advance and instead had to attend to an

orthogonal feature dimension such as color. In addition, in both prior tasks and the

current task there were an equal number of coherent events in the cued and uncued

color. If participants failed to attend to the cued color and instead responded to any

orientation event, their performance in the task would be at chance.

Behavioral performance in the reviewed studies ranged from $d' = 0.8$ to $d' = 3.25$

(Appendix T). In many studies, performance was quite high ($d' > 2$ or accuracy > 90%)

relative to performance in our study (Adamian et al., 2019; Andersen et al., 2008, 2009,

2012, 2013; Forschack et al., 2017; Müller et al., 2006; Quigley et al., 2010; Quigley &

Müller, 2014; Steinhauser & Andersen, 2019). However, there were several studies

where the authors found SSVEP attention effects despite overall lower behavioral

performance values more comparable to our study (d' between 1 and 1.5; Andersen et

al., 2015; Martinovic et al., 2018; Martinovic & Andersen, 2018). Sometimes, a more

difficult task may actually be associated with increased attention effects: Martinovic and

Andersen (2018) observed attention effects that were stronger in the subset of

conditions with lower behavioral performance ($d' = .8 - 1.5$) compared to conditions that

were easier ($d' > 2.25$).

869   **Discussion**

In this pre-registered study, we sought to test whether cuing participants to

ignore a particular color modulates the time-course of feature-based attention as

indexed by steady-state visually evoked potentials (SSVEPs). As a baseline point of

comparison, we also included a condition in which participants were cued to *attend* a

particular color. This "attend cue" condition was intended as a close replication of much

prior work showing that SSVEP amplitudes are modulated by attention (greater

amplitude for the attended feature; e.g., Andersen et al., 2008; Andersen & Müller,

2010; Müller et al., 2006). However, we failed to replicate this basic overall attention

effect; we found no difference in SSVEP amplitude as a function of attention in either

the attend cue or the ignore cue condition. Thus, because we found no overall SSVEP

attention effect, we were unable to test our hypotheses about how this attention effect

was modulated by being cued to attend versus cued to ignore. Despite the lack of an

SSVEP attention effect, positive control analyses indicated that that participants did

successfully select the cued target color (i.e., we observed a significantly larger P3

component for target events in the attended color than in the ignored color).

Given our failure to observe an effect of attention on SSVEP amplitude with our

task procedures, we performed a focused review of the literature to quantify key

methodological aspects of prior studies using SSVEPs to study feature-based attention.

Based on this review, we concluded that sample size and trial counts likely did not

explain our failure to find an effect; our sample size and trial counts were near the

maximum values found in the surveyed literature. Likewise, the range of accuracy

values found in the literature suggests that task difficulty does not explain our failure to

find an attention effect. However, two key, intentional design differences may have

hampered our ability to find an effect: (1) the number of colors in our stimulus set and

(2) the frequencies used to generate the SSVEP.

The first key design difference in our study was the number colors in our stimulus

set. We purposefully minimized the influence of inter-trial priming on our estimates of

feature-based attention (Theeuwes, 2013) by using 5 unique colors and randomly

choosing target and distractor colors on each trial. According to a priming account of

feature-based attention, a relatively high proportion of trials where the attended color is

repeated back-to-back could inflate or even entirely drive apparent feature-based

attention effects. Using 5 colors somewhat protects against this possibility, because it

ensures that the attended color is repeated on 20% of trials, and both the
attended/ignored colors are repeated on only 4% of trials. In the literature, we found that
most studies had back-to-back color repeats on at least 50% of trials. It is thus plausible
that inter-trial priming could contribute to observed attention differences in these
studies. Contrary to a priming account, however, one study found robust feature-based
attention effects using a set of 5 unique colors (Störmer & Alvarez, 2014), suggesting
that participants can use a cue to direct feature-based attention even when the
proportion of repeated trials is relatively low. To date, however, no study has directly
manipulated the proportion of repeated trials or the number of possible stimulus colors
in an SSVEP study. Given emerging evidence that history-driven effects play an
important role in shaping both spatial and feature-based attentional selection (Adam &
Serences, 2020; Awh et al., 2012; Failing et al., 2019; Geng et al., 2019; Kadel et al.,
2017; B. Wang & Theeuwes, 2018a, 2018b; B.-Y. Won & Geng, 2020), we think that
future work is needed to directly investigate whether and to what degree SSVEP
estimates of feature-based attention are modulated by inter-trial priming.

The second key design difference in our study was the chosen set of
frequencies. To ensure adequate temporal resolution to characterize time-course
effects, we chose to use slightly higher frequencies (24 and 30 Hz). We believed these
values would be reasonable, because an initial study of the time-course of spatial
attention used SSVEP frequencies in a similar range (20 and 28 Hz; Müller et al., 1998).
In addition, frequencies in the beta band (~15-30 Hz) have commonly been used in
other SSVEP studies of spatial attention (Garcia et al., 2013; Kashiwase et al., 2012;
Müller, Picton, et al., 1998; Müller & Hillyard, 2000; Toffanin et al., 2009; D.-O. Won et
al., 2016), and SSVEPs are overall robust using a wide array of frequencies (at least 1
to 50 Hz; Herrmann, 2001; Zhu et al., 2010). However, some spatial attention studies
have found no attentional modulation of SSVEPs in the beta band (Antonov et al.,
2020; Gulbinaite et al., 2019), or have found effects only for the second harmonic of
beta band frequencies (Garcia et al., 2013; Kim et al., 2007; Vissers et al., 2017).
Further, the SSVEP amplitude, estimated spatial extent of the SSVEP signal, and the
size of spatial attention effects vary with frequency (Ding et al., 2006; Gulbinaite et al.,

2019; Herrmann, 2001). Given differences in the cortical processing of locations and

features (M. R. Cohen & Maunsell, 2011; Haxby et al., 1994; Kastner & Ungerleider,

2000; Mishkin & Ungerleider, 1982; Owen et al., 1996), and differences in SSVEP

spatial extent and strength with frequency (Ding et al., 2006; Gulbinaite et al., 2019;

Lithari et al., 2016), it is plausible that feature-based attention can only be tracked with a

limited range of frequencies (e.g., frequencies near the alpha band). Future work will be

needed to systematically investigate the effect of SSVEP frequency on feature-based

attention.

It is perhaps puzzling that frequencies above 20 Hz have been commonly used in

the spatial attention literature but have not been used in the feature-based attention

literature. The truncation of the frequency distribution in the reviewed literature could be

a piece of the "file drawer" in action. It is possible that other researchers likewise

discovered that they were unable to track feature-based attention using certain

frequencies, but that these null results were never published due to journals' and

authors' biases toward publishing positive results (Cooper et al., 1997; Dickersin, 1990;

Dickersin et al., 1992; Dwan et al., 2008; Ferguson & Heene, 2012; Franco et al., 2014;

Rosenthal, 1979) and biases against publishing negative results (i.e., "censoring of null

results", Guan & Vandekerckhove, 2016; Sterling, 1959; Sterling et al., 1995). Thus, our

results highlight the practical and theoretical importance of regularly publishing null

results. On the practical side, if prior null results had been published, we may have

better known which frequencies to use or avoid, and we would have been able to test

our key hypotheses. On the theoretical side, our results highlight how seemingly

unimportant null results can have implications for theory when viewed in the context of

the broader literature. For example, if certain frequencies track spatial but not feature-

based attention, this may inform our understanding of the brain networks and cognitive

processes differentially modulated by flicker frequency (Ding et al., 2006; Srinivasan et

al., 2006).

In short, we found no evidence that SSVEPs track the deployment of feature-

based attention with our procedures, and future methodological work is needed to

determine constraints on generalizability of the SSVEP method for tracking feature-

based attention. We performed a focused review of prior studies using SSVEPs to study

feature-based attention, and from this review we identified two key factors (frequencies

used; likelihood of inter-trial feature priming) that should be systematically investigated

in future work.

References

966   Adam, K. C. S., & Serences, J. T. (2020). *History-driven modulations of population codes in*
967   *early visual cortex during visual search* [Preprint]. bioRxiv.
968   https://doi.org/10.1101/2020.09.30.321729
969   Adamian, N., Slaustaite, E., & Andersen, S. K. (2019). Top–Down Attention Is Limited Within but
970   Not Between Feature Dimensions. *Journal of Cognitive Neuroscience*, *31*(8), 1173–
971   1183. https://doi.org/10.1162/jocn_a_01383
972   Adrian, E. D., & Matthews, B. H. C. (1934). The Berger Rhythm: Potential Changes from the
973   Occiptial Lobes in Man. *Brain*, *57*(4), 355–385.
974   Allison, B. Z., McFarland, D. J., Schalk, G., Zheng, S. D., Jackson, M. M., & Wolpaw, J. R.
975   (2008). Towards an independent brain–computer interface using steady state visual
976   evoked potentials. *Clinical Neurophysiology*, *119*(2), 399–408.
977   https://doi.org/10.1016/j.clinph.2007.09.121
978   Andersen, S. K., Fuchs, S., & Müller, M. M. (2011). Effects of Feature-selective and Spatial
979   Attention at Different Stages of Visual Processing. *Journal of Cognitive Neuroscience*,
980   *23*(1), 238–246. https://doi.org/10.1162/jocn.2009.21328
981   Andersen, S. K., Hillyard, S. A., & Müller, M. M. (2008). Attention Facilitates Multiple Stimulus
982   Features in Parallel in Human Visual Cortex. *Current Biology*, *18*(13), 1006–1009.
983   https://doi.org/10.1016/j.cub.2008.06.030
984   Andersen, S. K., Hillyard, S. A., & Müller, M. M. (2013). Global Facilitation of Attended Features
985   Is Obligatory and Restricts Divided Attention. *Journal of Neuroscience*, *33*(46), 18200–
986   18207. https://doi.org/10.1523/JNEUROSCI.1913-13.2013
987   Andersen, S. K., & Müller, M. M. (2010). Behavioral performance follows the time course of
988   neural facilitation and suppression during cued shifts of feature-selective attention.
989   *Proceedings of the National Academy of Sciences*, *107*(31), 13878–13882.
990   https://doi.org/10.1073/pnas.1002436107
991   Andersen, S. K., Müller, M. M., & Hillyard, S. A. (2009). Color-selective attention need not be
992   mediated by spatial attention. *Journal of Vision*, *9*(6), 2–2. https://doi.org/10.1167/9.6.2
993   Andersen, S. K., Müller, M. M., & Hillyard, S. A. (2011). Tracking the allocation of attention in
994   visual scenes with steady-state evoked potentials. In M. I. Posner (Ed.), *Cognitive*
995   *Neuroscience of Attention* (2nd ed.). The Guilford Press.
996   Andersen, S. K., Müller, M. M., & Hillyard, S. A. (2015). Attentional Selection of Feature
997   Conjunctions Is Accomplished by Parallel and Independent Selection of Single Features.
998   *Journal of Neuroscience*, *35*(27), 9912–9919. https://doi.org/10.1523/JNEUROSCI.5268-
999   14.2015
1000 Andersen, S. K., Müller, M. M., & Martinovic, J. (2012). Bottom-Up Biases in Feature-Selective
1001   Attention. *Journal of Neuroscience*, *32*(47), 16953–16958.
1002   https://doi.org/10.1523/JNEUROSCI.1767-12.2012

Antonov, P. A., Chakravarthi, R., & Andersen, S. K. (2020). Too little, too late, and in the wrong place: Alpha band activity does not reflect an active mechanism of selective attention. *NeuroImage*, *219*, 117006. https://doi.org/10.1016/j.neuroimage.2020.117006

Appelbaum, L. G., & Norcia, A. M. (2009). Attentive and pre-attentive aspects of figural processing. *Journal of Vision*, *9*(11), 18–18. https://doi.org/10.1167/9.11.18

Arita, J. T., Carlisle, N. B., & Woodman, G. F. (2012). Templates for rejection: Configuring attention to ignore task-irrelevant features. *Journal of Experimental Psychology: Human Perception and Performance*, *38*(3), 580–584. https://doi.org/10.1037/a0027885

Awh, E., Belopolsky, A. V., & Theeuwes, J. (2012). Top-down versus bottom-up attentional control: A failed theoretical dichotomy. *Trends in Cognitive Sciences*, *16*(8), 437–443. https://doi.org/10.1016/j.tics.2012.06.010

Baker, D. H., Vilidaite, G., Lygo, F. A., Smith, A. K., Flack, T. R., Gouws, A. D., & Andrews, T. J. (2019). Power contours: Optimising sample size and precision in experimental psychology and human neuroscience. *ArXiv:1902.06122 [q-Bio, Stat]*. http://arxiv.org/abs/1902.06122

Bartsch, M. V., Loewe, K., Merkel, C., Heinze, H.-J., Schoenfeld, M. A., Tsotsos, J. K., & Hopf, J.-M. (2017). Attention to Color Sharpens Neural Population Tuning via Feedback Processing in the Human Visual Cortex Hierarchy. *The Journal of Neuroscience*, *37*(43), 10346–10357. https://doi.org/10.1523/JNEUROSCI.0666-17.2017

Beck, V. M., & Hollingworth, A. (2015). Evidence for negative feature guidance in visual search is explained by spatial recoding. *Journal of Experimental Psychology: Human Perception and Performance*, *41*(5), 1190–1196. https://doi.org/10.1037/xhp0000109

Becker, M. W., Hemsteger, S., & Peltier, C. (2015). No templates for rejection: A failure to configure attention to ignore task-irrelevant features. *Visual Cognition*, *23*(9–10), 1150–1167. https://doi.org/10.1080/13506285.2016.1149532

Boudewyn, M. A., Luck, S. J., Farrens, J. L., & Kappenman, E. S. (2018). How many trials does it take to get a significant ERP effect? It depends. *Psychophysiology*, *55*(6), e13049. https://doi.org/10.1111/psyp.13049

Boylan, M. R., Kelly, M. N., Thigpen, N. N., & Keil, A. (2019). Attention to a threat-related feature does not interfere with concurrent attentive feature selection. *Psychophysiology*, *56*(6), e13332. https://doi.org/10.1111/psyp.13332

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436. https://doi.org/10.1163/156856897X00357

Bridwell, D. A., & Srinivasan, R. (2012). Distinct Attention Networks for Feature Enhancement and Suppression in Vision. *Psychological Science*, *23*(10), 1151–1158. https://doi.org/10.1177/0956797612440099

Button, K. S., Ioannidis, J. P. A., Mokrysz, C., Nosek, B. A., Flint, J., Robinson, E. S. J., & Munafò, M. R. (2013). Power failure: Why small sample size undermines the reliability of neuroscience. *Nature Reviews Neuroscience*, *14*(5), 365–376. https://doi.org/10.1038/nrn3475

Button, K. S., & Munafò, M. R. (2017). Powering Reproducible Research. In S. O. Lilienfeld & I. D. Waldman (Eds.), *Psychological Science Under Scrutiny* (pp. 22–33). John Wiley & Sons, Inc. https://doi.org/10.1002/9781119095910.ch2

Canolty, R. T., Edwards, E., Dalal, S. S., Soltani, M., Nagarajan, S. S., Kirsch, H. E., Berger, M. S., Barbaro, N. M., & Knight, R. T. (2006). High Gamma Power Is Phase-Locked to Theta Oscillations in Human Neocortex. *Science*, *313*(5793), 1626–1628. https://doi.org/10.1126/science.1128115

Carlisle, N. B., & Nitka, A. W. (2019). Location-based explanations do not account for active
attentional suppression. *Visual Cognition*, *27*(3–4), 305–316.
https://doi.org/10.1080/13506285.2018.1553222
Chang, S., & Egeth, H. E. (2019). Enhancement and Suppression Flexibly Guide Attention.
Psychological Science, *30*(12), 1724–1732. https://doi.org/10.1177/0956797619878813
Chapman, A. F., Geweke, F., & Störmer, V. S. (2019). Feature-based attention resolves
differences in target-distractor similarity through multiple mechanisms. *Journal of Vision*,
19(10), 45a. https://doi.org/10.1167/19.10.45a
Chen, Y., Seth, A. K., Gally, J. A., & Edelman, G. M. (2003). The power of human brain
magnetoencephalographic signals can be modulated up or down by changes in an
attentive visual task. *Proceedings of the National Academy of Sciences*, *100*(6), 3501–
3506. https://doi.org/10.1073/pnas.0337630100
Chu, V. C., & D'Zmura, M. (2019). Tracking feature-based attention. *Journal of Neural
Engineering*, *16*(1), 016022. https://doi.org/10.1088/1741-2552/aaed17
Clayson, P. E., Carbine, K. A., Baldwin, S. A., & Larson, M. J. (2019). Methodological reporting
behavior, sample sizes, and statistical power in studies of event-related potentials:
Barriers to reproducibility and replicability. *Psychophysiology*, *56*(11).
https://doi.org/10.1111/psyp.13437
Clementz, B. A., Wang, J., & Keil, A. (2008). Normal Electrocortical Facilitation But Abnormal
Target Identification during Visual Sustained Attention in Schizophrenia. *Journal of
Neuroscience*, *28*(50), 13411–13418. https://doi.org/10.1523/JNEUROSCI.4095-08.2008
Cohen, M. R., & Maunsell, J. H. R. (2011). Using Neuronal Populations to Study the
Mechanisms Underlying Spatial and Feature Attention. *Neuron*, *70*(6), 1192–1204.
https://doi.org/10.1016/j.neuron.2011.04.029
Cohen, M. X., & Gulbinaite, R. (2017). Rhythmic entrainment source separation: Optimizing
analyses of neural responses to rhythmic sensory stimulation. *NeuroImage*, *147*, 43–56.
https://doi.org/10.1016/j.neuroimage.2016.11.036
Combrisson, E., & Jerbi, K. (2015). Exceeding chance level by chance: The caveat of
theoretical chance levels in brain signal classification and statistical assessment of
decoding accuracy. *Journal of Neuroscience Methods*, *250*, 126–136.
https://doi.org/10.1016/j.jneumeth.2015.01.010
Conci, M., Deichsel, C., Müller, H. J., & Töllner, T. (2019). Feature guidance by negative
attentional templates depends on search difficulty. *Visual Cognition*, *27*(3–4), 317–326.
https://doi.org/10.1080/13506285.2019.1581316
Cooper, H., DeNeve, K., & Charlton, K. (1997). Finding the missing science: The fate of studies
submitted for review by a human subjects committee. *Psychological Methods*, *2*(4), 447–
452. https://doi.org/10.1037/1082-989X.2.4.447
Cunningham, C. A., & Egeth, H. E. (2016). Taming the White Bear: Initial Costs and Eventual
Benefits of Distractor Inhibition. *Psychological Science*, *27*(4), 476–485.
https://doi.org/10.1177/0956797615626564
Dickersin, Kay. (1990). The Existence of Publication Bias and Risk Factors for Its Occurrence.
JAMA: The Journal of the American Medical Association, *263*(10), 1385.
https://doi.org/10.1001/jama.1990.03440100097014
Dickersin, Kay, Min, Y. I., & Meinert, C. L. (1992). Factors influencing publication of research
results. Follow-up of applications submitted to two institutional review boards. *JAMA*,
267(3), 374–378.
Ding, J., Sperling, G., & Srinivasan, R. (2006). Attentional Modulation of SSVEP Power
Depends on the Network Tagged by the Flicker Frequency. *Cerebral Cortex*, *16*(7),
1016–1029. https://doi.org/10.1093/cercor/bhj044

Donohue, S. E., Bartsch, M. V., Heinze, H.-J., Schoenfeld, M. A., & Hopf, J.-M. (2018). Cortical Mechanisms of Prioritizing Selection for Rejection in Visual Search. *The Journal of Neuroscience*, *38*(20), 4738–4748. https://doi.org/10.1523/JNEUROSCI.2407-17.2018

Dwan, K., Altman, D. G., Arnaiz, J. A., Bloom, J., Chan, A.-W., Cronin, E., Decullier, E., Easterbrook, P. J., Von Elm, E., Gamble, C., Ghersi, D., Ioannidis, J. P. A., Simes, J., & Williamson, P. R. (2008). Systematic Review of the Empirical Evidence of Study Publication Bias and Outcome Reporting Bias. *PLoS ONE*, *3*(8), e3081. https://doi.org/10.1371/journal.pone.0003081

Failing, M., Feldmann-Wüstefeld, T., Wang, B., Olivers, C., & Theeuwes, J. (2019). Statistical regularities induce spatial as well as feature-specific suppression. *Journal of Experimental Psychology: Human Perception and Performance*, *45*(10), 1291–1303. https://doi.org/10.1037/xhp0000660

Ferguson, C. J., & Heene, M. (2012). A Vast Graveyard of Undead Theories: Publication Bias and Psychological Science's Aversion to the Null. *Perspectives on Psychological Science*, *7*(6), 555–561. https://doi.org/10.1177/1745691612459059

Forschack, N., Andersen, S. K., & Müller, M. M. (2017). Global Enhancement but Local Suppression in Feature-based Attention. *Journal of Cognitive Neuroscience*, *29*(4), 619–627. https://doi.org/10.1162/jocn_a_01075

Franco, A., Malhotra, N., & Simonovits, G. (2014). Social science. Publication bias in the social sciences: Unlocking the file drawer. *Science (New York, N.Y.)*, *345*(6203), 1502–1505. https://doi.org/10.1126/science.1255484

Garcia, J. O., Srinivasan, R., & Serences, J. T. (2013). Near-Real-Time Feature-Selective Modulations in Human Cortex. *Current Biology*, *23*(6), 515–522. https://doi.org/10.1016/j.cub.2013.02.013

Geng, J. J. (2014). Attentional Mechanisms of Distractor Suppression. *Current Directions in Psychological Science*, *23*(2), 147–153. https://doi.org/10.1177/0963721414525780

Geng, J. J., Won, B.-Y., & Carlisle, N. B. (2019). Distractor Ignoring: Strategies, Learning, and Passive Filtering. *Current Directions in Psychological Science*, *28*(6), 600–606. https://doi.org/10.1177/0963721419867099

Geweke, F., Li, S.-C., & Störmer, V. (2018). Feature-based attention is constrained to attended locations in older adults. *Journal of Vision*, *18*(10), 306. https://doi.org/10.1167/18.10.306

Guan, M., & Vandekerckhove, J. (2016). A Bayesian approach to mitigation of publication bias. *Psychonomic Bulletin & Review*, *23*(1), 74–86. https://doi.org/10.3758/s13423-015-0868-6

Gulbinaite, R., Roozendaal, D. H. M., & VanRullen, R. (2019). Attention differentially modulates the amplitude of resonance frequencies in the visual cortex. *NeuroImage*, *203*, 116146. https://doi.org/10.1016/j.neuroimage.2019.116146

Hasan, R., Srinivasan, R., & Grossman, E. D. (2017). Feature-based attentional tuning during biological motion detection measured with SSVEP. *Journal of Vision*, *17*(9), 22. https://doi.org/10.1167/17.9.22

Haxby, J., Horwitz, B., Ungerleider, L., Maisog, J., Pietrini, P., & Grady, C. (1994). The functional organization of human extrastriate cortex: A PET-rCBF study of selective attention to faces and locations. *The Journal of Neuroscience*, *14*(11), 6336–6353. https://doi.org/10.1523/JNEUROSCI.14-11-06336.1994

Herrmann, C. S. (2001). Human EEG responses to 1?100 Hz flicker: Resonance phenomena in visual cortex and their potential correlation to cognitive phenomena. *Experimental Brain Research*, *137*(3–4), 346–353. https://doi.org/10.1007/s002210100682

Ipata, A. E., Gee, A. L., Gottlieb, J., Bisley, J. W., & Goldberg, M. E. (2006). LIP responses to a
popout stimulus are reduced if it is overtly ignored. *Nature Neuroscience*, *9*(8), 1071–
1076. https://doi.org/10.1038/nn1734
Itthipuripat, S., Deering, S., & Serences, J. T. (2019). When Conflict Cannot be Avoided:
Relative Contributions of Early Selection and Frontal Executive Control in Mitigating
Stroop Conflict. *Cerebral Cortex*, *29*(12), 5037–5048.
https://doi.org/10.1093/cercor/bhz042
Itthipuripat, S., Garcia, J. O., & Serences, J. T. (2013). Temporal dynamics of divided spatial
attention. *Journal of Neurophysiology*, *109*(9), 2364–2373.
https://doi.org/10.1152/jn.01051.2012
Jiang, Y., Wu, X., & Gao, X. (2017). A category-specific top-down attentional set can affect the
neural responses outside the current focus of attention. *Neuroscience Letters*, *659*, 80–
85. https://doi.org/10.1016/j.neulet.2017.07.029
Kadel, H., Feldmann-Wüstefeld, T., & Schubö, A. (2017). Selection history alters attentional
filter settings persistently and beyond top-down control: Selection history alters
attentional filter settings. *Psychophysiology*, *54*(5), 736–754.
https://doi.org/10.1111/psyp.12830
Kaiser, P. K. (1988). Sensation luminance: A new name to distinguish CIE luminance from
luminance dependent on an individual's spectral sensitivity. *Vision Research*, *28*(3),
455–456. https://doi.org/10.1016/0042-6989(88)90186-1
Kappenman, E. S., & Luck, S. J. (2010). The effects of electrode impedance on data quality and
statistical significance in ERP recordings. *Psychophysiology*.
https://doi.org/10.1111/j.1469-8986.2010.01009.x
Kashiwase, Y., Matsumiya, K., Kuriki, I., & Shioiri, S. (2012). Time Courses of Attentional
Modulation in Neural Amplification and Synchronization Measured with Steady-state
Visual-evoked Potentials. *Journal of Cognitive Neuroscience*, *24*(8), 1779–1793.
https://doi.org/10.1162/jocn_a_00212
Kastner, S., & Ungerleider, L. G. (2000). Mechanisms of Visual Attention in the Human Cortex.
Annual Review of Neuroscience, *23*(1), 315–341.
https://doi.org/10.1146/annurev.neuro.23.1.315
Keitel, C., & Müller, M. M. (2016). Audio-visual synchrony and feature-selective attention co-
amplify early visual processing. *Experimental Brain Research*, *234*(5), 1221–1231.
https://doi.org/10.1007/s00221-015-4392-8
Kim, Y. J., Grabowecky, M., Paller, K. A., Muthu, K., & Suzuki, S. (2007). Attention induces
synchronization-based response gain in steady-state visual evoked potentials. *Nature
Neuroscience*, *10*(1), 117–125. https://doi.org/10.1038/nn1821
Kiyonaga, A., & Egner, T. (2016). Center-Surround Inhibition in Working Memory. *Current
Biology*, *26*(1), 64–68. https://doi.org/10.1016/j.cub.2015.11.013
Kleiner, M., Brainard, D., & Pelli, D. (2007). *What's new in Psychtoolbox-3?* European
Conference on Visual Perception (ECVP), Arezzo, Italy.
https://pdfs.semanticscholar.org/04d4/7572cec08b7a582a9366e5ac61dcfd633f2a.pdf
Kravitz, D., & Mitroff, S. (2017). Estimates of a priori power and false discovery rates induced by
post-hoc changes from thousands of independent replications. *Journal of Vision*, *17*(10),
223. https://doi.org/10.1167/17.10.223
Lamy, D. F., Antebi, C., Aviani, N., & Carmel, T. (2008). Priming of Pop-out provides reliable
measures of target activation and distractor inhibition in selective attention. *Vision
Research*, *48*(1), 30–41. https://doi.org/10.1016/j.visres.2007.10.009
Lamy, D. F., & Kristjansson, A. (2013). Is goal-directed attentional guidance just intertrial
priming? A review. *Journal of Vision*, *13*(3), 14–14. https://doi.org/10.1167/13.3.14

Lithari, C., Sánchez-García, C., Ruhnau, P., & Weisz, N. (2016). Large-scale network-level
processes during entrainment. *Brain Research*, *1635*, 143–152.
https://doi.org/10.1016/j.brainres.2016.01.043
Luck, S. J. (2005). *An Introduction to the Event-Related Potential Technique* (1st ed.). MIT
Press.
Martinez-Trujillo, J. C., & Treue, S. (2004). Feature-Based Attention Increases the Selectivity of
Population Responses in Primate Visual Cortex. *Current Biology*, *14*(9), 744–751.
https://doi.org/10.1016/j.cub.2004.04.028
Martinovic, J., & Andersen, S. K. (2018). Cortical summation and attentional modulation of
combined chromatic and luminance signals. *NeuroImage*, *176*, 390–403.
https://doi.org/10.1016/j.neuroimage.2018.04.066
Martinovic, J., Wuerger, S. M., Hillyard, S. A., Müller, M. M., & Andersen, S. K. (2018). Neural
mechanisms of divided feature-selective attention to colour. *NeuroImage*, *181*, 670–682.
https://doi.org/10.1016/j.neuroimage.2018.07.033
Mishkin, M., & Ungerleider, L. G. (1982). Contribution of striate inputs to the visuospatial
functions of parieto-preoccipital cortex in monkeys. *Behavioural Brain Research*, *6*(1),
57–77. https://doi.org/10.1016/0166-4328(82)90081-X
Moher, J., & Egeth, H. E. (2012). The ignoring paradox: Cueing distractor features leads first to
selection, then to inhibition of to-be-ignored items. *Attention, Perception, &*
Psychophysics, *74*(8), 1590–1605. https://doi.org/10.3758/s13414-012-0358-0
Moher, J., Lakshmanan, B. M., Egeth, H. E., & Ewen, J. B. (2014). Inhibition Drives Early
Feature-Based Attention. *Psychological Science*, *25*(2), 315–324.
https://doi.org/10.1177/0956797613511257
Morgan, S. T., Hansen, J. C., & Hillyard, S. A. (1996). Selective attention to stimulus location
modulates the steady-state visual evoked potential. *Proceedings of the National*
Academy of Sciences of the United States of America, *93*(10), 4770–4774.
Müller, M. M., Andersen, S., Trujillo, N. J., Valdes-Sosa, P., Malinowski, P., & Hillyard, S. A.
(2006). Feature-selective attention enhances color signals in early visual areas of the
human brain. *Proceedings of the National Academy of Sciences*, *103*(38), 14250–14254.
https://doi.org/10.1073/pnas.0606668103
Müller, M. M., Gundlach, C., Forschack, N., & Brummerloh, B. (2018). It takes two to tango:
Suppression of task-irrelevant features requires (spatial) competition. *NeuroImage*, *178*,
485–492. https://doi.org/10.1016/j.neuroimage.2018.05.073
Müller, M. M., & Hillyard, S. (2000). Concurrent recording of steady-state and transient event-
related potentials as indices of visual-spatial selective attention. *Clinical*
Neurophysiology, *111*(9), 1544–1552. https://doi.org/10.1016/S1388-2457(00)00371-0
Müller, M. M., Picton, T. W., Valdes-Sosa, P., Riera, J., Teder-Sälejärvi, W. A., & Hillyard, S. A.
(1998). Effects of spatial selective attention on the steady-state visual evoked potential
in the 20–28 Hz range. *Cognitive Brain Research*, *6*(4), 249–261.
https://doi.org/10.1016/S0926-6410(97)00036-0
Müller, M. M., Teder-Sälejärvi, W., & Hillyard, S. A. (1998). The time course of cortical
facilitation during cued shifts of spatial attention. *Nature Neuroscience*, *1*(7), 631–634.
https://doi.org/10.1038/2865
Norcia, A. M., Appelbaum, L. G., Ales, J. M., Cottereau, B. R., & Rossion, B. (2015). The
steady-state visual evoked potential in vision research: A review. *Journal of Vision*,
15(6), 4. https://doi.org/10.1167/15.6.4
Nunez, M. D., Srinivasan, R., & Vandekerckhove, J. (2015). Individual differences in attention
influence perceptual decision making. *Frontiers in Psychology*, *8*, 18.
https://doi.org/10.3389/fpsyg.2015.00018

Owen, A. M., Milner, B., Petrides, M., & Evans, A. C. (1996). Memory for object features versus
1246   memory for object location: A positron-emission tomography study of encoding and
1247   retrieval processes. *Proceedings of the National Academy of Sciences*, *93*(17), 9212–
1248   9217. https://doi.org/10.1073/pnas.93.17.9212
1249   Painter, D. R., Dux, P. E., & Mattingley, J. B. (2015). Causal involvement of visual area MT in
1250   global feature-based enhancement but not contingent attentional capture. *NeuroImage*,
1251   *118*, 90–102. https://doi.org/10.1016/j.neuroimage.2015.06.019
1252   Painter, D. R., Dux, P. E., Travis, S. L., & Mattingley, J. B. (2014). Neural Responses to Target
1253   Features outside a Search Array Are Enhanced during Conjunction but Not Unique-
1254   Feature Search. *Journal of Neuroscience*, *34*(9), 3390–3401.
1255   https://doi.org/10.1523/JNEUROSCI.3630-13.2014
1256   Pei, F., Pettet, M. W., & Norcia, A. M. (2002). Neural correlates of object-based attention.
1257   *Journal of Vision*, *2*(9), 1. https://doi.org/10.1167/2.9.1
1258   Pelli, D. G. (1997). The VideoToolbox software for visual psychophysics: Transforming numbers
1259   into movies. *Spatial Vision*, *10*(4), 437–442. https://doi.org/10.1163/156856897X00366
1260   Quigley, C., Andersen, S. K., Schulze, L., Grunwald, M., & Müller, M. M. (2010). Feature-
1261   selective attention: Evidence for a decline in old age. *Neuroscience Letters*, *474*(1), 5–8.
1262   https://doi.org/10.1016/j.neulet.2010.02.053
1263   Quigley, C., & Müller, M. M. (2014). Feature-Selective Attention in Healthy Old Age: A Selective
1264   Decline in Selective Attention? *Journal of Neuroscience*, *34*(7), 2471–2476.
1265   https://doi.org/10.1523/JNEUROSCI.2718-13.2014
1266   Reeder, R. R., Olivers, C. N. L., & Pollmann, S. (2017). Cortical evidence for negative search
1267   templates. *Visual Cognition*, *25*(1–3), 278–290.
1268   https://doi.org/10.1080/13506285.2017.1339755
1269   Regan, D. (1977). Steady-state evoked potentials. *Journal of the Optical Society of America*,
1270   *67*(11), 1475. https://doi.org/10.1364/JOSA.67.001475
1271   Rosenthal, R. (1979). The file drawer problem and tolerance for null results. *Psychological
1272   Bulletin*, *86*(3), 638–641. https://doi.org/10.1037/0033-2909.86.3.638
1273   Rungratsameetaweemana, N., Itthipuripat, S., Salazar, A., & Serences, J. T. (2018).
1274   Expectations Do Not Alter Early Sensory Processing during Perceptual Decision-Making.
1275   *The Journal of Neuroscience*, *38*(24), 5632–5648.
1276   https://doi.org/10.1523/JNEUROSCI.3638-17.2018
1277   Sàenz, M., Buracas, G. T., & Boynton, G. M. (2002). Global effects of feature-based attention in
1278   human visual cortex. *Nature Neuroscience*, *5*(7), 631–632. https://doi.org/10.1038/nn876
1279   Sàenz, M., Buracâs, G. T., & Boynton, G. M. (2003). Global feature-based attention for motion
1280   and color. *Vision Research*, *43*(6), 629–637. https://doi.org/10.1016/S0042-
1281   6989(02)00595-3
1282   Sawaki, R., & Luck, S. J. (2010). Capture versus suppression of attention by salient singletons:
1283   Electrophysiological evidence for an automatic attend-to-me signal. *Attention,
1284   Perception, & Psychophysics*, *72*(6), 1455–1470. https://doi.org/10.3758/APP.72.6.1455
1285   Srinivasan, R., Bibi, F. A., & Nunez, P. L. (2006). Steady-State Visual Evoked Potentials:
1286   Distributed Local Sources and Wave-Like Dynamics Are Sensitive to Flicker Frequency.
1287   *Brain Topography*, *18*(3), 167–187. https://doi.org/10.1007/s10548-006-0267-4
1288   Steinhauser, M., & Andersen, S. K. (2019). Rapid adaptive adjustments of selective attention
1289   following errors revealed by the time course of steady-state visual evoked potentials.
1290   *NeuroImage*, *186*, 83–92. https://doi.org/10.1016/j.neuroimage.2018.10.059
1291   Sterling, T. D. (1959). Publication Decisions and their Possible Effects on Inferences Drawn
1292   from Tests of Significance—Or Vice Versa. *Journal of the American Statistical
1293   Association*, *54*(285), 30–34. https://doi.org/10.1080/01621459.1959.10501497

Sterling, T. D., Rosenbaum, W. L., & Weinkam, J. J. (1995). Publication Decisions Revisited:
The Effect of the Outcome of Statistical Tests on the Decision to Publish and Vice Versa.
The American Statistician, *49*(1), 108–112.
https://doi.org/10.1080/00031305.1995.10476125
Stilwell, B. T., & Vecera, S. P. (2019). Cued distractor rejection disrupts learned distractor
rejection. *Visual Cognition*, *27*(3–4), 327–342.
https://doi.org/10.1080/13506285.2018.1564808
Störmer, V. S., & Alvarez, G. A. (2014). Feature-Based Attention Elicits Surround Suppression
in Feature Space. *Current Biology*, *24*(17), 1985–1988.
https://doi.org/10.1016/j.cub.2014.07.030
Tallon-Baudry, C., Bertrand, O., Delpuech, C., & Pernier, J. (1996). Stimulus Specificity of
Phase-Locked and Non-Phase-Locked 40 Hz Visual Responses in Human. *The Journal*
of Neuroscience, *16*(13), 4240–4249. https://doi.org/10.1523/JNEUROSCI.16-13-
04240.1996
Talsma, D., Doty, T. J., Strowd, R., & Woldorff, M. G. (2006). Attentional capacity for processing
concurrent stimuli is larger across sensory modalities than within a modality.
Psychophysiology, *43*(6), 541–549. https://doi.org/10.1111/j.1469-8986.2006.00452.x
Theeuwes, J. (2013). Feature-based attention: It is all bottom-up priming. *Philosophical*
Transactions of the Royal Society B: Biological Sciences, *368*(1628), 20130055.
https://doi.org/10.1098/rstb.2013.0055
Thigpen, N. N., Kappenman, E. S., & Keil, A. (2017). Assessing the internal consistency of the
event-related potential: An example analysis: Assessing internal consistency of the ERP.
Psychophysiology, *54*(1), 123–138. https://doi.org/10.1111/psyp.12629
Thigpen, N. N., Petro, N. M., Oschwald, J., Oberauer, K., & Keil, A. (2019). Selection of Visual
Objects in Perception and Working Memory One at a Time. *Psychological Science*,
30(9), 1259–1272. https://doi.org/10.1177/0956797619854067
Toffanin, P., de Jong, R., Johnson, A., & Martens, S. (2009). Using frequency tagging to
quantify attentional deployment in a visual divided attention task. *International Journal of*
Psychophysiology, *72*(3), 289–298. https://doi.org/10.1016/j.ijpsycho.2009.01.006
Van Moorselaar, D., & Slagter, H. A. (2020). Inhibition in selective attention. *Annals of the New*
York Academy of Sciences, *1464*(1), 204–221.
Vatterott, D. B., & Vecera, S. P. (2012). Experience-dependent attentional tuning of distractor
rejection. *Psychonomic Bulletin & Review*, *19*(5), 871–878.
https://doi.org/10.3758/s13423-012-0280-4
Verghese, P., Kim, Y.-J., & Wade, A. R. (2012). Attention Selects Informative Neural
Populations in Human V1. *Journal of Neuroscience*, *32*(46), 16379–16390.
https://doi.org/10.1523/JNEUROSCI.1174-12.2012
Vialatte, F.-B., Maurice, M., Dauwels, J., & Cichocki, A. (2010). Steady-state visually evoked
potentials: Focus on essential paradigms and future perspectives. *Progress in*
Neurobiology, *90*(4), 418–438. https://doi.org/10.1016/j.pneurobio.2009.11.005
Vissers, M. E., Gulbinaite, R., van den Bos, T., & Slagter, H. A. (2017). Protecting visual short-
term memory during maintenance: Attentional modulation of target and distractor
representations. *Scientific Reports*, *7*(1), 4061. https://doi.org/10.1038/s41598-017-
03995-0
Wang, B., & Theeuwes, J. (2018a). Statistical regularities modulate attentional capture. *Journal*
of Experimental Psychology: Human Perception and Performance, 44(1), 13–17.
https://doi.org/10.1037/xhp0000472

Wang, B., & Theeuwes, J. (2018b). How to inhibit a distractor location? Statistical learning
versus active, top-down suppression. *Attention, Perception, & Psychophysics*.
https://doi.org/10.3758/s13414-018-1493-z
Wang, J., Clementz, B. A., & Keil, A. (2007). The neural correlates of feature-based selective
attention when viewing spatially and temporally overlapping images. *Neuropsychologia*,
45(7), 1393–1399. https://doi.org/10.1016/j.neuropsychologia.2006.10.019
Wang, Y., Miller, J., & Liu, T. (2015). Suppression effects in feature-based attention. *Journal of
Vision*, *15*(5), 15. https://doi.org/10.1167/15.5.15
White, A. L., & Carrasco, M. (2011). Feature-based attention involuntarily and simultaneously
improves visual performance across locations. *Journal of Vision*, *11*(6), 15–15.
https://doi.org/10.1167/11.6.15
Williams, R. S., Pratt, J., & Ferber, S. (2020). Directed avoidance and its effect on visual
working memory. *Cognition*, *201*, 104277.
https://doi.org/10.1016/j.cognition.2020.104277
Won, B.-Y., & Geng, J. J. (2020). Passive exposure attenuates distraction during visual search.
Journal of Experimental Psychology: General. https://doi.org/10.1037/xge0000760
Won, D.-O., Hwang, H.-J., Dähne, S., Müller, K.-R., & Lee, S.-W. (2016). Effect of higher
frequency on the classification of steady-state visual evoked potentials. *Journal of
Neural Engineering*, *13*(1), 016014. https://doi.org/10.1088/1741-2560/13/1/016014
Zar, J. H. (2010). *Biostatistical Analysis* (5th ed.). Prentice-Hall.
Zhang, D., Maye, A., Gao, X., Hong, B., Engel, A. K., & Gao, S. (2010). An independent brain–
computer interface using covert non-spatial visual selective attention. *Journal of Neural
Engineering*, *7*(1), 016010. https://doi.org/10.1088/1741-2560/7/1/016010
Zhang, W., & Luck, S. J. (2009). Feature-based attention modulates feedforward visual
processing. *Nature Neuroscience*, *12*(1), 24–25. https://doi.org/10.1038/nn.2223
Zhang, Z., Gaspelin, N., & Carlisle, N. B. (2020). Probing early attention following negative and
positive templates. *Attention, Perception, & Psychophysics*, *82*(3), 1166–1175.
https://doi.org/10.3758/s13414-019-01864-8
Zhu, D., Bieger, J., Garcia Molina, G., & Aarts, R. M. (2010). A Survey of Stimulation Methods
Used in SSVEP-Based BCIs. *Computational Intelligence and Neuroscience*, *2010*, 1–12.
https://doi.org/10.1155/2010/702357

1372

1373

1374

1375

1376

1377

1378

1379

1380

1381

1382

1383

1384

1385

Appendix A. Example frames during the stimulus presentation. Eight example frames (1-8) from the stimulus presentation period illustrate how the flicker was achieved (refresh rate was 120 Hz, so each frame was ~8.33 ms). In this example, the attended color is yellow, and the attended frequency is 24 Hz (3 frames on, 2 frames off). Blue is the unattended color (30 Hz; 2 frames on, 2 frames off). The white dot in the upper left-hand corner was used to record the attended frequency flicker using a photodiode (this corner of the screen was covered with thick, opaque black electrical tape so that it was not visible to the participants.

Appendix B. Accuracy for target-present trials as a function of the time between Cue Onset and the Target Onset. For short cue-target intervals (<= 275 ms), participants were more accurate for attend cues than ignore cues. This pattern suggests that participants were more quickly able to utilize the attend cue than the ignore cue. Shaded error bars indicate +/- 1 SEM. Small gray dots indicate p < .05 (uncorrected), large dots indicate p < .001 (uncorrected).

Appendix C. Power and SNR for each frequency. (A) Power for each frequency using the Gaussian wavelet filter analysis. (B) SNR for each frequency, calculated as the power at the frequency (e.g., 24 Hz) divided by the power at the average of the 2 neighboring 1-Hz frequencies on either side (e.g., average of 22, 23, 25, and 26 Hz). The theoretical chance level for SNR is 1 (dotted line), but because SNR is calculated with neighboring frequencies, frequencies that are adjacent to a significant "peak" may have values below 1. (C) Cohen's d for the comparison between SNR at each of the two target SSVEP frequencies (24 Hz, 30 Hz) relative to other baselined frequencies (3-33 Hz excluding frequencies within +/- 2 Hz of the target SSVEP frequencies).

Appendix D. Time-course of SNR for each frequency. The stimulus began flickering at -1,333 ms, and the cue indicating which color to attend appeared at 0 ms. Red lines show when 24 Hz was the attended frequency; Blue lines show when 30 Hz was the attended frequency. Solid lines show data from the "attend cue" condition; Dotted lines show the "ignore cue" condition.

Appendix E. Frequency spectra separately for each target/distractor presence condition. Trials were counterbalanced to have a 50% chance of having a target event (T1) and to have 50% chance of including a distractor event (D1). Thus, 25% of trials had neither a target nor distractor (T0D0), 25% of trials had a target only (T1D0), 25% of trials had a distractor only (T0D1), and 25% of trials had both a target and a distractor (T1D1). Frequency spectra for each sub-condition are shown (Rows: Attend Cue or Ignore Cue, Columns: Each combination of target and distractor present/absent).

Appendix F. Signal to noise ratio (SNR) values separately for each target/distractor presence condition. Trials were counterbalanced have a 50% chance of having a target event (T1) and to have 50% chance of including a distractor event (D1). Thus, 25% of trials had neither a target nor distractor (T0D0), 25% of trials had a target only (T1D0), 25% of trials had a distractor only (T0D1), and 25% of trials had both a target and a distractor (T1D1). Frequency spectra for each sub-condition are shown (Rows: Attend Cue or Ignore Cue, Columns: Each combination of target and distractor present/absent). The bottom row of asterisks shows post-hoc, uncorrected significance for overall SSVEP signal compared to a null value of 1. The SSVEP signal was overall highly significant (***, p<.001). The top row of asterisks shows post-hoc, uncorrected significance for the comparison between the two adjacent bars (n.s. p > .10, ~ p <.10, * p < .05). Note, no conditions showed an attention effect (attended frequency > ignored frequency); the only significant, uncorrected post-hoc comparison was in the wrong direction (ignored > attended).

1480 **Appendix G. Power by frequency separately for each color distance condition.**
1481 Target and distractor colors were randomly assigned on each trial from a pool of 5
1482 possible colors. Thus, the target and distractor colors could be either 72 degrees (Color
1483 1) or 144 degrees (Color 2) apart on a color wheel. We found no evidence of an
1484 attention effect in either color distance condition.

1485

Appendix H. An additional analysis variant for the main SNR measure: skipping the first bin for computing SNR. Rather than using the pre-registered frequencies of +/- 1 and +/- 2 Hz for computing SNR, we instead skipped the first 1 Hz bin. Since +/-1 Hz had greater than baseline power, we may have attenuated our ability to observe SSVEP-related differences by including this bin in our SNR subtraction. For this analysis variant, we instead calculated SNR as the peak frequency minus the average of all frequencies +/- 2 and +/- 3 Hz from the peak (e.g., to compute SNR for 24 Hz, we subtracted the mean power at 21, 22, 26, and 27 Hz). Although overall SNR was much higher across all conditions using this metric, the pattern across experimental conditions was unchanged (i.e., we found no significant attention effects).

Appendix I. FFT analysis with a wider x-axis to show both the fundamental and second harmonic frequencies. (Left) FFT for the 'attend cue' condition. (Right) FFT for the 'ignore cue' condition. X-axis values are frequency (Hz); Y-axis values are amplitude (microvolts).

Appendix J. Violin plot of the second harmonic frequencies 48 Hz and 60 Hz from
the FFT analysis. (Left) Violin plot of SNR for the second harmonic frequencies in the
'attend cue' condition; SNR for both harmonics was greater than 1, but there were no
attention effects. (B) Violin plot of SNR for the second harmonic frequencies in the
'ignore cue' condition; SNR for both harmonics was greater than 1, but there were no
attention effects.

Appendix K. Rhythmic Entrainment Source Separation (RESS) analysis likewise shows null attention effects. Following code associated with [1], we performed rhythmic entrainment source separation (RESS) on our data to ensure that our a priori choice of electrodes did not impede our ability to find an attention effect. We decided to stick very closely to the default settings for RESS code developed by others in order to take some 'researcher degrees of freedom' out of the equation. We obtained a highly consistent pattern of results despite using a data-driven, single-trial approach that differs substantially from our pre-registered trial-averaged approach. We also note that the SNR values from the RESS approach are lower than the trial-averaged FFT we present in the main analysis, but that RESS does still provide an SNR advantage when compared to a single-trial FFT approach, as in [1]. We first calculated the spatial filters using data from all trials and the full trial length (-1000 ms to 2000 ms). We then applied the spatial filters to calculate SNR for each condition of interest (e.g., "Attend 24 Hz, Attend Cue Condition", 24 Hz RESS time series; 500 ms to 2000 ms). For the analysis, we used a frequency resolution of 0.5 Hz, a full-width half maximum (FWHM) of 0.5 Hz for the center frequency, a FWHM of 1 Hz for the neighboring baseline frequencies +/- 2 Hz from the peak frequency. SNR was calculated as the ratio between each frequency of interest and the frequencies +/- 2 Hz away. (A) Normalized SNR by frequency and topography of the RESS time series optimized for 24 Hz (red) and 30 Hz (blue), collapsed across all conditions. (B) SSVEP response (computed as normalized SNR) for the 24 Hz-optimized RESS time series in the attend cue condition and ignore cue condition. (C) SSVEP response (computed as normalized SNR) for the 30 Hz-optimized RESS time series in the attend cue condition and ignore cue condition. We again found no significant effects of attention for either SSVEP frequency.'

Appendix L. Violin plots of values obtained from the Rhythmic Entrainment Source Separation (RESS) analysis. We found no effect of attention on RESS values in either the Attend Cue condition (left panel) or the Ignore Cue condition (right panel).

Appendix M. Violin plots of SNR values for each frequency, calculated from an FFT analysis on accurate trials only. Performing an FFT analysis on accurate trials only likewise yields null attention effects both in the attend cue condition (left panel) and the ignore cue condition (right panel).

Appendix N. Results of the phase-locking index (PLI) analysis. We performed an
FFT on single trials rather than on condition-averaged waveforms (time window: 333 ms
– 2000 ms), and we extracted single-trial phase values (*'angle.m'*). We calculated a
phase-locking index by computing mean-resultant vector length on histograms of single-
trial phase values (separate histograms for each condition, electrode, and frequency).
Mean-resultant vector length ranges from 0 (fully random values) to 1 (perfectly identical
values), for reference, see: Zar (2010). **(A)** Phase locking index (PLI) as indexed by
mean-resultant vector length, averaged across electrodes O1, O2, and Oz. Replicating
prior work, we found robust PLI values at the two SSVEP frequencies (24 and 30 Hz).
(B) However, we found no evidence that PLI values were modulated by attention in the
expected direction.

1622 **Appendix O. P3 component at electrodes Pz and POz, split by whether or not a**
1623 **response was made.** (A) No response made, "attend cue" condition. (B) No response
1624 made, "ignore cue" condition. (C) Response made, "attend cue" condition. (D)
1625 Response made, "ignore cue" condition. Shaded error bars represent standard error of
1626 the mean.

Appendix P. Study overview for studies employing a variant of the "competing
gratings" task. From left to right, columns indicate: "Exp." = Experiment number out of
those reviewed, "Ref" = Paper reference, "N" = number of subjects in the experiment,
"Total Trials" = total number of trials completed by the participant, "Trials Per Cond." =
The number of trials that could be analyzed per condition (i.e., after excluding target and
distractor onsets), "Stimulus Duration" = the duration, in seconds, that participants
attended the stimulus, "Freq." = Frequency, in Hertz (Hz), that the stimuli flickered at,
"Sig" = Qualitative code for the overall presence of a basic attention effects (when
expected); 1 = attended > ignored, -1 = attended < ignored, 0.5 = mixed effects across
conditions, 0 = null, n/a = statistical values for the basic attention effect not reported
directly. Notes: * Statistics were performed for individuals but not across subjects;
standard attention effect in one condition, reversed effect in the other. **Group level
statistics not reported. ***No attention effect for the main flicker frequencies (14.2, 17
Hz), but attention effect for the slow oscillating changes to the Gabor's features (3.14,
3.62 Hz).
1657

Exp.	Ref.	N	Total Trials	Trials Per Cond.	Stim. Dur. (s)	Freq. (Hz)	Sig.
1	(Chen et al., 2003)	11	16	8	100	7.41, 8.33	.5*
2	(Wang et al., 2007), Exp 1	12	16	8	120	7.14, 8.33	0
3	(Wang et al., 2007), Exp 2	12	16	8	120	7.69, 7.14, 8.33	-1
4	(Wang et al., 2007), Exp 3	12	16	8	120	6.67, 7.14, 7.69, 8.33	-1
5	(Allison et al., 2008)	14	8	4	60	10, 12	n/a**
6	(Keitel & Müller, 2016)	13	600	75	3.5	3.14, 3.62, 14.2, 17	.5***

1658
1659

Appendix Q. Study overview for studies employing a variant of the "whole-field flicker" task. From left to right, columns indicate: "Exp." = Experiment number out of those reviewed, "Ref" = Paper reference, "N" = number of subjects in the experiment, "Total Trials" = total number of trials completed by the participant, "Trials Per Cond." = The number of trials that could be analyzed per condition (i.e., after excluding target and distractor onsets), "Stimulus Duration" = the duration, in seconds, that participants attended the stimulus, "Freq." = Frequency, in Hertz (Hz), of the stimulus flicker, "Sig" = Qualitative code for the overall presence of a basic attention effects (when expected); 1 = attended > ignored, -1 = attended < ignored, 0.5 = mixed effects across conditions, 0 = null, n/a = statistical values for the basic attention effect not reported directly. Notes: *Analyzed harmonics (2F, 4F) but not the fundamental frequency. 2F but not 4F had a significant attention effect. ** Attention modulation scores were only compared across conditions, not to baseline; they are presumably overall significant, but this was not formally tested.

Exp.	Ref.	N	Total Trials	Trials Per Cond.	Stim. Dur. (s)	Freq. (Hz)	Sig.
7	(Pei et al., 2002)	11	20	20	8	2.4, 3	n/a*
8	(Müller et al., 2006)	11	450	153	4.114	7, 11.67	1
9	(Andersen et al., 2008)	15	600	90	3.092	10,12,15, 17.14	1
10	(Andersen et al., 2009)	15	432	72	3.042	10, 12	1
11	(Andersen & Müller, 2010)	16	480	240	2	11.98, 16.77	1
12	(Quigley et al., 2010)	10	440	110	2.2	8, 12	1
13	(Zhang et al., 2010)	18	300	300	4	10, 12	1
14	(Andersen et al., 2012)	16	300	60	8.5	10, 12	1
15	(Quigley & Müller, 2014)	20	320	90	4.167	15, 17	1
16	(Andersen et al., 2015)	15	192	96	15	8, 10, 12, 15	1
17	(Forschack et al., 2017)	23	480	120	1.783	10, 12.5, 15, 17.5	1
18	(Martinovic & Andersen, 2018)	9	768	23	6.5	10, 12	n/a**
19	(Martinovic et al., 2018) Exp 1	11	600	70	3.14	8.57, 10, 12, 15	1
20	(Martinovic et al., 2018) Exp 2	14	600	70	3.14	8.57, 10, 12, 15	1
21	(Steinhauser & Andersen, 2019)	17	1600	400	1	10, 15	1

Appendix R. Study overview for studies employing a variant of the "hemifield flicker" task. From left to right, columns indicate: "Exp." = Experiment number out of those reviewed, "Ref" = Paper reference, "N" = number of subjects in the experiment, "Total Trials" = total number of trials completed by the participant, "Trials Per Cond." = The number of trials that could be analyzed per condition (i.e., after excluding target and distractor onsets), "Stimulus Duration" = the duration, in seconds, that participants attended the stimulus, "Freq." = Frequency, in Hertz (Hz), of the stimulus flicker, "Sig" = Qualitative code for the overall presence of a basic attention effects (when expected); 1 = attended > ignored, -1 = attended < ignored, 0.5 = mixed effects across conditions, 0 = null, n/a = statistical values for the basic attention effect not reported directly.

Exp.	Ref.	N	Total Trials	Trials Per Cond.	Stim. Dur. (s)	Freq. (Hz)	Sig.
22	(Andersen et al., 2011)	19	600	100	3.05	8.46, 11.85, 14.81, 19.75	1
23	(Andersen et al., 2013) Exp 1	13	560	160	2.94	7.5, 8.57, 10, 12	1
24	(Andersen et al., 2013) Exp 2	11	560	320	2.94	7, 8.57, 10, 12	1
25	(Störmer & Alvarez, 2014)	16	640	160	2.6	7.1, 8.5, 10.7	1
26	(Müller et al., 2018)	23	480	120	1.783	6.5, 8.5, 11.5, 13.5	1
27	(Adamian et al., 2019)	16	672	128	2.94	7.5, 8.57, 10, 12	1

Appendix S. Study overview for studies employing a variant of the "attend
central, peripheral flicker" task. From left to right, columns indicate: "Exp." =
Experiment number out of those reviewed, "Ref" = Paper reference, "N" = number of
subjects in the experiment, "Total Trials" = total number of trials completed by the
participant, "Trials Per Cond." = The number of trials that could be analyzed per
condition (i.e., after excluding target and distractor onsets), "Stimulus Duration" = the
duration, in seconds, that participants attended the stimulus, "Freq." = Frequency, in
Hertz (Hz), of the stimulus flicker, "Sig" = Qualitative code for the overall presence of a
basic attention effects (when expected); 1 = attended > ignored, -1 = attended <
ignored, 0.5 = mixed effects across conditions, 0 = null, n/a = statistical values for the
basic attention effect not reported directly. Notes: *No attention effect at *a priori*
electrode; other electrodes were examined post-hoc, but statistics were not reported for
each. **Significant in 1 of 2 expected conditions.

Exp.	Ref.	N	Total Trials	Trials Per Cond.	Stim. Dur. (s)	Freq. (Hz)	Sig.
29	(Painter et al., 2014) Exp 1	20	288	144	7.2	12.5, 16.7	1
30	(Painter et al., 2014) Exp 2	20	216	216	8.4	7.6, 13.3, 17.8	1
31	(Painter et al., 2015, p. 2)	20	512	128	8	8, 12	0*
32	(Jiang et al., 2017)	23	288	144	8.4	12, 15	.5**
33	(Chu & D'Zmura, 2019) Exp 1	20	128	32	7	12.5, 18.75	1
34	(Chu & D'Zmura, 2019) Exp 2	21	128	32	9	12.5, 18.75	1

Appendix T. Accuracy and task variant for studies where participants detected a
target within the flickering stimulus (whole-field and hemifield flicker tasks). To
test if the difficulty of our task may have contributed to our null results, we examined
behavior from studies in which participants monitored for a target in the flickering
stimulus (i.e., whole-field and hemifield flicker tasks). We also noted the type of target
and how long it was on the screen. Notes: †Values were not listed in the text, so some
values were approximated based on the figures (e.g., hit rates or d' depicted in a bar
graph). *Analyzed harmonics (2F, 4F) but not the fundamental frequency. 2F but not 4F
had a significant attention effect. **Attention modulation scores were only compared
across conditions, not to baseline; they were presumably significant overall, but this was
not formally tested.

Exp.	Ref.	Behavior	Target type	Dur. (ms)	Sig.
7	(Pei et al., 2002)	n/a	No Targets	n/a	n/a*
8	(Müller et al., 2006)	d' = 1.95 – 2.89	75% Coherent Motion	586	1
9	(Andersen et al., 2008)	d' = 2.74 – 3.25	70% Coherent Motion	500	1
10	(Andersen et al., 2009)	d' = 2.67 – 3.23†	20% Luminance Decrement	200	1
11	(Andersen & Müller, 2010)	d' = 1.83	75% Coherent Motion	298	1
12	(Quigley et al., 2010)	d' = 2.665	85% Coherent Motion	556	1
13	(Zhang et al., 2010)	n/a	No Targets	n/a	1
14	(Andersen et al., 2012)	d' = 2.64	50% Coherent Motion	400	1
15	(Quigley & Müller, 2014)	Acc = 87.5% - 98%†	40% Coherent Oblique Motion	500	1
16	(Andersen et al., 2015)	d' = 1.3 – 1.75†	70% Coherent Motion	500	1
17	(Forschack et al., 2017)	d' = 2	60% Coherent Motion	300	1
18	(Martinovic & Andersen, 2018)	d' = 0.8 – 3.0†	50% Coherent Motion	400	n/a**
19	(Martinovic et al., 2018) Exp 1	d' = 1.05	50% Coherent Motion	400	1
20	(Martinovic et al., 2018) Exp 2	d' = 1.0	50% Coherent Motion	400	1
21	(Steinhauser & Andersen, 2019)	Acc = 90.3%	75% Coherent Motion	500	1
22	(Andersen et al., 2011)	d' = 0.95 – 2.8	75% Coherent Motion	500	1
23	(Andersen et al., 2013) Exp 1	d' = 2.133 – 3.111	20% Luminance Decrement	200	1
24	(Andersen et al., 2013) Exp 2	d' = 2.637	20% Luminance Decrement	200	1
25	(Störmer & Alvarez, 2014)	Acc = 78%	80% Coherent Motion	230	1
26	(Müller et al., 2018)	d' = 1.81	60% Coherent Motion	300	1
27	(Adamian et al., 2019)	d' = 2.8†	20% Luminance Decrement	200	1

1749

A

24 Hz
30 Hz

Baseline (1,333 ms)

Cue onset (Stimulus begins)

Target Event (333 ms)

Inter-Trial Interval (1,800 - 2,100 ms, jittered)

Stimulus Period (2000 ms)

B

72 deg

144 deg

C

Cue

Target Event (Respond)

Distractor Event (Do not respond)

'Attend Cue' Condition

'Ignore Cue' Condition