SARS-CoV-2 Cell Entry Factors ACE2 and TMPRSS2 are Expressed in the Pancreas but

are Not Enriched in Islet Endocrine Cells

Katie C. Coate[1,*], Jeeyeon Cha[1,*], Shristi Shrestha[1], Wenliang Wang[2], Luciana Mateus

Gonçalves[3], Joana Almaça[3], Meghan E. Kapp[4], Maria Fasolino[2], Ashleigh Morgan[2], Chunhua

Dai[1], Diane C. Saunders[1], Rita Bottino[5,6], Radhika Aramandla[1], Regina Jenkins[1], Roland Stein[7],

Klaus H. Kaestner[2], Golnaz Vahedi[2], HPAP consortium[8], Marcela Brissova[1], Alvin C.

Powers[1,7,9,10]

[1]Division of Diabetes, Endocrinology and Metabolism, Department of Medicine, Vanderbilt

University Medical Center, Nashville, TN, 37232

[2]Department of Genetics and Institute for Diabetes, Obesity, and Metabolism, University of

Pennsylvania Perelman School of Medicine, Philadelphia, PA, 19104

[3]Division of Endocrinology, Diabetes and Metabolism, Department of Medicine, University of

Miami Miller School of Medicine, Miami, FL, 33136

[4]Department of Pathology, Microbiology and Immunology, Vanderbilt University Medical Center,

Nashville, TN, 37232

[5]Institute of Cellular Therapeutics, Allegheny Health Network, Pittsburgh, PA, 15212

[6]Imagine Pharma, Devon, PA, 19333

[7]Department of Molecular Physiology and Biophysics, Vanderbilt University School of Medicine,

Nashville, TN, 37232

[8]The Human Pancreas Analysis Program (RRID:SCR_016202)

[9]VA Tennessee Valley Healthcare System, Nashville, TN, 37212

[10]Lead Contact

[*]Co-first authors

Running Title: ACE2 and TMPRSS2 are Not Enriched in Pancreatic β cells

Correspondence: al.powers@vumc.org; marcela.brissova@vumc.org

Alvin C. Powers or Marcela Brissova, Vanderbilt University Medical Center, 2215 Garland Ave,

Nashville, TN 37232-0475

Summary/Abstract

Reports of new-onset diabetes and diabetic ketoacidosis in individuals with COVID-19

have led to the hypothesis that SARS-CoV-2, the virus that causes COVID-19, is directly

cytotoxic to pancreatic islet β cells. This would require binding and entry of SARS-CoV-2 into

host β cells via cell surface co-expression of ACE2 and TMPRSS2, the putative receptor and

effector protease, respectively. To define ACE2 and TMPRSS2 expression in the human

pancreas, we examined six transcriptional datasets from primary human islet cells and

assessed protein expression by immunofluorescence in pancreata from donors with and without

diabetes. *ACE2* and *TMPRSS2* transcripts were low or undetectable in pancreatic islet

endocrine cells as determined by bulk or single cell RNA sequencing, and neither protein was

detected in α or β cells from these donors. Instead, ACE2 protein was expressed in the islet and

exocrine tissue microvasculature and also found in a subset of pancreatic ducts, whereas

TMPRSS2 protein was restricted to ductal cells. The absence of significant ACE2 and

TMPRSS2 co-expression in islet endocrine cells reduces the likelihood that SARS-CoV-2

directly infects pancreatic islet β cells through these cell entry proteins.

Introduction

In coronavirus disease of 2019 (COVID-19), elevated plasma glucose levels with or without pre-existing diabetes and BMI have been identified as independent risk factors of morbidity and mortality with severe acute respiratory syndrome-associated coronavirus-2 (SARS-CoV-2) infection (Barron et al., 2020; Cariou et al., 2020; Holman et al., 2020; Riddle et al., 2020; Wang, 2020; Zhou et al., 2020). Isolated cases of new-onset diabetes and diabetic emergencies such as ketoacidosis and hyperosmolar hyperglycemia have been reported with COVID-19 (Chee et al., 2020; Goldman et al., 2020; Hollstein et al., 2020; Kim et al., 2020; Li et al., 2020a; Rafique and Ahmed, 2020; Unsworth et al., 2020), leading to the hypothesis that SARS-CoV-2 has a diabetogenic effect mediated by direct cytotoxicity to pancreatic islet β cells (Koch, 2020).

In vitro studies have shown that SARS-CoV-2 entry into human host cells requires binding to the cell surface receptor angiotensin converting enzyme 2 (ACE2) as well as proteolytic cleavage of the viral spike (S) protein by transmembrane serine protease 2 (TMPRSS2) (Hoffmann et al., 2020; Lan et al., 2020; Shang et al., 2020; Wiersinga et al., 2020). In 2010, Yang and colleagues (Yang et al., 2010) examined autopsy samples from a single deceased patient infected by SARS-CoV-1, which uses similar machinery for binding and cellular entry, and reported expression of ACE2 in pancreatic islet cells. Though the identity of these islet cells was not assessed, the authors suggested that binding of ACE2 by SARS-CoV-1 damages islets and causes acute diabetes, which could be reversed after viral recovery (Yang et al., 2010). There have been occasional reports of other viral infections eliciting a diabetogenic effect (reviewed in (Filippi and von Herrath, 2008)). More recently, Yang and colleagues reported that β-like cells derived from human pluripotent stem cells (hPSCs) as well as β cells of primary human islets express ACE2, raising the possibility of direct infection and cytotoxicity of β cells by SARS-CoV-2 (Yang et al., 2020). Importantly, neither of these prior studies (Yang et al., 2010; Yang et al., 2020) characterized the expression and localization of TMPRSS2, an

obligate co-factor for SARS-CoV-2 cellular entry. Thus, a more detailed analysis of both ACE2

and TMPRSS2 expression and localization in human pancreatic tissue from normal donors and

those with diabetes is urgently needed. The purpose of this study was to test the hypothesis that

native pancreatic islet β cells possess the cellular machinery that could render them direct

targets of SARS-CoV-2. Importantly, we found that ACE2 and TMPRSS2 protein are not

detectable in human islet endocrine cells from normal donors or those with diabetes, making a

direct diabetogenic effect of SARS-CoV-2 via ACE2 and TMPRSS2 unlikely.

Results and Discussion

***ACE2* and *TMPRSS2* mRNA expression is minimal in human α or β cells.**

We first evaluated mRNA expression of *ACE2* and *TMPRSS2* from two existing bulk RNA-

sequencing (RNA-seq) datasets (Arda et al., 2016; Blodgett et al., 2015) where human islet

α and β cells were enriched by fluorescence activated cell sorting and compared their

expression to that of key islet-enriched genes, some of which are normally expressed at

relatively low levels in islet cells (e.g., transcription factors). Median expression level of *ACE2*

and *TMPRSS*2 mRNA was much less than transcripts of such key islet-enriched genes in

human α and β cells (~84% and 92% lower than α and β cell-enriched transcripts, respectively)

(**Figure 1A;** n=7-8 adult donors per study). In addition, analysis of four single-cell (sc) RNA-seq

datasets of human pancreatic cells (Baron et al., 2016; Camunas-Soler et al., 2020; Kaestner et

al., 2019; Segerstolpe et al., 2016) revealed that, in aggregate, less than 1.5% of β cells

expressed *ACE2* or *TMPRSS2*, and each transcript was minimally expressed or undetectable in

all other endocrine cell subsets (**Figure 1B, S4J** and **Table S1**). We note that this includes

analysis of the robust Human Pancreas Analysis Program (HPAP) dataset that includes more

than 25,000 cells from 11 normal donors (Kaestner et al., 2019), findings of which are confirmed

in the analyses of three previously reported, but smaller, datasets (**Figure 1B** and **S4J**).

Furthermore, given that *ACE2* and *TMPRSS2* co-expression is required for canonical SARS-

CoV-2 host cell entry (Hoffmann et al., 2020), we also evaluated this occurrence but found that

no β cells co-expressed *ACE2* and *TMPRSS2* in any of these four datasets (**Table S1**).

As for non-endocrine cells, the HPAP dataset (Kaestner et al., 2019) revealed that a

small subset (< 5%) of endothelial and stellate cells (which include pericytes) expressed

moderate to high levels of *ACE2*, whereas only ~1-3% of either population expressed *ACE2* in

the datasets by Baron and colleagues (Baron et al., 2016) and Segerstolpe and colleagues

(Segerstolpe et al., 2016) (**Figure 1B**). This difference most likely stems from the number of

cells analyzed in that the HPAP dataset contains ~2.5- and 20-fold more cells than that of Baron

et al. (Baron et al., 2016) or Segerstolpe et al. (Segerstolpe et al., 2016), respectively. In

addition, pooled analysis of these datasets (Baron et al., 2016; Camunas-Soler et al., 2020;

Kaestner et al., 2019) showed that less than 1% of acinar or ductal cells expressed *ACE2*,

whereas ~35% of both cell types expressed *TMPRSS2*. In the dataset by Segerstolpe and

colleagues (Segerstolpe et al., 2016), *ACE2* was expressed in ~20% of acinar and ductal cells,

whereas *TMPRSS2* was expressed in greater than 75% of both populations (**Figure 1B**).

Evaluation of *ACE2* and *TMPRSS2* co-expression in non-endocrine cells revealed that on

average, less than 1% of acinar, ductal, endothelial and stellate cells co-expressed both

transcripts in these datasets (Baron et al., 2016; Camunas-Soler et al., 2020; Kaestner et al.,

2019). In the Segerstolpe et al. dataset (Segerstolpe et al., 2016), ~5% of acinar cells and 15%

of ductal cells co-expressed *ACE2* and *TMPRSS2*, but this difference could represent

unintended cell selection bias as this study analyzed the fewest number of cells.

Altogether, these gene expression findings reduce the likelihood that SARS-CoV-2 can

bind and enter human β cells via the canonical pathway involving ACE2 and TMPRSS2.

However, they do not exclude the possibility of viral entry via non-canonical pathways involving

other suggested effector proteases of SARS-CoV-2, such as Cathepsin L (*CTSL*), ADAM

metallopeptidase domain 17 (*ADAM17*), *FURIN* and *TMPRSS4* (Breidenbach et al., 2020;

Schreiber et al., 2020; Seyedpour et al., 2020; Zang et al., 2020). We evaluated the expression

of these transcripts across all four scRNA-seq datasets and found that with the exception of

TMPRSS4, each is ubiquitously expressed to varying degrees in all cell types (**Figure 1B,**

Table S1). In aggregate, however, less than 1% of β cells co-expressed *ACE2* with *CTSL*,

ADAM17, *FURIN* or *TMPRSS4* (**Table S1**). An outstanding question is whether expression of

these transcripts in even 1% of β cells is sufficient to confer permissiveness to SARS-CoV-2

infection *in vivo*. Furthermore, heparan sulfate in the extracellular glycocalyx (Clausen et al.,

2020) and motile cilia (present on islet endocrine cells) (Lee et al., 2020) were mechanisms

proposed to assist ACE2-mediated viral entry. Future studies addressing these possibilities in β

cells are urgently needed. Notwithstanding, *CTSL*, *ADAM17*, *FURIN* and *TMPRSS4* were more

highly expressed in exocrine, endothelial and stellate cells, as was ACE2 (**Figure 1B**). These

gene expression patterns raise the possibility that infection of certain pancreatic cell types by

SARS-CoV-2 indirectly impacts β cell function.

Obesity is a key metabolic risk factor associated with mortality with COVID-19. A recent

report noted increased correlation of *TMPRSS2*, but not *ACE2*, expression with BMI (Taneera et

al., 2020). We evaluated the expression of *ACE2*, *TMPRSS2* and *ADAM17* in β cells from 11

donors according to their BMI in our largest sc-RNAseq dataset (Kaestner et al., 2019). We did

not observe changes in the expression of *ACE2* or *TMPRSS2* expression with increasing BMI

but observed a trend towards higher *ADAM17* expression (**Figure S1**). However, this dataset

included only one patient with an obese BMI. Studies with a larger number of donor islets are

needed to address the role of obesity and other pancreas-resident SARS-CoV-2 effector

molecules on β cell function in the context of COVID-19.

ACE2 and TMPRSS2 protein are not detected in adult or juvenile human islet α or β cells.

Recently, Yang and colleagues (Yang et al., 2020) reported that ACE2 protein was present in α

and β cells from primary human islets and in β-like cells derived from hPSCs. To investigate

ACE2 expression, we immunostained cryosections of intact isolated human islets using the

same ACE2 antibody as used by Yang and colleagues (Yang et al., 2020) (antibody validation

studies in **Figure S2**). In accordance with the transcriptomic data reported above, ACE2 was

not detected in insulin-positive β cells or glucagon-positive α cells in our samples; instead,

ACE2 signal was prominent in surrounding cells such as those of the microvasculature (**Figure

1C, top panels**).

Since hPSC-derived β-like cells differentiated *in vitro* are considered "juvenile-like" and not

functionally mature β cells (Nair et al., 2019), one possibility is that ACE2 is expressed in less

differentiated cells, or in the cultured, immortalized β cells (EndoCβH1) studied by Fignani et al.

(preprint: Fignani, 2020). We investigated such a possibility by staining for ACE2 in pancreatic

sections from juvenile human donors (age range 5 days to 5 years) using the same ACE2

antibody as used by Yang et al. (Yang et al., 2020). However, ACE2 was not detected in these

juvenile β or α cells (**Figure 1C, bottom panels**), making this explanation less likely.

To further characterize ACE2 and TMPRSS2 protein expression in the native pancreas, we

next analyzed human pancreatic tissue sections from normal donors (ND, n=14; age range 18-

59 years) and those with type 2 diabetes (T2D, n=12; age range 42-66 years) or type 1 diabetes

(T1D, n=11; age range 13-63 years) for these proteins on the same sections. ACE2 protein did

not co-localize with markers of α or β cells but was detected within the distinct islet areas

indicative of microvascular structures in all ND individuals (**Figures 2A-C", 2D-F"** and **S3A-D**)

as well as those with T2D (**Figures 2G-I", 2J-L"** and **S3I-L**) or T1D (**Figures 2M-O", 2P-R"** and

S3O-R). We noted similar findings with three additional ACE2-directed antibodies, including the

same antibodies and dilutions reported by Yang et al. (Yang et al., 2020) and by Fignani et al.

(preprint: Fignani, 2020), and one antibody validated by the Human Protein Atlas (HPA) (Hikmet

et al., 2020) (**Figure S2**). Three of these antibodies detect epitopes corresponding to the

extracellular domain of ACE2 (AF933; MAB933; HPA000288), while one detects an epitope in

the C-terminal domain (ab15348). Staining patterns were similar across all antibodies (**Figure**

S2). Furthermore, ACE2 antibody specificity (ab15348) was confirmed via peptide competition

with the commercially available immunizing peptide (**Figure S2F-I**).

TMPRSS2 protein was not detected within islets in ND (**Figures 2A-C", 2D-F"** and **S3E-H**),

T2D (**Figures 2G-I", 2J-L"** and **S3M-N**), or T1D donors (**Figures 2M-O", 2P-R"** and **S3S-V**).

The TMPRSS2-positive structures in exocrine tissue resembled intercalated and larger ducts as

indicated by the typical organization of their nuclei (**Figures 2C"**, **2F", 2I", 2L", 2O" and 2R"**).

We did not detect differences in the signal intensity or spatial distribution of ACE2 or TMPRSS2

between ND and T2D tissue (**Figures 2** and **S3**) but labeling for both proteins appeared to be

reduced in T1D pancreatic sections (**Figure 2M-R" and S3O-V**). Altogether, these observations

indicate that the cell surface proteins required for canonical SARS-CoV-2 host cell entry were

not detected on islet endocrine cells from normal or diabetic donors.

Recent in silico analyses by Vankadari et al. (Vankadari and Wilce, 2020) and Li et al. (Li et

al., 2020b) suggested that human dipeptidyl peptidase 4 (DPP4) may interact with SARS-CoV-2

and facilitate its entry into host cells. Therefore, we examined the distribution of DPP4 in human

pancreas and found that it localized to α cells, but not β cells, of ND, T2D and T1D islets

(**Figure S4A-I**). This finding is consistent with our analysis of four scRNA-seq datasets showing

DPP4-enriched α cells (**Figures 1B and S4J**). Given that we did not detect TMPRSS2 within

islets, these data suggest that DPP4 is an unlikely mediator of β cell entry by SARS-CoV-2

(Drucker, 2020).

The discrepancy between our findings and those of recent studies identifying ACE2 and

TMPRSS2 in human β cells (Yang et al., 2010) (Yang et al., 2020) (preprint: Fignani, 2020) is

likely explained by important differences in experimental approaches and contexts. Yang et al.

(Yang et al., 2010) showed ACE2 staining in a single donor islet but did not identify its cellular

identity with endocrine markers. Furthermore, the ACE2 antibody used in their study was not

reported, which precluded replication of their finding. Fignani et al. (preprint: Fignani, 2020)

identified ACE2 protein in presumed islet endocrine cells from seven non-diabetic donors;

however, ACE2 was primarily seen in subcellular compartments rather than the expected cell

surface location. Colocalization of ACE2 with insulin granules in this study raises the possibility

of a staining artifact. Furthermore, Yang et al. (Yang et al., 2020) examined ACE2 expression in

isolated dispersed human islet cells, whereas our study examined ACE2 in isolated intact islets

and in human islets of native pancreata. It is possible that dispersion of pancreatic cells may

have altered the expression of ACE2 in this context (van den Brink et al., 2017). Likewise,

sources of variability in immunostaining techniques can stem from differences in tissue fixation

procedures, antigen retrieval methods, antibodies used and their dilutions. Notwithstanding, the

remarkable concordance of our findings with those from a recent independent effort in native

pancreas (Kusmartseva, et al) suggest that differences in experimental conditions may explain

the discordance with other studies.

ACE2 is localized to islet and exocrine tissue capillaries.

The non-endocrine cell staining pattern of ACE2 in islets prompted us to examine whether

ACE2 was expressed in the microvasculature, as described in other organs (Hamming et al.,

2004). Indeed, staining of adult and juvenile human pancreatic tissue sections with CD31, an

endothelial cell marker, revealed that ACE2 labeling was localized to the perivascular

compartment of islet capillaries (**Figures 3A-D', I-L'** and **S5A-H'**). In addition, exocrine tissue

capillaries, similar to those in islets, showed perivascular ACE2 labeling (**Figures 3E-H'** and

S5I-P'). As ACE2-positive cells were scarce in T1D pancreatic tissues (**Figures 2M, 2O'-O",**

2P, 2R'-R", and **S3O-R**), the presence of ACE2 labeling in the perivascular compartment was

relatively rare in both islet (**Figure S5E-H'**) and exocrine (**Figure S5M-P'**) T1D tissues. The

structure and close relationship between ACE2-positive perivascular cells and CD31-positive

endothelial cells in islet and exocrine tissue capillaries raised the possibility that ACE2-positive

cells were in fact pericytes enveloping capillary endothelial cells of the vascular tube (Almaca et

al., 2018). In addition, we observed ACE2-positive cytoplasmic processes extending along

CD31-positive endothelial cells colocalized with the extracellular matrix marker collagen-IV

within the vascular basement membrane (**Figures 3D'**, **3H'**, **S5D'**, **S5H'**, **S5L' and S5P'**), a

pattern that is consistent with pericyte morphology (Almaca et al., 2018).

To address this observation more directly, we visualized ACE2 expression in relationship to

known pericyte markers and found that ACE2 was enriched in pericyte populations expressing

PDGFRβ (**Figure 4A-D**). Indeed, colocalization analysis revealed that ~60% of ACE2-positive

cells were PDGFRβ-positive (**Figure 4H-I**). Since PDGFRβ labels pericytes and perivascular

fibroblasts (Almaca et al., 2020), we also stained tissues with a more specific pericyte marker,

NG2, and found that ~30% of ACE2-positive cells were NG2-positive (**Figure 4E-G', 4H-I**).

These data agree with our scRNA-seq analysis of the HPAP dataset which identified a subset of

pericytes, marked by *PDGFRB*, that express moderate levels of *ACE2* (**Figure 1B**). In addition,

our findings support recent reports of pericyte-specific vascular expression of ACE2 in brain and

heart tissue (Chen et al., 2020; He, 2020). Emerging evidence from both pre-clinical models of

SARS-CoV-2 infection (Aid, 2020) and patients with COVID-19 indicate that cells of the

microvasculature (e.g. endothelial cells and pericytes) are requisite contributors to the initiation

and propagation of severe disease (Teuwen et al., 2020). Although we found that *TMPRSS2*

expression was negligible in endothelial and stellate cells (which includes pericytes) of human

islets, *CTSL*, *ADAM17*, *FURIN* and *ACE2* were moderately to highly expressed in these

populations (**Figure 1B**), raising the intriguing possibility of direct infection of cells in the islet

microvasculature by SARS-CoV-2. Such an occurrence could trigger β cell dysfunction and the

metabolic sequelae of COVID-19. Future studies addressing this possibility in human islets and

pancreatic tissue are urgently needed.

Both ACE2 and TMPRSS2 protein are present in pancreatic ducts but rarely are co-

expressed.

Our work indicated that TMPRSS2 protein was localized to the apical surface of intercalated

and larger ducts in ND (**Figures 2B, 2C", 2E, 2F"** and **S3E-H**), T2D (**Figures 2H, 2I", 2K, 2L"**

and **S3M-N**), and T1D pancreatic tissues (**Figures 2N, 2O", 2Q, 2R"** and **S3S-V**). To further

define ductal expression of TMPRSS2 and determine whether ACE2 was expressed beyond

exocrine tissue capillaries, we visualized expression of these two proteins in relationship to

cytokeratin-19 protein (KRT19), a ductal cell marker. TMPRSS2, but not ACE2, was expressed

by KRT19-positive cells on the apical surface of intercalated and larger ducts throughout the

exocrine compartment (**Figure 5A-D'**). Rarely, both ACE2 and TMPRSS2 were found on the

apical surface of ductal epithelial cells but were spatially distinct (**Figure 5E-H'**). These findings

agree with our scRNA-seq analysis in which *TMPRSS2* was more highly expressed in ductal

cells than *ACE2*, but cells positive for both *ACE2* and *TMPRSS2* were rare. In addition,

although *TMPRSS2* mRNA was detected in acinar cells by scRNA-seq (**Figure 1B**), TMPRSS2

protein was undetectable by immunofluorescence analysis (**Figure 5**), indicating that studies of

mRNA and protein may be discordant.

Because rare cells in the exocrine compartment co-express ACE2 and TMPRSS2, it is

possible that SARS-CoV-2 could infect those cells, induce pancreatitis, and consequently

impact islet function. While recent case reports of patients with COVID-19 have noted mild

elevations of amylase and lipase or frank pancreatitis (Karimzadeh et al., 2020; Meireles et al.,

2020; Rabice et al., 2020; Wang et al., 2020), the vast majority of patients with COVID-19 have

neither elevated pancreatic enzymes nor frank pancreatitis (Ashok et al., 2020; Bonney et al.,

2020; Bruno et al., 2020). To investigate this possibility, we analyzed the histology of pancreatic

tissue obtained at autopsy from seven COVID-19 positive patients, three of whom had diabetes.

We did not find signs of pancreatitis, interstitial edema, inflammatory infiltrate, hemorrhage or

necrosis (**Figure 5I**). It is important that exocrine inflammation and ACE2/TMPRSS2 expression

be analyzed in a larger number of COVID-19 autopsy samples, particularly in those with new-

onset diabetes. Such studies will be challenging since the pancreas rapidly undergoes autolysis

upon death.

In summary, recent reviews, commentaries, and clinical guidelines (Apicella et al., 2020;

Bornstein et al., 2020; Goldman et al., 2020; Gupta et al., 2020; Heaney et al., 2020) have

highlighted an attractive hypothesis of direct cytotoxicity by SARS-CoV-2 to β cells. Here, we

addressed the fundamental question of whether canonical SARS CoV-2 cell entry machinery is

present in β cells of human pancreatic tissue. By examining pancreatic ACE2 and TMPRSS2

expression in normal donors and in those with diabetes, we found that, **1)** β cells do not

coexpress *ACE2* and *TMPRSS2* transcripts by scRNA-seq analysis and neither protein is

detectable in α or β cells by immunofluorescence; **2)** ACE2 is primarily expressed in the islet

and exocrine tissue capillaries, including pericytes, as well as a subset of ductal cells; **3)**

TMPRSS2 is primarily expressed in ductal cells; **4)** ACE2 and TMPRSS2 are infrequently co-

expressed in pancreatic ducts; **5)** ACE2 and TMPRSS2 protein expression and distribution

appear similar in ND and T2D pancreata but studies on additional pancreata are needed to

definitively answer this question; and **6)** Alternative pathways facilitating SARS-CoV-2 viral

infection in β cells cannot be excluded. These findings align considerably with independent

observations from Kusmartseva and colleagues (Kusmartseva, 2020) and do not support the

hypothesis that SARS-CoV-2 binds and infects islet β cells via a canonical pathway mediated by

ACE2 and TMPRSS2, eliciting a diabetogenic effect. As outlined below in the "Limitations of the

Study" section, additional studies are needed to investigate whether direct SARS-CoV-2

infection of β cells occurs or is detrimental to β cell health or function by other mechanisms.

However, based on current data, it appears that the interaction of diabetes and SARS-CoV-2 is

mediated by systemic inflammation and/or metabolic changes in other organs such as liver,

muscle or adipose tissue.

Limitations of the Study

We did not directly measure SARS-CoV-2 binding or entry into human β cells but instead

assessed expression of canonical co-receptors, ACE2 and TMPRSS2, in the human pancreas.

While studies culturing SARS-CoV-2 and human islets *in vitro* will be of considerable interest,

such experiments may be challenging to interpret as the islet isolation process itself may impact

ACE2 or TMPRSS2 expression or susceptibility to viral infection. In addition, infection of islet

cells within an intact, isolated islet with a variety of viruses has proven difficult as is selection of

an appropriate ratio of viral particles and islet cells reflective of the *in vivo* environment.

Furthermore, it is not clear if infection of an isolated islet accurately models the physiology of a

vascularized, innervated islet within the context of the pancreas.

While the preponderance of research suggests that ACE2 and TMPRSS2 are the primary

means for SARS-CoV-2 entry into host cells, knowledge about additional pathways and/or

mechanisms of SARS-CoV-2 cell entry and infection is rapidly evolving. We sought, but did not

find evidence, suggesting that other proposed pathways (namely *DPP4*, *CTSL*, *ADAM17*,

FURIN or *TMPRSS4*) are involved in β cell entry.

We did not assess the presence of SARS-CoV-2, its proteins, or RNA in the pancreas from

COVID-19-infected individuals. It is possible that SARS-CoV-2 could infect pancreatic exocrine

cells, ductal cells, endothelial cells, or microvasculature, leading to pancreatic inflammation that

negatively impacts β cell health or function. We did not find evidence of inflammation in the

pancreas of seven COVID-19-infected patients (three of whom had diabetes). Future efforts are

needed to collect a large number of pancreata from COVID-19-infected individuals with and

without diabetes (long-standing and recent-onset) to search for presence of SARS-CoV-2 or

inflammation (islet or exocrine). Since the pancreas undergoes rapid autolysis after death, care

must be taken to collect the pancreas very soon after death for this analysis.

Acknowledgements

We are especially thankful to organ donors and their families. This research was supported by

funding provided by the National Institute of Diabetes and Digestive and Kidney Diseases, the

Human Islet Research Network (HIRN; RRID:SCR_014393; https://hirnetwork.org; DK112232,

330 DK123716, U01DK123594, DK104211, DK108120), and by DK106755, DK117147, DK117147-

331 01A1S1, DK111757, DK112217, DK20593, and the Functional Genomics Core of DK19525.

332 This manuscript used data acquired from and available at the Human Pancreas Analysis

333 Program (HPAP-RRID:SCR_016202) Database (https://hpap.pmacs.upenn.edu), a HIRN

334 consortium. This work was also supported by grants from the Doris Duke Charitable Foundation

335 (DDCF 4043516256), Human Islet Research Network New Investigator Pilot Award (UC4

336 DK104162), JDRF, The Leona M. and Harry B. Helmsley Charitable Trust, and the Department

337 of Veterans Affairs (BX000666). Human pancreatic islets were provided by the NIDDK-funded

338 Integrated Islet Distribution Program at the City of Hope (NIH Grant # 2UC4 DK098085 RRID:

339 SCR_014387; http://iidp.coh.org). Human kidney sections were provided by Dr. Agnes B. Fogo

340 at Vanderbilt University Medical Center. We thank Amber M. Bradley for technical assistance.

342 **Author Contributions:** K.C. J.C., M.B., and A.C.P. designed the experiments. K.C., J.C., M.B.,

343 and A.C.P. wrote the manuscript. K.C., J.C., S.S., W.W., L.G. J.A., M.E.K., M.F., A.M., C.D.,

344 D.S., R.B., R.J., R.W.S., K.L.H., G.V., M.B., and A.C.P. performed experiments or analyzed the

345 data. All authors reviewed and edited the final manuscript. As the lead contact, A.C.P. is

346 responsible for: 1) communication with the journal and co-authors; 2) addressing requests for

347 reagents and resources; and 3) overseeing decisions and disputes related to the manuscript.

349 **Declaration of Interests:** The authors declare no conflicts of interests.

Figure 1. *ACE2* and *TMPRSS2* Expression is Minimal in Isolated Human Islet α and

β Cells. (A) Relative expression of *ACE2* and *TMPRSS2* compared with select α (white bars)

and β (green bars) cell type-enriched genes in sorted human islet α and β cells from previously

published bulk RNA-seq datasets, reported as transcript per million mapped reads (TPM; n=7;

(Blodgett et al., 2015)) or reads per kilobase of transcript per million mapped reads (RPKM;

n=8; (Arda et al., 2016)). Mean expression values are presented as $\log_2$ (TPM+1) or $\log_2$

(RPKM+1) to account for negative values. Dotted line highlights *ACE2* and *TMPRSS2*

expression. (**B**) Dot plots of *ACE2*, *TMPRSS2*, *CTSL*, *ADAM17*, *FURIN*, *TMPRSS4*, and *DPP4*

expression compared with cell type-enriched genes from three single cell (sc) RNA-seq

datasets (Baron et al., 2016; Kaestner et al., 2019; Segerstolpe et al., 2016). Dot size indicates

percentage of cells in a given population expressing the gene; dot color represents scaled

average expression. Dotted lines highlight expression of the putative SARS-CoV-2 entry

machinery. Percentages of β cells expressing and co-expressing these genes are available in

Table S1. (**C**) Representative images showing an isolated islet from an adult human donor (top

panels) and a pancreatic section from a juvenile human donor (bottom panels) stained for ACE2

(red; antibody ab15348), insulin (INS; green), glucagon (GCG; white), and DAPI (blue). Dotted

yellow line denotes islet area. Scale bar is 50 µm. Human pancreatic donor information is

available in Table S2 (1C, donors I1-I3, J1-J5). See also Figures S1, S2, and S4.

Figure 2. ACE2 and TMPRSS2 Protein are Not Detected in Human Islet α or β Cells in

Adult Pancreas. Immunostaining does not detect SARS-CoV-2 cell entry markers ACE2 (**A**, **D**,

G, **J**, **M**, **P**; antibody AF933; in red) or TMPRSS2 (**B**, **E**, **H**, **K**, **N**, **Q;** in green), in islet β cells

(INS, blue; **C-C'**, **I-I'**, **O-O'**) or islet α cells (GCG; blue; **F-F'**, **L-L'**, **R-R'**) in native pancreatic

sections from adult donors without diabetes (**A-F"**) or from donors with type 2 (**G-L"**) or type 1

(**M-R"**) diabetes. Dotted yellow lines denote islets. Islet (i) and exocrine (e) inset areas are

marked by yellow boxes in MERGE column with DAPI counterstain. Pancreatic ducts, identified

structurally by rosette pattern of DAPI-labeled nuclei (white), are shown within dotted yellow

lines in exocrine INSET column. Scale bars are 50 μm (**A-R**) and 25 μm (Insets, **C'-R"**). Human

pancreatic donor information is available in Table S2 (**A-F"**, donor N9; **G-L"**, donor 2H; **M-R"**,

donor 1C). See also Figures S2, S3, and S4.

Figure 3. ACE2 Protein is Localized to Islet and Exocrine Capillaries in Adult and Juvenile

Human Pancreata. Representative images of endothelial cells (CD31, white) and ACE2-

positive perivascular cells (red; antibody ab15348) in the adult islet (**A-D'**) and exocrine (**E-H'**)

microvasculature, as well as in juvenile human islet (**I-L'**) microvasculature. Yellow arrowheads

point to CD31-positive endothelial cells, while magenta arrowheads point to perivascular ACE2-

positive cells. ACE2-positive perivascular cells abut the extracellular matrix marker collagen-IV

within the vascular basement membrane (collagen-IV [COL4], green; **D, D', H, H'**). Inset areas

(**A'-L'**) are marked by yellow boxes in **A-L**. In juvenile human pancreas (**I-L'**), ACE2 is not

detected in islet endocrine cells expressing insulin and glucagon (ENDO, green); DAPI (blue).

Scale bars are 50 μm (**A-L**) and 10 μm (Insets, **A'-L'**). Human pancreatic donor information is

available in Table S2 (**A-H'**, donor N2; **I-L'**: donor J4). See also Figure S5.

Figure 4. ACE2 Protein is Expressed by Islet Pericytes. (**A-C**) Maximum intensity projection

of representative confocal images labelled for platelet-derived growth factor receptor β-positive

cells (PDGFRβ, red) and ACE2 (green) in perivascular cells within a normal adult human islet;

nuclear DAPI (blue). (**D**) Image of the same islet shown in A-C with insulin labeling (blue).

Region within the yellow box in **D** is displayed in **A-C**. (**E-G**) Maximal intensity projection of

pericyte labeling with an antibody against the pericyte marker neuron-glial antigen 2 (**E**, NG2,

red) colocalized with ACE2 labeling (**F**, green) and nuclear DAPI (blue) in a normal adult human

islet (**E-F**). (**G**) Image of the same islet shown in E-F with insulin labeling (blue). Regions within

yellow boxes containing pericytes in **G** denote inset areas (**G'**) and represent a subset of islet

ACE2-expressing pericytes. (**H**) Colocalization of ACE2 with NG2- or PDGFRβ-positive cells. (**I**)

Quantification of ACE2-positive cells expressing pericyte markers NG2 and PDGFRβ. Shown

are average values obtained for 3 confocal planes per islet for a minimum of 5 islets per each

donor of a given age. Scale bars are 20 μm (**A-C**), 40 μm (**D-G**), and 10 μm (**G'**). Human

pancreatic donor information is available in Table S2 (**A-I,** donors HP1754, HP2041, HP2091).

See also Figure S5.

Figure 5. Analysis of ACE2 and TMPRSS2 Protein Expression in Pancreatic Ducts and

Pancreatic Histology. (**A-D'**) TMPRSS2 (green), but not ACE2 (red; antibody R&D AF933), is

expressed by KRT19-positive cells (blue) on the apical surface of intercalated and larger ducts

throughout the exocrine compartment. DAPI (white). Yellow arrowheads point to TMPRSS2 on

the apical surface of ductal cells (KRT19-positive). Magenta arrowheads point to ACE2-positive

non-ductal cells. (**E-H'**) Rarely, both TMPRSS2 (green) and ACE2 (red; antibody R&D AF933)

were expressed on the apical surface of ductal epithelial cells (KRT19, blue), but appeared to

be spatially distinct. Yellow and white arrowheads point to spatially distinct TMPRSS2 and

ACE2 labeling, respectively, localized to the apical surface of larger ducts (KRT19-positive

cells). Inset areas (**A'-H'**) are marked by yellow boxes in **A-H**. Scale bars are 50 μm (**A-H**) and

10 μm (**A'-H'**). (**I**), Representative H&E image of human pancreas upon autopsy after COVID-19

disease. None of the seven donor samples evaluated showed signs of pancreatitis as

determined by the histological absence of interstitial edema and inflammatory infiltrate,

hemorrhage or necrosis. BV, blood vessel; D, duct. Scale bar in green is 100 μm. Human

pancreatic donor information is available in Table S2 (**A-D'**, donor N8; **E-H'**, donor N9; **I**,

COVID-19 donor 1). See also Figure S3.

STAR METHODS

Resource Availability

Lead contact. Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Alvin C. Powers (al.powers@vumc.org).

Materials Availability. This study did not generate new unique reagents.

Data and Code Availability. All data generated or analyzed during this study are included in this published article or in the data repositories listed in the Key Resources Table. Original/source data for Figure 1A (bulk RNA-seq of FACS sorted human islet α and β cells) is available under NCBI GEO accession numbers GSE67543 (Blodgett et al., 2015) and GSE57973 (Arda et al., 2016). Original/source data for Figure 1B and Figure S4J (single cell RNA-seq of human islets) is available under NCBI GEO accession number GSE84133 (Baron et al., 2016), GSE124742 (Camunas-Soler et al., 2020), and ArrayExpress (EBI) accession number E-MTAB-5061 (Segerstolpe et al., 2016). This manuscript used data acquired from the Human Pancreas Analysis Program (HPAP) Database (https://hpap.pmacs.upenn.edu), a Human Islet Research Network consortium.

Experimental Model and Subject Details

Human Subjects. Pancreata and islets from juvenile, adult, T1D and T2D human donors were obtained through partnerships with the International Institute for Advancement of Medicine (IIAM), National Disease Research Interchange (NDRI), Integrated Islet Distribution Program

(IIDP), and local organ procurement organizations. Pancreata from normal donors were

processed either for islet isolation (Balamurugan et al., 2003) and/or histological analysis as

described previously (described below and in Table S2) (Brissova et al., 2018; Dai et al., 2017).

Pancreata from COVID-19 decedents after autopsy were obtained from the Translational

Pathology Shared Resource (TPSR) at Vanderbilt University Medical Center (Nashville, TN) and

and processed for histological analysis as according to standard procedures for clinical

diagnostics (VUMC Histology Peloris Processing Protocol). Samples from donors of both sexes

were used in our analyses. Donor demographic information is summarized in Table S2. The

Vanderbilt University Institutional Review Board does not consider de-identified human

pancreatic specimens to be human subjects research.

METHOD DETAILS

Human Pancreas Procurement and Preparation of Tissue for Histological Analysis.

Pancreata from juvenile, adult, T1D and T2D donors were obtained within 18 hours from cross-

clamp and maintained in cold preservation solution on ice until processing, as described

previously (Balamurugan et al., 2003; Brissova et al., 2018). Pancreata recovered from seven

COVID-19 decedants were obtained from Vanderbilt University Medical Center (Nashville, TN)

within 8-27 hours of death. Donor demographic information is summarized in Table S2. Human

kidney samples were provided by Dr. Agnes Fogo, Vanderbilt University. The Vanderbilt

University Institutional Review Board has declared that studies on de-identified human

pancreatic specimens do not quality as human subjects research.

Mouse Tissue Preparation for Histological Analysis. Mice were maintained on standard

rodent chow under a 12-hour light/dark cycle. Kidney from adult NOD.$Cg\text{-}Prkdc^{scid}Il2rg^{tm1Wjl}$/ Sz

(NSG) mice ages 12 to 18 weeks (Jackson Laboratory) were isolated. Tissue specimens were

processed for cryosections as described previously (Brissova et al., 2018).

Immunohistochemical Analysis. Immunohistochemical analysis was performed on 5-µm

cryosections (Figure 1C; Figure 3 and 4; Figure S2:A-I; S3A-D, I-L and O-R; S4 and S5) or

formalin-fixed paraffin embedded (FFPE) pancreatic sections (Figure 2 and 5; Figure S2:J-W;

S3E-H, M-N and S-V) as indicated and described previously (Brissova et al., 2014; Brissova et

al., 2018; Wright et al., 2020). FFPE sections were first deparaffinized in xylene and ethanol

followed by heat-induced epitope retrieval in a citrate-based antigen unmasking solution, pH 6.0

(see Key Resource Table). Immunofluorescence analysis of isolated islets was performed on 8-

µm cryosections of islets embedded in collagen gels as described previously (Brissova et al.,

2005; Brissova et al., 2018). Primary antibodies to all antigens and their working dilutions are

listed in the Key Resource Table. The antigens were visualized using appropriate secondary

antibodies listed in the Key Resource Table. Digital images were acquired with an Olympus

FLUOVIEW FV3000 laser scanning confocal microscope (Olympus Corporation).

Peptide Competition. We combined 1µg of anti-ACE2 antibody (ab15348) with or without 10µg

of the immunizing human ACE2 peptide (ab15352) in antibody buffer solution (0.1% Triton X-

100/1% BSA/1X PBS) and incubated overnight at 4°C with gentle agitation.

Immunofluorescence staining of 5-µm serial cryosections from the same donor was performed

as described above. One section was incubated with the neutralized antibody buffer while the

other section was incubated with the non-neutralized antibody buffer. ACE2 staining patterns

were visualized via confocal microscopy as described above.

H&E Staining. FFPE pancreas tissue sections (5-µm sections, n=7 donors) were processed

according to standard procedures for clinical diagnostics (Vanderbilt University Medical Center

Histology Peloris Processing Protocols). Slides were stained with hematoxylin & eosin and

reviewed by a clinical pathologist (M.E.K.) at Vanderbilt University Medical Center.

Bulk RNA-seq Data Acquisition. Transcripts Per Million (TPM) and Reads Per Kilobase Per

Million (RPKM) normalized counts were extracted from publicly available RNA-seq datasets

(Arda et al., 2016; Blodgett et al., 2015). The sources of the datasets are summarized in the Key

Resources Table. GraphPad Prism v8 was used to generate plots in Figure 1A.

Single Cell RNA-seq Data Acquisition. Raw gene count matrices were extracted from existing

single cell RNA-seq datasets (Baron et al., 2016; Segerstolpe et al., 2016; Yang et al., 2020)

and from the Human Pancreas Analysis Program (HPAP) Database

(https://hpap.pmacs.upenn.edu), a Human Islet Research Network consortium. The sources of

the datasets are summarized in the Key Resources Table. Gene count matrices were further

analyzed using the R package Seurat version 3.1 (Stuart et al., 2019). Gene count

measurements were normalized for each cell by library size and log-transformed using a size

factor of 10,000 molecules per cell. The data was further scaled to unit variance and zero mean

implemented in the "ScaleData" function. Cell types already annotated by the authors in the

original study were used and thus, no clustering was performed. Seurat's "DotPlot" function was

used to generate plots shown in Figures 1B and S4J to visualize *ACE2* and *TMPRSS*2 scaled

expression.

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistical analysis. Bulk RNA-seq data are expressed as mean ± standard error of mean

(Figures 1A). Sample size (n) is provided as the number of independent donor samples. A p-

value less than 0.05 was considered significant. Statistical analysis (unpaired t-test) was

performed using GraphPad Prism software. Mander's coefficients were determined to quantify

colocalization between pericyte markers (NG2 and PDGFRβ) and ACE2 in confocal images of

islets using the ImageJ plugin "Just Another Co-localization Plugin":

(https://imagej.nih.gov/ij/plugins/track/jacop2.html).

ADDITIONAL RESOURCES

None.

Figure S1. Related to Figure 1. Stratification of *ACE2*, *TMPRSS2*, and *ADAM17*

Expression in β cells by BMI. β cell gene expression from eleven donors (ages 1-39 years)

from the HPAP scRNA-seq dataset (Kaestner et al., 2019) are displayed according to increasing

BMI. Only one donor had an obese BMI. (**A-B**) *ACE2* and *TMPRSS2* expression in β cells did

not show correlation with increasing BMI. (**C**) A trend towards increased *ADAM17* expression

with BMI was identified. However, only one obese donor was available and analyzed in this

dataset. Human pancreatic donor information is available in Table S2.

Figure S2. Related to Figure 1 and 2. Testing and Characterization of Four ACE2-directed

Antibodies on Human Pancreatic Tissue. (**A-E**) Characterization of ACE2 antibody (red,

ab15348) used by Yang et al. (Yang et al., 2020). Antibody epitope encompasses the ACE2 C-

terminal domain (human aa 788-805). Mouse kidney tissue served as a positive control for

ACE2 (**A**), while normal adult human pancreatic tissue incubated with anti-rabbit-Cy3 secondary

antibody only served as a negative control (**B**). Normal adult human pancreas labeled for ACE2

(red), INS (green, β cells) and GCG (blue, α cells) (**C-E**). Inset area is marked by a yellow box in

MERGE column. (**F-I**) Competition with immunizing peptide neutralized ACE2 labeling,

demonstrating the antibody's antigen specificity. Scale bars are 100 μm (**A-E**) and 50 μm (Inset,

E and **F-I**). (**J-N**) Characterization of ACE2 antibody (red, R&D MAB933) at same dilution (1:33)

reported by Fignani et al. (preprint: Fignani, 2020). Antibody epitope encompasses the ACE2

extracellular domain (human aa 18-740). Human kidney tissue served as a positive control for

ACE2 (**J**), while normal adult human pancreatic tissue incubated with anti-mouse-Cy3

secondary antibody only served as a negative control (**K**). Normal adult human pancreas

labeled for ACE2 (red), INS (green, β cells) and GCG (blue, α cells) (**L-N**). Inset area is marked

by a yellow box in MERGE column. Scale bars are 50 μm (**J-N**) and 25 μm (Inset, **N**). (**O-S**)

Characterization of ACE2 antibody (red, R&D AF933) at same dilution (1:200) reported by Yang

et al. (Yang et al., 2020). Antibody epitope encompasses the ACE2 extracellular domain

(human aa 18-740). Human kidney served as a positive control for ACE2 (**O**), while normal adult

human pancreatic tissue incubated with anti-goat-Cy3 secondary antibody only served as a

negative control (**P**). Normal adult human pancreas labeled for ACE2 (red), INS (green, β cells)

and GCG (blue, α cells) (**Q-S**). Inset area is marked by a yellow box in MERGE column. Scale

bars are 50 μm (**O-R**) and 25 μm (Inset, **S**). (**T-W**) Characterization of ACE2 antibody (red,

HPA000288) used by the Human Protein Atlas (Uhlen et al., 2015) and Hikmet et al. (Hikmet et

al., 2020). Antibody epitope encompasses the ACE2 extracellular domain (human aa 1-111).

Human kidney tissue served as a positive control for ACE2 (**T**). Normal adult human pancreas

labeled for ACE2 (red) and INS (green, β cells) (**U-W**). Inset area is marked by a white dashed

box in MERGE column. Scale bars are 100 μm (**T-V**) and 50 μm (Inset, **W**). DAPI (white).

Human pancreatic donor information is available in Table S2 (**B**, donor N8; **C-E**, donor N4; **F-I**,

donors N6 and N2; **J-N**, donor N2; **O-S**, donor N7; **T-W**, donor N8).

Figure S3. Related to Figures 2 and 5. ACE2 and TMPRSS2 Protein are Not Detected by

Immunofluorescence in α or β Cells from Normal, T2D or T1D Adult Donors. SARS-CoV-2

cell entry markers ACE2 (antibody ab15348) and TMPRSS2, both shown in red, are not

detected in islet α cells (GCG, blue) or β cells (INS, green) in pancreatic sections from adult

donors without diabetes (**A-H**) or donors with type 2 (**I-N**) or type 1 (**O-V**) diabetes. Insets are

depicted by a yellow box. DAPI (white). Scale bars are 100 μm (**A-V**) and 25 μm (Insets).

Human islet and pancreatic donor information is available in Table S2 (**A-D**, donors N3, N7, N9,

N8; **E-H**, donors N14, N12, N11, N10; **I-L**, donors 2L, 2B, 2G, 2I; **M-N**, donors 2H, 2G; **O-R**,

donors 1B, 1D, 1C, 1A; **S-V**, donors 1H, 1K, 1J, 1G).

Figure S4. Related to Figures 1 and 2. DPP4 is Expressed in α cells of Adult Normal and

Diabetic Donors. DPP4 (red) is expressed in human pancreatic islet α cells (GCG, blue;

Merged, magenta) but not β cells (INS, green) in pancreatic sections from adult donors without

diabetes (**A-C**) or donors with type 2 (**D-F**) or type 1 (**G-I**) diabetes. Scale bars are 50 μm (**A-I**).

Human islet and pancreatic donor information is available in Table S2 (**A-C**, donor N8; **D-F**,

donor 2K; **G-I**, donor 1H). (**J**) Dot plots of *ACE2*, *TMPRSS2*, *CTSL*, *ADAM17*, *FURIN*,

TMPRSS4, and *DPP4* expression compared with cell type-enriched genes from a previously

published single cell (sc) RNA-seq datasets (Camunas-Soler et al., 2020). Dot size indicates

percentage of cells in a given population expressing the gene; dot color represents scaled

average expression. Dotted line highlights *ACE2*, *TMPRSS2*, *CTSL*, *ADAM17*, *FURIN*,

TMPRSS4, and *DPP4* expression.

Figure S5. Related to Figures 3 and 4. ACE2 is Localized to Islet and Exocrine Capillaries

in Adult Human Pancreas of T2D and T1D donors. Representative images of endothelial

cells (CD31, white) and ACE2-positive perivascular cells (red; antibody ab15348) in islet (**A-H'**)

and exocrine tissue microvasculature (**I-P'**) of individuals with type 2 and type 1 diabetes. DAPI

(white). Yellow arrowheads point to CD31-positive endothelial cells, while magenta arrowheads

point to perivascular ACE2-positive cells. Insets (**A'-P'**) are depicted by yellow boxes in **A-P**.

ACE2 positive perivascular cells abut extracellular matrix marker collagen-IV within the vascular

basement membrane (collagen-IV [COL4], green; **D, D', H, H', L, L', P, P'**). ACE2 labeling was

reduced in both islets (**E-H'**) and exocrine tissue of individuals with type 1 diabetes (**M-P'**)

compared to normal donors (Figure 3) and those with type 2 diabetes. Scale bars are 50 μm (**A-**

P) and 10 μm (Insets, **A'-P'**). Human pancreatic donor information is available in Table S2 (**A-**

D', donor 2E; **E-H'**, donor 1F; **I-L'**, donor 2E; **M-P'**, donor 1C).

Table S1. Related to Figure 1. Number and Percentage of β cells that Express and Co-

express Putative SARS-CoV-2 Cell Entry Genes Across Four Independent scRNA-seq

Datasets.

Genes	Droplet-based scRNA-seq				SMART-seq			
	HPAP[a] (β cell total, N=2828)		Baron et al.[b] (β cell total, N=2525)		Segerstolpe et al.[c] (β cell total, N=157)		Camunas-Soler et al.[c] (β cell total, N=357)	
	# β cell	% β cells	# β cell	% β cells	# β cell	% β cells	# β cell	% β cells
ACE2	17	0.6	4	0.2	3	1.9	12	3.4
TMPRSS2	60	2.1	7	0.3	4	2.5	2	0.6
TMPRSS4	0	0.0	0	0.0	0	0.0	9	2.5
CTSL	1421	50.2	977	38.7	132	84.1	291	81.5
FURIN	779	27.5	942	37.3	91	58.0	255	71.4
ADAM17	494	17.5	251	9.9	78	49.7	96	26.9
ACE2, TMPRSS2	0	0.0	0	0.0	0	0.0	0	0.0
ACE2,TMPRSS4	0	0.0	0	0.0	0	0.0	0	0.0
ACE2, CTSL	0	0.0	2	0.1	3	1.9	9	2.5
ACE2, FURIN	0	0.0	1	0.0	1	0.6	8	2.2
ACE2, ADAM17	0	0.0	0	0.0	1	0.6	3	0.8

614　[a]10x genomics; [b]InDrop (Klein et al., 2015); [c]SMART-seq2 (Picelli et al., 2014)

627
628

629 **Table S2. Related to STAR Methods. Demographic Information of Donors.**

Donor ID		Age	Ethnicity / Race	Diabetes Duration	Sex	BMI	Cause of Death	Tissue/Islet Source
Juvenile (Histology)	J1	5 days	Caucasian	--	F	14.9	Anoxia	IIAM
	J2	3 months	Caucasian	--	M	16.8	Anoxia	NDRI
	J3	10 months	Caucasian	--	F	15.4	CVA	NDRI
	J4	20 months	Caucasian	--	F	23.5	Anoxia	IIAM
	J5	5 years	Caucasian	--	M	16.2	Anoxia	IIAM
Normal Adult (Histology)	N1	42 years	Caucasian	--	M	32.2	Overdose	TNDS
	N2	45 years	Caucasian	--	F	29.7	Anoxia	OPO
	N3	46 years	Caucasian	--	F	32.9	CVA	IIAM
	N4	48 years	Caucasian	--	M	24.6	Anoxia	OPO
	N5	51 years	Caucasian	--	M	20.4	Anoxia	OPO
	N6	52 years	Black	--	M	29.2	ICH	TNDS
	N7	52 years	Caucasian	--	M	28.1	Head Trauma	OPO
	N8	55 years	Black	--	M	35.6	CVA	IIAM
	N9	59 years	Caucasian	--	M	32.7	Head Trauma	IIAM
	N10	20 years	Hispanic	--	M	19.4	Head Trauma	IIAM
	N11	24 years	Caucasian	--	M	35.5	ICH	IIAM
	N12	35 years	Caucasian	--	M	26.8	Head Trauma	IIAM
	N13	20 years	Caucasian	--	M	27.8	Head Trauma	NDRI
	N14	18 years	Caucasian	--	M	25.1	Head Trauma	IIAM
	HP1754	15 years	N/A	--	M	22.6	Head Trauma	IIAM
	HP2041	29 years	N/A	--	M	22.3	Head Trauma	IIAM
	HP2091	44 years	N/A	--	F	23.7	CVA	IIAM
Adult T1D (Histology)	1A	43 years	N/A	36 years	M	31.2	CVA	NDRI
	1B	45 years	Caucasian	43 years	M	25.0	Anoxia	IIAM
	1C	54 years	Caucasian	14 years	F	24.9	Anoxia	IIAM
	1D	57 years	Black	45 years	M	33.3	CVA	IIAM
	1E	58 years	Caucasian	31 years	M	21.8	Anoxia	NDRI
	1F	63 years	Caucasian	44 years	M	24.1	Anoxia	IIAM
	1G	35 years	Caucasian	23 years	M	26.9	Anoxia	NDRI
	1H	20 years	Caucasian	7 years	M	25.5	Anoxia	NDRI
	1I	27 years	Caucasian	17 years	M	18.4	Anoxia	NDRI
	1J	13 years	Caucasian	5 years	M	19.1	Anoxia	IIAM
	1K	30 years	Caucasian	20 years	M	29.8	Anoxia	NDRI
Adult T2D (Histology)	2A	44 years	Caucasian	7 years	M	44.4	CVA	IIAM
	2B	52 years	Caucasian	7 years	M	33.6	CVA	IIAM
	2C	52 years	Asian	10 years	F	21.9	CVA	NDRI
	2D	52 years	Caucasian	< 1 year	F	29.2	CVA	IIAM
	2E	42 years	Black	< 1 year	M	42.0	CVA	IIAM

	2F	43 years	Black	1 year	M	36.0	Head Trauma	IIAM
	2G	66 years	Caucasian	3 years	F	32.8	CVA	IIAM
	2H	47 years	Caucasian	3 years	M	31.3	CVA	IIAM
	2I	64 years	Caucasian	5 years	M	33.2	ICH	IIAM
	2J	59 years	Caucasian	6 years	F	27.5	CVA	IIAM
	2K	60 years	Caucasian	1 year	M	38.3	CVA	IIAM
	2L	49 years	Caucasian	3 years	F	33.8	CVA	IIAM
Normal Adult Islets (Gels and scRNA-Seq)	I1	40 years	Caucasian	--	F	30.8	Head Trauma	IIDP
	I2	41 years	N/A	--	M	20.3	N/A	IIDP
	I3	42 years	Caucasian	--	M	32.2	Overdose	IIDP
	HPAP022	39 years	Caucasian	--	F	34.7	Anoxia	HPAP
	HPAP026	24 years	Caucasian	--	M	20.8	Anoxia	HPAP
	HPAP034	13 years	Caucasian	--	M	18.6	Head Trauma	HPAP
	HPAP035	35 years	Caucasian	--	M	26.9	Anoxia	HPAP
	HPAP036	23 years	Caucasian	--	F	16	Head Trauma	HPAP
	HPAP037	35 years	Caucasian	--	F	21.9	CVA	HPAP
	HPAP039	5 years	Caucasian	--	F	16.3	Anoxia	HPAP
	HPAP040	35 years	Caucasian	--	M	23.9	CVA	HPAP
	HPAP042	1 year	Caucasian	--	M	17.9	Anoxia	HPAP
	HPAP044	3 years	Caucasian	--	F	12	Anoxia	HPAP
	HPAP047	8 years	Caucasian	--	M	16.8	CVA	HPAP
COVID-19 Patient Autopsy Samples (Histology)	1	82 years	Caucasian	--	M	26.8	ALI	VUMC Autopsy
	2	97 years	Caucasian	--	F	19.7	ALI	VUMC Autopsy
	3	81 years	Caucasian	>10 years[a]	M	23.3	ALI	VUMC Autopsy
	4	60 years	Hispanic	--	M	36.7	ALI	VUMC Autopsy
	5	51 years	Hispanic	23 years	M	29.4	ALI	VUMC Autopsy
	6	60 years	Caucasian	--	F	38.4	PE	VUMC Autopsy
	7	71 years	Black	Pre-existing[b]	M	31.5	ALI	VUMC Autopsy

ALI – acute lung injury; CVA, cerebrovascular accident; HPAP – Human Pancreas Analysis Program (Human Islet Research Network); ICH, intracerebral hemorrhage; IIAM – International Institute for the Advancement of Medicine; IIDP – Integrated Islet Distribution Program; N/A – not available; NDRI – National Disease Research Interchange; OPO – Organ Procurement Organization; PE – pulmonary embolism; T1D = type 1 diabetes; T2D – type 2 diabetes; TNDS – Tennessee Donor Services, Nashville; VUMC Autopsy – Vanderbilt University Medical Center Autopsy Pathology

[a]Oldest clinical patient note including diagnosis of diabetes mellitus was signed in 2010, suggesting disease duration of at least 10 years.

[b]Patient was prescribed an oral anti-diabetic medication confirming pre-existing diabetes diagnosis of unknown duration prior to admission with COVID-19.

630
631

REFERENCES

Aid, M., Busman-Sahay, K., Vidal, S., Maliga, Z., Bondoc, S., Starke, C., Terry, M., Jacobson,
C., Wrijil, L., Ducat, S., Brook, O., Miller, A., Porto, M., Pellegrini, K., Pino, M., Hoang, T.,
Chandrashekar, A., Patel, S., Stephenson, K., Bosinger, S., Andersen, H., Lewis, M., Hecht, J.,
Sorger, P., Martinot, A., Estes, J., Barouch, D. (2020). Vascular disease and thrombosis in
SARS-CoV-2 infected Rhesus macaques. Cell *In Press*.

Almaca, J., Caicedo, A., and Landsman, L. (2020). Beta cell dysfunction in diabetes: the islet
microenvironment as an unusual suspect. Diabetologia *63*, 2076-2085.

Almaca, J., Weitz, J., Rodriguez-Diaz, R., Pereira, E., and Caicedo, A. (2018). The pericyte of
the pancreatic islet regulates capillary diameter and local blood flow. Cell Metab *27*, 630-644
e634.

Apicella, M., Campopiano, M.C., Mantuano, M., Mazoni, L., Coppelli, A., and Del Prato, S.
(2020). COVID-19 in people with diabetes: understanding the reasons for worse outcomes.
Lancet Diabetes Endocrinol.

Arda, H.E., Li, L., Tsai, J., Torre, E.A., Rosli, Y., Peiris, H., Spitale, R.C., Dai, C., Gu, X., Qu, K.,
et al. (2016). Age-dependent pancreatic gene regulation reveals mechanisms governing human
beta cell function. Cell Metab *23*, 909-920.

Ashok, A., Faghih, M., and Singh, V.K. (2020). Mild pancreatic enzyme elevations in COVID-19
pneumonia: synonymous with injury or noise? Gastroenterology.

Balamurugan, A.N., Gu, Y., Miyamoto, M., Wang, W., Inoue, K., and Tabata, Y. (2003).
Isolation, culture, and functional characteristics of diabetic islets [corrected]. Pancreas *26*, 102-
103.

Baron, M., Veres, A., Wolock, S.L., Faust, A.L., Gaujoux, R., Vetere, A., Ryu, J.H., Wagner,
B.K., Shen-Orr, S.S., Klein, A.M., et al. (2016). A single-cell transcriptomic map of the human
and mouse pancreas reveals inter- and intra-cell population structure. Cell Syst *3*, 346-360
e344.

Barron, E., Bakhai, C., Kar, P., Weaver, A., Bradley, D., Ismail, H., Knighton, P., Holman, N.,
Khunti, K., Sattar, N., et al. (2020). Associations of type 1 and type 2 diabetes with COVID-19-
related mortality in England: a whole-population study. Lancet Diabetes Endocrinol *8*, 813-822.

Blodgett, D.M., Nowosielska, A., Afik, S., Pechhold, S., Cura, A.J., Kennedy, N.J., Kim, S.,
Kucukural, A., Davis, R.J., Kent, S.C., et al. (2015). Novel observations from next-generation
RNA sequencing of highly purified human adult and fetal islet cell subsets. Diabetes *64*, 3172-
3181.

Bonney, G.K., Gao, Y., Chew, C.A., and Windsor, J.A. (2020). SARS-COV-2 associated acute
pancreatitis: Cause, consequence or epiphenomenon? Pancreatology.

Bornstein, S.R., Rubino, F., Khunti, K., Mingrone, G., Hopkins, D., Birkenfeld, A.L., Boehm, B.,
Amiel, S., Holt, R.I., Skyler, J.S., et al. (2020). Practical recommendations for the management
of diabetes in patients with COVID-19. Lancet Diabetes Endocrinol *8*, 546-550.

Breidenbach, J.D., Dube, P., Ghosh, S., Abdullah, B.N., Modyanov, N.N., Malhotra, D., Dworkin,
L.D., Haller, S.T., and Kennedy, D.J. (2020). Impact of Comorbidities on SARS-CoV-2 Viral
Entry-Related Genes. J Pers Med *10*.

Brissova, M., Aamodt, K., Brahmachary, P., Prasad, N., Hong, J.Y., Dai, C., Mellati, M.,
Shostak, A., Poffenberger, G., Aramandla, R., et al. (2014). Islet microenvironment, modulated
by vascular endothelial growth factor-A signaling, promotes beta cell regeneration. Cell Metab
19, 498-511.

Brissova, M., Fowler, M.J., Nicholson, W.E., Chu, A., Hirshberg, B., Harlan, D.M., and Powers,
A.C. (2005). Assessment of human pancreatic islet architecture and composition by laser
scanning confocal microscopy. J Histochem Cytochem *53*, 1087-1097.

Brissova, M., Haliyur, R., Saunders, D., Shrestha, S., Dai, C., Blodgett, D.M., Bottino, R.,
Campbell-Thompson, M., Aramandla, R., Poffenberger, G., et al. (2018). alpha Cell Function
and Gene Expression Are Compromised in Type 1 Diabetes. Cell Rep *22*, 2667-2676.

Bruno, G., Fabrizio, C., Santoro, C.R., and Buccoliero, G.B. (2020). Pancreatic injury in the
course of coronavirus disease 2019: A not-so-rare occurrence. J Med Virol.

Camunas-Soler, J., Dai, X.Q., Hang, Y., Bautista, A., Lyon, J., Suzuki, K., Kim, S.K., Quake,
S.R., and MacDonald, P.E. (2020). Patch-seq links single-cell transcriptomes to human islet
dysfunction in diabetes. Cell Metab *31*, 1017-1031 e1014.

Cariou, B., Hadjadj, S., Wargny, M., Pichelin, M., Al-Salameh, A., Allix, I., Amadou, C., Arnault,
G., Baudoux, F., Bauduceau, B., et al. (2020). Phenotypic characteristics and prognosis of
inpatients with COVID-19 and diabetes: the CORONADO study. Diabetologia *63*, 1500-1515.

Chee, Y.J., Ng, S.J.H., and Yeoh, E. (2020). Diabetic ketoacidosis precipitated by Covid-19 in a
patient with newly diagnosed diabetes mellitus. Diabetes Res Clin Pract *164*, 108166.

Chen, L., Li, X., Chen, M., Feng, Y., and Xiong, C. (2020). The ACE2 expression in human
heart indicates new potential mechanism of heart injury among patients infected with SARS-
CoV-2. Cardiovasc Res *116*, 1097-1100.

Clausen, T.M., Sandoval, D.R., Spliid, C.B., Pihl, J., Perrett, H.R., Painter, C.D., Narayanan, A.,
Majowicz, S.A., Kwong, E.M., McVicar, R.N., et al. (2020). SARS-CoV-2 Infection Depends on
Cellular Heparan Sulfate and ACE2. Cell.

Dai, C., Hang, Y., Shostak, A., Poffenberger, G., Hart, N., Prasad, N., Phillips, N., Levy, S.E.,
Greiner, D.L., Shultz, L.D., et al. (2017). Age-dependent human beta cell proliferation induced
by glucagon-like peptide 1 and calcineurin signaling. J Clin Invest *127*, 3835-3844.

Drucker, D.J. (2020). Coronavirus infections and type 2 diabetes-shared pathways with
therapeutic implications. Endocr Rev *41*.

Filippi, C.M., and von Herrath, M.G. (2008). Viral trigger for type 1 diabetes: pros and cons.
Diabetes *57*, 2863-2871.

Goldman, N., Fink, D., Cai, J., Lee, Y.N., and Davies, Z. (2020). High prevalence of COVID-19-
associated diabetic ketoacidosis in UK secondary care. Diabetes Res Clin Pract *166*, 108291.

Gupta, A., Madhavan, M.V., Sehgal, K., Nair, N., Mahajan, S., Sehrawat, T.S., Bikdeli, B.,
Ahluwalia, N., Ausiello, J.C., Wan, E.Y., et al. (2020). Extrapulmonary manifestations of COVID-
19. Nat Med *26*, 1017-1032.

Hamming, I., Timens, W., Bulthuis, M.L., Lely, A.T., Navis, G., and van Goor, H. (2004). Tissue
distribution of ACE2 protein, the functional receptor for SARS coronavirus. A first step in
understanding SARS pathogenesis. J Pathol *203*, 631-637.

He, L., Mae, M., Sun, Y., Muhl, L, Nahar, K., et al (2020). Pericyte-specific vascular expression
of SARS-CoV-2 receptor ACE2 – implications for microvascular inflammation and
hypercoagulopathy in COVID-19 patients. Biorxiv.

Heaney, A.I., Griffin, G.D., and Simon, E.L. (2020). Newly diagnosed diabetes and diabetic
ketoacidosis precipitated by COVID-19 infection. Am J Emerg Med.

Hikmet, F., Mear, L., Edvinsson, A., Micke, P., Uhlen, M., and Lindskog, C. (2020). The protein
expression profile of ACE2 in human tissues. Mol Syst Biol *16*, e9610.

Hoffmann, M., Kleine-Weber, H., Schroeder, S., Kruger, N., Herrler, T., Erichsen, S.,
Schiergens, T.S., Herrler, G., Wu, N.H., Nitsche, A., et al. (2020). SARS-CoV-2 cell entry
depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. Cell
181, 271-280 e278.

Hollstein, T., Schulte, D.M., Schulz, J., Gluck, A., Ziegler, A.G., Bonifacio, E., Wendorff, M.,
Franke, A., Schreiber, S., Bornstein, S.R., et al. (2020). Autoantibody-negative insulin-
dependent diabetes mellitus after SARS-CoV-2 infection: a case report. Nat Metab.

Holman, N., Knighton, P., Kar, P., O'Keefe, J., Curley, M., Weaver, A., Barron, E., Bakhai, C.,
Khunti, K., Wareham, N.J., et al. (2020). Risk factors for COVID-19-related mortality in people
with type 1 and type 2 diabetes in England: a population-based cohort study. Lancet Diabetes
Endocrinol *8*, 823-833.

Kaestner, K.H., Powers, A.C., Naji, A., Consortium, H., and Atkinson, M.A. (2019). NIH Initiative
to Improve Understanding of the Pancreas, Islet, and Autoimmunity in Type 1 Diabetes: The
Human Pancreas Analysis Program (HPAP). Diabetes *68*, 1394-1402.

Karimzadeh, S., Manzuri, A., Ebrahimi, M., and Huy, N.T. (2020). COVID-19 presenting as
acute pancreatitis: Lessons from a patient in Iran. Pancreatology.

Kim, N.Y., Ha, E., Moon, J.S., Lee, Y.H., and Choi, E.Y. (2020). Acute hyperglycemic crises with
coronavirus disease-19: case reports. Diabetes Metab J *44*, 349-353.

Klein, A.M., Mazutis, L., Akartuna, I., Tallapragada, N., Veres, A., Li, V., Peshkin, L., Weitz,
D.A., and Kirschner, M.W. (2015). Droplet barcoding for single-cell transcriptomics applied to
embryonic stem cells. Cell *161*, 1187-1201.

Koch, M., Holmes, D., Bennet, N. (2020). COVID-19 and diabetes: a co-conspiracy? Lancet
Diabetes Endocrinol *8*, 801.

Kusmartseva, I., Wu, W., Syed, F., Van Der Heide, V., Jorgensen, M., Joseph, P., Tang, X.,
Candelario-Jalil, E., Yang, C., Nick, H., Harbert, J., Posgai, A., Lloyd, R., Cechin, S., Pugliese,
A., Campbell-Thompson, M., Vander Heide, R., Evans-Molina, C., Homann, D., Atkinson, M.
(2020). ACE2 and SARS-CoV-2 Expression in the Normal and COVID-19 Pancreas. Biorxiv.

Lan, J., Ge, J., Yu, J., Shan, S., Zhou, H., Fan, S., Zhang, Q., Shi, X., Wang, Q., Zhang, L., et
al. (2020). Structure of the SARS-CoV-2 spike receptor-binding domain bound to the ACE2
receptor. Nature *581*, 215-220.

Lee, I.T., Nakayama, T., Wu, C.T., Goltsev, Y., Jiang, S., Gall, P.A., Liao, C.K., Shih, L.C.,
Schurch, C.M., McIlwain, D.R., et al. (2020). Robust ACE2 protein expression localizes to the
motile cilia of the respiratory tract epithelia and is not increased by ACE inhibitors or angiotensin
receptor blockers. medRxiv.

Li, J., Wang, X., Chen, J., Zuo, X., Zhang, H., and Deng, A. (2020a). COVID-19 infection may
cause ketosis and ketoacidosis. Diabetes Obes Metab.

Li, Y., Zhang, Z., Yang, L., Lian, X., Xie, Y., Li, S., Xin, S., Cao, P., and Lu, J. (2020b). The
MERS-CoV receptor DPP4 as a candidate binding target of the SARS-CoV-2 spike. iScience
23, 101160.

Meireles, P.A., Bessa, F., Gaspar, P., Parreira, I., Silva, V.D., Mota, C., and Alvoeiro, L. (2020).
Acalculous acute pancreatitis in a COVID-19 patient. Eur J Case Rep Intern Med *7*, 001710.

Nair, G.G., Liu, J.S., Russ, H.A., Tran, S., Saxton, M.S., Chen, R., Juang, C., Li, M.L., Nguyen,
V.Q., Giacometti, S., et al. (2019). Recapitulating endocrine cell clustering in culture promotes
maturation of human stem-cell-derived beta cells. Nat Cell Biol *21*, 263-274.

Picelli, S., Faridani, O.R., Bjorklund, A.K., Winberg, G., Sagasser, S., and Sandberg, R. (2014).
Full-length RNA-seq from single cells using Smart-seq2. Nat Protoc *9*, 171-181.

preprint: Fignani, D., Licata, G., Brusco, N., Nigi, L., Grieco, G., Marselli, L., Overbergh, L.,
Gysemans, C., Colli, M., Marchetti, P., Mathieu, C., Eizirik, D., Sebastiani, G., Dotta, F. (2020).
SARS-CoV-2 receptor angiotensin I-converting enzyme type 2 is expressed in human
pancreatic islet β-cells and is upregulated by inflammatory stress. BioRxiv.

Rabice, S.R., Altshuler, P.C., Bovet, C., Sullivan, C., and Gagnon, A.J. (2020). COVID-19
infection presenting as pancreatitis in a pregnant woman: A case report. Case Rep Womens
Health *27*, e00228.

Rafique, S., and Ahmed, F.W. (2020). A Case of Combined Diabetic Ketoacidosis and
Hyperosmolar Hyperglycemic State in a Patient With COVID-19. Cureus *12*, e8965.

Riddle, M.C., Buse, J.B., Franks, P.W., Knowler, W.C., Ratner, R.E., Selvin, E., Wexler, D.J.,
and Kahn, S.E. (2020). COVID-19 in people with diabetes: urgently needed lessons from early
reports. Diabetes Care *43*, 1378-1381.

Schreiber, B., Patel, A., and Verma, A. (2020). Shedding Light on COVID-19: ADAM17 the
Missing Link? Am J Ther.

Segerstolpe, A., Palasantza, A., Eliasson, P., Andersson, E.M., Andreasson, A.C., Sun, X.,
Picelli, S., Sabirsh, A., Clausen, M., Bjursell, M.K., et al. (2016). Single-cell transcriptome
profiling of human pancreatic islets in health and type 2 diabetes. Cell Metab *24*, 593-607.

Seyedpour, S., Khodaei, B., Loghman, A.H., Seyedpour, N., Kisomi, M.F., Balibegloo, M.,
Nezamabadi, S.S., Gholami, B., Saghazadeh, A., and Rezaei, N. (2020). Targeted therapy
strategies against SARS-CoV-2 cell entry mechanisms: A systematic review of in vitro and in
vivo studies. J Cell Physiol.

Shang, J., Wan, Y., Luo, C., Ye, G., Geng, Q., Auerbach, A., and Li, F. (2020). Cell entry
mechanisms of SARS-CoV-2. Proc Natl Acad Sci U S A *117*, 11727-11734.

Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., 3rd, Hao, Y.,
Stoeckius, M., Smibert, P., and Satija, R. (2019). Comprehensive Integration of Single-Cell
Data. Cell *177*, 1888-1902 e1821.

Taneera, J., El-Huneidi, W., Hamad, M., Mohammed, A.K., Elaraby, E., and Hachim, M.Y.
(2020). Expression Profile of SARS-CoV-2 Host Receptors in Human Pancreatic Islets
Revealed Upregulation of ACE2 in Diabetic Donors. Biology (Basel) *9*.

Teuwen, L.A., Geldhof, V., Pasut, A., and Carmeliet, P. (2020). COVID-19: the vasculature
unleashed. Nat Rev Immunol *20*, 389-391.

Uhlen, M., Fagerberg, L., Hallstrom, B.M., Lindskog, C., Oksvold, P., Mardinoglu, A., Sivertsson,
A., Kampf, C., Sjostedt, E., Asplund, A., et al. (2015). Proteomics. Tissue-based map of the
human proteome. Science *347*, 1260419.

Unsworth, R., Wallace, S., Oliver, N.S., Yeung, S., Kshirsagar, A., Naidu, H., Kwong, R.M.W.,
Kumar, P., and Logan, K.M. (2020). New-Onset Type 1 Diabetes in Children During COVID-19:
Multicenter Regional Findings in the U.K. Diabetes Care.

van den Brink, S.C., Sage, F., Vertesy, A., Spanjaard, B., Peterson-Maduro, J., Baron, C.S.,
Robin, C., and van Oudenaarden, A. (2017). Single-cell sequencing reveals dissociation-
induced gene expression in tissue subpopulations. Nat Methods *14*, 935-936.

Vankadari, N., and Wilce, J.A. (2020). Emerging WuHan (COVID-19) coronavirus: glycan shield
and structure prediction of spike glycoprotein and its interaction with human CD26. Emerg
Microbes Infect *9*, 601-604.

Wang, F., Wang, H., Fan, J., Zhang, Y., Wang, H., and Zhao, Q. (2020). Pancreatic injury
patterns in patients with coronavirus disease 19 pneumonia. Gastroenterology *159*, 367-370.

Wang, S., Ma, P., Zhang, S., Song, S., Wang, Z., Ma, Y., Xu, J., Wu, F., Duan, L., Yin, Z., Luo,
H., Xiong, N., Xu, M., Zeng, T., Jin, Y. (2020). Fasting blood glucose at admission is an
independent predictor for 28-day mortality in patients with COVID-19 without previous diagnosis
of diabetes: a multi-centre retrospective study. Diabetologia.

Wiersinga, W.J., Rhodes, A., Cheng, A.C., Peacock, S.J., and Prescott, H.C. (2020).
Pathophysiology, Transmission, Diagnosis, and Treatment of Coronavirus Disease 2019
(COVID-19): A Review. JAMA.

Wright, J.J., Saunders, D.C., Dai, C., Poffenberger, G., Cairns, B., Serreze, D.V., Harlan, D.M.,
Bottino, R., Brissova, M., and Powers, A.C. (2020). Decreased pancreatic acinar cell number in
type 1 diabetes. Diabetologia *63*, 1418-1423.

Yang, J.K., Lin, S.S., Ji, X.J., and Guo, L.M. (2010). Binding of SARS coronavirus to its receptor
damages islets and causes acute diabetes. Acta Diabetol *47*, 193-199.

Yang, L., Han, Y., Nilsson-Payant, B.E., Gupta, V., Wang, P., Duan, X., Tang, X., Zhu, J., Zhao,
Z., Jaffre, F., et al. (2020). A human pluripotent stem cell-based platform to study SARS-CoV-2
tropism and model virus infection in human cells and organoids. Cell Stem Cell *27*, 125-136
e127.

Zang, R., Gomez Castro, M.F., McCune, B.T., Zeng, Q., Rothlauf, P.W., Sonnek, N.M., Liu, Z.,
Brulois, K.F., Wang, X., Greenberg, H.B., et al. (2020). TMPRSS2 and TMPRSS4 promote
SARS-CoV-2 infection of human small intestinal enterocytes. Sci Immunol *5*.

Zhou, F., Yu, T., Du, R., Fan, G., Liu, Y., Liu, Z., Xiang, J., Wang, Y., Song, B., Gu, X., et al.
(2020). Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan,
China: a retrospective cohort study. Lancet *395*, 1054-1062.
834

Figure 1

HPAP

A

ACE2

B

TMPRSS2

C

ADAM17

Donor BMI

CoV-2 Entry Markers INS GCG DAPI

DPP4	INS GCG	MERGE
A 55yM	**B**	**C**
D 60y_1yM	**E**	**F**
G 20y_7yM	**H**	**I**

Normal Adult (A, B, C)
Type 2 Diabetes (D, E, F)
Type 1 Diabetes (G, H, I)

J

Camunas-Soler et al.

Genes: ACE2, TMPRSS2, CTSL, ADAM17, FURIN, TMPRSS4, DPP4, GCG, IRX2, PCSK2, INS, IAPP, PDX1, SST, HHEX, LEPR, PPY, GHRL, PRSS1, REG1A, CPA2, KRT19, PROM1, CFTR, PECAM1, PLVAP, VWF, PDGFRB, TIMP1, COL1A1, HLA-DRA, CD68, CD74

Cell types: Alpha, Beta, Delta, Gamma, Acinar, Other

Average Expression: 2, 1, 0, -1
Percent Expressed: 0, 25, 50, 75, 100

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Mouse anti-Cytokeratin 19 (1:1000)	Abcam	Cat# ab194399
Rabbit anti-TMPRSS2 (1:400)	Sigma	Cat# HPA035787, RRID:AB_2674782
Mouse anti-Glucagon (1:500-1:1000)	Abcam	Cat# ab10988, RRID:AB_297642
Guinea pig anti-Glucagon (1:500)	LSBio	Cat# LS-C202759-50
Rat anti-C-peptide (1:1000)	DSHB	Cat# GN-ID4, RRID:AB_2255626
Guinea pig anti-Insulin (1:500-1:1000)	Dako	Cat# A0564, RRID:AB_2617169
Rabbit anti-ACE2 (1:500)	Atlas Antibodies	Cat# HPA000288, RRID:AB_1078160
Rabbit anti-ACE2 (1:1000)	Abcam	Cat# ab15348, RRID:AB_301861
Goat anti-ACE2 (1:200, Figure S1,3A-E; 1:400, Figure 4)	R&D	Cat# AF933, RRID:AB_355722
Mouse anti-ACE2 (1:33)	R&D	Cat# MAB933, RRID:AB_2223153
Mouse anti-CD31 (1:50)	BD Biosciences	Cat# 550389, RRID:AB_2252087
Goat anti-collagen IV alpha 1 (1:500)	Novus	Cat# NBP1-26549, RRID:AB_1853202
Goat anti-DPP4 (1:200)	R&D	Cat# AF1180, RRID:AB_354651
Mouse anti-NG2 [647] (1:50)	R&D	Cat# FAB2585R;
Goat anti-PDGFRβ (1:100)	R&D	Cat# AF385, RRID:AB_355339
Donkey anti-mouse-Cy2 (1:500)	Jackson ImmunoResearch	Cat# 715-225-150, RRID:AB_2340826
Donkey anti-mouse-Cy3 (1:500)	Jackson ImmunoResearch	Cat# 715-165-150, RRID:AB_2340813
Donkey anti-mouse-Cy5 (1:250 – 1:500)	Jackson ImmunoResearch	Cat# 715-175-151, RRID:AB_2340820
Donkey anti-rabbit-Cy3 (1:500)	Jackson ImmunoResearch	Cat# 711-165-152, RRID:AB_2307443
Donkey anti-rat-Cy2 (1:500)	Jackson ImmunoResearch	Cat# 712-225-153, RRID:AB_2340674
Donkey anti-guinea pig-Cy2 (1:500)	Jackson ImmunoResearch	Cat# 706-225-148, RRID:AB_2340467
Donkey anti-goat-Cy2 (1:500)	Jackson ImmunoResearch	Cat# 705-225-147, RRID:AB_2307341
Donkey anti-goat-Cy3 (1:500)	Jackson ImmunoResearch	Cat# 705-165-147, RRID:AB_2307351
Donkey anti-guinea pig-Cy5 (1:250-1:500)	Jackson ImmunoResearch	Cat# 706-175-148, RRID:AB_2340462
Normal Donkey Serum	Jackson ImmunoResearch	Cat# 017-000-121, RRID: AB_2337258
Bacterial and Virus Strains		
n/a		
Biological Samples		
Human pancreatic islets	Integrated Islet Distribution Program (IIDP)	http://iidp.coh.org, RRID: SCR_014387
Human pancreatic tissue	International Institute for the Advancement of Medicine (IIAM)	http://www.iiam.org, RRID: SCR_016172

	National Disease Research Interchange (NDRI)	http://ndriresource.org, RRID: SCR_000550
Human pancreatic tissue		
Chemicals, Peptides, and Recombinant Proteins		
Propidium Iodide	BD Biosciences	Cat# 556463
Antigen Unmasking Solution, Citrate-Based	Vector Laboratories	Cat# H-3300-250
1X PBS	Corning	Cat# 46-013-CM
Triton X-100	USB	Cat# 22686
BSA	Sigma	Cat# A6003-25G
Human ACE2 Peptide	Abcam	Cat# ab15352
Critical Commercial Assays		
RNAqueous-Micro Kit	Fisher Scientific (Invitrogen)	Cat# AM1931
High Capacity cDNA Reverse Transcription Kit	Fisher Scientific (Invitrogen)	Cat# 4368814
Deposited Data		
RNA-seq data for FACS purified human alpha and beta cells (Blodgett et al., 2015)	NCBI Gene Expression Omnibus	GEO: GSE67543
RNA-seq data for FACS purified human alpha and beta cells (Arda et al., 2016)	NCBI Gene Expression Omnibus	GEO: GSE57973
Single-cell RNA-seq data for human islets (Baron et al., 2016)	NCBI Gene Expression Omnibus	GEO: GSE84133
Single-cell RNA-seq data for human islets (Camunas-Soler et al., 2020)	NCBI Gene Expression Omnibus	GEO: GSE124742
Single-cell RNA-seq data for human islets (Segerstolpe et al., 2016)	ArrayExpress	E-MTAB-5061
Single-cell RNA-seq data for human islets (Human Pancreas Analysis Program [HPAP])	HPAP Database	https://hpap.pmacs.upenn.edu RRID:SCR_014393
Experimental Models: Cell Lines		
n/a		
Experimental Models: Organisms/Strains		
n/a		
Recombinant DNA		
n/a		
Software and Algorithms		
FV31S-SW Viewer 2.4.1.198	Olympus	https://www.olympus-lifescience.com/en/support/downloads/
Seurat 3.1 R package	(Butler et al., 2018) (Stuart et al., 2019)	N/A
Prism for macOS v.8.4.3	Graphpad Software	http://www.graphpad.com, RRID: SCR_002798
JACoP: Just Another Co-localization Plugin	ImageJ	https://imagej.nih.gov/ij/plugins/track/jacop2.html