A genetic screen identifies processes that couple oocyte maturation to proteostasis in the immortal *C. elegans* germ lineage

Madhuja Samaddar*,[1], Jérôme Goudeau*,[1], Melissa Sanchez[2], David H. Hall[3], K. Adam Bohnert[1,#], Maria Ingaramo[1] and Cynthia Kenyon[1,†]

[1]Calico Life Sciences LLC, South San Francisco, CA 94080, USA.

[2]Department of Molecular and Cellular Biology, University of California, Berkeley, CA 94720, USA.

[3]Dominick P. Purpura Department of Neuroscience, Albert Einstein College of Medicine, New York, NY 01461, USA.

[#]Present address: Department of Biological Sciences, Louisiana State University, Baton Rouge, LA 70803

* These authors contributed equally

† Correspondence to: cynthia@calicolabs.com

Abstract

Somatic cells age and die, but the germ-cell lineage is immortal. In *C. elegans,* oocyte-maturation signals from sperm trigger the clearance of carbonylated proteins and protein aggregates. Here, we explore the cell biology of this proteostasis renewal in the context of a whole-genome RNAi screen for knockdowns that interfere with aggregate clearance. Oocyte-maturation signals are known to trigger protein-aggregate removal via lysosome acidification, and our findings suggest that lysosomes are acidified as a consequence of changes in ER morphology and function that permit assembly of the lysosomal V-ATPase. Once lysosomes are acidified, our genetic findings support the model that they remove aggregates by microautophagy. We also define two functions for mitochondria in this proteostasis renewal, both of which appear to be independent of mitochondrial ATP generation. Finally, many genes from the screen also regulate lysosome acidification and age-dependent protein aggregation in the soma, suggesting a fundamental mechanistic link between proteostasis renewal in the germline and the maintenance of the soma.

Introduction

The germ cells of an organism not only give rise to the rich diversity of somatic lineages, they also generate new germ cells that, by seeding subsequent generations, confer species immortality. This sequential rejuvenation of youthfulness is one of the most interesting mysteries in biology. How is it achieved? Not surprisingly, a process so crucial to the survival of the species involves multiple levels of quality control. At one level, an entire oocyte can be eliminated via apoptosis; for example, following DNA damage (Hunter, 2017; Luo et al., 2010; Qiao et al., 2018; Schumacher et al., 2001). Oocytes can also eliminate defective organelles or age-related molecular damage. For example, in mice, mitochondria containing defective DNA are eliminated during oogenesis (Fan et al., 2008), and in yeast, nuclear senescence factors and protein aggregates are extruded during meiosis (King et al., 2019; Ünal et al., 2011). Mechanisms likely exist to counteract protein aggregation in *Drosophila* as well: relative to the soma, the eggs of *Drosophila* exhibit enhanced proteasome activity and resistance to protein aggregation (Fredriksson et al., 2012). Autophagy, a process known to eliminate damaged organelles and macromolecules, may play a role as well, as it is required for the survival of the mouse germ lineage (Gawriluk et al., 2011).

The nematode *C. elegans* is a valuable organism for studies of germline quality control. The animal is transparent and genetically accessible, allowing one to analyze the effects of perturbations over time in living animals using fluorescent reporters. *C. elegans* is a hermaphrodite, producing both sperm and oocytes in each of two symmetrical gonadal arms. Germ cells originate in stem-cell niches located at the distal ends of each arm and pass through successive stages of meiosis as they move proximally towards the spermatheca, where fertilization occurs (Crittenden et al., 1994; Hubbard and Greenstein, 2005; Mccarter et al., 1999). (Figure 1A). In 2010, Goudeau and Aguilaniu discovered that carbonylated proteins, known to accumulate during aging, are present throughout the distal germline, but are eliminated in proximal oocytes (Goudeau and Aguilaniu, 2010). Mutant worms that lack sperm

and develop as females retain carbonylated proteins throughout the entire germline, indicating

that sperm play a role in their elimination. In addition to carbonylated proteins, protein

aggregates also accumulate in the proximal oocytes of females but not in those of

hermaphrodites (Bohnert and Kenyon, 2017). Signals from sperm trigger the elimination of

protein aggregates in a process that is coupled to the broader choreography of oocyte

maturation, which involves dramatic changes in organelle morphology and function (Bluemink et

al., 1983; Carroll et al., 1996; Charbonneau and Grey, 1984; FitzHarris et al., 2007; Kobayashi

et al., 1991; Maller et al., 1977; Mehlmann et al., 1995; Wasserman et al., 1982; Yamashita,

2018). In *C. elegans,* oocyte maturation is triggered by actin-like major sperm proteins (MSPs)

released from sperm (Kosinki et al., 2005; Miller et al., 2003, 2001), and involves changes in

chromosome and nuclear dynamics linked to progression through meiosis, up-regulation of

translation, and changes in ER, lysosomal and mitochondrial function (Bohnert and Kenyon,

2017; Huelgas-Morales and Greenstein, 2018; Langerak et al., 2019). The shift in lysosomal

physiology that occurs during oocyte maturation is crucial for aggregate removal: the sperm-

derived signals trigger the acidification of lysosomes, which in turn engulf the aggregates by a

process that resembles microautophagy morphologically (Bohnert and Kenyon, 2017).

Lysosomal acidification also triggers a conversion in mitochondrial dynamics from what appears

to be a poised state, characterized by a high membrane potential, to an active state, in which

the potential is reduced. Conditions that prevent this mitochondrial membrane-potential shift,

such as inactivation of mitochondrial ATP synthase, prevent aggregate removal. However, how

lysosomes become acidified in response to sperm signals, how mitochondria contribute to

aggregate removal, and whether other organelles also play a role in aggregate clearance are

important outstanding questions.

To better understand mechanisms of proteostasis enhancement in the *C. elegans* germ

lineage, we combined cell biological approaches with a confocal-microscopy based genome-

wide RNAi screen for gene knockdowns that cause protein aggregates to accumulate even in

the presence of sperm. Our findings deepen our understanding of many aspects of protein

aggregation and clearance within the germline. Specifically, we find that (i) protein aggregates

accumulate in young, newly formed oocytes in the absence of sperm—their formation is an

integral part of oocyte development, rather than a consequence of prolonged oocyte

quiescence; (ii) inhibiting processes required for changes in ER morphogenesis and function

during oocyte maturation, including protein synthesis and actin dynamics, prevents lysosome

acidification by preventing the assembly of the V-ATPase lysosomal proton pump; (iii)

mitochondria elaborate an energy-independent checkpoint that gates protein-aggregate

removal; (iv) ESCRT vesicle-transport proteins are required for aggregate removal, consistent

with protein aggregates being removed via microautophagy; and, unexpectedly (v) as with

proteasome inhibition, inhibiting TRiC-complex chaperonins or HSP70 chaperones promotes

protein aggregation indirectly, by acting upstream in the pathway to block lysosome

acidification. Finally, we show that inhibiting lysosomal V-ATPase, vesicle-transport or

proteasome function also accelerates age-dependent protein aggregation in the soma, thus

bridging mechanisms that enhance proteostasis in the immortal germ lineage to those that

maintain proteostasis in the aging soma.

Results

A genome-wide RNAi screen identifies 84 genes that influence oocyte protein aggregation.

To undertake an unbiased genome-wide RNAi screen for components of the germline proteostasis network, we utilized the Ahringer RNAi feeding library (Kamath et al., 2003), which targets ~87% of annotated *C. elegans* genes, to identify gene inhibitions that cause proteins to aggregate even in the presence of sperm; that is, in hermaphrodites. A germline-specific GFP::RHO-1 translational fusion was selected as the reporter for the screen, as it forms dense patchy aggregates in female oocytes (Bohnert and Kenyon, 2017) and in unfertilized oocytes of aging hermaphrodites, which deplete their stores of sperm (David et al., 2010). Oocytes are formed in young adults, so RNAi was initiated at the beginning of the last larval stage (L4) to reduce the chance of missing genes with essential roles during development (Figure 1B). The animals were examined manually by confocal microscopy on day 2 of adulthood for the presence of GFP::RHO-1 aggregates in proximal oocytes (Figure 1A, 1B and S1A). The primary screen identified 367 genes whose knockdown led to GFP::RHO-1 aggregation in at least 50% of the animals. To eliminate false positives, the candidates were subjected to three independent rounds of validation using a similar workflow, with at least two independent rounds from two experimentalists (Figure 1C). We note that false negatives could result from failure of an RNAi bacterial strain to grow adequately, from insufficient or excessive gene inhibition, or because of functional redundancy.

The resulting set of 88 genes represented diverse cellular processes (Table S1), with highly enriched gene ontology (GO) categories including translation, RNA processing and cytoskeletal organization (Figure S1C). In addition, ATP-hydrolysis-coupled proton transport and lysosome acidification were both highly enriched, in agreement with our previous findings (Bohnert and Kenyon, 2017). We sorted the candidates according to relative representation in major biological processes (Figure 1D), after discarding candidates with known roles in male gonad and germ cell development (*fkh-6, spe-15*), in regulation of our proteostasis marker

RHO-1 (*cyk-4*), or with potential off-target effects (*adr-1* and *dhs-5*) (Table S1). Protein

synthesis and degradation, RNA processing, protein folding, and lysosome acidification were

strongly represented biological processes, as were genes involved in cytoskeletal organization,

including four of the five *C. elegans* actin genes. Among organelle systems, multiple candidates

were associated with ER homeostasis (*hsp-4, spcs-1 and srpa-68*) and vesicle-mediated

transport (*copb-2, sec-24.1, sar-1, rab-11.1* and *E03H4.8*). We also identified genes that could

not be classified into any of the highly-enriched biological-process categories and were

designated 'orphans' (Table S1 and Fig. 1D).

We subjected representative candidates from each group, as well as all the orphans, to

an orthogonal screening approach, this time utilizing a different aggregation reporter, a

germline-specific NMY-2::GFP fusion protein (Bohnert and Kenyon, 2017) that forms distinctive

punctate aggregates in the absence of sperm (Figure 1C and Table S1). Out of the 55

candidates tested, 53 RNAi knockdowns also caused NMY-2::GFP protein-aggregate

accumulation (Figure S1B). Thus, these genes are likely to have a general effect on protein-

aggregate clearance rather than a specific effect on GFP::RHO-1.

Figure 1

Figure 1. **Identification of genes that prevent protein aggregation in maturing *C. elegans* oocytes.** (A) Diagram of the *C. elegans* germline (B) Genome-wide RNAi-screen workflow. High resolution images from the validation rounds are represented here. Scale bars, 10 µm. (C) Schematic representing the procedure used for candidate validation and orthogonal verification using two independent germline-specific reporters. 45 candidates representing the major GO categories and all 10 orphans were subjected to orthogonal verification. (D) Common biological process GO categories enriched in the screen. Genes that were not members of these enriched categories were classified as orphans. (E) Very early oocytes of females and hermaphrodites (CF4557 and CF4552 strains, respectively). GFP::RHO-1 aggregates often appear in newly-formed female oocytes. Scale bars, 10 µm.

Aggregates form in young proximal oocytes.

We classified the RNAi phenotypes as strong, intermediate or subtle based on the

degree of GFP::RHO-1 aggregation (Table S1). Typically, the phenotypically strong

knockdowns led to sterility, whereas the subtle knockdowns generally did not, though some

reduced the brood size. In principle, one could imagine that aggregates accumulate in oocytes

from sterile hermaphrodites because oocytes that would normally undergo ovulation and pass

into the uterus do not, and instead simply "age in place". Likewise, aggregates could

accumulate in normal female oocytes because in the absence of sperm, they proceed much

more slowly through the spermatheca and into the uterus (Mccarter et al., 1999). As a

consequence, proximal oocytes of females are technically older than those of age-matched

hermaphrodites (Kim et al., 2013). This oocyte-age discrepancy raises the question of whether

aggregates accumulate only during long-term oocyte quiescence, or whether they are present in

newly formed female oocytes. To address this question, we collected age-matched late L4

('Christmas-tree' stage) females and hermaphrodites and followed the hermaphrodites until their

first oocytes were cellularized. We found that age-matched first-proximal oocytes of young

females often contained protein aggregates (Figure 1E and S4). Therefore, protein-aggregate

accumulation does not require prolonged oocyte quiescence; instead, it appears to be part of

the normal developmental sequence of oocytes that are not exposed to sperm.

The endoplasmic reticulum influences protein aggregation

Many genes we identified were associated with the ER, so we explored ER biology in more detail. The ER undergoes a major morphological rearrangement in response to oocyte maturation signals (Langerak et al., 2019). We visualized the ER in *C. elegans* oocytes using a germline-specific mCherry::SP12 reporter (Joseph-Strauss et al., 2012). SP12 is a signal-peptidase subunit (Poteryaev et al., 2005) and *spcs-1*, which encodes SP12, was identified in our screen. In hermaphrodite oocytes, the ER assumed a fine network appearance, whereas in female oocytes, the ER formed bright, distinctive patches (Figure 2A, 2C and S2A). We also examined ER morphology in the first, newly-formed female oocytes, using the L4 staging procedure described above. Pristine female oocytes also exhibited ER patches, indicating that the altered ER architecture is an intrinsic property of female oocytes and not a consequence of prolonged quiescence (Figure 2B and S2B). The fluorescent signal from the female ER was further enhanced as oocytes aged, possibly due to continued ER biogenesis coupled with oocyte compression caused by their packing within the gonad (Figure S2A). Using two fluorescent reporters, we visualized both the ER and GFP::RHO-1 simultaneously (Figure 2B, S2C, S3 and S4), and found that in female oocytes, the ER and GFP::RHO-1 aggregates segregated spatially into mutually-exclusive regions (Figure 2B, S2C and S4).

We also examined oocytes from young, age-matched hermaphrodites and females by transmission electron microscopy (TEM). The ER in pristine oocytes from both sexes appeared to be associated with ribosomes, indicative of active (or poised) protein synthesis, yet the female ER was distinctively clustered into parallel stacks (Figure 2D and S5).

To test whether the stacked ER architecture of female oocytes could be reversed by sperm, we visualized fluorescently-tagged ER and NMY-2::GFP aggregates simultaneously over time in female oocytes after mating. The presence of sperm triggered distinctive remodeling of the ER architecture and the clearance of aggregates, although with different kinetics (Figure 2E and S6A). The earliest changes in the ER were detected approximately 10

minutes after mating, and the morphology was significantly altered by 30 minutes (Figure 2E). In

contrast, the clearance of NMY-2 aggregates started a few minutes later, with aggregate

clearance in the most proximal oocytes typically occurring between 20 to 30 minutes after

mating (Figure S6A, S6B and S7A).

Next, we investigated whether the genes identified in our RNAi screen might contribute

to proteostasis by influencing a process associated with ER architecture. To this end, we

subjected mCherry::SP12-germline-labeled hermaphrodites to 35 RNAi treatments representing

the various biological process categories as well as the individual orphans. This subset of RNAi

clones, which we termed the "assay pool", was used for all subsequent large-scale phenotypic

analyses (Table 1). We visualized oocyte ER morphology in young adults subjected to the gene

knockdowns (Figure S7B and Table 1). Genes involved in cytoskeletal organization, protein

synthesis and protein folding were required for maintaining normal ER morphology and led to an

altered ER architecture when knocked down. Genes involved in RNA processing, protein

degradation and trafficking comprised a mixed class, with some knockdowns appearing to affect

ER morphology. [We note that our analysis does not distinguish between genes with direct

effects on ER morphogenesis, like the actin cytoskeleton (Poteryaev et al., 2005), and genes

with indirect effects, like the proteasome, which is required for the degradation of GLD-1

(Bohnert and Kenyon, 2017; Goudeau et al., 2020b; Spike et al., 2018), a translational

repressor involved in oocyte cell fate determination (Francis et al., 1995; Jones et al., 1996)].

Notably, disrupting the mitochondrial ATP synthase, the V-ATPase complex or ER function itself

did not visibly alter the ER network, suggesting that these gene functions were not required for

the sperm-induced ER morphology change, or more generally for ER morphogenesis.

Figure 2

A **mCherry::SP12** | Hermaphrodite / Female

B **mCherry::SP12** | **GFP::RHO-1** | **Merge** | Hermaphrodite / Female

C Animals with ER patches (%) — ******** — Herm / Female

D Hermaphrodite | Female — n

E **mCherry::SP12** — Time series, post mating

female; pre-mating | 10 min | 20 min
u* ; -7 -6 -5 -4 -3 -2 -1

30 min | 50 min | 70 min
fertilized | fertilized | fertilized / sp*
-7 -6 -5 -4 -3 -2 "-1" | -7 -6 -5 -4 -3 -2 "-1" | -6 -5 -4 -3 "-2" "-1"

Figure 2. ER morphology is regulated by sperm and correlates with protein aggregation.

(A) ER morphology in the germline of hermaphrodite (OCF15) and female (CF4542) animals, visualized using a mCherry::SP12 reporter. Orange arrows, ER patches. Scale bar, 10 μm. (B) GFP::RHO-1 localization in newly-formed oocytes from hermaphrodites (CF4552) and females (CF4557) also expressing mCherry::SP12. White arrows, GFP::RHO-1 aggregates. Scale bars, 10 μm. (C) Percentage of animals with ER patches in oocytes. n=12; mean +/- s.d., **** $P \leq 0.0001$. (D) Transmission electron micrographs from hermaphrodite (N2E) and female (CF4101) oocytes. Orange arrows, ER. Left panel, magnification 1,400X, scale bar, 1 μm; inset: magnification 13,000X, scale bar, 100 nm. Right panel, magnification 1,200X, scale bar, 2 μm; inset: magnification 18,500X, scale bar, 100 nm. (E) Reversal of ER patches (orange arrows) in female oocytes upon mating with CB1490 males. Female oocytes were imaged before and after mating at the indicated time points. Each oocyte is numbered according to its initial position relative to the spermatheca, and original numbers are maintained after fertilization. The figure shows a single representative animal, from 7 independent experiments. u*, unfertilized oocyte/debris in uterus in a different focal plane; sp*, an oocyte passing through spermatheca during fertilization. Scale bars, 10 μm.

Biological process	RNAi targeted against	Gene code	Normal ER network architecture	GFP::VHA-13 puncta (skewness)	Lysosome acidification	Reduction in mitochondrial membrane potential
Protein degradation	pbs-7	F39H11.5	No	No	No	No
	rpn-6.1	F57B9.10	Mixed	No	No	No
Protein synthesis (Translation)	rps-20	Y105E8A.16	No	No	No	No
	rpl-3	F13B10.2	No	No	No	No
Protein folding (Chaperones)	cct-1	T05C12.7	No	Reduced	No	No
	cct-5	C07G2.3	No	Reduced	No	No
Cytoskeleton-associated process	act-1	T04C12.6	No	No	No	No
	pat-3	ZK1058.2	No	No	No	No
	ani-2	K10B2.5	No	No	No	No
ER protein homeostasis	hsp-4	F43E2.8	Yes	No	No	No
	spcs-1	C34B2.10	Yes	No	No	No
	srpa-68	F55C5.8	Yes	No	No	No
Vesicle-mediated transport (Trafficking)	copb-2	F38E11.5	No	No	No	No
	sec-24.1	F12F6.6	Inconclusive	No	No	No
	sar-1	ZK180.4	Mixed	No	No	No
Lysosome acidification (V-ATPase)	vha-13	Y49A3A.2	Yes	No signal	No	No
	vha-12	F20B6.2	Yes	No	No	No
	vha-2	R10E11.2	Yes	No	No	No
ATP synthase	atp-3	F27C1.7	Yes	Yes	Yes	No
	atp-2	C34E10.6	Yes	Yes	Yes	No
RNA processing	ess-2	F42H10.7	No	Reduced	No	No
	cgh-1	C07H6.5	Mixed	No	No	No
	let-711	F57B9.2	Mixed	Reduced	No	No
Calcium ion transport	itr-1	F33D4.2	No	Reduced	No	No
	sca-1	K11D9.2	Yes	Reduced	No	No
Orphans	kin-2	R07E4.6	Mixed	No	No	No
	ttr-14	T05A10.3	Mixed	No	No	No
	par-5	M117.2	Yes	No	No	No
	xpo-1	ZK742.1	Yes	No	No	No
	srab-17	T11A5.2	Yes	Reduced	No	No
	rmd-2	C27H6.4	Yes	No	No	No
	dlg-1	C25F6.2a.1	No	No	No	No
	lgc-46	Y71D11A.5	Yes	Reduced	No	No
	aco-2	F54H12.1	Yes	No	No	No
	imb-1	F28B3.8	Inconclusive	No	No	No
ESCRT subunits*	vps-20	Y65B4A.3	Not tested	Yes	Yes	Not tested
	vps-28	Y87G2A.10	Not tested	Yes	Yes	Not tested
	vps-37	CD4.4	Not tested	Yes	Yes	Not tested
	vps-54	T21C9.2	Not tested	Yes	Yes	Not tested

Table 1. Functional gene categories from the proteostasis screen.
List of representative genes from each functional category that were selected for the assay pool and their effects on ER morphology, lysosome acidification, GFP::VHA-13 localization, and mitochondrial membrane potential. The ESCRT complex subunits (*) were not derived from the original screen but from subsequent candidate testing.

Sperm signals initiate V-ATPase assembly, enabling lysosomal acidification

Signals from sperm elicit the lysosomal acidification required for the clearance of protein

aggregates, and inhibiting the V-ATPase, which acidifies lysosomes, causes proteins to

aggregate in hermaphrodite oocytes (Bohnert and Kenyon, 2017). However, the mechanism by

which lysosome acidification occurs in response to signals from sperm is not known (Goudeau

et al., 2020b; Spike et al., 2018). To address this question with minimal physiological disruption,

we first attempted to fluorescently tag endogenously-expressed V-ATPase subunits using

CRISPR/Cas9. We picked four different subunits, two from the membrane-embedded V_0 domain

(VHA-4 and VHA-7) and two from the peripheral V_1 domain (VHA-11 and VHA-13) (Table S2).

Unexpectedly, although each one of these tagged proteins was visible in somatic tissues, we

were unable to detect any fluorescence in the germline in either intact animals (Figure S8) or in

their dissected gonads (Figure S9A). Germline fluorescence was also undetectable when

imaging with a more sensitive wide-field system, or following photobleaching of the strong

intestinal signal, which we reasoned might mask a faint signal from the germline (data not

shown). Endogenous genes tagged with fluorescent proteins can be silenced in the germline by

the process of epigenetic RNAe (Shirayama et al., 2012); however, this explanation seems

unlikely since, unlike V-ATPase RNAi-treated animals, these animals produced viable embryos.

We also considered the possibility that these V-ATPase proteins did not act cell-

autonomously in the gonad, though V-ATPase mRNAs previously had been identified as direct

targets of the germline-specific GLD-1 translational regulator (Wright et al., 2011). To

investigate cell autonomy, we carried out tissue-specific RNAi experiments using sterility as a

phenotypic readout. Three hermaphrodite strains [wild-type, globally RNAi-defective *rde-*

1(mkc36) mutants (DCL565), and germline-only RNAi-sensitive DCL569 animals (Zou et al.,

2019)] were subjected to RNAi from hatching (empty vector, *vha-12* and *vha-13*). Both wild-type

and germline-RNAi-permissive DCL569 animals exhibited sterility when fed *vha-12* and *vha-13*

RNAi bacteria (Figure S9B), whereas the whole-body RNAi-defective *rde-1* mutant did not.

Thus, these V-ATPase subunits are expressed and required in the germline, despite our inability

to detect them when tagged endogenously.

As a next-best approach for visualizing germline V-ATPase subunits, we obtained a

Mos1-mediated Single Copy Insertion (MosSCI) of GFP::VHA-13 driven by the germline-specific

pie-1 promoter. This construct expressed GFP::VHA-13 in a genetically-stable fashion, though

presumably at elevated levels. In this strain, we observed a striking, sperm-dependent pattern:

GFP::VHA-13 formed distinct foci in hermaphrodite oocytes (Figure 3A and S10A) that co-

localized with foci of LysoTracker, a dye that stains acidic lysosomes (Figure 3B and S11A). In

contrast, the proximal oocytes of females exhibited diffuse GFP::VHA-13 fluorescent signal

(Figure 3A and S10B), with a slight enrichment in the perinuclear region. Female oocytes

demonstrated very weak LysoTracker staining (Bohnert and Kenyon, 2017), and there was no

significant overlap in the two signals (Figure 3B and S11B). To better control for effects of

overexpression, we lowered the level of GFP::VHA-13 expression by serially diluting *gfp* RNAi

bacteria. Even at the limit of detection, hermaphrodites still exhibited GFP::VHA-13 foci, and

females exhibited a diffuse distribution (Figure S9C). Thus, the germline harbors a mechanism

by which VHA-13 is localized to discrete, acidic cellular locations, indicative of lysosomes, only

in the presence of sperm. Notably, knocking down the *vha-12* gene, which encodes another

peripheral V-ATPase V_1-domain subunit, also prevented GFP::VHA-13 foci formation in

hermaphrodite oocytes (Figure S9D), suggesting a coordinated assembly pathway.

To ask whether sperm signals regulate V-ATPase localization, we mated GFP::VHA-13-

expressing females with males and monitored their proximal oocytes. The diffuse fluorescent

signal began to change approximately 10 minutes after mating, organizing into distinct foci

within 20 to 30 minutes (Figure 3C and 3D). This change coincided with release from arrest and

oocyte maturation, ultimately leading to ovulation and fertilization. The time scales of ER

remodeling and GFP::VHA-13 relocalization (Figure 2E) were strikingly similar, suggesting a

possible causal relationship. Together these findings suggest that sperm signals lead to

lysosome acidification by initiating the recruitment of V-ATPase subunits to the surface of the

lysosome and enabling proton-pump assembly.

Figure 3

Figure 3. Sperm-dependent lysosomal localization of VHA-13.

(A) Distribution of GFP::VHA-13 in proximal oocytes of hermaphrodites (PHX1414) and females (CF4599). (B) LysoTracker-stained oocytes from *gfp::vha-13*-expressing hermaphrodites and females, for visualization of acidic lysosomes. The two signals colocalize in hermaphrodite oocytes. Dotted line; intestinal boundary, and solid line; oocytes. (C) Sperm signal-induced GFP::VHA-13 redistribution in female oocytes (CF4599). Oocytes were imaged before and after mating with CB1490 males. The GFP::VHA-13 signal shifts from diffuse to punctate following mating. Each oocyte is numbered according to its initial position relative to the spermatheca, and original numbers are maintained after fertilization. A representative individual is shown, from 5 independent mating experiments. (D) Increase in signal skewness (inhomogeneity) of GFP::VHA-13 as a function of time following mating. Unpaired t-test with Welch's correction was used to determine statistical significance, and P values are represented as $P \leq 0.05 = *$, $P \leq 0.01 = **$, and $P \leq 0.001 = ***$. Scale bars, 10 µm (A-C).

Most of the genes identified disrupt VHA-13 localization and lysosome acidification.

To ask which gene inhibitions blocked lysosome acidification, we subjected

LysoTracker-stained hermaphrodites to knockdowns of the 'assay-pool' genes (Table 1). In wild-

type hermaphrodites, lysosomes are acidified only in proximal oocytes (Bohnert and Kenyon,

2017). Therefore, to internally control for variations in dye uptake, we analyzed the ratios of

staining intensities between proximal oocytes and distal germline regions. LysoTracker staining

of hermaphrodites, females and animals subjected to empty-vector RNAi, were used as

controls, and the simultaneous presence of GFP::RHO-1 aggregates confirmed successful gene

knockdown. Unexpectedly, we found that the great majority of gene inhibitions prevented

lysosome acidification (Figure 4A and S12A). The ATP synthase genes *atp-2* and *atp-3* were

the only exceptions, consistent with previous observations suggesting that mitochondrial ATP

synthase activity acts downstream of lysosome acidification to prevent protein aggregation

(Bohnert and Kenyon, 2017).

Next, we asked whether reduced lysosomal acidification in the RNAi-treated

hermaphrodites might be caused by an inability to localize the V-ATPase subunits to lysosomes,

as occurs in females. To test this, we subjected GFP::VHA-13-expressing animals to assay-pool

RNAi and screened for the presence of GFP::VHA-13 puncta (Figure S12C). The extent of

puncta formation was quantified by measuring the skewness (non-homogeneity) of the

GFP::VHA-13 signal at multiple locations within the most proximal oocyte. Hermaphrodites and

the empty-vector RNAi-treated animals demonstrated significantly higher skewness values than

did female oocytes, which exhibited diffuse GFP::VHA-13 localization (Figure 4B). All of the

knockdowns that blocked lysosome acidification showed reduced GFP::VHA-13 skewness,

albeit to varying degrees. While most knockdowns resulted in a marked reduction in GFP::VHA-

13 skewness, similar to the female, some genes involved in protein folding (*cct-1, cct-5*), RNA

processing (*ess-2, let-711*) and calcium ion transport (*itr-1, sca-1*) demonstrated intermediate

values of skewness, indicative of at least a reduction of GFP::VHA-13 localization (Figure 4B).

Similarly, the orphan-gene inhibitions demonstrated reduced GFP skewness, with several

candidates showing only intermediate effects (*kin-2, par-5, xpo-1, rmd-2, lgc-46, aco2, imb-1*)

(Figure S12B). As anticipated, neither of the two ATP synthase gene knockdowns affected

GFP::VHA-13 puncta formation (Figure 4B). Together these data support the model that the

failure of the knockdowns to acidify lysosomes is due, at least in part, to a failure to assemble

the lysosomal V-ATPase.

Figure 4

Figure 4. Lysosome acidification and GFP::VHA-13 localization following knockdown of genes identified by the screen.

(A) LysoTracker staining in hermaphrodites (SA115) subjected to RNAi. The LysoTracker signal is represented as the ratio of proximal oocyte to distal germline staining. Staining ratios obtained from hermaphrodite (N2E) and female (CF4101) animals were used as controls (~1.3 and ~1.0, respectively). Each data point represents the average of values from three different locations in the most proximal oocyte of a single animal. The presence of GFP::RHO-1 aggregates in the same animals confirmed successful gene knockdown (not shown). (B) Analysis of GFP::VHA-13 localization, assessed by the degree of GFP skewness (inhomogeneity) in hermaphrodites (PHX1414) subjected to RNAi, and compared to control hermaphrodites and females (PHX1414 and CF4599). Each data point represents the average of ten skewness values from different regions within the first three proximal oocytes of a single animal. The presence of aggregates in GFP::RHO-1 animals (SA115) treated in parallel was used as a control to verify successful gene knockdown (not shown). For (A) and (B), Unpaired t-test with Welch's correction was used to determine the statistical significance. The top and bottom rows of significance indicators in each experiment compare the mean values from each process category to hermaphrodite and female animals, respectively. P values: $P > 0.05$ = n.s., $P \leq 0.05$ = *, $P \leq 0.01$ = **, $P \leq 0.001$ = *** and $P \leq 0.0001$ = ****.

Two distinct, mysterious roles for mitochondria in protein aggregation.

Mitochondria undergo a morphological and metabolic shift in response to signals from

sperm, a shift that is required for aggregate removal (Bohnert and Kenyon, 2017). Unfertilized

female oocytes exhibit a high mitochondrial membrane potential ($\Delta\Psi$) relative to hermaphrodite

oocytes or to female oocytes exposed to sperm by mating. Loss of the mitochondrial ATP

synthase, which discharges the proton gradient during the generation of ATP, elevates the $\Delta\Psi$

in hermaphrodites and causes protein aggregates to accumulate (Bohnert and Kenyon, 2017).

As expected, subunits from the mitochondrial ATP synthase complex (*atp-2, atp-3*) emerged

from the screen. Previously, we found that lysosomal V-ATPase activity was required for the

shift in mitochondrial membrane potential (Bohnert and Kenyon, 2017). To test the prediction

that our screen hits, which all inhibited lysosome acidification, would also block the $\Delta\Psi$ switch,

we stained hermaphrodites with the dye DiOC6(3), whose signal is sensitive to $\Delta\Psi$. Again, to

control for dye uptake, we measured the ratio of DiOC6(3) staining intensities between proximal

oocytes and the distal syncytial germline. As predicted, all of the knockdowns prevented the $\Delta\Psi$

decrease in proximal hermaphrodite oocytes, with their mitochondria exhibiting $\Delta\Psi$ values

similar to, or even higher than those of the distal germline (Figure 5A and S13A).

The observation that gene knockdowns from all the functional categories caused a

relatively high oocyte $\Delta\Psi$ value prompted us to question whether a high mitochondrial

membrane potential might be necessary for protein aggregate accumulation. To test this, we

asked whether exposing females expressing germline GFP::RHO-1 to conditions that lower $\Delta\Psi$;

namely, to electron-transport chain (ETC) RNAi knockdowns, might reduce protein aggregation.

However, neither *cyc-1* nor *nuo-2* RNAi prevented females from accumulating aggregates, even

though $\Delta\Psi$ was reduced (Figure 5B, S13B and S13C). Thus, a high mitochondrial membrane

potential is not a prerequisite for protein aggregation.

RNAi-inhibition of ATP synthase has been shown to reduce ATP levels in worms, as

does RNAi inhibition of electron transport (Dillin et al., 2002; Lee et al., 2003). To ask whether

inhibiting ATP synthase was likely to cause protein aggregation in hermaphrodites by limiting

ATP levels, we analyzed hermaphrodites subjected to ETC RNAi. The treatments reduced body

size and growth rate (data not shown), two known correlates of reduced ATP levels. However,

these perturbations did not cause protein aggregation in hermaphrodite oocytes (Figure 5B).

This finding suggests the interesting hypothesis that loss of ATP synthase causes aggregates to

accumulate in hermaphrodites not because it limits ATP levels, but for a different reason (see

Discussion).

Given the striking change in mitochondrial dynamics that takes place during oocyte

maturation, we were surprised that in our screen we recovered only one additional gene

encoding a mitochondrial-localized protein, *aco-2* (Krebs-cycle aconitase). We tested several

additional Krebs-cycle gene knockdowns (*pdha-1, idhg-1, ogdh-1, sucl-1,* and *sdhd-1*), and

found that α-ketoglutarate dehydrogenase (*ogdh-1*) and succinate dehydrogenase (*sdhd-1*)

RNAi also led to oocyte GFP::RHO-1 aggregation (Figure S13D). Notably, unlike ATP-synthase

inhibition, *aco-2* inhibition prevented lysosome acidification (Figure S12A), indicating that the

Krebs cycle and ATP synthase likely have distinct roles in the protein-aggregation pathway.

ESCRT-complex genes are required for the clearance of protein aggregates.

375   The direct mechanism by which protein aggregates are removed during oocyte

maturation is not known with certainty, although morphologically, the process resembles

lysosomal microautophagy (Bohnert and Kenyon, 2017). Microautophagy has not been

described previously in *C. elegans*; however, orthologs of genes encoding components of the

ESCRT machinery, which mediates microautophagy in yeast and mammals, are present (Sahu

et al., 2011; Williams and Urbé, 2007). We identified two ESCRT subunits (*vps-37* and *vps-54*)

in our primary screen, but they did not pass our initial validation cutoff (Table S3). To explore

the possibility that these were false negatives, we retested these RNAi clones along with RNAi

clones inhibiting additional ESCRT subunits (*tsg-101*, *vps-20* and *vps-28*). In these experiments,

the ESCRT subunits *vps-20*, *vps-28*, *vps-37*, and *vps-54* tested positive with subtle aggregation

phenotypes (although not observed in all animals tested) (Figure S14A), supporting the model

that aggregates are removed via microautophagy. Importantly, the degree of GFP::VHA-13

localization to puncta in these knockdowns was comparable to that of hermaphrodites (Figure

S14B, S14C), and they also permitted acidification of oocyte lysosomes (Figure 5C). Taken

together, these findings suggest that the ESCRT machinery operates downstream of V-ATPase

assembly and lysosome acidification to prevent aggregate accumulation.

Figure 5

Figure 5. Analysis of mitochondrial membrane potential and ESCRT-complex subunits

(A) Analysis of oocyte ΔΨ using DiOC6(3) staining in RNAi-treated hermaphrodites (OCF15), represented as the ratio of proximal oocyte to distal germline staining. Staining ratios obtained from N2E hermaphrodites (~0.8) and CF4101 females (~1) were used as controls for comparison. Each data point represents the average of values from three different locations in the most proximal oocyte of a single animal. When applicable, alteration in ER morphology was used as a control to determine successful gene knockdown in the same animals. The presence of GFP::RHO-1 aggregates in SA115 animals treated in parallel, was also used as an RNAi control (not shown). (B) Visualization of GFP::RHO-1 aggregates in oocytes of hermaphrodites (SA115) and females (CF4115) subjected to disruption of mitochondrial electron transport chain genes. Successful ETC disruption was confirmed by smaller body sizes, visible in the images and severely reduced brood sizes, similar to ATP synthase subunit knockdowns. Scale bars, 25 μm. (C) LysoTracker staining in hermaphrodites (SA115) subjected to ESCRT-complex RNAi, represented as the ratio of proximal oocyte to distal germline staining, and compared to hermaphrodites (N2E) and females (CF4101). Only animals exhibiting GFP::RHO-1 aggregates upon RNAi treatment (confirming knockdown; shown in Fig. S14A), were included in the quantification. For (A) and (C), unpaired t-test with Welch's correction was used to determine the statistical significance. The top and bottom rows of significance indicators in each experiment compare the mean values from each process category to hermaphrodite and female animals respectively. P values: $P > 0.05$ = n.s., $P \leq 0.05$ = *, $P \leq 0.01$ = **, $P \leq 0.001$ = *** and $P \leq 0.0001$ = ****.

Mechanisms enhancing germline proteostasis also operate in the soma.

Widespread proteostasis collapse is an inherent part of aging in *C. elegans* (Ben-Zvi et al., 2009; David et al., 2010; Taylor and Dillin, 2011; Walther et al., 2015), and mechanisms that extend lifespan invariably enhance cellular proteostasis. Understanding how the immortal germ lineage maintains cellular quality and proteostasis across the generations could suggest new ways to enhance proteostasis in the soma and delay organismal aging. Our lab and others have shown that many endogenous proteins become insoluble with age and can form visible aggregates when overexpressed with fluorescent tags (David et al., 2010; Huang et al., 2019; Reis-Rodrigues et al., 2012; Roux et al., 2016). To visualize age-related protein aggregation, we monitored the casein kinase subunit KIN-19, which forms insoluble aggregates in an age-dependent manner in wild-type somatic tissues (David et al., 2010). To do this, we fluorescently-tagged the endogenous *kin-19* gene specifically in the soma using split-wrmScarlet (Goudeau et al., 2020a) (Table S2). To avoid interference from the bright intestinal autofluorescence, we examined KIN-19::wrmScarlet in the head. The signal was mainly diffuse in young day-1 adults but began to exhibit a striking punctate appearance as early as day-4 of adulthood (Figure 6A, 6B, S15A, S15B and S15C). To the best of our knowledge, this tool is the first used to visualize age-dependent aggregation of an endogenously-expressed protein in *C. elegans*.

To ask whether the germline-proteostasis genes we identified in our screen might also affect somatic protein aggregation, we screened our germline-aggregation 'assay pool' for gene knockdowns that caused KIN-19 to aggregate prematurely in head tissues. We found that multiple RNAi knockdowns that impaired lysosomal acidification in the germline induced premature KIN-19::wrmScarlet aggregation in the soma, most notably RNAi of genes encoding lysosomal V-ATPase subunits (*vha-12*, *vha-13*, *vha-16* and *vha-19*), but also genes that, in the germline, were required for V-ATPase assembly (the actin gene *act-5* and the vesicle-mediated ER-to-Golgi transport gene *copb-2*) (Figure 6C and 6D). We were not able to visualize lysosome acidification in the head using LysoTracker, but these knockdowns did affect lysosome

418   acidification in the intestine, where the RNAi treatment led to fewer discrete acidic lysosome-

419   related organelles (LROs) thereby reducing the skewness of the LysoTracker signal (Figure 6E

420   and S16). We also recovered the proteasome subunits *pbs-7* and *rpn-6.1*, indicating a direct or

421   indirect role for proteasome function to maintain KIN-19 solubility in the aging soma (Figure 6C

422   and 6D).

Figure 6

A Endogenous KIN-19::wrmScarlet$_{11}$ + somatic wrmScarlet$_{1-10}$

day 1   day 4   day 7   day 11

min ▮▮▮ max

day 1   day 4   day 7   day 11

B

# of KIN-19 aggregate per animal

Days of adulthood

C

empty vector   *act-5* RNAi   *copb-2* RNAi   *pbs-7* RNAi   *rpn-6.1* RNAi

min ▮▮▮ max

empty vector   *act-5* RNAi   *copb-2* RNAi   *pbs-7* RNAi   *rpn-6.1* RNAi

vha-12 RNAi   *vha-13* RNAi   *vha-16* RNAi   *vha-19* RNAi

vha-12 RNAi   *vha-13* RNAi   *vha-16* RNAi   *vha-19* RNAi

D

Animals per category (%)

■ > 50 puncta
■ 20 to 50 puncta
□ < 20 puncta

empty vector, *act-5* RNAi, *copb-2* RNAi, *pbs-7* RNAi, *rpn-6.1* RNAi, *vha-12* RNAi, *vha-13* RNAi, *vha-16* RNAi, *vha-19* RNAi

E

Skewness of intestinal LysoTracker signal

RNAi treatment

Empty vector, *vha-12*, *vha-13*, *vha-19*, *act-5*, *copb-2*, *pbs-7*, *rpn-6.1*

Figure 6. Analysis of protein aggregation and lysosome acidification in somatic tissues.

(A) Representative images of CF4609 animals expressing KIN-19::wrmScarlet$_{11}$ and somatic wrmScarlet$_{1\text{-}10}$ at days-1, 4, 7 and 11 of adulthood. Maximum intensity projections of 3D stacks are shown in red (upper panels), and also using an intensity scale (lower panels). Scale bars, 10 μm. (B) Quantification of KIN-19::wrmScarlet aggregates per animal in the head region. Unpaired t-test with Welch's correction was used to determine the statistical significance. *$P \leq$ 0.05, ***$P \leq$ 0.001 and **** $P \leq$ 0.0001. (C) Representative images of CF4609 animals expressing KIN-19::wrmScarlet$_{11}$ and somatic wrmScarlet$_{1\text{-}10}$ subjected to RNAi or empty-vector negative control. Scale bars, 10 μm. (D) Quantification of fluorescent KIN-19 puncta in CF4609 animals subjected to RNAi or empty-vector control. (E) Skewness of the intestinal LysoTracker signal in SA115 animals subjected to RNAi (shown in Figure S16). Each data point represents the average of six independent skewness values from the anterior intestinal region, near the pharyngeal bulb, in a single animal. The simultaneous presence of GFP::RHO-1 aggregates in oocytes, was used as a control for successful gene knockdown in each animal (not shown). Unpaired t-test with Welch's correction was used to determine the statistical significance. P values: P ≤ 0.05 = *, P ≤ 0.01 = ** and P ≤ 0.001 = ***.

Discussion

Sperm-derived signals that trigger oocyte maturation in *C. elegans* initiate a global change in oocyte structure and function, including the rapid removal of protein aggregates. A key step in aggregate removal is the acidification of lysosomes, which subsequently engulf the aggregates (Bohnert and Kenyon, 2017). In addition, mitochondria, which undergo morphological and metabolic changes in response to maturation signals, appear to play an important regulatory role, as inhibition of mitochondrial ATP synthase prevents the downshift in mitochondrial membrane potential normally triggered by sperm, and causes aggregates to form even when sperm are present (Bohnert and Kenyon, 2017). In this study, we identified a key role for a third organelle, the ER, in proteostasis renewal, and, using genetic and cell biological approaches, we deepened our understanding of all three organelles, the ER, lysosomes and mitochondria, in oocyte protein-quality control.

Limits of our genetic screen

To identify additional genes and processes involved in protein-aggregate metabolism in oocytes, we carried out a genome-wide screen for RNA-inhibitions that cause proteins to aggregate even in the presence of sperm; namely, in hermaphrodites. We expected to identify genes and processes that are directly involved in the sperm-dependent proteostasis switch, as well as genes that function more indirectly in protein homeostasis and cellular quality control. Many of the genes we recovered affected large multiprotein complexes, such as the ribosome and proteasome, making a compelling argument for these complexes influencing protein aggregation, directly or indirectly. However, we may have missed processes carried out by only one or a few genes if the corresponding RNAi bacteria were missing from the RNAi library or otherwise problematic; or if their inhibition produced milder protein-aggregation phenotypes, like the ESCRT genes that failed our initial, stringent validation tests (Table S3).

To place the genes into a proteostasis pathway, we asked which hallmarks of the sperm-

triggered aggregate-clearance pathway (lysosome acidification, mitochondrial membrane-

potential shift, etc.) were blocked by the knockdowns (Figure 7). An intrinsic limitation of this

strategy is that it only identifies the first role of a gene that acts in multiple steps in the process.

For example, because inhibiting the proteasome prevents degradation of the proximal-oocyte

cell-fate determinant GLD-1 (Bohnert and Kenyon, 2017; Kisielnicka et al., 2018), and inhibiting

various other chaperones prevents lysosome acidification, we cannot determine whether these

proteins also play a direct role in the process of aggregate clearance itself.

Figure 7

Figure 7. Model: Biological processes implicated in the clearance of protein aggregates during oocyte maturation.

A role for the endoplasmic reticulum

Many of the genes we identified affected the endoplasmic reticulum (Figure 7; Table 1). All of

these gene inhibitions prevented lysosome acidification, arguing that ER function influences

aggregate formation indirectly, acting at a relatively early step in the proteostasis pathway

(though ER function could be required again, at later stages, as well). Using transmission

electron microscopy and fluorescent reporters, we validated and extended previous

observations of ER reorganization during *C. elegans* oocyte maturation (Langerak et al., 2019).

We found that the ER assumes a stacked arrangement, spatially distinct from GFP::RHO-1

protein aggregates, in the proximal oocytes of females. Importantly, we found that neither the

ER stacking, nor protein-aggregate accumulation, were a consequence of long-term oocyte

quiescence ("oocyte aging"), as both were present as soon as female oocytes appeared during

development. When oocyte maturation was triggered by mating, the ER underwent a rapid

reorganization that preceded aggregate removal, consistent with a model that ER activity might

be a precondition for aggregate clearance.

The ER-gene inhibitions fell into two classes. First was a set of genes with known ER

functions that did not appear to affect ER morphology: *hsp-4* (BIP)*, spcs-1* (signal peptidase)*,

and *srpa-68* (signal recognition protein). This finding argues that ER reorganization does not

require ER function itself. The second class did affect ER morphology, generally causing the ER

from hermaphrodites to exhibit a bright patchy appearance characteristic of (though not

necessarily identical to) that seen in female oocytes. Some of these gene knockdowns, such as

that of *itr-1,* which encodes the channel that releases calcium stored in the ER, likely cause both

sexes to adopt a female-specific, stacked ER morphology. In *C. elegans*, the major wave of

oocyte calcium release occurs after fertilization; this finding suggests that ER-dependent

calcium metabolism plays an important earlier role as well. Other RNAi knockdowns may simply

disrupt ER morphology more generally and might not identify genes that play an active role in

the oocyte proteostasis switch. For example, the actin cytoskeleton is known to influence ER

morphogenesis (Poteryaev et al., 2005), and we recovered a number of cytoskeletal gene hits.

Notably, TRiC-complex chaperonin genes fell into this class as well. The TRiC-complex

chaperonin can counteract protein aggregation directly; for example, by promoting correct

protein folding during protein synthesis. However, TRiC is known to assist in folding cytoskeletal

components including actin and tubulin (Llorca et al., 2000) and it was interesting to find that the

TRiC-complex genes we recovered, *cct-1* and *cct-5,* acted indirectly, by affecting ER

morphology and lysosome acidification. As mentioned above, the proteasome, too, affects

aggregate clearance indirectly. In general, it might be valuable to assay lysosome acidification

when analyzing the functions of these chaperone/chaperonin systems on the accumulation of

protein aggregates in other cellular systems, to test for the possibility of indirect effects. That

said, it is also possible that TRiC-complex and other chaperones act at multiple points in the

proteostasis pathway, with both indirect and direct effects on protein aggregation.

Finally, we identified several genes that, to our knowledge, were not previously known to

influence ER morphology; namely, *dlg-1*, *kin-2, and ttr-14* (Table 1). How these genes act,

directly or indirectly, to influence the ER could be interesting to explore in the future.

Regulation of lysosome acidification

A central goal of this study was to better understand how lysosomes are acidified during oocyte

maturation. To address this issue, we first generated functional, fluorescently-tagged V-ATPase

subunits expressed from their own loci. However, for reasons that we do not understand, none

of these proteins was visible in the germline. We then obtained a stable, single-copy MosSCI

strain in which *gfp::vha-13* was expressed from a germline-specific promoter. In these animals,

GFP::VHA-13 was visible, and we observed a striking sperm-dependent localization pattern. In

the presence of sperm (in hermaphrodites), GFP::VHA-13 was located on lysosomes; that is, it

assumed a punctate appearance that co-localized with LysoTracker signal. In the absence of

sperm (in unmated females), the protein was distributed throughout the cytoplasm (Figure 7).

The simplest interpretation of these findings is that signals from sperm trigger the localization of

GFP::VHA-13 to lysosomes, and that this localization enables lysosome acidification. In

principle, additional levels of control, such as translational repression of V-ATPase synthesis,

could operate as well, since our overexpressed construct could potentially override negative

regulators; though we observed no evidence of this with serially-diluted *gfp* RNAi treatments

that progressively reduced visible GFP::VHA-13 protein levels.

In yeast, peripheral, cytoplasmic V_1 subunits of the V-ATPase (such as VHA-13) localize

to lysosomes when the membrane-localized V_0 sector is trafficked there from the ER, where it is

assembled (Graham et al., 1998). Thus, we propose that the extensive rearrangement in ER

morphology triggered by sperm in *C. elegans* oocytes is coupled to movement of either pre-

existing or newly-synthesized membrane-localized V_0 subunits from the ER to lysosomes,

where they create docking sites for cytoplasmic, peripheral V_1 subunits. Consistent with this

idea, RNAi of the ER genes, as well as each of the three genes required for vesicle trafficking

from the ER to lysosomes, prevented GFP::VHA-13 localization to lysosomes, as did inhibiting

genes encoding the membrane-localized V-ATPase V_0 subunits. Unfortunately, our efforts to

observe the predicted sperm-activated movement of V_0 subunits from the ER to lysosomes were

thwarted by our inability to visualize either of two functional GFP-tagged V_0 subunits, even when

overexpressed using MosSCI technology. In the future, immuno-gold EM labeling of

endogenous V_0 subunits might enable this model to be tested more directly.

We note that an interesting puzzle remains unexplained: If the multi-subunit lysosomal

V-ATPase complexes are present in such low amounts in mature oocytes that they cannot be

seen when we fluorescently tag the endogenous *vha-13* gene, then why do they appear when

the tagged VHA-13 protein is overexpressed? How does the excess GFP::VHA-13 protein dock

onto lysosomes? Perhaps the ER supplies lysosomes with alternative docking sites for

unassembled V-ATPase subunits, or perhaps the stoichiometry change due to the excess

GFP::VHA-13 subunits triggers the production of entire new V-ATPase complexes that

incorporate the overexpressed GFP::VHA-13 subunits. This remains an unexplained but

intriguing mystery (see (Hughes and Gottschling, 2012)).

The gene knockdowns identified by the screen nearly all failed to acidify lysosomes,

reinforcing the central role of acidified lysosomes in clearing oocyte protein aggregates (Figure

7). Moreover, none of the knockdowns that failed to acidify lysosomes exhibited a punctate,

lysosomal GFP::VHA-13 pattern (Table 1). This finding suggests that V-ATPase activity may not

only be necessary but also sufficient for lysosome acidification; however, this interpretation

remains tentative, since we may simply have failed to identify the relevant genes in our screen.

Finally, several of the knockdowns from our screen did not appear to affect ER morphology (and

have not been reported to affect ER function), but did prevent V-ATPase localization, for

example *srab-17*, a chemosensory gene of the serpentine receptor class ab, and *lgc-46,*

involved in chloride ion transport (Table 1). How, directly or indirectly, these proteins might

influence V-ATPase localization and assembly is an interesting question to pursue in the future.

Two Roles for Mitochondria

Like the ER and lysosomes, mitochondria undergo morphological and functional changes during

C. elegans oocyte maturation. In response to sperm, the mitochondria of proximal oocytes

reduce their relatively high membrane potential and shift from a fragmented to a tubular

structure. Inhibiting mitochondrial ATP synthase preserves this high membrane potential even

when sperm are present and causes aggregates to accumulate. Injecting ADP, whose levels

would be expected to rise as maturing oocytes initiate protein synthesis and acidify lysosomes,

reduces the mitochondrial membrane potential in female oocytes, in an ATP-synthase

dependent fashion. Together these findings suggested the model that in the absence of sperm,

mitochondria are held in a quiescent, poised state that relaxes as metabolic activity and ATP

synthesis commences (Bohnert and Kenyon, 2017).

Whether *C. elegans* oocyte maturation is powered primarily by glycolysis or respiration is

unknown; however, in principle, the accumulation of protein aggregates caused by inhibiting

mitochondrial ATP synthase could reflect ATP limitation. However, if this were the case, then

inhibiting components of the electron transport chain should also cause protein aggregates to

accumulate in hermaphrodites, since this, too, limits ATP levels. However, though these

knockdowns affected body size, growth rates and brood size, this was not the case; ETC

knockdowns had no effect on protein aggregation. This unexpected finding suggests that ATP-

synthase knockdown prevents aggregate clearance not because of changes in ATP levels, but

instead either because a continuously high mitochondrial membrane potential acts like a

checkpoint to inhibit aggregate clearance (Figure 7), or because of another, unknown role of

ATP synthase in protein homeostasis.

Our screen revealed an additional requirement for mitochondria in protein homeostasis;

specifically, for Krebs cycle activity. Inhibiting mitochondrial aconitase (*aco-2)* produced a

different phenotype from inhibiting ATP synthase; namely, a failure to acidify lysosomes. Why

might this be? Again, the explanation seems unlikely to involve energy limitation, since inhibiting

ETC components or ATP synthase did not produce this phenotype. One possibility is that

inhibiting the Krebs cycle affects lysosome acidification by affecting redox potential, as several

Krebs cycle enzymes affecting protein aggregation (*ogdh-1* and *sdhd-1*) encode

dehydrogenases that produce the reducing agents NADH and $FADH_2$, and aconitase supplies

the substrate for another Krebs cycle dehydrogenase. Alternatively, other metabolites generated

via Krebs cycle intermediates could influence the events leading to lysosome acidification. It will

be interesting to explore this mechanism in the future.

Microautophagy

Previously, we observed lysosomes engulfing protein aggregates in a process reminiscent of

microautophagy (Bohnert and Kenyon, 2017). Before these observations, microautophagy had

not been described in *C. elegans.* In this study, using a candidate approach, we showed that

inhibiting ESCRT-complex members, which are known to mediate microautophagy in other

organisms, prevents aggregate accumulation in maturing *C. elegans* oocytes. Moreover, they

influence protein aggregation independently of lysosomal acidification, consistent with their

having a direct role. These findings strengthen the case that microautophagy is an active

proteostasis mechanism in *C. elegans,* and likely the mechanism by which aggregates are

cleared in response to oocyte-maturation signals (Figure 7).

Implications for somatic proteostasis and aging

Together our findings all point towards lysosomal acidification being a key, determinative event

in oocyte protein-aggregate clearance. Moreover, the screen hits that affected the lysosomal V-

ATPase and COPII (ER-to-lysosomal vesicle transport) had a striking ability to accelerate age-

dependent protein aggregation in somatic cells. Thus, the same mechanisms that enhance

oocyte proteostasis in the immortal germline likely enhance somatic proteostasis as well. These

findings may have implications for aging and longevity more broadly. In yeast, lysosomal acidity

declines with age, and overexpressing the yeast ortholog of the *C. elegans* V-ATPase gene

vha-13 delays this decline and extends replicative lifespan (Hughes and Gottschling, 2012). In

worms, lysosome acidification also declines with age, and this decline is delayed in long-lived

mutants (Baxi et al., 2017; Sun et al., 2020). Whether increasing V-ATPase function might be

sufficient to extend lifespan in *C. elegans* is unclear, but overproduction of the transcription

factor HLH-30/TFEB, which augments many lysosomal functions, has been shown to do so

(Lapierre et al., 2013; O'Rourke and Ruvkun, 2013), and in mammals, similar perturbations slow

the aging of the immune system (Zhang et al., 2019).

In summary, a genetic screen for proteostasis enhancers in the immortal germ lineage

has revealed a landscape of interconnected processes, spanning protein translation,

cytoskeletal dynamics, Krebs cycle activity, ER activation and vesicle transport, that together

enable lysosome acidification and clearance of protein aggregates during oocyte maturation.

Lysosomal acidification maintains proteostasis in the aging soma as well. In addition to raising

many new questions for future investigations, our findings support a growing body of evidence

that interventions that rejuvenate lysosomal activity during aging might yield new ways to

increase human health and longevity.

Acknowledgements:

We thank colleagues in the Kenyon lab and at Calico Life Sciences for advice and discussion,

and Daniel Gottschling and Calvin Jan for comments on the manuscript. Special thanks to

Andrew Dillin (University of California, Berkeley) for sharing his transmission electron

microscopy resources, and Peichuan Zhang for help with converting the RNAi collection format.

We thank Robert Keyser for help with liquid handling robotics for the primary screen, and Kayley

Hake for suggesting skewness as a way to capture aggregate formation in microscopy images.

Peichuan Zhang and Rex Kerr helped to analyze somatic KIN-19 aggregation. We thank Di

Chen for sharing the DCL565 and DCL569 strains. Some *C. elegans* strains were provided by

the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440).

The endogenously tagged V-ATPase *C. elegans* strains were generated by SunyBiotech,

China. Some illustrations in Figures 1 and 7 were created with BioRender.com. The study was

supported by Calico Life Sciences and David H. Hall was supported by NIH OD010943.

Author contributions:

Ma.S and J.G executed all experiments with the exception of transmission electron microscopy,

which was performed by Me.S, in Andrew Dillin's laboratory. D.H.H provided advice for the TEM

experiments and helped in the interpretation of the TEM images. K.A.B provided valuable inputs

in the early stages of the project and contributed to the ER and ETC analysis. M.I. provided help

and advice for fluorescence microscopy experiments and data analysis. The research and

analysis was designed by Ma.S, J.G and C.K, who also prepared the final manuscript.

Materials and Methods:

C. elegans **strains and maintenance**

All strains used in this study are listed in Table S2. Hermaphrodites expressing germline

transgenes were maintained at 25°C to delay silencing. Mutants carrying the thermosensitive,

feminizing *fem-1(hc17ts)* allele, which causes cells that would normally become sperm instead

to become oocytes, were maintained at the permissive temperature of 15°C and moved to 25°C

for feminization prior to hatching, or at the L1 stage. Animals were grown under standard

laboratory conditions (Brenner, 1974) and maintained on Nematode Growth Medium (NGM)

seeded with OP50 bacteria, unless otherwise indicated.

Generation of *C. elegans* **strains**

a. Strain generation: genetic crosses.

The following *C. elegans* strains were created by standard genetic crosses: CF4542 *fem-*

1(hc17ts) IV; ocfIs2[Ppie-1:mCherry::sp12::pie-1 3'UTR, Cbr-unc-119(+)], CF4552 tjIs1[Ppie-

1::gfp::rho-1, Cbr-unc-119(+)]; ocfIs2 [Ppie-1:mCherry::sp12::pie-1 3'UTR, Cbr-unc-119(+)],

CF4557 fem-1(hc17ts) IV; tjIs1[Ppie-1::gfp::rho-1, Cbr-unc-119(+)]; ocfIs2 [Ppie-

1:mCherry::sp12::pie-1 3'UTR, Cbr-unc-119(+)], CF4559 zuIs45[Pnmy-2::nmy-2::gfp, Cbr-unc-

119(+)] V; ocfIs2 [Ppie-1:mCherry::sp12::pie-1 3'UTR, Cbr-unc-119(+)], CF4560 fem-1(hc17ts)

IV; zuIs45[Pnmy-2::nmy-2::gfp, Cbr-unc-119(+)] V; ocfIs2 [Ppie-1:mCherry::sp12::pie-1 3'UTR,

Cbr-unc-119(+)] and CF4599 *sybSi35[Ppie-1::gfp::vha-13::vha-13 3'UTR + Cbr-unc-119(+)] II;*

unc-119(ed3/+) III; fem-1(hc17ts) IV.

b. Strain generation: Mos1-mediated single-copy insertion.

The strains PHX1414 *sybSi35[Ppie-1::gfp::vha-13::vha-13 3'UTR + Cbr-unc-119(+)] II; unc-*

119(ed3) III and PHX2690 *sybIs2690[Ppie-1::vha-7::wrmScarlet::vha-7 3'UTR, Cbr-unc-119(+)]*

I; unc-119(ed3) III were generated by SunyBiotech using MosSCI services.

c. Strain generation: CRISPR/Cas9-triggered homologous recombination.

WrmScarlet$_{11}$ was introduced at the C-terminus of KIN-19 using CRISPR/Cas9 and single-

stranded oligodeoxynucleotide template (ssODN) in the strain CF4582 expressing wrmScarlet$_{1-}$

$_{10}$ in somatic tissues (driven by the *eft-3* promoter and *unc-54* 3'UTR), to obtain the strain

CF4609. CRISPR insertion of wrmScarlet$_{11}$ was performed following published protocols

(Goudeau et al., 2020a; Paix et al., 2016, 2015). Briefly, ribonucleoprotein complexes (protein

Cas9, tracrRNA, crRNA) and ssODN were microinjected into the gonad of young adults using

standard methods (Evans, 2006). Injected worms were singled and placed at 25ºC overnight.

The crRNA and ssODN template sequences used to generate wrmScarlet$_{11}$ knock-in at the C

terminus of *kin-19* are the following:

ssDNA_kin-19C:

ACAGGGAGCTACCGTTCCATCAGCTGGAGTTCCAGCTGGAGTTGCACCAGGAGGAACTAC

TCCACAGGGAGGAGGATCCTACACCGTCGTCGAGCAATACGAGAAGTCCGTCGCCCGTCA

CTGCACCGGAGGATAAgacgttttgttgccgtcctgaggctttttaatccaaaaaagcccatgttaaatcatgtactatc

kin-19_C_crRNA: GGCAACAAAACGTCTTACTG(TGG).

KIN-19::wrmScarlet$_{11}$ integrants were identified by screening for fluorescent progeny in the

progeny of injected worms.

Strains PHX731 *vha-13(syb731[wrmScarlet::vha-13]) V*, PHX1049 *vha-13(syb1049[gfp::vha-*

13]) V, PHX1198 *ieSi65[sun-1p::TIR1::sun-1 3'UTR + Cbr-unc-119(+)] II; unc-119(ed3) III; vha-*

11(syb1198[vha-11::gfp::degron]/+) IV, PHX1363 *vha-4(syb1363[wrmScarlet::vha-4]/+) II*, and

PHX1317 *vha-7(syb1317[vha-7::wrmScarlet]) IV* were generated by SunyBiotech using CRISPR

services for the Kenyon lab.

d. Nucleic acid reagents. Synthetic nucleic acids were purchased from Integrated DNA

Technologies (IDT). For wrmScarlet$_{11}$ knock-in, a 200-mer HDR template was ordered in

ssODN form (synthetic single-stranded oligodeoxynucleotide donors) from IDT.

RNA interference

Unless otherwise indicated, RNA interference was initiated in the early L4 larval stage, prior to

formation of oocytes but following much of development. One exception was *fem-1* RNAi, which

was initiated at hatching to ensure feminization. RNAi-bacterial feeding was initiated and

maintained for 36-40 hours prior to imaging to allow sufficient time for knockdown.

a. Genome-wide screen

The Ahringer *C. elegans* RNAi collection containing 18,225 bacterial clones, targeting around

87% of currently annotated *C. elegans* genes (Source BioScience), was used for screening. A

workflow for screening in liquid (Lehner et al., 2006) was modified suitably for our needs. For a

single 96-well plate, SA115 (GFP::RHO-1) hermaphrodites were grown on NGM plates seeded

with thick OP50 lawns (from 2 medium sized plates) for ~4 days to obtain a high density of

healthy and gravid young adults.

Day 1: Animals were subjected to standard bleaching and the eggs were washed thoroughly

and then incubated in M9 buffer overnight at 20°C on a rotating mixer to obtain synchronized

L1-arrested animals.

Day 2: L1 animals were transferred to NGM plates spotted with HT115 bacteria and incubated

24-28 hrs at 25°C to reach the early L4 stage. The RNAi-library plate was aseptically replicated

into deep well plates (96-well, square; Greiner Bio-One) containing 1 mL LB (with 25 µg/ml

carbenicillin and 12.5 µg/ml tetracycline) per well. Positive (*vha-13* and *atp-2*) and negative

(empty L4440 vector) controls were inoculated manually into blank wells if available, or into

wells on a separate control plate. The cultures were grown overnight at 37°C with shaking (185

rpm).

Day 3: Fresh media (1 mL LB with 25 µg/ml carbenicillin) was added to the wells and incubated

at 37°C, 155 rpm for 2 hrs. Subsequently, 4 mM isopropyl β-d-1-thiogalactopyranoside (IPTG;

Promega) was added to each RNAi well for induction of dsRNA synthesis and plates were

incubated for 3-4 hours at 30°C and 155 rpm. Synchronized L4 animals were washed off the

HT115 plates with M9 and resuspended in NGM broth (with 25 µg/ml carbenicillin and 1 mM

IPTG). This worm suspension (20 µL, ~10-15 animals) was added to each well of a Cellvis flat-

bottom glass plate with high performance #1.5 cover glass. The deep-well plate with dsRNA-

induced RNAi bacteria was centrifuged and the cells were resuspended in the same NGM broth

as the worms. 35 µL of each cell suspension was transferred into the corresponding well of the

Cellvis plate already seeded with L4 animals for bacterial RNAi feeding. The feeding plate was

incubated at 20°C, 155 rpm with adequate humidity in the shaker to prevent worms from settling

or drying out.

Day 5: After checking the for presence of adults with fully developed germlines*,* 25 mM sodium

azide (Sigma-Aldrich) was added to each well and the plate was subjected directly to imaging

on a spinning-disk confocal microscope. Typically, ~6 such 96-well plates were handled

simultaneously in a given round of screening. The presence of aggregates in 50% or more

animals for a specific RNAi condition was considered a hit.

Notes: The ability to automate image acquisition was complicated by a combination of low

GFP::RHO-1 reporter expression and the close apposition of gonads with the intestine, an

organ with significant autofluorescence. The density of animals and temperature of incubation

determine developmental rate and were optimized by multiple pilot trials. Liquid handling was

performed using Multidrop Combi from Thermo Scientific (initial inoculation of RNAi bacteria;

Day 2) and Agilent Bravo (resuspension and mixing of worms and induced bacteria; Day 3).

Following the validation rounds, the identities of the RNAi clones emerging from the screen

were verified by sequencing (Quintara Biosciences). Investigators were blinded to identity of

RNAi clones, except controls, during screening.

b. RNAi feeding on agar plates

Individual cultures were inoculated in tubes containing 1mL LB (with 25 µg/ml carbenicillin and

12.5 µg/ml tetracycline) and incubated at 37°C with shaking (200 rpm). The cultures were

diluted 1:10 in fresh LB containing only carbenicillin and allowed to grow for a further 2 hrs, after

which IPTG was added (4 mM) and tubes moved to a 30°C shaker for dsRNA induction (3-4

hours). Induced cultures were spotted on NGM plates containing 50 µg/ml carbenicillin and 2

mM IPTG and the bacterial lawn was allowed to grow for 24-36 hrs at 30°C, prior to introducing

synchronized L4 animals.

c. Somatic KIN-19 protein-aggregation RNAi screen

KIN-19::wrmScarlet-expressing CF4609 animals were grown from a synchronized population of

L1 animals on HT115 bacteria until they reached the L3 stage and then were transferred to

plates seeded with RNAi bacteria until they reached day-2 of adulthood. The animals' heads

were imaged by confocal microscopy. Z-stacks were analyzed using ImageJ (Fiji); image

manipulations consisted of maximum intensity projections. The images of KIN-19::wrmScarlet

puncta were quantified by the investigator who acquired the images (Figure 6B), as well as by

three blinded investigators (Figure S15).

Fluorescence Microscopy

Confocal fluorescence images were acquired on a Nikon Eclipse Ti CSU-X1 equipped with 405, 488, 561 and 640 nm lasers. Emission was collected through 455/50, 525/36, 605/70 or 700/75 nm filters on an Andor iXon Ultra EMCCD camera, using the NIS-Elements software. Images were visualized using ImageJ Fiji. Maintenance and selection of transgenic worms expressing fluorescent markers were performed using a Leica M165 FC fluorescent stereo microscope equipped with a Sola SE-V light source. Photobleaching and wide-field experiments were performed using a Leica TIRF microscope equipped with an infinity scanner, using 488 nm laser excitation. Emission was collected through a GFP-T filter cube on a Hamamatsu ORCA-Flash4 camera.

a. Preparing worms for microscopy

Worms imaged in 96-well plates were immobilized by adding sodium azide to the liquid medium at a concentration of 25 mM (except when measuring mitochondrial $\Delta\Psi$, where the medium was carefully removed, and animals immobilized with levamisole). For all other experiments, animals were picked from plates and placed in a drop of 2 mM levamisole (Sigma) on a 4% agarose pad resting on a glass slide, and then secured with a coverslip.

b. Dye staining

For experiments involving several different RNAi-knockdown conditions, RNAi feeding was set up similar to the screen workflow. The dyes LysoTracker Red DND-99 and DiOC6(3) (both from Life Technologies) were added to the liquid NGM at 2 µM concentration prior to resuspension of dsRNA-induced RNAi-bacterial cells. After addition of synchronized L4 animals, feeding and dye uptake were allowed to proceed at 20°C for 36-40 hrs. The 96-well feeding plates were protected from light during this time. The plate was centrifuged, bacterial suspensions were carefully washed off, and the animals were resuspended in fresh M9 buffer for imaging. For

experiments involving fewer knockdown conditions or pairwise comparisons between

hermaphrodites and females, the dye-feeding was performed on NGM Agar plates as described

previously (Bohnert and Kenyon, 2017).

To analyze lysosome acidification in maturing oocytes, the ratio of signal intensities between

proximal oocytes and the distal germline were calculated. In case of intestinal LysoTracker

staining, signal intensities were not measured due to highly variable dye uptake and staining;

and signal skewness (inhomogeneity) was measured instead. In both cases, multiple regions of

oocytes or the anterior intestine were sampled for intensity or skewness values; and averaged

for a single animal.

Mating experiments

An individual female animal was immobilized on an agarose pad using 2 mM levamisole, and

the reporter of interest was imaged in the anterior gonad arm. Levamisole was washed off

immediately, and the animal was introduced to an NGM plate containing young CB1490 male

animals in a small spot of OP50 bacteria (5 males:1 female). The interactions were monitored

using a benchtop stereo microscope and when mating occurred, the female was transferred

immediately to the agarose pad and imaged again, ensuring that the same oocytes were

visualized before and after mating. Successful mating events were tracked by gonadal sheath

contractions in response to sperm and subsequently confirmed by successful ovulation.

Changes within individual oocytes upon mating were monitored up to 90 minutes at 10-minute

intervals. An identical procedure was performed for control females, except for exposure to

males. Specific regions within proximal oocytes of control female animals were compared at all

time points to determine the extent of signal intensity change due to photobleaching, with

increasing time (Figure S6B). In addition, regions of the distal germline were also monitored in

mated animals (data not shown) to verify that the disappearance of the signal representing ER

clusters and protein aggregates was not due to photobleaching.

KIN-19::wrmScarlet aggregate accumulation in somatic tissues

A synchronized population of KIN-19::wrmScarlet-expressing CF4609 animals were grown on OP50 from L1 at 20°C. Worms were transferred every other day to freshly-seeded NGM plates to remove progeny. Worms were imaged by confocal microscopy at day-1, day-4, day-7 and day-11 of adulthood.

Transmission electron microscopy

N2E hermaphrodites and *fem-1(hc17ts)* females were cultured for five generations at 15°C without starvation or contamination at any point. Animals from 12 small plates of each strain were bleached, and the synchronized eggs were collected and washed thoroughly in M9 and strained through 40 µm strainers to remove adult carcasses. Eggs were then resuspended in fresh M9 and incubated at 25°C overnight on a rotating mixer. Synchronized L1 animals were moved to NGM plates seeded with OP50 and incubated at 25°C (~50 animals per plate). Newly-developed day-1 adults were further processed according to standard *C. elegans* TEM techniques (Hall et al., 2012). Briefly, the animals were loaded into 100 µm-deep specimen carriers and high-pressure frozen in a Bal-tec HPM (approximately 20 worms per carrier). After freezing, the carriers were transferred to vials containing Freeze Substitution (FS) media (1% OsO4 and 0.1% uranyl acetate in acetone), while still submerged in liquid N_2. Frozen samples were slowly warmed to approximately -15°C while being agitated on a rotating platform (adapted from (McDonald and Webb, 2011)). Following freeze substitution, samples were infiltrated with Epon Resin/Acetone series [25% Resin (1hr), 50% Resin (1hr), 75% Resin (overnight), 100% resin (1hr), 100% resin (1hr) and the resin from last step replaced with fresh resin for embedding] and subjected to polymerization at 60°C for 48 hrs. For staining, thin sections (70 nm thick) were cut on an ultramicrotome, picked up on Formvar-coated 50 mesh grids and post-

stained for 7 min in 2% uranyl acetate (aq), followed by 7 min in lead citrate. Images were

acquired on a Tecnai 12 Transmission Electron Microscope.

Data analysis and statistics

ImageJ Fiji was used to quantify and analyze attributes from image data, and images from a

given experimental set were scaled and processed identically for comparison. Preparation of

graphs and statistical analyses were performed using GraphPad Prism 8. Statistical tests used

to determine significance and p-values are described in the figure legends of each experiment.

No statistical methods were used to predetermine sample sizes. Investigators were blinded to

identity of RNAi clones (except for controls) while imaging in 96-well plates.

References:

Baxi K, Ghavidel A, Waddell B, Harkness TA, de Carvalho CE. 2017. Regulation of lysosomal

function by the daf-16 forkhead transcription factor couples reproduction to aging in

caenorhabditis elegans. *Genetics*. doi:10.1534/genetics.117.204222

Ben-Zvi A, Miller EA, Morimoto RI. 2009. Collapse of proteostasis represents an early molecular

event in Caenorhabditis elegans aging. *Proc Natl Acad Sci U S A*.

doi:10.1073/pnas.0902882106

Bluemink JG, Hage WJ, van den Hoef MH, Dictus WJ. 1983. Freeze-fracture electron

microscopy of membrane changes in progesterone-induced maturing oocytes and eggs of

Xenopus laevis. *Eur J Cell Biol*.

Bohnert KA, Kenyon C. 2017. A lysosomal switch triggers proteostasis renewal in the immortal

C. Elegans germ lineage. *Nature* **551**:629–633. doi:10.1038/nature24620

Brenner S. 1974. The genetics of Caenorhabditis elegans. *Genetics*.

Carroll J, Jones KT, Whittingham DG. 1996. Ca2+ release and the development of Ca2+

release mechanisms during oocyte maturation: A prelude to fertilization. *Rev Reprod*

1:137–143. doi:10.1530/ror.0.0010137

Charbonneau M, Grey RD. 1984. The onset of activation responsiveness during maturation

coincides with the formation of the cortical endoplasmic reticulum in oocytes of Xenopus

laevis. *Dev Biol*. doi:10.1016/0012-1606(84)90177-5

Crittenden SL, Troemel ER, Evans TC, Kimble J. 1994. GLP-1 is localized to the mitotic region

of the C. elegans germ line. *Development* **120**:2901–2911.

David DC, Ollikainen N, Trinidad JC, Cary MP, Burlingame AL, Kenyon C. 2010. Widespread

protein aggregation as an inherent part of aging in C. elegans. *PLoS Biol* **8**:47–48.

doi:10.1371/journal.pbio.1000450

Dillin A, Hsu AL, Arantes-Oliveira N, Lehrer-Graiwer J, Hsin H, Fraser AG, Kamath RS, Ahringer

J, Kenyon C. 2002. Rates of behavior and aging specified by mitochondrial function during

development. *Science (80-)*. doi:10.1126/science.1077780

Evans T. 2006. Transformation and microinjection. *WormBook*. doi:10.1895/wormbook.1.108.1

Fan W, Waymire KG, Narula N, Li P, Rocher C, Coskun PE, Vannan MA, Narula J, MacGregor

GR, Wallace DC. 2008. A mouse model of mitochondrial disease reveals germline

selection against severe mtDNA mutations. *Science (80-)*. doi:10.1126/science.1147786

FitzHarris G, Marangos P, Carroll J. 2007. Changes in endoplasmic reticulum structure during

mouse oocyte maturation are controlled by the cytoskeleton and cytoplasmic dynein. *Dev*

Biol **305**:133–144. doi:10.1016/j.ydbio.2007.02.006

Francis R, Barton M, Kimble J, Schedl T. 1995. gld-1, a tumor suppressor gene required for

oocyte development in Caenorhabditis elegans. *Genetics* **139**:579–606.

Fredriksson Å, Johansson Krogh E, Hernebring M, Pettersson E, Javadi A, Almstedt A, Nyström

T. 2012. Effects of aging and reproduction on protein quality control in soma and gametes

of Drosophila melanogaster. *Aging Cell*. doi:10.1111/j.1474-9726.2012.00823.x

Gawriluk TR, Hale AN, Flaws JA, Dillon CP, Green DR, Rucker EB. 2011. Autophagy is a cell

survival program for female germ cells in the murine ovary. *Reproduction*.

doi:10.1530/REP-10-0489

Goudeau J, Aguilaniu H. 2010. Carbonylated proteins are eliminated during reproduction in C.

elegans. *Aging Cell* **9**:991–1003. doi:10.1111/j.1474-9726.2010.00625.x

Goudeau J, Paw J, Savy L, Leonetti MD, York A, Kenyon C, Ingaramo M. 2020a. Split-

wrmScarlet and split-sfGFP: tools for faster, easier fluorescent labeling of endogenous

proteins in Caenorhabditis elegans. *bioRxiv*. doi:10.1101/2020.07.02.185249

Goudeau J, Samaddar M, Bohnert KA, Kenyon C. 2020b. Addendum: A lysosomal switch

triggers proteostasis renewal in the immortal C. elegans germ lineage. *Nature* **580**:2020.

doi:10.1038/s41586-020-2108-0

Graham LA, Hill KJ, Stevens TH. 1998. Assembly of the Yeast Vacuolar H+-ATPase Occurs in

the Endoplasmic Reticulum and Requires a Vma12p/Vma22p Assembly Complex. *J Cell*

Biol **142**:39–49.

Hall DH, Hartwieg E, Nguyen KCQ. 2012. Modern Electron Microscopy Methods for C.

elegansMethods in Cell Biology. Academic Press Inc. pp. 93–149. doi:10.1016/B978-0-12-

394620-1.00004-7

Huang C, Wagner-Valladolid S, Stephens AD, Jung R, Poudel C, Sinnige T, Lechler MC,

Schörit N, Lu M, Laine RF, Michel CH, Vendruscolo M, Kaminski CF, Schierle GSK, David

DC. 2019. Intrinsically aggregation-prone proteins form amyloid-like aggregates and

contribute to tissue aging in caenorhabditis elegans. *Elife*. doi:10.7554/eLife.43059

Hubbard EJA, Greenstein D. 2005. Introduction to the germ line. *WormBook*.

doi:10.1895/wormbook.1.18.1

Huelgas-Morales G, Greenstein D. 2018. Control of oocyte meiotic maturation in C. elegans.

Semin Cell Dev Biol. doi:10.1016/j.semcdb.2017.12.005

Hughes AL, Gottschling DE. 2012. An early age increase in vacuolar pH limits mitochondrial

function and lifespan in yeast. *Nature* **492**:261–265. doi:10.1038/nature11654

Hunter N. 2017. Oocyte Quality Control: Causes, Mechanisms, and Consequences. *Cold Spring*

Harb Symp Quant Biol **82**:235–247. doi:10.1101/sqb.2017.82.035394

Jones AR, Francis R, Schedl T. 1996. GLD-1, a cytoplasmic protein essential for oocyte

differentiation, shows stage- ans sex-specific expression during Caenorhabditis elegans

germline development. *Dev Biol*. doi:10.1006/dbio.1996.0293

Joseph-Strauss D, Gorjánácz M, Santarella-Mellwig R, Voronina E, Audhya A, Cohen-Fix O.

2012. Sm protein down-regulation leads to defects in nuclear pore complex disassembly

and distribution in C. elegans embryos. *Dev Biol* **365**:445–457.

doi:10.1016/j.ydbio.2012.02.036

Kamath RS, Fraser AG, Dong Y, Poulin G, Durbin R, Gotta M, Kanapin A, Le Bot N, Moreno S,

Sohrmann M, Welchman DP, Zipperien P, Ahringer J. 2003. Systematic functional analysis

of the Caenorhabditis elegans genome using RNAi. *Nature* **421**:231–237.

doi:10.1038/nature01278

Kim S, Spike C, Greenstein D. 2013. Control of oocyte growth and meiotic maturation in

caenorhabditis elegans. *Adv Exp Med Biol* **757**:277–320. doi:10.1007/978-1-4614-4015-4-

10

King GA, Goodman JS, Schick JG, Chetlapalli K, Jorgens DM, McDonald KL, Ünal E. 2019.

Meiotic cellular rejuvenation is coupled to nuclear remodeling in budding yeast. *Elife* **8**.

doi:10.7554/eLife.47156

Kisielnicka E, Minasaki R, Eckmann CR. 2018. MAPK signaling couples SCF-mediated

degradation of translational regulators to oocyte meiotic progression. *Proc Natl Acad Sci U*

S A **115**:E2772–E2781. doi:10.1073/pnas.1715439115

Kobayashi H, Minshull J, Ford C, Golsteyn R, Poon R, Hunt T. 1991. On the synthesis and

destruction of A- and B-type cyclins during oogenesis and meiotic maturation in Xenopus

laevis. *J Cell Biol*. doi:10.1083/jcb.114.4.755

Kosinki M, McDonald K, Schwartz J, Yamamoto I, Greenstein D. 2005. C. elegans sperm bud

vesicles to deliver a meiotic maturation signal to distant oocytes. *Development* **132**:3357–

3369. doi:10.1242/dev.01916

Langerak S, Trombley A, Patterson JR, Leroux D, Couch A, Wood MP, Schisa JA. 2019.

Remodeling of the endoplasmic reticulum in Caenorhabditis elegans oocytes is regulated

by CGH-1. *Genesis* **57**:1–8. doi:10.1002/dvg.23267

Lapierre LR, De Magalhaes Filho CD, McQuary PR, Chu CC, Visvikis O, Chang JT, Gelino S,

Ong B, Davis AE, Irazoqui JE, Dillin A, Hansen M. 2013. The TFEB orthologue HLH-30

regulates autophagy and modulates longevity in Caenorhabditis elegans. *Nat Commun* **4**.

doi:10.1038/ncomms3267

Lee SS, Lee RYN, Fraser AG, Kamath RS, Ahringer J, Ruvkun G. 2003. A systematic RNAi

screen identifies a critical role for mitochondria in C. elegans longevity. *Nat Genet*.

doi:10.1038/ng1056

Lehner B, Tischler J, Fraser AG. 2006. RNAi screens in Caenorhabditis elegans in a 96-well

liquid format and their application to the systematic identification of genetic interactions.

Nat Protoc **1**:1617–1620. doi:10.1038/nprot.2006.245

Llorca O, Martín-Benito J, Ritco-Vonsovici M, Grantham J, Hynes GM, Willison KR, Carrascosa

JL, Valpuesta JM. 2000. Eukaryotic chaperonin CCT stabilizes actin and tubulin folding

intermediates in open quasi-native conformations. *EMBO J* **19**:5971–5979.

doi:10.1093/emboj/19.22.5971

Luo S, Kleemann GA, Ashraf JM, Shaw WM, Murphy CT. 2010. TGF-β and Insulin Signaling

Regulate Reproductive Aging via Oocyte and Germline Quality Maintenance. *Cell*

143:299–312. doi:10.1016/j.cell.2010.09.013

Maller J, Wu M, Gerhart JC. 1977. Changes in protein phosphorylation accompanying

maturation of Xenopus laevis oocytes. *Dev Biol*. doi:10.1016/0012-1606(77)90093-8

Mccarter J, Bartlett B, Dang T, Schedl T. 1999. On the Control of Oocyte Meiotic Maturation and

Ovulation in Caenorhabditis elegans.

McDonald KL, Webb RI. 2011. Freeze substitution in 3 hours or less. *J Microsc* **243**:227–33.

doi:10.1111/j.1365-2818.2011.03526.x

Mehlmann LM, Terasaki M, Jaffe LA, Kline D. 1995. Reorganization of the endoplasmic

reticulum during meiotic maturation of the mouse oocyte.

Miller MA, Nguyen VQ, Lee MH, Kosinski M, Schedl T, Caprioli RM, Greenstein D. 2001. A

sperm cytoskeletal protein that signals oocyte meiotic maturation and ovulation. *Science*

(80-) **291**:2144–2147. doi:10.1126/science.1057586

Miller MA, Ruest PJ, Kosinski M, Hanks SK, Greenstein D. 2003. An Eph receptor sperm-

sensing control mechanism for oocyte meiotic maturation in Caenorhabditis elegans.

Genes Dev **17**:187–200. doi:10.1101/gad.1028303

O'Rourke EJ, Ruvkun G. 2013. MXL-3 and HLH-30 transcriptionally link lipolysis and autophagy

to nutrient availability. *Nat Cell Biol* **15**:668–676. doi:10.1038/ncb2741

Paix A, Folkmann A, Rasoloson D, Seydoux G. 2015. High efficiency, homology-directed

genome editing in Caenorhabditis elegans using CRISPR-Cas9ribonucleoprotein

complexes. *Genetics*. doi:10.1534/genetics.115.179382

Paix A, Schmidt H, Seydoux G. 2016. Cas9-assisted recombineering in C. elegans: Genome

editing using in vivo assembly of linear DNAs. *Nucleic Acids Res* **44**:e128.

doi:10.1093/nar/gkw502

Poteryaev D, Squirrell JM, Campbell JM, White JG, Spang A. 2005. Involvement of the Actin

Cytoskeleton and Homotypic Membrane Fusion in ER Dynamics in Caenorhabditis elegans

□ V. *Mol Biol Cell* **16**:2139–2153. doi:10.1091/mbc.E04

Qiao H, Rao HBDP, Yun Y, Sandhu S, Fong JH, Sapre M, Nguyen M, Tham A, Van BW, Chng

TYH, Lee A, Hunter N. 2018. Impeding DNA Break Repair Enables Oocyte Quality Control.

Mol Cell **72**:211-221.e3. doi:10.1016/j.molcel.2018.08.031

Reis-Rodrigues P, Czerwieniec G, Peters TW, Evani US, Alavez S, Gaman EA, Vantipalli M,

Mooney SD, Gibson BW, Lithgow GJ, Hughes RE. 2012. Proteomic analysis of age-

dependent changes in protein solubility identifies genes that modulate lifespan. *Aging Cell*.

doi:10.1111/j.1474-9726.2011.00765.x

Roux AE, Langhans K, Huynh W, Kenyon C. 2016. Reversible Age-Related Phenotypes

Induced during Larval Quiescence in C. Elegans. *Cell Metab*.

doi:10.1016/j.cmet.2016.05.024

Sahu R, Kaushik S, Clement CC, Cannizzo ES, Scharf B, Follenzi A, Potolicchio I, Nieves E,

Cuervo AM, Santambrogio L. 2011. Microautophagy of Cytosolic Proteins by Late

Endosomes. *Dev Cell*. doi:10.1016/j.devcel.2010.12.003

Schumacher B, Hofmann K, Boulton S, Gartner A. 2001. The C. elegans homolog of the p53

tumor suppressor is required for DNA damage-induced apoptosis. *Curr Biol*.

doi:10.1016/S0960-9822(01)00534-6

Shirayama M, Seth M, Lee HC, Gu W, Ishidate T, Conte D, Mello CC. 2012. PiRNAs initiate an

epigenetic memory of nonself RNA in the C. elegans germline. *Cell*.

doi:10.1016/j.cell.2012.06.015

Spike CA, Huelgas-Morales G, Tsukamoto T, Greenstein D. 2018. Multiple mechanisms

inactivate the LIN-41 RNA-binding protein to ensure a robust oocyte-to-embryo transition in

caenorhabditis elegans. *Genetics* **210**:1011–1037. doi:10.1534/genetics.118.301421

Sun Y, Li M, Zhao D, Li X, Yang C, Wang X. 2020. Lysosome activity is modulated by multiple

longevity pathways and is important for lifespan extension in C. elegans. *Elife*.

doi:10.7554/eLife.55745

Taylor RC, Dillin A. 2011. Aging as an event of proteostasis collapse. *Cold Spring Harb*

Perspect Biol. doi:10.1101/cshperspect.a004440

Ünal E, Kinde B, Amon A. 2011. Gametogenesis eliminates age-induced cellular damage and

resets life span in yeast. *Science (80-)* **332**:1554–1557. doi:10.1126/science.1204349

Walther DM, Kasturi P, Zheng M, Pinkert S, Vecchi G, Ciryam P, Morimoto RI, Dobson CM,

Vendruscolo M, Mann M, Hartl FU. 2015. Widespread proteome remodeling and

aggregation in aging C. elegans. *Cell*. doi:10.1016/j.cell.2015.03.032

Wasserman WJ, Richter JD, Smith LD. 1982. Protein synthesis during maturation promoting

factor- and progesterone-induced maturation in Xenopus oocytes. *Dev Biol*.

doi:10.1016/0012-1606(82)90303-7

Williams RL, Urbé S. 2007. The emerging shape of the ESCRT machinery. *Nat Rev Mol Cell*

Biol. doi:10.1038/nrm2162

Wright JE, Gaidatzis D, Senften M, Farley BM, Westhof E, Ryder SP, Ciosk R. 2011. A

quantitative RNA code for mRNA target selection by the germline fate determinant GLD-1.

EMBO J **30**:533–545. doi:10.1038/emboj.2010.334

Yamashita YM. 2018. Subcellular specialization and organelle behavior in germ cells. *Genetics*

208:19–51. doi:10.1534/genetics.117.300184

Zhang H, Alsaleh G, Feltham J, Sun Y, Napolitano G, Riffelmacher T, Charles P, Frau L, Hublitz

P, Yu Z, Mohammed S, Ballabio A, Balabanov S, Mellor J, Simon AK. 2019. Polyamines

Control eIF5A Hypusination, TFEB Translation, and Autophagy to Reverse B Cell

Senescence. *Mol Cell* **76**:110-125.e9. doi:10.1016/j.molcel.2019.08.005

Zou L, Wu D, Zang X, Wang Z, Wu Z, Chen D. 2019. Construction of a germline-specific RNAi

tool in C. elegans. *Sci Rep* **9**:1–10. doi:10.1038/s41598-019-38950-8

1040

Figure S1

A

Controls | Hits from screen | | Negative (non-hits)

Controls
- Empty vector — Negative control
- vha-13 RNAi — Positive control

Hits from screen

Strong
- rps-15 RNAi
- pat-10 RNAi

Intermediate
- cct-1 RNAi
- sar-1 RNAi

Subtle
- skr-2 RNAi
- par-5 RNAi

Negative (non-hits)
- ZK994.1 RNAi
- fos-1 RNAi

GFP::RHO-1

B

NMY-2::GFP

- Negative control — Empty vector
- vha-12 RNAi
- act-1 RNAi
- rab-11.1 RNAi
- atp-3 RNAi
- rpn-6.1 RNAi
- rpl-17 RNAi
- itr-1 RNAi

C

GO biological process	Fold enrichment	P-value
Translation	5.01	5.53E-08
ATP-hydrolysis coupled proton transport	27.17	3.76E-08
Ovulation	13.66	6.19E-05
RNA processing	3.26	3.22E-05
Cortical actin cytoskeleton organization	20.11	7.73E-04
Vacuolar acidification	24.58	4.77E-04

Figure S1.

(A) RNAi clones causing GFP::RHO-1 aggregates in hermaphrodite oocytes (SA115), visualized at low magnification (20X objective) during primary screening. Scale bars, 20 µm. (B) Examples of NMY-2::GFP aggregates in oocytes of hermaphrodites (JJ1473) undergoing orthogonal verification with a different aggregation-prone protein. Scale bars,10 µm. (C) The highest enriched gene-ontology (GO) process categories represented by the validated genes identified in the screen.

Figure S2

A — Hermaphrodites / Females

Adult hermaphrodite oocytes | Stacked female oocytes

B — Youngest female oocytes

C — Stacked female oocytes

mCherry::SP12   GFP::RHO-1   Merge

Figure S2.

(A) Comparison of ER morphology in oocytes from hermaphrodites (OCF15) and females (CF4542) expressing the ER reporter mCherry::SP12 in the germline. Oocytes in the female gonad stack over time, as new meiotic cells make their way down towards the spermatheca, but mature ones pass less readily into the uterus. Orange arrows, ER patches. (B) Visualization of ER patches in young oocytes of female animals, staged to capture the earliest oocytes. (C) Simultaneous visualization of GFP::RHO-1 aggregates and the ER in females (CF4557) at high magnification (100X objective). Orange arrows: ER patches; white arrows: GFP::RHO-1 aggregates. Scale bars (A-C),10 μm.

Figure S3

Hermaphrodites (Very first oocytes)

mCherry::SP12 GFP::RHO-1 Merge

Figure S3.

Extended data related to Figure 2B.

ER morphology and GFP::RHO-1 localization in the first, newly-formed oocytes from mCherry::SP12-expressing hermaphrodites (CF4552). Scale bars,10 µm. Note that the GFP::RHO-1 image from the 1st row is also displayed in figure 1E.

Figure S4

mCherry::SP12   GFP::RHO-1   Merge

Female; very early oocytes

Figure S4.

Extended data related to Figure 2B.

ER morphology and GFP::RHO-1 localization in the first, newly-formed oocytes from mCherry::SP12 expressing females (CF4557). Orange arrows: ER patches; white arrows: GFP::RHO-1 aggregates. Scale bars, 10 μm. Note that the GFP::RHO-1 image from the 6th row is also displayed in figure 1E.

Figure S5

Females

A

B

Hermaphrodite

C

Figure S5.

Extended data related to Figure 2D.

Transmission electron micrographs from hermaphrodite (N2E) and female (CF4101) oocytes. (A) Female oocyte. Arrow, ER stacks. Left panel, magnification: 1,200X, scale bar, 2 μm. Right panel, magnification: 13,000X, scale bar, 100 nm. (B) Female oocyte. Arrow, ER. Left panel, magnification: 890X, scale bar, 2 μm. Right panel, magnification: 6,800X, scale bar, 200 nm. (C) Hermaphrodite oocyte. Arrow, ER. Left panel, magnification: 1,400X, scale bar, 1 μm. Right panel, magnification: 13,000X, scale bar, 100 nm.

Figure S6A

Figure S6B

Figure S6.

(A) Simultaneous visualization of the alteration in ER morphology and clearance of aggregates in female oocytes (CF4560) as a function of time, following mating. The earliest changes in the mCherry::SP12 signal (ER) were observed ~10 minutes after mating, whereas the first signs of aggregate (NMY-2::GFP) clearance appeared later, between 20 and 30 minutes. The ER morphology was significantly altered by 30 minutes. Orange arrows: Initial ER patches altered at ~10 minutes. The oocytes are numbered according to their initial positions relative to the spermatheca prior to mating, and the numbers are maintained as the oocytes enter the uterus following fertilization. The figure shows a single representative animal from 7 independent experiments. Scale bars, 10 µm. (B) The mCherry::SP12 and NMY-2::GFP signals in oocytes of CF4560 females visualized as in (A) but not subjected to mating, to control for photobleaching. A single individual is shown from a total of 4 independent experiments. Scale bars,10 µm. (C) Reduction in signal intensity of the two fluorophores with time due to photobleaching, without exposure to males (from Fig. S6B). The signal intensities from specified areas (white rectangles) of the most proximal oocyte were assessed for this purpose.

Figure S7

A

pre-mating

Test female (to be mated) — Control female (not mated)

Skewness of NMY-2::GFP signal

Oocyte number

30 min post-mating

Mated female — Control female (not mated)

NMY-2::GFP aggregates cleared

Oocyte number

50 min post-mating

Mated female — Control female (not mated)

NMY-2::GFP aggregates cleared

Oocyte number

B

mCherry::SP12

Empty vector / Negative control	*hsp-4* RNAi	*atp-2* RNAi	*vha-2* RNAi

Normal ER morphology

cct-5 RNAi	*pbs-7* RNAi	*rps-20* RNAi	*ess-2* RNAi

act-1 RNAi	*sar-1* RNAi	*itr-1* RNAi	*dlg-1* RNAi

Figure S7.

(A) NMY-2::GFP signal skewness as a function of time in the proximal oocytes of CF4560 females, either subjected to mating with CB1490 males (white), or not exposed to males (pink). The mated females exhibit a reduction in signal skewness with time, and in relation to the distance from spermatheca, indicative of aggregate clearance. (B) Examples of oocyte ER morphology in CF4552 hermaphrodites subjected to assay-pool RNAi treatment, demonstrating the range of morphologies we observed. GFP::RHO-1 aggregation in the same animals was used as a control to verify successful gene knockdown (not shown). Scale bars, 20 μm.

Figure S8

A wrmScarlet::VHA-4

B VHA-7::wrmScarlet

C VHA-11::GFP

D wrmScarlet::VHA-13

E GFP::VHA-13

F *Ppie-1::VHA-7::wrmScarlet::vha-7* 3'UTR

Figure S8.

Expression of endogenously tagged V-ATPase subunits from the V_0 domain (A and B) and the V_1 domain (C, D and E), and of a single-copy integrated germline transgene (F).

None of the strains exhibited detectable signal in the gonad. (A) wrmScarlet::VHA-4, PHX1363 *vha-4(syb1363[wrmScarlet::vha-4]/+) II*, (B) VHA-7::wrmScarlet, PHX1317 *vha-7(syb1317[vha-7::wrmScarlet]) IV*, (C) VHA-11::GFP, PHX1198 *ieSi65[sun-1p::TIR1::sun-1 3'UTR + Cbr-unc-119(+)] II; unc-119(ed3) III; vha-11(syb1198[vha-11::gfp::degron]/+) IV*, (D) wrmScarlet::VHA-13, PHX731 *vha-13(syb731[wrmScarlet::vha-13]) V*, (E) GFP::VHA-13, PHX1049 *vha-13(syb1049[gfp::vha-13]) V,* (F) *Ppie-1::vha-7::wrmScarlet::vha-7* 3'UTR, PHX2690 *sybIs2690[Ppie-1::vha-7::wrmScarlet::vha-7 3'UTR, Cbr-unc-119(+)] I; unc-119(ed3) III.* Solid line; gonad and dotted line; intestine. Scale bars, 50 µm. Strains were generated by SunyBiotech for the Kenyon lab.

Figure S9

A

germline

intestine

intestine

germline

germline

B

Fertile adults (%)

empty vector   *vha-13* RNAi

Fertile adults (%)

empty vector   *vha-12* RNAi

- ■ N2E (RNAi permissive)
- □ DCL565 (RNAi inhibited globally)
- ▨ DCL569 (Germline-only RNAi)

C

GFP::VHA-13

Hermaphrodite

Female

0   25   50   75   100

Concentration of GFP RNAi (%)

D

GFP::VHA-13

empty vector   *vha-12* RNAi

Figure S9.

(A) Attempts to visualize tagged endogenous *wrmScarlet::vha-13* expression in the germline of PHX731 animals by dissecting gonads, to distance them from somatic tissues where it is highly expressed. Scale bars, 25 µm. (B) Percentage of fertile adults in N2E, DCL565 *rde-1(mkc36)*, and DCL569 *rde-1(mkc36); Psun-1::rde-1(+)* strains upon exposure to empty vector RNAi control, *vha-13* RNAi or *vha-12* RNAi, from hatching. Results are averaged from 3 biological replicates with each n>10. (C) Reduction of GFP::VHA-13 expression by serial dilution of *gfp* RNAi bacteria. Even at the limit of detection, hermaphrodites (PHX1414) exhibit discrete GFP::VHA-13 foci, whereas females (CF4599) exhibit a diffuse distribution. Scale bars, 10 µm. (D) Expression of GFP::VHA-13 in the germline of hermaphrodites exposed to empty vector RNAi (left) or *vha-12* RNAi (right). Scale bars, 50 µm.

Figure S10

A   GFP::VHA-13   Hermaphrodites

B   GFP::VHA-13   Females

Figure S10.

Extended data related to Figure 3A.

Additional images showing the distribution of GFP::VHA-13 in proximal oocytes of (A) hermaphrodites (PHX1414) and (B) females (CF4599). Scale bars, 10 μm.

Figure S11

A

Hermaphrodites

GFP::VHA-13

LysoTracker

Merge

B

Females

GFP::VHA-13

LysoTracker

Merge

Figure S11.

Extended data related to Figure 3B.

Additional images of the LysoTracker-stained oocytes from GFP::VHA-13-expressing (A) hermaphrodites (PHX1414) and (B) females (CF4599), for visualization of acidic lysosomes. The two signals colocalize in hermaphrodite oocytes. Dotted line; intestinal boundary, and solid line; oocytes. Scale bars, 10 μm

Figure S12

Figure S12.

(A) Analysis of germline LysoTracker staining in hermaphrodites (SA115) subjected to RNAi for the orphan genes. The LysoTracker signal is represented as the ratio of proximal oocyte to distal germline staining, and the values obtained from hermaphrodites (N2E) and females (CF4101) were used as controls (~1.3 and ~1.0, respectively). Each data point represents the average of values from 3 different locations in the most proximal oocyte of a single animal. The presence of GFP::RHO-1 aggregates in the same animals was used as a control to determine successful gene knockdown (not shown). (B) Analysis of GFP::VHA-13 localization (in terms of signal skewness) in hermaphrodites (PHX1414) subjected to RNAi for the orphan genes, and compared to hermaphrodite and female animals (PHX1414 and CF4599). Each data point represents the average of 10 skewness values from independent regions within the first three oocytes of a single animal. The presence of GFP::RHO-1 aggregates in SA115 animals in a parallel experiment served as a control for successful gene knockdown (not shown). For (A) and (B), unpaired t-test with Welch's correction was used to determine the statistical significance. The top and bottom rows of significance indicators in each experiment compare a specific RNAi condition to hermaphrodite and female animals respectively. P values: $P > 0.05$ = n.s., $P \leq 0.05$ = *, $P \leq 0.01$ = **, $P \leq 0.001$ = *** and $P \leq 0.0001$ = ****. (C) Visualization of GFP::VHA-13 distribution and localization in PHX1414 hermaphrodites following RNAi. A few representative gene knockdowns from the assay pool are shown here to demonstrate the range of localization phenotypes observed. Scale bars, 10 μm.

Figure S13

Figure S13.

(A) Analysis of germline DiOC6(3) staining in hermaphrodite animals (OCF15) subjected to RNAi for the orphan genes, represented as the ratio of proximal oocyte to distal germline staining. Staining ratios obtained from N2E hermaphrodites (~0.8) and CF4101 females (~1) were used as controls for comparison. Each data point represents the average of values from 3 different locations within the most proximal oocyte of a single animal. When applicable, alteration in ER morphology was used as a control to determine successful gene knockdown in the same animals. The presence of aggregates in GFP::RHO-1 animals (SA115) treated in parallel was also used as a control to verify successful gene knockdown (not shown). The top and bottom rows of significance indicators compare a specific RNAi treatment to hermaphrodites and females, respectively. (B) DiOC6(3) staining expressed as the ratio between proximal oocytes and distal germline, in hermaphrodites (N2E; white) and females (CF4101; yellow) subjected to RNAi treatment for control (*vha-12* and *atp*-2) and electron transport chain (*cyc-1* and *nuo*-2) genes. Successful gene knockdown was determined by monitoring GFP::RHO-1 aggregation in parallel (shown in Figure 5B). (A-B) Unpaired t-test with Welch's correction was used to determine the statistical significance. P values: $P > 0.05$ = n.s., $P \leq 0.05$ = *, $P \leq 0.01$ = **, $P \leq 0.001$ = *** and $P \leq 0.0001$ = ****. (C) For the animals in B, DiOC6(3) signal intensity following RNAi treatment for electron transport chain genes in the distal (left panel) and proximal (right panel) germlines, respectively. *vha-12*-RNAi was used an independent positive control. (D) Identification of Krebs cycle genes that, when knocked down in hermaphrodites (SA115), cause GFP::RHO-1 aggregates to accumulate in oocytes. Scale bars, 10 μm.

Figure S14

Figure S14.

(A) GFP::RHO-1 aggregation and LysoTracker signal in hermaphrodites (SA115) subjected to ESCRT-complex RNAi. The proximal oocyte/distal germline ratio of LysoTracker intensity is indicated for each animal. Distal germline images from the same animal are shown as insets in the LysoTracker panels. LysoTracker staining ratios ranged from 1.18-1.45 and 0.85-1.15 in control hermaphrodites (N2E) and females (CF4101), respectively, across multiple experiments (not shown in this figure). Oocyte GFP::RHO-1 aggregates (confirming knockdown) are indicated with white arrows, and only oocytes containing aggregates were selected for calculating lysosome acidification in Figure 5C. Scale bars, 10 µm. (B) GFP::VHA-13 localization (signal skewness) in hermaphrodites (PHX1414) subjected to RNAi for ESCRT subunits, and compared to hermaphrodites and females (PHX1414 and CF4599 respectively). Each data point represents the average of ten skewness values from different regions within the three most proximal oocytes of a single animal. The top and bottom rows of significance indicators compare a specific RNAi treatment to hermaphrodite and female animals respectively. P values: $P > 0.05$ = n.s., $P \leq 0.001$ = *** and $P \leq 0.0001$ = ****. (C) Variability in GFP::VHA-13 localization and lysosome acidification phenotypes for individual PHX1414 animals treated with *vps-54* RNAi. The skewness value for the GFP::VHA-13 signal and LysoTracker staining ratio (proximal oocyte to distal germline) are indicated for each animal. Successful knockdown could not be verified in each animal (since VHA-13 and RHO-1 were both tagged with GFP). Instead, it was done in a parallel experiment with GFP::RHO-1-expressing animals (SA115) in which more than 75% of animals subjected to a specific RNAi treatment, exhibited detectable aggregation (not shown). Scale bars, 10 µm.

Figure S15

Figure S15.

Extended data related to Figure 6B.

The same images of KIN-19 aggregates shown in figure 6B were scored again, blindly, by three additional investigators (A, B and C).

Figure S16

Figure S16.

Lysosome acidification in intestines.

LysoTracker staining in the intestines of animals subjected to gene knockdowns that caused KIN-19 aggregation in the head. Due to variability in dye uptake and staining, the actual LysoTracker signal intensities within the intestines were not evaluated. Instead, the mean skewness values, s, indicative of inhomogeneity expected for acidic intestinal lysosomes (so-called "lysosome-related organelles"), are reported for each animal (compared in Figure 6E). Skewness values were derived from six different locations within the anterior intestinal region near the pharyngeal bulb. Scale bar, 50 μm.

Table S1. Germline proteostasis candidates captured in the screen.

List of candidates obtained from primary screening and validation rounds with GFP::RHO-1. The phenotypic strengths and fertility status are indicated for each. Candidates that were discarded on the basis of their known functions, possible off-target effects, or the failure to pass orthogonal verification with NMY-2::GFP are indicated accordingly.

Table S2. List of *C. elegans* strains

Description of all strains used in the study, including the corresponding genotypes and sources.

Table S3. Candidates from primary screen that did not pass validation

This list is included because some of these candidates could be false negatives, as was the case for the ESCRT-complex genes. Also included are the GO terms (biological process) they represent.