The actin nucleator Spir-1 is a virus restriction factor that promotes IRF3 activation

Alice Abreu Torres[1], Stephanie L. Macilwee[¶1], Amir Rashid[1], Sarah E. Cox[1], Jonas D. Albarnaz[1],

Claudio A. Bonjardim[2] & Geoffrey L. Smith[1*]

[1]Department of Pathology, University of Cambridge, Tennis Court Road, CB1 2QP, Cambridge,

United Kingdom

[2]Laboratório de Vírus, Departamento de Microbiologia, Instituto de Ciências Biológicas, Universidade

Federal de Minas Gerais, Belo Horizonte, MG, Brazil.

[¶]Current address: Algal Omega 3, Penrhyn Rd, Knowsley, Prescot L34 9HY

[*]Corresponding author: email, gls37@cam.ac.uk

Short title: Spir-1 is a virus restriction factor

Key words: Spir-1, virus restriction factor, vaccinia virus, Zika virus, innate immunity, IRF3 activation,

protein K7, immune evasion.

Abstract

Cellular proteins often have multiple and diverse functions. This is illustrated with protein Spir-1 that is an actin nucleator, but, as shown here, also functions to enhance IRF3 activation downstream of RNA sensing by RIG-I/MDA-5. In human and mouse cells lacking Spir-1, IRF3 activation is impaired, whereas Spir-1 overexpression enhanced IRF3 activation. Furthermore in Spir-1$^{-/-}$ cells, the infectious virus titres and sizes of plaques formed by two viruses that are sensed by RIG-I, vaccinia virus (VACV) and Zika virus, are increased. These observations demonstrate the biological importance of Spir-1 in the response to virus infection. Like cellular proteins, viral proteins also have multiple and diverse functions. Here, we also show that VACV virulence factor K7 binds directly to Spir-1 and that a diphenylalanine motif of Spir-1 is needed for this interaction and for Spir-1-mediated enhancement of IRF3 activation. Thus, Spir-1 is a new virus restriction factor and is targeted directly by an immunomodulatory viral protein that enhances virus virulence and diminishes IRF3 activation.

Author Summary

Infection of cells by viruses is sensed by host molecules called pattern recognition receptors (PRRs) that activate signalling pathways leading to an anti-viral response. In turn, viruses express proteins that negate these host responses to mediate escape from the anti-viral response. Here, we report that protein K7 from a large DNA virus called vaccinia virus (VACV), binds to a host cell protein called Spir-1. Spir-1 is known to regulate the assembly of actin filaments inside cells, but here we show that Spir-1 also functions to activate the host response to virus infection and to limit the replication and spread of both RNA and DNA viruses. Thus, this study has uncovered new functions of cellular protein Spir-1 as an activator of innate immunity and as a restriction factor for diverse viruses. Further, it shows that Spir-1 is targeted by a virus protein during infection.

Introduction

The host innate immune response to viral infection begins with the sensing of pathogen-associated molecular patterns (PAMPs) by pattern recognition receptors (PRRs), such as Toll-like receptors and retinoic acid-inducible gene-I (RIG-I)-like receptors (RLRs) (1, 2). The sensing of virus macromolecules, such as viral DNA or RNA, by PRRs triggers signalling cascades that culminate in the activation of the transcriptional factors interferon regulatory factor (IRF) 3, activator protein (AP) 1 and nuclear factor kappa B (NF-κB) (3). These transcriptional factors translocate to the nucleus where they induce the transcription of genes encoding interferons (IFN), cytokines and chemokines. Once secreted from the cell, IFNs, cytokines and chemokines promote inflammation to restrict virus replication and control the infection. IFNs bind to their receptors on the surface of infected or non-infected cells, and trigger signal transduction via the JAK-STAT pathway leading to expression of interferon-stimulated genes (ISGs) that induce an antiviral state (4).

During the co-evolution with their hosts, viruses have evolved strategies to evade, suppress or exploit the host response to infection by targeting multiple steps of the host immune response (5). Vaccinia virus (VACV), the prototypical orthopoxvirus, is well known as the live vaccine used to eradicate smallpox (6). VACV has a large dsDNA genome of approximately 190 kbp (7), replicates in the cytoplasm (8) and encodes scores of proteins that antagonise innate immunity (9). Interestingly, some cellular pathways, such as those leading to activation of IRF3 or NF-κB, or the JAK-STAT pathway downstream of IFNs binding to their receptors, are targeted by multiple different VACV proteins. Moreover, some of these viral antagonists are multifunctional and inhibit more than one host innate immune pathway (9).

VACV protein K7 is one such antagonist of innate immunity. K7 is a small, intracellular protein that is non-essential for virus replication in cell culture, yet contributes to virulence in both intradermal and intranasal mouse models of infection (10). Functionally, K7 was reported to suppress NF-κB activation by binding to interleukin-1 receptor-associated kinase-like 2 (IRAK2) and tumour necrosis factor (TNF) receptor-associated factor 6 (TRAF6) (11). K7 also inhibits IRF3 activation and binds to the DEAD-box RNA helicase 3 (DDX3) (11). Another study reported that K7 affected regulation of histone methylation during VACV infection, by an unknown mechanism (12). Two unbiased proteomic searches identified cellular binding partners of K7 (13, 14), suggesting that K7 may have additional functions. The identification of cellular proteins targeted by viral proteins has been a useful approach to identify cellular factors that function in the recognition and restriction of virus infections (15, 16). In this study, by investigating the interaction between K7 and the cellular protein Spir-1 (14), we identified new functions for Spir-1 as an activator of innate immunity and as a restriction factor for both DNA and RNA viruses.

The protein spire homolog 1 (Spir-1, also known as SPIRE1) was first described to affect *Drosophila* embryogenesis (17). Spir-1 has actin-binding domains (18) through which it nucleates actin filaments, an activity shared with the Arp2/3 complex and the formins (19). Spir-1 is organised in multiple functional domains. The N-terminal region contains the kinase non-catalytic C-lobe domain (KIND), which mediates Spir-1 interaction with other proteins, such as the formins (20). Spir-1 and the formins cooperate during actin nucleation (21-29). The KIND domain is followed by four actin-binding Wiskott-Andrich syndrome protein homology domain 2 (WH2) domains that are responsible for actin nucleation (19, 30, 31). The C-terminal region of Spir-1 contains a globular tail domain-binding motif (GTBM), which is responsible for binding to myosin V (32). Next, there is a Spir-box (SB) domain that is conserved within the Spir protein family. Due to its similarity to a helical region of the rabphilin-3A protein that interacts with the GTPase Rab3A, it is thought that the Spir-box domain is involved in the association of Spir-1 and Rab-GTPases (24, 33-35). Following the SB domain, there is a modified

FYVE zinc finger domain that interacts with negatively-charged lipids in membranes. The membrane

targeting specificity is mediated via the interaction of Spir-1 with other membrane-bound proteins such

as the Rab-GTPases (29).

In adult mice, Spir-1 is expressed preferentially in neuronal and hematopoietic cells (36, 37) and in

humans the brain also has high Spir-1 expression (38). In general, Spir-1 is involved in several actin-

dependent cellular functions such as vesicle trafficking (24, 34, 35, 39), DNA repair (40), mitochondrial

division (41), and development of germ cells (17, 23, 42) but a role in innate immunity has not been

described. Interestingly, a genome-wide association study found a single nucleotide polymorphism in

Spir-1 that correlated with a different antibody response to smallpox vaccination (43).

Here, Spir-1 is shown to promote IRF3 activation and this activity depends upon a diphenylalanine

motif that is also necessary for the direct interaction of Spir-1 with VACV virulence factor K7. Using

gain-of-function and loss-of-function cell lines, Spir-1 is shown to diminish VACV and ZIKV replication

and/or spread and is therefore a viral restriction factor.

Results

Vaccinia virus protein K7 co-precipitates with the C terminal region of Spir-1

A previous proteomic study identified Spir-1 as a cellular interacting partner of VACV protein K7

although this interaction was not validated (14). Mammals have two *Spire* genes, *Spire1* and *Spire2*

that encode closely related proteins (overall 42% identity and 58% similarity in humans), especially

within the WH2 and Spir-box domains (37, 44). The Uniprot database for human Spir-1 (Q08AE8)

describes five Spir-1 isoforms and splicing before the GTBM (Exon 9) and SB (Exon 13) domains has

been demonstrated (45). Among Spir-1 isoforms, Spir-1 known as isoform 2 that doesn't contain Exon

9 or Exon 13, is the most abundant form in the brain and small intestine tissues (45) and is the form

studied here. To confirm if K7 co-precipitates with Spir-1, HEK293T cells were transfected with

plasmids encoding Myc-tagged human Spir-1, Spir-2, β-TrCP, or Myc-GFP, together with plasmids

expressing FLAG-tagged, codon-optimised K7 or another Bcl-2-like VACV protein, A49 (46).

Immunoprecipitation with anti-Myc or anti-FLAG affinity resins showed that K7 was co-precipitated by

Myc-Spir-1 (Fig 1A), but not by Myc-Spir-2, Myc-β-TrCP or Myc-GFP. In contrast, A49 interacted with

β-TrCP as reported (47), but not with the other proteins. Reciprocally, FLAG-K7 co-precipitated Spir-

1 (Fig 1B), but not the other Myc-tagged proteins, whilst β-TrCP only interacted with A49.

To investigate if the actin-binding domains of Spir-1 were needed for interaction with K7, two Myc-
tagged truncations of Spir-1 were generated: an N-terminal region (amino acid residues 1-390),
containing the four actin-binding WH2 domains and the KIND domain; and a C-terminal region,
containing the SB and FYVE domains (amino acid residues 391-742, Fig 1C). These truncations,
Spir-1 and Spir-2 were co-expressed with FLAG-K7 and Myc-tagged immunoprecipitation showed
that the C terminus of Spir-1 was sufficient for interaction with K7, whereas the actin-binding domains
were dispensable (Fig 1D, middle panel). The reciprocal IP gave the same conclusion and FLAG-K7
co-precipitated full-length Spir-1 and its C-terminal region (Fig 1D, right panel). As controls, the N-
terminal region co-precipitated endogenous actin via its WH2 domains, and the C-terminal region
interacted with 14-3-3, another binding partner of Spir-1 (48, 49).

To determine if Spir-1 and K7 interacted at endogenous levels, cells were either mock-infected or
infected with VACV expressing HA-tagged K7 (vHA-K7) (10) or B14 (vHA-B14) (50). HA-
immunoprecipitation confirmed that K7 interacts with endogenous Spir-1, whilst B14 does not (Fig
1E). As reported, K7 also co-precipitated with DDX3 (11), and B14 interacted with IKKβ (51).
Altogether, these results indicate that Spir-1, but not Spir-2, interacts with K7 via its C-terminal region
and independent of its actin-binding domains.

Ectopic Spir-1 increases IRF3-dependent gene expression
Since the interaction of Spir-1 with K7 is independent of its actin-binding domains, and K7 is a VACV
immunomodulator and virulence factor, we hypothesised that Spir-1 might have an unknown function
in antiviral immunity. To test this, the impact of Spir-1 on innate immune signalling pathways was
assessed by luciferase reporter gene assays. First, cells were transfected with a reporter plasmid in
which the expression of firefly luciferase is driven by the IFNβ promoter. IFNβ expression was induced
by Sendai virus (SeV) infection, which is sensed by RIG-I (52). Myc-Spir-1 expression alone did not
affect IFNβ-dependent gene activation, however, Spir-1 augmented activation induced by SeV
infection (Fig 2A). In contrast, Myc-tagged GFP had no effect, and VACV protein C6 was inhibitory as
described (53). Spir-1 did not affect activation of ISRE-dependent gene expression downstream of
type I IFN, whereas C6, but not N1, was inhibitory (54) (Fig 2B). Similarly, Spir-1 did not affect NF-κB
activation in response to TNF-α, whilst B14 inhibited it (51) and VACV protein N2 had no effect (55)
(Fig 2C). Further analysis using an IRF3-specific reporter plasmid (*ISG56.1* or *IFIT1* promoter)
showed Spir-1 caused a dose-dependent increase in IRF3 activation induced by the CARD domain
of RIG-I (Fig 2D, 2E). As controls, VACV protein N2 inhibited IRF3 activation as described (55) while
A49, an NF-κB inhibitor (47), did not. Unlike Spir-1, Spir-2 did not enhance IRF3 activation whereas
DDX3 did, as described (11) (Fig 2E).

To map at which stage in the IRF3 pathway Spir-1 was acting, further *ISG56.1* reporter assays were

performed in which the pathway was stimulated by expression of its different components. Spir-1

activated the IRF3 pathway in a dose-response manner when MAVS (mitochondrial antiviral-

signalling protein) was used as stimulant (Fig 2F), but not when downstream components such as

TBK1 (Fig 2G), IKKε (S1 Fig) or the constitutively active IRF3-5D (Fig 2H) were expressed. These

findings indicate Spir-1 enhances IRF3 activation at or downstream of MAVS and upstream of IKKε

or TBK1. Finally, N- and C-terminal fragments of Spir-1 were tested following stimulation by RIG-I

CARD. Neither half of Spir-1 activated the pathway fully, but the C-terminal region had greater activity

than the N-terminal region (Fig 2I).

Spir-1 interaction with K7 is direct, independent of DDX3, and requires a diphenylalanine motif

K7 interacts directly with DDX3 (56, 57), an adaptor protein in the IRF3 pathway (58, 59). To

investigate whether K7 binding to Spir-1 was via DDX3, cells stably expressing an inducible shRNA

targeting DDX3 (shDDX3) (59) were used. Knockdown, rather than knockout, of DDX3 was utilised

because *DDX3* encodes an essential protein (60). Cells expressing a non-silencing control (NSC)

shRNA were used in parallel. DDX3 knockdown was induced by incubation with doxycycline (DOX)

for 48 h as described (59), prior to transfection with either FLAG-tagged K7 or A49 for 24 h. Viral

proteins were immunoprecipitated via their FLAG tag, followed by immunoblotting for endogenous

Spir-1. DDX3 levels were reduced greatly in shDDX3 cells following DOX treatment, but not in control

cells (Fig 3A). Despite this, Spir-1 co-precipitation by K7 was unaffected (Fig 3A). A49 did not co-

precipitate either DDX3 or Spir-1 (Fig 3A). Next, the shDDX3 cells were used to determine if DDX3

contributed to Spir-1-induced IRF3 activation. After DDX3 knockdown, there was no significant

difference in IRF3 stimulation by Spir-1 (Fig 3B). Notably, K7 inhibited IRF3 activation when DDX3

was knocked down, showing K7 had another IRF3 inhibitory mechanism independent of DDX3 (Fig

3B). Moreover, the presence of K7 reversed the activation of the pathway by Spir-1 (Fig 3B). In

summary, Spir-1 interaction with K7 and its function in the IRF3 pathway are independent of DDX3.

Since both DDX3 and Spir-1 bind to K7 and activate IRF3, their amino acid sequences were

compared. The structure of K7 bound to a DDX3 peptide and subsequent structure-based

mutagenesis showed that a diphenylalanine (FF) motif in DDX3 is essential for binding K7 and for

IRF3 activation (57). An alignment of the C terminus of Spir-1 and the minimal ten-amino acid peptide

from DDX3 (residues 81-90) needed for binding K7, showed a similar FF motif in Spir-1 (Fig 3C). To

determine if this was important for Spir-1 binding to K7, the phenylalanines were mutated to alanines

(FFAA) and the mutant was expressed in cells together with FLAG-tagged K7. Wild-type (WT) Myc-

Spir-1 and Myc-GFP were also expressed. Anti-Myc or anti-FLAG immunoprecipitation showed that

the FFAA mutation impaired Spir-1 interaction with K7 (Fig 3D), although it still precipitated both actin

and 14-3-3, indicating normal folding. Notably, the FFAA mutant no longer activated IRF3 (Fig 3E).

To determine if the K7 interaction with Spir-1 was direct, these proteins were expressed by *in vitro*

transcription and translation and immunoprecipitated alongside Myc-tagged TAB2 and FLAG-tagged

VACV protein F12 (61) as controls. Both Spir-1 and DDX3 interacted directly with K7, whereas Spir-

1 did not interact with F12, and K7 did not interact with TAB2 (Fig 3F, 3G). Taken together, this

demonstrated that Spir-1 binds directly to K7 and shares with DDX3 a conserved diphenylalanine

motif, which is required for its function in the IRF3 pathway.

K7 uses the same amino acid residues to target both Spir-1 and DDX3

K7 binds to the DDX3 peptide through a hydrophobic pocket within a negatively-charged face (56,

57). Thus, interactions between K7 and the DDX3 peptide involve hydrophobic contacts, hydrogen

bonds and electrostatic interactions, of which electrostatic contacts between R88 of DDX3 with both

D28 and D31 of K7 are notable. D31 also forms a hydrogen bond with DDX3 S83 (57). Using this

information, K7 mutants were generated in an attempt to distinguish between K7 binding to Spir-1

and DDX3. K7 D28 and D31 were changed singly or together to alanine (D28A, D31A) and expressed

alongside WT K7 or GFP, and together with Myc-tagged Spir-1 and HA-tagged DDX3.

Immunoprecipitation showed that K7 D28A still bound both Spir-1 and DDX3, whilst K7 D31A had

impaired binding to both proteins, especially DDX3 (Fig 4A). Furthermore, the double mutant (DDAA)

had further reduced binding to Spir-1 and no detectable binding to DDX3 (Fig 4A). Importantly, all

mutants still interacted with COPε (Fig 4A), another K7 binding partner (13), suggesting normal K7

folding. Co-precipitation of endogenous Spir-1 with FLAG-tagged K7 and mutants gave similar

conclusions (Fig 4B). Notably, the K7 D31A mutant inhibited IRF3 activation poorly compared to the

WT or D28A (Fig 4C). Altogether, these data indicate that K7 requires D31 for interacting with both

DDX3 and Spir-1 and via each interaction it inhibits the IRF3 pathway.

IRF3 activation is reduced in the absence of Spir-1

To investigate Spir-1-mediated activation of IRF3 further, Spir-1 knockout (KO) HEK293T cell lines

were generated by CRISPR-Cas9-mediated targeting of *SPIRE1* exon 3, which is conserved in all

Spir-1 isoforms (Fig 5A). After single cell selection, HEK293T cell lines were confirmed to lack Spir-1

by immunoblotting (Fig 5B) and by sequencing of both *Spire-1* alleles (Fig 5A). A clone that lacked

Spir-1 protein expression and in which the *Spire-1* open reading frame was disrupted by frameshift

mutations in both alleles, and that lacked WT sequence, was selected. Additionally, a vector

expressing Myc-tagged Spir-1 was used to rescue Spir-1 expression in the KO cell line following

lentiviral transduction. In parallel, both WT and KO cells were transduced with a control empty vector

(EV) lentivirus. Spir-1 WT cells transduced with EV, and Spir-1 KO cells transduced with EV or Myc-

Spir-1 were infected with SeV to activate the IRF3 pathway. Supernatants were collected for ELISA

and cells lysates were used for either RNA extraction or immunoblotting, which confirmed that cells
did or did not express Spir-1 (Fig 5B). The expression of Myc-Spir-1 increased in the presence of
SeV, but this was not observed for endogenous Spir-1 (Fig 5B). One possible explanation is that Myc-
Spir-1 expression is driven by the human cytomegalovirus immediate early promoter (62) that
contains binding sites for transcription factors such as NF-κB (63), which can also be activated during
SeV infection. The phosphorylation of IRF3 at S396 (p-IRF3), a hallmark of IRF3 activation (64),
increased greatly 24 h after SeV infection of WT cells (Fig 5B) but was reduced in the absence of
Spir-1 and restored in the KO cells expressing Myc-Spir-1 (Fig 5B). Moreover, after SEV infection,
Spir-1 KO cells produced lower levels of mRNAs for *ISG56/IFIT1* (Fig 5C), *IFNB1* (Fig 5D) and
CXCL10 (Fig 5E), and secreted less CXCL-10 (Fig 5F), as measured by RT-qPCR and ELISA,
respectively. Importantly, the absence of Spir-1 did not change the levels of CXCL-10 secreted in
response to TNF-α, a specific agonist of the NF-κB pathway (Fig 5G).

Immortalised mouse embryonic fibroblasts (MEFs) from mice with a terminator (gene trap) between
exons 3 and 4 of the *Spire1* gene (65) were used to investigate the contribution of Spir-1 to IRF3
activation in cells from a different mammal. After stimulation of IRF3 by either poly I:C or SeV infection,
MEFs were lysed for immunoblotting or RNA analysis and supernatants were collected for ELISA. A
reduction in p-IRF3 was seen in Spir-1 KO cells (Fig 6A, 6B) and this was quantified (bottom graphs
in Fig 6A, 6B). The Spir-1 KO MEFs also expressed less mRNAs for cytokines and chemokines upon
stimulation with poly I:C or SeV infection than their WT controls (Fig 6C, 6D, 6E, 6F, 6G, 6H). Likewise,
CXCL-10 and IL-6 secretion were lower from Spir-1 KO cells (Fig 6I, 6J, 6K, 6L). However, the levels
of CXCL-10 and IL-6 secreted after stimulation with IL-1β were not reduced in Spir-1 KO MEFs (Fig
6M, 6N); indeed, there was a small increase in IL-6 levels in KO cells (Fig 6N). In summary, in both
mouse and human cells lacking Spir-1 there is a defect in IRF3 activation downstream of RIG-I/MDA5,
whilst other immune signalling pathways were largely unaltered.

Spir-1 is a viral restriction factor
Given that Spir-1 interacts with VACV protein K7 and enhances IRF3 activation, which is critical for
the host response to viral infection, the impact of Spir-1 on VACV replication and spread was
assessed. Spir-1 WT and KO HEK293T cells and the derivative cell line in which Spir-1 expression
was restored in the KO cells, were infected with A5-GFP-VACV, a VACV strain expressing GFP fused
to capsid protein A5 (66), and 2 days (d) later plaque diameters were measured (Fig 7A, 7B). A
significant increase in plaque size was seen in the absence of Spir-1 compared to both WT and
rescued cells (Fig 7B), suggesting that Spir-1 restricts VACV spread and/or replication. Lastly, virus
growth on those cells was assessed. Cells were infected with 0.001 plaque-forming unit (PFU)/cell of
VACV strain WR and 2 d later the virus yield was determined by plaque assay. Spir-1 KO cells yielded

higher viral titres compared to both WT and Spir-1-complemented cells (Fig 7C).

The IRF3 pathway also restrict RNA viruses, therefore the impact of Spir-1 in restricting Zika virus

(ZIKV), a single-stranded RNA virus, was also assessed. HEK293T Spir-1 WT, KO and Spir-1-

rescued cells were infected with ZIKV-mCherry with 0.01 PFU/cell. Three days later, the mCherry

signal was visualised by live-cell imaging and showed a greater signal in the KO cells compared to

controls (Fig 7D), suggesting an increased ZIKV replication in these cells. To address that possibility,

the virus yield in the supernatant of infected HEK293T cells was determined by plaque assay in VERO

E6 cells (Fig 7E). In the absence of Spir-1, there is an increase in ZIKV titres compared to the WT

and Spir-1 restored KO cells. Collectively, loss of Spir-1 caused an increase in both virus plaque size

and yield of infectious virus, showing that Spir-1 functions as a restriction factor against both a DNA

(VACV) and an RNA (ZIKV) virus.

Discussion

During co-evolution with their hosts, viruses have evolved mechanisms to evade the host antiviral

responses and thereby replicate and spread efficiently. Such virus proteins are numerous and have

diverse functions and, therefore, can be used to gain insight into how the host innate immune system

functions. VACV is a poxvirus and encodes scores of immunomodulatory proteins many of which

target the innate immune system. In several cases, multiple VACV proteins engage the same

pathway, but despite this, these proteins each contributes to virulence indicating non-redundant

functions (9). Sometimes a single protein can inhibit multiple pathways (9) or an open reading frame

can encode multiple proteins with different functions (67). High-throughput proteomic approaches

have been used to identify binding partners of VACV proteins, or cellular proteins that are up or down-

regulated during virus infection, and such cellular proteins may have antiviral activity and function in

innate immunity (14, 16). For example, histone deacetylases (HDAC) 4 and 5 are each targeted for

proteasomal degradation by VACV protein C6 and function as a virus restriction factor (15, 16).

Here, Spir-1 is characterised as a new cellular protein that is bound by VACV protein K7 (Fig 1), a

virulence factor that inhibits NF-κB and IRF3 activation (10, 11). K7 binding to Spir-1 was identified in

a proteomic screen for binding partners of virus immunomodulatory proteins (14). Here, this

interaction is confirmed, demonstrated to occur at endogenous levels during VACV infection (Fig 1E),

and shown to be direct (Fig 3F, 3G).

Spir-1 belongs to a family of proteins involved in actin organisation (44) but K7 interaction with Spir-1

does not require its N-terminal actin-binding domain (Fig 1D). VACV induces polymerisation of actin

to facilitate virus dissemination from the cell surface and VACV mutants unable to polymerise actin

spread poorly and form small plaques (68). However, a mutant lacking K7 produces normal size

plaques (10). Given that K7 is an immunomodulatory protein, we hypothesised that Spir-1 might have

a function in innate immunity. In reporter gene assays, Spir-1 over-expression enhanced IRF3-

dependent gene expression (Fig 2D) but did not affect JAK-STAT signalling induced by type I IFN, or

NF-κB activation (Fig 2B, 2C). Conversely, human and murine Spir-1 knockout cell lines showed

diminished IRF3 signalling, with reduced IRF3 phosphorylation and reduced transcription and

expression of IRF3-responsive genes (Fig 5B, 6A, 6B). Importantly, rescue of Spir-1 expression in

Spir-1 KO cells restored phospho-IRF3 and cytokine expression levels (Fig 5).

To map where Spir-1 influences IRF3 activation, the pathway was activated by different agonists and

the influence of Spir-1 examined. Spir-1 enhanced activation induced by SeV infection (Fig 2A, 5 and

6), poly I:C transfection (Fig 6), and overexpression of both RIG-I CARD (Fig 2D, 2E) and MAVS (Fig

2F), but not overexpression of TBK1, IKKε or a constitutively active IRF3 (Fig 2G and S1). SeV RNA

is sensed by RIG-I (69) whilst high molecular weight (HMW) poly I:C is sensed by MDA-5 (52, 70).

This suggests Spir-1 acts at or downstream of MAVS, the adaptor molecule recruited downstream of

RIG-I and MDA5 activation (71). The association of RIG-I/MDA5 with MAVS results in the recruitment

of tumour necrosis factor receptor-associated factors (TRAF), such as TRAF3, leading to the

phosphorylation of IRF3 by TBK1/IKKε. Collectively, these data suggest Spir-1 functions at MAVS or

TRAF3 level.

K7 inhibition of IRF3 activation was attributed to direct binding to DDX3, which acts as a

multifunctional scaffolding adaptor culminating in IRF3 activation (58, 59). However, data presented

here show K7 has additional targets within the IRF3 pathway, since Spir-1 also contributes to IRF3

activation and K7 is able to inhibit IRF3 activation even in the absence of DDX3. Furthermore, DDX3

and Spir-1 act at different positions in the pathway; DDX3 acts at the TRAF3 and TBK1/IKKε level (9,

11, 59, 72), whilst Spir-1 acts upstream of TBK1/IKKε (Fig 2). Interestingly, both DDX3 and Spir-1

share a diphenylalanine motif that is responsible for both binding to K7 and their function in the IRF3

pathway (Fig 3) (57). Furthermore, the same amino acid residue in K7 (D31) is crucial for its binding

to both cellular proteins (Fig 4). This prevented assessment of the relative importance of K7 binding

to each cellular protein.

Nonetheless, Spir-1 has an important antiviral role because cells lacking Spir-1 show enhanced VACV

and ZIKV replication and/or spread (Fig 7). Comparable data for DDX3 are lacking, and obtaining

such data is complicated by DDX3 being an essential protein (60). Both K7 and Spir-1 localise in the

cytoplasm (10, 34). Interestingly, mutations in the FYVE domain and in the diphenylalanine motif in

Spir-1 change its sub-cellular localisation from a trans-Golgi network and post-Golgi vesicles

localisation to an even cytoplasmic distribution (29, 34). Neither the N- or C-terminal halves of Spir-1

were capable of activating the IRF3 pathway as well as full-length protein (Fig 2I), despite comparable

expression levels, suggesting that both halves are important. It is possible that Spir-1 is directed to

the correct location by its C-terminal domain, whist its N-terminal domain mediates Spir-1 engagement

with other proteins to promote IRF3 activation. This might explain why the FFAA Spir-1 mutant is no

longer able to activate the IRF3 pathway. Another possibility is that K7 may bind to the Spir-1

diphenylalanine motif and displace other proteins or sequester it and prevent it reaching its subcellular

location. Spir-1 has several isoforms (45), and one, Spire-1C (also known as Spire1-E13), differs from

the canonical isoform by encoding an extra exon sequence (ExonC or E13), which mediates its

mitochondrial localisation and regulates mitochondrial division (41). Even though this was not the

isoform used in the present study, K7 might regulate Spir-1 trafficking to the mitochondrion, where

MAVS is anchored (73).

RIG-I and MDA5 can sense RNA virus genomes directly, but also RNA generated by infection with

DNA viruses (74), leading to IRF3 activation. Therefore, RIG-I/MDA5-triggered antiviral response is

antagonised by several RNA and DNA viruses (71, 75). For instance, ZIKV has been shown to be

sensed by both RIG-I and MDA-5 depending on the infected cell type (reviewed by (76)). Furthermore,

it has long been known that dsRNA is generated during VACV infection (77-79), so it is not surprising

that VACV encodes proteins to interfere with cytosolic RNA sensing. For instance, VACV protein E3

binds to dsRNA and prevents RIG-I-dependent sensing of RNA products generated via the

transcription of AT-rich DNA by RNA polymerase III (80, 81), and the decapping enzymes D9 and

D10 prevent the accumulation of dsRNA from viral complementary RNA molecules (82). Downstream

of RNA sensing, VACV encodes proteins to inhibit IRF3 activation, such as C6, which acts at

TBK1/IKKε activation level (53), N2, which acts downstream of IRF3 translocation to the nucleus (55)

and K7, which targets DDX3 (11). Here, we show that Spir-1 also contributes to IRF3 activation, with

both Spir-1 and DDX3 sharing a similar motif which binds K7.

In summary, Spir-1 is a new cellular protein that affects IRF3 activation and is a virus restriction factor.

By exploring in detail the function of VACV protein K7, an additional cellular factor that regulates IRF3

activation and acts as a viral restriction factor has been characterised.

Material and Methods

Cells lines

Human embryonic kidney (HEK) 293T, HEK293T NSC or shDDX3 (kind gifts from Dr. Martina Schröder, Maynooth University, Ireland), murine embryonic fibroblasts (MEFs) Spir-1 KO and WT (kindly provided by Prof. Dr. Eugen Kerkhoff - University Hospital Regensburg, Germany), BSC-1 and VERO E6 (African Green Monkey) cells were cultured in Dulbecco's modified Eagle's medium (DMEM, Gibco) supplemented with 10% heat-treated (56 ºC, 1 h) foetal bovine serum (FBS, Pan Biotech), 100 U/mL penicillin and 100 μg/mL streptomycin (P/S, Gibco). RK13 cells were grown in minimal essential medium (MEM, Gibco) supplemented with 10% FBS and P/S.

Viruses

VACV strain Western Reserve (WR) and derivative strains expressing GFP fused to the capsid protein A5 (A5-GFP-VACV) (66), VACV expressing HA-tagged B14 (50) and VACV expressing HA-tagged K7 (10) were described. VACV strains were grown on RK13 cells and titrated by plaque assay on BSC-1 cells. Sendai virus (SeV) Cantell strain (Licence No. ITIMP17.0612A) was a gift from Prof. Steve Goodbourn (St George's Hospital Medical School, University of London). ZIKV engineered to express a mCherry marker (83) was a kind gift from Dr. Trevor Sweeney (Department of Pathology, University of Cambridge).

Plasmids

A plasmid encoding the Myc-tagged human Spir-1 isoform 2 and human Spir-2 (20) constructs were a kind gift of Prof. Dr. Eugen Kerkhoff. Spir-1 N- and C-terminal truncations were constructed by PCR amplification from pcDNA3.1-Myc-Spir-1 and inserted into plasmid pcDNA3.1-Myc. Codon-optimised FLAG-K7 was described (84) and Spir-1 and K7 plasmids were used as templates for site-directed mutagenesis according to the instructions of the QuikChange Site-Directed Mutagenesis Kit (Agilent). All mutations were confirmed by DNA sequencing. Myc-DDX3 and HA-DDX3 (11) were kindly provided by Dr. Martina Schröder. The IFNβ-firefly luciferase reporter plasmid was from T. Taniguchi (University of Tokyo, Japan), NF-κB-firefly luciferase was from R. Hofmeister (University of Regensburg, Germany) and ISG56.1-firefly luciferase was a gift from Ganes Sen (Cleveland Clinic, USA). ISRE-firefly luciferase and pTK-renilla luciferase (pRL-TK) plasmids were from Promega. Vectors expressing MAVS, IKKε, TBK1 and IRF3-5D (53), and RIG-I-CARD domain (85) were described. Lentivirus vector plasmid pLKO.DCMV.TetO.mcs (pLDT) (86) was a gift from Prof. Roger Everett (University of Glasgow, UK) and was used as backbone for sub-cloning the Myc-Spir-1. Plasmids pCMV.dR8.91 (expressing all necessary lentivirus helper functions) and pMD-G (expressing the vesicular stomatitis virus envelope protein G) were from Dr. Heike Laman (University of

Cambridge, UK). pF3A WG plasmid was from Promega. px459 CRISPR-Cas9 plasmid was
purchased from Addgene. More information about plasmids and oligonucleotide primers used are
given in the reagents table.

Antibodies and Reagents

Primary antibodies used were from the following sources: rabbit (Rb) anti-Myc (Cell Signaling, 2278),
mouse (Ms) anti-Myc (Cell Signaling, 9B11), Rb anti-Actin (Sigma, A2066), Ms anti-FLAG (Sigma
F1804), Rb anti-14-3-3 (Santa Cruz, sc-629), Ms anti -Spir-1 (Santa Cruz, sc-517039), Ms anti- Spir-
1 (Abcam, ab57463), Rb anti-DDX3 (Cell Signaling, 2635), Rb anti-IKKβ (Cell Signaling, 2684), Rb
anti-HA (Sigma, H6908), Ms anti-α-Tubulin (Millipore, 05-829), Ms anti-GAPDH (Sigma, G8795), Rb
anti-IRF3 (Cell Signaling, 4962), Ms anti-COPε (Santa Cruz, sc-133194), Rb anti-phospho-IRF3
Ser396 (Cell Signaling, 4947S) and Rb polyclonal anti-C6 (53). For dilutions used for the primary
antibodies, see reagents table. Secondary antibodies used (1:10,000 dilution) were IRDye 680RD-
conjugated goat anti-rabbit IgG or anti-mouse IgG and IRDye 800CW-conjugated goat anti-rabbit IgG
or anti-mouse IgG (LI-COR).

Reagents used in this study were: Anti-c-Myc Agarose from Santa Cruz Biotechnology, and
monoclonal Anti-HA-Agarose, clone HA-7, ANTI-FLAG M2 Affinity Gel and Poly-D-lysine
hydrobromide (all from Sigma Aldrich). Human IFNα, human TNF-α and mouse IL-1β were from
Peprotech, HMW poly(I:C) and puromycin were from InvivoGen, and doxycycline was from Melford.

Reporter Gene Assay

HEK293T cells were seeded in 96-well plates with 1.5×10^4 cells per well. After two days, cells were
transfected with 60 ng per well of the firefly luciferase reporter plasmids (IFNβ, ISRE, NF-κB or
ISG56.1), 10 ng per well of pTK-renilla luciferase and different amounts of the expression plasmid
under test or empty vector (EV) control using polyethylenimine (PEI, CellnTec, 2 µL per 1 µg DNA).
When necessary, EV plasmid was added to the transfection so that the final amount of DNA
transfected was kept constant. In cases where stimulation was done by transfecting another plasmid,
the same amount of EV was transfected to the non-stimulated (NS) control wells. Cells were
stimulated as shown in the Figures: (i) infection with SeV for 24 h (IFNβ Luc), (ii) 10 ng/mL of TNF-α
for 8 h (NF-κB Luc) or (iii) with 1000 U/mL of IFNα for 8 h (ISRE-Luc). After stimulation, cells were
washed with PBS, lysed with 100 uL/well of passive lysis buffer (Promega) and firefly and renilla
luciferase activities were measured using a FLUOstar luminometer (BMG). The firefly luciferase
activity in each sample was normalised to the renilla luciferase activity and fold inductions were
calculated relative to the non-stimulated controls for each plasmid. In all cases, data shown are
representative from at least three independent experiments with at least triplicate samples analysed

for each condition.

Immunoblotting

HEK293T (8×10^5) or MEFs (2×10^5) cells were seeded in 6-well plates and 24 h later cells were

stimulated by infection with SeV or transfection with 5 µg/mL of poly I:C using lipofectamine 2000 (Life

Technologies). After stimulation, cells were washed twice with ice-cold PBS, and scrapped into a cell

lysis buffer containing 50 mM Tris-HCl pH 8, 150 mM NaCl, 1 mM EDTA, 10% (v/v) glycerol, 1% (v/v)

Triton X-100 and 0.05% (v/v) NP-40, supplemented with protease (cOmplete Mini, Roche) and

phosphatase inhibitors (PhosSTOP, Roche). Protein concentration was determined using a

bicinchoninic acid protein assay kit (Pierce) before being boiled at 100 ºC for 5 min. Proteins were

then separated by SDS-polyacrylamide gel electrophoresis and transferred onto a nitrocellulose

Amersham Protran membrane (GE Healthcare). Membranes were blocked at room temperature with

either 5% (w/v) milk or 5% (w/v) bovine serum albumin (BSA, Sigma) in PBS containing 0.1% Tween

20. Then, membranes were incubated with a specific primary antibody diluted in blocking buffer at 4

ºC overnight. After washing, membranes were probed with LI-COR secondary antibodies at room

temperature followed by imaging using the LI-COR Odyssey imaging system, according to the

manufacturer's instructions. Where indicated, protein bands from at least two independent

experiments were quantified by using Odyssey software (LI-COR Biosciences).

Immunoprecipitation

HEK293T cells were seeded in 10-cm dishes (3.5×10^6 cells per dish) and transfected with the

plasmids indicated in the Figures using PEI. The following day, cells were washed twice with PSB,

lysed in immunoprecipitation (IP) lysis buffer (150 mM NaCl, 50 mM Tris-HCl pH 7.4, 0.5% (v/v)

Nonidet P-40 (NP-40) and protease (cOmplete Mini, Roche) and phosphatase inhibitors (PhosSTOP,

Roche)) and cleared by centrifugation at 21,000 *g* for 15 min at 4 ºC. Cleared lysates were then

incubated with 20 µL of ANTI-FLAG M2 Affinity Gel (Sigma Aldrich) or Anti-HA Agarose (Sigma

Aldrich) for 2 h, or with 50 uL of Anti-c-Myc Agarose overnight, at 4 ºC. Alternatively, proteins were

expressed by *in vitro* transcription/translation using the TNT SP6 High-Yield Wheat Germ Protein

Expression System (Promega) prior to incubation with the affinity resins. Immunoprecipitations were

washed 3 or 4 times in IP buffer and bound proteins were eluted in Laemmli SDS-PAGE loading buffer

and heated at 100 ºC for 5 min. Samples were then analysed by SDS-PAGE and immunoblotting.

RT-qPCR

HEK293T (4×10^5) or MEFs (1×10^5) cells were seeded in 12-well plates. The next day, cells were

stimulated by infection with SeV or transfection with 5 µg/mL of poly I:C using lipofectamine 2000 (Life

Technologies). RNA was extracted using the RNeasy kit (QIAGEN) and 500 ng of each RNA sample

was used to synthesise cDNA using Superscript III reverse transcriptase according to the

manufacturer's protocol (Invitrogen). mRNA was quantified by real-time PCR using a ViiA 7 Real-Time PCR System (Life Technologies), fast SYBR Green Master Mix (Applied Biosystems) and the following primers: human *CXCL10* (Fwd: GTGGCATTCAAGGAGTACCTC, Rev: GCCTTCGATTCTGGATTCAGA), human *IFNB1* (Fwd:ACATCCCTGAGGAGATTAAGCA, Rev: GCCAGGAGGTTCTCAACAATAG), human *IFIT1* (Fwd: CCTGAAAGGCCAGAATGAGG, Rev: TCCACCTTGTCCAGGTAAGT), human *GAPDH* (Fwd: ACCCAGAAGACTGTGGATGG, Rev: TTCTAGACGGCAGGTCAGGT), mouse *Ifit1* (Fwd: ACCATGGGAGAGAATGCTGAT, Rev: GCCAGGAGGTTGTGC), mouse *Il6* (Fwd: GTAGCTATGGTACTCCAGAAGAC. Rev: ACGATGATGCACTTGCAGAA), mouse *Cxcl10* (Fwd: ACTGCATCCATATCGATGAC, Rev: TTCATCGTGGCAATGATCTC) and mouse *Gapdh* (Fwd: ATGGTGAAGGTCGGTGTGAACGG, Rev: TTACTCCTTGGAGGCCATGTAGGC). Gene amplification was normalised to *GAPDH* (glyceraldehyde-3-phosphate dehydrogenase) amplification from the same sample, and the fold induction of genes in stimulated samples was calculated relative to the unstimulated control. Experiments were performed in at least biological duplicate and conducted at least twice.

ELISA

HEK293T (4×10^5) or MEFs (1×10^5) cells were seeded in 12-well plates. The following day, cells were stimulated by: (i) infection with SeV; (ii) transfection with 5 µg/mL of poly I:C using lipofectamine 2000 (Life Technologies), (iii) 50 ng/mL of human TNF-α or (iv) 50 ng/mL of mouse IL-1β. After stimulation, supernatants were assayed for human or murine CXCL-10 and IL-6 protein using Duoset enzyme-linked immunosorbent assay (ELISA) reagents (R&D Biosystems) according to the manufacturer's instructions.

CRISPR/Cas9-mediated genome editing

Two guide RNAs (gRNAs) were designed using online software (http://tools.genome-engineering.org) to target *SPIRE1* gene exon 3, which is shared by all Spir-1 isoforms. CRISPR/Cas9-mediated genome editing of HEK293T cells was performed as described (87). Briefly, px459 CRISPR/Cas9 plasmids with or without gRNA sequence were transfected into HEK293T using TransIT-LT1 transfection reagent (Mirus, MIR 2306). Puromycin (1 µg/mL) was added to transfected cells and, after 48 h, puromycin-resistant cells were serially diluted to obtain individual clones. Several clones were amplified, and a few potential knockout clones were selected by immunoblotting and confirmed by genomic DNA sequencing at the gRNA target sites. Only one gRNA was successful and one clonal cell line was confirmed to be knockout after PCR-amplified genomic DNA was cloned into bacterial plasmids and multiple colonies (n = 20) were sequenced. These all contained frameshift mutations. No wild type allele was identified.

Lentivirus transductions

Lentivirus particles for transduction were generated after transient co-transfection of HEK293T cells
seeded in 6-cm dishes. Cells were transfected with pCMV.dR8.91 and pMD-G vectors together with
either pLDT-EV or pLDT-Myc-Spir-1. After 48 h and 72 h, the supernatant was collected and passed
through a 0.45 μm filter. Lentivirus-containing supernatant was then used to infect HEK293T Spir-1
WT and KO cells. Transduced cells were selected with 1 μg/mL puromycin followed by clonal selection
by serial dilution. Spir-1 expression was assessed by immunoblotting.

Virus Infection
Plaque size analysis were performed in HEK293T Spir-1 WT and KO cells seeded in 6-well plates
coated with poly-D-lysine (Sigma). Once confluent, cells were infected with VACV-A5-GFP at 20 PFU
per well for 2 d. Virus plaques diameters (n=54) were measured using AxioVision 4.8 software and a
ZEISS Axio Vert.A1 fluorescent microscope.

Viral replication was measured by multi-step growth analyses of HEK293T Spir-1 WT and KO cells
infected with VACV-WR at 0.001 PFU / cell or ZIKV-mCherry at 0.01 PFU/cell. For VACV, at 48 h
post infection, cells were scraped in their medium and collected by centrifugation at 500 g for 5 min.
Cells were subjected to three rounds of freeze-thawing before the infectious viral titre was determined
by plaque assay on BSC-1 cells for 3 days. For ZIKV, at 72 h post infection, supernatants of infected
cells were collected, and virus infectivity was determined by plaque assay on Vero E6 cells for 5 days.
ZIKV-infected monolayers were also imaged using a Zeiss Axiovert 200M microscope. Images in
Figure 7D were processed using Adobe Photoshop 2020 to enhance linearly the mCherry
visualisation. SC-1 and VERO E6 cells were then fixed with 4% paraformaldehyde (PFA) and stained
with toluidine blue.

Statistical Analysis
Statistical analysis was carried out using one or two-way ANOVA test where appropriate with the
Bonferroni post-test, using the GraphPad Prism statistical software (Graph-Pad Software). Statistical
significance is expressed as follows: ns = not significant, *P < 0.05, **P < 0.01, ***P < 0.001, ****P <
0.0001.

Acknowledgements

We thank Dr. Martina Schröder, (Maynooth University, Ireland) and Prof. Dr. Eugen Kerkhoff
(University Hospital Regensburg, Germany) for providing cells lines and plasmids for this study. We
also thank Callum Talbot-Cooper for critical reading of the manuscript.

Conflicts of Interest

565 The authors declare that they have no conflict of interest.

Figure Legends

Fig 1: Spir-1 co-immunoprecipitates VACV protein K7 via its C-terminal region. HEK293T cells
were transfected (**A**, **B** and **D**) with Myc-tagged and FLAG-tagged plasmids overnight. Cell lysates
were immunoprecipitated using either Myc (**A** and **D** – right panel) or FLAG affinity resins (**B** and **D** –
middle panel) and analysed by SDS-PAGE and immunoblotting. (**C**) Schematic representations of
hSpir-1 isoform 2 full-length (top) and its C- and N-terminal truncations. (**E**) HEK293T cells were either
mock-infected or infected at 5 or 10 PFU/cell with vHA-K7 or vHA-B14 for 4 h. Lysates were
immunoprecipitated using HA-affinity resin and analysed by SDS-PAGE and immunoblotting. In (**A**),
(**B**), (**D**) and (**E**) the positions of molecular mass markers in kDa are shown on the left. Each
experiment was done 3 times and representative results are shown.

Fig 2: **Ectopic expression of Spir-1 increases IRF3-dependent gene expression induced by**
other stimuli, at or downstream of MAVS. HEK293T cells were transfected with IFNβ (**A**), ISRE (**B**)
and NF-κB (**C**) firefly luciferase reporter plasmids, together with TK-renilla luciferase and 40 ng of the
plasmids for expression of the indicated proteins. After overnight transfection, cells were stimulated
with SeV for 24 h (**A**), IFNα (**B**) or TNF-α (**C**) for 8 h. **D-I**) HEK293T cells were transfected with the
ISG56.1 firefly luciferase reporter plasmid, TK-renilla luciferase and plasmids for expression of the
indicated proteins. Cells were also co-transfected with EV as the non-stimulated (NS) controls or with
the 5 ng of CARD-domain of RIG-I (**D**, **E** and **I**), 40 ng of MAVS (**F**), 40 ng of TBK-1 (**G**) and 5 ng of
IRF3-5D (**H**) plasmids to activate the IRF3 pathway. EV was added to samples when necessary to
keep the final amount of DNA transfected as 40 ng in all samples. Cell lysates were prepared and
luciferase expression was measured and normalised to renilla luciferase. At least triplicate samples
were analysed for each condition. Data are expressed as the mean (± SEM) fold induction of the
firefly luciferase activity normalised to renilla values for the stimulated versus non-stimulated samples.
Data shown are representative of three independent experiments. Immunoblots underneath each
graph show the expression levels of the different proteins. The positions of molecular mass markers
in kDa are shown on the right and the antibodies used are shown on the left. ns = not significant; *P <
0.05; ****P < 0.0001.

Fig 3: Spir-1 and DDX3 share a conserved diphenylalanine motif that is required for IRF3 activation and direct binding to K7. (**A**) DDX3 knockdown was induced in HEK293T cells stably transfected with pTRIPZ-shDDX3 (or a NSC pTRIPZ vector) through incubation with doxycycline for 48 h. Twenty-four h after doxycycline addition, cells were transfected with FLAG-tagged plasmids overnight. Cell lysates were immunoprecipitated using FLAG-affinity resin and analysed by SDS-PAGE and immunoblotting. (**B**) DDX3 knockdown was induced in HEK293T shDDX3 cells as in (**A**) and cells were then transfected with ISG56.1-firefly luciferase reporter, TK-renilla luciferase, plasmids for expression of the indicated proteins together with the CARD-domain. Cell lysates were prepared and analysed as in Fig 2. Immunoblots underneath the graph show the expression levels of the different proteins. (**C**) Alignment of amino acid residues of Spir-1 and DDX3 showing the conserved diphenylalanine motif. (**D**) HEK293T cells were transfected overnight with Myc-tagged Spir-1 wild type, GFP or Spir-1 mutant FFAA together with FLAG-K7. Cell lysates were immunoprecipitated using either Myc (middle panel) or FLAG-affinity resins (right panel) and analysed by SDS-PAGE and immunoblotting. (**E**) HEK293T cells were transfected with ISG56.1 firefly luciferase reporter, TK-renilla luciferase, plasmids for expression of the indicated proteins together with the CARD-domain of RIG-I or EV as the NS control. Cell lysates were prepared and analysed as in (**B**). The panel underneath the graph shows immunoblots for the expression level of K7 and GAPDH. (**F**, **G**) Myc or FLAG-tagged proteins were synthesized by *in vitro* transcription/translation. Samples were immunoprecipitated using FLAG- (**F**) or Myc-affinity resins (**G**) and analysed by SDS-PAGE and immunoblotting. For all immunoblots, the positions of molecular mass markers in kDa are shown on the left and the antibodies used on the right. ns = not significant; ****P < 0.0001. HC/LC: IgG heavy chain or light chain, respectively.

Fig 4: K7 residue Asp31 is important for binding to Spir-1 and DDX3 and inhibition of IRF3 activation. HEK293T cells were transfected with FLAG-tagged GFP, K7 wild type or mutants, Myc-Spir-1 and HA-DDX3 (**A**) or with only FLAG-tagged plasmids overnight (**B**). Cell lysates were immunoprecipitated using FLAG-affinity resin and analysed by SDS-PAGE and immunoblotting. The positions of molecular mass markers in kDa are shown on the left and the antibodies used on the right. (**C**) HEK293T cells were transfected with ISG56.1-firefly luciferase reporter, TK-renilla luciferase, plasmids for expression of the indicated proteins together with the CARD-domain or EV as the NS control. Cell lysates were prepared and analysed as in Fig 2. Statistical analyses compared the fold induction of the mutant sample to its respective K7 wild type. The panel underneath the graph shows immunoblots for the expression levels of the different K7 proteins and α-tubulin. The positions of molecular mass markers in kDa are shown on the right and the antibodies used on the left. ns = not significant; ****P < 0.0001. LC: IgG light chain.

Fig 5: Spir-1 contributes to IRF3 phosphorylation and IRF3 stimulated gene expression after SeV infection in HEK293T cells. (**A**) Schematic of CRISPR-Cas9-mediated knockout strategy targeting *Spire1* exon 3 and single allele sequences of HEK293T Spir-1 knockout (KO) cells. (**B-F**) Spir-1 knockout (KO) HEK293T cells were transduced with empty vector (EV) or Myc-Spir-1 lentiviruses to rescue Spir-1 expression. Cells lines were either non-infected (NS) or infected with SeV for the indicated times. Cells were then either lysed and analysed by immunoblotting (**B**) or subjected to total RNA extraction followed by RT-qPCR (**C-E**). Supernatants were analysed by ELISA (**F**). (**G**) Spir-1 KO or wild type (WT) HEK293T cells were stimulated overnight with TNFα and supernatants were analysed by ELISA. ns = not significant, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Fig 6: IRF3 activation is reduced in Spir-1 KO MEFs. Spir-1 WT or KO MEF cells were transfected with poly I:C (**A, C-E, I-J**) or infected with SeV (**B, F-H, K-L**) for the indicated times. Cells were lysed and analysed by immunoblotting (**A, B**) and phospho-IRF3 bands intensity was quantified and normalised by the intensity of α-tubulin from at least two different experiments (**A**, **B**, bottom graphs). The positions of molecular mass markers in kDa are shown on the left and the antibodies used on the right. Cells were also subjected to total RNA extraction followed by RT-qPCR (**C-H**) and supernatants were analysed by ELISA (**I-L**). (**M, N**) Spir-1 WT or KO MEF cells were stimulated with IL-1β for 8 h and supernatants were analysed by ELISA. ns = not significant, *P < 0.05, **P < 0.01, ****P < 0.0001.

Fig 7: Spir-1 is a cellular restriction factor for VACV and ZIKV. (**A, B**) Spir-1 WT or KO and Spir-1 KO complemented HEK293T cells were infected with VACV-A5-GFP and plaque diameters were measured at 48 h p.i. (**A**) Representative plaques formed in each cell line. (**B**) Plaques diameter measurements (n=54). (**C**) Spir-1 WT or KO and Spir-1 KO complemented HEK293T cells were infected with VACV WR at 0.001 PFU/cell for 48 h and the virus yield was measured in BSC-1 cells. ns = not significant, *P<0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001. (**D and E**) Spir-1 WT or KO and Spir-1 KO complemented HEK293T cells were infected with ZIKV-mCherry at 0.01 PFU/cell for 72 h and ZIKV-infected monolayers were imaged (**D**) or the virus yield was measured in VERO E6 cells (**E**). ns = not significant, **P < 0.01, ***P < 0.001.

Supporting information

S1 Fig: Ectopic expression of Spir-1 doesn't affect IRF3-dependent gene expression induced by IKKε. HEK293T cells was transfected with the ISG56.1 firefly luciferase reporter plasmid, TK-renilla luciferase and plasmids for expression of the indicated proteins. Cells were also co-transfected with EV as the non-stimulated (NS) controls or with the 100 ng of IKKε plasmid to activate the IRF3

pathway. EV was added to samples when necessary to keep the final amount of DNA transfected as 40 ng in all samples. Cell lysates were prepared and luciferase expression was measured and normalised to renilla luciferase. At least triplicate samples were analysed for each condition. Data are expressed as the mean (± SEM) fold induction of the firefly luciferase activity normalised to renilla values for the stimulated versus non-stimulated samples. Data shown are representative of three independent experiments. Immunoblots underneath each graph show the expression levels of the different proteins. The positions of molecular mass markers in kDa are shown on the right and the antibodies used are shown on the left. ns = not significant; *P < 0.05; ****P < 0.0001.

S1 Table. Plasmids used in the study

S2 Table. Oligonucleotides used in this study

S3 Table. Primary antibodies used in the study

References

1. Janeway CA, Jr., Medzhitov R. Innate immune recognition. Annu Rev Immunol. 2002;20:197-216.
2. Pang IK, Iwasaki A. Control of antiviral immunity by pattern recognition and the microbiome. Immunol Rev. 2012;245(1):209-26.
3. Thompson MR, Kaminski JJ, Kurt-Jones EA, Fitzgerald KA. Pattern recognition receptors and the innate immune response to viral infection. Viruses. 2011;3(6):920-40.
4. Randall RE, Goodbourn S. Interferons and viruses: an interplay between induction, signalling, antiviral responses and virus countermeasures. J Gen Virol. 2008;89(Pt 1):1-47.
5. Garcia-Sastre A. Ten Strategies of Interferon Evasion by Viruses. Cell Host Microbe. 2017;22(2):176-84.
6. Fenner F, Henderson DA, Arita I, Jezek Z, Ladnyi ID. Smallpox and its eradication. World Health Organisation, Geneva. 1988.
7. Goebel SJ, Johnson GP, Perkus ME, Davis SW, Winslow JP, Paoletti E. The complete DNA sequence of vaccinia virus. Virology. 1990;179(1):247-66, 517-63.
8. Moss B, Smith GL. Poxviridae: the viruses and their replication. In: Howley PM, Knipe DM, editors. Fields Virology: DNA viruses. 2. 7th ed: Wolters Kluwer Inc; 2021. p. 573-613.
9. Smith GL, Benfield CTO, Maluquer de Motes C, Mazzon M, Ember SWJ, Ferguson BJ, et al. Vaccinia virus immune evasion: mechanisms, virulence and immunogenicity. J Gen Virol. 2013;94(Pt 11):2367-92.
10. Benfield CTO, Ren H, Lucas SJ, Bahsoun B, Smith GL. Vaccinia virus protein K7 is a virulence factor that alters the acute immune response to infection. J Gen Virol. 2013;94(Pt 7):1647-57.
11. Schroder M, Baran M, Bowie AG. Viral targeting of DEAD box protein 3 reveals its role in TBK1/IKKepsilon-mediated IRF activation. EMBO J. 2008;27(15):2147-57.
12. Teferi WM, Desaulniers MA, Noyce RS, Shenouda M, Umer B, Evans DH. The vaccinia virus K7 protein promotes histone methylation associated with heterochromatin formation. PLoS One. 2017;12(3):e0173056.
13. Li Y, Zhang L, Ke Y. Cellular interactome analysis of vaccinia virus K7 protein identifies three transport machineries as binding partners for K7. Virus Genes. 2017;53(6):814-22.

14. Pichlmair A, Kandasamy K, Alvisi G, Mulhern O, Sacco R, Habjan M, et al. Viral immune
modulators perturb the human molecular network by common and unique strategies. Nature.
2012;487(7408):486-90.
15. Lu Y, Stuart JH, Talbot-Cooper C, Agrawal-Singh S, Huntly B, Smid AI, et al. Histone
deacetylase 4 promotes type I interferon signaling, restricts DNA viruses, and is degraded via
vaccinia virus protein C6. Proc Natl Acad Sci U S A. 2019;116(24):11997-2006.
16. Soday L, Lu Y, Albarnaz JD, Davies CTR, Antrobus R, Smith GL, et al. Quantitative Temporal
Proteomic Analysis of Vaccinia Virus Infection Reveals Regulation of Histone Deacetylases by
an Interferon Antagonist. Cell Rep. 2019;27(6):1920-33 e7.
17. Manseau LJ, Schupbach T. cappuccino and spire: two unique maternal-effect loci required for
both the anteroposterior and dorsoventral patterns of the Drosophila embryo. Genes Dev.
1989;3(9):1437-52.
18. Wellington A, Emmons S, James B, Calley J, Grover M, Tolias P, et al. Spire contains actin
binding domains and is related to ascidian posterior end mark-5. Development.
1999;126(23):5267-74.
19. Quinlan ME, Heuser JE, Kerkhoff E, Mullins RD. Drosophila Spire is an actin nucleation factor.
Nature. 2005;433(7024):382-8.
20. Pechlivanis M, Samol A, Kerkhoff E. Identification of a short Spir interaction sequence at the C-
terminal end of formin subgroup proteins. J Biol Chem. 2009;284(37):25324-33.
21. Bosch M, Le KH, Bugyi B, Correia JJ, Renault L, Carlier MF. Analysis of the function of Spire in
actin assembly and its synergy with formin and profilin. Mol Cell. 2007;28(4):555-68.
22. Quinlan ME, Hilgert S, Bedrossian A, Mullins RD, Kerkhoff E. Regulatory interactions between
two actin nucleators, Spire and Cappuccino. J Cell Biol. 2007;179(1):117-28.
23. Pfender S, Kuznetsov V, Pleiser S, Kerkhoff E, Schuh M. Spire-type actin nucleators cooperate
with Formin-2 to drive asymmetric oocyte division. Curr Biol. 2011;21(11):955-60.
24. Schuh M. An actin-dependent mechanism for long-range vesicle transport. Nat Cell Biol.
2011;13(12):1431-6.
25. Vizcarra CL, Kreutz B, Rodal AA, Toms AV, Lu J, Zheng W, et al. Structure and function of the
interacting domains of Spire and Fmn-family formins. Proc Natl Acad Sci U S A.
2011;108(29):11884-9.
26. Zeth K, Pechlivanis M, Samol A, Pleiser S, Vonrhein C, Kerkhoff E. Molecular basis of actin
nucleation factor cooperativity: crystal structure of the Spir-1 kinase non-catalytic C-lobe
domain (KIND)*formin-2 formin SPIR interaction motif (FSI) complex. J Biol Chem.
2011;286(35):30732-9.
27. Quinlan ME. Direct interaction between two actin nucleators is required in Drosophila
oogenesis. Development. 2013;140(21):4417-25.
28. Montaville P, Jegou A, Pernier J, Compper C, Guichard B, Mogessie B, et al. Spire and Formin
2 synergize and antagonize in regulating actin assembly in meiosis by a ping-pong mechanism.
PLoS Biol. 2014;12(2):e1001795.
29. Tittel J, Welz T, Czogalla A, Dietrich S, Samol-Wolf A, Schulte M, et al. Membrane targeting of
the Spir.formin actin nucleator complex requires a sequential handshake of polar interactions. J
Biol Chem. 2015;290(10):6428-44.
30. Ducka AM, Joel P, Popowicz GM, Trybus KM, Schleicher M, Noegel AA, et al. Structures of
actin-bound Wiskott-Aldrich syndrome protein homology 2 (WH2) domains of Spire and the
implication for filament nucleation. Proc Natl Acad Sci U S A. 2010;107(26):11757-62.
31. Sitar T, Gallinger J, Ducka AM, Ikonen TP, Wohlhoefler M, Schmoller KM, et al. Molecular
architecture of the Spire-actin nucleus and its implication for actin filament assembly. Proc Natl
Acad Sci U S A. 2011;108(49):19575-80.
32. Pylypenko O, Welz T, Tittel J, Kollmar M, Chardon F, Malherbe G, et al. Coordinated
recruitment of Spir actin nucleators and myosin V motors to Rab11 vesicle membranes. Elife.
2016;5.
33. Alzahofi N, Welz T, Robinson CL, Page EL, Briggs DA, Stainthorp AK, et al. Rab27a co-
ordinates actin-dependent transport by controlling organelle-associated motors and track
assembly proteins. Nat Commun. 2020;11(1):3495.

34. Kerkhoff E, Simpson JC, Leberfinger CB, Otto IM, Doerks T, Bork P, et al. The Spir actin
organizers are involved in vesicle transport processes. Curr Biol. 2001;11(24):1963-8.
35. Lagal V, Abrivard M, Gonzalez V, Perazzi A, Popli S, Verzeroli E, et al. Spire-1 contributes to
the invadosome and its associated invasive properties. J Cell Sci. 2014;127(Pt 2):328-40.
36. Schumacher N, Borawski JM, Leberfinger CB, Gessler M, Kerkhoff E. Overlapping expression
pattern of the actin organizers Spir-1 and formin-2 in the developing mouse nervous system
and the adult brain. Gene Expr Patterns. 2004;4(3):249-55.
37. Pleiser S, Rock R, Wellmann J, Gessler M, Kerkhoff E. Expression patterns of the mouse Spir-2
actin nucleator. Gene Expr Patterns. 2010;10(7-8):345-50.
38. Fagerberg L, Hallstrom BM, Oksvold P, Kampf C, Djureinovic D, Odeberg J, et al. Analysis of
the human tissue-specific expression by genome-wide integration of transcriptomics and
antibody-based proteomics. Mol Cell Proteomics. 2014;13(2):397-406.
39. Morel E, Parton RG, Gruenberg J. Annexin A2-dependent polymerization of actin mediates
endosome biogenesis. Dev Cell. 2009;16(3):445-57.
40. Belin BJ, Lee T, Mullins RD. DNA damage induces nuclear actin filament assembly by Formin -
2 and Spire-(1/2) that promotes efficient DNA repair. [corrected]. Elife. 2015;4:e07735.
41. Manor U, Bartholomew S, Golani G, Christenson E, Kozlov M, Higgs H, et al. A mitochondria-
anchored isoform of the actin-nucleating spire protein regulates mitochondrial division. Elife.
2015;4.
42. Wen Q, Li N, Xiao X, Lui WY, Chu DS, Wong CKC, et al. Actin nucleator Spire 1 is a regulator
of ectoplasmic specialization in the testis. Cell Death Dis. 2018;9(2):208.
43. Ovsyannikova IG, Kennedy RB, O'Byrne M, Jacobson RM, Pankratz VS, Poland GA. Genome-
wide association study of antibody response to smallpox vaccine. Vaccine. 2012;30(28):4182-9.
44. Kerkhoff E. Cellular functions of the Spir actin-nucleation factors. Trends Cell Biol.
2006;16(9):477-83.
45. Kollmar M, Welz T, Straub F, Alzahofi N, Hatje K, Briggs DA, et al. Animal evolution coincides
with a novel degree of freedom in exocytic transport processes. bioRxiv.
2020;https://doi.org/10.1101/591974.
46. Neidel S, Maluquer de Motes C, Mansur DS, Strnadova P, Smith GL, Graham SC. Vaccinia
virus protein A49 is an unexpected member of the B-cell Lymphoma (Bcl)-2 protein family. J
Biol Chem. 2015;290(10):5991-6002.
47. Mansur DS, Maluquer de Motes C, Unterholzner L, Sumner RP, Ferguson BJ, Ren H, et al.
Poxvirus targeting of E3 ligase beta-TrCP by molecular mimicry: a mechanism to inhibit NF-
kappaB activation and promote immune evasion and virulence. PLoS Pathog.
2013;9(2):e1003183.
48. Benzinger A, Muster N, Koch HB, Yates JR, 3rd, Hermeking H. Targeted proteomic analysis of
14-3-3 sigma, a p53 effector commonly silenced in cancer. Mol Cell Proteomics. 2005;4(6):785-
95.
49. Ewing RM, Chu P, Elisma F, Li H, Taylor P, Climie S, et al. Large-scale mapping of human
protein-protein interactions by mass spectrometry. Mol Syst Biol. 2007;3:89.
50. Chen RA, Jacobs N, Smith GL. Vaccinia virus strain Western Reserve protein B14 is an
intracellular virulence factor. J Gen Virol. 2006;87(Pt 6):1451-8.
51. Chen RA, Ryzhakov G, Cooray S, Randow F, Smith GL. Inhibition of IkappaB kinase by
vaccinia virus virulence factor B14. PLoS Pathog. 2008;4(2):e22.
52. Kato H, Takeuchi O, Sato S, Yoneyama M, Yamamoto M, Matsui K, et al. Differential roles of
MDA5 and RIG-I helicases in the recognition of RNA viruses. Nature. 2006;441(7089):101-5.
53. Unterholzner L, Sumner RP, Baran M, Ren H, Mansur DS, Bourke NM, et al. Vaccinia virus
protein C6 is a virulence factor that binds TBK-1 adaptor proteins and inhibits activation of IRF3
and IRF7. PLoS Pathog. 2011;7(9):e1002247.
54. Stuart JH, Sumner RP, Lu Y, Snowden JS, Smith GL. Vaccinia Virus Protein C6 Inhibits Type I
IFN Signalling in the Nucleus and Binds to the Transactivation Domain of STAT2. PLoS Pathog.
2016;12(12):e1005955.
55. Ferguson BJ, Benfield CTO, Ren H, Lee VH, Frazer GL, Strnadova P, et al. Vaccinia virus
protein N2 is a nuclear IRF3 inhibitor that promotes virulence. J Gen Virol. 2013;94(Pt 9):2070-
81.

56.   Kalverda AP, Thompson GS, Vogel A, Schroder M, Bowie AG, Khan AR, et al. Poxvirus K7
protein adopts a Bcl-2 fold: biochemical mapping of its interactions with human DEAD box RNA
helicase DDX3. J Mol Biol. 2009;385(3):843-53.
57.   Oda S, Schroder M, Khan AR. Structural basis for targeting of human RNA helicase DDX3 by
poxvirus protein K7. Structure. 2009;17(11):1528-37.
58.   Fullam A, Schroder M. DExD/H-box RNA helicases as mediators of anti-viral innate immunity
and essential host factors for viral replication. Biochim Biophys Acta. 2013;1829(8):854-65.
59.   Gu L, Fullam A, McCormack N, Hohn Y, Schroder M. DDX3 directly regulates TRAF3
ubiquitination and acts as a scaffold to co-ordinate assembly of signalling complexes
downstream from MAVS. Biochem J. 2017;474(4):571-87.
60.   Wang T, Birsoy K, Hughes NW, Krupczak KM, Post Y, Wei JJ, et al. Identification and
characterization of essential genes in the human genome. Science. 2015;350(6264):1096-101.
61.   Carpentier DC, Gao WN, Ewles H, Morgan GW, Smith GL. Vaccinia virus protein complex
F12/E2 interacts with kinesin light chain isoform 2 to engage the kinesin-1 motor complex.
PLoS Pathog. 2015;11(3):e1004723.
62.   Everett RD, Parsy ML, Orr A. Analysis of the functions of herpes simplex virus type 1 regulatory
protein ICP0 that are critical for lytic infection and derepression of quiescent viral genomes. J
Virol. 2009;83(10):4963-77.
63.   Stinski MF, Meier JL. Immediate-early viral gene regulation and function. In: Arvin A,
Campadelli-Fiume G, Mocarski E, Moore PS, Roizman B, Whitley R, et al., editors. Human
Herpesviruses: Biology, Therapy, and Immunoprophylaxis. Cambridge2007.
64.   Servant MJ, Grandvaux N, tenOever BR, Duguay D, Lin R, Hiscott J. Identification of the
minimal phosphoacceptor site required for in vivo activation of interferon regulatory factor 3 in
response to virus and double-stranded RNA. J Biol Chem. 2003;278(11):9441-7.
65.   Pleiser S, Banchaabouchi MA, Samol-Wolf A, Farley D, Welz T, Wellbourne-Wood J, et al.
Enhanced fear expression in Spir-1 actin organizer mutant mice. Eur J Cell Biol. 2014;93(5-
6):225-37.
66.   Carter GC, Rodger G, Murphy BJ, Law M, Krauss O, Hollinshead M, et al. Vaccinia virus cores
are transported on microtubules. J Gen Virol. 2003;84(Pt 9):2443-58.
67.   Neidel S, Torres AA, Ren H, Smith GL. Leaky scanning translation generates a second A49
protein that contributes to vaccinia virus virulence. J Gen Virol. 2020;101(5):533-41.
68.   Smith GL, Vanderplasschen A, Law M. The formation and function of extracellular enveloped
vaccinia virus. J Gen Virol. 2002;83(Pt 12):2915-31.
69.   Kato H, Sato S, Yoneyama M, Yamamoto M, Uematsu S, Matsui K, et al. Cell type-specific
involvement of RIG-I in antiviral response. Immunity. 2005;23(1):19-28.
70.   Kato H, Takeuchi O, Mikamo-Satoh E, Hirai R, Kawai T, Matsushita K, et al. Length-dependent
recognition of double-stranded ribonucleic acids by retinoic acid-inducible gene-I and
melanoma differentiation-associated gene 5. J Exp Med. 2008;205(7):1601-10.
71.   Brisse M, Ly H. Comparative Structure and Function Analysis of the RIG-I-Like Receptors: RIG-
I and MDA5. Front Immunol. 2019;10:1586.
72.   Soulat D, Burckstummer T, Westermayer S, Goncalves A, Bauch A, Stefanovic A, et al. The
DEAD-box helicase DDX3X is a critical component of the TANK-binding kinase 1-dependent
innate immune response. EMBO J. 2008;27(15):2135-46.
73.   Seth RB, Sun L, Ea CK, Chen ZJ. Identification and characterization of MAVS, a mitochondrial
antiviral signaling protein that activates NF-kappaB and IRF 3. Cell. 2005;122(5):669-82.
74.   Weber F, Wagner V, Rasmussen SB, Hartmann R, Paludan SR. Double-stranded RNA is
produced by positive-strand RNA viruses and DNA viruses but not in detectable amounts by
negative-strand RNA viruses. J Virol. 2006;80(10):5059-64.
75.   Liu Y, Olagnier D, Lin R. Host and Viral Modulation of RIG-I-Mediated Antiviral Immunity. Front
Immunol. 2016;7:662.
76.   Serman TM, Gack MU. Evasion of Innate and Intrinsic Antiviral Pathways by the Zika Virus.
Viruses. 2019;11(10).
77.   Colby C, Duesberg PH. Double-stranded RNA in vaccinia virus infected cells. Nature.
1969;222(5197):940-4.

78. Rice AP, Roberts WK, Kerr IM. 2-5A accumulates to high levels in interferon-treated, vaccinia
virus-infected cells in the absence of any inhibition of virus replication. J Virol. 1984;50(1):220-
8.
79. Pichlmair A, Schulz O, Tan CP, Rehwinkel J, Kato H, Takeuchi O, et al. Activation of MDA5
requires higher-order RNA structures generated during virus infection. J Virol.
2009;83(20):10761-9.
80. Marq JB, Hausmann S, Luban J, Kolakofsky D, Garcin D. The double-stranded RNA binding
domain of the vaccinia virus E3L protein inhibits both RNA- and DNA-induced activation of
interferon beta. J Biol Chem. 2009;284(38):25471-8.
81. Valentine R, Smith GL. Inhibition of the RNA polymerase III-mediated dsDNA-sensing pathway
of innate immunity by vaccinia virus protein E3. J Gen Virol. 2010;91(Pt 9):2221-9.
82. Liu SW, Katsafanas GC, Liu R, Wyatt LS, Moss B. Poxvirus decapping enzymes enhance
virulence by preventing the accumulation of dsRNA and the induction of innate antiviral
responses. Cell Host Microbe. 2015;17(3):320-31.
83. Mutso M, Saul S, Rausalu K, Susova O, Zusinaite E, Mahalingam S, et al. Reverse genetic
system, genetically stable reporter viruses and packaged subgenomic replicon based on a
Brazilian Zika virus isolate. J Gen Virol. 2017;98(11):2712-24.
84. Torres AA, Albarnaz JD, Bonjardim CA, Smith GL. Multiple Bcl-2 family immunomodulators
from vaccinia virus regulate MAPK/AP-1 activation. J Gen Virol. 2016;97(9):2346-51.
85. Odon V, Georgana I, Holley J, Morata J, Maluquer de Motes C. Novel Class of Viral Ankyrin
Proteins Targeting the Host E3 Ubiquitin Ligase Cullin-2. J Virol. 2018;92(23).
86. Everett RD, Bell AJ, Lu Y, Orr A. The replication defect of ICP0-null mutant herpes simplex
virus 1 can be largely complemented by the combined activities of human cytomegalovirus
proteins IE1 and pp71. J Virol. 2013;87(2):978-90.
87. Ran FA, Hsu PD, Wright J, Agarwala V, Scott DA, Zhang F. Genome engineering using the
CRISPR-Cas9 system. Nat Protoc. 2013;8(11):2281-308.

904
905

Supporting information

907
**S1 Fig. Ectopic expression of Spir-1 does not affect IRF3-dependent gene expression induced
by IKKε (related to Figure 2).** HEK293T cells were transfected with the ISG56.1 firefly luciferase
reporter plasmid, TK-renilla luciferase and plasmids for expression of the indicated proteins. Cells
were also co-transfected with EV as the non-stimulated (NS) controls or with 100 ng of IKKε plasmid
to activate the IRF3 pathway. EV was added to samples when necessary to keep the final amount of
DNA transfected as 40 ng in all samples. Cell lysates were prepared and luciferase expression was
measured and normalised to renilla luciferase. At least triplicate samples were analysed for each
condition. Data are expressed as the mean (± SEM) fold induction of the firefly luciferase activity
normalised to renilla values for the stimulated versus non-stimulated samples. Data shown are
representative of three independent experiments. Immunoblots underneath each graph show the
expression levels of the different proteins. The positions of molecular mass markers in kDa are shown
on the right and the antibodies used are shown on the left. ns = not significant; *P < 0.05; ****P <
0.0001.
921
922
923 **S1 Table. Plasmids used in the study**
924

PLASMID	SOURCE
human nMyc-Spir-1-pcDNA3.1	Gift from Prof. Dr. E. Kerkhoff (University Hospital Regensburg, Germany)
human nMyc-Spir-2-pcDNA3.1	Gift from Prof. Dr. E. Kerkhoff (University Hospital Regensburg, Germany)
mouse nMyc-B-TrCP-pcDNA3.1	(Mansur *et al*, 2013)

GFP-Myc-pCMV	(Neidel *et al*, 2019)
nFlag-coK7-pcDNA4/TO	(Torres *et al*, 2016)
coA49-cFlag-pcDNA3.1	(Neidel *et al.*, 2019)
GFP-cFlag-pcDNA4/TO	Sarah Neidel (Prof. G.L. Smith laboratory)
human nMyc-Spir-1-NT-pcDNA3.1	This study
human nMyc-Spir-1-CT-pcDNA3.1	This study
coN1-cTAP-pcDNA4/TO	(Maluquer de Motes *et al*, 2011)
coC6-cTAP-pcDNA4/TO	(Unterholzner *et al*, 2011)
nTAP-coN2-pcDNA4/TO	(Ferguson *et al*, 2013)
nFlag-coB14-pcDNA4/TO	(Torres *et al.*, 2016)
pcDNA4/TO	Invitrogen
nMyc-DDX3-pCMV	Gift from Dr M. Schröder - (Gu *et al*, 2013)
nHA-DDX3-pCMV	Gift from Dr M. Schröder - (Gu *et al.*, 2013)
IFN-β-luc	Gift from Dr T. Taniguchi (University of Tokyo, Japan)
ISRE-luc	Gift from Dr. A. Bowie (Trinity College, Dublin, Ireland)
NF-κB-luc	Gift from Dr. A. Bowie (Trinity College, Dublin, Ireland)
ISG56.1-luc	Gift from G. Sen (Cleveland Clinic, Cleveland, OH, USA).
TK-Renilla	Promega
Flag-RIG-I-CARD	Gift from A. Garcia-Sastre (Mount Sinai, New York, USA)
HA-MAVS	(Unterholzner *et al.*, 2011)
Flag-TBK1	(Unterholzner *et al.*, 2011)
Flag-IKKe	(Unterholzner *et al.*, 2011)
IRF3-5D	(Unterholzner *et al.*, 2011)
human nMyc-Spir-1-FFAA-pcDNA3.1	This study
pF3A-nFlag-coK7	This study
pF3A-nMyc-Spir-1	This study
pF3A-nMyc-DDX3	This study
pF3A-F12-cFlagd	GLS - unpublished
pF3A-nMyc-TAB2	GLS - unpublished
nFlag-coK7-D28A-pcDNA4/TO	This study
nFlag-coK7-D31A-pcDNA4/TO	This study
nFlag-coK7-DDAA-pcDNA4/TO	This study
pSpCas9(BB)-2A-Puro (px459)	(Ran *et al*, 2013) - Addgene plasmid #62988
px459-Spir-1-sgRNA#2	This study
pLKO.DCMV.TetO.mcs	(Everett *et al*, 2013)
pLKO.DCMV.TetO.Myc-Spir-1	This study
pCMV.dR8.91	Gift from Dr H. Laman (Lomonosov *et al*, 2011)
pMD-G	Gift from Dr H. Laman (Lomonosov *et al.*, 2011)

926
927 **S2 Table. Oligonucleotides used in this study**
928

PLASMID CONSTRUCTION	PRIMER
human nMyc-Spir-1-CT-pcDNA3.1	AAAAGCGGCCGCCAGTGTGATGGATGGGTCAGATCTCACTGAT
	AAAAGGATCCGCCGCCGCCATGGAGCAGAAGCTGATCTCCGAGGAGGACCTGGCT
human nMyc-Spir-1-NT-pcDNA3.1	AAAAGCGGCCGCCAGTGTGATGGATGGGTCACTCCTCTGGTGATACAGGCCGCAGCT
	AAGCTTGGTACCGAGCTCGGATCCGCCG
pLKO.DCMV.TetO.Myc-Spir-1	AAAAACGCGTGCCGCCAGTGTGATGGATGGGTCAGATCTCACTGATCGTCCT
	AAAAGCTAGCGCCGCCGCCATGGAGCAGAAGCTGATCTCCGAGGAGGA
human nMyc-Spir-1-FFAA-pcDNA3.1	TTGCCGAACCAGGAGGTTTTCCGCCGCCACTTGGTCTTATACCTGTCAG
	CTGACAGGTATAAGACCAAGTGGCGGCGGAAAACCTCCTGGTTCGGCAA
nFlag-coK7-D28A-pcDNA4.1/TO	CGGGACAGCATCATCGCCCTGATCGACGAGTAC
	GTACTCGTCGATCAGGGCGATGATGCTGTCCCG
nFlag-coK7-D31A-pcDNA4.1/TO	GCATCATCGACCTGATCGCCGAGTACATCACCTGGCG
	CGCCAGGTGATGTACTCGGCGATCAGGTCGATGATGC
nFlag-coK7-DDAA-pcDNA4.1/TO	CGGGACAGCATCATCGCCCTGATCGCCGAGTAC
	GTACTCGGCGATCAGGGCGATGATGCTGTCCCG
pF3A-nFlag-coK7	AAAAGCGATCGCATGGACTACAAAGACGATGACGACAAG
	AAAATTTAAACTCAGTTCAGCTTCTTTTCCAGGAA
pF3A-nMyc-Spir-1	AAAAGCGATCGCATGGAGCAGAAGCTGATCTCCGA
	AAAATTTAAACTCAGATCTCACTGATCGTCCTCTC
pF3A-nMyc-DDX3	AAAAGTTTAAACTCAGTTACCCCACCAGTCAACCC
	AAAAGCGATCGCATGGAGCAGAAGCTGATCTCAGAGGA
CRISPR/Cas9 gRNA targeting region	**gRNA SEQUENCE**
Spire1 exon 3	AGAGCAGCTTATCGATCACA
SEQUENCING PRIMERS	
Forward primer for exon 3 of human *Spire1*	TGGGAAATGTGGAAAATCAACTCG
Reverse primer for exon 3 of human *Spire1*	AAAAATAACAGCTTGGACACAGTGG

929
930
931 **S3 Table. Primary antibodies used in the study**
932

ANTIBODIES	SOURCE	DILUTION
Rb anti-Myc	Cell Signaling, 2278	1 in 1000

Ms anti-Myc	Cell Signaling, 9B11	1 in 1000
Rb anti-Actin	Sigma, A2066	1 in 1000
Ms anti-Flag	Sigma, F1804	1 in 1000
Rb anti-14-3-3	Santa Cruz, sc-629	1 in 1000
Ms anti-Spir-1	Santa Cruz, sc-517039	1 in 500
Ms anti-Spir-1	Abcam, ab57463	1 in 1000
Rb anti-DDX3	Cell Signaling, 2635	1 in 1000
Rb anti-IKKb	Cell Signaling, 2684	1 in 1000
Rb anti-HA	Sigma, H6908	1 in 1000
Rb anti-C6	(Unterholzner *et al.*, 2011)	1 in 1000
Ms anti-alpha-Tubulin	Millipore, 05-829	1 in 5000
Ms anti-GAPDH	Sigma, G8795	1 in 1000
Rb anti-IRF3	Cell Signaling, 4962	1 in 1000
Ms anti-COPe	Santa Cruz, sc-133194	1 in 500
Rb anti-p-IRF3	Cell Signaling, 4947S	1 in 1000

933
934
935 **References**
936

Everett RD, Bell AJ, Lu Y, Orr A (2013) The replication defect of ICP0-null mutant herpes simplex
virus 1 can be largely complemented by the combined activities of human cytomegalovirus
proteins IE1 and pp71. *J Virol* 87: 978-990
Ferguson BJ, Benfield CTO, Ren H, Lee VH, Frazer GL, Strnadova P, Sumner RP, Smith GL (2013)
Vaccinia virus protein N2 is a nuclear IRF3 inhibitor that promotes virulence. *J Gen Virol* 94:
2070-2081
Gu L, Fullam A, Brennan R, Schroder M (2013) Human DEAD box helicase 3 couples IkappaB
kinase epsilon to interferon regulatory factor 3 activation. *Mol Cell Biol* 33: 2004-2015
Lomonosov M, Meziane el K, Ye H, Nelson DE, Randle SJ, Laman H (2011) Expression of Fbxo7 in
haematopoietic progenitor cells cooperates with p53 loss to promote lymphomagenesis. *PLoS*
One 6: e21165
Maluquer de Motes C, Cooray S, Ren H, Almeida GM, McGourty K, Bahar MW, Stuart DI, Grimes
JM, Graham SC, Smith GL (2011) Inhibition of apoptosis and NF-kappaB activation by vaccinia
protein N1 occur via distinct binding surfaces and make different contributions to virulence. *PLoS*
Pathog 7: e1002430
Mansur DS, Maluquer de Motes C, Unterholzner L, Sumner RP, Ferguson BJ, Ren H, Strnadova P,
Bowie AG, Smith GL (2013) Poxvirus targeting of E3 ligase beta-TrCP by molecular mimicry: a
mechanism to inhibit NF-kappaB activation and promote immune evasion and virulence. *PLoS*
Pathog 9: e1003183
Neidel S, Ren H, Torres AA, Smith GL (2019) NF-kappaB activation is a turn on for vaccinia virus
phosphoprotein A49 to turn off NF-kappaB activation. *Proc Natl Acad Sci U S A* 116: 5699-5704
Ran FA, Hsu PD, Wright J, Agarwala V, Scott DA, Zhang F (2013) Genome engineering using the
CRISPR-Cas9 system. *Nat Protoc* 8: 2281-2308
Torres AA, Albarnaz JD, Bonjardim CA, Smith GL (2016) Multiple Bcl-2 family immunomodulators
from vaccinia virus regulate MAPK/AP-1 activation. *J Gen Virol* 97: 2346-2351
Unterholzner L, Sumner RP, Baran M, Ren H, Mansur DS, Bourke NM, Randow F, Smith GL, Bowie
AG (2011) Vaccinia virus protein C6 is a virulence factor that binds TBK-1 adaptor proteins and
inhibits activation of IRF3 and IRF7. *PLoS Pathog* 7: e1002247

Fig 3

A

Input   IP:FLAG

	NSC	shDDX3	NSC	shDDX3

FLAG: A49   K7   A49   K7   A49   K7   A49   K7

Dox: − +   − +   − +   − +   − +   − +   − +   − +

18 —   IB:FLAG
72 —   IB:DDX3
93 —   IB:Spir-1
42 —   IB:Actin

B

ISG56.1-luc Fold Induction

EV   Spir-1   K7   Spir-1+ K7   EV   Spir-1   K7   Spir-1+ K7

w/o DOX   with DOX

75 — IB:DDX3
20 — IB:FLAG
100 — IB:Myc
42 — IB:Actin

C

Spir-1_601-610 RFSFFTWSYT
DDX3_81-90   KSSFFSDRGS
                : *** : :

D

Input   IP:Myc   IP: FLAG

Myc: Spir-1 GFP FFAA   Spir-1 GFP FFAA   Spir-1 GFP FFAA

FLAG-K7   FLAG-K7   FLAG-K7

100 —
75 —
63 —
48 —
35 —   IB:Myc

20 —
17 —   IB:FLAG

35 —
25 —   IB:14-3-3

48 —
35 —   IB:Actin

E

ISG56.1-luc Fold Induction

ns
****

EV   Spir-1   FFAA

NS   RIG-I-CARD

93 — IB:Myc
42 — IB:GAPDH

F

Input   IP:FLAG

FLAG: TAB2 + K7   DDX3+ K7   Spir-1 + K7   Spir-1 + F12

Myc: TAB2 + K7   DDX3+ K7   Spir-1 + K7   Spir-1 + F12

100 —   IB:Myc
75 —
63 —   HC
48 —   IB:FLAG
20 —   LC
17 —

G

Input   IP:Myc

FLAG: TAB2 + K7   DDX3+ K7   Spir-1 + K7   Spir-1 + F12

Myc: TAB2 + K7   DDX3+ K7   Spir-1 + K7   Spir-1 + F12

100 —   IB:Myc
63 —   IB:FLAG (F12)
20 —   IB:FLAG (K7)

*IgG heavy chain

973   Fig 4

975   Fig 5

979   Fig 7

980
981
982

983   S1 Fig

984