Title: Genome-wide identification of tomato xylem sap fitness factors for three

plant-pathogenic *Ralstonia* species

Authors: Stratton J. Georgoulis[a], Katie E. Shalvarjian[b], Tyler C. Helmann[b,c], Corri D.

Hamilton[d], Hans K. Carlson[e], Adam M. Deutschbauer[b,e], and Tiffany M. Lowe-Power[a,b]#

[a]Dept Plant Pathology, UC Davis, Davis, California, USA

[b]Dept Plant and Microbial Biology, UC Berkeley, Berkeley, California, USA

[c]Emerging Pests and Pathogens Research Unit, Robert W. Holley Center, Agricultural

Research Service, U.S. Department of Agriculture, Ithaca, NY, USA

[d]Dept Plant Pathology, UW-Madison, Madison, Wisconsin, USA

[e]Environmental Genomics and Systems Biology Division, Lawrence Berkeley National

Laboratory, Berkeley, California, USA

19 **Running Title:** *Ralstonia* fitness factors for xylem sap growth

20 **# Corresponding author**: Tiffany M. Lowe-Power, tlowepower@ucdavis.edu

22 **Abstract Word Count**: 214

23 **Text word count:** 3,892

Abstract

Plant pathogenic *Ralstonia* spp. colonize plant xylem and cause wilt diseases on a broad range of host plants. To identify genes that promote growth of diverse *Ralstonia* strains in xylem sap from tomato plants, we performed genome-scale genetic screens (random barcoded transposon mutant sequencing screens; RB-TnSeq) in three strains spanning the genetic, geographical, and physiological range of plant pathogenic *Ralstonia: R. solanacearum* IBSBF1503, *R. pseudosolanacearum* GMI1000, and *R. syzygii* PSI07. Contrasting mutant fitness phenotypes in culture media versus in xylem sap suggest that *Ralstonia* strains are adapted to *ex vivo* xylem sap and that culture media impose foreign selective pressures. Although wild-type *Ralstonia* grew in sap and in rich medium with similar doubling times and to a similar carrying capacity, more genes were essential for growth in sap than in rich medium. Each strain required many genes associated with envelope remodeling and repair processes for full fitness in xylem sap. These genes were associated with peptidoglycan peptide formation (*murI*), secretion of periplasmic proteins (*tatC*), periplasmic protein folding (*dsbA*), synthesis of osmoregulated periplasmic glucans (*mdoGH*), and LPS biosynthesis. Mutants in four genes had strong, sap-specific fitness defects in all strain backgrounds: *murI*, *thiC*, *purU*, and a lipoprotein (RSc2007*)*. Many amino acid biosynthesis genes were required for fitness in both minimal medium and xylem sap. Multiple mutants with insertions in virulence regulators had gains-of-fitness in culture media and neutral fitness in sap. Our genome-scale genetic screen identified *Ralstonia* fitness factors that promote growth in xylem sap, an ecologically relevant condition.

Importance

Traditional transposon mutagenesis genetic screens pioneered molecular plant pathology and identified core virulence traits like the type III secretion system. TnSeq approaches that

leverage next-generation sequencing to rapidly quantify transposon mutant phenotypes are

ushering in a new wave of biological discovery. Here we have adapted a genome-scale approach,

random barcoded transposon mutant sequencing (RB-TnSeq), to discover fitness factors that

promote growth of three related bacterial strains in a common niche, tomato xylem sap. Fitness

of wild-type and mutants show that *Ralstonia* spp. are adapted to grow well in xylem sap from

their natural host plant, tomato. Our screen identified multiple sap-specific fitness factors with

roles in maintaining the bacterial envelope. These factors include putative adaptations to resist

plant defenses that may include antimicrobial proteins and specialized metabolites that damage

bacterial membranes.

Introduction

Ralstonia solanacearum, R. pseudosolanacearum, and *R. syzygii* (hereafter, "*Ralstonia*")

comprise the monophyletic but diverse species complex of plant wilt pathogens (1, 2). *Ralstonia*

strains are adapted to plant hosts belonging to over 50 botanical families and are distributed

worldwide in in warm tropics and temperate subtropical highlands with year-round precipitation

(3–7). Most *Ralstonia* strains invade plant roots, gain entry to the water-transporting xylem

vasculature, and spread systemically, which disrupts xylem function and fatally wilts the host

plant (8).

Key molecular pathogenesis, virulence, and fitness traits have been identified by studying

model strains like *R. pseudosolanacearum* GMI1000. However, comparative genomics shows

that the *Ralstonia* species complex is genetically heterogeneous (1, 9–11). Average nucleotide

identity (ANI) between strains in the different species ranges between 90.8-92.7% (1), which is

less than the 95% ANI threshold widely used for species borders. An average *Ralstonia* genome

contains approximately 5,000 genes, but approximately 2,900 core gene families are core to the

R. solanacearum species complex (unpublished data from Vinatzer and Lowe-Power labs).

Accordingly, ecological niches, host ranges, and physiological traits vary widely among

Ralstonia strains (1, 12–14). Developing and studying new model strains (15) that capture the

phylogenetic diversity of the *Ralstonia* species complex will provide a more complete view of

the biology of these globally important pathogens.

We hypothesized that genes in diverse *Ralstonia* strains contribute to their growth in

xylem sap of tomato plants, a common host. In this study, we used a RB-TnSeq *mariner*

transposon library with over 10^8 possible barcoded Tns (16) to create genome-wide transposon

insertion mutant libraries of three tomato-pathogenic *Ralstonia* strains: *R. pseudosolanacearum*

GMI1000, *R. solanacearum* IBSBF1503, and *R. syzygii* PSI07. We mapped the transposon

insertion sites and predicted core essential genes. We performed genetic screens to identify

fitness factors that influence competitive growth of strains in *ex vivo* xylem sap from healthy

susceptible tomato, as well as minimal and rich culture media. Comparison of genes' fitness

contributions between conditions revealed sap-specific fitness factors and genes that contribute

to fitness in both sap and in culture media.

Results

Adapting RB-TnSeq for plant pathogenic *Ralstonia* isolates. To identify genes

required for *Ralstonia* fitness, we constructed barcoded transposon insertion mutant libraries in

three *Ralstonia* species complex backgrounds: *R. pseudosolanacearum* GMI1000, *R.*

solanacearum IBSBF1503, and *R. syzygii* PSI07. We selected these isolates because they are all

virulent on tomato and because there are high-quality, complete genomes available for each

strain. These diverse strains share 76-83% of their gene content with average nucleotide

identities of 91-93% (1), which we hypothesized would allow us to identify strain-specific and

conserved fitness factors required by the species complex.

The mutant libraries were created by conjugation with the pKMW3 *E. coli* donor library

(16), which carries a pool of over 10^8 *mariner* transposon delivery vectors in which each

transposon is marked by a unique 20 base pair sequence. These unique sequences function as

DNA "barcodes". After conjugation, we selected and pooled approximately 10^5 transposon

insertion mutants per strain library and mapped the insertion site of each barcoded transposon.

Table 1 shows detailed summary statistics of each mutant library. For barcode sequencing

(BarSeq) mutant fitness assays, we calculate fitness contributions using transposons that had

inserted into the central 80% of the open reading frame (ORF) because insertions at a gene's N-

and C-termini are less likely to impair protein function.

The barcoded mutant libraries cover 79-86% of the protein coding genes of each strain.

There are several reasons why a given gene may be unrepresented in the mutant libraries. The

mariner transposon we used inserts into TA dinucleotide sites. Strains GMI1000, IBSBF1503,

and PSI07 have 48, 42, and 101 genes that could not be mutagenized because they lack TA sites.

Table S1 lists genes that lack fitness data and the number of TA sites in the central 80% of each

gene. Additionally, it is not possible to uniquely map transposon insertions in repetitive,

indistinguishable genomic regions like the 31 kb tandem duplication in the GMI1000 genome.

Furthermore, if these genes contribute to fitness in a condition, they would be functionally

redundant with their identical paralogs, mitigating any potential impact on fitness. Finally, a

subset of the genes without transposon insertions are likely essential or near-essential for growth

in rich medium, resulting in their failure to grow during library construction and exclusion from

our libraries.

Predicting genes essential for *Ralstonia* growth in rich medium. We predicted

essentiality of genes in each strain (Table S1) using the Bio::TraDIS pipeline (17) and our

previously described analysis pipeline (18). Briefly, both methods predict gene essentiality based

on genes that lacked transposon insertions in the central 80% of the coding region. After setting

cut-offs to exclude short genes ("Above length cutoff" in Table S1), which are more likely to

lack transposon insertions, 450-500 genes were predicted to be essential per strain by at least one

pipeline, while 393-396 genes per strain were predicted to be essential by both pipelines (Table

S1). To compare results between strains, we identified orthologous genes in GMI1000, PSI07,

and IBSBF1503 using the JGI-IMG (19) Genome Gene Best Homologs tool. A core set of 244

genes was predicted to be essential by both pipelines in all three strains (Fig S1) (20). We

inspected the predicted essential gene set for a positive control, the *speC* ornithine

decarboxylase. We previously showed that a GMI1000 *speC* mutant (RSc2365) is a putrescine

auxotroph that cannot not grow in rich medium unless it is supplemented with putrescine (8);

both pipelines predicted *speC* was essential in GMI1000 and PSI07. Bio::TraDIS, but not our

own pipeline, predicted *speC* to be essential in IBSBF1503.

To visualize which cellular processes are predicted essential for growth in rich medium,

we used KEGG Mapper (21) to query the KEGG database with the GMI1000 locus tags of the

244 genes predicted to be essential in all three strains. As expected, many of these genes are

involved in central cellular functions, such as DNA replication, transcription, translation, mRNA

degradation, and lipid and peptidoglycan biosynthesis. Several genes from central carbon

metabolism and respiration are predicted to be essential, including components of the Entner-

Doudoroff pathway, the pentose phosphate pathway, the TCA cycle, NADH dehydrogenase, and

ATP synthase. Many genes involved in cofactor and vitamin biosynthesis (ubiquinone,

coenzyme A, thiamine, heme, and folate) are predicted to be essential. Genes that encode

biosynthetic steps of several amino acids, including threonine, arginine, lysine, and histidine,

also appear to be essential for growth in rich medium.

Growth of *Ralstonia* species representatives in xylem sap. To design an RB-TnSeq

experiment with a sufficient number of cell doublings to quantify mutant fitness defects, we

empirically determined the carrying capacity and doubling rate of *Ralstonia* in xylem sap. We

compared growth of *R. pseudosolanacearum* GMI1000 in rich medium, minimal medium, or

xylem sap harvested from greenhouse-grown Moneymaker tomato (Fig S2A-B). The carrying

capacities of xylem sap and rich medium were higher than the carrying capacity of minimal

medium: $3.1\text{-}3.5 \times 10^9$ CFU/ml in sap, $2.4\text{-}2.6 \times 10^9$ CFU/ml in rich medium, and $2.2\text{-}4.6 \times 10^8$

CFU/ml in minimal medium. Both xylem sap and rich medium cultures exceeded 1×10^8 CFU

within 24 hours, while minimal medium cultures had an extended lag time. We were surprised

by the high carrying capacity supported by xylem sap based on our previous experience growing

diverse *Ralstonia* in xylem sap from a different susceptible tomato (Bonny Best) grown in

environmentally controlled growth chambers (8, 22). To determine whether the growing

conditions or plant cultivar influenced the saps' carrying capacity, we collaborated with

researchers at University of Wisconsin—Madison to repeat this experiment with sap from

growth chamber-grown Moneymaker and Bonny Best tomato in all three of our study strains.

Each *Ralstonia* isolate grew to a higher carrying capacity in xylem sap from Moneymaker plants

($3.3\text{-}6.6 \times 10^9$ CFU/ml) than sap from Bonny Best plants ($3.6 \times 10^7\text{-}5.5 \times 10^8$ CFU/ml) (Fig S2C-

E).

To estimate the doubling rate in xylem sap and rich and minimal culture media, we

compared growth of GMI1000, PSI07, and IBSBF1503 with frequent sampling intervals (every 4

hours, Fig 1A). Strains grew slowest in minimal medium and the fastest in xylem sap (Fig 1B).

The maximum doubling rate in xylem sap was 120-131% higher than in rich medium (P<0.005

by ANOVA with Tukey's multiple comparison's test). The higher growth rate and carrying

capacity in xylem sap compared to rich medium indicate that *Ralstonia* strains GMI1000,

IBSBF1503, and PSI07 are well-adapted to growth in xylem sap from Bonny Best tomato plants.

RB-TnSeq experiments identify mutants with altered fitness in xylem sap and

minimal medium. We used RB-TnSeq to identify genes that influence fitness in *ex vivo* xylem

sap, minimal medium, and rich medium. Mutant abundance before and after selective

competitive growth was measured by BarSeq. In BarSeq, each gene is assigned a fitness score

("Fit" score), which is a weighted average of the $\log_2$ ratios of each barcoded mutant's final

abundance divided by its abundance before growth in the selective condition (the "time 0"

abundance) (16). In total, our BarSeq assays measured fitness contributions for approximately

4,000 genes per strain. Multiple independent transposon mutants in the same gene are used to

calculate each gene's Fit score (Table 1). We calculated Fit scores and t-like test statistics for

each gene in each condition (Table S2). Fitness data are also publicly available on the interactive

Fitness Browser (http://fit.genomics.lbl.gov/). Because Fit scores are on a $\log_2$ scale, a score of -

5 shows in principle that those mutants replicated for 5 fewer generations than the population. In

practice, we found that mutants with the strongest fitness defects had Fit scores that exceeded the

number of generations that we measured by dilution plating. For example, GMI1000 *murI*

(RSc1956) had a mean Fit score of –8.6 in xylem sap even though the population grew for only 7

generations. However, most Fit scores were in the expected ranges for each TnSeq trial.

There were trends in the number genes associated with gain or loss of fitness between

growth conditions (Fig 2A-C). There were more mutants with fitness defects in xylem sap

(n=37-55) and minimal medium (n=28-42) than in rich medium (n=8-28) (Fit<-1; t-like statistic

<-2.5 in all trials). Logically, mutants with strong fitness defects in rich medium are excluded

from the libraries because they would not have grown on the selection plates. Interestingly, we

observed that more mutants had gain-of-fitness phenotypes in minimal (4-20) and rich media (2-

19) than in xylem sap (0-1). We next determined which genes influence fitness in a single

condition and which genes have intersecting fitness contributions in two or more conditions (Fig

2D). The intersection analysis reveals that each strain has many genes that contribute to fitness in

xylem sap only, in minimal medium only, or in both xylem sap and minimal medium.

***Ralstonia* require multiple envelope-associated genes for fitness in xylem sap.** Our

intersection analysis identified envelope-associated genes as fitness factors in xylem sap in all

three strains, in addition to other strain-specific xylem sap fitness factors (Fig 3). The strongest

xylem sap fitness factor in all strains was the *murI* gene (RSc1956) with a Fit score of < -5 in all

strains. MurI is a glutamate racemase that converts L-Glutamate to D-Glutamate, which is

incorporated into peptide cross-linkages in peptidoglycan. The *murI* gene is required for multiple

Ralstonia strains to grow in tomato stems (23, 24). For all three study strains, other xylem sap

fitness factors included *tatC* (involved in Tat-dependent protein secretion), *dsbA* which catalyzes

disulfide bonds in periplasmic proteins, and a lipoprotein of unknown function (RSc2940,

RSc0285, and RSc2007, respectively). BLAST analysis reveals that this lipoprotein is present in

genomes of *Ralstonia*, *Cupriavidus*, *Paraburkholderia, Burkholderia*, which are all genera in the

Burkholderiaceae family.

IBSBF1503 requires 13 different LPS genes for full fitness in xylem sap, and several of

these genes are important for fitness in xylem sap in PSI07 and GMI1000 as well (Fig 3B).

Though conserved in all three strains, LPS genes colocalized in LPS cluster 1 had stronger

fitness phenotypes in IBSBF1503 (-4.2 < Fit < -1.1) than in GMI1000 (-0.9 < Fit < -0.3) and

PSI07 (-1.6 < Fit < -0.8). These results corroborate previous work in *Ralstonia* strain Pss4

indicating that mutants with structural defects in LPS also have reduced growth *in planta* (25)

while highlighting subtle interstrain differences in the exact orthologs involved.

Other strain-specific envelope proteins identified in our analysis include the finding that

GMI1000 requires *mdoGH* (RSc2908 and RSc2909) during growth in xylem sap (Fig 3C), which

putatively encode enzymes that synthesize periplasmic branched oligoglucans. Since periplasmic

oligoglucans can act as osmoprotectants to Gram-negative bacteria (26), and *mdoGH* contributes

to the growth of several plant pathogens the host (27), these enzymes may play a role in

protecting GMI1000 from environmental stress posed during growth in xylem sap.

Metabolic functions required for full fitness in xylem sap. Many of the xylem sap

fitness factor genes are associated with bacterial metabolism. Consistent with previous studies

that showed *Ralstonia scrA* sucrose catabolism mutants (RSp1285) have growth defects in

tomato stems (28, 29), *scrB* (RSp1284) sucrose catabolism mutants also had reduced fitness in

xylem sap in our TnSeq experiments. The fitness defect of *scrB* mutants was stronger in the

IBSBF1503 background than in the GMI1000 or PSI07 backgrounds (Fit = -2.0, -1.3, and -0.9,

respectively). Overall, our findings validate that sucrose is an important carbon source in xylem

sap although *Ralstonia* can catabolize other carbon sources present in sap (8, 30, 31).

All strains required amino acid biosynthesis genes for fitness in xylem sap and minimal

medium (Fig 4). Mutants lacking Ser, Cys, Met, Phe, and Trp biosynthesis genes had fitness

defects in both xylem sap and minimal medium. Although all strains required several Glu, Pro,

and branched chain AA (Leu/Ile) biosynthesis genes for growth in minimal medium, only

GMI1000 consistently required these genes for full growth in xylem sap. Our TnSeq results are

consistent with xylem sap metabolomic studies that indicate that multiple tomato cultivars lack

Cys and Trp in sap (8, 31) Although Ser, Met, and Phe are all present in xylem sap, our TnSeq

results indicate that *Ralstonia* requires many *de novo* amino acid synthesis enzymes for full

fitness in xylem sap.

Multiple cofactor biosynthesis mutants had xylem sap fitness defects in one or more

strains (Fig 5). All three strains required the thiamine biosynthesis gene *thiC* for full growth in

xylem sap (Fit -1.0 to -2.4). Similarly, all strains strongly required *purU* (RSc1873) for growth in

xylem sap (Fit -3.9 and -6.0), with weaker fitness defects in minimal medium (Fit -0.3 to -1.4).

PurU is a putative formyltetrahydrofolate deformylase that hydrolyzes 10-formyltetrahydrofolate

into formate and tetrahydrofolatic acid (THFA). Because THFA is a cofactor for biosynthesis of

many amino acids and nucleic acids, it is possible that PurU is necessary for *Ralstonia* to recycle

THFA in order to synthesize amino acids. Interestingly, only one of the three strains required

cobalamin biosynthesis to grow in xylem sap. Eighteen IBSBF1503 cobalamin biosynthesis

mutants had subtle but consistent fitness defects in xylem sap with Fit scores that ranged -0.9 to -

1.4 (Fig 5B).

Mutants with improved fitness. Many mutants with gains-of-fitness phenotypes had

transposon insertions in regulatory genes (Fig 6). Consistent with previous results showing that

the PhcA quorum sensing regulator mediates a trade-off between fast growth at low cell density

and expression of *in planta* fitness traits at high cell density (32, 33), GMI1000 and IBSBF1503

phcA mutants had improved growth in all tested conditions (Fig 6A). Unexpectedly, PSI07 *phcA*

mutants behaved differently. Although PSI07 *phcA* mutants had increased fitness in minimal

medium (Fit: +1.9), they had strongly reduced growth in xylem sap (Fit: -2.8). Contrasting

phenotypes of *phcA* mutants in different wildtype backgrounds highlights the genetic and

regulatory diversity within plant-pathogenic *Ralstonia.* At high cell densities, PhcA positively

regulates production of an energetically costly extracellular polysaccharide (EPS) among other

traits (33, 34). Transposon insertion in other regulatory genes that activate expression of *eps* and

other virulence genes (*phcBSR*, *vsrAD*, and *rpoS*) (35, 36) also increased fitness in one or more

strains (Fig 6A).

Strain PSI07 had the most mutants with gain-of-fitness phenotypes in culture media. Four

of these genes were associated with the type IV pilus, and another with a flagella-associated

chaperone, FlgN (Fig 6B). Three IBSBF1503 type IV pilus mutants showed the same trend as

PSI07, but with weaker phenotypes. This result suggests that these wildtype strains may express

energy-consuming motility systems in liquid rich medium but not in sap or minimal medium.

PSI07 had additional regulators that had gain-of-fitness phenotypes in culture media: a MarR-

family transcriptional regulator nested in a hydroxycinnamic acid degradation cluster

(RSp0224/RPSI07_RS00780) (13), a transcriptional regulator co-localized with an ABC

transport system (RSp1579/ RPSI07_RS07405) (37), a TetR family transcriptional regulator

(RSc3207/ RPSI07_RS15470), a GGDEF/EAL domain-encoding gene

(RPSI07_RS05560/RSp1208), and the RelA (p)ppGpp synthase (RPSI07_RS15685/RSc1576).

Discussion

Why might envelope genes contribute to *Ralstonia* fitness in *ex vivo* xylem sap? Recent

studies have shown that clinical antibiotic use can select for pathogens that remodel their

envelopes, enabling them to exclude and resist antibiotics (38). Because many envelope-

associated genes were xylem sap fitness factors, we hypothesize that xylem sap contains

preformed antimicrobial chemicals and proteins. Future studies should test the hypothesis that

periplasmic branched oligoglucans, LPS, and other envelop genes protect *Ralstonia* from *in*

planta stresses like reactive oxygen species and antimicrobial peptides (38).

There are benefits and limitations of predicting gene essentiality using transposon

mutagenesis. Here we identified sets of 454-487 essential genes in three *Ralstonia* strains, of

which more than half (n=244) were conserved in all three strains and predicted essential by both

analysis pipelines. Unsurprisingly, many of the essential genes are involved in carbon

metabolism, amino acid and cofactor biosynthesis, and central dogma processes. Identifying

absolutely essential genes of human pathogens aids in antibacterial drug development research

(39), but precision antimicrobial treatment for most plant pathogens is not financially viable and

poses environmental concerns. Nonetheless, identifying putative essential genes has fundamental

value, particularly to the *Ralstonia* research community. Creating mutants lacking these genes

might be impossible or require specialized selection media as was required for the Δ*speC::*Sm

mutant that had a putrescine auxotrophy (8). The essentiality of multiple amino acid biosynthesis

genes in the peptone- and casamino acid-containing CPG rich medium suggests *Ralstonia* may

lack effective transporters for some amino acids.

Recently, Su *et al.* predicted 464 essential genes in *Ralstonia* GMI1000 (40) using a

TnSeq approach, but our libraries contain mutants with transposon insertions in 126 of these

genes. Of these 126 genes, only three genes contributed to fitness more than two-fold (Fit = -1)

in any of our experimental conditions, including rich medium. These discrepancies highlight the

limitations of using randomly generated mutant libraries to predict gene essentiality. Targeted

knockdown experiments using techniques like CRISPRi (41) could test essentiality more

robustly. In a subsequent TnSeq experiment using their mutant library in living tomato plant

hosts, Su *et al.* identified 131 genes as being important for *in planta* fitness (42). Of the 36 genes

we identified as important for xylem sap fitness (average | Fit | > 1, | t | > 2.5 in three trials), Su *et*

al. identified 16 as also being important for *in planta* fitness. This overlap suggests that

Ralstonia experiences some of the same selective pressures in *ex vivo* xylem sap as it does in

living tomato hosts, though additional genes are necessary for full fitness in the more complex *in*

planta environment.

RB-TnSeq reveals inter-strain differences in fitness factors. Although model strains such

as GMI1000 are well-characterized and extensively studied, there are known physiological and

genetic differences between strains (1, 28, 43). To more fully capture the diversity of *Ralstonia*,

we constructed barcoded transposon mutant libraries in strains from each of the three plant-

pathogenic *Ralstonia* species: *R. solanacearum* (GMI1000), *R. solanacearum* (IBSBF1503), and

R. syzygii (PSI07). While RB-TnSeq found specific metabolism- and envelope-related fitness

factors in all three strains, the orthologs found to be fitness factors varied from strains to strain.

For instance, GMI1000 requires *mdoGH* genes for periplasmic oligoglucan production in xylem

sap, while these genes do not produce strong fitness phenotypes in the other two strains.

Additionally, several amino acid biosynthesis genes were required by some by not all strains

when grown in minimal media or xylem sap, including *cysU*, *leuB1*, *leuC*, *gltB*, and *proB* (Fig

4). These strain-to-strain differences in gene fitness profiles reveal subtle differences in the

genetics that underlie the ability of all these strains to robustly grow in tomato xylem sap.

Considering our results in *ex vivo* xylem sap, different genes may be important for virulence of

these diverse strains in living hosts as well.

Novel environments provide strong selective pressures for bacterial evolution (44). We

suggest that the number of gain-of-fitness mutations in a TnSeq screen may be an indicator of

how closely the test conditions match the environment to which the bacterial isolate is adapted.

For instance, transposon insertions in over two dozen *E. coli* genes increased its growth on

cheese agar (45), a foreign growth medium. Similarly, serial passage of *Ralstonia* GMI1000 in

Medicago nodules (a foreign environment) selected for more gain-of-fitness mutations than

serial passage in tomato stems (46, 47). In our study, few mutations increased fitness in sap, but

many mutants increased fitness in artificial culture media, suggesting that the *Ralstonia* isolates

are well-adapted to grow in tomato xylem sap. We observed that the model strain *Ralstonia*

GMI1000 had fewer gain-of-function mutations in culture media than the less-studied

IBSBF1503 and PSI07, suggesting that GMI1000 may be more domesticated and adapted to

culture media since its isolation in 1978. Meta-analyses of TnSeq studies across multiple bacteria

in novel and naturalistic conditions can test the hypothesis that gain-of-fitness phenotypes are

more common when mutants grow in novel conditions.

Traditional transposon mutagenesis genetic screens pioneered molecular plant pathology

and identified core virulence traits like the type III secretion system (48–50). TnSeq approaches

leverage next-generation sequencing to rapidly quantify mutant phenotypes. These genome-wide

fitness assays are a powerful approach to rapidly investigate basic bacterial biology and identify

pathogen, commensal, and mutualist fitness factors. TnSeq studies have identified genes that

promote bacterial fitness in their native environments including plant hosts (51–57) animal hosts

(58–60), soft-rind cheese (45), and lake water (61). The RB-TnSeq technique that we use here is

a powerful TnSeq methodology that has a low cost and technical barrier per sample, which

facilitates profiling mutant fitness across in multiple strain backgrounds across many conditions

(16, 62). Future studies will profile fitness of *Ralstonia* mutants *in planta* to test whether sap

fitness factors also contribute to fitness in living hosts.

Methods:

Bacterial Strains and Growth Conditions

This study uses three *Ralstonia* strains, one representative per plant pathogenic species: *R. pseudosolanacearum* GMI1000 (phylotype I sequevar 18), *R. syzygii* PSI07 (phylotype IV sequevar 10), and *R. solanacearum* IBSBF1503 (phylotype IIB sequevar 4). All isolates are pathogenic on tomato (11).

Ralstonia was routinely grown in CPG rich medium (per L: 1 g casamino acids, 10 g Bacto-peptone, 5 g glucose, 1 g yeast extract) at 28°C. Agar plates were supplemented with 1% tetrazolium chloride to confirm colony morphology. For minimal medium growth curves and fitness assays, strains were grown in quarter-strength M63 (per L: 3.4 g KH_2PO_4, 0.5 g $(NH_4)_2SO_4$, 10 μl of 1.25 mg/ml $FeSO_4 \cdot 7H_2O$, 51.7 μl of 1 M $MgSO_4$; adjusted to pH 7) with 10 mM glucose.

Construction of barcoded transposon mutant libraries

Barcoded *mariner* transposon mutant libraries were created in three wild-type strain backgrounds: GMI1000, PSI07, and IBSBF1503. The barcoded transposons were introduced via conjugation. Recipient strains (*Ralstonia* isolates) were grown overnight in 5 ml CPG at 28°C with shaking at 200 rpm. A 2 ml portion of the overnight culture was sub-cultured into 25 ml CPG and grown for 2 hours at 28°C. The donor *E. coli* strain (WM3064 background, a synthetic auxotroph requiring 300 μM diaminopimenlate [DAP]) carrying the pKMW3 (Kan[R]) barcoded *Mariner* transposon vector library (16) was thawed on ice and 1 ml was inoculated into 20 ml LB with 25 μg/ml kanamycin and 300 μM DAP and grown to mid-log phase at 37°C with shaking. Conjugations were carried out as previously described (63). Donor and recipient strains were centrifuged at 5000 x*g* for 10 min and resuspended in CPG + 300 μM DAP. Recipient strains and

the donor strain were adjusted to OD_{600} of 3.0 and 1.0, respectively, and mixed in equal volumes.

In total, 1.2 ml of the strain mixture was spotted onto 12 nitrocellulose filters (82 mm discs with

0.45 μM pore size cut into sixths) overlaid on four plates of CPG with 300 μM DAP and without

glucose. Mating was allowed to occur for 3-4 hours at 28°C. Filters were pooled in 20 ml CPG

broth and vortexed in a 50 ml tube for 30 s to dislodge bacteria. The suspension was adjusted to

80 ml and evenly spread over 400 CPG plates (200 μl per plate) with 25 μg/ml kanamycin. To

calculate transformation efficiencies, a portion of each suspension was dilution plated on CPG

with kanamycin (to quantify transformants) and without kanamycin (to quantify total *Ralstonia*

cells). Transformation efficiencies were: GMI1000, 3.9×10^{-5}; PSI07, 8.6×10^{-3}; IBSBF1503,

1.5×10^{-4}. Selection plates were incubated for 2 days at 28°C. Cells were harvested by scraping

and pooling approximately 2×10^{6} colonies in CPG broth with kanamycin. Density was adjusted

to $OD_{600} = 0.25$ in 200 ml and grown at 28°C with shaking until OD_{600} reached 1.0. Individual 1

ml aliquots were preserved in 20% v/v glycerol (final concentration) at -80°C.

Mapping transposon insertion sites.

We used a previously described protocol for mapping the transposon insertion locations

and to link these mutants to their unique DNA barcode sequences (16). Full details describing

this protocol are available (16). Briefly, genomic DNA was extracted using the Qiagen DNeasy

Blood and Tissue Kit, with RNase A treatment, according to the manufacturer's

recommendations. The DNA was sheared to ~300 bp using a Covaris S220, and the DNA was

repaired and A-tailed using the NEBNext DNA Library preparation kit for Illumina (New

England Biolabs). After ligating Illumina compatible adapters to the DNA fragments, we PCR

amplified the transposon insertion junctions using primers with Illumina P5 and P7 sequences.

The final PCR product was purified using AMPure XP beads and assessed for quantity and size

using an Agilent Bioanalyzer DNA1000 chip. Samples were sequenced on a HiSeq2500

instrument at the QB3 Berkeley Genomics Center using 2 x 150 reads. TnSeq reads were

analyzed with a custom Perl script, MapTnSeq.pl, which assigns each unique barcode sequence

to a corresponding location in the genome. Each barcode was mapped to a single insertion site by

identifying barcodes that consistently map to a unique location in the genome using

DesignRandomPool.pl. All scripts used are available at

https://bitbucket.org/berkeleylab/feba/src/master/bin/ (16, 62).

Essential gene calculations

Based on the TnSeq data, we used standard computational methods (62) to predict which

genes are likely essential for growth in CPG. Briefly, this analysis predicts gene essentiality

based on genes that lacked transposon insertions in the central 80% of the coding region. We

excluded two categories of genes from the essentiality calculation. Genes sharing high nucleotide

identity with other genes in the genome (measured via BLAT) (64) were excluded because we

could not map transposon insertions to highly similar genes. Second, we excluded genes below a

minimum length because they are less likely to be disrupted. Briefly, this cutoff is determined

using our previously described analysis pipeline (18) using saturation of *mariner* transposon

insertions to calculate the length at which genes have a 1% chance of being missed due to

random chance. Due to differences in insertion mutant coverage depth between the transposon

libraries, the minimum gene length considered was 450 bp for GMI1000, 325 bp for PSI07, and

400 bp for IBSBF1503. For comparison, we also used the Bio-Tradis pipeline

(github.com/sanger-pathogens/Bio-Tradis) (17) to predict essential genes in these strains, using

the same minimum gene length cutoffs and excluding insertions in the first and last 10% of the

coding region from analysis.

Orthologous gene predictions

412   Orthologous genes were predicted using the "Genome Gene Best Homologs" tool from

413 the JGI IMG database (19). We searched for homologs in PSI07 and IBSBF1503 against

414 GMI1000 as the reference genome using the JGI IMG default identity threshold of 60%. The

415 results, originally reported with IMG locus tags, were matched to the corresponding NCBI locus

416 tags by gene sequence.

Tomato xylem sap harvesting

Xylem sap was harvested from susceptible tomato plants (cv. Moneymaker and Bonny

Best). Most experiments were performed with sap from Moneymaker tomato plants grown in the

Oxford Tract Greenhouse in Berkeley, CA. Bacterial growth curves were independently

replicated with sap from tomato plants grown in a growth chamber (28°C with a 12 hr day/night

cycle) at UW-Madison.

Xylem sap was harvested by de-topping each plant at the cotyledon juncture with a razor

blade and allowing sap to pool on the stump (65). To reduce cytoplasmic contamination, the sap

that accumulated in the first few minutes was discarded; the stump was washed with diH_2O and

gently blotted dry. Sap was then pipetted and pooled into a 50 ml conical over a collection period

of 3 hours. Pooled sap was centrifuged at 5,000 *xg* RT for 10 minutes and the supernatant was

filter sterilized using filters with 0.22 μm pores (Thermo Scientific #725-2520). Sap was

aliquoted and stored at -20°C until use. Each batched pool of xylem sap was collected from

approximately 70 plants.

***R. solanacearum* growth in tomato xylem sap**

Colonies of each strain were inoculated into 6 ml of CPG and grown overnight at 28°C

with shaking at 200 rpm for a total of three biological replicates. Cells from stationary phase

cultures were pelleted at 13,000 xg and washed twice in 1 ml of ddH$_2$O. The washed pellet was

resuspended in ddH$_2$O and adjusted to a final OD$_{600}$ of 0.02. Each growth condition was

inoculated with 7 μl of each culture at a starting cell density of ~10^5 CFU/ml in 48-well plates

(Corning #353078), for a total of two technical replicates per biological replicate. Growth was

measured by dilution plating at 0-, 4-, 8-, 24-, 32-, and 48-hour time points. In parallel,

measurements for 12-, 16-, and 20-hour time points were taken from independent cultures that

were started 12 hours after the first, using the same cultures, growth media, and technique

described above.

Fitness experiments with transposon libraries

For genome wide fitness experiments, we adapted established methods (16). Per

condition (minimal medium, rich medium, or Moneymaker tomato xylem sap), three 1 ml

aliquots of each transposon library (GMI1000, PSI07, and IBSBF1503) were thawed on ice and

revived in separate 100 ml flasks of CPG with 12.5 mg/ml Kanamycin with shaking incubation

at 28°C for 16-20 hours. Once cultures reached an OD$_{600}$ of 0.2 to 0.5, cells were pelleted by

centrifuging for 10 min at 5000 xg RT and resuspended in 2 ml of rich or minimal medium or

sterile H$_2$O (xylem sap experiments). Cells were washed 3 times in media or xylem sap. An

aliquot of the washed cells (>10^9 cells) was retained (pelleted and frozen) for a "time 0" control.

For rich and minimal medium experiments, cells were seeded into 5 ml at OD$_{600}$ = 0.02 (~2-

5×10^7 total cells). Rich medium cultures were grown to saturation (20-24 hours; 5-6 cell

doublings) and minimal medium cultures were grown for over 48 hours (5-6 cell doublings for

PSI07 and GMI1000 and 4 doublings for IBSBF1503). For xylem sap experiments, cells were

resuspended in 20 ml of xylem sap at a starting cell density of 10^6 cells/ml (2×10^7 cells total)

and grown for 25 hours (7-8 doublings for GMI1000 and 8-9 doublings for PSI07). Genomic

DNA was extracted (Qiagen DNeasy Blood and Tissue Kit) from the "time 0" and cells

harvested after growing in differential media.

BarSeq and Fitness Calculations

Prior to PCR, the quality if total genomic DNA isolated from fitness experiments was

assessed on an agarose gel, while DNA concentration was determined by NanoDrop. Barcodes

were PCR amplified from each sample using the previously described reaction protocol (16). For

most experiments, we used indexed P2 oligos and a mix of non-indexed P1 oligos of variable

length (2-5 bases) to stagger the reads. For some experiments, we used indexed versions of both

the P1 and P2 oligos to minimize incorrectly assigned indexes in Illumina HiSeq4000 runs.

Following PCR, amplicon mixtures from each sample were pooled in equal volumes, purified

over a Zymo DNA Clean and Concentrator column, and eluted with water. Prior to sequencing,

the quality of cleaned and concentrated amplicon pools was assessed using a Bioanalyzer. qPCR

was used to determine how much of each sample to load for sequencing. Barcode amplicons

were sequenced on a HiSeq4000 at QB3 Berkeley Genomics Center with 96 samples

multiplexed (50 base pair reads, single end).

Sequencing data was analyzed using the BarSeq pipeline, available at

https://bitbucket.org/berkeleylab/feba/src/master/bin/. For each competitive fitness assay, fitness

score ("Fit scores") for each gene were calculated as the $\log_2$ ratio between barcode abundance

after outgrowth in a condition vs. its abundance in the time 0 sample. Each Fit score is the

weighted average of fitness values for all mutant strains with transposon insertions in a given

gene. These Fit scores are normalized across the genome such that a gene with neither a fitness

cost nor benefit has a value of 0. Significance was determined based on an absolute t-like test

statistic with a threshold of $|t| > 2.5$ in all three RB-TnSeq trials.. The t-like test statistic

considers the consistency of the fitness scores for all barcoded mutants for each gene in the

experiment as previously described in detail (16).

Data availability.

The raw reads used for TnSeq mapping is available in the NCBI SRA under accession

PRJNA629015. The fitness browser (http://fit.genomics.lbl.gov) offers a graphical user interface

for exploring the fitness data from these experiments, orthology of the genes between these

strains and other Gram negative bacteria, and cross-references to Kegg, Paperblast, and NCBI

databases.

Acknowledgements

We thank Steve Lindow, Caitilyn Allen, and Jeff Flynn for useful discussions. This work

used the Vincent J. Coates Genomics Sequencing Laboratory at UC Berkeley, supported by NIH

S10 OD018174 Instrumentation Grant. Oxford Tract Greenhouse Staff and the QB3 Berkeley

Genomics Center provided technical assistance.

This work was supported by USDA NIFA #2018-67012-31497 awarded to TLP; partial

support was provided by a UC Berkeley SURF Rose Hills Independent Fellowship awarded to

KES. The funders had no role in study design, data collection and interpretation, or the decision

to submit the work for publication.

Mention of trade names or commercial products in this publication is solely for the

purpose of providing specific information and does not imply recommendation or endorsement

by the U.S. Department of Agriculture. USDA is an equal opportunity provider and employer.

References

1.   Prior P, Ailloud F, Dalsing BL, Remenant B, Sanchez B, Allen C. 2016. Genomic and

proteomic evidence supporting the division of the plant pathogen *Ralstonia solanacearum*

into three species. BMC Genomics 17:1–11.

2.   Safni I, Cleenwerck I, De Vos P, Fegan M, Sly L, Kappler U. 2014. Polyphasic taxonomic

revision of the *Ralstonia solanacearum* species complex. Int J Syst Evol Microbiol

64:3087–3103.

3.   Hayward AC. 1991. Biology and epidemiology of bacterial wilt caused by *Pseudomonas*

solanacearum. Annu rev Phytopathol 29:65–87.

4.   Safni I, Subandiyah S, Fegan M. 2018. Ecology, epidemiology and disease management

of *Ralstonia syzygii* in Indonesia. Front Microbiol 9:1–11.

5.   Lopes CA, Rossato M. 2018. History and status of selected hosts of the *Ralstonia*

solanacearum species complex causing bacterial wilt in Brazil. Front Microbiol 9:1–6.

6.   Jiang G, Wei Z, Xu J, Chen H, Zhang Y, She X, Macho AP, Ding W, Liao B. 2017.

Bacterial wilt in China: history, current status, and future perspectives. Front Plant Sci

8:1–10.

7.   Lowe-Power T, Chipman K. 2020. A meta-analysis of the known global distribution and

host range of the *Ralstonia* species complex. bioRxiv 1–7.

8.   Lowe-Power TM, Hendrich CG, von Roepenack-Lahaye E, Li B, Wu D, Mitra R, Dalsing

BL, Ricca P, Naidoo J, Cook D, Jancewicz A, Masson P, Thomma B, Lahaye T, Michael

AJ, Allen C. 2018. Metabolomics of tomato xylem sap during bacterial wilt reveals

Ralstonia solanacearum produces abundant putrescine, a metabolite that accelerates wilt

disease. Environ Microbiol 20:1330–1349.

9.   Remenant B, Coupat-Goutaland B, Guidot A, Cellier G, Wicker E, Allen C, Fegan M,

Pruvost O, Elbaz M, Calteau A, Salvignol G, Mornico D, Mangenot S, Barbe V, Médigue

C, Prior P. 2010. Genomes of three tomato pathogens within the *Ralstonia solanacearum*

species complex reveal significant evolutionary divergence. BMC Genomics 11:1–16.

10.   Remenant B, de Cambiaire JC, Cellier G, Jacobs JM, Mangenot S, Barbe V, Lajus A,

Vallenet D, Medigue C, Fegan M, Allen C, Prior P. 2011. *Ralstonia syzygii*, the Blood

Disease bacterium and some Asian *R. solanacearum* strains form a single genomic species

despite divergent lifestyles. PLoS One 6:1–10.

11.   Ailloud F, Lowe T, Cellier G, Roche D, Allen C, Prior P. 2015. Comparative genomic

analysis of *Ralstonia solanacearum* reveals candidate genes for host specificity. BMC

Genomics 16:1–11.

12.   Hayward AC. 1964. Characteristics of *Pseudomonas solanacearum*. J Appl Bacteriol

27:265–277.

13.   Lowe TM, Ailloud F, Allen C. 2015. Hydroxycinnamic acid degradation, a broadly

conserved trait, protects *Ralstonia solanacearum* from chemical plant defenses and

contributes to root colonization and virulence. Mol Plant-Microbe Interact 28:286–297.

14.   Lowe-Power TM, Jacobs JM, Ailloud F, Fochs B, Prior P, Allen C. 2016. Degradation of

the plant defense signal salicylic acid protects *Ralstonia solanacearum* from toxicity and

enhances virulence on tobacco. MBio 7:1–12.

15.   Liu H, Deutschbauer AM. 2018. Rapidly moving new bacteria to model-organism status.

Curr Opin Biotechnol 51:116–122.

16.   Wetmore KM, Price MN, Waters RJ, Lamson JS, He J, Hoover CA, Blow MJ, Bristow J,

Butland G, Arkin AP, Deutschbauer A. 2015. Rapid quantification of mutant fitness in

diverse bacteria by sequencing randomly bar-coded transposons. MBio 6:1–15.

17.   Barquist L, Mayho M, Cummins C, Cain AK, Boinett CJ, Page AJ, Langridge GC, Quail

MA, Keane JA, Parkhill J. 2016. The TraDIS toolkit: sequencing and analysis for dense

transposon mutant libraries. Bioinformatics 32:1109–1111.

18.   Rubin BE, Wetmore KM, Price MN, Diamond S, Shultzaberger RK, Lowe LC, Curtin G,

Arkin AP, Deutschbauer A, Golden SS. 2015. The essential gene set of a photosynthetic

organism. Proc Natl Acad Sci U S A 112:e6634–e6643.

19.   Chen I-MA, Chu K, Palaniappan K, Pillay M, Ratner A, Huang J, Huntemann M,

Varghese N, White JR, Seshadri R, Smirnova T, Kirton E, Jungbluth SP, Woyke T, Eloe-

Fadrosh EA, Ivanova NN, Kyrpides NC. 2019. IMG/M v.5.0: an integrated data

management and comparative analysis system for microbial genomes and microbiomes.

Nucleic Acids Res 47:D666–D677.

20.   Hulsen T, Vlieg J De, Alkema W. 2008. BioVenn–a web application for the comparison

and visualization of biological lists using area-proportional Venn diagrams 9:1–6.

21.   Kanehisa M, Goto S. 2000. KEGG: Kyoto Encyclopedia of Genes and Genomes. Nucleic

Acids Res 28:27–30.

22.   Dalsing BL, Allen C. 2014. Nitrate assimilation contributes to *Ralstonia solanacearum*

root attachment, stem colonization, and virulence. J Bacteriol 196:949–960.

23.   Choi K, Son GJ, Ahmad S, Lee SY, Lee HJ, Lee S-W. 2020. Contribution of the *murI*

gene encoding glutamate racemase in the motility and virulence of *Ralstonia*

solanacearum. Plant Pathol J 36:355–363.

24.   Lin YM, Chou IC, Wang JF, Ho FI, Chu YJ, Huang PC, Lu DK, Shen HL, Elbaz M,

Huang SM, Cheng CP. 2008. Transposon mutagenesis reveals differential pathogenesis of

Ralstonia solanacearum on tomato and *Arabidopsis*. Mol Plant-Microbe Interact 21:1261–

1270.

25.   Li CH, Wang KC, Hong YH, Chu TH, Chu YJ, Chou IC, Lu DK, Chen CY, Yang WC,

Lin YM, Cheng CP. 2014. Roles of different forms of lipopolysaccharides in *Ralstonia*

solanacearum pathogenesis. Mol Plant-Microbe Interact 27:471–478.

26.   Bohin JP. 2000. Osmoregulated periplasmic glucans in Proteobacteria. FEMS Microbiol

Lett 186:11–19.

27.   Bontemps-Gallo S, Bohin J-P, Lacroix J-M. 2017. Osmoregulated periplasmic glucans.

EcoSal Plus 7.

28.   Jacobs JM, Babujee L, Meng F, Milling A, Allen C. 2012. The *in planta* transcriptome of

Ralstonia solanacearum: conserved physiological and virulence strategies during bacterial

wilt of tomato. MBio 3:1–11.

29.   Hamilton CD, Steidl O, MacIntyre AM, Allen C. 2021. *Ralstonia solanacearum* depends

on catabolism of myo-inositol, sucrose, and trehalose for virulence in an infection stage-

dependent manner. Mol Plant-Microbe Interact 1–18.

30.   Xian L, Yu G, Wei Y, Li Y, Zhang H, Xue H, Morcillo RJL, Macho AP. 2020. A bacterial

effector protein hijacks plant metabolism to support bacterial nutrition. Cell Host Microbe

28:1–10.

31.   Gerlin L, Escourrou A, Cassan C, Maviane Macia F, Peeters N, Genin S, Baroukh C.

2021. Unravelling physiological signatures of tomato bacterial wilt and xylem metabolites

exploited by *Ralstonia solanacearum*. Environ Microbiol https://doi.org/10.1111/1462-

2920.15535.

32.   Khokhani D, Lowe-Power TM, Tran TM, Allen C. 2017. A single regulator mediates

strategic switching between attachment/spread and growth/virulence in the plant pathogen

Ralstonia solanacearum. MBio 8:1–20.

33.   Peyraud R, Cottret L, Marmiesse L, Gouzy J, Genin S. 2016. A resource allocation trade-

off between virulence and proliferation drives metabolic versatility in the plant pathogen

Ralstonia solanacearum. PLOS Pathog 12:1–25.

34.   Brumbley SM, Denny TP. 1990. Cloning of wild-type *Pseudomonas solanacearum phcA*,

a gene that when mutated alters expression of multiple traits that contribute to virulence. J

Bacteriol 172:5677–5685.

35.   Huang J, Carney BF, Denny TP, Weissinger AK, Schell MA. 1995. A complex network

regulates expression of *eps* and other virulence genes of *Pseudomonas solanacearum*. J

Bacteriol 177:1259–1267.

36.   Flavier AB, Schell MA, Denny TP. 1998. An RpoS (σS) homologue regulates

acylhomoserine lactone-dependent autoinduction in *Ralstonia solanacearum*. Mol

Microbiol 28:475–486.

37.   Brown DG, Allen C. 2004. *Ralstonia solanacearum* genes induced during growth in

tomato: an inside view of bacterial wilt. Mol Microbiol 53:1641–1660.

38.   Powers MJ, Trent MS. 2018. Phospholipid retention in the absence of asymmetry

strengthens the outer membrane permeability barrier to last-resort antibiotics. Proc Natl

Acad Sci U S A 115:E8518–E8527.

39.   Poulsen BE, Yang R, Clatworthy AE, White T, Osmulski SJ, Li L, Penaranda C, Lander

ES, Shoresh N, Hung DT. 2019. Defining the core essential genome of *Pseudomonas*

aeruginosa. Proc Natl Acad Sci U S A 116:10072–10080.

40.   Su Y, Xu Y, Li Q, Yuan G, Zheng D. 2020. The essential genome of *Ralstonia*

solanacearum. Microbiol Res 238:1–6.

41.   Larson MH, Gilbert LA, Wang X, Lim WA, Weissman JS, Qi LS. 2013. CRISPR

interference (CRISPRi) for sequence-specific control of gene expression. Nat Protoc

8:2180–2196.

42.   Su Y, Xu Y, Liang H, Yuan G, Wu X, Zheng D. 2021. Genome-wide identification of

Ralstonia solanacearum genes required for survival in tomato plants. mSystems 6.

43.   Kai K, Ohnishi H, Shimatani M, Ishikawa S, Mori Y, Kiba A, Ohnishi K, Tabuchi M,

Hikichi Y. 2015. Methyl 3-hydroxymyristate, a diffusible signal mediating *phc* quorum

sensing in *Ralstonia solanacearum*. ChemBioChem 16:2309–2318.

44.   Lenski RE, Rose MR, Simpson SC, Tadler SC. 1991. Long-term experimental evolution

in *Escherichia coli*. I . Adaptation and divergence during 2,000 generations. Am Nat

138:1315–1341.

45.   Morin M, Pierce EC, Dutton RJ. 2018. Changes in the genetic requirements for microbial

interactions with increasing community complexity. Elife 7:1–26.

46.   Capela D, Marchetti M, Clérissi C, Perrier A, Guetta D, Gris C, Valls M, Jauneau A,

Cruveiller S, Rocha EPC, Masson-Boivin C. 2017. Recruitment of a lineage-specific

virulence regulatory pathway promotes intracellular infection by a plant pathogen

experimentally evolved into a legume symbiont. Mol Biol Evol 34:2503–2521.

47.   Guidot A, Jiang W, Ferdy J, Thébaud C, Barberis P, Gouzy J, Genin S. 2014. Multihost

experimental evolution of the pathogen *Ralstonia solanacearum* unveils genes involved in

adaptation to plants. Mol Biol Evol 31:2913–2928.

48.   Arlat M, Gough CL, Zischek C, Barberis PA, Trigalet A, Boucher CA. 1992.

Transcriptional organization and expression of the large *hrp* gene cluster of *Pseudomonas*

solanacearum. Mol Plant-Microbe Interact 5:187–193.

49. Boucher CA, Barberis PA, Trigalet APH, Demery DA. 1985. Transposon mutagenesis of *Pseudomonas solanacearum*: isolation of Tn5-induced avirulent mutants. J Gen Microbiol 131:2449–2457.

50. Staskawicz BJ, Dahlbeck D, Keent NT. 1984. Cloned avirulence gene of *Pseudomonas syringae* pv. *glycinea* determines race-specific incompatibility on *Glycine max* (L.) Merr. PNAS 81:6024–6028.

51. Cole BJ, Feltcher ME, Waters RJ, Wetmore KM, Mucyn TS, Ryan EM, Wang G, Ul-Hasan S, McDonald M, Yoshikuni Y, Malmstrom RR, Deutschbauer AM, Dangl JL, Visel A. 2017. Genome-wide identification of bacterial plant colonization genes. PLoS Biol 15:1–24.

52. Duong DA, Jensen R V., Stevens AM. 2018. Discovery of *Pantoea stewartii* ssp. *stewartii* genes important for survival in corn xylem through a Tn-Seq analysis. Mol Plant Pathol 19:1929–1941.

53. Gonzalez-Mula A, Lachat J, Mathias L, Naquin D, Lamouche F, Mergaert P, Faure D. 2019. The biotroph *Agrobacterium tumefaciens* thrives in tumors by exploiting a wide spectrum of plant host metabolites. New Phytol 222:455–467.

54. Helmann TC, Deutschbauer AM, Lindow SE. 2019. Genome-wide identification of *Pseudomonas syringae* genes required for fitness during colonization of the leaf surface and apoplast. PNAS 116:18900–18910.

55. Helmann TC, Ongsarte CL, Lam J, Deutschbauer AM, Lindow SE. 2019. Genome-wide transposon screen of a *Pseudomonas syringae mexB* mutant reveals the substrates of efflux transporters. MBio 10:1–15.

56.   Liu Z, Beskrovnaya P, Melnyk RA, Hossain SS, Khorasani S, O'Sullivan LR, Wiesmann

CL, Bush J, Richard JD, Haney CH. 2018. A genome-wide screen identifies genes in

rhizosphere-associated *Pseudomonas* required to evade plant defenses. MBio 9:1–17.

57.   Royet K, Parisot N, Rodrigue A, Gueguen E, Condemine G. 2019. Identification by Tn-

seq of *Dickeya dadantii* genes required for survival in chicory plants. Mol Plant Pathol

20:287–306.

58.   Brooks, John F I, Gyllborg MC, Cronin DC, Quillin SJ, Mallama CA, Foxall R, Whistler

C, Goodman AL, Mandel MJ. 2014. Global discovery of colonization determinants in the

squid symbiont *Vibrio fischeri*. PNAS 111:17284–17289.

59.   Hava DL, Camilli A. 2002. Large-scale identification of serotype 4 *Streptococcus*

pneumoniae virulence factors. Mol Mic 45:1389–1406.

60.   Lewin GR, Stacy A, Michie KL, Lamont RJ, Whiteley M. 2019. Large-scale identification

of pathogen essential genes during coinfection with sympatric and allopatric microbes.

PNAS 116:19685–19694.

61.   Hentchel KL, Ruiz LMR, Curtis PD, Fiebig A, Coleman ML, Crosson S. 2019. Genome-

scale fitness profile of *Caulobacter crescentus* grown in natural freshwater. ISME J

13:523–536.

62.   Price MN, Wetmore KM, Waters RJ, Callaghan M, Ray J, Liu H, Keuhl J V, Melnyk RA,

Lamson JS, Suh Y, Carlson HK, Esquivel Z, Sadeeshkumar H, Chakraborty R, Zane GM,

Rubin BE, Wall JD, Visel A, Bristow J, Blow MJ, Arkin AP, Deutschbauer AM. 2018.

Mutant phenotypes for thousands of bacterial genes of unknown function. Nature

557:503–509.

63.   Perrier A, Barberis P, Genin S. 2018. Introduction of genetic material in *Ralstonia*

solanacearum through natural transformation and conjugation, p. 201–207. *In* Methods in

Molecular Biology. Humana Press, New York, NY.

64.   Kent WJ. 2002. BLAT—the BLAST-like alignment tool. Genome Res 12:656–664.

65.   Khokhani D, Tran TM, Lowe-Power TM, Allen C. 2018. Plant assays for quantifying

Ralstonia solanacearum virulence. Bio-protocol 8:1–19.

66.   Lex A, Gehlenborg N, Strobelt H, Vuillemot R, Pfister H. 2014. UpSet□: visualization of

intersecting sets. IEEE Trans Vis Comput Graph 20:1983–1992.

Figures and Tables

Table 1: Properties of barcoded *mariner* mutant libraries

Mutant Library	Total genes in genome	Genes with Fit scores: # (% total)	Library Size[a]: # mutants	Mutants per gene in fitness calculations: Median (Mean)	Estimated Essential Genes[b]: # (%)
GMI1000	5,204	4,003 (77%)	90,016	11 (17.1)	456-470 (8.7-9.0%)
PSI07	4,968	4,057 (82%)	190,155	22 (35.7)	454-487 (9.5-9.8%)
IBSBF1503	4,897	3,809 (78%)	104,510	12 (20.4)	454-487 (9.3-10.0%)

697 [a]Measured by # of DNA barcodes mapped to a specific insertion site.

698 [b] Two pipelines (described in Methods) were used to predict gene essentiality based on the

699 distribution of transposon insertions in open reading frames.

Fig 1. Growth of *Ralstonia* strains in xylem sap is similar to growth in rich medium.

Ralstonia strain GMI1000, strain PSI07, or strain IBSBF1503 were grown in rich medium

(CPG), quarter-strength M63 minimal medium (MM) or xylem sap from Moneymaker tomato

plants. (**A**) Strains were inoculated at 10^5 CFU/ml into CPG, MM, or xylem sap. All cultures

were set up with 3 biological replicates (individual overnight cultures and individual batches of

xylem sap) and were performed in technical duplicate. Solid lines indicate time points sampled

from the same culture; the 12-24 hour timepoints were from an independent culture, but 0-8 and

24-48 hour timepoints were taken from the same culture. Error bars indicate standard deviation.

(**B**) The minimum doubling time between contiguous 4 hour measurements was calculated per

biological replicate. Lines indicate mean. Per strain, the doubling times in all conditions were

713   significantly different from each other (adjusted P<0.05 by ANOVA with Tukey's multiple

714   comparisons).

A. *R. pseudosolanacearum* GMI1000

B. *R. solanacearum* IBSBF1503

C. *R. syzygii* PSI07

D. Intersection analysis: Genes with co-fitness or condition-specific fitness impacts

Fig 2. Summary of RB-TnSeq data after growth of mutant libraries in xylem sap, minimal medium (MM), or rich medium (CPG). (A-C) Ranked Fit scores for all measured genes in CPG, MM, and xylem sap for (A) *R. pseudosolanacearum* GMI1000, and (B) *R. solanacearum* IBSBF1503, and (C) *R. syzygii* PSI07. The number of genes that influence fitness in each condition is indicated (average Fit > 1 or <-1; t-like test statistic >|2.5| all three RB-TnSeq trials). (D) Intersection analysis quantifying number of genes that influence fitness in a single condition or that have co-fitness in multiple conditions. Green bars indicate genes that influence fitness in sap and grey bars indicate genes that only influence fitness in culture media. The intersection visualization is modeled on UpSet Plots (66), an alternative to Venn Diagrams.

A. Fitness Factors in all strains

B. Inter-strain variation in fitness contributions of LPS genes

IBSBF1503

GMI1000

PSI07

LPS cluster 1   LPS cluster 2

C. *mdoGH* osmoregulated periplasmic glucan (OPG) genes are a GMI1000-specific xylem sap fitness factor

Fig 3. Envelope-associated genes contribute to fitness in xylem sap. Fit Scores of xylem sap

fitness factors in one or more *Ralstonia* strains. Graphs show fitness contributions of (A) Four

envelope-associated genes that are sap fitness factors in all three strains, (B) LPS genes that are

xylem sap fitness factors for at least one strain, and (C) periplasmic glucan biosynthesis genes

that are xylem sap fitness factors for GMI1000. Graphs show Fit scores of n=3 TnSeq trials and

bars show the mean. Full Fit scores and t-like statistics are shown in Table S2.

Fig 4. Amino acid auxotroph mutants have fitness defects in minimal medium (MM) and/or

xylem sap. Fit scores of amino acid biosynthesis mutants with reduced fitness in sap, MM, or

both. Gene name and locus tag for the GMI1000 gene are displayed. Full Fit scores and t-like

test statistics are shown in Table S2.

A. Cofactor biosynthesis genes universally required for growth in xylem sap

B. IBSBF1503 required two cobalamin (vitamin B12) gene clusters for growth in xylem sap

Fig 5. Mutants lacking cofactor biosynthesis genes have fitness defects in xylem sap. (A) A

thiamine and tetrahydrofolic acid gene were both xylem sap fitness factors in all three strains.

Graphs show Fit scores of n=3 TnSeq trials and bars show the mean. (B) *Ralstonia* IBSBF1503

required cobalamin biosynthesis for growth in xylem sap. The xylem sap fit scores of each gene

are written below the genes. "n.d." indicates fitness was not determined because mutants were

not present in the RB-TnSeq library. Full Fit scores and t-like test statistics are shown in Table

S2.

Fig 6. Mutants with gain-of-fitness in one-or-more strain and condition. (A) Regulatory mutants with known defects in extracellular polysaccharide (EPS) production relative to wild type: *phc* quorum sensing mutants, *vsrAD* mutants, and *rpoS* mutants. (B) Motility mutants lacking type IV pilus genes and the *flgN* flagellar chaperone genes had increased fitness in rich media in one or more strains. Graphs show Fit scores of n=3 TnSeq trials and bars show the mean. Full Fit scores and t-like test statistics are shown in Table S2.

754   **Supplementary Data**

756   **Fig S1. Venn diagram of putative essential genes in *Ralstonia* strains GMI1000,**

757   **IBSBF1503, and PSI07.** Orthologous predicted essential genes are represented proportionally in

758   Venn diagram intersections. Genes were included in the Venn diagram only if they were

759   predicted to be essential by both analysis pipelines used in this study (17, 18).

Fig S2. Growth of *Ralstonia* in rich medium, minimal medium, and xylem sap from

multiple susceptible tomato cultivars. (A-B) To determine the carrying capacity of sap and

culture media, *Ralstonia* strain GMI1000 was inoculated into rich CPG medium, quarter strength

M63 medium, or xylem sap from Moneymaker tomato at 10^4, 10^5, or 10^6 CFU/ml starting

density. (C-E). Wildtype *Ralstonia* strains GMI1000, PSI07, and IBSBF1503 were grown in sap

from Moneymaker and Bonny Best tomato. All strains were inoculated in biological triplicate

into three pools of xylem sap incubated at 28°C with shaking. Cell density was measured by

dilution plating.

Table S1: Gene essentiality predictions for GMI1000, PSI07, and IBSBF1503

Table S2: Genome-wide Fit scores for GMI1000, PSI07, and IBSBF1503

A. *R. pseudosolanacearum* GMI1000

ex vivo xylem sap Minimal medium (MM) Rich medium (CPG)

B. *R. solanacearum* IBSBF1503

ex vivo xylem sap Minimal Medium (MM) Rich Medium (CPG)

C. *R. syzygii* PSI07

ex vivo xylem sap Minimal medium (MM) Rich medium (CPG)

D. Intersection analysis: Genes with co-fitness or condition-specific fitness impacts

GMI1000 IBSBF1503 PSI07

A. Fitness Factors in all strains

B. Inter-strain variation in fitness contributions of LPS genes

C. *mdoGH* osmoregulated periplasmic glucan (OPG) genes are a GMI1000-specific xylem sap fitness factor

A. Cofactor biosynthesis genes universally required for growth in xylem sap

thiC
RSc0113

purU
RSc1873

CPG (Rich Medium)
Minimal Medium
Xylem sap

Fit Score

GMI1000 IBSBF1503 PSI07

B. IBSBF1503 required two cobalamin (vitamin B12) gene clusters for growth in xylem sap

hxoN	*cbiW*	*cobN*	*chlID*	*cbiE*	*cbiXC*	*cbiD*	*cbiL*	*cbiF*	*cbiG*	*cbiJH*	*cobW*	*btuR*	*cbiA*	*bluB*
-1.1	*n.d.*	*-1.1*	*-1.1*	*-1.1*	*-1.1*	*-0.9*	*-1.0*	*-1.3*	*-1.2*	*-1.1*	*-1.4*	*-0.9*	*-1.2*	*-0.1*

cbiP	*cobP*	*cbiB*	*cobD*	*btuF*	*cobC*	*cobS*	*cobT*
-1.1	*-1.0*	*-1.3*	*-0.9*	*0*	*n.d.*	*-1.2*	*n.d.*

1 kb

A. Regulatory Mutants

B. Motility Mutants