Fine-scale spatial patterns of wildlife disease are common and understudied

Gregory F Albery*[1], Amy R Sweeny[2], Daniel J Becker[3], Shweta Bansal[1]

1. Department of Biology, Georgetown University, Washington, DC, USA
2. Institute of Evolutionary Biology, University of Edinburgh, Edinburgh, Scotland
3. Department of Biology, Indiana University, Bloomington, Indiana, USA.

*Corresponding author: gfalbery@gmail.com

Abstract

1. All parasites are heterogeneous in space, yet little is known about the prevalence and scale of this spatial variation, particularly in wild animal systems. To address this question, we sought to identify and examine spatial dependence of wildlife disease across a wide range of systems.

2. Conducting a broad literature search, we collated 31 such datasets featuring 89 replicates and 71 unique host-parasite combinations, only 51% of which had previously been used to test spatial hypotheses. We analysed these datasets for spatial dependence within a standardised modelling framework using Bayesian linear models, and we then meta-analysed the results to identify generalised determinants of the scale and magnitude of spatial autocorrelation.

3. We detected spatial autocorrelation in 48/89 model replicates (54%) across 21/31 datasets (68%), spread across parasites of all groups. Even some very small study areas (under $0.01km^2$) exhibited substantial spatial variation.

4. Despite the common manifestation of spatial variation, our meta-analysis was unable to identify host-, parasite-, or sampling-level determinants of this heterogeneity across systems. Parasites of all transmission modes had easily detectable spatial patterns, implying that structured contact networks and susceptibility effects are potentially as important in spatially structuring disease as are environmental drivers of transmission efficiency.

5. Our findings demonstrate that fine-scale spatial patterns of infection manifest frequently and across a range of wild animal systems, and many studies are able to investigate them whether or not the original aim of the study was to examine spatially varying processes. Given the widespread nature of these findings, studies should more frequently record and analyse spatial data, facilitating development and testing of spatial hypotheses in disease ecology. Ultimately, this may pave the way for an *a priori* predictive framework for spatial variation in novel host-parasite systems.

Keywords: Wildlife disease; parasite transmission; spatial analysis; meta-analysis

Acknowledgements

We would like to extend our profound gratitude to the researchers who generously provided us with their data, and to all those who withstood our numerous emails regardless. GFA's postdoctoral fellowship is funded by NSF grant number 1414296.

Author contributions

GFA and SB conceived of the study, GFA and ARS collected the datasets, GFA conducted the INLA analyses, GFA and DJB conducted the meta-analysis, and GFA wrote the manuscript, with comments and edits offered by all authors.

Data Availability

All code and some datasets will be made available upon publication on Github, and archived on Zenodo.org.

Introduction

The maintenance and spread of parasites are inherently spatially structured (Cross, Lloyd-Smith, Johnson, & Getz, 2005; Kirby, Delmelle, & Eberth, 2017; Pullan, Sturrock, Soares Magalhaes, Clements, & Brooker, 2012), which holds important ramifications for epidemiological dynamics and disease control efforts (Becker et al., 2020; Cross et al., 2005; Plowright, Becker, McCallum, & Manlove, 2019). Spatial structure can arise through a wide variety of processes: for example, many parasites are transmitted from one host individual to another via direct contact, which requires a degree of spatiotemporal coincidence between

individuals (Manlove et al., 2018), so that infections are spatiotemporally staggered in waves

of transmission across the population. Other parasites transmit through persistent

environmental stages or arthropod vectors whose viability depends on spatially varying

abiotic conditions, creating spatial patterns of exposure and therefore of infection (Altizer et

al., 2006; Jamison, Tuttle, Jensen, Bierly, & Gonser, 2015; Patz, Graczyk, Geller, & Vittor,

2000). Finally, host immunity and susceptibility can be influenced by environmentally

varying factors like resource availability and climatic conditions, with knock-on impacts on

parasite burden and transmission (Becker et al., 2020, 2018). These diverse processes should

produce spatial patterns of infection across a wide range of wildlife systems, yet many

wildlife disease studies examine coarse spatial scales or assume that spatial patterns will be

negligible compared to other hypothesised drivers. As such, it is unclear how often infection

is spatially structured in these systems, at what range this variation can manifest, and how

host and parasite traits might alter its manifestation.

For logistical reasons, many studies of spatial drivers of infectious disease focus on discrete

between-population differences across large distances, often using a limited number of

discrete sampling locations rather than distributing their sampling locations continuously in

space (Plowright et al., 2019). Nevertheless, recent work suggests that spatial patterns of

infection may manifest at surprisingly fine spatial scales, within kilometres or even metres

(Abolins et al., 2018; Albery, Becker, Kenyon, Nussey, & Pemberton, 2019; Brooker et al.,

2006; Wood et al., 2007). This observation begs the question: what is the lower bound for the

range at which spatial effects can act? Identifying the range of spatial dependence (or

autocorrelation, meaning that data points that are closer together in space tend to be more

similar) is important for many reasons, including designing sampling regimes (Nusser, Clark,

Otis, & Huang, 2008; Plowright et al., 2019; Vidal-Martínez, Pech, Sures, Purucker, &

Poulin, 2010), building mechanistic models of parasite evolution over space (Best, Webb,

White, & Boots, 2011; Débarre, Hauert, & Doebeli, 2014), examining how disease risk

responds to anthropogenic activities like urbanisation (Saito & Sonoda, 2017), and directing

public health and conservation schemes (Brooker et al., 2006; Gilbertson et al., 2016).

Identifying the range of spatial dependence can also help examining how parasites spread

over landscapes and determining their transmission mechanisms (Reynolds, 1988). For

example, spatial dependence across large distances might suggest the influence of major

climatic correlates, while spatial dependence between nearby locations suggests a highly

localised infection process (Pullan et al., 2012). In human disease systems, such work has

shown that neighbouring districts of Thailand have more similar human malaria incidence,

suggesting local similarities in abiotic conditions or vector control programs that could limit

mosquito survival (Zhou et al., 2005). Similar analyses of wildlife disease could help pinpoint

transmission routes and guide disease control effort: for example, if researchers find that a

zoonotic disease has a long range of dependence in its wildlife reservoir, this could motivate

the use of widely placed sampling locations when trying to identify environmental drivers

(Becker, Crowley, Washburne, & Plowright, 2019; Plowright et al., 2019). Lastly, the scale

of spatial dependence has implications for more general theoretical understanding of

infectious disease dynamics. For example, links between biodiversity and disease dynamics

(e.g. "dilution effects") are dependent on the spatial scale of sampling (Cohen et al., 2016;

Rohr et al., 2020), and several rodent systems have identified contrasting spatial trends for

zoonotic diseases dependent on sampling scale (Luis, Kuenzi, & Mills, 2018; Morand et al.,

2019).

The strength and range of spatial dependence are also likely to depend on the traits of the

hosts and parasites involved. For example, parasites that persist for longer in the environment

are likely to experience stronger influences of environmental gradients than directly

transmitted counterparts (Satterfield, Altizer, Williams, & Hall, 2017). Similarly, highly

mobile species such as large carnivores or nomadic bats may more efficiently disseminate

parasites through the environment, reducing spatial autocorrelation (Gilbertson et al., 2016;

Peel et al., 2013). However, the relative contribution of host and parasite traits to shaping

spatial variation in infection remains unknown. The range of spatial dependence is most

commonly identified using spatial autocorrelation models (e.g. Albery et al., 2019; Becker,

Nachtmann, et al., 2019; Brooker et al., 2006; Gilbertson et al., 2016; Wood et al., 2007) or

analyses that quantify the spatial buffer regions in which environmental variables are best-

correlated with disease (e.g. Saito & Sonoda, 2017). Unfortunately, these approaches are

almost always reactive rather than proactive and they occur on a case-by-case basis rather

than being based on general rules or *a priori* understanding. To establish general factors

influencing the scale of spatial dependence in wildlife disease, a variety of host-parasite

systems must be analysed using comparable techniques and then synthesised. As well as

revealing fundamental drivers of spatial heterogeneity, identifying general rules in this way

could facilitate the development of predictive models for spatial structuring in host-parasite

systems with relatively poorly understood epidemiology (Gilbertson et al., 2016).

Researchers could then predict how within- and between-population processes will differ *a*

priori, before using empirical methods such as long-term studies at multiple scales (e.g. Luis

et al., 2018; Morand et al., 2019).

Prescriptive rules for examining geographic variation in wildlife disease are rare and hard to

generalise. For example, where studies seek to quantify the impact of environmental drivers

on parasitism, larger study extents may allow sampling the widest range of different

environmental factors and thus increasing spatial variation (Becker et al., 2020; Cohen et al.,

2016). Part of this methodological vacuum for generalisable rules emerges from the

analytical complexity of identifying them. A recent systematic review of ecoimmunology

studies uncovered a surprising lack of spatial methods, with most studies fitting discrete fixed

or random effects to control for spatial autocorrelation rather than directly examining

continuous patterns in space or using spatially explicit statistics (Becker et al., 2020).

Although the statistical competence of ecologists is high and increasing, particularly with

regards to areas like movement ecology and network analysis (Albery, Kirkpatrick, Firth, &

Bansal, 2021; Dougherty, Seidel, Carlson, Spiegel, & Getz, 2018; Jacoby & Freeman, 2016;

Webber & Vander Wal, 2019), no empirical framework exists for establishing the presence or

range of spatial variation in wildlife disease. Establishing general factors shaping spatial

variation across wildlife disease systems could lay the basis for this framework, improving

mechanistic understanding of parasite transmission, spatial sampling designs, and control

efforts.

Here, we conducted a synthesis of spatially distributed wildlife disease datasets across a wide

range of different host and parasite taxa, geographic contexts, and sampling regimes. We

analysed these datasets individually using a standardised modelling procedure, identifying

how generalised host-, parasite-, and sampling-level factors affect the prevalence and range

of spatial dependence. Specifically, we expected that studies would be most vulnerable to

strong spatial effects in larger study areas, with greater sampling efforts, and when parasites

exhibit indirect transmission mechanisms with extended environmental stages. We aimed to

provide important general estimates for the range of spatial autocorrelation from a wide range

of different host-parasite systems, laying the groundwork for *a priori* predictions about host-

parasite systems with unknown spatial properties.

Materials and methods

Data collection

To obtain a wide set of raw datasets we carried out a literature search, emailed authors to request data, and searched data repositories for publicly available datasets (Supplementary Figure 1). Our literature search used Web of Science to identify potential datasets published between 2009 and 28th August 2019, with the following terms: "(parasit* OR infect* OR disease) AND (wild OR natural) AND (mammal)". We restricted the search to mammals to increase the generalisability of our findings within this group of animals, and because of their importance for human and livestock health (Han, Kramer, & Drake, 2016).

We first screened a random subset of studies based on their abstracts, excluding studies of captive animals, review papers, and meta-analyses; publications without parasite data; studies without hosts (i.e., only sampling parasites in the environment; and studies of non-mammals. Because our downstream analyses relied upon a standard spatial modelling procedure, we also excluded studies with few samples (N<35), very low prevalence (<10%), or very high prevalence (>90%), owing to likely failure in model convergence.

If a study had openly available datasets we downloaded them, and for those that included binary infection data in map figures, we derived approximate spatial locations and associated infection status (i.e., "heads up digitisation", HUD). We also searched the Dryad data repository (https://datadryad.org) using the same search terms to find publicly available datasets.

For all other studies, we contacted corresponding authors using a standardised email template in September-December 2019 to request data. We classified the authors' responses into the following categories (Supplementary Figure 1): System not suitable: the system was poorly suited to our questions (e.g., migratory host population). No parasitology: the system did not include disease measures. No spatial data collected: no sources of spatial data (grid references, GPS locations) were collected and associated with individuals or samples. Privacy concerns: researchers were unable to share the data because they were collected on private land. Data not suitable: once data were inspected, the genre of spatial data was found to be unsuitable (e.g. too few spatial replicates), or it was deemed unlikely that models would run (e.g., points very unevenly distributed, sample sizes too low).

Some of the datasets contained multiple spatial sites that were each defined as a contiguous
population. Therefore, within the datasets, each replicate was defined as a unique host-
parasite-locality combination examining a contiguous population. We excluded replicates
with under 100 samples, to ensure convergence of our standardised spatial models (see
below).

Although we principally aimed to quantify fine-scale, within-population spatial effects, we
included several studies employing continuous or semi-continuous sampling at national and
county levels, to investigate whether the methods we used would operate well at these scales
and to establish an upper bound for sampling effects.

Statistical Analysis

Data standardisation

Data were manipulated and analysed using R version 3.6.3 (R Development Core Team,
2011). All code is available at github.com/gfalbery/libra. Our data cleaning procedure aimed
to minimise the probability of false positives and to restrict the data pool to a continuous
spatial distribution of samples. All spatial coordinates were converted to the scale of
kilometres or metres to allow comparison across systems. We removed spatial outliers and
parasite count outliers; if parasite counts were very overdispersed and/or highly zero-inflated
they were analysed as binomial (0/1) infection data rather than negative binomial. Categories
with low replication (generally <10 samples) were removed. We removed specific classes
that exhibited very low prevalence: e.g., adult Soay sheep and red deer had a very low
prevalence of *Nematodirus* sp., which is primarily a parasite of young ungulates (Hoberg,
Kocan, & Rickard, 2001); hence only lambs/calves were analysed. Individual identity was
fitted as a random effect if the dataset involved repeat measurements of the same individuals.
209

INLA Models

We based our analysis on a framework previously used in a study of spatial patterns of
disease in wild red deer (Albery et al., 2019). Integrated Nested Laplace Approximation
(INLA) models were fitted to each spatial dataset using the `inla` package. INLA is a
deterministic Bayesian algorithm that allows fitting of a Stochastic Partial Differentiation
Equation (SPDE) random effect to quantify and control for patterns of the response variable
in space. This relies on detection of spatial autocorrelation, where samples closer in space are

more similar than those further apart (Kirby et al., 2017; Tobler, 1970). The model estimates

how much variance is accounted for by autocorrelation, and models with and without the

SPDE effect can be compared to assess how it affects the fit of the model (Lindgren & Rue,

2015; Zuur, Ieno, & Saveliev, 2017). The model also provides a "range" parameter, which

estimates the distance at which samples are autocorrelated. We took this parameter to

represent a combination of sampling, transmission, and immune processes determining the

scale of spatial variation in the focal population.

We first fitted a "base" model with parasite burden (Gaussian or negative binomial) or

presence/absence (binary) as a response variable and with any fixed and random covariates.

To simplify our analyses, covariates usually included only temporal variables (month, year,

both as categorical variables), age category, and sex. We then fitted a model featuring an

SPDE random effect, with a penalised complexity prior (Fuglstad, Simpson, Lindgren, &

Rue, 2019). We compared the base model with the SPDE model, identifying whether the

latter had a lower Deviance Information Criterion (DIC), indicating improved model fit. We

took a change in DIC (ΔDIC) of 2 to distinguish between the two models and calculated the

DIC weight for the base and SPDE model, giving a proportion (0-1) that can be

conceptualised as "confidence that the spatial model was the best-fitting" (Wagenmakers &

Farrell, 2004). We also extracted the INLA range parameters. In total, we fitted INLA models

to 89 host-locale-parasite combinations, each of which comprised a different host-locale-

parasite combination, generated from 31 different study systems.

Meta-analysis of INLA models

To identify factors driving general trends of spatial variation, we conducted a meta-analysis

treating each unique parasite-system-site combination as a replicate, including parasite-, host-,

and sampling-level traits as fixed effects. We constructed hierarchical models using the

`metafor` package. Generally, meta-analyses typically focus on synthesizing effect sizes and

their variances across multiple systems (e.g. Sánchez *et al.* 2018). However, as generalised

spatial variation does not have a directional effect, we instead analysed measures of model fit,

predictive capacity, and the autocorrelation range, which is bounded at 0 and infinity. To give

a coarse measure of model predictive capacity that was easily standardised across all models,

we calculated the Spearman's Rank correlation between the observed and predicted values

for the model, using only the SPDE effect to predict (henceforth referred to as R). The

measures of model fit give an impression of the detectability and importance of spatial

patterns, while comparisons of the range estimate across systems will inform whether

different host and parasite traits cause spatial patterns to vary more sharply in space. We used

the *escalc* function to derive sampling variances for DIC weight and the INLA range (using

the point estimate and 95% confidence interval).

Our hierarchical models included each replicate nested within study as a random effect to

account for within- and between-study heterogeneity (Konstantopoulos, 2011). We also

included a random effect for host family, for which the covariance structure used the

phylogenetic correlation matrix (Nakagawa & Santos, 2012); we obtained our phylogeny

from the Open Tree of Life with the rotl and ape packages (Michonneau, Brown, & Winter,

2016; Paradis, Claude, & Strimmer, 2004). All models used the `rma.mv` function and

weighting by sampling variance. We first assessed heterogeneity in each of our response

variables by fitting a random-effects model (REM; intercept only) with restricted maximum

likelihood and then used Cochran's Q to test if such heterogeneity was greater than expected

by sampling error alone (Borenstein, Hedges, Higgins, & Rothstein, 2009).

We next used mixed-effects models (MEMs) to test how sampling-, host-, and parasite-level

factors affected our INLA model outputs. Sampling variables included: Number of samples;

Sampling area (total rectangular extent between the furthest points on the X- and Y-

coordinates, in km^2); Sampling method (3 levels: trapping, censusing, and

necropsy/convenience sampling); Spatial encoding method (4 levels: GPS; trapping grid;

locality; Easting/Northing); Spatial hypothesis testing (binary – i.e., did the study aim to

quantify spatial variation in some way?). We interpreted this latter variable as a combination

of study design and publication bias, where studies that are intended to pick up spatial

variation are both more likely to identify spatial patterns because of their sampling design,

and then more likely to be published if they do. Parasite traits included Transmission mode (4

levels: direct; faecal-oral; vector-borne; environmentally transmitted) and Taxon (8 levels:

arthropod, nematode, trematode, cestode, protozoan, bacterium, virus, other). Host traits

included: Home Range size (in km^2; log-transformed); Body Mass (in grams; log-

transformed); Host order (5 levels: Carnivora, Chiroptera, Ungulates, Glires, Proboscidea).

There was only one lagomorph, so rodents and lagomorphs were lumped together into the

"glires" clade. The same was true of odd-toed ungulates (Perissodactyla), so they were

lumped with Artiodactyla into an "ungulates" clade. For species for which a phenotypic

measure (e.g. body mass) was unavailable, we used the value for the closest relative for

which the data were available, according to a mammalian supertree (Fritz, Bininda-Emonds,
& Purvis, 2009).

To identify important drivers among these many correlated drivers, we conducted a model
addition process using maximum likelihood and Akaike Information Criterion corrected for
sample size (AICc) to determine model fit. Each of our meta-analytical explanatory variables
was added in turn, and the best-fitting variable (i.e., the one that most decreased AICc) was
kept for the following round. This process was repeated with the remaining variables, until no
variables improved model fit by more than 2 AICc. We report the final model, with the
minimal number of variables that improved model fit.

Spatiotemporal INLA models

Finally, we constructed spatiotemporal INLA models to assess the consistency of spatial
hotspots from year to year, and to investigate evidence of ephemeral waves of transmission
across the study systems. Of our 89 replicates, 44 replicates had more than one year of
sampling, with more than 100 spatial points per year, facilitating fitting spatiotemporal
models. For these replicates, we first reran the original models with the reduced dataset that
only included years with more than 100 replicates. We then fitted a spatiotemporal model
with a different spatially distributed effect (i.e. "spatial field") for each year, with no
autocorrelation between the fields. Improved model fit for this model would imply that the
spatial distribution of the parasite varied notably from year to year. Second, we fitted a similar
spatiotemporal model with an "exchangeable" autocorrelation specification between years.
This model format allows correlation between spatial fields, but without enforcing a time
sequence: that is, all fields were correlated by the same parameter ("Rho") regardless of how
far apart in time they were. The Rho parameter, which is bounded between -1 and 1, was then
interpreted to give an impression of the spatiotemporal consistency of the parasite distribution.
Parasites with high rho coefficients had very similar hotspots from year to year, while those
with low coefficients did not.

Results

Our literature review returned 3399 studies, and we screened a random selection of 1993
abstracts (over two weeks) to expedite data collection. 1151 of these were unsuitable because
they were in the wrong environment, host, or subject area, or had no data. This left 496

studies, for which we assessed data availability and emailed 432 authors to request data if

unavailable (Supplementary Figure 1).

Very few studies publicly archived continuous, within-population spatial data. Only 3/496

studies (0.6%) had such data ready to download, and 4 further studies had maps of samples

from which we could easily digitise sufficient data (Supplementary Figure 1). We also

already owned 3 datasets. When we emailed the corresponding authors of the remaining

studies, 92 responded, 22 of which (23.9%) indicated that they had not collected any within-

population spatial data as part of their study (Supplementary Figure 1). After navigating a

number of other obstacles to data sharing, followed by initial data triage, 26 authors kindly

offered to provide us with spatial data, resulting in 36 total viable datasets. Of these 36

datasets, 31 had at least one contiguous spatial population with >100 samples to which we

could apply INLA models.

Figure 1: The geographic and taxonomic distribution of the 31 datasets that we included in our final meta-analysis. Our data were evenly spread across the earth (Panel A), although with a notable cluster in Western Europe (see inset map in pink rectangle, Panel B). Sampling areas greater than 5000 km^2 are displayed as rectangles; smaller sample areas are represented by dots. Study system names correspond to the names in Supplementary Table 1. The datasets also included a wide range of different mammal orders and families (Panel C). The inset phylogeny represents order-level summaries for studies that were not carried out at the species level. Dots next to species' names in the phylogenies denote that multiple datasets included samples from that species. Different colours correspond to different taxonomic groups used for meta-analysis: ungulates, carnivores, glires, elephants, and carnivorous marsupials.

Most authors that responded (and had collected spatial data) were happy to share data with us, and the vast majority of studies for which we did not receive data were due to a lack of response or secondary response (Figure 1). 15 authors responded but declined to share data

due to privacy concerns, ongoing data usage, or authorship concerns. Comparing this to the

22 responders that had not collected spatial data implies that the main reason researchers do

not share spatial data is that they did not collect it; however, given that >300 researchers did

not respond (and they may not have been a random subset of the total), our ability to infer

this confidently is diminished. Notably, studies that investigated spatial variation tended to be

larger than those that did not (Supplementary Figure 2), implying that larger study areas

motivate researchers to more often consider spatial variation in their analyses.

We concluded data collection with 31 datasets, including 89 spatial replicates and 90 host

species (Figure 1). 67 replicates were species-level; the rest were conducted on selections of

species in the same order (e.g., rodent trapping, bat sampling, carnivore faecal sampling). The

datasets were distributed across five continents (Figure 1), and included 7 different

mammalian orders (Figure 1). The studies examined 41 different parasites, across a diverse

selection including viruses (N=6), bacteria (N=10), helminths (N=25), arthropods (N=14),

and one transmissible cancer (N=8). Infection measures included counts of parasites or

immune markers (N=30), binary assessment of infection status using observation or

seropositivity (N=52), and one study used parasite-associated mortality as a proxy (Myanmar

elephants, *Elephas maximus* (Lynsdale, Mumby, Hayward, Mar, & Lummaa, 2017)). Study

systems included, for example: rodent trapping studies examining flea burdens and their

associated parasites (e.g. rodents trapped in the Arizona hills (Kosoy et al., 2017) and

chipmunks in Yosemite National Park (Hammond et al., 2019)); long-term studies with

parasite data collected over the course of several decades (e.g. the Soay sheep of St Kilda

(Hayward et al., 2014), the Isle of Rum red deer (Albery et al., 2019), and the badgers of

Wytham Wood (Albery et al., 2020)); and studies examining seropositivity of mammals

across a geographic range to identify endemic areas (e.g. British otters infected with

Toxoplasma gondii (Smallbone et al., 2017)). See Supplementary Table 1 for a description of

each study system and the associated references and researchers that provided us with the

data. The area of the study systems varied widely, from 0.02 to 10^6 km^2 (Figure 2A).

Figure 2: The spatial autocorrelation term (SPDE) improved models across host-parasite systems and sampling regimes. The Y axis displays the degree of confidence that the spatial autocorrelation term improved model fit (Deviance Information Criterion weight), where models at the top of the figure fitted better than those at the bottom. The dashed line at DIC weight=0.5 denotes the point at which spatial and non-spatial models were equally supported. A: larger study areas more often revealed spatial patterns. B: most of our 31 study systems exhibited at least one spatially structured host-parasite combination. Study systems have been assigned arbitrary letters to anonymise them, and are arranged in order of increasing DIC weight. C: multiple mammalian host taxa exhibited spatial effects. D: multiple parasite taxa exhibited spatial effects. The points in panels C and D are sized according to the number of samples in the replicate. None of the terms displayed here had significant effects in our meta-analysis.

Our INLA models applied across datasets consistently revealed strong spatial patterns of

disease (Figure 2-3). The mean DIC change across all study systems was -14.5 (median -3.3),

and the spatial model fit better than the base model for 65/89 models (73%; DIC weight>0.5).

Using a conventional change of 2ΔDIC as a cutoff for improved model fit, 54% of models

across 21 study systems displayed detectable spatial patterns (Figure 2). Cochran's Q

revealed very low heterogeneity between systems in terms of their DIC weight ($Q(df=86) =$

46.89, P=0.9998), but extreme heterogeneity in terms of the range of autocorrelation

($Q(df=86) = 3823$, P<.0001).

Although half of the systems were spatially structured, our meta-analyses revealed that few

host-, parasite-, or sampling factors were predictive of spatial effects (see Supplementary

Table 2). The best-fitting model for DIC weight included only the study duration (years),

revealing that long-term studies were slightly more likely to uncover spatial effects

(ΔAIC=3.38; for all other variables ΔAIC<1.56). The INLA range parameter increased with

study area (ΔAIC=74.44) but was not affected by any other variables (ΔAIC<0.09). No

variation was accounted for by host or parasite taxon, host size, or host ranging behaviour.

Most notably, there was no significant variation in spatial range or DIC changes across

parasite transmission modes (Figure 3A-B).

Figure 3: Parasites of diverse transmission modes exhibit spatial autocorrelation effects. We
display A) spatial model DIC (deviance information criterion) weight, with points
representing the outcome of each replicate INLA (integrated nested laplace approximation)
model. Boxplots represent range, interquartile range, and median for parasites of each
transmission mode. The dashed line at DIC weight=0.5 denotes the point at which spatial and
non-spatial models were equally supported; points above the line display host-parasite
systems for which the spatial model was better supported than the non-spatial model. B)
INLA autocorrelation ranges; each line represents the autocorrelation decay of a different
replicate INLA model. The colours correspond to different transmission modes,
demonstrating substantial mixing of the range estimates for parasites of different transmission
modes. C) Temporal autocorrelation (Rho) component demonstrating inter-annual
correlations between spatial fields, for the subset of model replicates that had multiple
sampling years. Points represent a different replicate INLA model; black dots represent
means, and error bars represent standard errors. The dashed line at Rho=0 represents the point
of no correlation; points above the line had a positive correlation, while points below the line
had a negative correlation. D) Mosaic plot displaying the proportions of best-fitting models
according to DIC changes, across our spatiotemporal replicates.

Spatiotemporal models examining a subset of multi-year studies consistently improved model fit over static equivalents. The best-fitting model for many examined replicates was a spatiotemporal model, but the findings did not differ notably across transmission modes (Figure 3D). Rho (temporal autocorrelation of the spatial field) estimates for these models were moderate, and did not vary notably across transmission modes (Figure 3C). Most (36/44, 82%) had 95% credibility intervals that overlapped with zero, and 8 (18%) were significantly positive.

Discussion

We uncovered strong, pervasive spatial heterogeneity manifesting within an expansive diversity of mammal-parasite systems. Contrary to expectations, spatial heterogeneity was equally common and short-ranged for all transmission modes, despite our prediction that parasites with longer environmental stages would be more likely to exhibit spatial patterns. There are therefore three main takeaways from our findings: first, many study systems are spatially structured, likely by a combination of drivers, whether or not the study in question aims to quantify spatial variation or environmental drivers. Second, these drivers are relatively rarely investigated, but many systems currently have the spatial power and ability to investigate them if they wish, irrespective of the host-parasite system involved. Third, we were unable to develop a predictive framework for spatial dependence using the data available, but given more data across a wider range of host-parasite systems, such a framework may be possible to develop in the future. We therefore recommend that wild animal studies in disease ecology more regularly collect and share data on spatial behaviours and sampling locations where possible, regardless of host, parasite, or sampling regime.

Our methodology differed from that used in many other studies by investigating generalised spatial dependence rather than by quantifying specific environmental drivers that might drive this dependence. The only similar study that we know of (Gilbertson et al., 2016) used 48 parasite-locality replicates of cougar (*Puma concolor*) and bobcat (*Lynx rufus*) populations and found little evidence of spatial autocorrelation in parasite infection. In contrast to their approach, we used a wide set of different hosts, and our replicates all had between 100 and 10,000 samples (Supplementary Table 1), whereas only a few of their replicates had >100 samples, and none had >200 (Gilbertson et al., 2016). Additionally, they used Mantel tests,

which do not account for fixed covariates, while the INLA analyses we employed are more

suited to controlling for this variation. As such, we interpret our contrasting findings to

represent a difference in the power of our analyses, and the absence of large carnivores from

our dataset. Owing to its generality, similar methodology could be used in a range of

ecological contexts as a useful hypothesis-generating exercise: after uncovering strong spatial

structuring, researchers could follow up on this finding by investigating possible biotic or

abiotic drivers. We hope that more disease ecology studies in wild animals will make use of

similar methodology to bolster our understanding of disease dynamics in wild settings.

Surprisingly, neither larger study systems nor those that had previously been used to study

spatial hypotheses were more likely to exhibit detectable spatial patterns. Some very small

spatial replicates exhibited strong spatial effects, and the smallest area demonstrating a strong

spatial trend was 0.002km^2 (Figure 2). Similarly, some very large, well-sampled areas

showed no detectable spatial patterns: anti-*Toxoplasma gondii* antibodies in almost 200

Pennsylvania black bears (*Ursus americanus*) did not (Dubey et al., 2016), while prevalence

of *T. gondii* exhibited very strong spatial patterns in otters (*Lutra lutra*) across the United

Kingdom (Smallbone et al., 2017), and in house mice (*Mus musculus*) within the Senegalese

city of Dakar (Galal et al., 2019). However, larger study extents unsurprisingly exhibited

more long-range spatial autocorrelation effects. These areas inevitably contain within them a

multitude of smaller spatial effects and gradients, so that the findings of a specific study will

depend critically on the spatial sampling scale it employs (Cohen et al., 2016; Luis et al.,

2018; Morand et al., 2019; Pullan et al., 2012). Notably, the studies that did attempt to

quantify spatial variation tended to have substantially larger spatial extent than those that did

not (Supplementary Figure 2); this may represent a perception bias, where researchers

operating in larger study areas tend to anticipate spatial variation as being more important to

account for – or, *vice versa*, researchers asking spatial questions tend to sample across a

wider range to incorporate as much testable variation as possible (Becker, Nachtmann, et al.,

2019). The finding that larger study systems do not tend to more commonly exhibit

detectable spatial patterns in disease demonstrates that this perception bias is perhaps

unwarranted, and researchers at all scales should be able to incorporate spatial components

and hypotheses about infection processes.

Despite the ubiquity of spatial effects, we discovered a very low frequency of spatial data

collection and sharing: only 3 publicly available datasets included spatial data, and 22/92

responders said they had not collected any spatial data. The responses that we received
implied that alongside concerns about privacy and the understandable desire to control the
data associated with one's study system, the main reason for not sharing spatial data was that
the data were not collected in the first place. Location data may evade collection in some
contexts where GPS signals are hard to receive, precluding spatial data collection and
investigation of spatial questions. GPS instruments that function in remote environments can
be expensive, and for studies that do not explicitly aim to identify spatial patterns this may
seem an unnecessary expenditure. However, smartphones that can receive GPS data are now
widely available and can be used in all but the most remote locations. As many researchers
carry the means to collect spatial data in their pocket on a daily basis, it might take little
alteration to collection protocols to include location data in many cases. Future studies should
capitalise on the increasing availability of spatial telemetry and biologging technology, and
associated analytical capacity (Kays, Crofoot, Jetz, & Wikelski, 2015; Long, Nelson, Webb,
& Gee, 2014; Williams et al., 2020) to more frequently record, analyse, and share spatial data
in disease ecology (Albery et al., 2019; Kirby et al., 2017). This practice will facilitate easier
testing of the hypotheses that we outline above, as well as informing sampling regimes and
mechanistic models of disease dynamics, and allowing *a priori* prediction of host-parasite
systems' spatial properties.
505
Privacy is an issue of considerable ethical concern in epidemiology (Kirby et al., 2017), and
we contend that this concern may be contributing to a lack of open data sharing in wildlife
disease ecology. Sharing spatial data risks connecting individuals with their disease status,
which is particularly unwelcome in the case of stigmatised diseases such as HIV/AIDS;
indeed, although we did not examine human diseases, several of the researchers we contacted
opted not to share data because they were concerned that their results could be traced to
specific households or individuals. Researchers could overcome this issue by jittering points,
or by masking the actual GPS locations, replacing them with relative locations which are the
same distance away (Kirby et al., 2017). Unfortunately, the first option will reduce precision
and the latter may preclude investigation of specific geographic hypotheses or environmental
drivers, but this is a small price to pay in the cases where data are potentially sensitive.
517
518 We foresee a range of potential uses for curated datasets like ours. Although we quantified
519 some ecological and sampling-level drivers here, the dataset was still relatively small, and
520 subject to covarying factors: for example, most analyses of nematode infection were

conducted on even-toed ungulates, so that it was difficult to disentangle their implications for spatial variation. Future data collection and kind contributions from researchers may allow us to bolster this dataset to include a greater number of replicates, increasing the power and diversity of our analyses, bringing predictive models of spatial variation within our grasp. Further analysis on this dataset could investigate a number of general drivers such as population density or environmental heterogeneity, informing how they drive spatial patterns of infection within and across systems. Similar methodology could be applied to other animal groups such as birds and reptiles, whose nest and burrow locations offer ideal spatial context (e.g. Wood et al., 2007), or to marine mammals like dolphins that are regularly subject to behavioural censuses and disease surveillance (e.g. Leu, Sah, Krzyszczyk, Jacoby, & Mann, 2020). This approach could also be applied to intensively and widely spatially distributed sampling locations, either for smaller animals such as insects (Wallace et al., 2021) or for immobile organisms like plants (Halliday et al., 2020). Finally, immunity is often quantified alongside parasite burden and prevalence, and it would be interesting to see whether spatial variation in immunity manifests on the same scale, and whether it predicts disease risk (Becker et al., 2020). Given these diverse and widespread opportunities, popularising the breadth and frequency of open spatial data sharing is likely to open the door to a wide range of interesting studies and, ultimately, to the development of predictive *a priori* frameworks for spatial processes in disease ecology.

References

Abolins, S., Lazarou, L., Weldon, L., Hughes, L., King, E. C., Drescher, P., … Viney, M. (2018). The ecology of immune state in a wild mammal, Mus musculus domesticus. *PLoS Biology*, *16*(4), e2003538. doi: 10.1371/journal.pbio.2003538

Albery, G. F., Becker, D. J., Kenyon, F., Nussey, D. H., & Pemberton, J. M. (2019). The fine-scale landscape of immunity and parasitism in a wild ungulate population. *Integrative and Comparative Biology*, *icz016*(5), 1–11. doi: 10.1093/icb/icz016

Albery, G. F., Kirkpatrick, L., Firth, J. A., & Bansal, S. (2021). Unifying spatial and social network analysis in disease ecology. *Journal of Animal Ecology*, *90*, 1–17. doi: 10.1111/1365-2656.13356

Albery, G. F., Newman, C., Ross, J. B. G. B., Macdonald, D. W., Bansal, S., Buesching, C. D., … Buesching, C. D. (2020). Negative density-dependent parasitism in a group-living carnivore. *Proceedings of the Royal Society B: Biological Sciences*, *287*(1941), 20202655. doi:

553   10.1101/2020.06.15.153726

554 Altizer, S., Dobson, A., Hosseini, P., Hudson, P., Pascual, M., & Rohani, P. (2006). Seasonality and
555   the dynamics of infectious diseases. *Ecology Letters*, *9*(4), 467–484. doi: 10.1111/j.1461-
556   0248.2005.00879.x

557 Becker, D. J., Albery, G. F., Kessler, M. K., Lunn, T. J., Falvo, C. A., Czirják, G. Á., … Plowright, R.
558   K. (2020). Macroimmunology: the drivers and consequences of spatial patterns in wildlife
559   immune defense. *Journal of Animal Ecology*, *89*(4), 972–995. doi: 10.1111/1365-2656.13166

560 Becker, D. J., Crowley, D. E., Washburne, A. D., & Plowright, R. K. (2019). Temporal and spatial
561   limitations in global surveillance for bat filoviruses and henipaviruses. *Biology Letters*, *15*,
562   20190423.

563 Becker, D. J., Czirják, G. Á., Volokhov, D. V., Bentz, A. B., Carrera, J. E., Camus, M. S., …
564   Streicker, D. G. (2018). Livestock abundance predicts vampire bat demography, immune
565   profiles, and bacterial infection risk. *Philosophical Transactions of the Royal Society B*, *373*,
566   20170089. doi: 10.1098/rstb.2017.0089

567 Becker, D. J., Nachtmann, C., Argibay, H. D., Botto, G., Escalera-Zamudio, M., Carrera, J. E., …
568   Streicker, D. G. (2019). Leukocyte Profiles Reflect Geographic Range Limits in a Widespread
569   Neotropical Bat. *Integrative and Comparative Biology*, *icz007*. doi: 10.1093/icb/icz007

570 Best, A., Webb, S., White, A., & Boots, M. (2011). Host resistance and coevolution in spatially
571   structured populations. *Proceedings of the Royal Society B: Biological Sciences*, *278*(1715),
572   2216–2222. doi: 10.1098/rspb.2010.1978

573 Borenstein, M., Hedges, L. V., Higgins, J. P. T., & Rothstein, H. (2009). *Introduction to meta-*
574   *analysis*. Retrieved from https://www.wiley.com/en-us/Introduction+to+Meta+Analysis-p-
575   9780470057247

576 Brooker, S., Alexander, N., Geiger, S., Moyeed, R. A., Stander, J., Fleming, F., … Bethony, J. (2006).
577   Contrasting patterns in the small-scale heterogeneity of human helminth infections in urban and
578   rural environments in Brazil. *International Journal for Parasitology*, *36*(10–11), 1143–1151.
579   doi: 10.1016/j.ijpara.2006.05.009

580 Cohen, J. M., Civitello, D. J., Brace, A. J., Feichtinger, E. M., Ortega, C. N., Richardson, J. C., …
581   Rohr, J. R. (2016). *Spatial scale modulates the strength of ecological processes driving disease*
582   *distributions*. 3359–3364. doi: 10.1073/pnas.1521657113

583 Cross, P. C., Lloyd-Smith, J. O., Johnson, P. L. F., & Getz, W. M. (2005). Duelling timescales of host
584   movement and disease recovery determine invasion of disease in structured populations.

Ecology Letters, *8*(6), 587–595. doi: 10.1111/j.1461-0248.2005.00760.x

Débarre, F., Hauert, C., & Doebeli, M. (2014). Social evolution in structured populations. *Nature*
Communications, *5*, 1–7. doi: 10.1038/ncomms4409

Dougherty, E. R., Seidel, D. P., Carlson, C. J., Spiegel, O., & Getz, W. M. (2018). Going through the
motions : incorporating movement analyses into disease research. *Ecology Letters*, *21*(4), 588–
604. doi: 10.1111/ele.12917

Dubey, J. P., Brown, J., Ternent, M., Verma, S. K., Hill, D. E., Cerqueira-Cézar, C. K., …
Humphreys, J. G. (2016). Seroepidemiologic study on the prevalence of Toxoplasma gondii and
Trichinella spp. infections in black bears (Ursus americanus) in Pennsylvania, USA. *Veterinary*
Parasitology, *229*, 76–80. doi: 10.1016/j.vetpar.2016.09.013

Fritz, S. A., Bininda-Emonds, O. R. P., & Purvis, A. (2009). Geographical variation in predictors of
mammalian extinction risk: big is bad, but only in the tropics. *Ecology Letters*, *12*(6), 538–549.
doi: 10.1111/j.1461-0248.2009.01307.x

Fuglstad, G. A., Simpson, D., Lindgren, F., & Rue, H. (2019). Constructing Priors that Penalize the
Complexity of Gaussian Random Fields. *Journal of the American Statistical Association*,
114(525), 445–452. doi: 10.1080/01621459.2017.1415907

Galal, L., Schares, G., Stragier, C., Vignoles, P., Brouat, C., Cuny, T., … Mercier, A. (2019).
Diversity of Toxoplasma gondii strains shaped by commensal communities of small mammals.
International Journal for Parasitology, *49*(3–4), 267–275. doi: 10.1016/j.ijpara.2018.11.004

Gilbertson, M. L. J., Carver, S., Vandewoude, S., Crooks, K. R., Lappin, M. R., & Craft, M. E.
(2016). Is parasite exposure spatially autocorrelated? Patterns of parasites in puma (Puma
concolor) and bobcat (Lynx rufus). *Ecosphere*, *7*(11), 1–12. doi: 10.1002/ecs2.1558

Halliday, F. W., Penczykowski, R. M., Barrès, B., Eck, J. L., Numminen, E., & Laine, A. L. (2020).
Facilitative priority effects drive parasite assembly under coinfection. *Nature Ecology and*
Evolution, *4*(11), 1510–1521. doi: 10.1038/s41559-020-01289-9

Hammond, T. T., Hendrickson, C. I., Maxwell, T. L., Petrosky, A. L., Palme, R., Pigage, J. C., &
Pigage, H. K. (2019). Host biology and environmental variables differentially predict flea
abundances for two rodent hosts in a plague-relevant system. *International Journal for*
Parasitology: Parasites and Wildlife, *9*(April), 174–183. doi: 10.1016/j.ijppaw.2019.04.011

Han, B. A., Kramer, A. M., & Drake, J. M. (2016). Global Patterns of Zoonotic Disease in Mammals.
Trends in Parasitology, *32*(7), 565–577. doi: 10.1016/j.pt.2016.04.007

Hayward, A. D., Garnier, R., Watt, K. a, Pilkington, J. G., Grenfell, B. T., Matthews, J. B., …

Graham, A. L. (2014). Heritable, heterogeneous, and costly resistance of sheep against
618   nematodes and potential feedbacks to epidemiological dynamics. *The American Naturalist*, *184*
619   *Suppl*, S58-76. doi: 10.1086/676929

620   Hoberg, E. P., Kocan, A. A., & Rickard, L. G. (2001). Gastrointestinal strongyles in wild ruminants.
621   *Diseases of Wild Mammals*, 193–227. doi: 10.1002/9780470377000.ch8

622   Jacoby, D. M. P., & Freeman, R. (2016). Emerging Network-Based Tools in Movement Ecology.
623   *Trends in Ecology and Evolution*, *31*(4), 301–314. doi: 10.1016/j.tree.2016.01.011

624   Jamison, A., Tuttle, E., Jensen, R., Bierly, G., & Gonser, R. (2015). Spatial ecology, landscapes, and
625   the geography of vector-borne disease: A multi-disciplinary review. *Applied Geography*, *63*,
626   418–426. doi: 10.1016/j.apgeog.2015.08.001

627   Kays, R., Crofoot, M. C., Jetz, W., & Wikelski, M. (2015). Terrestrial animal tracking as an eye on
628   life and planet. *Science*, *348*(6240), aaa2478. doi: 10.1126/science.aaa2478

629   Kirby, R. S., Delmelle, E., & Eberth, J. M. (2017). Advances in spatial epidemiology and geographic
630   information systems. *Annals of Epidemiology*, *27*(1), 1–9. doi:
631   10.1016/j.annepidem.2016.12.001

632   Konstantopoulos, S. (2011). Fixed effects and variance components estimation in three-level meta-
633   analysis. *Research Synthesis Methods*, *2*(1), 61–76. doi: 10.1002/jrsm.35

634   Kosoy, M., Reynolds, P., Bai, Y., Sheff, K., Enscore, R. E., Montenieri, J., … Gage, K. (2017).
635   Small-Scale Die-Offs in Woodrats Support Long-Term Maintenance of Plague in the U.S.
636   Southwest. *Vector-Borne and Zoonotic Diseases*, *17*(9), 635–644. doi: 10.1089/vbz.2017.2142

637   Leu, S. T., Sah, P., Krzyszczyk, E., Jacoby, A., & Mann, J. (2020). Sex, synchrony, and skin contact:
638   integrating multiple behaviors to assess parasite transmission risk. *Behavioral Ecology*, 1–10.
639   doi: 10.1093/beheco/araa002

640   Lindgren, F., & Rue, H. (2015). Bayesian Spatial Modelling with R-INLA. *Journal of Statistical*
641   *Software*, *63*(19), 1–25. doi: 10.18637/jss.v063.i19

642   Long, J. A., Nelson, T. A., Webb, S. L., & Gee, K. L. (2014). A critical examination of indices of
643   dynamic interaction for wildlife telemetry studies. *Journal of Animal Ecology*, *83*(5), 1216–
644   1233. doi: 10.1111/1365-2656.12198

645   Luis, A. D., Kuenzi, A. J., & Mills, J. N. (2018). Species diversity concurrently dilutes and amplifies
646   transmission in a zoonotic host–parasite system through competing mechanisms. *Proceedings of*
647   *the National Academy of Sciences*, *115*(31), 7979–7984. doi: 10.1073/pnas.1807106115

Lynsdale, C. L., Mumby, H. S., Hayward, A. D., Mar, K. U., & Lummaa, V. (2017). Parasite-

associated mortality in a long-lived mammal: Variation with host age, sex, and reproduction.

Ecology and Evolution, *7*(24), 10904–10915. doi: 10.1002/ece3.3559

Manlove, K., Aiello, C., Sah, P., Cummins, B., Hudson, P. J., & Cross, P. C. (2018). The ecology of

movement and behaviour: a saturated tripartite network for describing animal contacts.

Proceedings. Biological Sciences, *285*(1887), 20180670. doi: 10.1098/rspb.2018.0670

Michonneau, F., Brown, J. W., & Winter, D. J. (2016). rotl: an R package to interact with the Open

Tree of Life data. *Methods in Ecology and Evolution*, *7*(12), 1476–1481. doi: 10.1111/2041-

210X.12593

Morand, S., Blasdell, K., Bordes, F., Buchy, P., Carcy, B., Chaisiri, K., … Tran, A. (2019). Changing

landscapes of Southeast Asia and rodent-borne diseases: decreased diversity but increased

transmission risks. *Ecological Applications*, *29*(4), 1–15. doi: 10.1002/eap.1886

Nakagawa, S., & Santos, E. S. A. (2012). Methodological issues and advances in biological meta-

analysis. *Evolutionary Ecology*, *26*(5), 1253–1274. doi: 10.1007/s10682-012-9555-5

Nusser, S. M., Clark, W. R., Otis, D. L., & Huang, L. (2008). Sampling Considerations for Disease

Surveillance in Wildlife Populations. *Journal of Wildlife Management*, *72*(1), 52–60. doi:

10.2193/2007-317

Paradis, E., Claude, J., & Strimmer, K. (2004). APE: Analyses of phylogenetics and evolution in R

language. *Bioinformatics*, *20*(2), 289–290. doi: 10.1093/bioinformatics/btg412

Patz, J. A., Graczyk, T. K., Geller, N., & Vittor, A. Y. (2000). Effects of environmental change on

emerging parasitic diseases. *International Journal for Parasitology*, *30*(12–13), 1395–1405. doi:

10.1016/S0020-7519(00)00141-7

Peel, A. J., Sargan, D. R., Baker, K. S., Hayman, D. T. S., Barr, J. A., Crameri, G., … Cunningham,

A. A. (2013). Continent-wide panmixia of an African fruit bat facilitates transmission of

potentially zoonotic viruses. *Nature Communications*, *4*. doi: 10.1038/ncomms3770

Plowright, R. K., Becker, D. J., McCallum, H., & Manlove, K. R. (2019). Sampling to elucidate the

dynamics of infections in reservoir hosts. *Philosophical Transactions of the Royal Society B:*

Biological Sciences, *374*(1782). doi: 10.1098/rstb.2018.0336

Pullan, R. L., Sturrock, H. J. W., Soares Magalhaes, R. J., Clements, A. C. A., & Brooker, S. J.

(2012). Spatial parasite ecology and epidemiology: a review of methods and applications.

Parasitology, *139*(14), 1870–1887. doi: 10.1017/S0031182012000698

R Development Core Team, R. (2011). R: A Language and Environment for Statistical Computing (R.

D. C. Team, Ed.). *R Foundation for Statistical Computing*, p. 409. doi: 10.1007/978-3-540-

74686-7

Reynolds, K. M. (1988). Analysis of Epidemics Using Spatio-Temporal Autocorrelation.
Phytopathology, *78*(2), 240. doi: 10.1094/phyto-78-240

Rohr, J. R., Civitello, D. J., Halliday, F. W., Hudson, P. J., Lafferty, K. D., Wood, C. L., & Mordecai,
E. A. (2020). Towards common ground in the biodiversity– disease debate. *Nature Ecology &*
Evolution, *4*(1), 24–33. doi: 10.1038/s41559-019-1060-6

Saito, M. U., & Sonoda, Y. (2017). Symptomatic Raccoon Dogs and Sarcoptic Mange Along an
Urban Gradient. *EcoHealth*, *14*(2), 318–328. doi: 10.1007/s10393-017-1233-1

Sánchez, C. A., Becker, D. J., Teitelbaum, C. S., Barriga, P., Brown, L. M., Majewska, A. A., …
Altizer, S. (2018). On the relationship between body condition and parasite infection in wildlife:
a review and meta-analysis. *Ecology Letters*, *21*(12), 1869–1884. doi: 10.1111/ELE.13160

Satterfield, D. A., Altizer, S., Williams, M. K., & Hall, R. J. (2017). Environmental persistence
influences infection dynamics for a butterfly parasite. *PLoS ONE*, *12*(1), 1–16. doi:
10.1371/journal.pone.0169982

Senior, A. M., Grueber, C. E., Kamiya, T., Lagisz, M., O'Dwyer, K., Santos, E. S. A., & Nakagawa,
S. (2016). Heterogeneity in ecological and evolutionary meta-analyses: Its magnitude and
implications. *Ecology*, *97*(12), 3293–3299. doi: 10.1002/ecy.1591

Smallbone, W. A., Chadwick, E. A., Francis, J., Guy, E., Perkins, S. E., Sherrard-Smith, E., & Cable,
J. (2017). East-West Divide: Temperature and land cover drive spatial variation of Toxoplasma
gondii infection in Eurasian otters (Lutra lutra) from England and Wales. *Parasitology*, *144*(11),
1433–1440. doi: 10.1017/S0031182017000865

Tobler, W. R. (1970). A Computer Movie Simulating Urban Growth in the Detroit Region. *Economic*
Geography, *46*, 234. doi: 10.2307/143141

Vidal-Martínez, V. M., Pech, D., Sures, B., Purucker, S. T., & Poulin, R. (2010). Can parasites really
reveal environmental impact? *Trends in Parasitology*, *26*(1), 44–51. doi:
10.1016/j.pt.2009.11.001

Wagenmakers, E. J., & Farrell, S. (2004). AIC model selection using Akaike weights. *Psychonomic*
Bulletin and Review, *11*(1), 192–196. doi: 10.3758/BF03206482

Wallace, M. A., Coffman, K. A., Gilbert, C., Ravindran, S., Albery, G. F., Abbott, J., … Obbard, D. J.
(2021). The discovery, distribution and diversity of DNA viruses associated with *Drosophila*
melanogaster in Europe. *Virus Evolution*. doi: 10.1093/ve/veab031

Webber, Q. M. R., & Vander Wal, E. (2019). Trends and perspectives on the use of animal social

network analysis in behavioural ecology: a bibliometric approach. *Animal Behaviour*, *149*, 77–

87. doi: 10.1016/j.anbehav.2019.01.010

Williams, H. J., Taylor, L. A., Benhamou, S., Bijleveld, A. I., Clay, T. A., de Grissac, S., … Börger,

L. (2020). Optimizing the use of biologgers for movement ecology research. *Journal of Animal

Ecology*, *89*(1), 186–206. doi: 10.1111/1365-2656.13094

Wood, M. J., Cosgrove, C. L., Wilkin, T. A., Knowles, S. C. L., Day, K. P., & Sheldon, B. C. (2007).

Within-population variation in prevalence and lineage distribution of avian malaria in blue tits,

<i>Cyanistes caeruleus</ i>. *Molecular Ecology*, *16*(15), 3263–3273. doi: 10.1111/j.1365-

294X.2007.03362.x

Zhou, G., Sirichaisinthop, J., Sattabongkot, J., Jones, J., Bjørnstad, O. N., Yan, G., & Cui, L. (2005).

Spatio-temporal distribution of Plasmodium falciparum and p. Vivax malaria in Thailand. *The

American Journal of Tropical Medicine and Hygiene*, *72*(3), 256–262. Retrieved from

http://www.ncbi.nlm.nih.gov/pubmed/15772317

Zuur, A. F., Ieno, E. N., & Saveliev, A. A. (2017). *Beginner's guide to spatial, temporal, and spatial-

temporal ecological data analysis with R-INLA*. Retrieved from

https://searchworks.stanford.edu/view/12089113

Supplement to Albery et al.: *Fine-scale spatial patterns of wildlife disease are common and*

understudied.

Study system	Country	Location	Species	Host order	N	Parasites	Samples	Sampling method	Tested spatial?	Contributors	Example Reference
AlpineIbex	Italy	Gran Paradiso National Park	*Capra ibex*	Ungulates	4	Helminths; *Nematodirus*; *Marshallagia*; Coccidia	145	Census	N	Alice Brambilla	(Brambilla *et al.* 2015)
BerlinFoxes	Germany	Berlin	*Vulpes vulpes*	Carnivora	1	Canine distemper virus	762	Opportunistic	Y	Downloaded	(Gras *et al.* 2018)
BrazilianBats	Brazil		5+	Chiroptera	1	*Bartonella*	162	Noninvasive	Y	Keyla deSousa; Marcos Andre	Unpublished
BritishOtters	UK		*Lutra lutra*	Carnivora	1	*Toxoplasma gondii*	583	Opportunistic	Y	Derived	(Smallbone *et al.* 2017)
ConnecticutMice	USA	Connecticut	*Peromyscus leucopus*	Glires	1	Ticks	105	Trapping	N	Danielle Tufts; Maria Diuk-Wasser	(Tufts & Diuk-Wasser 2018)
ChileTriatomines	Chile		5+	Glires	4	Trypanosomes	230	Trapping	Y	Antonella Bacigalupo	(Ihle-Soto *et al.* 2019)
DakarRodents	Senegal	Dakar	*Mus musculus*	Glires	1	*Toxoplasma gondii*	424	Trapping	N	Lokman Galal	(Galal *et al.* 2019)
ElDoRodents	USA	El Dorado	5+	Glires	2	Fleas; *Bartonella*	1612	Trapping	Y	Michael Kosoy	(Kosoy *et al.* 2017)
FalkirkMice	Scotland	Callendar Park	*Apodemus sylvaticus*	Glires	8	*Heligmosomoides polygyrus*; *Eimeria*; *Capillaria*; Ticks; Mites; Fleas	596	Trapping	N	Amy Sweeny; Amy Pedersen	(Sweeny *et al.* 2019)
FoleyRodents	United States	California	5+	Glires	3	*Anaplasma*; *Borrelia*; Ectoparasites	948	Trapping	N	Janet Foley	(Foley *et al.* 2016)
GermanRaccoons	Germany		*Procyon lotor*	Carnivora	1	*Toxoplasma gondii*	458	Opportunistic	Y	Mike Heddergott	(Heddergott *et al.* 2017)
GreekHares	Greece		*Lepus europaeus*	Glires	1	European brown hare syndrome virus	209	Hunted	Y	Derived	(Sokos *et al.* 2018)

GrenadaMongooses	Grenada		*Herpestes javanicus*	Carnivora	2	Rabies; *Salmonella*	157	Trapping	Y	Ulrike Ziegler; Bruno Chomel; David Jaffe	(Jaffe *et al.* 2018)
HendyRodents	USA	California	*Neotoma fuscipes; Peromyscus maniculatus; Tamias ochrogenys*	Glires	3	Ticks; Aphag; Borrelia	451	Trapping	N	Janet Foley	Unpublished
HighlandDeer	Scotland	Highlands	*Cervus elaphus*	Ungulates	1	*Fasciola hepatica*	103	Hunted	Y	Andrew French	(French *et al.* 2019)
KielderVoles	England	Kielder Woods, Liverpool	*Microtus agrestis*	Glires	12	Cowpoxvirus; *Anaplasma*; *Babesia*	3020	Trapping	N	Mike Begon; Sandra Telfer	(Davis *et al.* 2015)
MultimammateMice	Tanzania	Morogoro	*Mastomys natalensis*	Glires	1	MORV	5547	Trapping	Y	Lucinda Kirkpatrick	(Berkvens *et al.* 2019)
MyanmarElephants	Myanmar		*Elephas maximus*	Proboscidea	1	Parasite-associated mortality	1626	Opportunistic	N	Carly Lynsdale; Virpi Lummaa	(Lynsdale *et al.* 2017)
PantanalMammals	Brazil	Pantanal Wetlands	*Cerdocyon thous*	Carnivora	1	*Hepatozoon*	115	Trapping	N	Marcos Andre; Keyla de Sousa	(de Sousa *et al.* 2017)
PennsylvaniaBears	USA	Pennsylvania	*Ursus americanus*	Carnivora	1	*Toxoplasma gondii*	173	Hunted	N	Jitender Dubey; Justin Brown	(Dubey *et al.* 2016)
RomanianFoxes	Romania		*Vulpes vulpes*	Carnivora	1	Fleas	268	Opportunistic	Y	Janet Foley; Andrei Mihalca	(Foley *et al.* 2017)
RumDeer	Scotland	Isle of Rum	*Cervus elaphus*	Ungulates	8	Strongyles; *Nematodirus*; *Capillaria*; Coccidia; *Moniezia*; *Fasciola hepatica*; *Dictyocaulus*; *Elaphostrongylus cervi*	2068	Census	N	Greg Albery; Josephine Pemberton; Daniel Nussey	(Albery *et al.* 2019)
SoaySheep	Scotland	Isle of St. Kilda	*Ovis aries*	Ungulates	4	Strongyles; Coccidia; *Nematodirus*; Keds	7197	Census	N	Josephine Pemberton; Daniel Nussey	(Hayward *et al.* 2014)
SpanishIbex	Spain	Sierra Nevada	*Capra ibex*	Ungulates	1	Mange	746	Hunted	Y	Joao Carvalho; José Granados	(Carvalho *et al.* 2015)
StrasbourgPrimate	France	Strasbourg Primatology Centre	5+	Carnivora	1	Taeniids	103	Noninvasive	N	Valentin Greigert	(Greigert *et al.* 2019)

SwedenVoles	Sweden	Revinge	*Myodes glareolus*	Glires	1	*Borrelia*	1999	Trapping	N	Lars Raberg	(Råberg *et al.* 2017)
TasmanianDevils	Tasmania		*Sarcophilus harrisii*	Dasyuro-morphia	7	Devil facial tumour disease	822	Trapping	Y	Downloaded	(Lazenby *et al.* 2018)
TurnerUngulates	Namibia	Etosha National Park	*Loxodonta africana; Oryx gazella; Connochaetes taurinus; Antidorcas marsupialis; Equus quagga*	Proboscidea; Ungulates; Artiodactyla; Proboscidea	10	Strongyles; *Eimeria melis*; *Strongyloides* sp.	1132	Census	Y	Wendy Turner; Wayne Getz	(Turner *et al.* 2014)
WoodchesterBadgers	England	Woodchester Park, Gloucestershire	*Meles meles*	Carnivora	1	Bovine Tuberculosis	3319	Trapping	N	Matthew Silk; David Hodgson; Dez Delahay	(Rozins *et al.* 2018)
WythamBadgers	England	Wytham Wood	*Meles meles*	Carnivora	4	Fleas; Lice; Ticks; *Eimeria melis*	7220	Trapping	Y	Chris Newman; Christina Buesching; David MacDonald	(Albery *et al.* 2020)
YosemiteChipmunks	USA	Yosemite National Park, California	*Tamias speciosus*	Glires	1	Fleas	1126	Trapping	Y	Tali Hammond	(Hammond *et al.* 2019)

736   Supplementary Table 1: Summary table depicting the study systems used in the meta-
737 analysis, including names, locations, host species, and sampling traits. The column "Tested
738 spatial" denotes whether or not one of the study's stated aims was to investigate spatial or
739 geographic variation, e.g. in environmental drivers. The column "N" refers to the number of
740 spatial replicates and INLA models associated with the study system. The "contributors"
741   column refers to the individuals who generously provided data.

745   Supplementary Figure 1: The data collection, cleaning, and analysis pipeline.

748   Supplementary Figure 2: Datasets that were previously used to test spatial hypothesis were

749   larger. The y axis is in log-km^2. The boxplots represent the range, the interquartile range, and

750   the median.

	Round	Variable	Delta	Kept
DIC Weight	1	hOrder	7.774202	0
		Mass	1.239772	0
		HR	1.430737	0
		ParasiteTaxon	5.63204	0
		TransmissionMode	2.617334	0
		Coord_System	3.75258	0
		SpatialTested	1.712346	0
		Family	1.075977	0
		Area	-1.35223	0
		UniqueSites	-3.18133	0
		Years	-3.38106	1
	2	hOrder	8.646439	0
		Mass	1.70461	0
		HR	1.821279	0
		ParasiteTaxon	7.057377	0
		TransmissionMode	4.62246	0
		Coord_System	6.067934	0
		SpatialTested	1.598549	0
		Family	1.216243	0
		Area	-1.56394	0
		UniqueSites	0.554708	0
Range	1	hOrder	-1.38091	0
		Mass	1.121613	0
		HR	-4.46747	0
		ParasiteTaxon	3.735021	0
		TransmissionMode1	-0.66759	0
		Family	-2.07499	0
		Area	-74.4449	1
		UniqueSites	1.539401	0
		Years	1.959999	0

2	hOrder	4.89838	0
	Mass	1.823107	0
	HR	0.039442	0
	ParasiteTaxon	2.174064	0
	TransmissionMode	0.65543	0
	Family	-0.09311	0
	UniqueSites	1.694263	0
	Years	0.85906	0

Supplementary Table 2: Akaike Information Criterion (AIC) changes associated with variable addition to meta-analysis models. Variables with asterisks were kept to the following round of model addition.

Supplementary Table 1 References

Albery, G.F., Becker, D.J., Kenyon, F., Nussey, D.H. & Pemberton, J.M. (2019). The fine-scale landscape of immunity and parasitism in a wild ungulate population. *Integr. Comp. Biol.*, icz016, 1–11.

Albery, G.F., Newman, C., Ross, J.B.G.B., Macdonald, D.W., Bansal, S., Buesching, C.D., *et al.* (2020). Negative density-dependent parasitism in a group-living carnivore. *Proc. R. Soc. B Biol. Sci.*, 287, 20202655.

Berkvens, R., Olivares, I.H., Mercelis, S., Kirkpatrick, L. & Weyn, M. (2019). Contact Detection for Social Networking of Small Animals. In: *Advances on P2P, Parallel, Grid, Cloud and Internet Computing*. Springer International Publishing, pp. 405–414.

Brambilla, A., Biebach, I., Bassano, B., Bogliani, G., von Hardenberg, A., Hardenberg, A. Von, *et al.* (2015). Direct and indirect causal effects of heterozygosity on fitness-related traits in Alpine ibex. *Proc. Biol. Sci.*, 282, 20141873.

Carvalho, J., Granados, J.E., López-Olvera, J.R., Cano-Manuel, F.J., Pérez, J.M., Fandos, P., *et al.* (2015). Sarcoptic mange breaks up bottom-up regulation of body condition in a large herbivore population. *Parasites and Vectors*, 8, 1–9.

Davis, S., Abbasi, B., Shah, S., Telfer, S., Begon, M. & Davis, S. (2015). Spatial analyses of wildlife contact networks. *J. R. Soc. Interface*, 12, 20141004.

Dubey, J.P., Brown, J., Ternent, M., Verma, S.K., Hill, D.E., Cerqueira-Cézar, C.K., *et al.* (2016). Seroepidemiologic study on the prevalence of Toxoplasma gondii and Trichinella spp. infections in black bears (Ursus americanus) in Pennsylvania, USA. *Vet. Parasitol.*, 229, 76–80.

Foley, J., Rejmanek, D., Foley, C. & Matocq, M. (2016). Fine-scale genetic structure of woodrat populations (Genus: Neotoma) and the spatial distribution of their tick-borne pathogens. *Ticks Tick. Borne. Dis.*, 7, 243–253.

Foley, P., Foley, J., Sándor, A.D., Ionică, A.M., Matei, I.A., D'Amico, G., *et al.* (2017). Diversity of flea (siphonaptera) parasites on red foxes (Vulpes vulpes) in Romania. *J. Med. Entomol.*, 54, 1243–1250.

French, A.S., Zadoks, R.N., Skuce, P.J., Mitchell, G., Gibbs, D.K.G. & Taggart, M.A. (2019). Habitat and host factors associated with liver fluke (*Fasciola hepatica*) diagnoses in wild red deer (*Cervus elaphus*) in the Scottish Highlands. *Parasit. Vectors*, 1–14.

Galal, L., Schares, G., Stragier, C., Vignoles, P., Brouat, C., Cuny, T., *et al.* (2019). Diversity of Toxoplasma gondii strains shaped by commensal communities of small mammals. *Int. J. Parasitol.*, 49, 267–275.

Gras, P., Knuth, S., Börner, K., Marescot, L., Benhaiem, S., Aue, A., *et al.* (2018). Landscape structures affect risk of canine distemper in urban wildlife. *Front. Ecol. Evol.*, 6, 1–16.

Greigert, V., Brion, N., Lang, C., Regnard, P., Pfaff, A.W., Abou-Bacar, A., *et al.* (2019). Cestode infections in non-human primates suggest the existence of zoonotic cycles in the area surrounding the Strasbourg primatology center. *Parasite*, 26.

Hammond, T.T., Hendrickson, C.I., Maxwell, T.L., Petrosky, A.L., Palme, R., Pigage, J.C., *et*

al. (2019). Host biology and environmental variables differentially predict flea
abundances for two rodent hosts in a plague-relevant system. *Int. J. Parasitol. Parasites*
Wildl., 9, 174–183.

Hayward, A.D., Nussey, D.H., Wilson, A.J., Berenos, C., Pilkington, J.G., Watt, K. a., *et al.*
(2014). Natural Selection on Individual Variation in Tolerance of Gastrointestinal
Nematode Infection. *PLoS Biol.*, 12, 1–13.

Heddergott, M., Frantz, A.C., Stubbe, M., Stubbe, A., Ansorge, H. & Osten-Sacken, N.
(2017). Seroprevalence and risk factors of Toxoplasma gondii infection in invasive
raccoons (Procyon lotor) in Central Europe. *Parasitol. Res.*, 116, 2335–2340.

Ihle-Soto, C., Costoya, E., Correa, J.P., Bacigalupo, A., Cornejo-Villar, B., Estadella, V., *et*
al. (2019). Spatio-temporal characterization of Trypanosoma cruzi infection and discrete
typing units infecting hosts and vectors from non-domestic foci of chile. *PLoS Negl.*
Trop. Dis., 13, 1–19.

Jaffe, D.A., Chomel, B.B., Kasten, R.W., Breitschwerdt, E.B., Maggi, R.G., McLeish, A., *et*
al. (2018). Bartonella henselae in small Indian mongooses (Herpestes auropunctatus)
from Grenada, West Indies. *Vet. Microbiol.*, 216, 119–122.

Kosoy, M., Reynolds, P., Bai, Y., Sheff, K., Enscore, R.E., Montenieri, J., *et al.* (2017).
Small-Scale Die-Offs in Woodrats Support Long-Term Maintenance of Plague in the
U.S. Southwest. *Vector-Borne Zoonotic Dis.*, 17, 635–644.

Lazenby, B.T., Tobler, M.W., Brown, W.E., Hawkins, C.E., Hocking, G.J., Hume, F., *et al.*
(2018). Density trends and demographic signals uncover the long-term impact of
transmissible cancer in Tasmanian devils. *J. Appl. Ecol.*, 55, 1368–1379.

Lynsdale, C.L., Mumby, H.S., Hayward, A.D., Mar, K.U. & Lummaa, V. (2017). Parasite-
associated mortality in a long-lived mammal: Variation with host age, sex, and
reproduction. *Ecol. Evol.*, 7, 10904–10915.

Råberg, L., Hagström, Andersson, M., Bartkova, S., Scherman, K., Strandh, M., *et al.* (2017).
Evolution of antigenic diversity in the tick-transmitted bacterium Borrelia afzelii: a role
for host specialization? *J. Evol. Biol.*, 30, 1034–1041.

Rozins, C., Silk, M.J., Croft, D.P., Delahay, R.J., Hodgson, D.J., McDonald, R.A., *et al.*
(2018). Social structure contains epidemics and regulates individual roles in disease
transmission in a group-living mammal. *Ecol. Evol.*, 1–12.

Smallbone, W.A., Chadwick, E.A., Francis, J., Guy, E., Perkins, S.E., Sherrard-Smith, E., *et*
al. (2017). East-West Divide: Temperature and land cover drive spatial variation of
Toxoplasma gondii infection in Eurasian otters (Lutra lutra) from England and Wales.
Parasitology, 144, 1433–1440.

Sokos, C., Touloudi, A., Giannakopoulos, A., Papaspyropoulos, K.G., Tsokana, C.N.,
Birtsas, P., *et al.* (2018). Population density, roads and altitude influences on spatial
distribution of hares positive to EBHSV. *Basic Appl. Ecol.*, 31, 82–91.

de Sousa, K.C.M., Fernandes, M.P., Herrera, H.M., Benevenute, J.L., Santos, F.M., Rocha,
F.L., *et al.* (2017). Molecular detection of Hepatozoon spp. in domestic dogs and wild
mammals in southern Pantanal, Brazil with implications in the transmission route. *Vet.*
Parasitol., 237, 37–46.

Sweeny, A.R., Clerc, M., Pontifes, P.A., Venkatesan, S., Babayan, S.A. & Pedersen, A.B.
(2019). Supplemented nutrition increases immunity and drug efficacy in a natural host-
helminth system, 1–43.

Tufts, D.M. & Diuk-Wasser, M.A. (2018). Transplacental transmission of tick-borne Babesia
microti in its natural host Peromyscus leucopus. *Parasites and Vectors*, 11, 1–9.

Turner, W.C., Kausrud, K.L., Krishnappa, Y.S., Cromsigt, J.P.G.M., Ganz, H.H., Mapaure,
I., *et al.* (2014). Fatal attraction: Vegetation responses to nutrient inputs attract
herbivores to infectious anthrax carcass sites. *Proc. R. Soc. B Biol. Sci.*, 281.

850