# Structural insights into mRNA reading frame regulation by tRNA modification and slippery codon-anticodon pairing

Eric D. Hoffer[1], Samuel Hong[1], S. Sunita[1], Tatsuya Maehigashi[1], Ruben L. Gonzalez, Jr.[2]

Paul Whitford[3] and Christine M. Dunham[1]

[1]Department of Biochemistry, Emory University School of Medicine, Atlanta, GA, USA

[2]Department of Chemistry, Columbia University, New York, NY, USA

[3]Department of Physics, Northeastern University, Boston, MA, USA

Contact: Christine M. Dunham, christine.m.dunham@emory.edu, phone +1-404-712-1756, fax +1-404-727-2738.

**ABSTRACT**

Modifications in the tRNA anticodon, adjacent to the three-nucleotide anticodon, influence translation fidelity by stabilizing the tRNA to allow for accurate reading of the mRNA genetic code. One example is the N1-methylguaonosine modification at guanine nucleotide 37 ($m^1G37$) located in the anticodon loop, immediately adjacent to the anticodon nucleotides 34-36. The absence of $m^1G37$ in $tRNA^{Pro}$ causes +1 frameshifting on polynucleotide, slippery codons. Here, we report structures of the bacterial ribosome containing $tRNA^{Pro}$ bound to either cognate or slippery codons to determine how the $m^1G37$ modification prevents mRNA frameshifting. The structures reveal that certain codon-anticodon contexts and $m^1G37$ destabilize interactions of $tRNA^{Pro}$ with the peptidyl site, causing large conformational changes typically only seen during EF-G mediated translocation of the mRNA-tRNA pairs. These studies provide molecular insights into how $m^1G37$ stabilizes the interactions of $tRNA^{Pro}$ with the ribosome and the influence of slippery codons on the mRNA reading frame.

26 **IMPACT STATEMENT**

27 Chemical modifications near the tRNA anticodon and specific mRNA-tRNA pairs combine to control the

28 ribosomal three-nucleotide mRNA reading frame, essential for the sequential addition of amino acids into

29 polypeptide chains.

## INTRODUCTION

Post-transcriptionally modified RNAs, including ribosomal RNA (rRNA), transfer tRNA (tRNA) and messenger RNA (mRNA), stabilize RNA tertiary structures during ribonucleoprotein biogenesis, regulate mRNA metabolism, and influence other facets of gene expression. RNA modifications located in the three-nucleotide anticodon of tRNAs contribute to accurate protein synthesis and expand tRNA coding capacity by facilitating Watson-Crick-like base-pairs with mRNA codons. The most commonly modified tRNA position is at nucleotide 34 and the modifications to this nucleotide are functionally important for gene expression by facilitating base-pairing interactions with the mRNA codon to expand the tRNA coding capacity (*1-3*). Other tRNA nucleotides also contain highly conserved nucleotide modifications, but their precise roles in protein synthesis are not fully understood. For example, nucleotide 37 is adjacent to the anticodon (Figure 1A) and is modified in >70% of all tRNAs (*4*). The two most common modifications at nucleotide 37 are 6-threonylcarbamoyladenosine ($t^6A$) and 1-methylguanosine ($m^1G$), which together account for more than 60% of nucleotide 37 modifications (*2*). Both the $t^6A37$ modification in $tRNA^{Lys}$ and the $m^1G37$ modification in $tRNA^{Pro}$ are thought to stabilize the tertiary structure of the anticodon loop for high-affinity binding to their cognate codons (*5-7*). The $m^1G37$ modification also prevents the ribosome from shifting out of the three-nucleotide mRNA codon frame ('frameshifting') to ensure the correct polypeptide is expressed (*8-11*). However, the molecular basis for how the $m^1G37$ modification maintains the mRNA reading frame is unknown.

The $m^1G37$ modification is present in >95% of all three $tRNA^{Pro}$ isoacceptors across all three domains of life (*2, 8*). The absence of $m^1G37$ in $tRNA^{Pro}$ causes high levels of +1 frameshifting on so-called 'slippery' or polynucleotide mRNA sequences where four nucleotides encode for a single proline codon (*8*). In the case of the $tRNA^{Pro}_{CGG}$ isoacceptor, a +1 slippery codon consists of the CCC proline codon and either an additional C or U to form a four-nucleotide codon, CCC-(U/C) (*12, 13*) (codons depicted 5'- to 3'). $tRNA^{Pro}_{CGG}$ lacking $m^1G37$ results in the ribosome being unable to distinguish a correct from an incorrect codon-anticodon interaction during decoding at the aminoacyl (A) site (*7, 14*). However, this miscoding event does not cause the shift in the mRNA frame in the A site despite the $tRNA^{Pro}_{CGG}$ anticodon stem-loop

nucleotides become more mobile (*14*). A post-decoding frameshift event is further supported by detailed

kinetic analyses (*11*). The absence of the $m^1G37$ modification in tRNA[Pro] causes both a ~5% frameshifting

frequency during translocation of the mRNA-tRNA pair from the A to the peptidyl (P) site and ~40%

frameshifting frequency once the mRNA-tRNA pair has reached the P site. When tRNA$_{CGG}^{Pro}$ lacks $m^1G37$

and decodes a +1 slippery codon, both the process of translocation and the unique environment of the P site

appear to contribute to the inability of the ribosome to maintain the mRNA frame.

62           In circumstances where mRNA frameshifting is caused by changes in the tRNA such as the absence

of modifications or changes in the size of the anticodon loop as found in frameshift suppressor tRNAs,

primer extension assays demonstrated that the shift into the +1 frame is observed upon direct tRNA binding

at the P site (*15, 16*). These studies clearly demonstrate that the nature of the interactions between the

frameshift-prone mRNA-tRNA pair and the ribosomal P site directly permit frameshifting. Furthermore,

the presence of a nascent polypeptide chain, the acylation status of the tRNA (deacylated or aminoacylated),

and even the presence of a tRNA in the A site do not contribute to the ability of the tRNA to cause

frameshifting. Structural studies of frameshift-prone or suppressor tRNAs have provided molecular insights

into how such tRNAs may dysregulate mRNA frame maintenance (*14, 17, 18*). Frameshift suppressor

tRNA[SufA6] contains an extra nucleotide in its anticodon loop and undergoes high levels of +1 frameshifting

on slippery proline codons (*8, 13*). The structure of the anticodon stem-loop (ASL) of the tRNA[SufA6] bound

directly at the P site revealed that its anticodon engages the slippery CCC-U proline codon in the +1 frame

(*18*). Further, the full-length tRNA[SufA6] bound at the P site reveals that the small 30S subunit head domain

swivels and tilts, a similar movement to that caused by GTPase elongation factor-G (EF-G) upon

translocation of the tRNAs through the ribosome (*19-22*). The process of mRNA-tRNA translocation is

coupled to this head domain swivel and tilt which is distinct from other conformational changes the

ribosome undergoes including intersubunit rotation (*21, 23-27*). However, the 70S-tRNA[SufA6] structure is

in the absence of EF-G, suggesting that a +1 frameshift event caused by frameshift-prone tRNAs

dysregulates some aspect of translocation. Although these studies demonstrate that +1 frameshift-prone

tRNAs are good model systems to uncover mechanisms by which both the ribosome and the mRNA-tRNA

pair contribute to mRNA frame maintenance, it is unclear whether normal tRNAs lacking modifications

result in +1 frameshifting in the same manner. To address this question, here we solved six 70S structures

of tRNA$_{CGG}^{Pro}$ in the presence or absence of the m$^1$G37 modification and bound to either cognate or slippery

codons in the P site. Our results define how the m$^1$G37 modification stabilizes the interactions of tRNA$_{CGG}^{Pro}$

with the ribosome. We further show that ribosomes bound to mRNA-tRNA pairs that result in +1

frameshifts promoted by tRNA$_{CGG}^{Pro}$ result in large conformational changes of the 30S head domain,

consequently biasing the tRNA-mRNA pair towards the E site.

**RESULTS**

**A near-cognate interaction between ASL$^{Pro}$ and the slippery CCC-U proline codon *alone* causes a**

**shift into the +1 frame.** To address whether tRNA$_{CGG}^{Pro}$ induces a +1 frameshift by a similar mechanism as

tRNA$^{SufA6}$, we determined two X-ray structures of ASL$^{Pro}$ decoding either a cognate CCG or a near-cognate,

slippery CCC-U codon bound at the P site (Figure 1 and Tables 1 and 2). We used chemically synthesized

ASL$^{Pro}$ (17 nucleotides) to ensure G37 is fully methylated at the N1 position (Figure 1A). The structure of

P-site ASL$^{Pro}$ interacting with a cognate CCG proline codon was solved to a resolution of 3.1 Å and reveals

that the three nucleotides of the anticodon (nucleotides G36-G35-C34) form three Watson-Crick base-pairs

with the C$_{+1}$-C$_{+2}$-G$_{+3}$ mRNA codon, respectively (Figure 1A) (mRNA nucleotides are numbered starting at

+1 from the P-site codon). The anticodon interacts with the codon in the canonical or 0 mRNA frame

indicating a frameshift has not occurred.

We next asked how ASL$^{Pro}$ interacts in the P site with a near-cognate CCC-U codon, a slippery

codon known to facilitate +1 frameshifts (*8, 11*). In this X-ray structure solved to a resolution of 3.4 Å, we

find two different conformations of the codon-anticodon interaction in the two molecules of the

crystallographic asymmetric unit (Figures 1C and D and Table 2 and Video 1). Although it is common for

one 70S ribosome to contain better electron density than the other in the asymmetric unit in this particular

crystal form (*28*), to our knowledge, it is not common to obtain two different conformational states of the

same functional complex. In the first ribosome molecule, nucleotides G36 and G35 of ASL$^{Pro}$ form Watson-

Crick base-pairs with the first two nucleotides of the 0-frame mRNA codon (Figure 1C). A mismatch

$C_{+3} \cdot C34$ forms at the third base-pair position of the codon-anticodon interaction and contains a single

hydrogen bond between the Watson-Crick face of $C_{+3}$ (the N3 position) and C34 (the N4 position). This

codon-anticodon interaction is thus defined as near cognate because of the single mismatch. The distance

between the anticodon nucleotide C34 and the mRNA nucleotide $C_{+3}$ is increased in the P site as compared

to this same mRNA-tRNA pair in the A site (3.6 Å versus 3.1 Å (*14*)), indicating that the interaction has

weakened. Although this interaction is at the distance limit of a hydrogen bond, C34 and $C_{+3}$ are positioned

to form a *cis*-Watson-Crick pair, similar to the orientation observed in the A site (*14*) (Figure 1C-figure

supplement 1). The codon-anticodon interaction is in the 0 frame as indicated by the clear phosphate density

of the next mRNA nucleotide, $U_{+4}$ (Figure 1C-figure supplement 1). The mRNA used in these studies only

contained a single nucleotide after the three-nucleotide codon programmed in the P site to allow for the

unambiguous identification of the reading frame.

Strikingly, in the other ribosome molecule in the crystallographic asymmetric unit, the codon-

anticodon interaction is in the +1 frame (Figure 1D and Video 1). The first two nucleotides of the anticodon

(G36-G35) form Watson-Crick base-pairs with $C_{+2}$-$C_{+3}$ nucleotides of the mRNA codon, respectively. The

first nucleotide of the proline codon, C+1, no longer interacts with the tRNA and does not appear to make

any interactions with the ribosome. At the third position of the codon-anticodon interaction, a *cis*-Watson-

Crick $U_{+4} \cdot C34$ pair forms that contains a single hydrogen bond between the Watson-Crick face of $U_{+4}$ (the

O4 position) and C34 (the N4 position) (Figure 1D-figure supplement 1C). The absence of electron density

for a subsequent nucleotide in the mRNA confirms the +1 frame of the codon-anticodon interaction as this

is the last nucleotide of the mRNA (Figure 1D-figure supplement 1C). Therefore, although the m$^1$G37

modification is present in ASL$^{Pro}$, the shift in the +1 frame can still occur. These data indicate that the

presence of a near-cognate, slippery codon is sufficient to promote a +1 shift in the mRNA frame.

Consistent with our observations, biochemical studies of other near-cognate mRNA-tRNA pairs in the P

site that also show frameshifting in both directions on the mRNA (*29*).

**The absence of m$^1$G37 in tRNA$_{CGG}^{Pro}$ bound to a cognate CCG codon does not destabilize its interactions**

**with the ribosome.** We next asked how full-length tRNA$_{CGG}^{Pro}$ bound at the P site interacts with the mRNA.

Our recent structure of the ribosome with +1 frameshift suppressor tRNA$^{SufA6}$ indicated that although the

tRNA was positioned at the P site, when engaged with mRNA that causes +1 frameshifts, the tRNA moves

towards the E site and induces large conformational changes of the 30S subunit, specifically the head

domain (*18*). We wanted to directly compare these structures given that the +1 frameshift is induced by

different signals: an extra nucleotide in the anticodon loop in tRNA$^{SufA6}$ and the absence of m$^1$G37 in tRNA

$_{CGG}^{Pro}$. It is possible that tRNA$^{SufA6}$ causes this destabilization and thus conformational changes of the 30S head

domain because of the extra nucleotide in its ASL; such a mechanism would likely not apply for m$^1$G37 in

tRNA$_{CGG}^{Pro}$. As a control, we solved a structure of tRNA$_{CGG}^{Pro}$ containing the m$^1$G37 modification bound to a

cognate CCG codon in the ribosomal P site to a resolution of 3.2 Å (Figure 2A and Table 3 and Video 2).

tRNA$_{CGG}^{Pro}$ binds in the classical P/P state (bound to the P site on the 50S and the 30S) and interacts with the

proline codon in the 0 frame. Specifically, the G36-G35-C34 anticodon nucleotides form three Watson-

Crick base-pairs with the C$_{+1}$-C$_{+2}$-G$_{+3}$ proline codon nucleotides, respectively (Figure 2E-figure supplement

1A and 1B). The methyl group at position 1 of the nucleobase of G37 stacks with the nucleobase of

anticodon nucleotide G36 to form a canonical four-nucleotide stack between nucleotides 34-37. This

anticodon stack is important for productive interactions with both mRNA and the ribosome (*14, 30, 31*).

The P/P location of tRNA$_{CGG}^{Pro}$ is thus consistent with functional assays demonstrating that tRNA$^{Pro}$

isoacceptors do not appear to spontaneously move into the +1 frame on non-slippery codons (*8, 11*).

We next solved a 3.9 Å structure of tRNA$_{CGG}^{Pro}$ lacking the m$^1$G37 modification (G37 Δm$^1$) and

interacting with a cognate CCG proline codon in the P site (Figure 2B-figure supplements 1C and 1D and

Table 3 and Video 2). Although the m$^1$G37 modification was previously shown to be critical for high-

affinity binding and decoding of the cognate CCG codon at the A site (*7*), its influence on the overall

conformation of tRNA$_{CGG}^{Pro}$ in the P site appears to be minimal as the tRNA adopts a P/P orientation (Figure

2B). Notably, the mRNA remains in the 0 frame with three Watson-Crick base-pair interactions between

the codon and the anticodon (Figure 2F). Similar to the structural studies of the ASL$^{Pro}$ in the 0 frame

(Figures 1B and D), the mRNA used in these studies contains an additional nucleotide after the P-site CCG

codon (Figures 1; Figure 2B and Table 1). The mRNA has clear phosphate density for the single A-site

nucleotide indicating the mRNA remains in the 0 frame (Figure 2-figure supplement 1D). The absence of

the $m^1G37$ modification correlates with a slightly weaker U32-A38 pairing when compared to modified

$tRNA^{Pro}_{CGG}$ (Figure 2-figure supplement 2B). These structures suggest that the absence of 1-methyl

modification on G37 alone is not sufficient to destabilize the tRNA in the P site of the ribosome. This

observation is also consistent with toeprint analyses demonstrating that $tRNA^{Pro}_{GGG}$ lacking $m^1G37$ on a non-

slippery codon remained in the 0 frame (*11*).

**A near-cognate, slippery proline codon in the P site causes $tRNA^{Pro}_{CGG}$ destabilization and 30S head**

**movement.** In an attempt to reconcile the exact role of the $m^1G37$ modification in $tRNA^{Pro}_{CGG}$ in the context

of a near-cognate, slippery codon, we next solved a ribosome structure in the presence of $m^1G37$ modified

$tRNA^{Pro}_{CGG}$ in the P site bound to a near-cognate, slippery CCC-U codon to a resolution of 3.2 Å (Figure 2C

and Table 3 and Video 2). Unexpectedly, $tRNA^{Pro}_{CGG}$ moves from the P site towards the E site, adopting a

position on the ribosome which we refer to as e*/E (*18*) (where lowercase e* signifies an intermediate

position between the 30S subunit P and the E sites and a 50S E-site position) (Figure 2C-figure supplement

1E). The e*/E position of $tRNA^{Pro}_{CGG}$ when bound to this near-cognate, slippery codon is different from the

classical P/P position that the same tRNA adopts when it interacts with a cognate CCG codon (compare

Figure 2A with 2C). The classical P/P $tRNA^{Pro}_{CGG}$ location on its cognate codon is consistent with dozens of

other P-site tRNA-ribosome structures. Therefore, the ability of $tRNA^{Pro}_{CGG}$ to move towards the E site when

bound to a slippery proline codon suggests that it is this near-cognate codon-anticodon interaction alone

that is sufficient to destabilize $tRNA^{Pro}_{CGG}$ (Figure 2C).

The e*/E position of the $tRNA^{Pro}_{CGG}$-mRNA pair is additionally coupled to large conformational

changes of the 30S head domain (Figure 3 and Video 2). 30S head motions accompanying canonical

translocation or caused by a frameshift-prone tRNAs have been previously described by our group and

others in terms of the degree of swivel and tilt (*18, 23-25, 32, 33*), but these two distinct movements have

never been analyzed separately (Figure 3). To more precisely describe this multidimensional motion, we

used previously reported procedures to define "swivel" as the movement of the head relative to the body

within a plane (*i.e.* pure rotation about a single axis defined by structures of the unrotated and swiveled

conformations (PDB codes 4V9D and 4V4Q, respectively)), while "tilt" describes any deviations from pure

rotation (*34*). Together, these more accurate calculations provide an unambiguous description of the full,

three-dimensional orientation of the head during translocation, where the tRNAs are found in intermediate

states (*24, 25*). The head domain swivels in a ~18° counterclockwise direction while also undergoing a ~5°

tilt away from the ribosome and perpendicular to the mRNA path (*23-25*) (the counterclockwise swivel is

defined with the ribosome viewed with E, P, A tRNA sites oriented from left to right as depicted in Figure

2). The head domain is comprised of 16S rRNA nucleotides 930-1380 and seven ribosomal proteins (S3,

S7, S9, S10, S13, S14, and S19) that collectively move as a rigid body during EF-G-mediated translocation

of the mRNA-tRNA pairs on the 30S (*23-25*). The same swivel and tilting of the head domain of e*/E tRNA

$_{CGG}^{Pro}$ occurs but in our structure lacks EF-G. This structure indicates that even with the m$^1$G37 modification,

interactions of tRNA$_{CGG}^{Pro}$ with the P site are destabilized when bound to a near-cognate codon, promoting its

movement towards the E site along with 30S head domain swiveling and tilting.

**The mRNA located in the 30S E and P sites is constricted when tRNA$_{CGG}^{Pro}$ adopts an e*/E position.** The

path of the mRNA on the ribosome accommodates three nucleotides, or a single codon, in each of the A, P

and E sites (Figures 4A and B). The ribosome interacts extensively with the mRNA in the A site to ensure

accurate decoding but interacts less substantially with the P-site codon, while the E-site mRNA codon is

the least monitored. The ribosome interacts with mRNA in the E site at two positions through non-sequence

specific interactions: 16S rRNA nucleotide G693 stacks with the first nucleotide of the E-site mRNA codon

and the G926 nucleobase contacts the phosphate of the third mRNA codon nucleotide (Figures 4B and C)

(*28*). The e* position of tRNA$_{CGG}^{Pro}$ signifies that the tRNA is closer to the E site than the P site on the 30S,

however it is similar to the pe/E tRNA position seen in translocation intermediate ribosome structures

containing EF-G (Figure 4D) (*25*). Only two nucleotides of the codon-anticodon interaction of the pe/E

tRNA are positioned in the E site leaving a pocket for an additional nucleotide to occupy upon full

translocation of the mRNA-tRNA pair (*24, 25*) (compare Figures 4C with 4D). In our structure containing

an e*/E tRNA$_{CGG}^{Pro}$ bound to a near-cognate, slippery codon, the first three nucleotides of the codon fully

occupy the E site (Figure 4E), despite the codon-anticodon adopting an intermediate position on the 30S

between the P and the E sites. This compaction of the mRNA means there is no space for the entire four

nucleotide codon in the E site upon translocation of the mRNA-tRNA pair. Additionally, the mRNA path

5' from the E site turns sharply by ~100° as the mRNA transits to the outside of the ribosome (Figure 4C).

In the case of e*/E tRNA$_{CGG}^{Pro}$, upon full translocation of the mRNA-tRNA pair to the E site, the first

nucleotide of the codon would be displaced from the E site to accommodate the three-nucleotide codon or

the last three nucleotides of the CCC-U codon. This placement would position the mRNA in the +1 frame.

The mRNA boundary between the P and the E sites on the 30S subunit appears to be demarcated

by 16S rRNA nucleotide G926 (Figures 4B and 4C) (*28*). G926 interacts with the phosphate of the +3

nucleotide of the E-site codon thus defining its 3'-end (Figure 4C). In our structure of tRNA$_{CGG}^{Pro}$ interacting

with a near-cognate, slippery codon, G926 instead interacts with U$_{+4}$ of the CCC-U codon rather than the

third nucleotide, thereby defining U$_{+4}$ as the end of the E-site codon (Figure 4E). Even in translocation

intermediate ribosome structures containing EF-G, G926 interacts with the +3 phosphate of the mRNA,

although this ribosome complex contains a cognate mRNA-tRNA pair (Figure 4D) (*25*). Comparison of

post-translocation (*35*), translocation intermediate (*24, 25*), and our structure presented here reveals that the

position of G926 in all three is very similar, while the position of the mRNA changes substantially (Figures

4C, D and E). In summary, although tRNA$_{CGG}^{Pro}$ is fully translocated to the E site in our new structure, the

four-nucleotide mRNA codon is compacted in the E site as if translocation has already taken place (Figure

4E).

**The G966-C1400 bridge connecting the 30S head and body domains is broken in the presence of a**

**near-cognate codon-anticodon interaction.** The ribosome minimally interacts with the mRNA-tRNA pair

in the P site: 16S rRNA C1400 contacts G966 and stacks with the third base-pair of the codon-anticodon

(Figure 5A) (*28*). The G966-C1400 pair remains intact during translocation of the mRNA-tRNA pair from

the P to the E site as part of the head domain, as evidenced by ribosome structures in translocation

intermediate states containing EF-G (Figure 5B). (*24, 25*). In our structure containing e*/E tRNA$_{CGG}^{Pro}$ bound

to a near-cognate, slippery codon, the G966-C1400 interaction is ablated (Figure 5C). Since the G966-

C1400 interaction is effectively the bridge between the head and body domains, this disruption can be

attributed to a dysregulation caused by this spontaneous translocation event.

**The absence of the m$^1$G37 modification in tRNA$_{CGG}^{Pro}$ coupled with a near-cognate, slippery codon-**

**anticodon interaction causes complete disordering of the 30S head domain.** Finally, we solved a 4.1 Å

structure of tRNA$_{CGG}^{Pro}$ lacking m$^1$G37 ($\Delta$m$^1$) interacting with a near-cognate, slippery CCC-U codon at the P

site. The electron density for majority of the tRNA$_{CGG}^{Pro}$ on the 50S subunit is well-resolved, indicating that,

like the analogous structure solved in the presence of the m$^1$G37 modification (Figure 2C), the tRNA has

also moved towards the E site on the 50S subunit (Figure 2D). In contrast, the electron density is weak the

tRNA$_{CGG}^{Pro}$ ASL and E-site mRNA indicating these regions are dynamic in the context of slippery codon and

in the absence of m$^1$G37 modification (Figure 2D-figure supplement 2G). Disordering of tRNA$^{Pro}$ G37

($\Delta$m$^1$) starts approximately at the beginning of the anticodon loop at nucleotides U32-A38.

Correspondingly, the 30S head domain region is also disordered (Figure 2D-figure supplement 3D),

whereas the only region of the 30S body domain with poor electron density is the P/E loop nucleotides

G1338 and A1339. Taken together, in this mRNA-tRNA pairing that causes high levels of frameshifting,

interactions with the tRNA are destabilized in the P site resulting in the codon-anticodon adopting an e*

location and high mobility of the 30S head domain.

**DISCUSSION**

The importance of tRNA modifications in protein synthesis has been recognized for decades, yet their

precise roles outside the anticodon have been elusive. Modifications in the anticodon loop at position 37 of

tRNAs have been implicated in mRNA frame maintenance (*10, 36*), but how absence of a single

modification can dysregulate the mRNA frame remained unclear. Here, we elucidated the role of the $m^1G37$

modification in $tRNA_{CGG}^{Pro}$, which was known to be important in stabilizing stacking interactions with

anticodon nucleotides (*14*) and in the prevention of frameshifting (*8, 37*). We find that this single methyl

group influences the overall stability of $tRNA_{CGG}^{Pro}$ in an unexpected manner that causes large conformational

changes between the tRNA and the 30S head domain, a domain known to move extensively during

translocation of the tRNAs (*21, 23-25*). However, the methylation *alone* does not stabilize $tRNA_{CGG}^{Pro}$ on the

ribosome and, instead, its position is heavily influenced by interactions with a distinct slippery proline

codon. Our structures of different mRNA-$tRNA_{CGG}^{Pro}$ pairs on the ribosome allows the first detailed

mechanistic insight into how mRNA frame maintenance is regulated by both the $m^1G37$ modification and

the stability of mRNA-tRNA interaction.

The location on the ribosome where frameshifting can occur is likely dependent on the type of

frameshift (in the positive or negative mRNA direction) and whether tRNA or mRNA causes the frameshift

(*38-41*). The ribosome itself can prevent the frameshift through inherent differences in how the ribosome

interacts with the tRNA-mRNA complex at the different tRNA binding sites and such differences could

potentially influence frameshifting by relaxing interactions, for example, between the codon and anticodon.

While the codon-anticodon interaction located in the 30S A site is strictly monitored by conserved 16S

rRNA nucleotides to select for cognate tRNA (*42*), there are comparatively few interactions with the codon-

anticodon when positioned within the P and E sites, underscoring their different functional roles during the

translation cycle. This relative absence of interactions in the P and E sites provides an opportunity for the

mRNA to shift out of frame. $tRNA^{Pro}$ isoacceptors frameshift either during translocation from the A to the

P site or after translocation in the P site (*11, 43*). Therefore we sought to capture the interactions of $tRNA^{Pro}$

during a frameshift event. The structure of $ASL^{Pro}$ bound to a slippery codon in the P site reveals the codon-

anticodon interaction has shifted into the +1 frame indicating that the anticodon stem-loop interaction with

the mRNA codon alone is important (Figure 1D). This frameshift  is likely possible because interactions

with the P-site tRNA are limited to only 16S rRNA nucleotides G1338 and A1339 (called the P/E loop

nucleotides) with the anticodon stem and 16S rRNA C1400 with the anticodon nucleotide 34 (Figure 5A-

figure supplement 1). The P/E loop appears to indirectly enforce the mRNA frame by gripping the P-site

tRNA and these interactions are maintained as the tRNA moves from the P site to a hybrid P/E state (*44*)

and then to a translocation intermediate pe/E state (*24*). In the structures of ASL$^{Pro}$ interacting with the

mRNA in the 0 frame, there is well-resolved density for G1338 and A1339 (Figures 1B and C-figure

supplement 1A,1D,1G), while there is some disordering of these nucleotides in the context of ASL$^{Pro}$ bound

to the near-cognate, slippery codon in the +1 frame (Figure 1-figure supplements 1C,1F,1I). The inability

for G1338 and A1339 to grip the anticodon stem when the codon-anticodon is in the +1 frame likely results

in the destabilization of full-length tRNA$^{Pro}$ in these frameshift competent contexts (Figures 2C and D).

Simulations of the ribosome undergoing large movements of the head domain during translocation

implicate nucleotides G1338 and A1339 in the coupling of 30S head dynamics and displacement of the P-

site tRNA towards the E site (*34*). Therefore, the lack of gripping by these nucleotides when a frameshift

event is located in the P site appears to bias tRNA$^{Pro}$ towards the E site which, in turn, causes the 30S head

domain to swivel and tilt as if EF-G were bound.

In addition to P/E loop nucleotides G1338 and A1339, 16S rRNA nucleotide G966 is also part of

the 30S head domain and stacks with wobble nucleotide 34 of the P-site tRNA anticodon (Figure 5A). The

30S head is connected to the body domain via the interactions of G966 with nucleotide C1400; the G966-

C1400 pair remains stacked beneath the P-site tRNA and follows the tRNA as it moves to a hybrid P/E

state and to a translocation pe/E intermediate state upon EF-G binding (Figure 5B) (*24, 25, 44*). In the case

of P-site ASL$^{Pro}$ interacting with a near-cognate, slippery codon in the 0 or +1 frame, *cis*-Watson-Crick

interactions form between C$_{+3}$•C34 and U$_{+4}$•C34, respectively (Figure 1-figure supplements 1K,1L). Both

interactions appear to result in reduced stacking with C1400. In structures of full-length tRNA$^{Pro}$-mRNA

pairs that cause +1 frameshifting and destabilization at the P site, the G966-C1400 interaction is broken

(Figure 5C). Together, these results suggest a connection between the reduced interactions of G1338-A1339

and G966-C1400 with the tRNA that appear to influence the movement of the P-site tRNA towards the E

site.

Although neither C1400 nor G966 have previously been implicated in mRNA frame maintenance,

there is a functional link between the P/E loop nucleotides G1338 and A1339, and C1400-G966. 16S rRNA

nucleotides in the P site are generally more tolerable of mutations than the A-site 16S rRNA, although

substitution of G966 substantially reduces ribosome activity to ~10% despite the mutation not being lethal

(*45*). This G966 mutant is suppressed by a G1338A mutation, indicating that the G1338A mutation can

stabilize interactions with the P-site tRNA even in the absence of the C1400-G966 interaction.

The interactions observed between the codon and anticodon of ASL$^{Pro}$ bound to either cognate or

near-cognate codons reveal the process of shifting into the +1 frame (Figure 1D). In the context of full-

length mRNA-tRNA pairs that cause +1 frameshifting, both pairs move from the P site to occupy a position

between the E and the P site (Figures 2C and D). This mRNA-tRNA placement is similar to the translocation

intermediate containing EF-G (*24*). In the structure presented containing an e*/E tRNA$^{Pro}$ with the EF-G

containing structures containing an pe/tRNA, the tRNA moves slightly away from the mRNA. In the e*/E

tRNA$^{Pro}_{CGG}$ structure, the first position of the codon-anticodon forms a Watson-Crick interaction but the

second and third nucleotides of the codon-anticodon are not within hydrogen bonding distances (Figure

2G). Interestingly, in the EF-G bound translocation intermediate structure, the codon-anticodon also is not

within bonding distance (*24*). These results suggest that it is the disruptive nature of moving between tRNA

binding sites that perturbs the interactions between the codon and anticodon and these interactions likely

reform once the transition to the next tRNA binding site is completed.

30S head domain conformational changes were first captured in ribosome structures of tRNAs

moving between the A to the P sites (ap/ap state) and between the P and the E sites (pe/E state) on the 30S

in the presence of EF-G (*24, 25, 44*). Upon EF-G binding, the head domain is predicted to first tilt and then

swivel in a counterclockwise manner (as viewed when looking at the ribosome with E, P, A tRNA sites

from left to right as depicted in Figure 2) to translocate the two tRNAs to final E/E and P/P positions (*21,*

*34*). The departure of EF-G and reverse tilting of the head back towards the intersubunit space is followed

by clockwise swivel, the rate-limiting step for translocation (*21*). Frameshift-prone tRNAs, such as tRNA

$^{Pro}_{CGG}$, cause spontaneous head swiveling and tilting once the tRNA occupies the P site after the mRNA

frameshift has occurred. The inability of the ribosome to hold the P-site tRNA is influenced by weakening

of the interactions of the P/E loop with the anticodon stem and by disruption of the C1400-G966 interaction,

both events likely major contributors to the dysregulation of the 30S head domain. In other words, the head

domain is unable to maintain extensive interactions with the P-site tRNA and this failure leads to

spontaneous translocation. Single molecule FRET (smFRET) studies of translocation also show that upon

tRNA movement from the P to the E site, there is increased dissociation of the tRNA from the ribosome,

bypassing the post-translocation E/E state (*21*). These data seem to indicate that even during canonical

translocation from the P to the E site, this is a highly dynamic process which leads to destabilized tRNA.

Consistent with these observations, we would expect that once EF-G fully translocates $tRNA_{CGG}^{Pro}$ to the post

E/E state, there may be further decrease in interaction between the tRNA and ribosome. Additionally, our

structures show 30S head swiveling and tilting, in the absence of EF-G, concurrent with the +1 frameshift

event would likely prevent EF-G from binding. EF-G residues located in the tip of domain IV interact with

the anticodon stem-loop of the A-site tRNA during translocation (*24, 35*) and mutations in this domain slow

the rate of translocation (*19, 46*). It is therefore an enticing hypothesis that EF-G may have an additional

role in mRNA frame maintenance in addition to its established function in translocation.

Although all three prokaryotic $tRNA^{Pro}$ isoacceptors have the capacity to frameshift, whether they

use similar mechanisms of action is unknown (*8, 11, 13*). The $m^1G37$ modification minimally influences

frameshifting in proline isoacceptor $tRNA_{GGG}^{Pro}$ (*proL*) in contrast to $tRNA_{cmo^5UGG}^{Pro}$ (*proM*) (*11*). An additional

difference is the dependency on elongation factor-P (EF-P) to reconcile +1 frameshifts. EF-P binds at the

E site to overcome ribosome stalling induced by poly-proline codons (*47, 48*) and is critical in suppressing

frameshifts on poly-proline stretches (*11*). While EF-P reduces +1 frameshifts with $tRNA_{GGG}^{Pro}$ G37 ($\Delta m^1$) to

an equivalent frequency as native $tRNA_{GGG}^{Pro}$, there is little influence on isoacceptor $tRNA_{cmo^5UGG}^{Pro}$ in the absence

or presence of $m^1G37$ (*11*). These mechanistic differences may be due to a combination of the codon-

anticodon pairings on slippery codons ($tRNA_{cmo^5UGG}^{Pro}$ and $tRNA_{GGG}^{Pro}$ are cognate with slippery codons while

$tRNA_{CGG}^{Pro}$ is near-cognate (*49*)) and/or the influence of other modifications such as the $cmo^5$-U34

modification in $tRNA_{cmo^5UGG}^{Pro}$ (*50*). Our structures show that regardless of the $m^1G37$ modification status of

tRNA$_\text{CGG}^\text{Pro}$, the near-cognate slippery codon causes the tRNA to be destabilized, biasing its position towards

the E site, which we predict is indicative of high levels of frameshifting. The possible synergistic effects of

cmo$^5$-U34 and m$^1$G37 in tRNA$_\text{cmo}^\text{Pro}{}_\text{5UGG}$ in preventing frameshifts is unclear, but tRNA$_\text{UGG}^\text{Pro}$ lacking all

modifications exhibits higher levels of frameshifting than tRNA$_\text{UGG}^\text{Pro}$ lacking only the m$^1$G37 modification

(*43*). Also, m$^1$G37 in tRNA$_\text{cmo}^\text{Pro}{}_\text{5UGG}$ may prevent movement to the e*/E position where it may not be an optimal

substrate for EF-P, which would be consistent with kinetic analyses (*11*). Since unmodified tRNA$_\text{UGG}^\text{Pro}$ would

be near-cognate on a slippery sequence without the cmo$^5$-U34, it is reasonable to hypothesize that the major

species of unmodified tRNA$_\text{CGG}^\text{Pro}$ would also frameshift, during translocation to the P site or in the P site, in

agreement with the structural work presented here.

**MATERIALS AND METHODS**

**mRNA, ASL and ribosome purification.** ASL$^\text{Pro}$ containing a m$^1$G37 modified anticodon stem-loop (18

nucleotides) and mRNA (19 nucleotides) were purchased from GE Healthcare Dharmacon and dissolved

in 10 mM Tris-HCl, pH 7.0 and 5 mM MgCl$_2$. We used a chemically synthesized 17-nucleotide ASL to

ensure complete m$^1$G modification at position 37, as previously used to examine interactions in the A site

(*14*). Purification of *Thermus thermophilus* 70S ribosomes was performed as previously described (*51*).

**tRNA$_\text{CGG}^\text{Pro}$ ligation and purification.** To ensure that tRNA$_\text{CGG}^\text{Pro}$ was methylated at tRNA nucleotide 37 (m$^1$G),

the 5' half of the tRNA (nucleotides 1-39) was chemically synthesized (GE Healthcare Dharmacon) and

enzymatically ligated to the chemically synthesized 3' half following established protocols (*52, 53*). Briefly,

T4 polynucleotide kinase (NEB) was used to phosphorylate the 5' end of the 3' tRNA half and was heat

inactivated. The 5' and 3' halves of each tRNA were then mixed and annealed in the T4 RNA ligase buffer

by heating to 80 ºC for 5 mins and slow cooling on the heat block to room temperature. T4 RNA ligase

(NEB) was added to the reaction at 37 ºC for 18 hours. The ligation reaction was run on a 12% denaturing

8M urea-polyacrylamide gel and the ligated fragment was excised and purified using a modified crush and

soak method (*53*). The RNA was ethanol precipitated, the pellet was thoroughly air dried and resuspended

in 10 mM Tris-HCl, pH 7.0 and 5 mM $MgCl_2$, followed by annealing at 70 ºC for 2 min and slow cooled

to room temperature on the benchtop. The purified full-length RNA was aliquoted and stored at -20 ºC.

**Structural studies.** $ASL^{Pro}$ complexes were formed with 3.5 μM 70S ribosomes programmed with 7 μM

mRNA for 6 min at 37 ºC. Then 22 μM $ASL^{Pro}$ was added and incubated for 30 min at 55 ºC. $tRNA^{Pro}_{CGG}$ and

$tRNA^{Pro}_{CGG}$ $\Delta m^1G$ complexes were formed with 3.5-4.4 μM of 70S ribosomes programmed with 7-10.5 μM

mRNA for 6 min at 37 ºC. Then 7.7-10.5 μM $tRNA^{Pro}_{CGG}$ or $tRNA^{Pro}_{CGG}$ $\Delta m^1G$ were incubated for 30 min at 55

ºC. Each ASL and tRNA were programmed in the ribosomal P site by designing the Shine-Dalgarno

sequence five nucleotides upstream of the P-site codon. Crystals were grown by sitting-drop vapor diffusion

in a 1:1 drop ratio of 100 mM Tris–acetate pH 7.6, 12–13% 2-methyl-2,4-pentanediol (MPD), 2.9-3.0%

polyethylene glycol (PEG) 20K, 100-150 mM L-arginine-HCl and 0.5-1 mM β-mercaptoethanol (β-Me).

$ASL^{Pro}$ crystals were cryoprotected step-wise in 100 mM Tris–acetate pH 7.6, 10 mM $NH_4Cl$, 50 mM KCl,

3.1% PEG 20K, 10 mM $Mg(CH_3COO)_2$, 6 mM β-Me and 20%-40% MPD, with the final cryoprotection

solution containing 30 μM $ASL^{Pro}$. Cryoprotection of $tRNA^{Pro}_{CGG}$ and $tRNA^{Pro}_{CGG}$ $\Delta m^1G$ was accomplished

similarly to $ASL^{Pro}$, except with 3% PEG 20K and 20 mM $Mg(CH_3COO)_2$. Additionally, the final

cryoprotection solution for both $tRNA^{Pro}_{CGG}$ and $tRNA^{Pro}_{CGG}$ $\Delta m^1G$ did not contain ligand. All crystals were flash

frozen in liquid nitrogen for data collection.

X-ray diffraction data were collected at the Southeast Regional Collaborative Access Team (SER-

CAT) 22-ID beamline line and the Northeastern Collaborative Access Team (NE-CAT) ID24-C and ID24-

E beamlines at the Advanced Photon Source (APS). Data were integrated and scaled using the program

XDS (54). All structures of 70S-$ASL^{Pro}$ and $tRNA^{Pro}_{CGG}$ containing cognate mRNA were solved by molecular

replacement in PHENIX using coordinates from a 70S structure containing mRNA and tRNAs (Table S1)

(PDB code 4V6G) (55, 56). For the 70S complex containing an e*/E tRNA and the mRNA, the start model

was changed to $tRNA^{SufA6}$ bound to slippery sequence (PDB code 5VPP) (18). In this structure, the 30S

head domain is swiveled ~18º relative to the 50S. The structure was solved by molecular replacement in

PHENIX followed by iterative rounds of manual building in Coot (*57*). All figures were prepared in PyMOL

(*58*).

**Calculating 30S head movement.** The following protocol was used to describe the 30S head and body

domains in terms of Euler angles ($\phi,\psi,\theta$). Here, $\phi+\psi$ defines the net swivel angle, tilting is described by $\theta$

and the tilt direction is defined by $\phi$. Euler angles were calculated separately for the 16S body and head

domains as previously defined (*34*). Briefly, for all calculations, the P atoms of the "core" residues of the

23S rRNA and 16S head/body were considered. The core residues are defined as the set of atoms that were

found to have spatial root mean squared fluctuations of less than 1 Å in a 1 microsecond explicit-solvent

simulation (*59*). This includes 1351, 442 and 178 residues in the 23S rRNA, 16S body and 16S head,

respectively. To ensure that the calculated angles reflect global domain orientations, rather than local

deformations, reference configurations of the 23S, as well as the 16S head and body, were aligned to each

group of core residues, separately. To calculate 16S body swivel angles, each model was first aligned to a

crystallographic model of a classical unrotated ribosome (PDB codes 4V9D:DA and 4V9D:BA), where

alignment was based on the 23S rRNA core atoms. To define the orientation of the body, axes were

constructed based on the aligned orientations of residues 41, 127 and 911, which are used to define the

plane of rotation. The Euler angles are then calculated based on the orientational difference of the fitted

axes in the reference classical configuration and the structural model of interest. To describe the orientation

of the head, the 16S body was first aligned to the reference model (alignment based on core atoms) and

axes were defined based on residue 984, 940 and 1106. According to these definitions, models (4V9D:DA,

4V9D:BA) and (4V9D:CA, 4V9D:AA) define pure (tilt-free) body rotation, and models (4V9D:DA,

4V9D:BA) and (4V4Q:DB, 4V4Q:CA) define pure (tilt-free) head swivel. The classical model (4V9D:DA,

4V9D:BA) is defined as the zero-tilt, zero-rotation orientation.

**ACKNOWLEDGMENTS**

We thank Ha An Nguyen, Jacob Mattingly, Pooja Srinivas, Dr. Alexandra Kuzmishin Nagy and Dr. Graeme

L. Conn for critical reading of the manuscript, and the staff members of the NE-CAT and SER-CAT

beamlines for assistance during data collection. This work was supported by the NIH (R01GM093278 (to

CMD), NIH R01GM119386 (to RLG), and NSF MCB-1915843 (PW). CMD is a Burroughs Wellcome

Fund Pathogenesis of Infectious Diseases Fellow. The x-ray crystallography datasets were collected at the

NE-CAT beamlines, which are funded by the National Institute of General Medical Sciences from the NIH

(P41 GM103403), and at the SER-CAT beamlines. The Pilatus 6M detector on 24-IDC beamline is funded

by a NIH-ORIP HEI grant (S10 RR029205). This research used resources of the Advanced Photon Source,

a US Department of Energy (DOE) Office of Science User Facility operated for the DOE Office of Science

by Argonne National Laboratory under contract no. DE-AC02-06CH11357.

**FIGURE LEGENDS**

**Figure 1. The ability of ASL$^{Pro}$ to +1 frameshift in the peptidyl (P) site is dependent on the near-cognate mRNA codon.** (A) Secondary structure of the anticodon stem-loop (ASL) of tRNA$^{Pro}$. The m$^1$G37 modified nucleotide is shown in orange and the anticodon nucleotides (C34, G35, and G36) and the U32-A38 pairing are labeled. (B) Structure of 70S-ASL$^{Pro}$ bound to a cognate CCG codon in the P site with the codon in the 0 or canonical frame. All 2F$_o$-F$_c$ electron density maps shown in panels B-D are contoured at 1.0 sigma. (C,D) Structure of 70S-ASL$^{Pro}$ bound to a +1 slippery, near-cognate CCC-U codon shows the mRNA position is either in the 0 (panel C) or +1 frame (panel D) in the two 70S molecules in the crystallographic asymmetric unit. In either the 0 or +1 frame, a *cis* Watson-Crick interaction at the third base-pair forms (C$_{+3}$•C34 or U$_{+4}$•C34). mRNA numbering starts at +1 according to the first position in the P site.

**Figure 2. Identity of the mRNA proline codon regulates 30S head domain movement.** Overview of 70S ribosome-tRNA$^{Pro}$ complex structures: (A) tRNA$^{Pro}$ m$^1$G37 on a cognate CCG codon adopts a P/P orientation (in the P site on the 30S and 50S subunits); (B) tRNA$^{Pro}$ lacking the m$^1$G37 modification (G37 Δm$^1$) on a cognate CCG codon also adopts a P/P orientation; (C) tRNA$^{Pro}$ on a slippery, near-cognate CCC-U codon adopts an e*/E orientation; and (D) tRNA$^{Pro}$ lacking the m$^1$G37 (G37 Δm$^1$) on a slippery, near-cognate CCC-U codon adopts an e*/E orientation. In this complex (panel D), the 30S head domain and anticodon-codon interaction are disordered. In panels A-D, the 16S rRNA of the 30S head domain is removed for clarity. Zoomed in view of 2F$_o$-F$_c$ density of (E) the codon-anticodon for tRNA$^{Pro}$ on a cognate CCG codon, (F) tRNA$^{Pro}$ G37 Δm$^1$ on a cognate CCG codon in the P site, and (G) tRNA$^{Pro}$ G37 Δm$^1$ on a near-cognate CCC-U codon in position between the E and the P sites (e*). All 2F$_o$-F$_c$ electron density maps shown in panels E-G are contoured at 1.0 sigma.

**Figure 3. 30S head domain movement in the presence of a +1 frameshift-prone tRNA.** (A) Overview of the 70S ribosome containing an e*/E tRNA$^{Pro}$ bound to a slippery, near-cognate CCC-U codon. Shifts in

phosphate atom positions of the 30S head domain (16S rRNA nucleotides 930-1380) in this structure as compared to the unrotated 70S (PDB code 4V5C). (B) *Top*, same view as in panel A but showing only the tilt of the head domain . *Bottom*, a 90° rotated view showing tilting is downward resulting in movement of the head domain away from the body domain. (C) *Left*, the same view as in panel A with only the counterclockwise swivel of the head domain indicated (left). *Right*, a 90° horizontal rotated view indicates the swivel is greatest towards the subunit interface, close to e*/E tRNA$^{Pro}$ and on the surface.

**Figure 4. Frame-dependent conformations of the mRNA in the E and P sites.** (A) Overview of the 70S ribosome with (B) a zoomed in view of the tRNA-mRNA interaction in the A, P and E sites (PDB code 4V6F). 16S rRNA nucleotides G693 and G926 interact with the E-site codon-anticodon. (C) The normal path of the mRNA (grey) in a ribosome structure containing P/P and E/E tRNAs demonstrates only a three-nucleotide codon (nucleotides +1, +2 and +3) is accommodated in the E site (PDB code 4V5F). 16S rRNA G693 defines the starts of the E-site codon and interacts with the first nucleotide. As the mRNA leaves the E site, there is a 100° kink between the first nucleotide of the E-site codon (+1) and the -1 nucleotide (shaded in purple). Panel C is rotated ~180° relative to the view in panel B. (D) A translocation intermediate state induced by EF-G contains a tRNA positioned between the P and the E sites on the 30S (denoted "pe"). The pe/E tRNA has not undergone full translocation to the E site and thus only two nucleotides (+1 and +2) are located in the E site (PDB code 4W29). In this translocation intermediate state, there is space available to accommodate an additional nucleotide of the codon (shaded in red). (E) tRNA$^{Pro}$ bound to a +1 slippery CCC-U codon reveals that although the codon-anticodon pair has not been fully translocated, this placement of the mRNA is different as compared to normal translocation intermediates structures as shown in panel D. The additional nucleotide (+4) of the four-nucleotide codon is shown in green.

**Figure 5. The 16S rRNA G966-C1400 gate between the 30S head and body domains is disrupted during a frameshift event.** (A) In a post-translocation state containing E/E, P/P and A/A tRNAs, 16S

rRNA nucleotides G966 and C1400 interaction forms beneath the P-site tRNA (PDB code 4V5F). (B) The G966 and C1400 interaction persists during EF-G-mediated translocation as the position of 30S head domain nucleotide G966 shifts while 30S body nucleotide C1400 remains constant (PDB code 4W29). (C) In a ribosome undergoing a +1 frameshift induced by tRNA$^{Pro}$ and a slippery CCC-U codon, the G966-C1400 interaction is broken. The additional nucleotide (+4) of the four nucleotide codon is shown in green.

**TABLES**

**Table 1: RNAs used in this study.**

| tRNA$^{Pro}$ 5' half | 5'-CGGUGAUUGGCGCAGCCUGGUAGCGCACUUCGUUCGGm$^1$GA-3' |
|---|---|
| tRNA$^{Pro}$ 3' half | 5'-CGAAGGGGUCGGAGGUUCGAAUCCUCUAUCACCGACCA-3' |
| mRNA_cognate | 5'- GGCA<u>AGGAG</u>GUAAAA **CCGG**-3' |
| mRNA_slippery | 5'- GGCA<u>AGGAG</u>GUAAAA **CCCU**-3' |

The underlined nucleotides indicates the Shine-Dalgarno region while the bold nucleotides are the P-site codons.

**Table 2. Data collection and refinement statistics for 70S-ASL$^{Pro}$ structures.**

|  | 70S-ASL$^{Pro}$-CCG codon | 70S-ASL$^{Pro}$ CCC-U codon |
|---|---|---|
| **Data collection** | | |
| Space group | P2$_1$2$_1$2$_1$ | P2$_1$2$_1$2$_1$ |
| Wavelength (Å) | 0.9791 | 0.9791 |
| Cell dimensions | | |
| a, b, c (Å) | 209.79 451.91 621.58 | 210.12451.80 622.96 |
| α, β, γ (°) | 90, 90, 90 | 90, 90, 90 |
| Resolution (Å) | 49.70 - 3.10 (3.21 - 3.10) | 49.70 - 3.40 (3.52 - 3.40) |
| R$_{pim}$ (%) | 21.2(80.5) | 14.5(65.5) |
| I/σI | 4.7(1.1) | 5.9(1.3) |
| Completeness (%) | 97.99 (87.41) | 98.85 (94.87) |
| Redundancy | 3.7(3.1) | 4.2(3.4) |
| CC1/2 | 0.968(0.250) | 0.987(0.359) |
| | | |
| **Refinement** | | |
| Reflections | 1034968 (91757) | 797600 (75973) |
| R$_{work}$/R$_{free}$ (%) | 24.1/27.2 | 19.8/23.7 |
| No. atoms | 289313 | 290047 |
| B-factors (Å$^2$) | | |
| Overall | 90.37 | 103.79 |
| Macromolecule | 90.6 | 104.05 |
| Ligand/ion | 39.43 | 43.84 |
| R.m.s deviations | | |
| Bond lengths (Å) | 0.004 | 0.010 |
| Bond angles (°) | 0.83 | 0.98 |
| PDB ID | 6NTA | 6NSH |

Highest resolution shell is shown in parentheses.

**Table 3. Data collection and refinement statistics for 70S-tRNA$^{Pro}$ structures**

| | tRNA$^{Pro}$ m$^1$G37 CCG codon | tRNA$^{Pro}$ m$^1$G37 CCC-U codon | tRNA$^{Pro}$ G37 ($\Delta$m$^1$) CCG codon | tRNA$^{Pro}$ G37 ($\Delta$m$^1$) CCC-U codon |
|---|---|---|---|---|
| **Data collection** | | | | |
| Space group | P2$_1$2$_1$2$_1$ | P2$_1$2$_1$2$_1$ | P2$_1$2$_1$2$_1$ | P2$_1$2$_1$2$_1$ |
| Wavelength (Å) | 0.9792 | 0.9792 | 0.9792 | 0.9792 |
| Cell dimensions | | | | |
| a, b, c (Å) | 210.20 451.47 620.21 | 210.74 450.26 626.11 | 209.97 450.71 619.40 | 210.09 450.32 622.89 |
| α, β, γ (°) | 90, 90, 90 | 90, 90, 90 | 90, 90, 90 | 90, 90, 90 |
| Resolution (Å) | 49.20 - 3.20 (3.31 - 3.20) | 49.93 - 3.50 (3.63 - 3.50) | 49.82 - 3.97 (4.11 - 3.97) | 49.83 - 4.14 (4.288 - 4.14) |
| R$_{pim}$ (%) | 11.5(51.7) | 9.00(44.1) | 8.60(86.0) | 9.807 (95.5) |
| I/σI | 6.7(1.5) | 7.8(1.7) | 6.3(1.0) | 5.1 (1.0) |
| Completeness (%) | 99.11 (98.45) | 97.55 (89.72) | 98.60 (95.72) | 98.39 (95.63) |
| Redundancy | 5.9(4.4) | 4.2(2.2) | 14.1(10.0) | 6.5 (3.6) |
| CC1/2 | 0.991(0.377) | 0.996(0.464) | 0.998(0.313) | 0.998 (0.37) |
| | | | | |
| **Refinement** | | | | |
| Reflections | 951115 (93932) | 723555 (66052) | 4959167(47742) | 439195 (42360) |
| R$_{work}$/R$_{free}$ (%) | 22.8/25.6 | 23.4/25.6 | 22.8/25.5 | 24.8/29.4 |
| No. atoms | 291966 | 292039 | 291793 | 291185 |
| B-factors (Å$^2$) | | | | |
| Overall | 103.87 | 113.6 | 177.39 | 247.09 |
| Macromolecule | 104.12 | 113.9 | 177.77 | 247.54 |
| Ligand/ion | 42.04 | 37.71 | 72.42 | 123.12 |
| R.m.s deviations | | | | |
| Bond lengths (Å) | 0.011 | 0.010 | 0.007 | 0.007 |
| Bond angles (°) | 1.00 | 1.23 | 0.94 | 1.38 |
| PDB ID | 6NUO | 6NWY | 6O3M | 6OSI |

Highest resolution shell is shown in parentheses.

**VIDEO LEGENDS**

**Video 1. mRNA frameshifting of ASL$^{Pro}$ to +1 mRNA reading frame in the peptidyl (P) site is dependent on the slippery mRNA codon.** An overview of the 70S ribosome (50S is light cyan and the 30S in grey) showing how the anticodon stem-loop (ASL) of tRNA$^{Pro}$ (blue) interacts with either a cognate CCG codon or a slippery CCC-U codon (black). The m$^1$G37 modified nucleotide is shown in orange. A zoomed-in view of the anticodon nucleotides (G36, G35, and C34) interacting with the cognate C$_{+1}$, C$_{+2}$ and G$_{+3}$ mRNA respectively, in the 0 frame. All 2F$_o$-F$_c$ electron density maps shown are contoured at 1.0 sigma. A morph from the 0 frame structure to the structure of ASL$^{Pro}$ interacting with a slippery CCC-U codon is shown to demonstrate the movement in the mRNA into the +1 frame. The fourth nucleotide of the mRNA codon is shown in green. In either the 0 or +1 frame, a *cis* Watson-Crick interaction at the third base-pair forms (C$_{+3}$•C34 or U$_{+4}$•C34). mRNA numbering starts at +1 according to the first position in the P site.

**Video 2. Influence of m$^1$G37 and the slippery codon on +1 frameshifting and conformational changes 30S head domain.** An overview of the 70S ribosome (50S is light cyan and the 30S in grey) showing how tRNA$^{Pro}$ (blue) (+/- m$^1$G37) interacts with either a cognate CCG codon or a slippery CCC-U codon (black). A zoomed-in view of the anticodon nucleotides (G36, G35, and C34) interacting with the cognate C$_{+1}$, C$_{+2}$ and G$_{+3}$ mRNA respectively, in the 0 frame. All 2F$_o$-F$_c$ electron density maps shown are contoured at 1.0 sigma. The m$^1$G37 modified nucleotide is shown in orange and the U32-A38 pairing is indicated. A morph of the changes of the U32-A38 pairing when the tRNA interacts with either a cognate CCG or a slippery CCC-U codon is shown. An overview and a morph of P/P tRNA$^{Pro}$ bound to a cognate CCG codon to the e*/E site is shown along with movement of the 30S head domain is shown. Lastly, a zoomed in view shows a morph from a 0 frame interaction that forms in the P/P site to the codon-anticodon interaction as the pairing is in the e*/E site.

**REFERENCES**

1.  P. F. Agris, A. Narendran, K. Sarachan, V. Y. P. Vare, E. Eruysal, The Importance of Being Modified: The Role of RNA Modifications in Translational Fidelity. *Enzymes* **41**, 1-50 (2017).
2.  P. Boccaletto, M. A. Machnicka, E. Purta, P. Piatkowski, B. Baginski, T. K. Wirecki, V. de Crecy-Lagard, R. Ross, P. A. Limbach, A. Kotter, M. Helm, J. M. Bujnicki, MODOMICS: a database of RNA modification pathways. 2017 update. *Nucleic Acids Res* **46**, D303-D307 (2018).
3.  P. F. Agris, E. R. Eruysal, A. Narendran, V. Y. P. Vare, S. Vangaveti, S. V. Ranganathan, Celebrating wobble decoding: Half a century and still much is new. *RNA biology* **15**, 537-553 (2018).
4.  M. A. Machnicka, A. Olchowik, H. Grosjean, J. M. Bujnicki, Distribution and frequencies of post-transcriptional modifications in tRNAs. *RNA biology* **11**, 1619-1629 (2014).
5.  F. V. t. Murphy, V. Ramakrishnan, A. Malkiewicz, P. F. Agris, The role of modifications in codon discrimination by tRNA(Lys)UUU. *Nat Struct Mol Biol* **11**, 1186-1191 (2004).
6.  M. Sundaram, P. C. Durant, D. R. Davis, Hypermodified nucleosides in the anticodon of tRNA(Lys) stabilize a canonical U-turn structure. *Biochemistry* **39**, 15652 (2000).
7.  H. A. Nguyen, E. D. Hoffer, C. M. Dunham, Importance of a tRNA anticodon loop modification and a conserved, noncanonical anticodon stem pairing in tRNACGGProfor decoding. *J Biol Chem* **294**, 5281-5291 (2019).
8.  G. R. Bjork, P. M. Wikstrom, A. S. Bystrom, Prevention of translational frameshifting by the modified nucleoside 1-methylguanosine. *Science* **244**, 986-989 (1989).
9.  J. Li, B. Esberg, J. F. Curran, G. R. Bjork, Three modified nucleosides present in the anticodon stem and loop influence the in vivo aa-tRNA selection in a tRNA-dependent manner. *Journal of molecular biology* **271**, 209-221 (1997).
10. J. Urbonavicius, Q. Qian, J. M. Durand, T. G. Hagervall, G. R. Bjork, Improvement of reading frame maintenance is a common function for several tRNA modifications. *Embo J* **20**, 4863-4873 (2001).
11. H. B. Gamper, I. Masuda, M. Frenkel-Morgenstern, Y. M. Hou, Maintenance of protein synthesis reading frame by EF-P and m(1)G37-tRNA. *Nature communications* **6**, 7226 (2015).
12. G. E. Sroga, F. Nemoto, Y. Kuchino, G. R. Bjork, Insertion (sufB) in the anticodon loop or base substitution (sufC) in the anticodon stem of tRNA(Pro)2 from Salmonella typhimurium induces suppression of frameshift mutations. *Nucleic Acids Res* **20**, 3463-3469 (1992).
13. Q. Qian, J. N. Li, H. Zhao, T. G. Hagervall, P. J. Farabaugh, G. R. Bjork, A new model for phenotypic suppression of frameshift mutations by mutant tRNAs. *Mol Cell* **1**, 471-482 (1998).
14. T. Maehigashi, J. A. Dunkle, S. J. Miles, C. M. Dunham, Structural insights into +1 frameshifting promoted by expanded or modification-deficient anticodon stem loops. *Proc Natl Acad Sci U S A* **111**, 12740-12745 (2014).
15. S. S. Phelps, C. Gaudin, S. Yoshizawa, C. Benitez, D. Fourmy, S. Joseph, Translocation of a tRNA with an extended anticodon through the ribosome. *Journal of molecular biology* **360**, 610-622 (2006).
16. S. E. Walker, K. Fredrick, Recognition and positioning of mRNA in the ribosome by tRNAs with expanded anticodons. *Journal of molecular biology* **360**, 599-609 (2006).
17. C. E. Fagan, T. Maehigashi, J. A. Dunkle, S. J. Miles, C. M. Dunham, Structural insights into translational recoding by frameshift suppressor tRNASufJ. *RNA* **20**, 1944-1954 (2014).
18. S. Hong, S. Sunita, T. Maehigashi, E. D. Hoffer, J. A. Dunkle, C. M. Dunham, Mechanism of tRNA-mediated +1 ribosomal frameshifting. *Proc Natl Acad Sci U S A* **115**, 11226-11231 (2018).
19. M. V. Rodnina, A. Savelsbergh, V. I. Katunin, W. Wintermeyer, Hydrolysis of GTP by elongation factor G drives tRNA movement on the ribosome. *Nature* **385**, 37-41 (1997).
20. D. N. Ermolenko, H. F. Noller, mRNA translocation occurs during the second step of ribosomal intersubunit rotation. *Nat Struct Mol Biol* **18**, 457-462 (2011).

21. M. R. Wasserman, J. L. Alejo, R. B. Altman, S. C. Blanchard, Multiperspective smFRET reveals rate-determining late intermediates of ribosomal translocation. *Nat Struct Mol Biol* **23**, 333-341 (2016).

22. R. Belardinelli, H. Sharma, N. Caliskan, C. E. Cunha, F. Peske, W. Wintermeyer, M. V. Rodnina, Choreography of molecular movements during ribosome progression along mRNA. *Nat Struct Mol Biol* **23**, 342-348 (2016).

23. A. H. Ratje, J. Loerke, A. Mikolajka, M. Brunner, P. W. Hildebrand, A. L. Starosta, A. Donhofer, S. R. Connell, P. Fucini, T. Mielke, P. C. Whitford, J. N. Onuchic, Y. Yu, K. Y. Sanbonmatsu, R. K. Hartmann, P. A. Penczek, D. N. Wilson, C. M. Spahn, Head swivel on the ribosome facilitates translocation by means of intra-subunit tRNA hybrid sites. *Nature* **468**, 713-716 (2010).

24. J. Zhou, L. Lancaster, J. P. Donohue, H. F. Noller, Crystal structures of EF-G-ribosome complexes trapped in intermediate states of translocation. *Science* **340**, 1236086 (2013).

25. J. Zhou, L. Lancaster, J. P. Donohue, H. F. Noller, How the ribosome hands the A-site tRNA to the P site during EF-G-catalyzed translocation. *Science* **345**, 1188-1191 (2014).

26. W. Holtkamp, C. E. Cunha, F. Peske, A. L. Konevega, W. Wintermeyer, M. V. Rodnina, GTP hydrolysis by EF-G synchronizes tRNA movement on small and large ribosomal subunits. *EMBO J* **33**, 1073-1085 (2014).

27. R. Belardinelli, H. Sharma, F. Peske, W. Wintermeyer, M. V. Rodnina, Translocation as continuous movement through the ribosome. *RNA biology* **13**, 1197-1203 (2016).

28. M. Selmer, C. M. Dunham, F. V. t. Murphy, A. Weixlbaumer, S. Petry, A. C. Kelley, J. R. Weir, V. Ramakrishnan, Structure of the 70S ribosome complexed with mRNA and tRNA. *Science* **313**, 1935-1942 (2006).

29. H. S. Zaher, R. Green, Quality control by the ribosome following peptide bond formation. *Nature* **457**, 161-166 (2009).

30. H. Grosjean, D. G. Soll, D. M. Crothers, Studies of the complex between transfer RNAs with complementary anticodons. I. Origins of enhanced affinity between complementary triplets. *Journal of molecular biology* **103**, 499-519 (1976).

31. E. M. Gustilo, F. A. Vendeix, P. F. Agris, tRNA's modifications bring order to gene expression. *Curr Opin Microbiol* **11**, 134-140 (2008).

32. S. Mohan, J. P. Donohue, H. F. Noller, Molecular mechanics of 30S subunit head rotation. *Proc Natl Acad Sci U S A* **111**, 13325-13330 (2014).

33. J. Zhou, L. Lancaster, J. P. Donohue, H. F. Noller, Spontaneous ribosomal translocation of mRNA and tRNAs into a chimeric hybrid state. *Proc Natl Acad Sci U S A* **116**, 7813-7818 (2019).

34. K. Nguyen, P. C. Whitford, Steric interactions lead to collective tilting motion in the ribosome during mRNA-tRNA translocation. *Nature communications* **7**, 10586 (2016).

35. Y. G. Gao, M. Selmer, C. M. Dunham, A. Weixlbaumer, A. C. Kelley, V. Ramakrishnan, The structure of the ribosome with elongation factor G trapped in the posttranslocational state. *Science* **326**, 694-699 (2009).

36. C. Yarian, H. Townsend, W. Czestkowski, E. Sochacka, A. J. Malkiewicz, R. Guenther, A. Miskiewicz, P. F. Agris, Accurate translation of the genetic code depends on tRNA modified nucleosides. *J Biol Chem* **277**, 16391-16395 (2002).

37. T. G. Hagervall, T. M. Tuohy, J. F. Atkins, G. R. Bjork, Deficiency of 1-methylguanosine in tRNA from Salmonella typhimurium induces frameshifting by quadruplet translocation. *Journal of molecular biology* **232**, 756-765 (1993).

38. J. D. Dinman, Control of gene expression by translational recoding. *Advances in protein chemistry and structural biology* **86**, 129-149 (2012).

39. J. A. Dunkle, C. M. Dunham, Mechanisms of mRNA frame maintenance and its subversion during translation of the genetic code. *Biochimie* **114**, 90-96 (2015).

40. J. F. Atkins, G. Loughran, P. R. Bhatt, A. E. Firth, P. V. Baranov, Ribosomal frameshifting and transcriptional slippage: From genetic steganography and cryptography to adventitious use. *Nucleic Acids Res*, (2016).

41. N. Korniy, E. Samatova, M. M. Anokhina, F. Peske, M. V. Rodnina, Mechanisms and biomedical implications of -1 programmed ribosome frameshifting on viral and bacterial mRNAs. *FEBS letters* **593**, 1468-1482 (2019).

42. J. M. Ogle, F. V. Murphy, M. J. Tarry, V. Ramakrishnan, Selection of tRNA by the ribosome requires a transition from an open to a closed form. *Cell* **111**, 721-732 (2002).

43. H. B. Gamper, I. Masuda, M. Frenkel-Morgenstern, Y. M. Hou, The UGG Isoacceptor of tRNAPro Is Naturally Prone to Frameshifts. *Int J Mol Sci* **16**, 14866-14883 (2015).

44. J. A. Dunkle, L. Wang, M. B. Feldman, A. Pulk, V. B. Chen, G. J. Kapral, J. Noeske, J. S. Richardson, S. C. Blanchard, J. H. Cate, Structures of the bacterial ribosome in classical and hybrid states of tRNA binding. *Science* **332**, 981-984 (2011).

45. N. M. Abdi, K. Fredrick, Contribution of 16S rRNA nucleotides forming the 30S subunit A and P sites to translation in Escherichia coli. *Rna* **11**, 1624-1632 (2005).

46. A. Savelsbergh, N. B. Matassova, M. V. Rodnina, W. Wintermeyer, Role of domains 4 and 5 in elongation factor G functions on the ribosome. *Journal of molecular biology* **300**, 951-961 (2000).

47. S. Ude, J. Lassak, A. L. Starosta, T. Kraxenberger, D. N. Wilson, K. Jung, Translation elongation factor EF-P alleviates ribosome stalling at polyproline stretches. *Science* **339**, 82-85 (2013).

48. P. Huter, S. Arenz, L. V. Bock, M. Graf, J. O. Frister, A. Heuer, L. Peil, A. L. Starosta, I. Wohlgemuth, F. Peske, J. Novacek, O. Berninghausen, H. Grubmuller, T. Tenson, R. Beckmann, M. V. Rodnina, A. C. Vaiana, D. N. Wilson, Structural Basis for Polyproline-Mediated Ribosome Stalling and Rescue by the Translation Elongation Factor EF-P. *Mol Cell* **68**, 515-527 e516 (2017).

49. S. J. Nasvall, P. Chen, G. R. Bjork, The modified wobble nucleoside uridine-5-oxyacetic acid in tRNAPro(cmo5UGG) promotes reading of all four proline codons in vivo. *RNA* **10**, 1662-1673 (2004).

50. I. Masuda, R. Takase, R. Matsubara, M. J. Paulines, H. Gamper, P. A. Limbach, Y. M. Hou, Selective terminal methylation of a tRNA wobble base. *Nucleic Acids Res* **46**, e37 (2018).

51. Y. Zhang, S. Hong, A. Ruangprasert, G. Skiniotis, C. M. Dunham, Alternative Mode of E-Site tRNA Binding in the Presence of a Downstream mRNA Stem Loop at the Entrance Channel. *Structure* **26**, 437-445 e433 (2018).

52. L. D. Sherlin, T. L. Bullock, T. A. Nissan, J. J. Perona, F. J. Lariviere, O. C. Uhlenbeck, S. A. Scaringe, Chemical and enzymatic synthesis of tRNAs for high-throughput crystallization. *RNA* **7**, 1671-1678 (2001).

53. M. R. Stark, S. D. Rader, Efficient splinted ligation of synthetic RNA using RNA ligase. *Methods Mol Biol* **1126**, 137-149 (2014).

54. W. Kabsch, Xds. *Acta crystallographica* **66**, 125-132 (2010).

55. L. B. Jenner, N. Demeshkina, G. Yusupova, M. Yusupov, Structural aspects of messenger RNA reading frame maintenance by the ribosome. *Nat Struct Mol Biol* **17**, 555-560 (2010).

56. P. D. Adams, P. V. Afonine, G. Bunkoczi, V. B. Chen, I. W. Davis, N. Echols, J. J. Headd, L. W. Hung, G. J. Kapral, R. W. Grosse-Kunstleve, A. J. McCoy, N. W. Moriarty, R. Oeffner, R. J. Read, D. C. Richardson, J. S. Richardson, T. C. Terwilliger, P. H. Zwart, PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta crystallographica* **66**, 213-221 (2010).

57. P. Emsley, B. Lohkamp, W. G. Scott, K. Cowtan, Features and development of Coot. *Acta crystallographica* **66**, 486-501 (2010).

58. Schrodinger, LLC. (2010).

59. P. C. Whitford, S. C. Blanchard, J. H. Cate, K. Y. Sanbonmatsu, Connecting the kinetics and energy landscape of tRNA translocation on the ribosome. *PLoS Comput Biol* **9**, e1003003 (2013).