Impact of Rap-Phr system abundance on adaptation of *Bacillus subtilis*

Ramses Gallegos-Monterrosa[1,*], Mathilde Nordgaard Christensen[2,*], Tino Barchewitz[1], Sonja Köppenhöfer[1,6], B. Priyadarshini[2], Balázs Bálint[3], Gergely Maróti[4], Paul J Kempen[5], Anna Dragoš[2], Ákos T. Kovács[1,2,#]

[1] Terrestrial Biofilms Group, Institute of Microbiology, Friedrich-Schiller-University Jena, Jena, Germany

[2] Bacterial Interactions and Evolution Group, DTU Bioengineering, Technical University of Denmark, Kongens Lyngby, Denmark

[3] Seqomics Biotechnology Ltd., Mórahalom, Hungary

[4] Institute of Plant Biology, Biological Research Centre, Hungarian Academy of Sciences, Szeged, Hungary

[5] Department of Health Technology, Technical University of Denmark, 2800 Kongens Lyngby, Denmark

* Shared first authors

corresponding author

Present address:

[6] Department of Biology, Memorial University of Newfoundland, St. John's, NL, Canada

Short title: Role of Rap-Phr in *Bacillus subtilis* evolution

Keywords: *Bacillus subtilis*, Rap, sporulation, biofilm, phage, selection

Abstract

Microbes commonly display great genetic plasticity, which has allowed them to colonize all ecological niches on Earth. The Gram-positive model bacterium *Bacillus subtilis* is a soil-dwelling organism that can be isolated from a wide variety of environments. An interesting characteristic of this bacterium is its ability to form biofilms that display complex population heterogeneity: individual, clonal cells develop diverse phenotypes in response to different environmental conditions within the biofilm. Here, we scrutinized the impact that the number and variety of the Rap-Phr family of regulators and cell-cell communication modules of *B. subtilis* has on genetic adaptation and evolution. We examine how the Rap family of phosphatase regulators impacts sporulation in diverse niches using a library of single and double *rap-phr* mutants in competition under 4 distinct growth conditions. Using specific DNA barcodes and whole-genome sequencing, population dynamics were followed, revealing the impact of individual Rap phosphatases and arising mutations on the adaptability of *B. subtilis* to thrive, produce spores, and outcompete other strains in various environments. Furthermore, we demonstrate that the used selection regimes can drive the activation of the phi3Ts-SPβ prophage, which contains a Rap homologue and further impacts spore production.

Author Summary

Free-living bacteria inhabit constantly changing environments where they face a multitude of challenges. To survive these challenges bacteria have evolved various adaptive strategies, such a spore production and biofilm formation. *Bacillus subtilis* is a Gram-positive bacterium that regularly uses both of these mechanisms to adapt and thrive in various ecological niches, including drastic metabolic changes or even the interruption of reproduction in the case of spore production. Thus, an efficient regulatory process is required to ensure that cells have maximum ecological adaptability and fitness. Here, we use a competition approach to examine in detail the role that the entire family of Rap phosphatases plays in the adaptability of *B. subtilis,* and show that, although initially they seem to play redundant regulatory roles, each Rap phosphatase participates in fine-tuning of the genetic regulatory network of this bacterium, providing it with a high adaptability to diverse environments.

Introduction

In recent years, *Bacillus subtilis* has become a model organism for the study of biofilms and population differentiation due to its ability to develop diverse phenotypes within an isogenic population [1,2]. Even when grown in liquid mixed cultures, where environmental conditions are assumed to be homogeneous, *B. subtilis* cells can be found as single flagellated cells or as non-flagellated chained cells due to stochastic variation in the expression of motility-related genes [3]. This population heterogeneity further increases when *B. subtilis* develops a biofilm, i.e., cells commit to particular functions, such as biofilm matrix production, exoenzyme secretion, or spore formation [1,4]. The development of these different cell types is partially triggered by the variation in the environmental conditions that exist in diverse parts of the biofilms, which can then be seen as a collection of ecological microniches, each with its own type of specialized inhabitant [5–7].

B. subtilis possesses a complex regulatory network that leads the cells within the biofilm to generate a phenotypically heterogeneous population. This network is described to be mainly controlled by the master transcriptional regulators Spo0A, DegU, and ComA. The activity of these regulators depends on their phosphorylation status, which is controlled by the activity of specific kinases that can sense a wide array of environmental and intracellular signals, and phosphorylate their corresponding response regulators accordingly. DegU and ComA are phosphorylated by kinases DegS and ComP, respectively, while Spo0A can be activated by five different kinases that act through a phosphorelay formed by the response regulators Spo0F and Spo0B. Furthermore, the regulatory network includes multiple cross-talk mechanisms and regulatory feedback loops that contribute to its modulation by constantly monitoring the general metabolic state of each cell within the biofilm [2,8,9].

The population-heterogeneity regulatory network of *B. subtilis* is further controlled by a family of response regulator aspartyl-phosphate (Rap) phosphatases and their cognate phosphatase-regulator (Phr) peptides. The cytoplasmic Rap proteins exert their regulatory function by inhibiting the activity of their target regulator (Spo0F, DegU, or ComA) via dephosphorylation, or by directly blocking DNA binding. The Rap proteins are in turn inhibited by their cognate Phr peptides, which are produced as pre-Phr proteins that are exported to the extracellular milieu and cleaved to produce mature 5-6 amino acid Phr peptides. The Phr peptides are

imported back into the cell upon reaching threshold concentrations at high cell density and bind to their cognate Rap phosphatase, inducing conformational changes that inhibit its activity [10,11]. The *rap* and *phr* genes are usually found as pairs in the same genetic loci, with the *phr* genes following and slightly overlapping the corresponding *rap* genes, and expression of both genes being transcriptionally coupled [12–14].

The Rap-Phr regulatory pairs are highly prevalent in the *Bacillus* genus, with ca. 2700 *rap* genes recently reported to be distributed among 346 *Bacilli* genomes; from those, ca. 80 different putative *rap-phr* alleles were found in the *B. subtilis* group alone [15]. Only a small minority of the *B. subtilis* Rap phosphatases has been characterized, finding that they have high redundancy in their regulatory function: most of them target Spo0F, ComA, or both; and only one (RapG) has been described to act on DegU [11]. Interestingly, *B. subtilis* shows great genomic plasticity regarding *rap-phr* gene pairs; 127 recently compared strains of the *B. subtilis* group showed to have multiple and diverse *rap-phr* gene pairs, with an average of 11 *rap* genes per strain [15]. This genetic variation among *B. subtilis* strains is not superfluous: since the Rap phosphatases modulate the activity of the main regulators of population heterogeneity, it has been proposed that the Rap-Phr pairs serve to adjust this regulatory network of *B. subtilis* to the needs of particular ecological niches [16,17]. As an example, it has been shown that *B. subtilis* strains isolated from gastrointestinal tracts of diverse animals have diverse sporulation initiation rates, with some being able to start sporulating already during logarithmic growth phase. This variation is correlated to the presence or absence of specific Rap-Phr pairs, and thus it has been suggested that the precise combination of *rap-phr* gene pairs matches the particular sporulation needs of a given niche [16].

Since the Phr peptides function as quorum sensing molecules regulating production of costly public goods [10], bacterial social interactions and evolutionary dynamics have been suggested to influence the emergence and persistence of multiple Rap-Phr systems with redundant functions in *B. subtilis* [15,18]. This idea is supported by the fact that up to 75% of *rap-phr* gene pairs are located in sections of the genome related to mobile genetic elements (such as prophages and transposons), suggesting that Rap-Phr systems are commonly acquired by horizontal gene transfer among *Bacillus* strains [15]; moreover, at least one *B. subtilis* Rap phosphatase (RapI) is known to promote the propagation of the mobile genetic element that

contains it [19]. The genetic variation in Rap-Phr systems among *B. subtilis* strains complicates the understanding of the role that the whole set of Rap phosphatases plays in modulating the population-heterogeneity regulatory network of any particular *B. subtilis* strain. Furthermore, the best-known Rap-Phr systems have usually been studied independently from each other, in diverse genetic backgrounds, and using different experimental conditions [10,11]. Likewise, previous investigations have focused on different aspects of Rap-Phr regulation, e.g.: RapA and RapB have been mainly studied for their role in sporulation regulation [20], while RapC and RapF are known to regulate competence development [21,22]. Conspicuously, Rap-Phr systems have not been thoroughly investigated with regards to biofilm formation [11]; to the best of our knowledge, only RapP has been previously shown to affect their formation [23,24].

Here, we aimed at revealing the influence of cultivation time (2 versus 5 days) and condition (planktonic versus biofilm) on selection towards particular *rap-phr* mutants or combined double mutations. To address this complexity, we used a barcoding approach that allowed us to follow the relative abundance of 79 strains, wild type, single, and double *rap-phr* deletion mutants, within the populations subjected to various selective conditions.

Results

Multi-strain experimental conditions to determine fitness differences among *rap-phr* mutants of *B. subtilis*. We were interested in systematically determining the impact that each Rap-Phr system has on the population fitness of *B. subtilis*, particularly on sporulation within biofilms, and how absence or presence of different Rap phosphatases would affect the adaptability of *B. subtilis* to different growth conditions. We used *B. subtilis* DK1042 (hereafter WT), which is a transformable strain derived from the isolate NCIB 3610. Strains NCIB 3610 and DK1042 have not undergone the domestication process that other commonly used *B. subtilis* strains have. This domestication can lead to loss of genetic functions and regulation changes, including modifications to Rap-Phr systems [25,26]. The WT has 11 *rap* genes encoded in its genome (*rapA* to *rapK*), additionally, it also possesses one more Rap-Phr system (*rapP-phrP*) encoded in its pBS32 plasmid [11,23]. We created single knock-out mutant strains of all the *rap-phr* genes, and double knock-out mutant strains that lack two *rap-phr* gene pairs in all possible combinations. In all cases both the *rap* and *phr* genes were deleted from the genome or plasmid. All created strains and the WT were further tagged with a pre-identified specific DNA barcode: a randomly-generated 12 bp nucleotide sequence that was integrated into the *amyE* locus of each strain. We used an experimental competition approach to analyze how the different Rap-Phr system combinations would impact the adaptability of all mutant strains (and WT as control) to 2 different growth conditions: shaken liquid cultures, where cells would multiply in a planktonic state; or static liquid cultures, where cells would form a pellicle biofilm on the air-liquid interface. All studied strains (78 mutants + WT) were mixed together in equal ratios, and the mix was used to inoculate bottles (for planktonic cultures), or microplate wells (for pellicle cultures) with MSgg liquid medium. We introduced further variability to our studied conditions by using two different incubation times: 2 or 5 days, at 30°C, for both culture conditions. After each incubation period we collected spores from the resulting cultures and used them to reinoculate bottles (for spores obtained from planktonic cultures) or microplate wells (for spores obtained from pellicle cultures), that were then incubated under the same conditions. The competition experiment was maintained for a total of nine reinoculation cycles for each culture condition. Full details of the competition experiment can be found in the Material and Methods. We purified total DNA from the cultures obtained after the 1st, 3rd, 5th, 7th, and 9th culture cycle, and from the mix used to initiate the competition. The DNA was used to PCR amplify the *amyE* locus containing the strain-specific DNA

barcodes. Using high-throughput sequencing, we were able to dissect the population dynamics of all the used strains throughout the competition experiment by analyzing their representation ratios within the competing population.

Incubation time alters selection pressure on both planktonic and biofilm populations. Experiments performed under all four studied growth conditions created distinct selection regimes, where different strains were favored during the competition (Fig 1 and S1 Fig). We expected that due to the specific culture reinoculation regime, cells that had formed mature spores within the available time (2 versus 5 days) would be transferred to the next culture iteration, and that those that multiplied for as long as possible prior to sporulation (thus delaying the spore maturation process, but still creating spores before the applied bottleneck), would face a selection advantage due to higher spore numbers. We expected that this would reinforce the competitive selection of each culture condition, and thus amplify the effects of small advantages that the absence or presence of particular Rap-Phr systems would confer in regards to spore formation. Indeed, we observed that after the first culture transfer no strain had increased their total population representation beyond 10% in most experimental replicates (from initial average population representations of 1.26%), in contrast, at the ninth culture transfer the majority of the experimental replicates showed at least one strain that represented more than 30% of the total population. Interestingly, strains that slightly increased their relative abundance in the population after the first culture transfer did not necessarily maintain this trend throughout the competition experiment, in both planktonic and pellicle forming conditions, e.g. Δ*rapA*Δ*rapD* or Δ*rapA*Δ*rapF* mutants (Fig 1). This is probably caused by the evolutionary adaptation to the experimental conditions of other strains in the same population, and genetic drift, which may confer them a selective advantage independent from the Rap-Phr systems. However, we note that our experimental setup offers very limited time for evolutionary adaptations to entirely explain these population dynamics. Alternatively, initial population mix conditions, where all strains are present in similar ratios, may confer slight advantages to specific strains that are overcome by others during the competition. In both planktonic and pellicle forming conditions incubation time was a critical selective parameter: populations that were incubated for 2 days showed greater variation in their final population composition than populations incubated for 5 days (Fig 2). Thus, 5 days incubated populations were under a stronger selection pressure, where only a limited number of *rap-phr* mutant or double mutant strains were able to increase in

relative abundance. Spore formation in *B. subtilis* begins with the detection of starvation conditions, however, this is not an homogeneous process in a population: cells use a bet-hedging strategy to avoid sporulation synchronization [27], furthermore, *B. subtilis* strains that lack specific Rap-Phr systems show temporal differences in sporulation initiation of several hours compared to strains that have those Rap phosphatases [16]. The 2 days incubation period seemed to be insufficient to trigger wide-spread sporulation, and thus only early spores of most strains were transferred into the next culture cycle. In contrast, the 5 days incubation seemed to allow specific strains to sporulate efficiently and thus be overrepresented at the start of each sequential culture cycle.

Fig 1. Heat map representation of the population dynamics of *B. subtilis rap-phr* mutants in competition. Color intensities of the boxes represent the population percentage of strains. Text columns at far-left and far-right indicate which *rap-phr* genes have been deleted (A indicates a Δ*rapA* mutant, AB

indicates a ΔrapAΔrapB mutant, and so on), WT indicates *B. subtilis* DK1042. Text rows on top indicate type of culture (planktonic or pellicle), incubation period (2d= 2 days, 5d= 5 days), transfer number of represented population (t1 to t9), and mix and replicate number. Competition populations were started from four population mixes (A to D), with six replicates per mix. The first two box columns indicate the population representation of tested strains in the competition starter mixes. Both panels show the same data, although using different scales (shown in the top-right corner of each panel): **a)** logarithmic increment of percentage **b)** linear increment of percentage.

Fig 2. Bubble graph representation of mutant frequencies in the evolved populations. Bubble sizes represent the number of populations (from n=24) in which the given *rap-phr* mutant has a frequency higher than the putative input % (i.e. 1.28).

Competition fitness does not depend on growth or biofilm formation exclusively. Our experimental methodology and population dynamics analysis allows examination of fitness impact of each Rap-Phr deletion or combined deletion on sporulation under planktonic and biofilm conditions. Since population size determines the maximum number of possible spores, we also examined how the *rap-phr* mutations affect growth after 16 hours of incubation (Fig 3). We observed drastically different effects on growth depending on the mutated *rap* genes. Interestingly, some *rap* mutations had a consistently negative effect on growth, but this effect was rescued in specific double *rap* mutants, e.g. a Δ*rapI* mutation had a strong negative effect on growth by itself or combined with any other *rap* mutation except Δ*rapG*, since a Δ*rapG*Δ*rapI* double mutant was able to grow almost as efficiently as WT (Fig 3). On the other hand, certain single *rap* mutations that showed a mild effect or no effect on growth, such as Δ*rapA* and Δ*rapK*, resulted in a drastic growth defect when combined in a double mutant strain. Likewise, we examined the effect of the *rap* mutations during pellicle formation after 2 and 5 days of incubation (S2 and S3 Fig). Again, drastic differences could be observed in the impact of single and double *rap* mutations upon the capacity of *B. subtilis* to form pellicles. Nevertheless, there was no correlation between growth and pellicle formation (compare Fig 4, S2 and S3 Fig). Some strains that showed poor 16-hour growth, such as Δ*rapB*Δ*rapI,* were able to form stronger pellicles compared with strains that grew more efficiently, such as Δ*rapB*Δ*rapH* (Fig 4, S2 and S3 Fig).

Furthermore, mutations in diverse Rap phosphatases that target the same regulator can have drastically diverse effects on *B. subtilis* competitiveness in our experimental conditions, both individually and epistatically: RapA and RapB regulate Spo0F, however, Δ*rapA* or Δ*rapA*Δ*rapB* strains became nearly extinct in all tested conditions, while a Δ*rapB* strain increased in frequency, especially when combined with mutations in other Rap proteins that also target Spo0F, such as Δ*rapB*Δ*rapE* and Δ*rapB*Δ*rapH* mutants. Efficient growth of a given mutant strain does not directly correlate with higher fitness in our competition experiments, e.g. a Δ*rapA*Δ*rapG* mutant strain shows better growth compared with a Δ*rapA*Δ*rapF* strain, however, the later shows better competitiveness and increased population representation ratio under all studied conditions already after the first culture transfer. Importantly, the population obtained from the first growth cycle does not depend on viable spores yet, while all the subsequent cultures do. In accordance, all mutant strains that showed a drastic decrease in population representation after the first culture cycle also showed poor growth (Fig 4). Thus, the regulatory and fitness impact of any individual Rap phosphatase cannot be understood solely by knowing its target transcriptional regulator; rather, the whole set of Rap-Phr proteins must be considered in order to explain the regulation of population heterogeneity in *B. subtilis*.

Fig 3. Growth of *B. subtilis* DK1042 and *rap-phr* mutants (as OD_{590} increment over 16 hours). X-axis indicates which *rap-phr* genes have been deleted (A indicates a Δ*rapA* mutant, AB indicates a Δ*rapA*-Δ*rapB* mutant, and so on), WT indicates *B. subtilis* DK1042. Bars represent the average of 4 independent replicates. Error bars represent standard deviation.

Fig 4. Heat map comparing growth of *B. subtilis rap-phr* mutants (16 hours) to their population percentage after the first culture cycle (t1). Yellow boxes represent 16-hour growth on MSgg medium as increase in O.D.$_{590}$. The yellow intensity scale bar (bottom) indicates O.D.$_{590}$ increment. Blue boxes represent the population percentage of all tested strains. The blue intensity scale bar (top) indicates increment of percentage. Each blue box represents the average percentage of 6 population replicates per starter mix. Text rows on top indicate type of culture (planktonic or pellicle), incubation period (2d= 2 days, 5d= 5 days), and starter mix (A or B). Text columns at far-left indicate which *rap-phr* genes have been deleted (A indicates a Δ*rapA* mutant, AB indicates a Δ*rapAB* mutant, and so on), WT indicates *B. subtilis* DK1042.

Selection is defined both by deletion of selected *rap-phr* modules and acquired mutations. The multi-strain competition experiment revealed no clear single winner strains for any of the four different growth conditions at the end of the competition experiment (Fig 1). Furthermore, strains that increased their population representation after the first transfer did not in all cases maintain this trend throughout the experimental competition (Fig 1). Based on these observations, we speculated that the population dynamics were not solely driven by the *rap-phr* deletions. To scrutinize the genomic changes, ten evolved isolates representing all four competition conditions were randomly selected for complete genome resequencing at the end of the selection experiment. Resequencing data showed that eight out of ten isolated evolved strains had gained mutations in one or more genes involved in the regulatory network controlling population heterogeneity (S1 File). Non-synonymous mutations were detected in *degU* (V278G in 5 days planktonic culture) and the biofilm matrix regulator *sinR* (L74S in 2 days pellicle culture) genes, the *comA* gene contained a frame shift mutation in one of the isolates from 2 days pellicle culture. In addition, mutations were identified in genes related to sporulation and germination, including *gerKA* (3 isolates), *gerKC* (2 isolates), *gerAB* (1 isolate), and also in the non-coding region upstream of *yhaX,* a gene that codes for a SigE-dependent sporulation protein. Finally, the pBS32 plasmid, therefore also the *rapP-phrP* genes, was lost in one of the 5 days cultured pellicle isolates that contained *rapB*, *rapD-phrD* deletions. Additional mutations were also detected unrelated to sporulation or with unknown function (S1 File).

Importantly, nine out of the ten isolated evolved strains spontaneously released phage particles that showed lytic activity towards the ancestral strain (S4A,B Fig) – a phenomenon, recently described to be associated with amplification of cryptic phi3Ts and its recombination with indigenous SPβ (into phi3Ts-SPβ hybrids) [28]. Indeed, testing the culture supernatants of these evolved isolates against the indicator strain showed the presence of plaque formation (S4A Fig). In addition, the culture supernatant contained phage particles reminiscent of phi3Ts or its hybrids (S4B Fig), and testing phi3Ts-specific gene PCR verified the presence of the *rapX* gene in these 9 isolates (data not shown). We recently demonstrated that the cryptic phage, phi3Ts, a very close relative of phi3T (KY030782.1; 99.98% sequence identity), increases in copy number in various 3610 or domesticated 168-derivatives of *B. subtilis* when a sporulation bottleneck is used for selection, probably driven by the presence of a Rap-homologue in phi3T that promotes sporulation or production of high-quality spores that germinate faster [28]. However, we hypothesized that in addition to

such a phi3T-driven genetic adaptation that easily spreads around in the whole population, selection might be still driven by certain *rap-phr* deletions.

To test the hypothesis of the population dynamics being affected both by the original *rap-phr* mutations, as well as by the adaptation of some strains to the growth conditions (either by acquiring additional mutations, or by phi3T-driven gene expression), two different competition experiments were performed. To test the impact of the original *rap-phr* mutations on fitness and sporulation, ancestors of the *Δrap* mutant strains that during the competition under 2 days pellicle conditions that persisted after transfer 9 in several of the replicate populations, i.e. that showed higher final population representation than the input level (> 1.28 %, Fig 1), were selected for competition against the WT. Furthermore, the evolved strains isolated at the end of the competition were individually competed against their cognate ancestor in order to test whether the extra mutations arisen during the experiment affected fitness and sporulation. Importantly, we note that these control experiments do not represent the competition experiment of the 79 strains, however, these competitions provided novel knowledge about the individual impact of the *rap-phr* mutations as well as the extra mutations identified in the evolved strains on fitness and sporulation. The ancestral *rap-phr* mutants were competed against the WT, while the evolved *rap-phr* mutants were tested against their corresponding ancestors. The competitions were performed in a 1:1 starting ratio for 48 hours to obtain pellicle biofilms. Afterwards, total cell suspensions as well as heat-treated cell suspensions (to select for spores) were plated for CFU counting, and the relative fitness (of all cells) and sporulation fitness of ancestor versus WT, and of evolved versus ancestor was calculated (see Materials and methods for calculations).

The *ΔrapB*, *ΔrapBΔrapD*, *ΔrapBΔrapE*, *ΔrapBΔrapH*, *ΔrapC*, *ΔrapEΔrapF*, *ΔrapEΔrapH*, *ΔrapHΔrapI* and *ΔrapHΔrapP* ancestors showed significantly increased fitness and/or sporulation fitness compared to the WT (Fig 5A,B). The *ΔrapBΔrapJ* ancestor was not affected in fitness nor sporulation fitness compared to the WT, while the *ΔrapDΔrapJ*, *ΔrapFΔrapJ*, *ΔrapHΔrapJ*, *ΔrapJ* and *ΔrapJΔrapK* ancestors showed reduced pellicle fitness or sporulation fitness compared to the WT. Of the evolved isolates, *ΔrapBΔrapD*, *ΔrapC*, *ΔrapEΔrapF*, *ΔrapHΔrapI* and *ΔrapJΔrapK* demonstrated significantly enhanced spore fitness compared to their ancestor (Fig 5C,D). The evolved *ΔrapBΔrapE* exhibited enhanced pellicle fitness but reduced spore fitness, while the evolved *ΔrapEΔrapH* and *ΔrapHΔrapP* were not affected in either, compared to their ancestor.

Fig 5. The impact of *rap-phr* deletions and additional mutations arisen during the experiment on fitness and sporulation. To test the impact of the *rap-phr* deletions on fitness and sporulation, mutants showing persistence after transfer 9 (>1.28 % abundance) were competed against the WT in MSgg medium to allow pellicle development. Similarly, to test the effect of the extra mutations arisen during the experiment on fitness and sporulation, the isolated evolved strains were competed against their cognate ancestor in MSgg medium. After 2 days, the developed pellicles were harvested, sonicated and exposed to heat treatment in order to kill vegetative cells. Total cell suspensions as well as heat-treated cell suspensions were plated to quantify total cell number and number of spores. A: Relative fitness of the ancestor *rap-phr* mutant strain compared to WT. B: Sporulation fitness of the ancestor *rap-phr* mutant strain compared to WT. C: Relative fitness of evolved *rap-phr* mutant strain compared with its cognate ancestor. D: Sporulation fitness of the evolved strains compared to its cognate ancestor. Data is represented with technical replicates (n=4) as squares and the mean of the replicates are indicated as a black line (n = 3-4). The red horizontal line denotes the WT (A and B) or the cognate *rap-phr* mutant ancestor (C and D). For statistical analysis, the relative (sporulation) fitness was log-transformed and a One Sample T-test was performed to test whether the mean was significantly different from 0. *: P<0.05, **: P<0.005, ***: P<0.001.

The results of these competition experiments reveal that both the original *rap-phr* deletion mutations as well

as the extra mutations arisen during the nine transfers have an impact on the fitness and sporulation of the

strains and thereby also on the observed dynamics of the 79-strain population competition under sporulation selection regime.

Discussion

In this work we have examined how variability in the number and function of Rap-Phr pairs allows *B. subtilis* to adapt to certain selection pressures. Our experimental competition approach, paired with high-throughput sequencing, allowed us to analyze the population dynamics during a multi-strain competition and assess the impact of each Rap-Phr system. Furthermore, the use of single and double *rap-phr* mutants revealed epistatic effects that the presence or absence of specific Rap-Phr systems may have upon population differentiation of *B. subtilis*, including biofilm development and spore formation. Recent investigations indicate that Rap-Phr systems are readily transferred among *B. subtilis* strains, possibly helped by the natural development of competence by this bacterium and by the ability of some Rap-Phr systems to promote their own genetic mobility [15,18]. Our results indicate that variability in Rap-Phr systems among *B. subtilis* strains fine-tunes their ability to compete in diverse environments, and that Rap-Phr systems are particularly important if adaptation depends on differentiation of particular cell types, such as spores. Over evolutionary time, and paired with the genetic plasticity of this bacterium, this could explain the genomic diversity of Rap-Phr systems and the ecological success of this widely spread bacterium [15,18].

Our initial hypothesis was that the strains initially showing increased population representation in the first culture and subsequent sporulation, would be the ones adapting most efficiently to the growth-sporulation cycles applied during our competition, and thereby be the ones winning (having a large increase in their population proportion) at the end of the experiment. Interestingly, no single strain was detected to be a transcendent winner for all of the four different conditions at the end of the competition experiment. Instead, various *rap-phr* mutant strains show an increase in population abundance under specific competition conditions, e.g. multiple population replicates grown under planktonic culture conditions (for 2 and 5 days) show strains Δ*rapD*Δ*rapJ and* Δ*rapH*ΔrapJ to increase their population abundance, but this is not the case for the populations grown under pellicle-forming conditions for 2 and 5 days. On the other hand, mutants in specific Rap-Phr systems do show consistently decreased population abundance across all tested conditions, e.g. most double *rap-phr* mutants with a Δ*rapK* deletion. Our competition approach allows us to

compare the effect of particular *rap-phr* mutations across different conditions, and thus analyze their regulatory function: RapK has been previously suggested to inhibit ComA activity [14,19], here we show that various Δ*rapK* strains suffer a loss of competition fitness when selecting for spores. We hypothesize that the Δ*rapK* mutation leads to an overexpression of ComA-controlled genes related to production of antimicrobials [29], exoenzymes [30], and competence development [31], this in turn may direct metabolic resources away from spore production and maturation, thus resulting in the observed loss of competition fitness. Another interesting behavior was that *rap-phr* mutants the increased their population representation after the first transfer did not necessarily maintain their abundance throughout the multi-strain competition experiment (Fig 1). Under 2 days pellicle conditions, strains Δ*rapA*Δ*rapD*, Δ*rapB*Δ*rapE*, Δ*rapD*, Δ*rapD*Δ*rapK*, Δ*rapF*Δ*rapI*, Δ*rapG*Δ*rapI* and Δ*rapH*Δ*rapI* showed an initial increase in abundance (i.e. during the first transfer). However, of those only the Δ*rapB*Δ*rapE*, Δ*rapF*Δ*rapI*, Δ*rapG*Δ*rapI* and Δ*rapH*Δ*rapI* strains kept this trend throughout the experiment and showed increased population representation in most replicates after transfer 9. On the other hand, Δ*rapB*Δ*rapE*, Δ*rapD*Δ*rapJ*, Δ*rapF*Δ*rapJ*, Δ*rapH*Δ*rapJ*, Δ*rapJ* and Δ*rapJ*Δ*rapK* mutants did not show increased abundance after transfer 1, but were increased after transfer 9. This highlights the versatility that the regulatory effect of Rap-Phr systems confers to *B. subtilis*' ability to generate a phenotypically diverse population. Certain *rap-phr* mutant strains display an initially increased fitness when competing under conditions where all strains are present in the same ratio (starting conditions of the competition), while other strains show an increase in abundance over time. Their particular *rap-phr* mutations allows them to capitalize on hypothetically small fitness advantages when producing mature spores. Furthermore, during the sequential competition transfers, evolutionary adaptation to the experimental conditions of other strains in the population may take place, resulting in those strains gaining an increased sporulation fitness, and possibly taking over the strains initially showing an increase in population abundance. In support of this, resequencing of 10 clones isolated from random populations at the end of the competition experiment revealed that eight of the 10 strains had gained mutations in different genes, including genes involved in the population heterogeneity regulatory network, such as sporulation. This indicates that the strains can quickly acquire additional mutations that help them further adapt to the growth-sporulation cycles applied during the competition experiment.

Competition experiments quantifying fitness and spore fitness were employed to compare *rap-phr* mutants with WT and the evolved *rap-phr* mutants with their respective ancestors. For example, the Δ*rapB*Δ*rapE* ancestor showed enhanced fitness and sporulation fitness compared to the WT after 2 days pellicle formation, while the evolved strain isolated from a 2 days pellicle culture at the end of the competition experiment (transfer 9) showed increased pellicle but reduced spore fitness compared to the ancestor. This evolved Δ*rapB*Δ*rapE* strain had gained mutations in two genes, *gerKC* and *yqcG* encoding a membrane receptor involved in spore germination [32] and a toxin [33], respectively. The Δ*rapB*Δ*rapE* mutant that maintained the increased abundance in the 2 days pellicle conditions throughout the nine transfers is therefore presumably a result of both deletion in Δ*rapB* and Δ*rapE*, as well as the mutations arising during the experiment. Other *rap-phr* mutants that lacked positive fitness or spore fitness compared to the ancestor WT also acquired mutations that may have increased their fitness during the mutli-strain competition experiment, e.g. mutation in *degS* in the Δ*rapH*Δ*rapI* background. The observation that the majority of the isolated evolved strains shows enhanced sporulation fitness compared to the cognate ancestor, indicates that the mutations arising during the experiment and the acquired awakening of the phi3T phage have a dramatic impact on the sporulation fitness, thereby allowing the strains to adapt to the growth-sporulation cycles applied in the competition experiment. This, as well as the observation that most of the *rap* mutations have an effect on fitness and sporulation, indicate that the population dynamics during the experiment are a result of both the original *rap-phr* mutations present in the 78 mutant strains, as well as the additional mutations and phi3T arising during the experiment.

The differences in the variance of surviving *rap-phr* strains between 2- and 5-days culture conditions, and between planktonic and pellicle-forming conditions points to dissimilar selection pressures generated by these growth conditions that further reveal the role that Rap-Phr systems play in environmental adaptability, focusing on the ability of cells to generate mature spores, and to form pellicles. Additionally, genetic drift and negative frequency-dependent selection may occur under the used conditions.

Our results indicate that variability in Rap-Phr systems among *B. subtilis* strains strongly impacts their ability to compete in diverse environments, and that these systems are particularly important if adaptation depends on differentiation of particular cell types such as spores. We propose that the diversity of Rap-Phr regulatory

systems allows *B. subtilis* to fine-tune its genetic regulatory network in order to quickly adapt to new ecological niches, possibly accompanied by the acquisition of additional mutations that provide the bacterium with increased fitness. Further, genetic exchange of elements of a family of regulatory proteins could be a general mechanism for genetically-related bacteria to more efficiently and quickly adapt to new environments.

Materials and methods

Media and cultivation methods. When fresh cultures were needed, strains were pre-grown overnight in Lysogeny broth (LB) medium (LB-Lennox, Carl Roth; 10 g L^{-1} tryptone, 5 g L^{-1} yeast extract, and 5 g L^{-1} NaCl) at 37°C and shaken at 225 r.p.m. LB medium was used for all *B. subtilis* and *Escherichia coli* transformations. MSgg medium (5 mM potassium phosphates buffer (pH 7), 100 mM MOPS, 2 mM MgCl$_2$, 700 µM CaCl$_2$, 100 µM MnCl$_2$, 50 µM FeCl$_3$, 1 µM ZnCl$_2$, 2 µM thiamine, 0.5% glycerol, 0.5% glutamate, 50 µM L-tryptophan, and 50 µM L-phenylalanine, adapted from [34]) was used for the competition experiment and to examine growth kinetics and pellicle formation. GCHE medium (1% glucose, 0.2% glutamate, 100 mM potassium phosphate buffer (pH: 7), 3 mM trisodium citrate, 3 mM MgSO$_4$, 22 mg L^{-1} ferric ammonium citrate, 50 mg L^{-1} L-tryptophan, and 0.1% casein hydrolysate) was used to induce natural competence in *B. subtilis* [35]. Gallegos Rich medium was used to grow *Lactococcus lactis* MG1363*, in order to purify pMH66: 21 g L^{-1} tryptone, 5 g L^{-1} yeast extract, 8.3 g L^{-1} NaCl, 3 g L^{-1} soya peptone, 2.6 g L^{-1} glucose, and 2.5 g L^{-1} MgSO$_4$·7H$_2$O [36]. Media were supplemented with Bacto agar 1.5 % when media were needed for preparing plates. Antibiotics were used at the following final concentrations: kanamycin, 10 µg mL^{-1}; chloramphenicol, 5 µg mL^{-1}; erythromycin-lincomycin (MLS), 0.5 µg mL^{-1} and 12.5 µg mL^{-1} respectively; ampicillin, 100 µg mL^{-1}; spectinomycin, 100 µg mL^{-1}; tetracycline, 10 µg mL^{-1}.

Strain and Plasmid construction. All strains used in this study are listed in S1 Table. To create the single and double *rap-phr* mutant strains, plasmids were first designed that allowed to create clean-deletion mutants of all *rap-phr* gene pairs. All plasmids used in this study are listed in S1 Table, and they were created using standard molecular biology techniques. Briefly, upstream and downstream DNA fragments of ~600 bp flanking the *rap-phr* genes to be mutated were PCR amplified from genomic DNA of *B. subtilis* NCIB 3610. Afterwards, these DNA fragments were sequentially cloned into plasmids pTB120, pTB233, or pTB234, all of which are pBluescript SK(+)-derived plasmids containing an antibiotic resistance cassette (kanamycin, spectinomycin, and MLS, respectively) flanked by *Cre*-recombinase recognition sites *lox66* and *lox71*. Thus, the obtained plasmids contain an antibiotic resistance cassette flanked by the upstream and downstream regions of a given *rap-phr* gene pair. All plasmids were created and maintained in *E. coli* MC1061.

B. subtilis mutants of a single *rap-phr* pair were created via transformation of DK1042 with the corresponding plasmid containing the flanking regions of the target *rap-phr* pair. Double *rap-phr* mutants were created by transforming clean-deletion mutants of single *rap-phr* pairs with genomic DNA of strains that had the desired *rap-phr* mutation still with the corresponding antibiotic resistance cassette. All *B. subtilis* strains generated in this work were obtained via natural competence transformation [35]. Briefly, overnight cultures of the receiver strains grown in LB medium were diluted to a 1:50 ratio with GCHE medium, these cultures were incubated at 37°C for 4 h with shaking at 225 r.p.m. After this incubation period, 5–10 μg of genomic or plasmid DNA were mixed with 500 μL of competent cells and further incubated for 2h before plating on LB plates added with selection antibiotics. *B. subtilis* clean-deletion mutants of single *rap-phr* gene pairs were obtained by using the Cre recombinase expressed from plasmid pMH66 to eliminate their corresponding antibiotic resistance cassette, and subsequently curing pMH66 via thermal elimination of the plasmid [37]. Briefly, strains were transformed with 10 μg of pMH66, selecting transformants via incubation at 37°C on LB plates added with tetracycline. Candidates were then screened for their capacity to grow at 37°C on LB plates added with the antibiotic to which their parental strains (prior to transformation with pMH66) were resistant, those that were not able to grow were further incubated on LB plates at 43°C for 18 h to induce the loss of pMH66. Candidates that were then unable to grow at 37°C on LB plates added with tetracycline were considered to have lost pMH66.

In order to track each strain during the competition experiment, *B. subtilis* DK1042 and all single and double *rap-phr* mutants were marked with a randomly-generated DNA 12 bp barcode in their *amyE* locus. To do this, plasmid pTB666 was created by cloning a chloramphenicol resistance cassette (*cat*) into pTB16 [38], substituting its original kanamycin resistance cassette. The *cat* cassette was amplified from pNW33n using primers oTB118 and oTB119 (see S2 Table). Primer oTB119 has a 12 nt-long random sequence after the binding site of the primer, the DNA barcode. Thus, pTB666 carries a barcoded *cat* which is flanked by the 5´- and 3´-end of the *B. subtilis amyE* gene. 80 different clones of *E. coli* carrying pTB666 were isolated during creation of this plasmid. Each version of pTB666 from these clones was isolated and sequenced with oBC_rev in order to identify it. The various versions of pTB666 were used to transform *B. subtilis* DK1042 and all single and double *rap-phr* mutants using natural transformation as described above.

Successful construction of all used strains and plasmids was validated via PCR, sequencing, and restriction pattern analysis; and by the lack of amylase activity on LB plates added with 1% starch for the case of

barcoded strains [39]. All PCR primers used in this study are listed in S2 Table. Primer pairs were used to amplify the indicated loci in order to confirm the proper mutation of the corresponding gene.

Multi-strain experimental competition. The experimental competition was done using the barcoded versions of *B. subtilis* DK1042 and the single and double *rap-phr* mutants. The experiment was initiated from 4 different starter mixes. Each mix was obtained by mixing overnight cultures of all the competing strains in similar ratios after adjusting their O.D.$_{600}$ to 1.0. Each starting mix was used to inoculate 100 ml bottles (0.5 ml of mix + 9.5 ml of MSgg medium) and 2 ml microplate wells (100 µl of mix + 1900 µl of MSgg medium). The experimental competition used 4 growth conditions: planktonic growth (10 ml culture in bottles shaken at 200 rpm) or pellicle development (static 2 ml in 24-well microplate), and incubation for 2 or 5 days. 24 replicate populations were used for each growth condition (6 replicates from each starting mix). All cultures were incubated at 30°C throughout the experiment.

After each incubation period, spores obtained from each population replicate were used to inoculate the next iteration of the same population. For this, pellicles obtained from the microplate cultures were collected in Eppendorf tubes with 1 ml of MSgg medium and sonicated until the pellicles were completely dispersed; afterwards, 100 µl aliquots from the dispersed pellicles, or 500 µl aliquots from the planktonic culture bottles were incubated at 80°C for 20 minutes in order to kill all vegetative cells. After the incubation period, the heat-treated aliquots were used to inoculate new 100 ml bottles or 2 ml microplate well using the same volumes as during the initiation of the experiment. This regime was followed during 9 culture reinoculation cycles. Additionally, DNA was obtained from aliquots of the starter mixes and from aliquots of each population replicate obtained before the heat treatment during the 1st, 3rd, 5th, 7th, and 9th culture reinoculations. The DNA extraction was performed with the GeneMatrix Bacterial and Yeast Genomic DNA Purification Kit (EURx Ltd., Poland) with the following modifications to the manufacturer's instructions: step 2, added 10 µl of lysozyme (10 mg ml^{-1}); step 3, extended the incubation time to 25 min; step 6, extended the incubation time to 45 min.

48-plex high-throughput barcode sequencing. The *B. subtilis amyE* locus containing the barcodes was PCR amplified from the DNA samples obtained from the competition experiment using primers oBC1 to oBC4 and oBC5 to oBC16. These primers contain distinct 5-bp (for oBC1-4 primers) or 7-bp (for oBC5-16

primers) sequences that allowed us to identify up to 48 individual replicate populations per Illumina sequencing run. Data analysis was carried out using the R statistics environment [40]. PCR products, each represented by one R1-R2 Illumina sequence pair, were looked up for the presence of the 79 barcodes that differentiate between the 79 bacterial strains used in the study (see S3 Table for the barcode sequences of each strain). We linked a PCR product to a given barcode if at least one of its paired-end reads displayed 100% sequence identity over the entire length of the barcode. PCR products that gave ambiguous results (i.e. hits against multiple barcodes) were excluded from the study. Fig 1 and 3 were prepared with Genesis [41].

Growth kinetics measurements. We examined the ability of all barcoded strains to grow on MSgg medium in order to assess the impact of the *rap-phr* mutations. Overnight LB liquid cultures of all barcoded strains were adjusted to O.D.$_{600}$ 0.2. 10 µl of the O.D. adjusted cultures were added to 190 µl of MSgg liquid medium in 200 µl microplate wells. These cultures were incubated at 30°C for 17 hours with shaking. Cell growth was measured as O.D.$_{590}$ change every 15 min using a Tecan Infinite 200Pro microplate reader (Tecan Group Ltd., Switzerland).

Pellicle formation. We examined the ability of all barcoded strains to form pellicles on MSgg medium. Overnight LB liquid cultures of all barcoded strains were adjusted to O.D.$_{600}$ 0.1. 20 µl of the O.D. adjusted cultures were added to 2 ml of MSgg liquid medium in 2 ml microplate wells. These cultures were incubated at 30°C for 5 days. After 2 days of incubation and at the end of the incubation period the obtained pellicles were examined with an Axio Zoom V16 stereomicroscope (Carl Zeiss, Germany) equipped with a Zeiss CL 9000 LED light source, a PlanApo Z 0.5× objective, and AxioCam MRm monochrome camera (Carl Zeiss, Germany).

Resequencing of selected clones. 10 populations from the end of the competition experiment were selected randomly (representing at least 2 replicates of each growth condition). Aliquots of the selected populations were used to inoculate LB plates that were incubated overnight at 30°C in order to isolate clones from each population. Overnight LB liquid cultures of the isolated clones were used to extract genomic DNA using the GeneMatrix Bacterial and Yeast Genomic DNA Purification Kit (EURx Ltd., Poland). Paired-end

libraries were prepared using the NEBNext® Ultra™ II DNA Library Prep Kit for Illumina. Paired-end fragment reads were generated on an Illumina NextSeq sequencer using TG NextSeq® 500/550 High Output Kit v2 (300 cycles). Primary data analysis (base-calling) was carried out with "bcl2fastq" software (v2.17.1.14, Illumina). All further analysis steps were done in CLC Genomics Workbench Tool 9.5.1. Reads were quality-trimmed using an error probability of 0.05 (Q13) as the threshold. In addition, the first ten bases of each read were removed. Reads that displayed ≥80% similarity to the reference over ≥80% of their read lengths were used in mapping. Non-specific reads were randomly placed to one of their possible genomic locations. Quality-based SNP and small In/Del variant calling was carried out requiring ≥8× read coverage with ≥25% variant frequency. Only variants supported by good quality bases (Q ≥ 20) were considered and only when they were supported by evidence from both DNA strands.

Competition experiments. For competition experiments in pellicles, overnight LB liquid cultures of ancestors, evolved strains and WT labelled with mKATE were adjusted to same OD_{600}. None-barcoded ancestors were competed against WT, whereas the evolved, barcoded strains, isolated after transfer 9, were mixed with their cognate (none-barcoded) ancestor strain in a 1:1 ratio. From this mix, 15 µL was added to 1.5 mL MSgg liquid medium in 2 mL microplate wells in a 24-well plate. The cultures were grown under static conditions at 30°C for 48 hours. After incubation, the pellicles were harvested and transferred into Eppendorf tubes containing 1 mL NaCl 0.9 % (Carl Roth) and subjected to standard sonication protocol allowing proper disruption of the biofilm without damaging the cells. Afterwards, 200 µL of the obtained cell suspensions were incubated at 80 °C for 20 minutes, killing all vegetative cells leaving only spores left. CFU assays were performed for cell suspensions (total cells) and heat-treated aliquots (spores only). For the ancestor versus WT competition, obtained cell suspensions were plated on LB plates for CFU counting. For the evolved versus ancestor competition assays, cell suspensions were plated on two different types of plates: LB plates and LB + chloramphenicol (chl; 5 µg ml^{-1}). LB medium allowed growth of both evolved and ancestors, while LB + chl only allows growth of evolved strains due to the presence of a *chl* resistance gene next to the barcode gene in the evolved strains. Thereby, CFU of evolved strains equals the CFU obtained from the LB + chl plates, while CFU of ancestors equals the CFU on the LB plates (evolved + ancestors) subtracted the CFU of LB + chl plates (evolved only).

Relative fitness was calculated for ancestor strains relative to the mKATE-labelled WT, and for evolved strains relative to their cognate ancestors. Relative fitness for strain A, W_A, in competition with strain B was calculated as follows:

$$W_A = \frac{\ln\left(CFU_{A_{48h}}\right)/\ln(CFU_{A\_start})}{\ln\left(CFU_{B_{48h}}\right)/\ln(CFU_{B\_start})}$$

In addition, relative sporulation percentage (SP) and sporulation fitness (SF) of ancestor strains relative to WT and of evolved strains relative to ancestors were calculated. Relative SP and SF for strain A (SP_A and SF_A, respectively) in competition with strain B were calculated as follows:

$$SP_A = \frac{\ln\left(CFU\ spores_{A_{48h}}\right)/\ln(CFU\ cells + spores_{A\_48h})}{\ln\left(CFU\ spores_{B_{48h}}\right)/\ln\left(CFU\ cells + spores_{B\_48h}\right)}$$

$$SF_A = \frac{\ln\left(CFU\ spores_{A_{48h}}\right)/\ln\left(CFU\ cells + spores_{A\_start}\right)}{\ln\left(CFU\ spores_{B_{48h}}\right)/\ln\left(CFU\ cells + spores_{Bstart}\right)}$$

Statistical analysis

Statistical analysis was performed using OriginPro 2018b (OriginLab, US). The relative fitness values calculated from the competition experiments were log-transformed, and statistical differences from $\log^{10}(W)$ = 0 (corresponding to statistical differences of the relative fitness values from W = 1) were calculated using one-sample T-test with test mean = 0 and a significance level of 0.05.

Plaque assay and phage purification. Supernatants, obtained from overnight cultures of the evolved strains, were spotted on soft-agar (0.3%) lawn of the ancestor strain with knockout of biofilm genes (NCBI 3610 Δ*eps*Δ*tasA*). Biofilm mutant was used as a lawn to improve the visibility of plagues. Phage particles were purified like described previously [42]. Specifically, culture supernatants were collected, mixed at a 1:4 ratio with PEG-8000 solution (PEG-8000 20%, 2 M NaCl), incubated on ice for at least 90 min and precipitated by centrifugation (20 min, 7600 rpm). The pellet was resuspended in 10% of the original supernatant volume in TBS solution (50 mM Tris-HCl, 150 mM NaCl, pH 7), incubated on ice for 90 min and centrifuged (20 min, 7600 rpm). Supernatant was carefully transferred to clean Eppendorf tubes and phages were visualized using electron microscopy.

Transmission electron microscopy. Before use, 400 mesh nickel grids with a 3-4 nm thick carbon film, CF400-Ni-UL EMS Diasum, were hydrophilized by 30 sec of electric glow discharging. Next, 5µl of purified phage solutions were applied onto the grids and allowed to adsorb for 1 minute. The grids were rinsed 3 times on droplets of milliQ water and subjected to staining with 2% uranyl acetate. Specifically, with a help of EM grid-grade tweezers, the grids were placed sequentially on droplets of 2% uranyl acetate solution for 10 sec, 2 sec and 20 sec. Excess uranyl acetate was wicked away using filter paper and the grids were allowed to dry overnight and stored in a desiccator until analysis. Transmission electron microscopy was performed utilizing a FEI Tecnai T12 Biotwin TEM operating at 120 kV located at the Center for Electron Nanoscopy at the Technical University of Denmark, and images were acquired using a Bottom mounted CCD, Gatan Orius SC1000WC.

Supporting information

S1 Fig. Heat map representation of the population dynamics of *B. subtilis rap-phr* mutants in competition. Boxes represent the population percentage of strains. Text columns at far-left and far-right indicate which *rap-phr* genes have been deleted (A indicates a Δ*rapA* mutant, AB indicates a Δ*rapA*Δ*rapB* mutant, and so on), WT indicates *B. subtilis* DK1042. Text rows on top indicate type of culture (planktonic or pellicle), incubation period (2d= 2 days, 5d= 5days), transfer number of represented population (t1, t3, t5, t7, and t9), and mix and replicate number. Competition populations were started from 4 population mixes (A to D), with 6 replicates per mix. The first two box columns indicate the population representation of tested strains in the competition starter mixes. Left and right panels show the same data, although using different scales (shown in the top-right corner of each panel): linear increment of percentage (left) and logarithmic increment of percentage (right).

S2 Fig. 2 days pellicles of *B. subtilis* DK1042 and *rap-phr* mutants. Bright-field images are shown after 2 days of incubation on MSgg medium at 30°C. WT indicates *B. subtilis* DK1042, A indicates a Δ*rapA* mutant, AB indicates a Δ*rapA*Δ*rapB* mutant, and so on. The images shown here have an adjusted contrast, so that the pellicles can be easily appreciated. The same image of *B. subtilis* DK1042 is presented twice (top-left and bottom-right) to facilitate pellicle comparison. The scale bars represent 5

mm.

S3 Fig. 5 days pellicles of *B. subtilis* DK1042 and *rap-phr* mutants. Bright-field images are shown after 5 days of incubation on MSgg medium at 30°C. WT indicates *B. subtilis* DK1042, A indicates a Δ*rapA* mutant, AB indicates a Δ*rapA*Δ*rapB* mutant, and so on. The images shown here have an adjusted contrast, so that the pellicles can be easily appreciated. The same image of *B. subtilis* DK1042 is presented twice (top-left and bottom-right) to facilitate pellicle comparison. The scale bars represent 5 mm.

S4 Fig. Presence of infective phage particles in the isolated evolved linages. (A) Supernatants, obtained from overnight cultures of the evolved strains, were spotted on soft-agar (0.3%) lawn of the ancestor strain with knockout of biofilm genes (NCBI 3610 Δ*eps*Δ*tasA*). (B) Electron microscopy images of isolated phage particles.

S1 Table. Strains and plasmids used in this study.

S2 Table. Primers used in this study

S3 Table. DNA barcoded strains used in this study.

S1 File. List of mutations detected in genomes of sequenced population clones. Mutations detected in each sequenced clone are indicated in separate tabs, while the last two tabs enlist mutations found in multiple and in single evolved isolates, respectively.

Acknowledgements

R.G.M. was supported by Consejo Nacional de Ciencia y Tecnología (CONACyT), German Academic Exchange Service (DAAD). Á.T.K. was supported for this project by a Marie Skłodowska Curie Career Integration Grant (PheHetBacBiofilm), DFG Graduate School Jena School of Microbial Communication (JSMC), and a DTU Bioengineering Start-up fund. This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 713683 (H.C. Ørsted COFUND to A.D.).

Competing Interests

The authors declare that there are no competing financial interests in relation to the work described.

Authors contributions

Á.T.K. and R.G.M. conceived the project; R.G.M., M.N.C., T.B., S.K., B.P., P.B., and P.K. performed experiments; B.B., G.M., and A.D. contributed to data analysis; R.G.M., M.N.C., and Á.T.K. wrote the manuscript, with all authors correcting the final version.

Data Availability Statement: Raw sequencing data has been deposited to NCBI Sequence Read Archive (SRA) database. Bioproject accession number: PRJNA626081

References.

1. Lopez D, Vlamakis H, Kolter R. Generation of multiple cell types in *Bacillus subtilis*. FEMS Microbiol Rev. 2009;33: 152–163. doi:10.1111/j.1574-6976.2008.00148.x

2. Vlamakis H, Chai Y, Beauregard P, Losick R, Kolter R. Sticking together: building a biofilm the *Bacillus subtilis way*. Nat Rev Microbiol. 2013;11: 157–168. doi:10.1038/nrmicro2960

3. Mukherjee S, Kearns DB. The structure and regulation of flagella in *Bacillus subtilis*. Annu Rev Genet. 2014;48: 319–40. doi:10.1146/annurev-genet-120213-092406

4. Shank EA, Kolter R. Extracellular signaling and multicellularity in *Bacillus subtilis*. Curr Opin Microbiol. 2011;14: 741–7. doi:10.1016/j.mib.2011.09.016

5. Costerton JW, Lewandowski Z, DeBeer D, Caldwell D, Korber D, James G. Biofilms, the customized microniche. J Bacteriol. 1994;176: 2137–2142. doi:10.1128/jb.176.8.2137-2142.1994

6. Costerton JW, Caldwell DE, Korber DR, Lappin-Scott HM. Microbial biofilms. Annu Rev Microbiol. 1995;49: 711–745. doi:10.1007/978-1-4939-0467-9

7. Davey ME, O'toole GA. Microbial biofilms: from ecology to molecular genetics. Microbiol Mol Biol Rev. 2000;64: 847–67. doi:10.1128/MMBR.64.4.847-867.2000

8. Mielich-Süss B, Lopez D. Molecular mechanisms involved in *Bacillus subtilis* biofilm formation. Environ Microbiol. 2015;17: 555–565. doi:10.1111/1462-2920.12527

9. Mhatre E, Monterrosa RG, Kovács ÁT. From environmental signals to regulators: Modulation of biofilm development in Gram-positive bacteria. J Basic Microbiol. 2014;54: 616–632. doi:10.1002/jobm.201400175

10. Lazazzera BA. The extracellular PHR peptide-Rap phosphatase signaling circuit of *Bacillus subtilis*. Front Biosci. 2003;8: 913. doi:10.2741/913

11. Perego M. Forty years in the making: Understanding the molecular mechanism of peptide regulation in bacterial development. PLoS Biol. 2013;11: 1–5. doi:10.1371/journal.pbio.1001516

12. Reizer J, Reizer A, Perego M, Saier MH. Characterization of a family of bacterial response regulator aspartyl-phosphate (RAP) phosphatases. Microb Comp Genomics. 1997;2: 103–111. doi:10.1089/omi.1.1997.2.103

13. Michna RH, Zhu B, Mäder U, Stülke J. SubtiWiki 2.0--an integrated database for the model

organism *Bacillus subtilis*. Nucleic Acids Res. 2016;44: D654-62. doi:10.1093/nar/gkv1006

14. Sol F del, Marina A, Gallego del Sol F, Marina A. Structural basis of Rap phosphatase inhibition by Phr peptides. PLoS Biol. 2013;11: e1001511. doi:10.1371/journal.pbio.1001511

15. Even-Tov E, Omer Bendori S, Pollak S, Eldar A. Transient duplication-dependent divergence and horizontal transfer underlie the evolutionary dynamics of bacterial cell–cell signaling. PLOS Biol. 2016;14: e2000330. doi:10.1371/journal.pbio.2000330

16. Serra CR, Earl AM, Barbosa TM, Kolter R, Henriques AO. Sporulation during growth in a gut isolate of *Bacillus subtilis*. J Bacteriol. 2014;196: 4184–4196. doi:10.1128/JB.01993-14

17. Budde I. Adaptation of *Bacillus subtilis* to growth at low temperature: a combined transcriptomic and proteomic appraisal. Microbiology. 2006;152: 831–853. doi:10.1099/mic.0.28530-0

18. Even-Tov E, Omer Bendori S, Valastyan J, Ke X, Pollak S, Bareia T, et al. Social evolution selects for redundancy in bacterial quorum sensing. PLOS Biol. 2016;14: e1002386. doi:10.1371/journal.pbio.1002386

19. Auchtung JM, Lee CA, Monson RE, Lehman AP, Grossman AD. Regulation of a *Bacillus subtilis* mobile genetic element by intercellular signaling and the global DNA damage response. Proc Natl Acad Sci U S A. 2005;102: 12554–9. doi:10.1073/pnas.0505835102

20. Perego M, Hoch JA. Cell-cell communication regulates the effects of protein aspartate phosphatases on the phosphorelay controlling development in *Bacillus subtilis*. Proc Natl Acad Sci U S A. 1996;93: 1549–53. doi:10.1073/pnas.93.4.1549

21. Core L, Perego M. TPR-mediated interaction of RapC with ComA inhibits response regulator-DNA binding for competence development in *Bacillus subtilis*. Mol Microbiol. 2003;49: 1509–1522. doi:10.1046/j.1365-2958.2003.03659.x

22. Bongiorni C, Ishikawa S, Stephenson S, Ogasawara N, Perego M. Synergistic regulation of competence development in *Bacillus subtilis* by two Rap-Phr Systems. J Bacteriol. 2005;187: 4353–4361. doi:10.1128/JB.187.13.4353-4361.2005

23. Omer Bendori S, Pollak S, Hizi D, Eldar A. The RapP-PhrP quorum-sensing system of *Bacillus subtilis* strain NCIB3610 affects biofilm formation through multiple targets, due to an atypical signal-insensitive allele of RapP. J Bacteriol. 2015;197: 592–602. doi:10.1128/JB.02382-14

24. Parashar V, Konkol M a, Kearns DB, Neiditch MB. A plasmid-encoded phosphatase regulates

Bacillus subtilis biofilm architecture, sporulation, and genetic competence. J Bacteriol. 2013;195: 2437–48. doi:10.1128/JB.02030-12

25. McLoon AL, Guttenplan SB, Kearns DB, Kolter R, Losick R. Tracing the domestication of a biofilm-forming bacterium. J Bacteriol. 2011;193: 2027–2034. doi:10.1128/JB.01542-10

26. Maughan H, Nicholson WL. Increased fitness and alteration of metabolic pathways during *Bacillus subtilis* evolution in the laboratory. Appl Environ Microbiol. 2011;77: 4105–4118. doi:10.1128/AEM.00374-11

27. Tan IS, Ramamurthi KS. Spore formation in *Bacillus subtilis*. Environ Microbiol Rep. 2014;6: 212–225. doi:10.1111/1758-2229.12130

28. Dragos A, B P, Hasan Z, Lenz-Strube M, Kempen PJ, Maroti G, et al. Phage recombination drives evolution of spore-forming *Bacilli*. bioRxiv [Preprint]. 2020. Available from: https://www.biorxiv.org/content/10.1101/2020.05.06.055103v1. doi:10.1101/2020.05.06.055103

29. Comella N, Grossman AD. Conservation of genes and processes controlled by the quorum response in bacteria: Characterization of genes controlled by the quorum-sensing transcription factor ComA in *Bacillus subtilis*. Mol Microbiol. 2005;57: 1159–1174. doi:10.1111/j.1365-2958.2005.04749.x

30. Msadek T, Kunst F, Klier A, Rapoport G. DegS-DegU and ComP-ComA modulator-effector pairs control expression of the *Bacillus subtilis* pleiotropic regulatory gene *degQ*. J Bacteriol. 1991;173: 2366–2377. doi:10.1128/jb.173.7.2366-2377.1991

31. Nakano MM, Magnuson R, Myers A, Curry J, Grossman AD, Zuber P. *srfA* is an operon required for surfactin production, competence development, and efficient sporulation in *Bacillus subtilis*. J Bacteriol. 1991;173: 1770–1778. doi:10.1128/JB.173.5.1770-1778.1991

32. Zheng L, Abhyankar W, Ouwerling N, Dekker HL, van Veen H, van der Wel NN, et al. *Bacillus subtilis* spore inner membrane proteome. J Proteome Res. 2016;15: 585–594. doi:10.1021/acs.jproteome.5b00976

33. Bloom-Ackermann Z, Steinberg N, Rosenberg G, Oppenheimer-Shaanan Y, Pollack D, Ely S, et al. Toxin-Antitoxin systems eliminate defective cells and preserve symmetry in *Bacillus subtilis* biofilms. Environ Microbiol. 2016;18: 5032–5047. doi:10.1111/1462-2920.13471

34. Branda SS, González-Pastor JE, Ben-Yehuda S, Losick R, Kolter R. Fruiting body formation by

Bacillus subtilis. Proc Natl Acad Sci U S A. 2001;98: 11621–6. doi:10.1073/pnas.191384198

35. Kunst F, Rapoport G. Salt stress is an environmental signal affecting degradative enzyme synthesis in *Bacillus subtilis*. J Bacteriol. 1995;177: 2403–2407.

36. Gallegos-Monterrosa R, Kankel S, Götze S, Barnett R, Stallforth P, Kovács ÁT. *Lysinibacillus fusiformis* M5 induces increased complexity in *Bacillus subtilis* 168 colony biofilms via hypoxanthine. J Bacteriol. 2017;199: JB.00204-17. doi:10.1128/JB.00204-17

37. Yan X, Yu H-JJ, Hong Q, Li S-PP. Cre/lox system and PCR-based genome engineering in *Bacillus subtilis*. Appl Environ Microbiol. 2008;74: 5556–5562. doi:10.1128/AEM.01156-08

38. Grau RR, de Oña P, Kunert M, Leñini C, Gallegos-Monterrosa R, Mhatre E, et al. A duo of potassium-responsive histidine kinases govern the multicellular destiny of *Bacillus subtilis*.MBio. 2015;6: e00581-15. doi:10.1128/mBio.00581-15

39. Harwood CR, Cutting SM. Molecular biological methods for *Bacillus*. Chichester, England: John Wiley & Sons Ltd.; 1990.

40. R Development Core Team. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; 2013. Available from: https://www.r-project.org

41. Sturn A, Quackenbush J, Trajanoski Z. Genesis: cluster analysis of microarray data. Bioinformatics. 2002;18: 207–8. doi:10.1093/bioinformatics/18.1.207

42. Martin M, Dragoš A, Hölscher T, Maróti G, Bálint B, Westermann M, et al. De novo evolved interference competition promotes the spread of biofilm defectors. Nat Commun. 2017;8: 15127. doi:10.1038/ncomms15127