Rapid adaptation to malaria facilitated by admixture in the human population of Cabo Verde

Iman Hamid[1], Katharine Korunes[1], Sandra Beleza[2], Amy Goldberg[1*]

[1]Department of Evolutionary Anthropology, Duke University, 130 Science Drive, Durham, NC, 27708
[2]Department of Genetics and Genome Biology, University of Leicester, Leicester, United Kingdom

*corresponding author, amy.goldberg@duke.edu

Abstract:

Humans have undergone large migrations over the past hundreds to thousands of years, exposing ourselves to new environments and selective pressures. Yet, evidence of ongoing or recent selection in humans is difficult to detect. Many of these migrations also resulted in gene flow between previously separated populations. These recently admixed populations provide unique opportunities to study rapid evolution in humans. Developing methods based on distributions of local ancestry, we demonstrate that this sort of genetic exchange has facilitated detectable adaptation to a malaria parasite in the admixed population of Cabo Verde within the last ~22 generations. We estimate the selection coefficient is approximately 0.08, one of the highest inferred in humans. Notably, we show that this strong selection at a single locus has likely affected patterns of ancestry genome-wide, potentially biasing demographic inference. Our study provides evidence of adaptation in a human population on historical timescales.

Introduction:

Genetic studies have demonstrated the important role of adaptation throughout human evolution, including classic examples such as loci underlying pigmentation, and adaptation to high-altitude lifestyles and infectious disease (Sabeti et al., 2002; Ohashi et al., 2004; Lamason et al., 2005; Voight et al., 2006; Nielsen et al., 2007; Norton et al., 2007; Pickrell et al., 2009; Yi et al., 2010; Fumagalli et al., 2011; Grossman et al., 2013; Lachance and Tishkoff, 2013). Yet, we have a limited understanding of adaptation in human populations on historical timescales; that is, during the last tens of generations. The ongoing selective pressures shaping human genomic variation, and how quickly humans can adapt to strong selective pressures, remain unclear. Adaptation on these short timescales is of particular importance because large-scale migrations within the past thousands to hundreds of years have exposed human populations to new environments and diseases, acting as new selective pressures (Hellenthal et al., 2014; Busby et al., 2016; Mathias et al., 2016; Nielsen et al., 2017; Laso-Jadart et al., 2017; Patin et al., 2017; Fernandes et al., 2019).

Admixture—gene flow between previously diverged populations to form a new population with ancestry from both source populations—provides a particularly rapid mechanism for selection to act in a population by introducing alleles previously adapted in a source population into the admixed population (Huerta-Sánchez et al., 2014; Jeong et al., 2014; Racimo et al., 2015; Norris et al., 2020). Additionally, in recent human admixture, ancestry contributions from each source population are often large enough to introduce alleles at intermediate frequencies, potentially avoiding loss from drift (Ruiz-Linares et al., 2014; Mathias et al., 2016; Fernandes et al., 2019; Fortes-Lima et al., 2019). More generally, admixture is ubiquitous in human history (Moorjani et al., 2011; Hellenthal et al., 2014; Busby et al., 2015, 2016; Triska et al., 2015; Laso-Jadart et al., 2017; Patin et al., 2017); therefore, understanding the effects of selection in these often understudied populations is essential to the study of human evolution.

The admixture process may obscure the signals commonly used to detect selection by increasing linkage disequilibrium and changing the distribution of allele frequencies (Lohmueller et al., 2011, 2010; Gravel, 2012; Racimo et al., 2015). Further, common signatures of selection, such as deviation from neutral expectations of the allele frequency spectrum, may not be sensitive to adaptation on the scale of tens of generations (Sabeti et al., 2006; Field et al., 2016). Recent progress has given insight into human adaptation during the past few thousand years by using allele frequency trajectories from ancient DNA (Lindo et al., 2016) or the distribution of singletons in extremely large datasets (Field et al., 2016). We focus on admixed populations as an opportunity to detect rapid adaptation using modern populations and moderate sample sizes, allowing broader sets of populations to be studied. That is, within-genome ancestry patterns across multiple nearby loci may be easier to detect than single allele frequency shifts (Tang et al. 2007). Further, ancestry-based methods constrain the timing of potential selection to post-admixture, providing concrete information about the timing of selection, whereas non-ancestry-based summary statistics may detect selection in the source populations.

We test this concept developing new ancestry-based methods to characterize adaptation to malaria during the ~20 generations since the founding of the admixed human population of Cabo Verde. The Republic of Cabo Verde is an archipelago off the coast of Senegal and was uninhabited before settlement in ~1460 by Portuguese colonizers and enslaved peoples from the

Senegambian region of West Africa (Fernandes et al., 2003; Beleza et al., 2012; Verdu et al., 2017), henceforth referred to as "European" and "West African" source populations, respectively. We focus on three major island regions of Cabo Verde: Santiago, Fogo, and the Northwest Cluster (**Fig. 1A**).

The malaria parasites *Plasmodium vivax*, *P. falciparum*, and *P. malariae* have been reported across the islands of Cabo Verde since settlement; recurrent malaria epidemics have primarily occurred in highly populated regions (World Health Organization et al., 2012; DePina et al., 2019). Santiago, which has consistently been the most densely populated of the Cabo Verde islands, has experienced the most substantial burden of malaria transmission since settlement ~20 generations ago. Personal and historical accounts of malaria incidence within Cabo Verde described the largest and most populous island, Santiago, as the most "sickly" and "malarious" (Patterson, 1988). In the last century, malaria epidemics of both *P. vivax* and *P. falciparum* have occurred primarily in Santiago (Snow et al., 2012; World Health Organization et al., 2012; Ferreira et al., 2017; DePina et al., 2018). Whereas the other two island regions we consider share ancestry components with Santiago, but largely lacked the selective pressure of recurrent malaria transmission, providing a unique opportunity to compare related populations with and without malaria as a selective pressure.

We hypothesized that admixture has facilitated rapid adaptation to the malaria parasite *Plasmodium vivax* via the malaria-protective *DARC* locus (*Duffy antigen receptor for chemokines*; also known as *Atypical Chemokine Receptor 1* (*ACKR1*)) in Santiago. The protective allele is almost fixed in West African populations and rare elsewhere (Howes et al., 2011; Gething et al., 2012). The malaria parasite *P. vivax* uses the chemokine receptor encoded by the *DARC* gene to enter and infect red blood cells. The Duffy-null allele (also known as FY*O, rs2814778) is protective against *P. vivax* infection via a SNP that disrupts binding of an erythroid-specific transcription factor in the promoter region (Mercereau-Puijalon and Ménard, 2010; Gething et al., 2012). Thus, individuals carrying the null allele have reduced expression of Duffy antigens on the surface of the blood cell, protecting against *P. vivax* infection. Duffy-null is a classic example of strong selection in the human lineage, and it has been estimated to be under one of the strongest selective pressures in human history (Hamblin and Di Rienzo, 2000; Hamblin et al., 2002; Kwiatkowski, 2005; McManus et al., 2017), suggesting it is a plausible selective pressure in Cabo Verde.

Interestingly, this hypothesis goes back to a voyage in 1721 in which Captain George Roberts reported that a disease in Santiago is "dangerous to strangers" during the rainy season (Roberts, 1745). Consistent with ancestry-mediated protection from malaria, the record has been interpreted by medical historians to suggest that "foreign visitors and residents of European descent seem to have suffered more than the African and Afro-Portuguese majority" from malaria in Santiago (Patterson, 1988).

In this study, we combine ancestry-based summary statistics and simulations to identify and characterize selection at the malaria-protective *DARC* locus on the island of Santiago during the ~20 generations since the onset of admixture. Importantly, we also consider the genome-wide consequences of this mode of selection. That is, we find that strong selection at a single locus may shift genome-wide ancestry patterns, with potential to bias demographic inference. The

results of this study provide evidence for rapid adaptation in human populations and advance our ability to detect and characterize selection in recently admixed populations.

Figure 1. Enrichment of West African ancestry at *DARC* locus in Santiago, Cabo Verde. **A)** Map of Cabo Verde islands, and sample sizes for number of individuals from each island region. **B)** The distribution of West African-related local ancestry proportion across the genome by SNP (n=881,279) by island, with the *DARC* locus marked by vertical red lines. The *DARC* locus is an outlier for high West African-related ancestry in Santiago, but not Fogo or the Northwest cluster.

Results:

Enrichment of West African Ancestry at the *DARC* locus on Santiago

Empirical studies of selection in admixed populations often look for regions of the genome that deviate from genome-wide patterns of genetic ancestry (Tang et al., 2007; Jin et al., 2012; Jeong et al., 2014; Rishishwar et al., 2015; Triska et al., 2015; Zhou et al., 2016; Laso-Jadart et al., 2017; Patin et al., 2017; Fernandes et al., 2019; Lopez et al., 2019; Norris et al., 2019; Vicuña et al., 2020). Regions of the genome with substantially higher ancestry from one source than present on average in the rest of the genome are hypothesized to be enriched for genes under selection. For an allele at different frequencies in the source populations, selection will increase the frequency of the ancestry on which the adaptive allele occurs at that locus. We tested if the *DARC* locus was an outlier within the genome for West African ancestry. **Figure 1B** shows the distribution of West African ancestry for over 880,000 SNPs by island, with the value for the Duffy-null SNP position marked in red. Within Santiago, this locus has the highest frequency of West African ancestry in the population, occurring at 0.834 frequency compared to mean West African ancestry across SNPs for individuals from Santiago of 0.730. In contrast, the *DARC* locus is not an outlier in its frequency of West African ancestry on the other island regions, occurring at the 75[th] and 65[th] percentiles for Fogo and the NW Cluster, respectively. High West African local ancestry proportion at the *DARC* locus in Santiago is consistent with the expectation that the Duffy-null allele rapidly increased in frequency following admixture, simultaneously increasing the proportion of individuals with West African ancestry at that locus relative to the genome-wide average.

Long, high-frequency West African ancestry tracts span the *DARC* locus on Santiago

The distribution of the lengths of ancestry tracts spanning a selected locus can provide information for detecting and characterizing selection beyond single-locus outlier tests. In the case of recent admixture and strong selection, we might generally expect to see a parallel increase in local ancestry proportion in the regions surrounding the beneficial locus because the beneficial allele increases before recombination can break up large surrounding ancestry blocks. This is analogous to the increase in linkage disequilibrium and homozygosity in non-admixed populations (Sabeti et al., 2002; Kim and Nielsen, 2004; Voight et al., 2006). **Figure 2A** plots ancestry tracts that span the *DARC* locus for individuals from Santiago. As expected from source population allele frequencies, the Duffy-null allele is contained on all West African ancestry tracts spanning the locus and not found on any European ancestry tracts. Consistent with recent selection, West African ancestry tracts are longer and in higher frequency than European ancestry tracts covering the region. The median West African ancestry tract length spanning the locus is ~85 Mb, while the median European tract length is ~39 Mb.

In order to test if the observed local ancestry patterns are suggestive of selection beyond genome-wide ancestry proportion, we developed a summary statistic based on the length and frequency of the tract-length surrounding a locus. The integrated Decay in Ancestry Tract (*iDAT*) score compares the rate of decay of ancestry tract lengths as a function of distance from a site of interest (**Fig. 2B**; **Methods**). The statistic is analogous to the commonly used integrated haplotype score (*iHS*) (Voight et al., 2006), but considers local ancestry tracts instead of haplotypes. Negative values for *iDAT* indicate longer West African ancestry tracts at higher frequencies compared to European ancestry tracts. Positive values indicate longer European ancestry tracts at higher frequencies compared to West African ancestry tracts. Windows that contain multiple extreme values of *iDAT* provide stronger evidence for recent selection. **Figure 2C** plots *iDAT* values along the genome for individuals from Santiago. Values are calculated by averaging over 20 Mb sliding windows (step size of 1 Mb) for 10000 random standardized *iDAT* scores. Notably, in Santiago, the *DARC* gene is contained in the window with the lowest *iDAT* score in the genome (**Fig. 2C**; window *iDAT* score = -2.602). *iDAT* scores near the *DARC* locus are not outliers in other islands regions (**Fig. S1**).

Ancestry-based signatures for Santiago cannot be explained by drift alone

Under a simple population genetic model, we expect the frequency of a neutral allele in an admixed population to be a linear combination of the allele frequencies in each source population and their relative ancestry contributions. The Duffy-null allele is nearly fixed in the West African source population and largely absent in the European source population; therefore, under neutrality, the expected frequency of the allele in each Cabo Verdean population is approximately equal to the West African ancestry contribution. Using the observed global ancestry proportion inferred with ADMIXTURE (Alexander and Lange, 2011) as an estimate of the ancestry contribution from West Africa to each island, we found that the Duffy-null allele is at a higher frequency than expected under neutrality for the island of Santiago, but not the other regions of Cabo Verde (**Table S1**, binomial test, Santiago: $p = 2.193 \times 10^{-5}$; Fogo: $p = 0.1915$; NW Cluster: $p = 0.8172$).

Figure 2. Long, high-frequency West African ancestry tracts span the *DARC* locus in Santiago. **A)** The distribution of West African (purple) and European (green) ancestry tract lengths spanning the *DARC* locus (dashed line). Each horizontal line represents a single chromosome in the population (n=343, one chromosome was excluded due to having unknown ancestry at the *DARC* locus). **B)** Decay in Ancestry Tract (*DAT*) as function of absolute distance from the Duffy-null allele for West African (purple) and European (green) ancestry tracts. **C)** Mean standardized integrated *DAT* (*iDAT*) score for 20 Mb sliding windows (step size = 1Mb), using standardized *iDAT* for 10000 random positions across the genome. Horizontal solid grey line indicates mean windowed standardized *iDAT* score (-0.196), and horizontal dashed grey lines indicates three standard deviations from the mean windowed score. The red dot is the most extreme windowed standardized *iDAT* score (-2.602), indicative of a larger area under the curve for West African *DAT* compared to European *DAT*. This 20 Mb window contains the Duffy-null SNP.

In order to estimate the expected distribution of ancestry within the population and test if the values of various summary statistics for the *DARC* locus on Santiago can be explained by drift alone, we conducted neutral simulations in SLiM (Haller and Messer, 2019) **(Methods)**. We calculated the following five summary statistics for each simulated population: the West African local ancestry proportion at *DARC*, the variance in West African local ancestry proportion across chromosome 1, the median and mean West African ancestry tract length containing the Duffy-null allele, and the unstandardized *iDAT* score for the Duffy-null SNP. Studies of demographic history and selection in recently admixed populations often assume constant population size and a single admixture event to simplify simulations. In order to confirm that assumptions about demographic history do not change our expectations, we considered multiple scenarios of population growth, differences in population size, and models of both constant contributions and single admixture events (**Table S2**). The values for the summary statistics for Santiago generally lie outside our expectations for all models (**Fig. S2**).

Together, these summary statistics provide suggestive evidence that the *DARC* locus has been under positive selection on the island of Santiago since admixture started ~20 generations ago. To formally test this hypothesis, we extended the SWIF(r) framework developed by (Sugden et al., 2018). SWIF(r) is a machine learning classification framework that explicitly learns the joint distributions for a set of features and returns a posterior probability of a selective sweep at a site of interest. It is particularly useful for handling summary statistics that are correlated, such as the length and frequency of ancestry tracts. We trained SWIF(r) using data simulated in SLiM and estimated the posterior probability of a selective sweep at the *DARC* locus using the five ancestry-based measures (**Methods**). SWIF(r) returned a high posterior probability of a sweep at *DARC* on Santiago starting 20 generations ago ($P > 0.999$).

Strong selection inferred at *DARC* locus in Santiago

Beyond identifying selection, inference of the strength of selection is informative about the evolutionary processes shaping human genomes. We used two complementary approaches to infer the strength of selection at the *DARC* locus. First, we considered a deterministic classical population-genetic model of selection based on the trajectory of allele frequencies over time on a grid of possible dominance and selection coefficients (**Methods**). The estimate of the selection coefficient depends on dominance; past studies have modeled Duffy-null as recessive (Hodgson et al., 2014), dominant (Pierron et al., 2018), and additive (McManus et al., 2017) when estimating selection strength in other human populations. **Figure 3A** plots the selection strength (s) as a function of the dominance coefficient (h) of the Duffy-null allele for a set of three realistic initial frequencies, assuming 20 generations of constant selection strength. Functional studies suggest that heterozygotes have at least partial protection against *P. vivax* infection (Cavasini et al., 2007; Gething et al., 2012; Kano et al., 2018; Sousa et al., 2007); the Duffy-null allele is unlikely to be fully recessive or fully dominant. Taking the mean of selection coefficients for $0.2 \leq h \leq 0.8$, we estimate the selection coefficient for each initial frequency, $s_{0.65} = 0.106$, $s_{0.70} = 0.082$, and $s_{0.75} = 0.056$, where s_{p} is the inferred selection coefficient for initial allele frequency p_{o}.

Second, we used a simulation and rejection framework, approximate Bayesian computation (ABC), to jointly infer the selection coefficient and initial West African contribution while allowing for drift (**Methods**). We used the five ancestry-based summary statistics described previously. We assumed an additive model, a single admixture event, and exponential growth in the population. Taking the median of the posterior distribution as the point estimate for selection coefficient, we estimated $s = 0.0795$ (**Fig. 3B,** see **Fig. S3** for estimates of s when modeling Duffy-null as either a dominant or a recessive mutation). This estimate of selection coefficient is consistent with those estimated under the deterministic population-genetic model.

Figure 3. Strong selection inferred at *DARC* locus in Santiago. **A)** Pairs of *s* and *h* that result in a small difference in final allele frequency calculated under the model and the allele frequency observed in the Santiago genetic data, $\left|p_{20}\text{-}p_{\text{Duffy}}\right|$ <0.01 under a deterministic population genetic model. Colors indicate the initial Duffy-null frequency: p_o=0.65, black; p_o=0.70, dark grey; p_o=0.75, light grey. **B)** ABC estimates of the selection coefficient for Duffy-null on Santiago. Shaded grey area shows prior distribution of selection coefficient [$s \sim U(0, 0.2)$]. Dark grey histogram shows posterior distribution for selection coefficient (median = 0.0795), constructed from regression-adjusted values from accepted simulations.

Selection at a single locus impacts genome-wide ancestry estimates

Mean global ancestry proportion is often used as an estimate for initial ancestry contributions for admixed populations (Moreno-Estrada et al., 2013; Hellenthal et al., 2014; Bryc et al., 2015; Mathias et al., 2016; Laso-Jadart et al., 2017; Patin et al., 2017; Fernandes et al., 2019). However, our ABC estimates of the initial contributions from West Africa are lower than the mean ancestry currently observed in Santiago. The median of the posterior of initial contributions from West Africa is 0.690, with the middle fifty percentile of observed values in [0.682,0.697] (**Fig. 4A**). In contrast, the observed mean ancestry in Santiago (**Fig. 1B**) is 0.737. While this particular difference in observed and inferred founding contributions may be due to sampling biases or other neutral processes, it raises the more general question of how strong selection at a single locus may impact genome-wide ancestry patterns. We hypothesized that selection at *DARC* may have increased the genome-wide West African ancestry proportions in the current population of Santiago.

To test the genomic consequences of post-admixture selection at a single locus, we simulated whole human autosomes under a model of exponential growth and a single admixture event with selection at a single locus. We first considered a model based on the history of Santiago, using the posterior distributions of selection coefficient and initial West African ancestry contribution as parameters for the simulations (**Methods**). **Figure 4A** plots the estimated posterior distribution of initial West African contribution in dark grey and the simulated distribution of global ancestry after 20 generations in white. The distribution of global ancestry in the populations simulated with selection (median 0.723, white) is noticeably higher than the initial contributions specified in the simulations (median 0.690, dark grey). This demonstrates that

selection at a single locus is a plausible mechanism to increase mean global ancestry in an admixed population under a scenario similar to Santiago.

To explore the mechanism and relationship between selection strength at a single locus and genome-wide ancestry patterns, we simulated whole autosomes with initial West African ancestry contribution at 0.65 with selection coefficient varying from 0 to 0.2 at a single locus on chromosome 1. **Figure 4B** plots the mean global ancestry in each simulated population after 20 generations as a function of the selection coefficient at a single locus. The global ancestry proportion increases with selection strength (black), beyond the contribution of higher ancestry on the selected chromosome alone (grey). Together, this evidence suggests that strong selection at the *DARC* locus over 20 generations may have skewed global ancestry in Santiago and raises potential biases with a statistic that is often used to infer neutral demographic histories.

Figure 4. Selection at a single locus impacts genome-wide ancestry proportion. **A)** Inferred (dark grey), simulated (white), and observed (red) mean of global ancestry in Santiago over time. The dark grey histogram plots the posterior distribution for initial ($g = 1$) West African ancestry contribution inferred using ABC (median, 0.690); the prior distribution [$m\sim U(0.1, 0.9)$] is in light grey. The red line plots the mean global ancestry estimated by ADMIXTURE from modern genetic data from Santiago, 0.737. The observed global ancestry is higher than most values of the initial contributions inferred in dark grey. The white histogram plots the distribution of West African global ancestry proportion calculated after 20 generations in populations simulated with selection coefficients and initial ancestries drawn from the ABC-inferred values (median, 0.723). The global ancestry calculated after 20 generations of simulated selection (white) more closely matches that observed from Santiago genetic data (red line). **B)** West African mean global ancestry proportion calculated for 500 simulated populations after 20 generations under varying single-locus selection coefficients, *s*. We set the initial West African ancestry contribution to 0.65. Black circles indicate whole autosome simulations. Grey circles indicate global ancestry calculated if selection only affects ancestry on chromosome 1, fixing the rest of the genome at 0.65 West African ancestry proportion. The higher values for the black circles than grey circles demonstrate that selection impacts global ancestry beyond the local effects of the chromosome under selection.

Discussion:

Using adaptation to malaria in the admixed population of Cabo Verde as a case study, we have demonstrated that admixture can facilitate adaptation in merely tens of generations in humans. Developing methods to identify and characterize post-admixture selection, we found that this

rapid adaptation leaves detectable genomic signatures (**Fig. 2**), with potential genome-wide consequences (**Fig. 4**). Combining inference under two complementary methods, and under a range of possible dominance coefficients and initial allele frequencies, we estimated selection strength of $s \approx 0.08$ for the Duffy-null allele in Santiago (**Fig. 3**). This is among the highest inferred for a locus in any human population.

Introgression of an adaptive allele can facilitate adaptation on short timescales, particularly for traits with large effects from single loci. When ancestry contributions from multiple sources are high, such as is common in recent human admixture, selection post-admixture can be a faster mode of adaptation, similar to selection on standing variation (Hedrick, 2013; Hermisson and Pennings, 2005).

Commonly-used ancestry outlier approaches have identified candidate regions for admixture-enabled adaptation in many populations; for example, at the *HLA* and *LCT* regions in Bantu speaking populations (Patin et al., 2017), the *MHC* locus in Mexicans (Zhou et al., 2016), and adaptation to high-altitude at *EGLN1* and *EPAS1* in Tibetans (Jeong et al., 2014). However, results from this framework alone can be difficult to interpret because drift post-admixture may substantially change allele frequencies and the distribution of local ancestry within and between individuals (Bhatia et al., 2014; Belbin et al., 2018; Calfee et al., 2020). Further, recessive deleterious variation masked by heterosis can similarly cause a signal of increased introgressed ancestry, especially in regions of low recombination (Kim et al., 2018). As a result, outlier approaches may have increased rates of false-positive detection of regions under selection. Outlier approaches also discard local haplotype information and are not informative about the selection strength or timing. Instead, we developed a suite of methods to identify and characterize selection post-admixture: the *iDAT* summary statistic, application of the SWIF(r) framework to estimate the probability a locus is under selection in an admixed population, and an ABC framework to jointly infer selection strength and initial admixture contributions (**Methods**).

Methods to detect adaptation driven by alleles introduced through gene flow in human populations have typically focused on ancient admixture between highly diverged populations, often with small contributions from one of the sources (Jagoda et al., 2018; Racimo et al., 2017; Setter et al., 2020). Recent advances leverage patterns of ancestry to consider recent admixture, though perform best for events at least hundreds of generations ago (Svedberg et al., 2020). Instead, we emphasize admixed populations as a model for adaptation on historical timescales, with selection dramatically changing genomic variation within tens of generations. This timescale is important for elucidating human history and has implications for conservation genetics and ecology in other organisms. Additionally, our summary statistic approach can be flexibly applied in a variety of inference methods. For example, our implementation in a likelihood-free ABC framework allows for flexible population history models fit to the population of interest. This approach moves beyond identification of loci under selection, allowing joint inference of selection and population history parameters.

We apply these methods to characterize post-admixture adaptation in the Cabo Verdean island of Santiago. The Cabo Verdean populations have a number of advantages for identifying and interpreting selection over the last ~500 years. First, the island geography minimizes within-population structure and provides comparison island populations with shared ancestry

components to partially account for demography. Second, historical records give a clear boundary for the earliest onset of selection in the admixed population, based on the initial occupation and admixture in the 1460s (approximately 22 generations ago assuming a 25-year human generation time).

We inferred that the Duffy-null allele rapidly increased in frequency after admixture as a result of its adaptive resistance to *P. vivax* infection. While we consider this to be the strongest candidate locus, given the large ancestry tracts, it is possible that selection at other nearby loci is responsible for the observed ancestry patterns. The Duffy-null allele shows extreme geographic differentiation, being nearly fixed in sub-Saharan African populations and mostly absent in non-African populations (Mercereau-Puijalon and Ménard, 2010; Howes et al., 2011; Gething et al., 2012), and multiple populations with ancestry from sub-Saharan Africa show evidence for admixture-enabled adaptation throughout human history at the *DARC* locus (Hodgson et al., 2014; Triska et al., 2015; Laso-Jadart et al., 2017; Pierron et al., 2018 Fernandes et al., 2019). Further, the Duffy-null allele has well-characterized functional protection against *P. vivax*, a malaria parasite with a documented record of recurrent transmission in Santiago since settlement.

We demonstrated how selection at the *DARC* locus may have affected patterns of ancestry genome-wide (**Fig. 4**). The initial admixture contributions inferred under our model were lower than those observed in Santiago today, and we confirmed this pattern more generally in simulations. Population-genetic studies often use loci far from potentially selected sites as putatively neutrally-evolving loci, or treat many dispersed loci as neutral based on the assumption that a few selected sites will not dramatically change genome-wide distributions of summary statistics. However, we found that selection on a single locus may shape patterns of ancestry genome-wide. On short timescales, individuals with higher genome-wide proportions of ancestry from the source population carrying the beneficial allele are more likely to have a selective advantage in early generations post-admixture as recombination has yet to uncouple global ancestry proportion and the local ancestry of the selected allele (Pierron et al., 2018). Therefore, since global ancestry is often used to infer initial admixture contributions, these and related demographic inferences may be biased under the common assumption that genome-wide patterns of ancestry reflect demography alone.

Difference in observed and simulated global ancestry may be caused by a variety of statistical and evolutionary processes beyond selection, including sampling, estimation method, demographic model misspecification, or drift. Regions aside from *DARC* may also have been under selection post-admixture in Santiago, and therefore affected the global ancestry patterns. For example, we also observed a cluster of extreme negative *iDAT* values on chromosome 16 [~chr16: 35000000-65000000] that may be of interest for future study.

The framework developed here is broadly applicable to detect and characterize selection in other recently admixed human and non-human populations. For most analyses, we assumed a single admixture event 20 generations in the past. The simulation software we use, SLiM, makes it straightforward to consider other models of population history specific to populations of interest for future studies. Further research into the interaction of selection and demography will refine inference. Additionally, while the timing of admixture in Cabo Verde is well documented, our ABC approach may be extended to infer the timing of selection and admixture as well, as the

tract length distribution is informative about these parameters (Gravel, 2012; Liang and Nielsen, 2014). Indeed, we demonstrated the need to jointly infer selection and demographic histories.

Code & data availability. Scripts for analyses, simulations, and to reproduce figures can be found at https://github.com/agoldberglab/CV_DuffySelection. Sampling consent forms do not allow for public release of genotype data.

Acknowledgements. We thank Joshua Schraiber for useful discussions. We acknowledge support from NIH grant R35 GM133481 to AG, and NIH grant F32 GM139313 to KLK. We thank Hua Tang and Greg Barsh for generating genetic data used in this study.

Author Contributions. IH, KK, and AG designed and performed research. SB provided genetic data. IH and AG wrote the paper, with input from all authors. All authors discussed results and reviewed and commented on the manuscript.

Methods

Genetic data & ancestry inference

For this study, we used SNP array data from (Beleza et al., 2013) which included 564 admixed individuals across the island regions of Cabo Verde. From this dataset, we filtered individuals with greater than 5% missing calls overall or greater than 10% missing calls on a single chromosome. This resulted in removal of one individual with high missingness on chromosome 14 (11.36%). We merged genotypes for the remaining 563 individuals with genotypes from 107 IBS (Iberian Population in Spain) and 107 GWD (Gambian in Western Division - Mandinka) samples from high-coverage resequencing data released through the International Genome Sample Resource (Clarke et al., 2017; Fairley et al., 2020). Our analyses considered autosomal chromosomes only. We selected biallelic SNPs occurring in both the Cabo Verde samples and the reference samples. The final merged dataset contained 884,656 autosomal SNPs. Average missingness by SNP was 0.0017.

Using the 884,656 autosomal SNP dataset, we performed phasing with SHAPEIT2 using the Phase 3, NCBI build 37 (hg19) reference panel of haplotypes and associated genetic map in IMPUTE2 format (Delaneau et al., 2013). Following the SHAPEIT documentation, we first ran SHAPEIT -check to exclude sites not contained within the reference map, followed by SHAPEIT phasing to yield phased genotypes at 881,279 SNPs. For local ancestry inference, we ran RFMix v1.5 on the phased samples using a two-way admixture model (Maples et al., 2013). We used the RFMix PopPhased program with default window size, the --use-reference-panels-in-EM option, -e = 2 (2 EM iterations), and --forward-backward. Ancestry references for European and West African source populations, respectively, were IBS and GWD.

Global ancestry inferred in ADMIXTURE (Alexander and Lange, 2011) was averaged over ten independent runs with randomly chosen seeds using supervised cluster mode with the GWD and IBS individuals specified as the reference populations.

The island of Boa Vista was excluded from analyses due to our small sample from the region (26 individuals), leaving a final set of 537 individuals across the three island regions considered in this study (Santiago: 172, Fogo: 129, NW Cluster: 236; **Fig. 1A**).

Integrated Decay in Ancestry Tract (*iDAT*) score

In order to account for global ancestry patterns that contribute to the ancestry tract-length distribution when identifying loci under selection post-admixture, we developed the integrated Decay in Ancestry Tract (*iDAT*) score. This statistic quantifies the length and homozygosity of the tract-length distribution by comparing the decay of tract lengths from alternate ancestries at increasing distance from a site of interest.

First, we describe the Decay in Ancestry Tract (DAT) feature, which is calculated similarly to expected haplotype homozygosity (EHH) (Sabeti et al., 2002), using local ancestry tracts rather than haplotypes. For each source population i we calculate:

$$DAT_i(x) = \left(\frac{n_x}{n}\right)^2,$$

where n_x is the number of ancestry tracts that extend some absolute distance x from a position of interest, and n is the total number of ancestry tracts (extending in either direction) that contain the site of interest. We calculate DAT at increasing distances from the site of interest. Similar to integrated haplotype homozygosity (iHH) (Voight et al., 2006), we then calculate the area under the curve for DAT as a function of distance from the position of interest, producing an integrated DAT ($iDAT$) (**Fig. 2B**). For this study, we calculated $iDAT$ only for distances where $DAT \geq 0.25$; that is, where at least half of the ancestry tracts extend that absolute distance from the site of interest. We compare the difference in order of magnitude between $iDAT$ for each ancestry, analogous to the integrated haplotype score (iHS), we have,

$$iDAT\ score = \ln\left(\frac{iDAT_2}{iDAT_1}\right),$$

where $iDAT_i$ is the $iDAT$ calculated for source population i. The length of ancestry tracts, and therefore the $iDAT$ score, will be influenced by relative ancestry contributions from each source population. So, when possible, we standardized the $iDAT$ score using the empirical distribution of $iDAT$ scores for 10,000 random positions across the genome. In this way, we deviate from the calculation of standardized iHS, which is standardized by the empirical distribution of SNPs with the same allele frequency because the length of haplotypes will be affected by the age of a variant (Voight et al., 2006). In the case of admixed populations, we instead need to account for the effect of global ancestry proportion on ancestry tract lengths, where the majority ancestry will tend to have longer contiguous tracts. We standardized by the genome-wide empirical distribution of $iDAT$ scores, rather than $iDAT$ scores for variants with the same local ancestry proportion, because the variance in local ancestry across the genome can be heavily influenced by drift. That is, to standardize the $iDAT$ score, we calculate

$$\frac{\ln\left(\frac{iDAT_2}{iDAT_1}\right) - E\left[\ln\left(\frac{iDAT_2}{iDAT_1}\right)\right]}{SD\left[\ln\left(\frac{iDAT_2}{iDAT_1}\right)\right]}.$$

Single-chromosome simulations

We used SLiM forward simulations with tree-sequence recording to track local ancestry (Haller et al., 2019; Haller and Messer, 2019). We considered eight different demographic scenarios: combinations of initial population size (N=10000 or N=1000) with either a constant population size or exponential growth at a rate of 0.05 per generation, and either a single pulse of admixture at the start of simulation or continuous admixture at 1% total new migrants per generation

(**Table S2**). The proportion of new migrants from each source population was weighted by the respective initial admixture contributions.

For each demographic scenario, we generated 1000 simulations of the human chromosome 1. For realistic recombination rates, we used the human genetic map from IMPUTE2 (https://mathgen.stats.ox.ac.uk/impute/1000GP_Phase3.html). We simulated admixture from two source populations to form a third admixed population (similar to recipe 17.5 in the SLiM manual) (Haller and Messer, 2016). One source population, representing the West African source population, was fixed for a neutral variant at the same position as the Duffy-null allele (chr1:159174683; GRCh37 coordinates in accordance with genetic map). Because the true West African ancestry contribution is unknown, for neutral simulations this parameter was drawn from a uniform distribution with lower and upper bounds at 0.65 and 0.75, respectively. We simulated the admixed population for 20 generations.

Using the tree-sequence files output from the SLiM simulations, we calculated the five ancestry-based summary statistics for each simulation: West African local ancestry proportion at *DARC*, variance in West African local ancestry proportion across the chromosome, mean and median West African ancestry tract length containing Duffy-null, and unstandardized *iDAT* score for the Duffy-null variant. We used the unstandardized *iDAT* score because there is no genome-wide distribution of *iDAT* scores for single-chromosome simulations. *iDAT* scores also could not be standardized using the distribution of simulated Duffy-null *iDAT* scores because each simulation had a different starting admixture proportion.

SWIF(r) implementation

We incorporated ancestry-based summary statistics and admixture simulations into the SWIF(r) framework developed by (Sugden et al., 2018). For the simulated training set, we followed the single-chromosome simulation framework described above. We considered a single realistic demographic scenario for training: starting population size N=10000, exponential growth at a rate of 0.05 per generation, and single-pulse admixture with starting West African ancestry contribution randomly drawn from a uniform distribution from 0.65 to 0.75.

For simulations with selection at a single locus, we assumed an additive selection model (*h*=0.5). Selection coefficient was randomly drawn from a uniform distribution from 0 to 0.2. We calculated the five ancestry-based summary statistics for each simulation. Under this model, we generated 50000 neutral simulations and 100 sweep simulations for training.

To validate this extension of SWIF(r), we generated a new set of 1000 neutral simulations and 1000 sweep simulations. **Figure S4** shows a precision-recall plot for a SWIF(r) implementation using the five ancestry-based statistics, a prior sweep probability of 0.2% (reflecting the training set proportions above), and two classes (neutral or sweep). **Figure S5** demonstrates SWIF(r) performance for each class and across values of admixture contribution and selection coefficient. This SWIF(r) implementation had a low rate of false-negative classification of neutral simulations as sweeps.

Inference of selection under a deterministic population-genetic model

To estimate the selection coefficient for the Duffy-null allele based on the dominance and the allele frequency trajectory over 20 generations, we used a deterministic population-genetic model of selection. We used the following recursion equation (Coop, 2020):

$$p_{t+1} = p_t + \frac{p_t hs + q_t s(1-h)}{1 - 2p_t q_t sh - q_t^2 s} p_t q_t,$$

where p_t is the frequency of Duffy-null in a given generation t, q_t is the frequency of the alternate allele in generation t, s is the selection coefficient which is constant over time, and h is the dominance coefficient (0 if Duffy-null is dominant, 1 if recessive, and 0.5 in an additive selective model). We calculated the allele frequency over a grid of values for pairs of $h \in [0,1]$ and $s \in [0,0.2]$ in 0.005 and 0.001 increments, respectively. **Figure 3A** shows the combination of h and s that produce $\left| p_{20} - p_{\text{Duffy}} \right| < 0.01$, where p_{20} is the calculated frequency of the selected allele after 20 generations and p_{Duffy} is the observed frequency of Duffy-null in Santiago (0.834). Because the true initial allele frequency p_o is not known, we performed the analysis for three reasonable starting allele frequencies: $p_o = 0.65$, $p_o = 0.70$, and $p_o = 0.75$.

Inference of selection coefficient using approximate Bayesian computation

We used an approximate Bayesian computation (ABC) framework with non-linear regression (neural network) adjustment to jointly estimate the posterior distributions for selection strength acting on the Duffy-null allele and the initial West African ancestry contribution (*R* package '*abc*', with 'neuralnet' method) (Csilléry et al., 2012).

We followed the same simulation framework we previously described for our SWIF(r) sweep simulations of chromosome 1, with the following demographic scenario for training: starting population size N=10000, exponential growth at a rate of 0.05 per generation, and single-pulse admixture. We assumed an additive selection model (*h*=0.5). The selection coefficient (*s*) and the initial West African ancestry contribution (*m*) were drawn from uniform prior distributions, $s \sim U(0,0.2)$ and $m \sim U(0.1,0.9)$. We generated 10000 simulations for ABC inference and calculated the five ancestry-based summary statistics for each simulation.

We chose the tolerance and hidden layer sizes for ABC estimation based on the best combination of *RMSE* and R^2 values for leave-one-out cross-validation under different combinations of these hyperparameters. We performed cross-validation for 1000 simulations. Estimation accuracy for this dataset was similar under a variety of hyperparameter choices. We set the number of units in the hidden layer to 'sizenet=2' and the acceptance rate to 'tol=0.05'. For the cross-validation, our estimates of the selection coefficient had an *RMSE* = 0.0083 and R^2 = 0.9785; our estimates of starting admixture proportion had an *RMSE* = 0.0090 and R^2 = 0.9985 (**Fig. S6**).

Global Ancestry Simulations

To assess how selection at a single locus impacts genome-wide patterns of ancestry, we simulated changes in global ancestry over 20 generations (**Fig. 4**). Whole autosome (22 chromosome) simulations with a realistic recombination map is computationally intensive. Instead, we took two complementary approaches.

First, we performed whole autosome simulations (22 independently segregating chromosomes) in SLiM by specifying a "crossover rate" between chromosomes of 0.5 per generation. We used the total lengths of autosomal chromosomes from the Human Genome Assembly GRCh37.p13 (https://www.ncbi.nlm.nih.gov/grc/human/data?asm=GRCh37.p13). We used a uniform recombination rate within each chromosome of 1×10^{-8} crossovers per base position per generation. We considered the following demographic scenario: starting population size N=10000, exponential growth at a rate of 0.05 per generation, and single-pulse admixture. Global ancestry was calculated by taking the mean West African ancestry proportion across the autosome.

Second, we performed two-chromosome simulations (modeled based on human chromosomes 1 and 2) in order to incorporate the human genetic map for variable recombination rates. Two-chromosome simulations were performed under the same demographic model as whole autosome simulations. Global ancestry was calculated by taking a weighted mean of ancestry for the two chromosomes, where chromosome 2 represented the contribution for the ~92% of the genome that segregates independently from chromosome 1.

To test how selection at a single locus affects global ancestry, we considered both whole autosome simulations and two-chromosome simulations, using a West African ancestry contribution of 0.65 for all simulations in the founding generation. We varied the selection coefficient for the simulated Duffy-null variant on chromosome 1 from 0 to 0.2. We assumed an additive selection model (*h*=0.5). The results are similar across simulation methods (**Fig. S7**).

To determine how our ABC estimates of initial West African ancestry contribution differ from final global ancestry after 20 generations of selection at the Duffy-null locus, we next simulated whole autosomes admixing for 20 generations drawing parameters from the previously inferred posterior distributions of the selection coefficient and initial ancestry contribution. Specifically, we used the paired estimates of selection coefficient and West African admixture contribution from each accepted simulation from the ABC analysis and passed those parameters as input for each whole autosome simulation; this produced a distribution of estimates of global ancestry for populations simulated with realistic selection and initial ancestry contributions (**Fig. 4A**).

References

Alexander, D.H., Lange, K., 2011. Enhancements to the ADMIXTURE algorithm for individual ancestry estimation. BMC Bioinformatics 12, 246. https://doi.org/10.1186/1471-2105-12-246

Belbin, G.M., Nieves-Colón, M.A., Kenny, E.E., Moreno-Estrada, A., Gignoux, C.R., 2018. Genetic diversity in populations across Latin America: implications for population and medical genetic studies. Current Opinion in Genetics & Development, Genetics of Human Origins 53, 98–104. https://doi.org/10.1016/j.gde.2018.07.006

Beleza, S., Campos, J., Lopes, J., Araújo, I.I., Almada, A.H., Silva, A.C. e, Parra, E.J., Rocha, J., 2012. The Admixture Structure and Genetic Variation of the Archipelago of Cape Verde and Its Implications for Admixture Mapping Studies. PLOS ONE 7, e51103. https://doi.org/10.1371/journal.pone.0051103

Beleza, S., Johnson, N.A., Candille, S.I., Absher, D.M., Coram, M.A., Lopes, J., Campos, J., Araújo, I.I., Anderson, T.M., Vilhjálmsson, B.J., Nordborg, M., Silva, A.C. e, Shriver, M.D., Rocha, J., Barsh, G.S., Tang, H., 2013. Genetic Architecture of Skin and Eye Color in an African-European Admixed Population. PLOS Genetics 9, e1003372. https://doi.org/10.1371/journal.pgen.1003372

Bhatia, G., Tandon, A., Patterson, N., Aldrich, M.C., Ambrosone, C.B., Amos, C., Bandera, E.V., Berndt, S.I., Bernstein, L., Blot, W.J., Bock, C.H., Caporaso, N., Casey, G., Deming, S.L., Diver, W.R., Gapstur, S.M., Gillanders, E.M., Harris, C.C., Henderson, B.E., Ingles, S.A., Isaacs, W., De Jager, P.L., John, E.M., Kittles, R.A., Larkin, E., McNeill, L.H., Millikan, R.C., Murphy, A., Neslund-Dudas, C., Nyante, S., Press, M.F., Rodriguez-Gil, J.L., Rybicki, B.A., Schwartz, A.G., Signorello, L.B., Spitz, M., Strom, S.S., Tucker, M.A., Wiencke, J.K., Witte, J.S., Wu, X., Yamamura, Y., Zanetti, K.A., Zheng, W., Ziegler, R.G., Chanock, S.J., Haiman, C.A., Reich, D., Price, A.L., 2014. Genome-wide Scan of 29,141 African Americans Finds No Evidence of Directional Selection since Admixture. The American Journal of Human Genetics 95, 437–444. https://doi.org/10.1016/j.ajhg.2014.08.011

Bryc, K., Durand, E.Y., Macpherson, J.M., Reich, D., Mountain, J.L., 2015. The Genetic Ancestry of African Americans, Latinos, and European Americans across the United States. The American Journal of Human Genetics 96, 37–53. https://doi.org/10.1016/j.ajhg.2014.11.010

Busby, G.B., Band, G., Si Le, Q., Jallow, M., Bougama, E., Mangano, V.D., Amenga-Etego, L.N., Enimil, A., Apinjoh, T., Ndila, C.M., Manjurano, A., Nyirongo, V., Doumba, O., Rockett, K.A., Kwiatkowski, D.P., Spencer, C.C., Malaria Genomic Epidemiology Network, 2016. Admixture into and within sub-Saharan Africa. eLife 5, e15266. https://doi.org/10.7554/eLife.15266

Busby, G.B.J., Hellenthal, G., Montinaro, F., Tofanelli, S., Bulayeva, K., Rudan, I., Zemunik, T., Hayward, C., Toncheva, D., Karachanak-Yankova, S., Nesheva, D., Anagnostou, P., Cali, F., Brisighelli, F., Romano, V., Lefranc, G., Buresi, C., Ben Chibani, J., Haj-Khelil, A., Denden, S., Ploski, R., Krajewski, P., Hervig, T., Moen, T., Herrera, R.J., Wilson, J.F., Myers, S., Capelli, C., 2015. The Role of Recent Admixture in Forming the Contemporary West Eurasian Genomic Landscape. Current Biology 25, 2518–2526. https://doi.org/10.1016/j.cub.2015.08.007

Calfee, E., Agra, M.N., Palacio, M.A., Ramírez, S.R., Coop, G., 2020. Selection and hybridization shaped the Africanized honey bee invasion of the Americas. bioRxiv 2020.03.17.994632. https://doi.org/10.1101/2020.03.17.994632

Cavasini, C.E., de Mattos, L.C., Couto, Á.A.D., Couto, V.S.D., Gollino, Y., Moretti, L.J., Bonini-Domingos, C.R., Rossit, A.R., Castilho, L., Machado, R.L., 2007. Duffy blood group gene polymorphisms among malaria vivax patients in four areas of the Brazilian Amazon region. Malaria Journal 6, 167. https://doi.org/10.1186/1475-2875-6-167

Clarke, L., Fairley, S., Zheng-Bradley, X., Streeter, I., Perry, E., Lowy, E., Tassé, A.-M., Flicek, P., 2017. The international Genome sample resource (IGSR): A worldwide collection of genome variation incorporating the 1000 Genomes Project data. Nucleic Acids Res 45, D854–D859. https://doi.org/10.1093/nar/gkw829

Coop, G., 2020. Population and Quantitative Genetics. Github. URL https://github.com/cooplab/popgen-notes/blob/master/popgen_notes.pdf (accessed 6.12.20).

Csilléry, K., François, O., Blum, M.G.B., 2012. abc: an R package for approximate Bayesian computation (ABC). Methods in Ecology and Evolution 3, 475–479. https://doi.org/10.1111/j.2041-210X.2011.00179.x

Delaneau, O., Zagury, J.-F., Marchini, J., 2013. Improved whole-chromosome phasing for disease and population genetic studies. Nature Methods 10, 5–6. https://doi.org/10.1038/nmeth.2307

DePina, A.J., Dia, A.K., de Ascenção Soares Martins, A., Ferreira, M.C., Moreira, A.L., Leal, S.V., Pires, C.M., Moreira, J.M.G., Tavares, M.F., da Moura, A.J.F., Pereira, J.M., Faye, O., Seck, I., Niang, E.H.A., 2019. Knowledge, attitudes and practices about malaria in Cabo Verde: a country in the pre-elimination context. BMC Public Health 19, 850. https://doi.org/10.1186/s12889-019-7130-5

DePina, A.J., Niang, E.H.A., Barbosa Andrade, A.J., Dia, A.K., Moreira, A., Faye, O., Seck, I., 2018. Achievement of malaria pre-elimination in Cape Verde according to the data collected from 2010 to 2016. Malaria Journal 17, 236. https://doi.org/10.1186/s12936-018-2376-4

Fairley, S., Lowy-Gallego, E., Perry, E., Flicek, P., 2020. The International Genome Sample Resource (IGSR) collection of open human genomic variation resources. Nucleic Acids Res 48, D941–D947. https://doi.org/10.1093/nar/gkz836

Fernandes, A.T., Velosa, R., Jesus, J., Carracedo, A., Brehm, A., 2003. Genetic Differentiation of the Cabo Verde Archipelago Population Analysed by STR Polymorphisms. Annals of Human Genetics 67, 340–347. https://doi.org/10.1046/j.1469-1809.2003.00029.x

Fernandes, V., Brucato, N., Ferreira, J.C., Pedro, N., Cavadas, B., Ricaut, F.-X., Alshamali, F., Pereira, L., 2019. Genome-Wide Characterization of Arabian Peninsula Populations: Shedding Light on the History of a Fundamental Bridge between Continents. Mol Biol Evol 36, 575–586. https://doi.org/10.1093/molbev/msz005

Ferreira, M., Moreira, A., Correia, A., De Pina, A., 2017. Malaria in Cabo Verde: Pre-elimination versus the Elimination. https://doi.org/10.13140/RG.2.2.28878.72004

Field, Y., Boyle, E.A., Telis, N., Gao, Z., Gaulton, K.J., Golan, D., Yengo, L., Rocheleau, G., Froguel, P., McCarthy, M.I., Pritchard, J.K., 2016. Detection of human adaptation during the past 2000 years. Science 354, 760–764. https://doi.org/10.1126/science.aag0776

Fortes-Lima, C.A., Laurent, R., Thouzeau, V., Toupance, B., Verdu, P., 2019. Complex genetic admixture histories reconstructed with Approximate Bayesian Computations. bioRxiv 761452. https://doi.org/10.1101/761452

Fumagalli, M., Sironi, M., Pozzoli, U., Ferrer-Admettla, A., Pattini, L., Nielsen, R., 2011. Signatures of Environmental Genetic Adaptation Pinpoint Pathogens as the Main Selective Pressure through Human Evolution. PLOS Genetics 7, e1002355. https://doi.org/10.1371/journal.pgen.1002355

Gething, P.W., Elyazar, I.R.F., Moyes, C.L., Smith, D.L., Battle, K.E., Guerra, C.A., Patil, A.P., Tatem, A.J., Howes, R.E., Myers, M.F., George, D.B., Horby, P., Wertheim, H.F.L., Price, R.N., Müeller, I., Baird, J.K., Hay, S.I., 2012. A Long Neglected World Malaria Map: Plasmodium vivax Endemicity in 2010. PLOS Neglected Tropical Diseases 6, e1814. https://doi.org/10.1371/journal.pntd.0001814

Gravel, S., 2012. Population Genetics Models of Local Ancestry. Genetics 191, 607–619. https://doi.org/10.1534/genetics.112.139808

Grossman, S.R., Andersen, K.G., Shlyakhter, I., Tabrizi, S., Winnicki, S., Yen, A., Park, D.J., Griesemer, D., Karlsson, E.K., Wong, S.H., Cabili, M., Adegbola, R.A., Bamezai, R.N.K., Hill, A.V.S., Vannberg, F.O., Rinn, J.L., Lander, E.S., Schaffner, S.F., Sabeti, P.C., 2013. Identifying Recent Adaptations in Large-Scale Genomic Data. Cell 152, 703–713. https://doi.org/10.1016/j.cell.2013.01.035

Haller, B.C., Galloway, J., Kelleher, J., Messer, P.W., Ralph, P.L., 2019. Tree-sequence recording in SLiM opens new horizons for forward-time simulation of whole genomes. Molecular Ecology Resources 19, 552–566. https://doi.org/10.1111/1755-0998.12968

Haller, B.C., Messer, P.W., 2019. SLiM 3: Forward Genetic Simulations Beyond the Wright–Fisher Model. Mol Biol Evol 36, 632–637. https://doi.org/10.1093/molbev/msy228

Haller, B.C., Messer, P.W., 2016. SLiM: An Evolutionary Simulation Framework.

Hamblin, M.T., Di Rienzo, A., 2000. Detection of the Signature of Natural Selection in Humans: Evidence from the Duffy Blood Group Locus. The American Journal of Human Genetics 66, 1669–1679. https://doi.org/10.1086/302879

Hamblin, M.T., Thompson, E.E., Di Rienzo, A., 2002. Complex Signatures of Natural Selection at the Duffy Blood Group Locus. The American Journal of Human Genetics 70, 369–383. https://doi.org/10.1086/338628

Hedrick, P.W., 2013. Adaptive introgression in animals: examples and comparison to new mutation and standing variation as sources of adaptive variation. Molecular Ecology 22, 4606–4618. https://doi.org/10.1111/mec.12415

Hellenthal, G., Busby, G.B.J., Band, G., Wilson, J.F., Capelli, C., Falush, D., Myers, S., 2014. A Genetic Atlas of Human Admixture History. Science 343, 747–751. https://doi.org/10.1126/science.1243518

Hermisson, J., Pennings, P.S., 2005. Soft Sweeps: Molecular Population Genetics of Adaptation From Standing Genetic Variation. Genetics 169, 2335–2352. https://doi.org/10.1534/genetics.104.036947

Hodgson, J.A., Pickrell, J.K., Pearson, L.N., Quillen, E.E., Prista, A., Rocha, J., Soodyall, H., Shriver, M.D., Perry, G.H., 2014. Natural selection for the Duffy-null allele in the recently admixed people of Madagascar. Proceedings of the Royal Society B: Biological Sciences 281, 20140930. https://doi.org/10.1098/rspb.2014.0930

Howes, R.E., Patil, A.P., Piel, F.B., Nyangiri, O.A., Kabaria, C.W., Gething, P.W., Zimmerman, P.A., Barnadas, C., Beall, C.M., Gebremedhin, A., Ménard, D., Williams, T.N.,

Weatherall, D.J., Hay, S.I., 2011. The global distribution of the Duffy blood group. Nat Commun 2, 1–10. https://doi.org/10.1038/ncomms1265

Huerta-Sánchez, E., Jin, X., Asan, Bianba, Z., Peter, B.M., Vinckenbosch, N., Liang, Y., Yi, X., He, M., Somel, M., Ni, P., Wang, B., Ou, X., Huasang, Luosang, J., Cuo, Z.X.P., Li, K., Gao, G., Yin, Y., Wang, W., Zhang, X., Xu, X., Yang, H., Li, Y., Wang, Jian, Wang, Jun, Nielsen, R., 2014. Altitude adaptation in Tibetans caused by introgression of Denisovan-like DNA. Nature 512, 194–197. https://doi.org/10.1038/nature13408

Jagoda, E., Lawson, D.J., Wall, J.D., Lambert, D., Muller, C., Westaway, M., Leavesley, M., Capellini, T.D., Mirazón Lahr, M., Gerbault, P., Thomas, M.G., Migliano, A.B., Willerslev, E., Metspalu, M., Pagani, L., 2018. Disentangling Immediate Adaptive Introgression from Selection on Standing Introgressed Variation in Humans. Mol Biol Evol 35, 623–630. https://doi.org/10.1093/molbev/msx314

Jeong, C., Alkorta-Aranburu, G., Basnyat, B., Neupane, M., Witonsky, D.B., Pritchard, J.K., Beall, C.M., Rienzo, A.D., 2014. Admixture facilitates genetic adaptations to high altitude in Tibet. Nat Commun 5, 1–7. https://doi.org/10.1038/ncomms4281

Jin, W., Xu, S., Wang, H., Yu, Y., Shen, Y., Wu, B., Jin, L., 2012. Genome-wide detection of natural selection in African Americans pre- and post-admixture. Genome Res. 22, 519–527. https://doi.org/10.1101/gr.124784.111

Kano, F.S., de Souza, A.M., de Menezes Torres, L., Costa, M.A., Souza-Silva, F.A., Sanchez, B.A.M., Fontes, C.J.F., Soares, I.S., de Brito, C.F.A., Carvalho, L.H., Sousa, T.N., 2018. Susceptibility to Plasmodium vivax malaria associated with DARC (Duffy antigen) polymorphisms is influenced by the time of exposure to malaria. Scientific Reports 8, 1–14. https://doi.org/10.1038/s41598-018-32254-z

Kim, B.Y., Huber, C.D., Lohmueller, K.E., 2018. Deleterious variation shapes the genomic landscape of introgression. PLOS Genetics 14, e1007741. https://doi.org/10.1371/journal.pgen.1007741

Kim, Y., Nielsen, R., 2004. Linkage Disequilibrium as a Signature of Selective Sweeps. Genetics 167, 1513–1524. https://doi.org/10.1534/genetics.103.025387

Kwiatkowski, D.P., 2005. How Malaria Has Affected the Human Genome and What Human Genetics Can Teach Us about Malaria. The American Journal of Human Genetics 77, 171–192. https://doi.org/10.1086/432519

Lachance, J., Tishkoff, S.A., 2013. Population Genomics of Human Adaptation. Annu. Rev. Ecol. Evol. Syst. 44, 123–143. https://doi.org/10.1146/annurev-ecolsys-110512-135833

Lamason, R.L., Mohideen, M.-A.P.K., Mest, J.R., Wong, A.C., Norton, H.L., Aros, M.C., Jurynec, M.J., Mao, X., Humphreville, V.R., Humbert, J.E., Sinha, S., Moore, J.L., Jagadeeswaran, P., Zhao, W., Ning, G., Makalowska, I., McKeigue, P.M., O'Donnell, D., Kittles, R., Parra, E.J., Mangini, N.J., Grunwald, D.J., Shriver, M.D., Canfield, V.A., Cheng, K.C., 2005. SLC24A5, a Putative Cation Exchanger, Affects Pigmentation in Zebrafish and Humans. Science 310, 1782–1786. https://doi.org/10.1126/science.1116238

Laso-Jadart, R., Harmant, C., Quach, H., Zidane, N., Tyler-Smith, C., Mehdi, Q., Ayub, Q., Quintana-Murci, L., Patin, E., 2017. The Genetic Legacy of the Indian Ocean Slave Trade: Recent Admixture and Post-admixture Selection in the Makranis of Pakistan. The American Journal of Human Genetics 101, 977–984. https://doi.org/10.1016/j.ajhg.2017.09.025

Liang, M., Nielsen, R., 2014. The Lengths of Admixture Tracts. Genetics 197, 953–967. https://doi.org/10.1534/genetics.114.162362

Lindo, J., Huerta-Sánchez, E., Nakagome, S., Rasmussen, M., Petzelt, B., Mitchell, J., Cybulski, J.S., Willerslev, E., DeGiorgio, M., Malhi, R.S., 2016. A time transect of exomes from a Native American population before and after European contact. Nature Communications 7, 13175. https://doi.org/10.1038/ncomms13175

Lohmueller, K.E., Bustamante, C.D., Clark, A.G., 2011. Detecting Directional Selection in the Presence of Recent Admixture in African-Americans. Genetics 187, 823–835. https://doi.org/10.1534/genetics.110.122739

Lohmueller, K.E., Bustamante, C.D., Clark, A.G., 2010. The Effect of Recent Admixture on Inference of Ancient Human Population History. Genetics 185, 611–622. https://doi.org/10.1534/genetics.109.113761

Lopez, M., Choin, J., Sikora, M., Siddle, K., Harmant, C., Costa, H.A., Silvert, M., Mouguiama-Daouda, P., Hombert, J.-M., Froment, A., Le Bomin, S., Perry, G.H., Barreiro, L.B., Bustamante, C.D., Verdu, P., Patin, E., Quintana-Murci, L., 2019. Genomic Evidence for Local Adaptation of Hunter-Gatherers to the African Rainforest. Current Biology 29, 2926-2935.e4. https://doi.org/10.1016/j.cub.2019.07.013

Maples, B.K., Gravel, S., Kenny, E.E., Bustamante, C.D., 2013. RFMix: A Discriminative Modeling Approach for Rapid and Robust Local-Ancestry Inference. The American Journal of Human Genetics 93, 278–288. https://doi.org/10.1016/j.ajhg.2013.06.020

Mathias, R.A., Taub, M.A., Gignoux, C.R., Fu, W., Musharoff, S., O'Connor, T.D., Vergara, C., Torgerson, D.G., Pino-Yanes, M., Shringarpure, S.S., Huang, L., Rafaels, N., Boorgula, M.P., Johnston, H.R., Ortega, V.E., Levin, A.M., Song, W., Torres, R., Padhukasahasram, B., Eng, C., Mejia-Mejia, D.-A., Ferguson, T., Qin, Z.S., Scott, A.F., Yazdanbakhsh, M., Wilson, J.G., Marrugo, J., Lange, L.A., Kumar, R., Avila, P.C., Williams, L.K., Watson, H., Ware, L.B., Olopade, C., Olopade, O., Oliveira, R., Ober, C., Nicolae, D.L., Meyers, D., Mayorga, A., Knight-Madden, J., Hartert, T., Hansel, N.N., Foreman, M.G., Ford, J.G., Faruque, M.U., Dunston, G.M., Caraballo, L., Burchard, E.G., Bleecker, E., Araujo, M.I., Herrera-Paz, E.F., Gietzen, K., Grus, W.E., Bamshad, M., Bustamante, C.D., Kenny, E.E., Hernandez, R.D., Beaty, T.H., Ruczinski, I., Akey, J., Barnes, K.C., 2016. A continuum of admixture in the Western Hemisphere revealed by the African Diaspora genome. Nature Communications 7, 1–10. https://doi.org/10.1038/ncomms12522

McManus, K.F., Taravella, A.M., Henn, B.M., Bustamante, C.D., Sikora, M., Cornejo, O.E., 2017. Population genetic analysis of the DARC locus (Duffy) reveals adaptation from standing variation associated with malaria resistance in humans. PLOS Genetics 13, e1006560. https://doi.org/10.1371/journal.pgen.1006560

Mercereau-Puijalon, O., Ménard, D., 2010. Plasmodium vivax and the Duffy antigen: A paradigm revisited. Transfusion Clinique et Biologique, "Role of blood group antigens and associated molecules in red cell biology: from molecular approach to clinical applications" organized by Institut National de la Transfusion Sanguine, September 15th to 17th 2010 17, 176–183. https://doi.org/10.1016/j.tracli.2010.06.005

Moorjani, P., Patterson, N., Hirschhorn, J.N., Keinan, A., Hao, L., Atzmon, G., Burns, E., Ostrer, H., Price, A.L., Reich, D., 2011. The History of African Gene Flow into Southern Europeans, Levantines, and Jews. PLOS Genetics 7, e1001373. https://doi.org/10.1371/journal.pgen.1001373

Moreno-Estrada, A., Gravel, S., Zakharia, F., McCauley, J.L., Byrnes, J.K., Gignoux, C.R., Ortiz-Tello, P.A., Martínez, R.J., Hedges, D.J., Morris, R.W., Eng, C., Sandoval, K., Acevedo-Acevedo, S., Norman, P.J., Layrisse, Z., Parham, P., Martínez-Cruzado, J.C., Burchard, E.G., Cuccaro, M.L., Martin, E.R., Bustamante, C.D., 2013. Reconstructing the Population Genetic History of the Caribbean. PLOS Genetics 9, e1003925. https://doi.org/10.1371/journal.pgen.1003925

Nielsen, R., Akey, J.M., Jakobsson, M., Pritchard, J.K., Tishkoff, S., Willerslev, E., 2017. Tracing the peopling of the world through genomics. Nature 541, 302–310. https://doi.org/10.1038/nature21347

Nielsen, R., Hellmann, I., Hubisz, M., Bustamante, C., Clark, A.G., 2007. Recent and ongoing selection in the human genome. Nat Rev Genet 8, 857–868. https://doi.org/10.1038/nrg2187

Norris, E.T., Rishishwar, L., Chande, A.T., Conley, A.B., Ye, K., Valderrama-Aguirre, A., Jordan, I.K., 2020. Admixture-enabled selection for rapid adaptive evolution in the Americas. Genome Biology 21, 29. https://doi.org/10.1186/s13059-020-1946-2

Norris, E.T., Rishishwar, L., Chande, A.T., Conley, A.B., Ye, K., Valderrama-Aguirre, A., Jordan, I.K., 2019. Admixture-enabled selection for rapid adaptive evolution in the Americas. bioRxiv 783845. https://doi.org/10.1101/783845

Norton, H.L., Kittles, R.A., Parra, E., McKeigue, P., Mao, X., Cheng, K., Canfield, V.A., Bradley, D.G., McEvoy, B., Shriver, M.D., 2007. Genetic Evidence for the Convergent Evolution of Light Skin in Europeans and East Asians. Mol Biol Evol 24, 710–722. https://doi.org/10.1093/molbev/msl203

Ohashi, J., Naka, I., Patarapotikul, J., Hananantachai, H., Brittenham, G., Looareesuwan, S., Clark, A.G., Tokunaga, K., 2004. Extended Linkage Disequilibrium Surrounding the Hemoglobin E Variant Due to Malarial Selection. The American Journal of Human Genetics 74, 1198–1208. https://doi.org/10.1086/421330

Patin, E., Lopez, M., Grollemund, R., Verdu, P., Harmant, C., Quach, H., Laval, G., Perry, G.H., Barreiro, L.B., Froment, A., Heyer, E., Massougbodji, A., Fortes-Lima, C., Migot-Nabias, F., Bellis, G., Dugoujon, J.-M., Pereira, J.B., Fernandes, V., Pereira, L., Veen, L.V. der, Mouguiama-Daouda, P., Bustamante, C.D., Hombert, J.-M., Quintana-Murci, L., 2017. Dispersals and genetic adaptation of Bantu-speaking populations in Africa and North America. Science 356, 543–546. https://doi.org/10.1126/science.aal1988

Patterson, K.D., 1988. Epidemics, Famines, and Population in the Cape Verde Islands, 1580-1900. The International Journal of African Historical Studies 21, 291–313. https://doi.org/10.2307/219938

Pickrell, J.K., Coop, G., Novembre, J., Kudaravalli, S., Li, J.Z., Absher, D., Srinivasan, B.S., Barsh, G.S., Myers, R.M., Feldman, M.W., Pritchard, J.K., 2009. Signals of recent positive selection in a worldwide sample of human populations. Genome Res. 19, 826–837. https://doi.org/10.1101/gr.087577.108

Pierron, D., Heiske, M., Razafindrazaka, H., Pereda-loth, V., Sanchez, J., Alva, O., Arachiche, A., Boland, A., Olaso, R., Deleuze, J.-F., Ricaut, F.-X., Rakotoarisoa, J.-A., Radimilahy, C., Stoneking, M., Letellier, T., 2018. Strong selection during the last millennium for African ancestry in the admixed population of Madagascar. Nat Commun 9, 1–9. https://doi.org/10.1038/s41467-018-03342-5

Racimo, F., Marnetto, D., Huerta-Sánchez, E., 2017. Signatures of Archaic Adaptive Introgression in Present-Day Human Populations. Mol Biol Evol 34, 296–317. https://doi.org/10.1093/molbev/msw216

Racimo, F., Sankararaman, S., Nielsen, R., Huerta-Sánchez, E., 2015. Evidence for archaic adaptive introgression in humans. Nature Reviews Genetics 16, 359–371. https://doi.org/10.1038/nrg3936

Rishishwar, L., Conley, A.B., Wigington, C.H., Wang, L., Valderrama-Aguirre, A., Jordan, I.K., 2015. Ancestry, admixture and fitness in Colombian genomes. Sci Rep 5, 1–16. https://doi.org/10.1038/srep12376

Roberts, G. Account of a Voyage to the Islands of the Canaries, Cape de Verde, and Barbados, in 1721, Vol. I in Thomas Astley, ed., A New General Collection of Voyages and Travels (reprinted London, 1968; 1st printed 1745-47), pg. 643

Ruiz-Linares, A., Adhikari, K., Acuña-Alonzo, V., Quinto-Sanchez, M., Jaramillo, C., Arias, W., Fuentes, M., Pizarro, M., Everardo, P., Avila, F. de, Gómez-Valdés, J., León-Mimila, P., Hunemeier, T., Ramallo, V., Cerqueira, C.C.S. de, Burley, M.-W., Konca, E., Oliveira, M.Z. de, Veronez, M.R., Rubio-Codina, M., Attanasio, O., Gibbon, S., Ray, N., Gallo, C., Poletti, G., Rosique, J., Schuler-Faccini, L., Salzano, F.M., Bortolini, M.-C., Canizales-Quinteros, S., Rothhammer, F., Bedoya, G., Balding, D., Gonzalez-José, R., 2014. Admixture in Latin America: Geographic Structure, Phenotypic Diversity and Self-Perception of Ancestry Based on 7,342 Individuals. PLOS Genetics 10, e1004572. https://doi.org/10.1371/journal.pgen.1004572

Sabeti, P.C., Reich, D.E., Higgins, J.M., Levine, H.Z.P., Richter, D.J., Schaffner, S.F., Gabriel, S.B., Platko, J.V., Patterson, N.J., McDonald, G.J., Ackerman, H.C., Campbell, S.J., Altshuler, D., Cooper, R., Kwiatkowski, D., Ward, R., Lander, E.S., 2002. Detecting recent positive selection in the human genome from haplotype structure. Nature 419, 832–837. https://doi.org/10.1038/nature01140

Sabeti, P.C., Schaffner, S.F., Fry, B., Lohmueller, J., Varilly, P., Shamovsky, O., Palma, A., Mikkelsen, T.S., Altshuler, D., Lander, E.S., 2006. Positive Natural Selection in the Human Lineage. Science 312, 1614–1620. https://doi.org/10.1126/science.1124309

Setter, D., Mousset, S., Cheng, X., Nielsen, R., DeGiorgio, M., Hermisson, J., 2020. VolcanoFinder: Genomic scans for adaptive introgression. PLOS Genetics 16, e1008867. https://doi.org/10.1371/journal.pgen.1008867

Snow, R.W., Amratia, P., Kabaria, C.W., Noor, A.M., Marsh, K., 2012. The changing limits and incidence of malaria in Africa: 1939-2009. Adv. Parasitol. 78, 169–262. https://doi.org/10.1016/B978-0-12-394303-3.00010-4

Sousa, T.N., Sanchez, B. a. M., Cerávolo, I.P., Carvalho, L.H., Brito, C.F.A., 2007. Real-time multiplex allele-specific polymerase chain reaction for genotyping of the Duffy antigen, the Plasmodium vivax invasion receptor. Vox Sanguinis 92, 373–380. https://doi.org/10.1111/j.1423-0410.2007.00902.x

Sugden, L.A., Atkinson, E.G., Fischer, A.P., Rong, S., Henn, B.M., Ramachandran, S., 2018. Localization of adaptive variants in human genomes using averaged one-dependence estimation. Nature Communications 9, 1–14. https://doi.org/10.1038/s41467-018-03100-7

Svedberg, J., Shchur, V., Reinman, S., Nielsen, R., Corbett-Detig, R., 2020. Inferring Adaptive Introgression Using Hidden Markov Models. bioRxiv 2020.08.02.232934. https://doi.org/10.1101/2020.08.02.232934

Tang, H., Choudhry, S., Mei, R., Morgan, M., Rodriguez-Cintron, W., Burchard, E.G., Risch, N.J., 2007. Recent Genetic Selection in the Ancestral Admixture of Puerto Ricans. The American Journal of Human Genetics 81, 626–633. https://doi.org/10.1086/520769

Triska, P., Soares, P., Patin, E., Fernandes, V., Cerny, V., Pereira, L., 2015. Extensive Admixture and Selective Pressure Across the Sahel Belt. Genome Biol Evol 7, 3484–3495. https://doi.org/10.1093/gbe/evv236

Verdu, P., Jewett, E.M., Pemberton, T.J., Rosenberg, N.A., Baptista, M., 2017. Parallel Trajectories of Genetic and Linguistic Admixture in a Genetically Admixed Creole Population. Current Biology 27, 2529-2535.e3. https://doi.org/10.1016/j.cub.2017.07.002

Vicuña, L., Klimenkova, O., Norambuena, T., Martinez, F.I., Fernandez, M.I., Shchur, V., Eyheramendy, S., n.d. Post-Admixture Selection on Chileans Targets Haplotype Involved in Pigmentation and Immune Defense Against Pathogens. Genome Biol Evol. https://doi.org/10.1093/gbe/evaa136

Voight, B.F., Kudaravalli, S., Wen, X., Pritchard, J.K., 2006. A Map of Recent Positive Selection in the Human Genome. PLOS Biology 4, e72. https://doi.org/10.1371/journal.pbio.0040072

World Health Organization, Global Malaria Programme, University of California, S.F., Global Health Sciences, Global Health Group, Cabo Verde, Ministério da Saúde, 2012. Moving towards sustainable elimination in Cape Verde. World Health Organization, Geneva.

Yi, X., Liang, Y., Huerta-Sanchez, E., Jin, X., Cuo, Z.X.P., Pool, J.E., Xu, X., Jiang, H., Vinckenbosch, N., Korneliussen, T.S., Zheng, Hancheng, Liu, T., He, W., Li, K., Luo, R., Nie, X., Wu, H., Zhao, M., Cao, H., Zou, J., Shan, Y., Li, Shuzheng, Yang, Q., Asan, Ni, P., Tian, G., Xu, J., Liu, X., Jiang, T., Wu, R., Zhou, G., Tang, M., Qin, J., Wang, T., Feng, S., Li, G., Huasang, Luosang, J., Wang, W., Chen, F., Wang, Y., Zheng, X., Li, Z., Bianba, Z., Yang, G., Wang, X., Tang, S., Gao, G., Chen, Y., Luo, Z., Gusang, L., Cao, Z., Zhang, Q., Ouyang, W., Ren, X., Liang, H., Zheng, Huisong, Huang, Y., Li, J., Bolund, L., Kristiansen, K., Li, Y., Zhang, Y., Zhang, X., Li, R., Li, Songgang, Yang, H., Nielsen, R., Wang, Jun, Wang, Jian, 2010. Sequencing of 50 Human Exomes Reveals Adaptation to High Altitude. Science 329, 75–78. https://doi.org/10.1126/science.1190371

Zhou, Q., Zhao, L., Guan, Y., 2016. Strong Selection at MHC in Mexicans since Admixture. PLOS Genetics 12, e1005847. https://doi.org/10.1371/journal.pgen.1005847

Supplemental Material

Table S1. Expected and observed Duffy-null allele frequencies for each island and source population. Expected Duffy-null frequencies are approximated by mean West African global ancestry proportion for each island, calculated using the ADMIXTURE software.

population	n (sampled individuals)	expected frequency	observed frequency	binomial test p-value
Santiago	172	0.737	0.834	2.193×10^{-5}
Fogo	129	0.498	0.539	0.192
NW Cluster	236	0.552	0.557	0.817
GWD	107	0.997	1.000	-
IBS	107	0.002	0.019	-

Table S2. Demographic models used for single-chromosome simulations.

initial population size (N)	population growth model	population growth rate (per generation)	admixture type	proportion of new migrants (per generation)	scenario number
1000	constant size	-	single-pulse	-	1
			continuous	0.01	2
	exponential	0.05	single-pulse	-	3
			continuous	0.01	4
10000	constant size	-	single-pulse	-	5
			continuous	0.01	6
	exponential	0.05	single-pulse	-	7
			continuous	0.01	8

Figure S1. Mean standardized integrated *DAT* (*iDAT*) score for 20 Mb sliding windows (step size = 1Mb), using standardized *iDAT* for 10000 random positions across the genome for **A)** Fogo and **B)** the Northwest Cluster. Solid grey lines indicate mean windowed standardized *iDAT* score for each island (Fogo, 0.006; NW Cluster, -0.024) and dashed grey lines indicate three standard deviations from the mean. *DARC* locus is indicated by vertical dashed red lines.

Figure S2. Density distributions for five ancestry-based statistics under eight neutral models. Summary statistics were calculated from a random sample of 172 individuals from each simulated population. High population size models correspond to initial N=10000, low population size (high drift) models correspond to initial N=1000. Exponential growth model corresponds to a rate of 0.05 per generation. Continuous migration refers to 1% total new migrants each generation, at the same proportions as initial admixture contributions for each source population. Vertical red line represents each measure's true value for Santiago.

Figure S3. Results of ABC estimation of posterior distributions for **A)** selection coefficient for Duffy-null and **B)** initial West African ancestry contribution for Santiago. Duffy-null allele was modeled as additive (blue; h=0.5), dominant (yellow; h=1 in SLiM), or recessive (pink; h=0 in SLiM). Posterior median estimates for selection coefficient: s_{rec}= 0.052, s_{add}= 0.0795, s_{dom}= 0.183; initial ancestry contribution: m_{rec}= 0.697, m_{add}= 0.690, m_{dom}= 0.665. Prior distributions were $s{\sim}U(0, 0.2)$ and $m{\sim}U(0.1, 0.9)$.

Figure S4. Precision-Recall curve for validation of SWIF(r) classification of neutral and sweep variants, using 1000 neutral and 1000 sweep simulations. With our ancestry-based measures, SWIF(r) achieved an area under the curve (AUC) of 0.966, where an AUC of 1 represents a classifier with perfect skill. Horizontal dashed line indicates the no-skill classifier for this dataset.

Figure S5. SWIF(r) classification results for 1000 neutral and 1000 sweep simulations used for the test set based on Santiago's demographic history. **A)** Confusion matrix with threshold P(sweep) > 0.5. There are no false positives in test set and a high rate of false negatives. **B)** Scatterplot of initial admixture contribution vs selection coefficient. Points colored by P(sweep) as estimated by SWIF(r). The majority of false-negative classifications (i.e. classifying sweeps as neutral) occurred with low starting admixture proportion or low selection strength. If we consider ABC estimates for admixture proportion (~0.7) and selection strength (~0.08), Duffy-null on Santiago would sit around the transition from high to low rate of false negatives. SWIF(r) returned high probability for Duffy-null, P(sweep) > 0.999.

Figure S6. 4 Results of leave-one-out cross-validation of ABC joint estimation **A)** selection coefficient (*RMSE* = 0.0083, R^2=0.9785) and **B)** initial West African admixture contribution (*RMSE* = 0.0090, R^2=0.9985).

Figure S7. Pink circles indicate West African mean global ancestry after 20 generations versus selection coefficient for whole autosome (22 chromosome) simulations, using a uniform recombination rate within each chromosome. Green triangles represent mean weighted ancestry for chromosome 1 and chromosome 2, with chromosome 2 representing the 92% of the genome that segregates independently from chromosome 1, using a human genetic map for recombination rates. We performed 500 simulations for each model, and all simulations started with West African ancestry contribution of 0.65. The estimated slope and intercept for the two methods of simulating global ancestry are highly similar. ANCOVA results suggest there is a significant effect of selection coefficient on global ancestry: $F(1,997)=1.0519\times10^4$, $p < 2\times10^{-6}$, but there is no significant effect of recombination rate and simulation model on global ancestry estimate after controlling for selection coefficient: $F(1,997) = 1.6350\times10^{-1}$, $p = 0.686$.