Response to Kozlov *et al.*: Inaccurate estimation of biases in herbarium specimen data

Emily K. Meineke[1], Charles C. Davis[2], T. Jonathan Davies[3,4]

[1] Department of Entomology and Nematology, University of California, Davis

[2] Department of Organismic and Evolutionary Biology, Harvard University Herbaria

[3] Departments of Botany, Forest & Conservation Sciences, University of British Columbia,

Vancouver

[4] African Centre for DNA Barcoding, University of Johannesburg

*Emily K. Meineke, ekmeineke@ucdavis.edu, phone: (+1) 919-308-3267, One Shields Avenue,

Davis, CA 95616

Running title: Response to Kozlov *et al.*

Keywords: herbivore, species interactions, insect, herbivory, museum specimens, herbarium

Summary. Kozlov and colleagues[1] call into question the application of herbarium specimens to quantify historical patterns of herbivory[2–5]. It is already widely appreciated that collectors of herbarium specimens may tend to avoid insect damage, thus making herbivory estimates from herbarium specimens potentially down-biased[2]. However, Kozlov *et al.* additionally suggest that variation in sampling selectivity among collectors and curators may lead herbarium specimens to misrepresent patterns of herbivory in nature. The authors sought to quantify these biases by collecting and contrasting insect herbivory data across 17 plant species from herbarium versus standard field ecological sampling procedures, and then assessed the selection of these specimens by curators. They concluded that herbivory estimates from herbarium specimens are highly variable, rendering them an inaccurate representation of herbivory in nature. Our re-analysis of Kozlov *et al.*'s data suggests that, in contrast with their results, herbarium specimens indeed provide a useful record of herbivory as long as sample sizes are appropriate. In addition, we assert that by arguing that herbarium specimens are "distorting mirrors", Kozlov *et al.*'s conclusions fundamentally overstep their data, which narrowly assesses biases across species. Kozlov *et al.* argue that herbarium specimens are inaccurate data sources, but fail to characterize the specific circumstances under which assumed biases would apply. Thus, Kozlov *et al.*'s data do not support their main premise, and the authors extrapolate beyond the specific biases investigated in their study; we believe their contribution does a disservice to researchers interested in exploring the potential value of herbarium specimens for studying herbivory through time.

Main text. Kozlov *et al.* [1] assert that estimates of herbivory damage derived from herbarium specimens are unreliable. Assessing leaf area removed by insects to infer herbivory is a common

practice[6]. However, it is well established that insect damage is distributed unevenly; some plants

experience extensive damage, whilst others exhibit little or no signs of herbivory (e.g.[7]). It is

critical, therefore, to sample multiple individuals when estimating species means, especially

when assessing single plant species from a given locality (e.g.[8,9]). Kozlov *et al.* gathered too-few

and too-variable specimens (n=1 to n=10 for ecological samples, and n=3 to n=20 for herbarium

samples) to infer reliable estimates of herbivory. The pathologies resulting from such under-

sampling are evident in Fig. 1 presented by Kozlov *et al*. Herbivory estimates derived from

ecological versus herbarium sampling vary widely when inferred from only a few herbarium

samples, but converge when sample sizes are higher and comparable between sampling

strategies. This trend is observable when species are sorted by total sample size (Fig. 1). For

example, herbivory estimates for *Betula pubescens* and *Rubus idaeus*, the only focal species in

which sample size is high for both categories, align closely (Fig. 1; Appendix S1).

Using Kozlov *et al.*'s published data, we show that herbivory estimated from ecological and

herbarium specimens are highly correlated as long as species have more than 10 data points (Fig.

2) or if we simply weight by sample size (Table 1). By recreating Fig. 3 from Kozlov *et al*., we

demonstrate that herbivory estimates from these two sampling methodologies exhibit a broadly

linear relationship which is disrupted by a few plant species represented by small sample sizes

(Fig. 2). We re-test the hypothesis that herbivory measurements extracted using the two

methodologies are correlated with three sets of models (Table 1). First, we performed weighted

linear models using the *lm* function R[10] with mean herbivory measured on herbarium samples as

the independent variable (*x*) and mean herbivory measured using ecological metrics (*y*) as the

dependent variable. The data points were weighted by the total number of herbivory samples

Kozlov *et al.* collected for a given species (total n= n of herbarium samples + n of ecological

samples). Because these weights allow us to account for uneven sample sizes, we restricted our

analyses to arithmetic means, not least-squared means (which the authors claim are another way

to reduce effects of uneven sampling across species; see Appendix S1). The resulting models—

one of which contains a transformed response variable to improve model fit—display broad

agreement between measurements of herbivory derived ecological and herbarium sampling

procedures (Table 1, Models 1 and 2; $p= 0.02$, $r^2 = 0.27$ and $p<0.01$, $r^2 = 0.33$). If we restrict

Kozlov *et al.*'s data to include only species for which n≥10—still a small sample size for noisy

ecological data—the relationship between these arithmetic means calculated from herbarium and

ecological sampling is even stronger in a simple linear regression (Table 1, Models 3 and 5; $p=$

0.01, $r^2 = 0.39$ and $p= 0.01$, $r^2 = 0.41$), and is nearly significant in a simple linear regression

focusing on the marginal means Kozlov *et al*. report (Table 1, Models 4 and 6; Fig. 2b: $p= 0.06$,

$r^2 = 0.21$, and $p= 0.05$, $r^2 = 0.23$). Finally, we ran Deming regressions including plant species

with n≥10. The Deming modeling procedure is arguably more appropriate for these data than

simple linear regression because it allows for errors in both the *x* and the *y* variables (in this case

herbivory measured from herbarium and ecological sampling procedures, which both carry

error). Both of these regressions were "significant" in that 95% confidence intervals did not

overlap zero (Table 1, Models 7 and 8; fits are plotted in Fig S1). We emphasize the plant

species not included in Models 3-8 with n<10 are not statistical outliers in terms of herbivory –

e.g., points of high leverage that might be identified using Cook's distance or other outlier

detection approaches. Rather, they are instances in which sample size is low and thus herbivory

estimates predictably do not match between ecological and herbarium samples because the data

are noisy. In summary, the patterns in the data (Figs. 2 and S1) and the statistical models (Table

1) illustrate that herbarium and ecologically derived data are highly correlated when sample sizes

are adequate. Therefore, while Kozlov *et al.*'s data provide evidence that estimates of herbivory

on herbarium specimens are down-biased, as has been shown previously[2], differences between

species' estimates are highly consistent when sampling effort is sufficient, contrary to the

author's published conclusion.

It might be argued that, like Kozlov et al.'s data, herbarium specimens provide few data points if

the sampling unit is a given species in a given place at a given time. However, the ecological

concept of sample size relates to the independent variable whose effects are being assessed. For

example, while just one herbarium specimen from a specific place-species-time combination

might be available, if we aim to assess the effects of temperature on herbivory, we can use

herbarium specimens from many place-species-time locations with matched temperature data

form that location and time period. Further, if the goal is to evaluate the general relationship

between temperature on herbivory, we can use hierarchical or mixed effects models that allow us

to borrow statistical strength from our better-informed estimates (e.g., species or sites with

higher sampling density). Thus, in contrast to most ecological field studies, herbarium specimens

offer the opportunity to expand sample sizes vastly across multiple axes, and the effects of

individual sites can be captured independently in appropriately fit statistical models that include

relevant co-factors. Kozlov *et al*. do not describe any systemic such biases for fellow ecologists

to beware of in these herbarium data; however, should they have done so, this information could

have been used to usefully inform future model fitting.

Kozlov *et al*. suggest that herbivory estimated from herbarium specimens produces inaccuracies

that preclude comparing these metrics among species. However, we know of few studies that use

herbarium specimens for this purpose; the focus has instead been on quantifying variability in

herbivory over years and space *within* species, and inferring general trends across species. Even

if there are idiosyncratic species differences between observed and estimated herbivory from

herbarium data, such that herbarium records tend to underestimate herbivory more in some

species compared to others, as Kozlov *et al.* suggest, specimens may still serve as an accurate

source data on herbivory variation across axes relevant to global change (years, latitude,

temperature). For instance, in a comparison of seasonal flowering sensitivity to temperature

between observational and herbarium data, Davis *et al.*[11] demonstrated that while phenological

sensitivity measurements from herbarium specimens did not predict phenological sensitivity

measured in the field, the effects of seasonal temperature on flowering still aligned. Therefore,

differences in the precision with which herbarium data capture herbivory (or phenology) between

species does not preclude their use to infer changes in herbivory (or phenology) across

environmental gradients or over time. Thus, Kozlov's criticism, which addresses bias when

comparing estimates among plant species, not over time, is not relevant to how most such data

are used, including in our study[2] examining changes in herbivory in the eastern United States

over time.

While we have shown that the conclusions drawn by Kozlov *et al*. from the comparison of

herbivory estimates on ecological versus herbarium samples are mistaken, better quantifying the

true relationships between sampling approaches would be a useful contribution to the literature

given the growing number of publications in this area, including some of our own work

(e.g.[2,3,12]). In a second analysis, Kozlov *et al*. provide one step toward this goal, polling 17

herbarium curators to assess their biases against the accession of specimens with insect damage.

Here they show that herbarium curators are differentially biased against damaged specimens, and

thus conclude that herbarium specimens cannot provide reliable estimates of herbivory in nature.

We do not dispute that curators likely differ in their selectivity of specimens for vouchering;

however, we again disagree the conclusions drawn by Kozlov *et al*.

If curators are inconsistent in their tolerance for damage, and assuming curator bias does not

itself vary systematically across the axis of interest, individuals' preferences would simply add a

further source of noise in the data. Although not a source of bias, added noise in herbivory

estimates could be problematic in statistical modelling fitting given the natural variability in

herbivory damage across individual plants[13]. New machine learning methods provide an exciting

path forward by allowing herbivory data to be collected rapidly from greater numbers of

specimens[14], decreasing the noise-to-signal ratio in datasets. We note that, given the inherent

differences in collector and curator preferences, it is perhaps more impressive that it is still

possible to extract expected patterns in herbivory across space and time from herbarium

specimens[3]. However, Kozlov *et al*.'s observation of variability in tolerance to herbivory

exhibited by curators does raise an important point regarding differences in cultural practices

among herbaria. Though, we are unaware of any studies that explicitly compare herbivory rates

across regions, Kozlov *et al*.'*s* analysis suggests that if ecologists want to use herbaria for this

purpose, collectors should be polled to ensure that collector biases do not vary systematically

between locations.

Finally, we suggest that Kozlov *et al.*'s conclusion that herbarium specimens may serve as

"distorting mirrors" is unjustified because it reaches beyond the limited quantifications of bias in

their study. Like any historical reconstruction from fossils[15] or modern specimens[11], it is

important to identify potential sources of bias in the data. However, a bias or source of noise

along any one axes does not preclude the value of these data for exploring variation across other

axes. Instead of explaining how the biases they detected across species and curators might affect

herbarium-based data analyses, the authors instead appear suggest that these data are too

inaccurate to be of use. Kozlov *et al*. do not provide any additional evidence to support this

statement. We here draw parallels to the use of paleontological or archeological data. If the

detection of taphonomic biases in fossil collections had halted the fields of archeology and

paleontology, the vast knowledge we have gleaned from fossil data would never have been

collected. We suggest that ecologists approach biases in modern collections data in the same way

through the robust assessments of particular biases that are most likely to affect conclusions,

rather than a wholesale rejection of such data.

Conclusion: While biases in herbarium data are well recognized[16,17], there remains tremendous

value in these historical specimens[3,18]. Our re-analysis of Kozlov *et al.*'s data reveals that

herbarium specimens provide a useful record of historical herbivory, contrary to the authors

published conclusions. We fully support efforts to validate herbivory estimates derived from

herbarium samples with gold standard ecological sampling but urge the use of power analyses

and/or rarefaction curves to ensure samples sizes sufficient to yield reliable estimates. We also

note that while largescale efforts to better quantify collector bias in specimen data across species

are useful, we are often more interested in temporal and spatial trends within species to capture

the fingerprint of global change (e.g.[19]). We hope that Kozlov *et al.*'s disproportionate and at

times unjustified criticism of herbarium specimens as a source of species interactions data will

not discourage ecologists from leveraging the information captured within these specimens,

which are increasingly being mobilized online and represent a rich resource for inferring past

ecologies[20].

References

1. Kozlov, M. V. *et al.* Biases in estimation of insect herbivory from herbarium specimens. *Sci.*

Rep. **10**, 12298 (2020).

2. Meineke, E. K., Classen, A. T., Sanders, N. J. & Jonathan Davies, T. Herbarium specimens

reveal increasing herbivory over the past century. *J. Ecol.* **107**, 105–117 (2019).

3. Meineke, E. K., Davis, C. C. & Davies, T. J. The unrealized potential of herbaria for global

change biology. *Ecol. Monogr.* **88**, 505–525 (2018).

4. Beaulieu, C., Lavoie, C. & Proulx, R. Bookkeeping of insect herbivory trends in herbarium

specimens of purple loosestrife (Lythrum salicaria). *Philos. Trans. R. Soc. B* **374**, 20170398

(2019).

5. Schilthuizen, M. *et al.* Incorporation of an invasive plant into a native insect herbivore food

web. *PeerJ* **4**, e1954 (2016).

6. Johnson, M. T., Bertrand, J. A. & Turcotte, M. M. Precision and accuracy in quantifying

herbivory. *Ecol. Entomol.* **41**, 112–121 (2016).

7. Atwater, D. Z., Butler, J. L. & Ellison, A. M. Spatial distribution and impacts of moth

herbivory on northern pitcher plants. *Northeast. Nat.* **13**, 43–56 (2006).

8. Kent, D. R. *et al.* Weak latitudinal gradients in insect herbivory for dominant rangeland

grasses of North America. *Ecol. Evol.* (2020).

9. Alexandre, N. M. *et al.* Habitat preference of an herbivore shapes the habitat distribution of its

host plant. *Ecosphere* **9**, e02372 (2018).

10.   R Core Team. *R: A language and environment for statistical computing.* (R Foundation

for Statistical Computing, Vienna, Austria., 2019).

11.   Davis, C. C., Willis, C. G., Connolly, B., Kelly, C. & Ellison, A. M. Herbarium records

are reliable sources of phenological change driven by climate and provide novel insights into

species' phenological cueing mechanisms. *Am. J. Bot.* **102**, 1599–1609 (2015).

12.   Meineke, E. K. & Davies, T. J. Museum specimens provide novel insights into changing

plant–herbivore interactions. *Philos. Trans. R. Soc. B Biol. Sci.* **374**, 20170393 (2018).

13.   Turcotte, M. M., Davies, T. J., Thomsen, C. J. & Johnson, M. T. Macroecological and

macroevolutionary patterns of leaf herbivory across vascular plants. *Proc. R. Soc. B Biol. Sci.*

281, 20140555 (2014).

14.   Meineke, E. K., Tomasi, C., Yuan, S. & Pryer, K. M. Applying Machine Learning to

Investigate Long Term Insect-Plant Interactions Preserved on Digitized Herbarium

Specimens. *bioRxiv* 790899 (2019).

15.   Darroch, S. A. F., Fraser, D. & Casey, M. M. The preservation potential of terrestrial

biogeographic patterns. *Proc. R. Soc. B Biol. Sci.* **288**, 20202927 (2021).

16.   Daru, B. H. *et al.* Widespread sampling biases in herbaria revealed from large-scale

digitization. *New Phytol.* **217**, 939–955 (2018).

17.   Meyer, C., Weigelt, P. & Kreft, H. Multidimensional biases, gaps and uncertainties in

global plant occurrence information. *Ecol. Lett.* **19**, 992–1006 (2016).

18.   Meineke, E. K., Davies, T. J., Daru, B. H. & Davis, C. C. Biological collections for

understanding biodiversity in the Anthropocene. *Philos. Trans. R. Soc. B Biol. Sci.* **374**,

20170386 (2019).

19.   Wolkovich, E. M. *et al.* Temperature-dependent shifts in phenology contribute to the

success of exotic species with climate change. *Am. J. Bot.* **100**, 1407–1421 (2013).

20.   Hedrick, B. P. *et al.* Digitization and the future of natural history collections. *BioScience*

70, 243–251 (2020).

Author contributions

E.K.M. and T.J.D. contributed to statistical analyses. E.K.M., T.J.D., and C.C.D. wrote the
paper.

Additional information

The authors declare no competing interests.

237
Figure 1. Effects of sample size on herbivory estimates. a) Raw data describing leaf area lost to

insect herbivores detected using different sampling methods in Kozlov *et al*.[5,6]. Note that sample

sizes differ slightly from those reported in Kozlov *et al*. because of a technical error that caused

minor data misreporting in the original paper's supplement and in the main text figures. b) The

same plot with least-squared (marginal) means. This figure includes the exact data from Fig. 1 in

Kozlov *et al.*, wherein they report marginal means for which calculation was not included in the

original manuscript. For more discussion on why we chose to include panel b), please see

Appendix S1.

Figure 2. The relationship between herbivory estimated from ecological vs. herbarium specimens. Each data point represents a species, and numbers indicate total sample sizes (ecological + herbarium). For a), the model was derived from the *lm* function in r[10] and represents the simple linear relationship between mean leaf area loss per species measured in

herbarium specimens (x) and ecological specimens (y). The model fitted to species with ≥ 10

specimens collected (species in black) is shown in blue (±SE in grey). Associated model

parameters are presented in the lower righthand corner ($F_{1,11}= 8.56$, $p= 0.01$). The r^2 value

reported is adjusted r^2. In b), we show similar patterns in the marginal means calculated by

Kozlov et al. for the model that included species for which $n>10$ ($F_{1,11}= 4.27$, $p= 0.06$). In both

plots, the black line represents a 1:1 agreement between herbivory estimates derived from

herbarium and ecological samples. Thus, herbarium specimens represent down-biased estimates

of herbivory as expected, but these data are nevertheless linearly correlated with herbivory data

from ecological sampling of the sample species.

Table 1. Various models of the relationships between herbivory estimated using herbarium

and ecological sampling methodologies. In all models, herbivory estimates derived from

herbarium specimens is the independent variable (x) and herbivory estimates derived from

ecological specimens is the independent variable (y). a) As described in the main text, Model 1

has arithmetic means as the response variable, and the data points are weighted by total sample

size. Model 2 is identical to Model 1 except that herbivory estimates are treated as proportion

data, and the response variable was arcsine(sqrt) transformed to improve model fit. Models 3 and

4 are simple linear regressions, one with arithmetic and one of least-squared mean herbivory

estimates as the response variables. Models 5 and 6 are transformed as described above. b)

Deming regressions of species with n≥10 data points. Listed values represent 95% confidence

intervals. In Deming regression, slope estimates are considered "significant" if the confidence

limits do not cross zero, as is the case in our models.

a) Linear regressions

Model number	Model type	Response	Estimate	F-value	DF	p	Adjusted-R^2
1	Weighted linear regression	Arithmetic means	0.89	6.83	1, 15	**0.02***	0.27
2	Weighted linear regression	Arcsine(sqrt) transformed Arithmetic means	1.84	8.89	1, 15	**<0.01****	0.33
3	Simple linear regression with n≥10	Arithmetic means	1.13	8.56	1, 11	**0.01***	0.39
4	Simple linear regression with n≥10	Least-squared means	0.83	4.27	1, 11	0.06 .	0.21
5	Simple linear regression	Arcsine(sqrt) transformed Arithmetic	2.27	9.36	1, 11	**0.01***	0.41

	with n≥10	means					
6	Simple linear regression with n≥10	Arcsine(sqrt) transformed Least-squared means	1.66	4.65	1, 11	0.05 .	0.23

278   **b) Deming regressions**

Model number	Model type	Response	Slope Estimate	Lower CI	Upper CI	Pearson's *r*
7	Deming	Arithmetic means	2.16	**0.98**	**3.76**	0.661
8	Deming	Least-squared means	2.24	**0.68**	**4.43**	0.529

Appendix S1: A discussion of least-squared vs. arithmetic means

In Fig. 1, and where appropriate in statistical analyses, we report results from both arithmetic

means (Fig. 1a) and the marginal (least-squared) means reported in Kozlov *et al.*'s manuscript

(Fig. 1b). The model structure for the calculation of the least-squared means was not described in

the original manuscript, and in subsequent conversations, the authors were not able to explain the

reason for including these least-squared means (rather than arithmetic means) to us beyond that

they account for "uneven sample sizes". Least-squared means do not necessarily correct for

uneven sample sizes, but they can help account for uneven sampling across a particular variable.

Given no information, we initially assumed Kozlov and colleagues collected herbivory

measurements from multiple sites, and that perhaps these means may have been derived to

correct for the uneven application of sampling methods across these sites. However, the vast

majority of their species (14/17) were only collected at one site.

        We suggest our weighted regression model on the arithmetic means provides the best and

most transparent representation of the data. Nonetheless, we provide analyses based on the

authors' reported least-squared means alongside the arithmetic means derived from their raw

data. In taking this approach, we aimed to be as inclusive as possible of the authors preferences

for their least-squared means estimates, which we struggled to replicate.

Figure S1. Deming regression fits described in Table 1, Models 7-8.

The 0.95–confidence bounds are calculated with the bootstrap(quantile) method.