Vertical transmission at the pathogen-symbiont interface: *Serratia symbiotica* and aphids

Julie Perreau[a,b], Devki J. Patel[a], Hanna Anderson[a], Gerald P. Maeda[a], Katherine M. Elston[b],

Jeffrey E. Barrick[b], Nancy A. Moran[a]

[a] Department of Integrative Biology, University of Texas at Austin, Austin, TX 78712

[b] Department of Molecular Biosciences, University of Texas at Austin, Austin, TX 78712

Running title: *Serratia symbiotica* at the pathogen-symbiont interface

Address correspondence to Julie Perreau, jperreau@utexas.edu

<u>Word count</u>

Abstract: 236

Text (excluding the references, table footnotes, and figure legends): 5,098

Abstract

Many insects possess beneficial bacterial symbionts that occupy specialized host cells and are maternally transmitted. As a consequence of their host-restricted lifestyle, these symbionts often possess reduced genomes and cannot be cultured outside hosts, limiting their study. The bacterial species *Serratia symbiotica* was originally described by noncultured strains that live as mutualistic symbionts of aphids and are vertically transmitted through transovarial endocytosis within the mother's body. More recently, culturable strains of *S. symbiotica* were discovered that retain a larger set of ancestral *Serratia* genes, are gut pathogens in aphid hosts, and are principally transmitted via a fecal-oral route. We find that these culturable strains, when injected into pea aphids, replicate in the hemolymph and are pathogenic. Unexpectedly, they are also capable of maternal transmission via transovarial endocytosis: using GFP-tagged strains, we observe that pathogenic *S. symbiotica*, but not *Escherichia coli*, are endocytosed into early embryos. Furthermore, pathogenic *S. symbiotica* strains are compartmentalized into specialized aphid cells in a similar fashion to mutualistic *S. symbiotica* strains during later stages of embryonic development. Thus, cultured, pathogenic strains of *S. symbiotica* have the latent capacity to transition to lifestyles as mutualistic symbionts of aphid hosts. This capacity is blocked by pathogenicity: their hosts die before infected progeny are born. To transition into stably inherited symbionts, culturable *S. symbiotica* strains may need to adapt to regulate their titer, limit their pathogenicity, and/or provide benefits to aphids that outweigh their cost.

Keywords: Secondary symbiont, *Buchnera aphidicola*, bacteriocyte, sheath cells

Importance

Insects have evolved various mechanisms to reliably transmit their beneficial bacterial symbionts to the next generation. Sap-sucking insects, including aphids, transmit symbionts by endocytosis of the symbiont into cells of the early embryo within the mother's body. Experimental studies of this process are hampered by the inability to culture or genetically manipulate host-restricted, symbiotic bacteria. *Serratia symbiotica* is a bacterial species that includes strains ranging from obligate, heritable symbionts to culturable gut pathogens. We demonstrate that culturable *S. symbiotica* strains, that are aphid gut pathogens, can be maternally transmitted by endocytosis. Cultured *S. symbiotica* therefore possess a latent capacity for evolving a host-restricted lifestyle and can be used to understand the transition from pathogenicity to beneficial symbiosis.

Introduction

Host-associated bacteria can be placed on a continuum ranging from parasitic to mutualistic. Mutualistic symbionts often arise from pathogenic ancestors and rarely revert back to pathogenicity (1). This one-way transition is expected when vertical transmission, usually from mother to offspring, replaces horizontal transmission as the dominant route of new infection, causing symbionts to benefit from host reproduction and thereby to face strong selection for avirulence (2–4). In insects, the reliable vertical transmission of mutualistic symbionts can be accomplished by mechanisms that are external, including the placement of symbionts or symbiont-containing capsules on the egg surface, or internal, via transovarial transmission (5, 6).

Transovarial transmission, the transfer of maternal symbionts to eggs or embryos within the mother's body, is common in obligate symbioses where symbionts occupy special host cells (bacteriocytes) within the body cavity (7). Individual bacteria may be transferred, as in aphids, cicadas, leafhoppers, cockroaches, and bedbugs, or entire bacteriocytes may be transferred, as in whiteflies (7–10). In ancient symbioses with exclusively maternal transmission, hosts appear to control transmission, as the symbionts involved often possess reduced genomes devoid of pathogenicity factors that would allow them to invade host cells (11). However, host mechanisms for transmission may depend on bacterial factors for inter-partner recognition, and thereby limit which bacterial species or strains can make the transition from pathogenicity to commensal or mutualistic symbioses.

Aphids (Hemiptera: Sternorrhyncha: Aphidoidea) are a clade of roughly 5,000 species that feed exclusively on nutrient-poor plant sap and depend on the primary bacterial symbiont *Buchnera aphidicola* for biosynthesis of essential amino acids missing from their diet (12–16). *B. aphidicola* have been transovarially transmitted in aphids for over 160 million years (17).

They occupy bacteriocytes, possess highly reduced genomes, and cannot persist outside of their

hosts (11, 18). In addition to *B. aphidicola*, many aphids also host secondary symbionts, such as

Serratia symbiotica, Candidatus Hamiltonella defensa, *Candidatus* Regiella insecticola, and

Candidatus Fukatsuia symbiotica (19–22). Due to their more recent associations with aphids,

secondary symbionts provide an opportunity to understand the transition to host-restricted

lifestyles.

S. symbiotica strains have evolved diverse associations with aphids. They range from

pathogens and facultative mutualists to obligate mutualists that co-reside with *B. aphidicola* (23–

32). The genome sizes and gene contents of *S. symbiotica* strains reflect this variation in lifestyle,

ranging from larger genomes similar to those of free-living *Serratia* (31, 33), to highly reduced

genomes similar to those of *B. aphidicola* and other obligate symbionts (24, 26, 27, 32). The first

descriptions of *S. symbiotica*, initially named Pea Aphid Secondary Symbiont (PASS) or the "R-

type" symbiont, were of strains that occupied secondary bacteriocytes, sheath cells, and

hemolymph (34, 19). Unlike *B. aphidicola,* these *S. symbiotica* strains are not required by their

hosts; they provide secondary benefits, such as protection against heat stress (23, 35), and against

parasitoid wasps (36). However, similar to *B. aphidicola,* they are host-restricted and vertically

transmitted by a transovarial route (8, 37, 38). A detailed study showed that a mutualistic *S.*

symbiotica strain, present in the hemolymph, migrates to early embryos and is endocytosed with

B. aphidicola into the syncytium, i.e., a specialized, multinucleated cell of the early embryo (8).

In the pea aphid, *B. aphidicola* and *S. symbiotica* are later segregated into distinct bacteriocytes.

Recently, strains of pathogenic *S. symbiotica* have been discovered living in the guts of

Aphis species collected in Belgium and Tunisia (25, 29–31). These strains are hypothesized to

resemble ancestors of facultative and co-obligate *S. symbiotica* strains (39, 40). In contrast to

previously studied *S. symbiotica* strains, their primary transmission route appears to be

horizontal, through honeydew (feces) and host plant phloem (39, 40). Also, in contrast to

previously studied strains, these *S. symbiotica* can be cultured axenically. *S. symbiotica* CWBI-

2.3^T, from the black bean aphid (*Aphis fabae*) (25), retains a larger gene set and larger genome

(3.58 Mb) (33) than do facultative *S. symbiotica* strains Tucson and IS (2.78 and 2.82 Mb,

respectively) (41, 42).

In this work, we investigate whether cultured *S. symbiotica* strains are capable of vertical

transmission similar to facultative or obligate symbionts. We isolated a new *S. symbiotica* strain,

HB1, that shares many features with CWBI-2.3^T but is notably less pathogenic. We examined the

capacity of each of these strains to colonize hemolymph of the pea aphid (*Acyrthosiphon pisum*)

and access embryos through the transovarial route described for *B. aphidicola* (8). Both strains

are endocytosed into embryos, but pathogenic strains kill their hosts before the birth of offspring

infected by the transovarial route. Using GFP-tagged strains, we addressed whether transovarial

transmission is open to any bacterial cell that comes into contact with the embryo, or whether

this process involves specific partner recognition. We find that *Escherichia coli* cannot colonize

embryos, despite achieving a high titer within hosts. Thus, the endocytosis step required for

transovarial transmission limits the taxonomic range of bacteria that can readily evolve to

become aphid symbionts.

Results

Pathogenic *S. symbiotica* form a distinct group closely related to mutualistic strains

We examined the evolutionary relationships of all *S. symbiotica* with publicly available complete

genome sequences. To this list, we added a recently isolated and sequenced strain, designated *S.*

symbiotica HB1, from the melon aphid (*Aphis gossypii*). Our phylogenetic analysis was based on

176 shared orthologous genes and was rooted with outgroups that included other *Serratia* and

more distant *Enterobacterales* species (Figure 1, Figure S1, Table S1). *S. symbiotica* is split into

two clades, as previously reported (43). Clade A is comprised of strains that act as pathogens,

mutualists, or co-obligate symbionts in aphids from across the family Aphididae, while clade B

is composed of strains that live only as co-obligate symbionts in aphids of the subfamily

Lachninae. Clade A strains possess a range of genome sizes (1.54–3.58 Mb), GC content (47.9–

52.5%), host species, and lifestyles. Within clade A, the cultured, gut pathogen strains (CWBI-

2.3^T, HB1, Apa8A1, and 24.1) form a clade closely related to vertically transmitted mutualist

strains (Tucson, IS, MCAR-56S, AURT-53S). Although average nucleotide identities are high

(96.0–98.6%) across these strains (Table S2), the gut pathogen strains retain larger genomes

(3.09–3.58 Mb) and more *Serratia*-specific marker genes than non-pathogenic, maternally

transmitted *S. symbiotica* (0.65–2.82 Mb) (Figure 1, Table S1). Together, these observations

suggest that both lifestyles have recently emerged from a common, presumably pathogenic

ancestor.

Cultured *S. symbiotica* strains are pathogenic when injected into pea aphid hemolymph

Based on previous studies, *S. symbiotica* CWBI-2.3^T acts as a gut pathogen in its original host,

the black bean aphid (40), but appears to be less pathogenic in the gut of pea aphids (44). In both

hosts, *S. symbiotica* CWBI-2.3^T appears to be restricted to the gut and is not observed infecting

hemolymph. To determine whether *S. symbiotica* CWBI-2.3^T and HB1 can persist and act as

pathogens in the hemolymph of pea aphids, and to determine if these strains are capable of

vertical transmission to offspring, we injected fourth instar pea aphids with tagged strains CWBI-

2.3^T-GFP or HB1-GFP. Then, we tracked aphid survival, vertical transmission rates, and

bacterial titer over time. For comparison, we simultaneously performed injections of *S.*

marcescens Db11*,* a known insect pathogen (45), injections of hemolymph from pea aphids

infected with facultative, vertically transmitted *S. symbiotica* Tucson (41), and injections of

buffer as a negative control. We found that *S. symbiotica* CWBI-2.3^T-GFP and HB1-GFP act as

pathogens in pea aphid hemolymph. The survival rates of aphids injected with *S. symbiotica*

CWBI-2.3^T-GFP, *S. symbiotica* HB1-GFP, or *S. marcescens* Db11 were much lower than those

of aphids injected with buffer or with *S. symbiotica* Tucson ($P < 0.001$, Cox Proportional

Hazards Model) (Figure 2A).

Typically, *S. symbiotica* CWBI-2.3^T infects by a fecal-oral route or from plants (40). To

determine if *S. symbiotica* CWBI-2.3^T-GFP and HB1-GFP can be transmitted to offspring via the

transovarial route, as for mutualistic symbiotic strains, we collected newborn offspring of

surviving aphids every 24 hours and screened them for *S. symbiotica* by plating aphids and

noting presence or absence of colony-forming units (CFU). As transovarial transmission of

symbionts occurs early in embryonic development, there is a delay between injection and the

birth of infected offspring (34). To determine when after injection we should begin to observe

offspring infected by transovarial transmission, we injected mutualistic *S. symbiotica* Tucson and

monitored its transmission by sampling offspring and using PCR to screen for the presence of *S.*

symbiotica. Transmission of *S. symbiotica* Tucson was identified in newborn offspring starting at

10 days post-injection (DPI) (Figure 2B). The proportion of infected offspring per mother

increased over time, reaching 100% for most mothers by 15 DPI (Figure 2B). In contrast to

aphids injected with *S. symbiotica* Tucson, aphids injected with *S. symbiotica* CWBI-2.3^T did not

survive past 9 DPI (Figure 2A), and aphids injected with *S. symbiotica* HB1 showed reduced

fecundity and produced only a few, non-viable offspring after 9 DPI. Therefore, successful

vertical transmission of these strains appears to be limited by their pathogenicity.

We hypothesized that the virulence of *S. symbiotica* CWBI-2.3^{T}-GFP and HB1-GFP was

related to their titer in hemolymph. To test this hypothesis, aphids were sacrificed every 24 hours

to measure the CFU per aphid. *S. symbiotica* CWBI-2.3^{T}-GFP and HB1-GFP grow exponentially

within aphids, plateauing at ~10^{9} CFU per aphid by 5-6 DPI (Figure 2C). The significant

difference in the growth rate of CWBI-2.3^{T} and HB1 at 3 and 4 DPI may explain the difference

in aphid survival rate across these treatment groups (Figure 2A). As *S. symbiotica* Tucson is not

culturable, qPCR was used to compare titers of CWBI-2.3^{T}-GFP and HB1-GFP to those reached

by Tucson at 5 DPI, and to the titers at which other mutualist *S. symbiotica* strains persist in pea

aphids reared in the lab (Figure 2D). Relative titer was calculated as the copy number of a single-

copy *Serratia* gene (*dnaK*) normalized by a single-copy pea aphid gene (*ef1a*). Both *S.*

symbiotica CWBI-2.3^{T}-GFP and HB1-GFP reach higher titers than *S. symbiotica* Tucson at 5

DPI (Figure 2D). The titer of *S. symbiotica* Tucson at 5 DPI is comparable to the steady-state

titer of *S. symbiotica* strains maintained in naturally infected, clonal pea aphids established as

laboratory lines (Figure 2D).

Cultured *S. symbiotica* strains are capable of colonizing *Ac pisum* embryos

Intergenerational transmission of *S. symbiotica* CWBI-2.3^{T} and HB1 was prevented because the

pathogenicity of these strains killed mothers before any infected embryos could be born. In order

to determine whether they were, nevertheless, capable of transovarial transmission, we injected

aphids with *S. symbiotica* CWBI-2.3^{T}, dissected ovarioles at 7 DPI, and used fluorescence *in-situ*

hybridization to visualize its infection pattern relative to the primary symbiont *B. aphidicola*

during embryonic development. To compare these results to the transmission pattern of a

mutualist strain, we injected aphids with hemolymph from pea aphids infected with *S. symbiotica*

Tucson (Figure 3A-D).

Embryos injected with *S. symbiotica* Tucson display regular growth and development,

reaching 400 μm in length by the time of katatrepsis (37) (Figure 3A). As previously described,

B. aphidicola and mutualistic *S. symbiotica* are transmitted to the syncytium of stage 7 blastula

from hemolymph via an endocytic process (8, 37, 46). *S. symbiotica* Tucson appears to be unable

to infect embryos at later developmental stages, as embryos that were beyond stage 7 at the time

of injection lack *S. symbiotica* (Figure 3B). This observation is supported by the absence of *S.*

symbiotica in offspring born the first 10 days after injection (Figure 2B) (34). Those embryos

exposed to *S. symbiotica* Tucson at stage 7 possess *S. symbiotica* in sheath cells, but Tucson does

not invade primary bacteriocytes that contain *B. aphidicola* (Figure 3C)*. The endocytosis of *S.*

symbiotica Tucson into stage 7 embryos occurs at the posterior end of the embryo (Figure 3D),

as previously described for the closely related strain *S. symbiotica* IS (8).

Despite its pathogenicity, the infection pattern of CWBI-2.3^T is similar to that of the non-

pathogenic Tucson strain (Figures 3E, 3F). Cells of CWBI-2.3^T are attached to the embryonic

surface but do not infect embryos that are beyond stage 7 (Figure 3G). Following infection,

CWBI-2.3^T is packaged similarly to the Tucson strain; both are sorted into sheath cells and

cannot colonize the primary bacteriocytes that house *B. aphidicola* (Figures 3C, 3H). CWBI-2.3^T

is endocytosed into early embryos with *B. aphidicola* (Figure 3I), as is *S. symbiotica* HB1

(Movie S1). Remarkably, at 7 DPI, CWBI-2.3^T greatly outnumbers *B. aphidicola* in the syncytial

cell (Figure 3I). In contrast to *S. symbiotica* Tucson, infection with cultured *S. symbiotica* CWBI-

2.3^T stunts embryonic growth, though it does not prevent progression through characteristic

developmental stages (Figures 3E, 3F).

Transovarial transmission is a specific capability of *S. symbiotica* strains

To determine if endocytosis is selective at the level of bacterial species, we tested the

transmission capability of *E. coli* K-12 strain BW25113. *E. coli* is related to *B. aphidicola, S.*

symbiotica, and several other mutualistic symbionts of aphids which are all within

Enterobacterales (20). We chose strain BW25113 because it can infect the gut and hemolymph

of pea aphids and kills aphids a few days post infection (47). We created the tagged *E. coli* strain

BW25113-GFP and injected it into fourth instar aphids as described above. For comparison, we

injected *S. symbiotica* CWBI-2.3^T-GFP into a separate set of fourth instar aphids. *E. coli*

BW25113-GFP forms a robust infection in pea aphid hemolymph, reaching comparable titers to

S. symbiotica CWBI-2.3^T-GFP at 5 DPI (Figure 4A). We dissected single ovarioles from ten

aphids in each treatment group at 5 DPI and observed early embryos to determine infection

status. Using this approach, *S. symbiotica* CWBI-2.3^T-GFP could be seen infecting early

embryos (Figures 4B, 4C, Movies S2, S3). In contrast, *E. coli* BW25113-GFP attaches to the

embryonic surface, sometimes coating the entire exterior of the embryo, but was never observed

endocytosing into embryos (Figures 4D, 4E, Movies S4, S5).

Discussion

Transovarial transmission is a key feature of many insect-bacterial symbioses wherein bacteria

provide a benefit to their host. This transmission route is linked to irreversible bacterial

transitions from pathogenicity to mutualism (2). However, many relationships that rely on

transovarial transmission are ancient, and their early stages cannot be experimentally

recapitulated, leaving unanswered if and how pathogenic bacteria access this transmission route.

Focusing on secondary symbionts may be more useful for understanding these early transitions,

as some secondary symbionts or their close relatives are culturable, genetically tractable, and can

be removed from or introduced to hosts without dramatically compromising host fitness (48–50).

For example, *Sodalis praecaptivus,* a close relative to *Sodalis* species found as host-restricted

symbionts across diverse insects, uses quorum sensing to attenuate virulence and gain access to

vertical transmission in a non-native host, the tsetse fly (51). The bacterial species *S. symbiotica*

was first known as a vertically transmitted mutualist, but pathogenic strains were subsequently

cultured from aphids collected in Europe and Africa (25, 29–31), and, in this study, in North

America. These strains have provided a new opportunity to dissect the early steps involved in the

transition to a host-restricted lifestyle. Knowing that vertical transmission is a key to this

transition, we aimed to determine whether culturable, pathogenic *S. symbiotica* could access this

pathway and what, if any, limitations are faced by *S. symbiotica* in this transition.

Cultured *S. symbiotica* strains are close relatives to non-culturable, vertically transmitted

mutualists. However, several lines of evidence suggest these strains lack a history of maternal

transmission in aphids. For one, persistent vertical transmission generally leads to the irreversible

loss of genes such that they can no longer access a free-living or pathogenic lifestyle (52).

Relative to vertically transmitted strains, CWBI-2.3^T, HB1, Apa8A1, and 24.1 are culturable,

maintain larger genomes, and possess more ancestral genes common to free-living *Serratia*

(Figure 1). Second, these strains do not appear to undergo vertical transmission in natural

infections. Following ingestion by black bean or pea aphids, *S. symbiotica* CWBI-2.3^T is not

subsequently detected in hemolymph or embryos, but is present in the gut and in honeydew,

suggesting that the dominant route of transmission for this strain is fecal-oral (29, 39, 44). We

injected CWBI-2.3^T and HB1 into hemolymph to determine if they are capable of vertical

transmission in pea aphids. Vertical transmission is theorized to be the primary force driving

permanent bacterial transitions from pathogenicity to mutualism, but to do so, vertical

transmission must precede mutualism (2). That pathogenic *S. symbiotica* CWBI-2.3^T and HB1

are endocytosed into the syncytial cell of early embryos along with *B. aphidicola* provides

empirical evidence for the precedence of vertical transmission in this system. Together, these

results suggest that the aphid gut has served as an access point for environmental or plant-

associated *Serratia* to infect aphids and that gut pathogenicity was an ancestral lifestyle for

strains that are now intracellular and mutualistic (29).

S. symbiotica is common to natural populations of pea aphids (34, 53), and previous 16S

rRNA gene surveys have identified *S. symbiotica* in aphid tribes from across the Aphidoidea (29,

54). However, pathogenic and mutualistic strains have near-identical 16S rRNA sequences, so it

is unclear how many cases represent *S. symbiotica* pathogens. To date, pathogenic strains have

been cultured only from *Aphis* species. However, *S. symbiotica* CWBI-2.3^T can horizontally

transmit across aphids feeding on the same plant (39), and can infect the guts of alternative aphid

species, including the pea aphid (44), suggesting that pathogenic *S. symbiotica* may be more

widespread across aphid genera in nature. The global distribution of *Aphis*-associated strains,

along with their ability to transmit using the same transovarial route as *B. aphidicola,* suggests

that related gut-associated strains may serve as a source pool for the evolution of commensal or

mutualistic strains. Mutualism may have arisen several times independently in *S. symbiotica,*

and, along with the subsequent horizontal transfer, would contribute to the phylogenetic

discordance between facultative symbionts and their aphid hosts (54). The acquisition and

replacement of secondary symbionts has occurred during the evolution of many ancient insect-

microbe symbioses and may help hosts to escape the "evolutionary rabbit hole" of dependence

on a primary symbiont that has become an ineffective mutualist due to genome decay (55, 56).

Transovarial transmission in aphids occurs by bacterial endocytosis into the syncytial cell of

early embryos (8). What host and bacterial factors are involved in this pathway are unclear, but

insights may be gained from other systems in which hosts are genetically tractable. In

Drosophila, knockout of yolk proteins or the Yolkless receptor results in reduced localization to

and/or endocytosis of *Spiroplasma* in embryos, suggesting that the vitellogenin pathway is

involved in transovarial transmission (57). While parthenogenetic aphids do not undergo

vitellogenesis or produce visible yolk, it is possible that similar receptor-mediated processes are

used for *B. aphidicola* and *S. symbiotica* transmission, and that specific bacterial ligands are

required. If this is the case, *S. symbiotica* strains that normally live in the aphid gut appear to

possess the requisite molecular determinants, as they display an innate potential for endocytosis

into embryos. Furthermore, the inability of *E. coli* BW25113 to endocytose into the syncytial cell

of embryos suggests that the endocytic step of transovarial transmission contributes to selectivity

in this system. While many bacterial taxa occasionally infect pea aphids (30, 58), few are found

as long-term mutualists. The primary symbiont, *B. aphidicola*, is stably maintained in most aphid

lineages, though rare replacements exist (e.g., (59)). Additionally, few species are found as

secondary symbionts*,* and most are members of the *Enterobacterales*, including *S. symbiotica,*

Ca. H. defensa*, Ca.* R. insecticola*, Ca.* F. symbiotica*,* and *Arsenophonus,* and, less commonly

other bacterial groups, including *Wolbachia, Rickettsia*, and *Spiroplasma* (21, 22).

Aphids that are co-infected with *B. aphidicola* and mutualistic secondary symbionts possess

several known mechanisms that limit the competition between these bacteria. For one, hosts can

sort symbionts into distinct cell types, with *B. aphidicola* in primary bacteriocytes and *S.*

symbiotica relegated to secondary bacteriocytes and sheath cells (60). Despite its pathogenicity,

we observed that CWBI-2.3^T is not able to invade primary bacteriocytes with *B. aphidicola* and

is compartmentalized into sheath cells in a similar manner as mutualistic strains (Figure 3). Pea

aphid genotypes may vary in their ability to associate with secondary symbionts. In contrast to

the results obtained with *Acyrthosiphon pisum* LSR1 in our study, when the facultative, host-

restricted strain *S. symbiotica* IS was transferred to *Acyrthosiphon pisum* AIST, it showed a

disordered localization, invading primary bacteriocytes with *B. aphidicola* (61). In these cases, *S.*

symbiotica IS was trapped in primary bacteriocytes, unable to exocytose during transmission (8)*.*

The specific exocytosis of *B. aphidicola* also likely plays a role in limiting competition between

B. aphidicola and secondary symbionts across multiple generations.

The vertical transmission of CWBI-2.3^T and HB1 in pea aphids is limited by their virulence

in hemolymph, as infected aphids do not live long enough to birth offspring infected by

transovarial transmission. Both CWBI-2.3^T and HB1 are more pathogenic in hemolymph than

facultative *S. symbiotica* Tucson, but also notably far less pathogenic than *S. marcescens* Db11.

Possibly, adaptation to the gut selects for reduced *Serratia* virulence by allowing *Serratia* the

time to form a robust gut infection and transmit to other aphids, including offspring, via

honeydew (40). The virulence of CWBI-2.3^T and HB1 coincides with unregulated titer, as both

of these strains attain 100 to 1,000-fold higher titers relative to mutualistic *S. symbiotica* when

injected into hemolymph. This enormous difference in titer, and the constancy of the low titers

observed for the symbiotic strains, suggests that mutualistic *S. symbiotica* growth is regulated.

The regulation of virulence and titer is important in the establishment of vertical transmission.

Self-regulation of both titer and virulence through quorum sensing has been demonstrated in

Sodalis praecaptivus, and allows this species to establish vertically transmitted infections in

weevil and tsetse fly hosts (51, 62)*.* Whether mutualistic *S. symbiotica* have relied on similar

mechanisms to establish persistent vertical transmission in aphids is unclear. Alternatively, *S.*

symbiotica virulence may be attenuated by the loss of one or several key virulence factors before

the establishment of vertical transmission. The experimental tractability of these strains will

allow for future investigations focused on these attenuation mechanisms and the role of vertical

transmission in the transition to a host-restricted lifestyle in aphids.

Materials and Methods

Additional details are provided in SI Appendix.

Isolation and culture of *S. symbiotica.* S. *symbiotica* strain CWBI-2.3^T (DSM 23270) was

obtained from the DSMZ-German Collection of Microorganisms and Cell Cultures and was

grown on tryptic soy agar (TSA) plates at 27 °C (25). *S. symbiotica* strain HB1 was isolated from

the melon aphid (*Aphis gossypii*)*,* collected in August 2018 from HausBar Farms in Austin,

Texas. Details are provided in SI Appendix.

***S. symbiotica* HB1 genome sequencing.** *S. symbiotica* HB1 was grown in TSB at room

temperature, harvested at an $OD_{600} \sim 1.0$, and DNA was extracted with the DNeasy Blood &

Tissue Kit (Qiagen). A paired-end sequencing library with dual barcodes was prepared using the

Illumina Nextera XT DNA kit, and sequencing was performed on an Illumina iSeq 100. Raw

reads were trimmed using Trimmomatic (63) and assembled using the SPAdes algorithm (64) via

Unicycler (65). Genome contamination and completeness were assessed using CheckM (66).

Phylogenetic analysis. *S. symbiotica* and outgroup genomes used for phylogenetic analysis are

listed in Table S1. Genomes were downloaded from the NCBI Assembly Database on March 2,

2020. All outgroup genomes were filtered for completeness > 95% and contamination < 5%

using CheckM (66). Annotations were obtained using Prokka (67), and 176 single-copy

orthologs were identified by OrthoFinder (68). These single-copy orthologs were aligned with

MAFFT (69), trimmed using a BLOSUM62 matrix in BMGE (70), and concatenated using an in-

house script, producing an alignment with 56,881 total amino acid positions. A tree was

constructed by maximum likelihood with a JTT+R10 model and 100 bootstraps, using IQ-Tree

(71). The complete phylogeny is available as Figure S1. The presence of *Serratia* marker genes

was determined using CheckM with the *Serratia* marker gene set provided with CheckM. The

average nucleotide identity for *S. symbiotica* genomes was calculated using FastANI (72).

Chromosomal integration of sfGFP. Superfolder GFP (sfGFP) was integrated into the

chromosomes of *S. symbiotica* CWBI-2.3^T, *S. symbiotica* HB1, and *E. coli* BW25113 through

mini-Tn7 tagging, as described by Choi *et al.* (73). Details are provided in SI Appendix.

Tracking aphid survival, fecundity, and transmission after injection with *S. marcescens, S.*

***symbiotica,* and injection buffer.** Fourth instar pea aphids were injected with *S. marcescens*

Db11*,* recombinant *S. symbiotica* CWBI-2.3^T-GFP, recombinant *S. symbiotica* HB1-GFP,

hemolymph from pea aphids infected with *S. symbiotica* Tucson, or injection buffer, as described

above. Every 24 h, survival was recorded, offspring were collected, and surviving adults were

moved to a fresh dish. Adults were collected at death or at the end of the experiment at 15 DPI

and screened for the presence or absence of *S. symbiotica*. Details are provided in SI Appendix.

Bacterial titer by spot-plating and quantitative PCR (qPCR). Fourth instar pea aphids were

injected with recombinant *S. symbiotica* CWBI-2.3^T-GFP, recombinant *S. symbiotica* HB1-GFP,

or with hemolymph from pea aphids infected with *S. symbiotica* Tucson, as described above. At

24 hours, aphids were transferred in sets of 15 to seedlings of *V. faba* and stored under long-day

conditions (16H light, 8H dark) in incubators held at constant 20 °C. At each timepoint, aphids

were collected in separate tubes, surface sterilized in 10% bleach for 1 minute, rinsed in

deionized water for 1 min, then crushed and resuspended in 100 µL PBS. For aphids injected

with culturable *S. symbiotica* CWBI-2.3^T-GFP or HB1-GFP, 50 µL of this homogenate was used

for spot plating and 50 µL frozen for DNA extraction and qPCR. For aphids injected with *S.*

symbiotica Tucson, all 100 µL of homogenate was frozen and used for DNA extraction and

qPCR. Details are provided in SI Appendix.

Statistical analyses. All statistical analyses and graphing were performed in the R programming

language (version 3.6.3) (74). Survival rates for each treatment group were visualized as Kaplan-

Meier survival curves, and comparisons of rates across treatment groups were performed using

the Cox Proportional Hazards Model. Bacterial titers across treatment groups were compared

using Kruskal-Wallis analysis of variance, followed by Dunn's multiple comparisons test.

Fluorescence *in-situ* hybridization (FISH) microscopy. Fourth instar pea aphids were injected

with wild-type *S. symbiotica* CWBI-2.3^T, wild-type *S. symbiotica* HB1, or with hemolymph from

pea aphids infected with *S. symbiotica* Tucson, as described above. Embryos were dissected at 4

DPI (Movie S1) or 7 DPI (Figure 3) in 70% ethanol. FISH was performed as in Koga *et al.* (8)

with slight modifications. Details are provided in SI Appendix.

Live imaging of *E. coli* and *S. symbiotica* in pea aphids. For live imaging, fourth instar

Acyrthosiphon pisum LSR1 were injected with recombinant *S. symbiotica* CWBI-2.3^T-GFP or

with recombinant *E. coli* BW25113-GFP, as described above. At 24 hours, aphids were

transferred in sets of 15 to seedlings of *V. faba* and stored under long-day conditions (16H light,

8H dark) in incubators held at constant 20 °C. At 5 DPI, a subset of aphids from each treatment

group were used to obtain titer counts via spot-plating, as described above, and remaining aphids

were dissected in TC-100 insect medium. Single ovarioles from ten infected aphids per treatment

group were observed under a Zeiss LSM 710 confocal microscope.

Acknowledgements

We thank Kim Hammond for maintaining aphid lines, Dorsey Barger for access to insect

collection sites at Hausbar Farms, and Margaret Steele, Travis Wiles, Peng Geng, and Sean

Leonard for providing bacterial strains. We thank Ryuichi Koga for training and advice on FISH

microscopy. We thank Daniel Deatherage for sequencing *S. symbiotica* HB1. We thank Anna

Webb at the Microscopy and Imaging Facility of the Center for Biomedical Research Support at

UT-Austin and Nicholas Kocian at Leica Microsystems for assistance with microscopy. We

acknowledge the Texas Advanced Computing Center (TACC) for providing HPC resources. We

thank members of Moran, Howard Ochman, and Barrick labs for helpful comments. pTNS2 was

a gift from Herbert Schweizer (Addgene plasmids 64968). pTn7xKS-sfGFP was a gift from

Karen Guillemin (Addgene plasmid 117394).

Data availability

This Whole Genome Shotgun project for *S. symbiotica* HB1 has been deposited at

DDBJ/ENA/GenBank under the accession JACBGK000000000. The version described in this

paper is version JACBGK010000000.

Competing interests

The authors declare that they have no competing interests.

Funding

JP was supported in part by the University of Texas at Austin Provost Graduate Excellence

Fellowship. This work was funded by the Defense Advanced Projects Agency (HR0011-17-2-

0052 to JEB and NAM) and by the National Science Foundation (DEB-1551092 to NAM).

References

1.   McCutcheon JP, Boyd BM, Dale C. 2019. The life of an insect endosymbiont from the

cradle to the grave. Curr Biol 29:R485–R495.

2.   Ewald PW. 1987. Transmission modes and evolution of the parasitism-mutualism

continuum. Ann NY Acad Sci 503:295–306.

3.   Maynard Smith J, Szathmáry E. 1998. The major transitions in evolution. Oxford

University Press, Oxford, United Kingdom.

4.   Bull JJ, Molineux IJ, Rice WR. 1991. Selection of benevolence in a host-parasite system.

Evolution 45:875–882.

5.   Bright M, Bulgheresi S. 2010. A complex journey: transmission of microbial symbionts.

Nat Rev Microbiol 8:218–230.

6.   Fukatsu T, Hosokawa T. 2002. Capsule-transmitted gut symbiotic bacterium of the

Japanese common plataspid stinkbug, *Megacopta punctatissima*. Appl Environ Microbiol

68:389–396.

7.   Buchner P. 1965. Endosymbiosis of animals with plant microorganisms. Interscience, New

York.

8.   Koga R, Meng X-Y, Tsuchida T, Fukatsu T. 2012. Cellular mechanism for selective

vertical transmission of an obligate insect symbiont at the bacteriocyte–embryo interface.

Proc Natl Acad Sci USA 109:E1230–E1237.

9.   Hosokawa T, Koga R, Kikuchi Y, Meng X-Y, Fukatsu T. 2010. *Wolbachia* as a

bacteriocyte-associated nutritional mutualist. Proc Natl Acad Sci USA 107:769–774.

10. Luan J, Sun X, Fei Z, Douglas AE. 2018. Maternal inheritance of a single somatic animal

cell displayed by the bacteriocyte in the whitefly *Bemisia tabaci*. Curr Biol 28:459-465.e3.

11. Shigenobu S, Watanabe H, Hattori M, Sakaki Y, Ishikawa H. 2000. Genome sequence of

the endocellular bacterial symbiont of aphids *Buchnera* sp. APS. Nature 407:81–86.

12. Hansen AK, Moran NA. 2011. Aphid genome expression reveals host–symbiont

cooperation in the production of amino acids. Proc Natl Acad Sci USA 108:2849–2854.

13. Douglas AE, Prosser WA. 1992. Synthesis of the essential amino acid tryptophan in the pea

aphid (*Acyrthosiphon pisum*) symbiosis. J Insect Physiol 38:565–568.

14. Lai CY, Baumann L, Baumann P. 1994. Amplification of trpEG: adaptation of *Buchnera*

aphidicola to an endosymbiotic association with aphids. Proc Natl Acad Sci USA 91:3819–

3823.

15. Sasaki T, Ishikawa H. 1995. Production of essential amino acids from glutamate by

mycetocyte symbionts of the pea aphid, *Acyrthosiphon pisum*. J Insect Physiol 41:41–46.

16. Douglas AE. 1998. Nutritional interactions in insect-microbial symbioses: aphids and their

symbiotic bacteria *Buchnera*. Annu Rev Entomol 43:17–37.

17. Moran NA, Munson MA, Baumann P, Ishikawa H. 1993. A molecular clock in

endosymbiotic bacteria is calibrated using the insect hosts. P R Soc B 253:167–171.

18. Chong RA, Park H, Moran NA. 2019. Genome evolution of the obligate endosymbiont

Buchnera aphidicola. Mol Biol Evol 36:1481–1489.

19. Sandström JP, Russell JA, White JP, Moran NA. 2001. Independent origins and horizontal

transfer of bacterial symbionts of aphids. Mol Ecol 10:217–228.

20. Moran NA, Russell JA, Koga R, Fukatsu T. 2005. Evolutionary relationships of three new

species of *Enterobacteriaceae* living as symbionts of aphids and other insects. Appl

Environ Microbiol 71:3302–3310.

21. Oliver KM, Degnan PH, Burke GR, Moran NA. 2010. Facultative symbionts in aphids and

the horizontal transfer of ecologically important traits. Annu Rev Entomol 55:247–266.

22. Manzano-Marín A, Szabó G, Simon J-C, Horn M, Latorre A. 2017. Happens in the best of

subfamilies: establishment and repeated replacements of co-obligate secondary

endosymbionts within Lachninae aphids. Environ Microbiol 19:393–408.

23. Burke G, Fiehn O, Moran N. 2010. Effects of facultative symbionts and heat stress on the

metabolome of pea aphids. ISME J 4:242–252.

24. Lamelas A, Gosalbes MJ, Manzano-Marín A, Peretó J, Moya A, Latorre A. 2011. *Serratia*

symbiotica from the aphid *Cinara cedri*: a missing link from facultative to obligate insect

endosymbiont. PLOS Genet 7:e1002357.

25. Sabri A, Leroy P, Haubruge E, Hance T, Frère I, Destain J, Thonart P. 2011. Isolation, pure

culture and characterization of *Serratia symbiotica* sp. nov., the R-type of secondary

endosymbiont of the black bean aphid *Aphis fabae*. Int J Syst Evol Micr 61:2081–2088.

26. Manzano-Marín A, Latorre A. 2014. Settling down: The genome of *Serratia symbiotica*

from the aphid *Cinara tujafilina* zooms in on the process of accommodation to a

cooperative intracellular life. Genome Biol Evol 6:1683–1698.

27. Manzano-Marín A, Simon J-C, Latorre A. 2016. Reinventing the wheel and making it

round again: evolutionary convergence in *Buchnera–Serratia* symbiotic consortia between

the distantly related Lachninae aphids *Tuberolachnus salignus* and *Cinara cedri*. Genome

Biol Evol 8:1440–1458.

28.   Meseguer AS, Manzano-Marín A, d'Acier AC, Clamens A-L, Godefroid M, Jousselin E.

496   2017. *Buchnera* has changed flatmate but the repeated replacement of co-obligate

497   symbionts is not associated with the ecological expansions of their aphid hosts. Mol Ecol

498   26:2363–2378.

499   29.   Renoz F, Pons I, Vanderpoorten A, Bataille G, Noël C, Foray V, Pierson V, Hance T. 2018.

500   Evidence for gut-associated *Serratia symbiotica* in wild aphids and ants provides new

501   perspectives on the evolution of bacterial mutualism in insects. Microb Ecol 78:159–169.

502   30.   Grigorescu AS, Renoz F, Sabri A, Foray V, Hance T, Thonart P. 2018. Accessing the

503   hidden microbial diversity of aphids: an illustration of how culture-dependent methods can

504   be used to decipher the insect microbiota. Microb Ecol 75:1035–1048.

505   31.   Renoz F, Ambroise J, Bearzatto B, Baa-Puyoulet P, Calevro F, Gala J-L, Hance T. 2020.

506   Draft genome sequences of two cultivable strains of the bacterial symbiont *Serratia*

507   *symbiotica*. Microbiol Resour Announc 9:e01579-19.

508   32.   Monnin D, Jackson R, Kiers ET, Bunker M, Ellers J, Henry LM. 2020. Parallel evolution in

509   the integration of a co-obligate aphid symbiosis. Curr Biol 30:1949–1957.

510   33.   Foray V, Grigorescu AS, Sabri A, Haubruge E, Lognay G, Francis F, Fauconnier M-L,

511   Hance T, Thonart P. 2014. Whole-genome sequence of *Serratia symbiotica* strain CWBI-

512   2.3^{T}, a free-living symbiont of the black bean aphid *Aphis fabae*. Genome Announc

513   2:e00767-14.

514   34.   Chen D-Q, Purcell AH. 1997. Occurrence and transmission of facultative endosymbionts in

515   aphids. Current Microbiology 34:220–225.

35.   Montllor CB, Maxmen A, Purcell AH. 2002. Facultative bacterial endosymbionts benefit

pea aphids *Acyrthosiphon pisum* under heat stress. Ecol Entomol 27:189–195.

36.   Oliver KM, Russell JA, Moran NA, Hunter MS. 2003. Facultative bacterial symbionts in

aphids confer resistance to parasitic wasps. Proc Natl Acad Sci USA 100:1803–1807.

37.   Miura T, Braendle C, Shingleton A, Sisk G, Kambhampati S, Stern DL. 2003. A

comparison of parthenogenetic and sexual embryogenesis of the pea aphid *Acyrthosiphon*

pisum (Hemiptera: Aphidoidea). J Exp Zool Part B 295B:59–81.

38.   Wilkinson TL, Fukatsu T, Ishikawa H. 2003. Transmission of symbiotic bacteria *Buchnera*

to parthenogenetic embryos in the aphid *Acyrthosiphon pisum* (Hemiptera: Aphidoidea).

Arthropod Struct Dev 32:241–245.

39.   Pons I, Renoz F, Noël C, Hance T. 2019. Circulation of the cultivable symbiont *Serratia*

symbiotica in aphids Is mediated by plants. Front Microbiol 10:764.

40.   Pons I, Renoz F, Noël C, Hance T. 2019. New insights into the nature of symbiotic

associations in aphids: infection process, biological effects, and transmission mode of

cultivable *Serratia symbiotica* bacteria. Appl Environ Microbiol 85:e02445-18.

41.   Burke GR, Moran NA. 2011. Massive genomic decay in *Serratia symbiotica*, a recently

evolved symbiont of aphids. Genome Biol Evol 3:195–208.

42.   Nikoh N, Koga R, Oshima K, Hattori M, Fukatsu T. 2019. Genome sequence of

"*Candidatus* Serratia symbiotica" strain IS, a facultative bacterial symbiont of the pea aphid

Acyrthosiphon pisum. Microbiol Resour Announc 8:e00272-19.

43.   Lamelas A, Pérez-Brocal V, Gómez-Valero L, Gosalbes MJ, Moya A, Latorre A. 2008.

Evolution of the secondary symbiont "*Candidatus* Serratia symbiotica" in aphid species of

the subfamily Lachninae. Appl Environ Microbiol 74:4236–4240.

44.   Renoz F, Noël C, Errachid A, Foray V, Hance T. 2015. Infection dynamic of symbiotic

bacteria in the pea aphid *Acyrthosiphon pisum* gut and host immune response at the early

steps in the infection process. PLOS One 10:e0122099.

45.   Flyg C, Kenne K, Boman HG. 1980. Insect pathogenic properties of *Serratia marcescens*:

phage-resistant mutants with a decreased resistance to cecropia immunity and a decreased

virulence to *Drosophila*. Microbiology 120:173–181.

46.   Braendle C, Miura T, Bickel R, Shingleton AW, Kambhampati S, Stern DL. 2003.

Developmental origin and evolution of bacteriocytes in the aphid–*Buchnera* symbiosis.

PLOS Biol 1:70–76.

47.   Altincicek B, ter Braak B, Laughton AM, Udekwu KI, Gerardo NM. 2011. *Escherichia coli*

K-12 pathogenicity in the pea aphid, *Acyrthosiphon pisum*, reveals reduced antibacterial

defense in aphids. Dev Comp Immunol 35:1091–1097.

48.   Koga R, Tsuchida T, Sakurai M, Fukatsu T. 2007. Selective elimination of aphid

endosymbionts: effects of antibiotic dose and host genotype, and fitness consequences.

FEMS Microbiol Ecol 60:229–239.

49.   Dale C, Young SA, Haydon DT, Welburn SC. 2001. The insect endosymbiont *Sodalis*

glossinidius utilizes a type III secretion system for cell invasion. Proc Natl Acad Sci USA

98:1883–1888.

50. Nadal-Jimenez P, Griffin JS, Davies L, Frost CL, Marcello M, Hurst GDD. 2019. Genetic

manipulation allows in vivo tracking of the life cycle of the son-killer symbiont,

Arsenophonus nasoniae, and reveals patterns of host invasion, tropism and pathology.

Environ Microbiol 21:3172–3182.

51. Medina Munoz M, Spencer N, Enomoto S, Dale C, Rio RVM. 2020. Quorum sensing sets

the stage for the establishment and vertical transmission of *Sodalis praecaptivus* in tsetse

flies. PLOS Genet 16:e1008992.

52. Moran NA, McCutcheon JP, Nakabachi A. 2008. Genomics and evolution of heritable

bacterial symbionts. Annu Rev Genet 42:165–190.

53. Tsuchida T, Koga R, Shibao H, Matsumoto T, Fukatsu T. 2002. Diversity and geographic

distribution of secondary endosymbiotic bacteria in natural populations of the pea aphid,

Acyrthosiphon pisum. Mol Ecol 11:2123–2135.

54. Russell JA, Latorre A, Sabater-Muñoz B, Moya A, Moran NA. 2003. Side-stepping

secondary symbionts: widespread horizontal transfer across and beyond the Aphidoidea.

Mol Ecol 12:1061–1075.

55. Husnik F, McCutcheon JP. 2016. Repeated replacement of an intrabacterial symbiont in the

tripartite nested mealybug symbiosis. Proc Natl Acad Sci USA 113:E5416–E5424.

56. Bennett GM, Moran NA. 2015. Heritable symbiosis: the advantages and perils of an

evolutionary rabbit hole. Proc Natl Acad Sci USA 112:10169–10176.

57. Herren JK, Paredes JC, Schüpfer F, Lemaitre B. 2013. Vertical transmission of a

Drosophila endosymbiont via cooption of the yolk transport and internalization machinery.

mBio 4:e00532-12.

58. Nakabachi A, Ishikawa H, Kudo T. 2003. Extraordinary proliferation of microorganisms in

aposymbiotic pea aphids, *Acyrthosiphon pisum*. J Invertebr Pathol 82:152–161.

59. Chong RA, Moran NA. 2018. Evolutionary loss and replacement of *Buchnera*, the obligate

endosymbiont of aphids. ISME J 12:898–908.

60. Fukatsu T, Nikoh N, Kawai R, Koga R. 2000. The secondary endosymbiotic bacterium of

the pea aphid *Acyrthosiphon pisum* (Insecta: Homoptera). Appl Environ Microbiol

66:2748–2758.

61. Koga R, Tsuchida T, Fukatsu T. 2003. Changing partners in an obligate symbiosis: a

facultative endosymbiont can compensate for loss of the essential endosymbiont *Buchnera*

in an aphid. P R Soc B 270:2543–2550.

62. Enomoto S, Chari A, Clayton AL, Dale C. 2017. Quorum sensing attenuates virulence in

Sodalis praecaptivus. Cell Host Microbe 21:629-636.e5.

63. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics 30:2114–2120.

64. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM,

Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G,

Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its

applications to single-cell sequencing. J Comput Biol 19:455–477.

65.   Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome

assemblies from short and long sequencing reads. PLOS Comput Biol 13:e1005595.

66.   Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM:

assessing the quality of microbial genomes recovered from isolates, single cells, and

metagenomes. Genome Res 25:1043–1055.

67.   Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics 30:2068–

2069.

68.   Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative

genomics. Genome Biol 20:238.

69.   Katoh K, Misawa K, Kuma K, Miyata T. 2002. MAFFT: a novel method for rapid multiple

sequence alignment based on fast Fourier transform. Nucleic Acids Res 30:3059–3066.

70.   Criscuolo A, Gribaldo S. 2010. BMGE (Block Mapping and Gathering with Entropy): a

new software for selection of phylogenetic informative regions from multiple sequence

alignments. BMC Evol Biol 10:210.

71.   Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and effective

stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol

32:268–274.

72.   Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. 2018. High

throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. Nat

Commun 9:5114.

73.   Choi K-H, Schweizer HP. 2006. Mini-Tn*7* insertion in bacteria with single *att*Tn*7* sites:

example *Pseudomonas aeruginosa*. Nat Protoc 1:153–161.

74.   R Core Team. 2020. R: a language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria.

Figure Legends

Figure 1. Phylogeny and gene content evolution of *S. symbiotica*. (A) Maximum likelihood

phylogeny of *S. symbiotica,* based on concatenation of 176 single-copy orthologs (56,881 amino

acid positions) shared across *S. symbiotica,* other *Serratia* species, *E. coli, Yersinia pestis,* and

Salmonella enterica. Genomes used for the phylogeny and relevant references for *S. symbiotica*

genome features are listed in Table S1. Bootstrap values are indicated with symbols on nodes.

The complete phylogeny is presented in Figure S1.

Figure 2. Infection dynamics of *S. symbiotica* strains CWBI-2.3$^\text{T}$ and HB1 injected into

hemolymph of fourth instar pea aphids. (A) Survival curves of pea aphids injected with cultured

S. marcescens Db11, recombinant *S. symbiotica* CWBI-2.3$^\text{T}$-GFP, or recombinant *S. symbiotica*

HB1-GFP, with hemolymph from *S. symbiotica* Tucson, or with injection buffer. *** $P < 0.001$,

Cox Proportional Hazards Model. (B) Fecundity and transmission of *S. symbiotica* for adults

injected with *S. symbiotica* Tucson. Boxplots in gray depict the total offspring, including infected

and uninfected offspring, per adult. Boxplots in orange depict total offspring infected with *S.*

symbiotica Tucson per adult. Filled curves represent a LOESS regression and display the

increase in offspring infected with *S. symbiotica* Tucson from 10 DPI to 15 DPI. (C) Bacterial

titer of recombinant *S. symbiotica* strains CWBI-2.3^T-GFP and HB1-GFP, obtained by spot-

plating dilutions from whole aphids. ** $p < 0.01$, Wilcoxon rank-sum test. (D) Relative titer of *S.*

symbiotica across treatment groups at 5 DPI (left) and for 7-day-old pea aphids from lab-reared

clonal lines that are naturally infected with vertically transmitted *S. symbiotica* strains,

established from collections in Austin in 2014 (Austin), Tucson in 2019 (Tuc19), Tucson in 1999

(Tucson), and Wisconsin in 2017 (WIG5) (right). Relative titer was calculated as the copy

number of *S. symbiotica dnaK* normalized by copy number of the single copy *Acyrthosiphon*

pisum gene *ef1a*. CWBI-2.3^T and HB1 samples used for qPCR are the same as those used to

determine bacterial titer by spot-plating. Letters a and b denote two groups of strains that have

significantly different titers from one another and indistinguishable titers within groups at $\alpha =$

0.01 by the Kruskal-Wallis test followed by Dunn's multiple comparison test.

Figure 3. Transovarial transmission of the pathogenic strain *S. symbiotica* CWBI-2.3^T and the

mutualistic strain *S. symbiotica* Tucson in stage 7 embryos of pea aphids. (A) *S. symbiotica*

Tucson infection across embryonic stages at 7 DPI. (B) *S. symbiotica* Tucson does not infect

embryos that are beyond stage 7 of development at the time of injection, as indicated by a lack of

S. symbiotica Tucson in late-stage embryos. (C) In infected, mid-stage embryos, *S. symbiotica*

Tucson cannot infect primary bacteriocytes that contain *B. aphidicola* but does infect sheath

cells. (D) *S. symbiotica* Tucson enters the syncytium of an early-stage blastula with *B. aphidicola*

but is underrepresented relative to *B. aphidicola* during this infection. (E-F) *S. symbiotica*

CWBI-2.3^T infection across embryonic stages at 7 DPI. Embryos depicted in (E) and (F) derive

from the same ovariole but were separated during dissection. Infection with this strain results in

reduced embryonic growth, as indicated by scale bars. (G) *S. symbiotica* CWBI-2.3^T does not

infect embryos that are beyond stage 7 of development at the time of injection, as indicated by a

lack of *S. symbiotica* CWBI-2.3^T in late-stage embryos. (H) In infected, mid-stage embryos, *S.*

symbiotica CWBI-2.3^T cannot infect primary bacteriocytes that contain *B. aphidicola* but does

infect sheath cells. (I) *S. symbiotica* CWBI-2.3^T enters the syncytium of an early-stage blastula

with *B. aphidicola* and is overrepresented relative to *B. aphidicola* during this infection. Blue,

red, and green signal are used for host nuclei, *S. symbiotica,* and *B. aphidicola,* respectively.

Figure 4. Ability of *S. symbiotica* CWBI-2.3^T but not *E. coli* BW25113 to achieve transovarial

transmission to stage 7 embryos of pea aphids. (A) Recombinant *S. symbiotica* CWBI-2.3^T-GFP

and *E. coli* BW25113-GFP reach comparable titers 5 DPI, as determined by spot-plating. (B, C)

Recombinant *S. symbiotica* CWBI-2.3^T-GFP enters the posterior region of stage 7 blastula.

Embryos were dissected and visualized at 5 DPI. Dotted lines outline *S. symbiotica* present

within the embryo. (D, E) Recombinant *E. coli* BW25113-GFP does not enter into stage 7

blastula. Embryos were dissected and visualized at 5 DPI. Arrows indicate bacteria attached to

the embryonic surface. Scale bars represent 20 μM.

Supplementary Figures, Tables, and Movies

Figure S1. Full phylogeny of *S. symbiotica* and outgroup. Maximum likelihood phylogeny with

a JTT+R10 model and 100 bootstraps*,* based on concatenation of 176 single-copy orthologs

(56,881 amino acid positions) shared across *S. symbiotica, Serratia*, *E. coli, Yersinia pestis,* and

Salmonella enterica. Full list of genomes used for the phylogeny is provided in Table S1.

Table S1. Accession numbers, genome information, and references for analyzed *S. symbiotica*

strains (Figures 1, S1).

Table S2. Average nucleotide identities for pairwise combinations of *S. symbiotica* strains.

Movie S1. Z-stack showing *S. symbiotica* HB1 entering the syncytium of a stage 7 embryo with

B. aphidicola at 4 DPI. Blue, red, and green signal are used for host nuclei, *S. symbiotica,* and *B.*

aphidicola, respectively.

Movie S2. Z-stack showing *S. symbiotica* CWBI-2.3^{T}-GFP entering a live stage 7 embryo. Z-

stack corresponds to the embryo depicted in Figure 4B.

Movie S3. Z-stack showing *S. symbiotica* CWBI-2.3^{T}-GFP entering a live stage 7 embryo. Z-

stack corresponds to the embryo depicted in Figure 4C.

Movie S4. Z-stack showing *E. coli* BW25113-GFP surrounding, but not entering, a live stage 7

embryo. Z-stack corresponds to the embryo depicted in Figure 4D.

Movie S5. Z-stack showing *E. coli* BW25113-GFP surrounding, but not entering, a live stage 7

embryo. Z-stack corresponds to the embryo depicted in Figure 4E.

Strain	Size (Mb)	*Serratia* marker genes (985)	% GC	Host species	Lifestyle
PCOLA-89S	1.66	659 (66.9%)	51.6	*Periphyllus acericola*	
PERIS-14S	1.54	599 (60.8%)	48.6	*Periphyllus aceris*	Co-obligate
PLYR-94S	2.58	779 (79.1%)	51.7	*Periphyllus lyropictus*	
CWBI-2.3ᵀ	3.58	870 (88.3%)	52.1	*Aphis fabae*	
Apa8A1	3.23	878 (89.1%)	51.9	*Aphis passeriniana*	
24.1	3.09	877 (89.0%)	51.4	*Aphis fabae*	Gut pathogen
HB1	3.19	879 (89.2%)	51.9	*Aphis gossypii*	
AURT-53S	2.03	657 (66.7%)	51.6	*Aphis urticata*	
MCAR-56S	2.17	758 (77.0%)	52.5	*Microlophium carnosum*	Vertically-transmitted mutualist
IS	2.82	828 (84.1%)	51.9	*Acyrthosiphon pisum*	
Tucson	2.78	816 (82.8%)	47.9	*Acyrthosiphon pisum*	
SCt-VLC	2.50	762 (77.4%)	51.9	*Cinara tujafilina*	Co-obligate

Figure 1. Phylogeny and gene content evolution of *S. symbiotica*. (A) Maximum likelihood phylogeny of *S. symbiotica*, based on concatenation of 176 single-copy orthologs (56,881 amino acid positions) shared across *S. symbiotica*, other *Serratia* species, *E. coli*, *Yersinia pestis*, and *Salmonella enterica*. Genomes used for the phylogeny and relevant references for *S. symbiotica* genome features are listed in Table S1. Bootstrap values are indicated with symbols on nodes. The complete phylogeny is presented in Figure S1.

Figure 2. Infection dynamics of *S. symbiotica* strains CWBI-2.3[T] and HB1 injected into hemolymph of fourth instar pea aphids. (A) Survival curves of pea aphids injected with cultured *S. marcescens* Db11, recombinant *S. symbiotica* CWBI-2.3[T]-GFP, or recombinant *S. symbiotica* HB1-GFP, with hemolymph from *S. symbiotica* Tucson, or with injection buffer. *** P < 0.001, Cox Proportional Hazards Model. (B) Fecundity and transmission of *S. symbiotica* for adults injected with *S. symbiotica* Tucson. Boxplots in gray depict the total offspring, including infected and uninfected offspring, per adult. Boxplots in orange depict total offspring infected with *S. symbiotica* Tucson per adult. Filled curves represent a LOESS regression and display the increase in offspring infected with *S. symbiotica* Tucson from 10 DPI to 15 DPI. (C) Bacterial titer of recombinant *S. symbiotica* strains CWBI-2.3[T]-GFP and HB1-GFP, obtained by spot-plating dilutions from whole aphids. ** p < 0.01, Wilcoxon rank-sum test. (D) Relative titer of *S. symbiotica* across treatment groups at 5 DPI (left) and for 7-day-old pea aphids from lab-reared clonal lines that are naturally infected with vertically transmitted *S. symbiotica* strains, established from collections in Austin in 2014 (Austin), Tucson in 2019 (Tuc19), Tucson in 1999 (Tucson), and Wisconsin in 2017 (WIG5) (right). Relative titer was calculated as the copy number of *S. symbiotica* dnaK normalized by copy number of the single copy *Acyrthosiphon pisum* gene *ef1a*. CWBI-2.3[T] and HB1 samples used for qPCR are the same as those used to determine bacterial titer by spot-plating. Letters a and b denote two groups of strains that have significantly different titers from one another and indistinguishable titers within groups at α = 0.01 by the Kruskal-Wallis test followed by Dunn's multiple comparison test.

Figure 3. Transovarial transmission of the pathogenic strain *S. symbiotica* CWBI-2.3^T and the mutualistic strain *S. symbiotica* Tucson in stage 7 embryos of pea aphids. (A) *S. symbiotica* Tucson infection across embryonic stages at 7 DPI. (B) *S. symbiotica* Tucson does not infect embryos that are beyond stage 7 of development at the time of injection, as indicated by a lack of *S. symbiotica* Tucson in late-stage embryos. (C) In infected, mid-stage embryos, *S. symbiotica* Tucson cannot infect primary bacteriocytes that contain *B. aphidicola* but does infect sheath cells. (D) *S. symbiotica* Tucson enters the syncytium of an early-stage blastula with *B. aphidicola* but is underrepresented relative to *B. aphidicola* during this infection. (E-F) *S. symbiotica* CWBI-2.3^T infection across embryonic stages at 7 DPI. Embryos depicted in (E) and (F) derive from the same ovariole but were separated during dissection. Infection with this strain results in reduced embryonic growth, as indicated by scale bars. (G) *S. symbiotica* CWBI-2.3^T does not infect embryos that are beyond stage 7 of development at the time of injection, as indicated by a lack of *S. symbiotica* CWBI-2.3^T in late-stage embryos. (H) In infected, mid-stage embryos, *S. symbiotica* CWBI-2.3^T cannot infect primary bacteriocytes that contain *B. aphidicola* but does infect sheath cells. (I) *S. symbiotica* CWBI-2.3^T enters the syncytium of an early-stage blastula with *B. aphidicola* and is overrepresented relative to *B. aphidicola* during this infection. Blue, red, and green signal are used for host nuclei, *S. symbiotica*, and *B. aphidicola*, respectively.

Figure 4. Ability of *S. symbiotica* CWBI-2.3^T but not *E. coli* BW25113 to achieve transovarial transmission to stage 7 embryos of pea aphids. (A) Recombinant *S. symbiotica* CWBI-2.3^T-GFP and *E. coli* BW25113-GFP reach comparable titers 5 DPI, as determined by spot-plating. (B, C) Recombinant *S. symbiotica* CWBI-2.3^T-GFP enters the posterior region of stage 7 blastula. Embryos were dissected and visualized at 5 DPI. Dotted lines outline *S. symbiotica* present within the embryo. (D, E) Recombinant *E. coli* BW25113-GFP does not enter into stage 7 blastula. Embryos were dissected and visualized at 5 DPI. Arrows indicate bacteria attached to the embryonic surface. Scale bars represent 20 µM.