Dynein/Sunday driver mediate disorders of dynamic profiles of Golgi outposts induced by amyloid precursor protein

Authors: Qianqian Du[1,2], Guo Cheng[1,2], Jin Chang[1,2], Wei Zhou[1,2*]

Affiliations

[1] Britton Chance Center for Biomedical Photonics, Wuhan National Laboratory for Optoelectronics-Huazhong University of Science and Technology, Wuhan, Hubei 430074, China

[2] MoE Key Laboratory for Biomedical Photonics, Collaborative Innovation Center for Biomedical Engineering, School of Engineering Sciences, Huazhong University of Science and Technology, Wuhan, Hubei 430074, China

*Correspondence to:

Wei Zhou, wzhou@mail.hust.edu.cn

Tel: + 86 27 87792033-303; + 86 15827641600;

Fax: +86 27 87792034

Abstract

Dendritic abnormality is the pathological hallmark of Alzheimer's disease (AD), but its underlying mechanism remains elusive. Here we showed that amyloid precursor protein (APP), the precursor of β-amyloid, caused dendritic defects in dendritic arborization neurons of *Drosophila* accompanied by Golgi outposts (GOPs) disorders. Half of the dendritic APP co-localized with GOPs, impairing their distribution and compartmental organization; in particular, the dynamic behaviors. By analyzing the state of anterograde/retrograde movements, we found that the motor protein dynein and its adapter protein Sunday driver (Syd) mediated APP-induced dynamic alterations of multi-compartment GOPs (mcGOPs) rather than single-compartment GOPs (scGOPs). Moreover, the loss of *Syd* rescued the defects of dendritic complexity caused by APP. Our findings indicate that APP resident in GOPs disrupts the equilibrium of anterograde/retrograde movements of mcGOPs maintained by dynein/Syd complex, resulting in dendritic abnormalities. This offers new insight into dendritic pathology in AD.

Keywords: APP, dendritic Golgi outposts, Sunday driver, dendritic pathology

Introduction

Alzheimer's disease (AD), characterized by the accumulation of β-amyloid (Aβ) plaque and neurofibrillary tangles in the brain, is one of the most common age-dependent neurodegenerative disorders (Terry, 1996; Terry & Katzman, 1983). Notably, growing evidence shows that dendritic abnormalities, including dystrophic neurites, reduction of dendritic complexity, and loss of dendritic spines, are prominent early-stage phenotypes of AD (SJ Baloyannis, 2009). However, the underlying mechanism remains elusive.

Amyloid precursor protein (APP) is the β-amyloid precursor and plays a significant role in the pathogenesis of AD (J. Hardy & Selkoe, 2002; J. A. Hardy & Higgins, 1992). Mutations of the *APP* gene can induce early-onset familial AD (Golde, Estus, Younkin, Selkoe, & Younkin, 1992), and excess expression of APP is also considered a pathological factor of sporadic AD (Sisodia & Price, 1995). Overexpression of APP results in morphological changes in dendrites, such as reduced dendrite length, bifurcation nodes, and surface area as well as spine density (Alpar et al., 2006; Luebke et al., 2010; Šišková et al., 2014). However, although both APP and related processing enzymes have been found in dendrites (Das et al., 2013; Das et al., 2016; Saraceno et al., 2014), it remains unclear how APP induces the dendritic defects.

Impairments of the structure and function of the Golgi apparatus are implicated in AD. Fragmented Golgi apparatus has been observed in cerebellar Purkinje neurons and hippocampal neurons of AD patients (S Baloyannis, 2002; S. J. Baloyannis, 2014). Meanwhile, the accumulation of Aβ produced by amyloidogenic APP processing has been shown to induce Golgi fragmentation by triggering phosphorylation of Golgi structural proteins such as GRASP65 (Gunjan & Yanzhuang, 2015; Joshi,

Chi, Huang, & Wang, 2014).

In neurons, the Golgi apparatus is present not only in the soma but also in the dendrites as Golgi

outposts (GOPs) (Horton & Ehlers, 2003; Horton et al., 2005; Pierce, Mayer, & Mccarthy, 2001).

Functioning as local secretory stations (Horton & Ehlers, 2003; Pierce et al., 2001; Valenzuela & Perez,

2015) and acentrosomal microtubule nucleation sites (Ori-McKenney, Jan, & Jan, 2012) in the

dendrites, GOPs are required for proper dendrite growth and maintenance. Disruptions to the

distribution of GOPs are always accompanied by abnormal dendritic events, such as atypical branching

patterns (Ye et al., 2007) and terminal dendrite elongation (Chung et al., 2017). Mutation of Golgi

structural protein dGM130, a regulator of the compartmental organization of GOPs, leads to a

reduction in high-order dendritic branches (Zhou et al., 2014). Furthermore, time-lapse imaging has

revealed that GOPs moving towards dendritic ends and the soma are tightly linked to the extension

and retraction of dendritic branches, and disruption of GOPs transport is likely to cause restrained

dendrite development and arborization (Arthur, Yang, Abellaneda, & Wildonger, 2015; Lin et al., 2015;

Ye et al., 2007). These converging evidence suggests that GOPs contribute to dendritic morphology,

an indication that GOPs are involved in APP-induced dendritic defects in AD.

To determine how GOPs contribute to dendritic aberrations in the associated pathological events,

we investigated the structural and dynamic profiles of GOPs in class III dendritic arborization neurons

(C3da neurons) of *Drosophila,* which expressed APP ectopically. First, the colocalization between

APP and GOPs was measured and their dynamic profiles were analyzed. Subsequently, dual-color live

imaging with *medial-* and *trans*-Golgi markers was used to analyze the compartmental organization

and dynamics of GOPs in wild-type and APP-neurons. We further defined the dynamic properties of

GOPs affected by APP by manipulating the dynein and kinesin function. Finally, the adaptor protein

associated with GOPs dynamic alterations was found by colocalization with GOPs and the rescue of

both GOPs dynamic and dendritic shapes.

Results

APP induces dendritic abnormalities in C3da neurons

To mimic the dendritic pathology in AD, we expressed APP in C3da neurons, a classic model for

dendrite development (Grueber, Jan, & Jan, 2002). Firstly, we observed the distribution of APP in

neurons by ectopically expressed APP-GFP. With the anti-HRP labeling of the entire neuron shape,

APP appeared as punctate patterns and was localized in both dendrites and axon (Figures 1a and b).

The quantification of APP-GFP revealed 45.9% ± 4.8% in the dendrites, as well as 54.1% ± 4.8% in

the axon (Figure 1c). This implies that nearly half of APP settles as puncta in the dendrites.

We then examined how APP influenced the dendritic morphology of C3da ddaA neurons. This

showed significant differences in dendritic features between APP-expressing neurons (APP neurons)

and wild-type neurons (WT), including the total dendrite length, dendritic branching patterns, as well

as the density and length of dendritic spikes (a dendritic spine-like structure in sensory neuron,

(Grueber et al., 2002)) (Figure 1d). Specifically, the total dendrite length (Figure 1e; WT: 1,914.7 ±

79.7 μm; APP: 1,626.9 ± 54.0 μm) and the number of dendritic branch points (Figure 1f; WT: 14.6 ±

1.0; APP: 10.4 ± 0.3) in APP neurons were reduced, which suggest that APP suppresses growth in

dendrites. In comparison, most of the reduction for dendritic branch points occurred in high-order

branch points (the fourth-order and up; Figure 1g; WT: 4.0 ± 0.5; APP: 2.4 ± 0.4). This means that due

to the reduced branching complexity APP restrains dendritic arbor maintenance. Despite the significant

decrease in the density of dendritic spikes (Figure 1h; WT: 27.7 ± 1.0 per 100 μm; APP: 17.4 ± 0.8 per

100 μm), their length remained similar (Figure 1i; WT: 5.4 ± 0.2 μm; APP: 5.5 ± 0.2 μm). Taken

together, these morphological defects indicate that dendrite growth and maintenance are seriously

disrupted in APP neurons. Such reductions are consistent with the commonly reported dendritic

pathology in AD studies (Alpar et al., 2006; Luebke et al., 2010; Šišková et al., 2014). It implies that

APP induces severe dendritic defects in C3da neurons.

APP alters the compartmental organization and distribution of GOPs

To find out why APP caused dendritic defects in C3da neurons, we focused on the dendritic Golgi

outposts (GOPs), which are important for the growth and branching of dendrites (Ori-McKenney et

al., 2012; Ye et al., 2007). We first examined the correlations between APP and GOPs in C3da neurons

using fluorescent protein-tagged APP (APP-GFP) and GalT (GalT-TagRFPt, a *trans*-Golgi marker). In

both soma and dendrites, APP puncta were colocalized with Golgi units (Figure 2a). Quantification

indicated that the overlapped GalT/APP occupied the majority of GalT (Figure 2b; soma: 90.9% ±

2.5%; dendrites: 84.6% ± 3.3%) and half of APP (Figure 2c; soma: 67.6% ± 3.2%; dendrites: 49.6% ±

3.4%). In addition, we tested APP-GFP with the *medial*-Golgi marker (ManII-TagRFPt). Similar to

GalT results, APP was highly colocalized with ManII (Figures 2b and c; In soma, ManII: 87.0% ± 2.8%

and APP: 68.2% ± 3.1%; in dendrites, ManII: 83.0% ± 0.9% and APP: 57.7% ± 4.8%). The

colocalization between APP and GOPs points to the fact that the dendritic APP resides in GOPs. This

implies that APP may influence dendritic patterning by altering GOPs.

We characterized the structure and dynamic of GOPs in APP neurons to determine whether APP

modulated GOPs. Considering the coexistence of stacked multi-compartment GOPs (mcGOPs) and

punctate single-compartment GOPs (scGOPs) in which the *medial-* and *trans-* cisternae are separated

from each other (Zhou et al., 2014), the fluorescently tagged *medial-* and *trans*-Golgi markers (ManII-

GFP and GalT-TagRFPt) were simultaneously used to distinguish GOPs with different compartmental

organization in dendrites (Figure 2d). We observed that the total number of GOPs in APP neurons,

including both mcGOPs and scGOPs reduced by almost half compared with the wild-type (Figure 2e;

WT: 16.2 ± 0.8 per 100 μm; APP: 8.0 ± 0.7 per 100 μm). Moreover, the proportions of mcGOPs and

scGOPs were inversely associated, with the former increasing (Figure 2f; WT: 32.0% ± 4.0%; APP:

48.4% ± 2.9%), while the latter decreased (Figure 2f; WT: 68.0% ± 4.0%; APP: 51.6% ± 2.9%). That

implies that APP induces alteration of the compartmental organization of GOPs.

We further examined the distribution of GOPs in dendrites. To describe the distribution, the

dendrites were divided into three segments depending on the distance to soma: proximal (0 -100 μm),

medial (100 - 200 μm), and distal (> 200 μm). Overall, the distribution tendency of mcGOPs showed

a pronounced shift between wild-type and APP neurons (Figures 2g and h). Specifically, the mcGOPs

were more enriched in the proximal part of dendrite in APP neurons (Figure 2g; proximal: 56.0% ±

5.0%; medial:31.7% ± 5.6%; distal:12.3% ± 3.5%), whereas they dominated the distal part in wild-

type (Figure 2g; proximal: 23.8% ± 2.7%; medial: 24.4% ± 4.8%; distal:51.8% ± 6.4%). In summary,

the compartmental organization and distribution of GOPs are severely disrupted in APP neurons. In

particular, the proportion and distribution tendency of mcGOPs change significantly.

APP disrupts the equilibrium of anterograde/retrograde movements of GOPs

GOPs are highly mobile in dendrites (Horton & Ehlers, 2003; Ye et al., 2007), and that could

influence the distribution. From the co-expression of APP-GFP and GalT-TagRFPt in C3da neurons,

we revealed three different types of puncta in dendrites namely: APP-alone, GOPs-alone, and colocalized APP/GOPs puncta. Using *in vivo* time-lapse imaging, we found that all of them can move bi-directionally (Figure 3a). We also quantified the proportion of mobile GOPs. In APP neurons, the proportion of mobile GOPs decreased from $73.3\% \pm 4.8\%$ of GOPs-alone puncta to $54.8\% \pm 5.0\%$ of colocalized APP/GOPs ones (Figure 3b). Besides, the proportion of GOPs-alone puncta was not statistically different between wild-type and APP neurons (Figure 3b), indicating that APP resides in GOPs and alters the movements of GOPs.

To clarify how APP affected the dynamic of GOPs, we analyzed the trajectories of GOPs labeled by *medial*- and *trans*-Golgi markers in APP neurons simultaneously (Figures 3c and d), which enabled us to access the compartmental organization of dynamic GOPs. Similar to Figure 3b, the percentage of dynamic GOPs was reduced in APP neurons (Figure 3e; WT: $77.6\% \pm 2.1\%$; APP: $64.9\% \pm 3.1\%$). We also characterized the behaviors of dynamic GOPs, including direction, displacement, and velocity. GOPs in dendrites have been reported to be able to move bi-directionally: anterograde (from soma to distal dendrite, called as "anterograde GOPs") and retrograde (from distal dendrite to soma, called as "retrograde GOPs"). In wild-type neurons, the proportion of anterograde versus retrograde movements of GOPs was $31.9\% \pm 1.9\%$ for anterograde to $68.1\% \pm 1.9\%$ for retrograde, whereas in APP neurons it was $47.1\% \pm 3.0\%$ to $52.9\% \pm 3.0\%$ (Figure 3f). This suggests that APP alters the equilibrium of anterograde/retrograde movements of GOPs. We also divided total GOPs into mcGOPs and scGOPs to distinguish the features of dynamic GOPs with different compartmental organizations. It turned out that the change of equilibrium of anterograde/retrograde movements came from both mcGOPs (Figure 3g; WT: anterograde, $38.7\% \pm 2.4\%$; retrograde, $61.3\% \pm 2.4\%$. APP:

anterograde, 49.5% ± 3.1%; retrograde, 50.5% ± 3.1%) and scGOPs (Figure 3h; WT: anterograde,

33.8% ± 2.2%; retrograde, 66.2% ± 2.2%. APP: anterograde, 45.5% ± 3.1%; retrograde, 54.5% ±

3.1%). We also measured the displacements and velocity of GOPs movements in APP neurons and

found only the retrograde displacements were shorter (Figure 3i; WT: mcGOPs / scGOPs, 9.4 ± 1.1 /

8.0 ± 0.9 μm; APP: mcGOPs / scGOPs, 5.3 ± 0.6 / 4.7 ± 0.6 μm). Nevertheless, there were no

alterations in anterograde displacements and velocity (Figures 3i and j). This showed that more GOPs

stayed stationary in APP neurons, whereas the dynamic GOPs tended to have less retrograde

movements with shorter displacements.

Dynein mediates the alteration of mcGOPs dynamic induced by APP

The motor proteins are well known to be involved in the positioning of the Golgi complex (Allan,

Thompson, & McNiven, 2002; Yadav & Linstedt, 2011). There are two primary motor proteins namely

Kinesin and Dynein. Kinesin transports the cargo towards microtubule plus ends, while Dynein

mediates the cargo towards minus-ends microtubule (Gibbons, 1981; Hirokawa, 1998; Vale, Reese, &

Sheetz, 1985). To find out how motor proteins were involved in dynamic of GOPs, we used RNA-

interference to knockdown the motor proteins, including *Dhc (Dhc-64c*, a *dynein heavy chain subunit)*

and *Khc* (*kinesin heavy chain*), which have been reported in movements and localization of GOPs in

C4da neurons (Kelliher et al., 2018; Lin et al., 2015). The results showed that both *Dhc* RNAi and *Khc*

RNAi reduced the proportion of dynamic GOPs (Figures 4a and b; WT: 74.8% ± 2.2%; *Dhc* RNAi:

65.5% ± 3.6%; *Khc* RNAi: 53.0 ± 2.8%). Further, we analyzed the direction of movement for dynamic

GOPs. To describe the state of anterograde/retrograde movements, we defined the direction index (DI)

by calculating the ratio of anterograde to retrograde movements. The results showed that loss of *Dhc*

and *Khc* had opposite effects on dynamic GOPs with *Dhc* RNAi decreasing the DI, and *Khc* RNAi

increasing it (Figures 4c and d; WT: 0.43 ± 0.03; *Dhc* RNAi: 0.30 ± 0.03; *Khc* RNAi: 0.92 ± 0.13). We

also observed that similar to total GOPs, DI for dynamic mcGOPs was downregulated in *Dhc* RNAi

and upregulated in *Khc* RNAi (Figure 4e; WT: 0.54 ± 0.07; *Dhc* RNAi: 0.27± 0.03; *Khc* RNAi: 1.20

± 0.24). However, DI for scGOPs dynamic was only regulated by *Khc* RNAi rather than *Dhc* RNAi

(Figure 4f; WT: 0.46 ± 0.04; *Dhc* RNAi: 0.52± 0.13; *Khc* RNAi: 1.15 ± 0.25). In addition, under *Dhc*

RNAi and *Khc* RNAi, there were no significant changes in the displacements of both mcGOPs (Figure

4g; WT: anterograde / retrograde, 6.0 ± 0.8 μm / 7.9 ± 0.8 μm; *Dhc* RNAi: anterograde / retrograde,

4.7 ± 0.6 μm / 8.2 ± 0.9 μm; *Khc* RNAi: anterograde / retrograde: 5.3 ± 0.7 μm / 7.8 ± 0.7 μm) and

scGOPs (Figure 4h; WT: anterograde / retrograde, 5.4 ± 0.8 μm / 8.7 ± 1.0 μm; *Dhc* RNAi: anterograde

/ retrograde, 4.5 ± 0.9 μm / 7.2 ± 1.0 μm; *Khc* RNAi: anterograde / retrograde, 5.4 ± 0.9 μm / 8.0 ±

1.2μm). This implies that Dynein and Kinesin work together to drive the dynamic of mcGOPs, and

maintain the equilibrium between anterograde and retrograde movements.

We then examined the dynamic behaviors of GOPs in APP neurons to determine the roles of motor

proteins in alterations of GOPs dynamic induced by APP. After the knockdown of Dynein and Kinesin

with RNAi, we analyzed the movement directions of GOPs in APP neurons. The results showed that

Dhc RNAi could rescue the DI of total dynamic GOPs in APP neurons compared with wild-type

neurons (Figures 4a and c-d; APP:0.87 ± 0.10; APP+*Dhc* RNAi: 0.52 ± 0.04). Further analyses

indicated that *Dhc* RNAi specifically rescued the directional equilibrium of mcGOPs (Figure 4e;

APP:1.06 ± 0.07; APP+*Dhc* RNAi: 0.62 ± 0.13) rather than scGOPs (Figure 4f; APP: 0.87 ± 0.09;

APP+*Dhc* RNAi: 0.76 ± 0.12). Similarly, *Dhc* RNAi could only rescue the retrograde displacement of

dynamic mcGOPs, but not for scGOPs in APP neurons (Figures 4g and h; APP: mcGOPs / scGOPs,

5.4 ± 0.6 / 6.0 ± 0.5 μm; APP+*Dhc* RNAi: mcGOPs / scGOPs, 9.1 ± 1.5 /5.8 ± 0.5 μm). Meanwhile,

Khc RNAi did not affect the alterations of GOPs dynamic induced by APP although it regulated GOPs

dynamic (Figures 4a-h). Neither *Dhc* RNAi nor *Khc* RNAi can recover the proportion of dynamic

GOPs in APP neurons as shown in Figure 4b. These data suggest that dynein is essential to the pattern

of directional movements of mcGOPs induced by APP.

Therefore, Kinesin and Dynein jointly regulate the direction of mcGOPs dynamic, of which,

dynein mediates the alterations of mcGOPs dynamic induced by APP. The alterations can only be

recovered via the loss of dynein.

Sunday driver (Syd) is required for abnormalities of mcGOPs dynamic and dendritic

arborization induced by APP

The anchoring of motor proteins to the Golgi apparatus is known to be regulated by adaptor

proteins (Papoulas, Hays, & Sisson, 2005; Yadav, Puthenveedu, & Linstedt, 2012), which

contribute to the various cellular functions. To identify the adaptor protein which was involved in

GOPs dynamic in APP neurons, we tested the Sunday driver (Syd), which is located in the post-Golgi

vesicles and interacts with both kinesin-I and dynein-dynactin complex (Bowman et al., 2000; Cavalli,

Kujala, Klumperman, & Goldstein, 2005).

To assess the role of Syd in neurons, we checked the localization of Syd and its function on the

density and dynamic of GOPs. First, we generated transgenic flies (UAS::EGFP-Syd) to observe the

location of Syd protein in neurons by fusing EGFP with *Syd* cDNA. After the expression of EGFP-Syd

in C3da neurons, it was identified in soma and dendrites as puncta and colocalized with Golgi units in

soma and dendrites (Figures 5a and b). The density of GOPs was then measured from Syd knockdown

(*Syd*-RNAi) and overexpression (OE). Both led to a significant reduction of GOPs density in the

dendrites (Figure 5c; density per 100 μm, WT: 9.0 ± 0.6; *Syd* OE: 5.4 ± 0.4; *Syd* RNAi: 7.3 ± 0.4).

Further, the properties of GOPs dynamic were examined with *Syd*-RNAi. Results revealed a

decline, including proportion of dynamic GOPs (Figures 5d and e; WT: 76.9% ± 2.5%; *Syd* RNAi:

55.1% ± 2.7%), and the direction index (DI) of dynamic GOPs (Figures 5f and g; WT: 0.49 ± 0.07;

Syd RNAi: 0.32 ± 0.02). Notably, those behaviors were dependent on changes of the dynamic of

mcGOPs rather than scGOPs (Figures 5h and i; WT: mcGOPs / scGOPs, 0.58 ± 0.11 / 0.53 ± 0.07; *Syd*

RNAi: mcGOPs / scGOPs, 0.29 ± 0.03 / 0.67 ± 0.13), which were consistent with those in *Dhc* RNAi

(Figures 3d-f). In addition, there were no alterations of the displacements of mcGOPs and scGOPs in

Syd RNAi (Figures 5j and k; WT: anterograde / retrograde, 4.7 ± 1.3 / 9.1 ± 1.1 μm in mcGOPs and

anterograde / retrograde, 6.2 ± 1.2 / 9.5 ± 1.0 μm in scGOPs; *Syd* RNAi: anterograde / retrograde, 5.3

± 0.6 / 7.8 ± 0.6 μm in mcGOPs and anterograde / retrograde, 5.5 ± 1.3 / 7.7 ±1.3 μm in scGOPs).

Therefore, Syd plays a critical role in regulating the directional equilibrium of GOPs dynamic,

especially for mcGOPs.

To determine the contribution of Syd on the alterations of GOPs dynamic induced by APP, we

examined the behaviors of dynamic GOPs in APP neurons with knockdown of *Syd*. This showed that

the directional equilibrium for dynamic GOPs was recovered in APP neurons where *Syd* was knocked

down (Figures 5f and g; APP: 1.06 ± 0.13; APP+ *Syd* RNAi: 0.57 ± 0.04). Besides, this rescue

primarily occurred in mcGOPs but not scGOPs (Figures 5h and i; APP: mcGOPs / scGOPs, 1.25 ±

0.24 / 1.01 ± 0.13; APP+ *Syd* RNAi: mcGOPs / scGOPs, 0.51 ± 0.07 / 0.76 ± 0.07). However, *Syd*

RNAi could not rescue the percentage of dynamic GOPs (Figure 5e; APP: 60.6% ± 2.3%; APP+ *Syd*

RNAi: 51.0% ± 3.3%) and the retrograde displacements of GOPs (Figures 5j and k; APP: mcGOPs /

scGOPs, 5.2 ± 0.7/ 5.2 ± 0.6 μm; APP+*Syd* RNAi: mcGOPs / scGOPs, 6.1 ± 0.7 / 5.4 ± 0.5 μm). Taken

together, Syd is required for the abnormal dynamic behaviors of GOPs induced by APP, especially the

equilibrium state of directional mcGOPs dynamic in dendrites.

Finally, to clarify the role of Syd on APP-induced dendritic pathology, we observed dendritic

morphology in APP neurons with Syd downregulation. Overall, the loss of Syd partially rescued the

dendrites phenotype in APP neurons (Figure 6a). In detail, *Syd*-RNAi recovered the number of total

branch points (Figure 6c; WT: 12.3 ± 0.3; *Syd* RNAi: 18.5 ± 1.1; APP: 10.3 ± 0.8; APP + *Syd* RNAi:

11.4 ± 0.8) and the dendritic complexity (Figure 6d; WT: 3.7 ± 0.3; *Syd* RNAi: 8.5 ± 1.0; APP: 2.1 ±

0.4; APP + *Syd* RNAi: 3.6 ± 0.4). However, Syd knockdown did not rescue the total dendrite length in

APP neurons (Figure 6b; WT: 1,852.3 ± 48.9 μm; *Syd* RNAi: 2,044.6 ± 80.5 μm; APP: 1,556.6 ± 78.4;

APP + *Syd* RNAi: 1,563.8 ± 80.0 μm), and had no recovery on spike density (Figure 6e; density per

100 μm: 30.6 ± 0.9 in WT; 17.0 ± 1.0 in *Syd* RNAi; 15.5 ± 2.1 in APP;13.2 ± 0.8 in APP + *Syd* RNAi).

Thus, these findings suggest that Syd mediates APP-induced dendritic branching defects by regulating

the dynamic behaviors of mcGOPs.

Discussion

In this study, we investigate the cellular mechanism of dendritic pathology in AD by ectopically

expressing APP in the C3da neurons of *Drosophila*. The findings suggest that dendritic APP is

transported through GOPs and modifies the dynamic profiles of GOPs. GOPs-resident APP breaks the

balance of anterograde/retrograde movements of mcGOPs maintained by dynein/Syd complex,

resulting in abnormalities of dendritic branching.

Here we show that GOPs are involved in APP transportation, and are regulated by APP entry.

Previous studies have found that APP secretases such as ADAM10 and BACE-1 exist in GOPs (Das

et al., 2013; Das et al., 2016; Saraceno et al., 2014). These suggest that GOPs are involved in the

trafficking and processing of APP. Our results show that GOPs co-localize and co-traffick with APP

puncta, which confirm that GOPs serve as the carriers of APP trafficking in dendrites. Furthermore, as

the local microtubule-organizing centers (Ori-McKenney et al., 2012) and secretory stations (Horton

& Ehlers, 2003; Pierce et al., 2001; Valenzuela & Perez, 2015) in the dendrites, GOPs play critical

roles in dendrite growth and neurodegenerative diseases (Chung et al., 2017; Lin et al., 2015). Here,

we observe that APP induces dendritic defects and severe GOPs deficiencies, including abnormal

density, distribution, and dynamic. Collectively, this indicates the involvement of GOPs in neuronal

degeneration.

The dynamic properties of GOPs are related to neurodegenerative diseases. One of the unique

features of GOPs is mobility. A previous study reported that the Parkinson's disease-associated protein

Lrrk regulates GOPs dynamic, which is manifested by inhibiting anterograde GOPs movements (Lin

et al., 2015). Here, we find that the equilibrium state of the movement direction is a key feature of APP

regulated GOPs dynamic. First, we observed that APP residing in GOPs can change their movements,

especially the balance of anterograde/retrograde movements. Second, we set up an index DI to quantify

the state of GOPs movements, which is equivalent to the ratio of anterograde to retrograde of GOPs

movements. We accessed the molecular mechanism of GOPs movements using DI. DI for mcGOPs is

downregulated or upregulated when *Dhc* or *Khc* is knocked down. It is worth noting that only the loss

of *Dh*c can recover APP-induced DI changes in mcGOPs. Therefore, we conclude that the change of

mcGOPs movements caused by APP is dynein-dependent.

We consider Syd to be a key molecule in APP regulating mcGOPs dynamic. Previous studies have

shown that Syd acts as a bidirectional adaptor interacting with both dynein and kinesin, and regulates

the intracellular transport by binding the *kinesin-1* or *dynein-dynactin* complex to cargos (Bowman et

al., 2000; Cavalli et al., 2005). In our study, both overexpression and knockdown of Syd decreased the

density of GOPs, and loss of it reduced dynamic GOPs and DI of GOPs in dendrites, which implicates

the regulation of Syd on the trafficking of GOPs. Moreover, *in vivo* imaging indicated that loss of Syd

could rescue DI of mcGOPs in APP neurons, suggesting that Syd is required to the abnormal mcGOPs

dynamic induced by APP. Besides, Syd contributes to the suppression of high-order dendrite arbors in

APP neurons. Taken together, the dynamic regulation of mcGOPs by Syd is the path for the dendritic

defects in APP neurons, providing new insight to understand the dendritic pathology mechanism in

AD.

Our results also demonstrate the functional differences between mcGOPs and scGOPs. After two

types of compartmental organization of GOPs in dendrites were reported (Zhou et al., 2014),

Valenzuela and Perez put forward a hypothesis that mcGOPs and scGOPs play different roles in the

secretory pathway with mcGOPs responsible for long-distance transport of proteins to dendritic branch

points and synaptic junctions, while scGOPs deliver the proteins to the destination, thereby promoting

local protein transport to the membrane (Valenzuela & Perez, 2015). Here, we found mcGOPs play

important role in neuronal development but scGOPs do not. Our results show that mcGOPs regulate

dendritic branching and complexity through the adaptor protein, Syd. Moreover, the results from the

loss function of *Dh*c and *Khc* also point that the movement mechanisms between mcGOPs and scGOPs

are different. Both *Dhc* and *Khc* can regulate the direction of movement of mcGOPs, while that of

scGOPs can only be regulated by *Khc*. Besides, this distinction is also confirmed by the fact that Syd

only regulates the movement direction of mcGOPs, but not scGOPs.

Materials and Methods

Fly Stocks and Transgenic Line

The stocks of flies published in this study included: GAL4[19-12] (Xiang et al., 2010); UAS-

GalT::TagRFPt and UAS-ManII::TagRFPt/GFP (Ye et al., 2007; Zhou et al., 2014); UAS-CD8::GFP

(Lee & Luo, 2001); UAS-APP::GFP[CG]; UAS-APP695[N-myc] and UAS-APP695-Swedish[N-myc]

(Fossgreen et al., 1998); UAS-Dhc64C[RNAi], UAS-Khc[RNAi] and Syd[RNAi] (Perkins et al., 2015). To

generate the UAS-Syd:: EGFP transgenic line, the *Syd* cDNA (#RE24340, BDGP) and *EGFP* cDNA

were amplified by PCR and transferred into the vector pJFRC2-10×UAS-IVS-mCD8-GFP (plasmid #:

26214, Addgene, Cambridge, MA). The construct was then injected into embryos of PBac{y[+]-attP-

3B}VK00033 to generate transgenic flies. All stocks and crosses were maintained in a 25°C incubator

with 40-60% humidity.

Dissection, Immunostaining, and Imaging

Third instar larvae were dissected in insect saline and fixed with 4% formaldehyde (PFA) for 40

minutes; permeabilized and washed in wash buffer (1X PBS containing 0.15% Triton X-100), and then

blocked in 5% normal serum for 30 minutes. After washing, larval fillets were incubated overnight at

4°C with the primary antibody and 3 hours with the second antibody at room temperature, avoiding

light. The antibodies included: chicken anti-GFP (1:5000, Aves Labs, Inc., Tigard, OR); Alexa647

anti-HRP (1:1000; Jackson ImmunoResearch Labs Inc., West Grove, PA); Alexa488 anti-chicken

(1:250; Jackson ImmunoResearch Labs Inc., West Grove, PA). The larval fillets were then dehydrated

by alcohol and transparentized with xylene before mounting on the slide with DPX (Sigma-Aldrich,

St Louis, MO) for imaging. The C3da neurons in the dorsal cluster of the fourth to sixth abdominal

segments of the larvae were imaged using a confocal microscope (FV1000, Olympus, Japan) under a

60x oil immersion objective (NA=1.42, Olympus, Japan). The images were captured at a resolution of

1024*1024 pixels and a z step of 0.4 μm in XYZ mode with sequential scanning. Images of neurons

requiring multiple fields were stitched together.

Morphological Analysis

The image analyses were performed by ImageJ (NIH). We measured the fluorescence intensity of

APP-GFP to quantify the distribution of APP in the C3da neurons (DeBoer, Dolios, Wang, & Sisodia,

2014). The images were corrected for uneven background illumination based on the "rolling ball"

algorithm (radius = 50 pixels). To determine the distribution of APP in axon and dendrites, we traced

the axon and dendritic arbors separately using line tool with 10-pixels-wide to fit the diameters of axon

and dendrites. Then we summed the fluorescence intensity separately in these regions and calculated

their proportion. The background intensity was subtracted (calculated in neighbouring area without

signal). As the axons of two adjacent C3da neurons (ddaA and ddaF) were closely bundled, we

measured both axons and halved the fluorescence intensity value for quantification.

For dendritic morphology analyses, we manually traced the dendritic branches by the NeuronJ

plugin (Meijering et al., 2004). The tracings were used to measure the length of the dendrite and count

the number of dendritic branch points. The number and length of spikes were similarly obtained.

To quantify the colocalization between Golgi and APP (or Syd), we calculated the proportion of

colocalized puncta. To count the number of total Golgi and total APP (or Syd) as well as the colocalized

ones, the Golgi and APP puncta were separately cataloged in ROIs manager, and the puncta that

overlapped with the puncta on the other channel were counted as a colocalization (as shown in yellow).

For the purposes of analyzing the compartmental organization of GOPs, we classified the colocalized

GalT::TagRFPt and ManII::GFP puncta as mcGOPs, while others as scGOPs.

In Vivo **Imaging**

In vivo imaging was performed as previously described (Zhou et al., 2014). Firstly, the 3rd instar

larvae were collected and anesthetized with ether. The orientation of larvae was adjusted until the

ventral trachea was located in the middle of the view field, the larvae were mounted in a drop of

halocarbon oil. Larvae were further immobilized by pressing a coverslip mounted on top of vacuum

grease. Time-lapse imaging of fluorescently tagged GOPs in C3da neurons was acquired using a

confocal microscope (FV1000, Olympus, Japan), 60x oil immersion objective. The settings for the two

channels were: ManII (488nm laser for GFP, emission filter: 500-525 nm) and GalT (543 nm laser for

RFP, emission filter: BA560IF). Multichannel time-lapse images were captured at zoom 2 with a

resolution of 512*512 pixels and collected in XYZT mode for 10 min at 6 s intervals.

Quantification of GOPs Dynamic

To quantify the dynamic behaviors of GOPs, we generated kymograph images (as X-T images

shown) using ImageJ. The kymograph images were generated in four steps: stabilization, straightening,

reslicing, and timing projection. First, images were stabilized using the Stabilizer plugin (Li, Feb. 2008)

to avoid fluctuation from larvae movements. We straightened the dendritic shafts to accurately measure

the distances of GOPs movements in the dendrites. The dendrites beyond the first branch points were

traced and straightened over the entire stack. The straightened images were resliced to reconstruct the

corresponding slices across the time series. Finally, the resliced images were time projected to produce

a kymograph of GOPs in the dendrites.

Further, we determined the number of dynamic GOPs and analyzed their properties, including

direction, displacement, and velocity. We compared the position of GOPs at the start and end of the

movement, and classified the movement as anterograde when the final position of GOPs puncta

relative to the starting point was towards dendritic ends, and retrograde when the final position of

GOPs was towards soma. The coordinates of GOPs trajectories on the kymograph were measured to

obtain displacement and velocity information during movements. Meanwhile, the mcGOPs and

scGOPs were distinguished based on colocalization analysis for ManII and GalT. Finally, the GOPs

trajectories were traced and generated combined tracks of GOPs by aligning the starting points of the

tracings, as shown in figures 3d, 4c, 5f. Besides, we defined the difference between dynamic and

stationary GOPs and classified them as stationary if they moved less than 0.5 μm in any direction,

otherwise as dynamic. To circumvent the influence of the physiological state of neurons and the

accuracy of calculation, only kymograph images with at least one moving GOPs were analyzed.

Statistical Analysis

Data were analyzed using a two-tailed Student's *t*-test throughout and p-value was used to

determine whether a significant difference existed. Significance levels were presented as follows: *, P

< 0.05; **, P < 0.01; ***, P < 0.001; n.s., no significance. Data were presented as mean ± SEM.

Data Availability

The data that support the findings of this study are available from the corresponding author upon

reasonable request. The source data underlying Figs. 1c, e-i, 2b-c, e-h, 3b, e-j, 4b, d-h, 5b-c, e, g-k, 6b-

f are provided as a source data file.

Acknowledgements

We thank Lei Xue (Tongji University, China), Bing Ye (University of Michigan, Ann Arbor), and

Yang Xiang (University of Massachusetts Medical School) for sharing fly stocks and and the Optical

Bioimaging Core Facility of WNLO-HUST for support with imaging systems. This work was

supported by the National Natural Science Foundation of China (Grant No. 31871027), Science Fund

for Creative Research Group of China (Grant No. 61721092), China Postdoctoral Science Foundation

founded project (2015T80788), Fundamental Research Funds for the Central Universities/HUST

(2017KFXKJC001, 2016YXMS034, 2014TS015), and the Director Fund of WNLO.

Conflict of interests

The authors declare no competing or financial interests.

Authors' contributions

W.Z. and J.C. conceived this project. W.Z., J.C. and QQ.D. designed the experiments. QQ.D.

conducted the experiments and analyzed the data. G.C. carried out molecular experiments. W.Z., J.C.

and QQ.D. wrote the paper.

References

Allan, V. J., Thompson, H. M., & McNiven, M. A. (2002). Motoring around the Golgi. *Nature cell*

biology, 4(10), E236-E242. doi:10.1038/ncb1002-e236

Alpar, A., Ueberham, U., Bruckner, M. K., Seeger, G., Arendt, T., & Gartner, U. (2006). Different

dendrite and dendritic spine alterations in basal and apical arbors in mutant human amyloid

precursor protein transgenic mice. *Brain Research, 1099*, 189-198.

doi:10.1016/j.brainres.2006.04.109

Arthur, A. L., Yang, S. Z., Abellaneda, A. M., & Wildonger, J. (2015). Dendrite arborization requires

the dynein cofactor NudE. *J Cell Sci, 128*(11), 2191-2201. doi:10.1242/jcs.170316

Baloyannis, S. (2002). The Golgi apparatus of Purkinje cells in Alzheimer's disease. *Neuropathology*

Back to the Roots, 1-10.

Baloyannis, S. (2009). Dendritic pathology in Alzheimer's disease. *Journal of the Neurological*

Sciences, 283(1), 153-157. doi:10.1016/j.jns.2009.02.370

Baloyannis, S. J. (2014). Golgi apparatus and protein trafficking in Alzheimer's disease. *Journal of*

Alzheimers Disease Jad, 42 Suppl 3, S153. doi:10.3233/JAD-132660

Bowman, A. B., Kamal, A., Ritchings, B. W., Philp, A. V., McGrail, M., Gindhart, J. G., & Goldstein,

L. S. (2000). Kinesin-dependent axonal transport is mediated by the sunday driver (SYD)

protein. *Cell, 103*(4), 583-594. doi:10.1016/S0092-8674(00)00162-8

Cavalli, V., Kujala, P., Klumperman, J., & Goldstein, L. S. (2005). Sunday Driver links axonal

transport to damage signaling. *The Journal of cell biology, 168*(5), 775-787.

doi:10.1083/jcb.200410136

Chung, C. G., Kwon, M. J., Jeon, K. H., Han, M. H., Park, J. H., Cha, I. J., . . . Kim, G. R. (2017).

Golgi outpost synthesis impaired by toxic polyglutamine proteins contributes to dendritic

pathology in neurons. *Cell reports, 20*(2), 356-369. doi:10.1016/j.celrep.2017.06.059

Das, U., Scott, D. A., Ganguly, A., Koo, E. H., Tang, Y., & Roy, S. (2013). Activity-induced

convergence of APP and BACE-1 in acidic microdomains via an endocytosis-dependent

pathway. *Neuron, 79*(3), 447-460. doi:10.1016/j.neuron.2013.05.035

Das, U., Wang, L., Ganguly, A., Saikia, J. M., Wagner, S. L., Koo, E. H., & Roy, S. (2016).

Visualizing APP and BACE-1 approximation in neurons yields insight into the amyloidogenic

pathway. *Nature neuroscience, 19*(1), 55. doi:10.1038/nn.4188

DeBoer, S. R., Dolios, G., Wang, R., & Sisodia, S. S. (2014). Differential release of beta-amyloid

from dendrite- versus axon-targeted APP. *J Neurosci, 34*(37), 12313-12327.

doi:10.1523/JNEUROSCI.2255-14.2014

Fossgreen, A., Bruckner, B., Czech, C., Masters, C. L., Beyreuther, K., & Paro, R. (1998).

Transgenic Drosophila expressing human amyloid precursor protein show gamma-secretase

activity and a blistered-wing phenotype. *Proc Natl Acad Sci U S A, 95*(23), 13703-13708.

doi:10.1073/pnas.95.23.13703

Gibbons, I. R. (1981). Cilia and flagella of eukaryotes. *J Cell Biol, 91*(3 Pt 2), 107s-124s.

doi:10.1083/jcb.91.3.107s

Golde, T. E., Estus, S., Younkin, L. H., Selkoe, D. J., & Younkin, S. G. (1992). Processing of the

amyloid protein precursor to potentially amyloidogenic derivatives. *Science, 255*(5045), 728-

730. doi:10.1126/science.1738847

Grueber, W. B., Jan, L. Y., & Jan, Y. N. (2002). Tiling of the Drosophila epidermis by multidendritic

sensory neurons. *Development, 129*(12), 2867-2878.

Gunjan, J., & Yanzhuang, W. (2015). Golgi defects enhance APP amyloidogenic processing in

Alzheimer's disease. *Bioessays, 37*(3), 240-247. doi:10.1002/bies.201400116

Hardy, J., & Selkoe, D. J. (2002). The Amyloid Hypothesis of Alzheimer's Disease: Progress and

Problems on the Road to Therapeutics. *Science, 297*(5580), 353-356.

doi:10.1126/science.1072994

Hardy, J. A., & Higgins, G. A. (1992). Alzheimer's disease: the amyloid cascade hypothesis. *Science,*

256(5054), 184-185. doi:10.1126/science.1566067

Hirokawa, N. (1998). Kinesin and dynein superfamily proteins and the mechanism of organelle

transport. *Science, 279*(5350), 519-526. doi:10.1126/science.279.5350.519

Horton, A. C., & Ehlers, M. D. (2003). Dual modes of endoplasmic reticulum-to-Golgi transport in

dendrites revealed by live-cell imaging. *Journal of Neuroscience, 23*(15), 6188-6199.

doi:10.1523/JNEUROSCI.23-15-06188.2003

Horton, A. C., Rácz, B., Monson, E. E., Lin, A. L., Weinberg, R. J., & Ehlers, M. D. (2005).

Polarized secretory trafficking directs cargo for asymmetric dendrite growth and

morphogenesis. *Neuron, 48*(5), 757-771. doi:10.1016/j.neuron.2005.11.005

Joshi, G., Chi, Y., Huang, Z., & Wang, Y. (2014). Aβ-induced Golgi fragmentation in Alzheimer's

disease enhances Aβ production. *Proceedings of the National Academy of Sciences of the*

United States of America, 111(13), 1230-1239. doi:10.1073/pnas.1320192111

Kelliher, M. T., Yue, Y., Ng, A., Kamiyama, D., Huang, B., Verhey, K. J., & Wildonger, J. (2018).

471   Autoinhibition of kinesin-1 is essential to the dendrite-specific localization of Golgi outposts.

472   *J Cell Biol, 217*(7), 2531-2547. doi:10.1083/jcb.201708096

473   Lee, T., & Luo, L. (2001). Mosaic analysis with a repressible cell marker (MARCM) for Drosophila

474   neural development. *Trends Neurosci, 24*(5), 251-254. doi:10.1016/s0166-2236(00)01791-4

475   Li, K. (Feb. 2008). The Image Stabilizer Plugin for ImageJ [Online].   Retrieved from

476   http://www.cs.cmu.edu/~kangli/code/Image_Stabilizer.html

477   Lin, C. H., Li, H., Lee, Y. N., Cheng, Y. J., Wu, R. M., & Chien, C. T. (2015). Lrrk regulates the

478   dynamic profile of dendritic Golgi outposts through the golgin Lava lamp. *J Cell Biol,*

479   *210*(3), 471-483. doi:10.1083/jcb.201411033

480   Luebke, J. I., Weaver, C. M., Rocher, A. B., Rodriguez, A., Crimins, J. L., Dickstein, D. L., . . . Hof,

481   P. R. (2010). Dendritic vulnerability in neurodegenerative disease: insights from analyses of

482   cortical pyramidal neurons in transgenic mouse models. *Brain Structure and Function, 214*(2-

483   3), 181-199. doi:10.1007/s00429-010-0244-2

484   Meijering, E., Jacob, M., Sarria, J. C., Steiner, P., Hirling, H., & Unser, M. (2004). Design and

485   validation of a tool for neurite tracing and analysis in fluorescence microscopy images.

486   *Cytometry A, 58*(2), 167-176. doi:10.1002/cyto.a.20022

487   Ori-McKenney, K. M., Jan, L. Y., & Jan, Y.-N. (2012). Golgi outposts shape dendrite morphology by

488   functioning as sites of acentrosomal microtubule nucleation in neurons. *Neuron, 76*(5), 921-

489   930. doi:10.1016/j.neuron.2012.10.008

490   Papoulas, O., Hays, T. S., & Sisson, J. C. (2005). The golgin Lava lamp mediates dynein-based Golgi

491   movements during Drosophila cellularization. *Nat Cell Biol, 7*(6), 612-618.

doi:10.1038/ncb1264

Perkins, L. A., Holderbaum, L., Tao, R., Hu, Y., Sopko, R., McCall, K., . . . Shim, H.-S. (2015). The

transgenic RNAi project at Harvard Medical School: resources and validation. *Genetics,*

201(3), 843-852. doi:10.1534/genetics.115.180208

Pierce, J. P., Mayer, T., & Mccarthy, J. B. (2001). Evidence for a satellite secretory pathway in

neuronal dendritic spines. *Current Biology Cb, 11*(5), 351-355. doi:10.1016/S0960-

9822(01)00077-X

Saraceno, C., Marcello, E., Di Marino, D., Borroni, B., Claeysen, S., Perroy, J., . . . Di Luca, M.

(2014). SAP97-mediated ADAM10 trafficking from Golgi outposts depends on PKC

phosphorylation. *Cell Death Dis, 5*, e1547. doi:10.1038/cddis.2014.492

Šišková, Z., Justus, D., Kaneko, H., Friedrichs, D., Henneberg, N., Beutel, T., . . . von der Kammer,

H. (2014). Dendritic structural degeneration is functionally linked to cellular

hyperexcitability in a mouse model of Alzheimer's disease. *Neuron, 84*(5), 1023-1033.

doi:10.1016/j.neuron.2014.10.024

Sisodia, S. S., & Price, D. (1995). Role of the beta-amyloid protein in Alzheimer's disease. *The*

FASEB Journal, 9(5), 366-370. doi:10.1096/fasebj.9.5.7896005

Terry, R. D. (1996). The pathogenesis of Alzheimer disease: an alternative to the amyloid hypothesis.

Journal of Neuropathology & Experimental Neurology, 55(10), 1023-1025.

doi:10.1097/00005072-199655100-00001

Terry, R. D., & Katzman, R. K. (1983). Senile dementia of the Alzheimer type. *Annals of Neurology,*

14(5), 497-506. doi:10.1002/ana.410140502

Vale, R. D., Reese, T. S., & Sheetz, M. P. (1985). Identification of a novel force-generating protein,

kinesin, involved in microtubule-based motility. *Cell, 42*(1), 39-50. doi:10.1016/s0092-

8674(85)80099-4

Valenzuela, J. I., & Perez, F. (2015). Diversifying the secretory routes in neurons. *Frontiers in*

Neuroscience, 9. doi:10.3389/fnins.2015.00358

Xiang, Y., Yuan, Q., Vogt, N., Looger, L. L., Jan, L. Y., & Jan, Y. N. (2010). Light-avoidance-

mediating photoreceptors tile the Drosophila larval body wall. *Nature, 468*(7326), 921-926.

doi:10.1038/nature09576

Yadav, S., & Linstedt, A. D. (2011). Golgi positioning. *Cold Spring Harb Perspect Biol, 3*(5).

doi:10.1101/cshperspect.a005322

Yadav, S., Puthenveedu, M. A., & Linstedt, A. D. (2012). Golgin160 recruits the dynein motor to

position the Golgi apparatus. *Dev Cell, 23*(1), 153-165. doi:10.1016/j.devcel.2012.05.023

Ye, B., Zhang, Y., Song, W., Younger, S. H., Jan, L. Y., & Jan, Y. N. (2007). Growing dendrites and

axons differ in their reliance on the secretory pathway. *Cell, 130*(4), 717-729.

doi:10.1016/j.cell.2007.06.032

Zhou, W., Chang, J., Wang, X., Savelieff, M. G., Zhao, Y., Ke, S., & Ye, B. (2014). GM130 is

required for compartmental organization of dendritic golgi outposts. *Current Biology, 24*(11),

1227-1233. doi:10.1016/j.cub.2014.04.008

Fig. 1. APP induces dendritic abnormalities in C3 dendritic arborization (da) ddaA neurons.

(a) The distribution of APP-GFP in C3da neuron of 3rd instar larva in *Drosophila*. The insets indicate dendrite (red) and axon (blue) respectively. Scale bar: 50 µm. **(b)** Magnified insets in **(a)** exhibit APP-GFP in the dendrite and axon. Scale bar for dendrite and axon: 5 µm. **(c)** The proportion of APP-GFP in the axon and dendrite. **(d)** The morphologies of wild-type (WT) and APP C3da ddaA neurons. Raw images (left) and tracings of dendritic branches (right, the numerous F-actin-based "dendritic spikes" of C3da neurons are not included). The red insets show magnified dendritic spikes. The red arrowhead points to the soma and blue to the branch points of the 4th order and up. Scale bar: 50 µm. **(e-j)** Quantification of dendritic morphologies in wild-type and APP neurons. **(e)** The total dendrite length, **(f)** the number of total branch points, **(g)** the number of branch points in different orders; **(h-i)** The characteristics of dendritic spikes: **(h)** density / 100 µm, **(i)** average length.

Fig.2. APP alters the compartmental organization and distribution of GOPs.

(a) Examples of APP (green) and Golgi complex (red) in the soma and dendrites. Golgi complex is labeled by GalT

(top, *trans*-Golgi) or ManII (bottom, *medial*-Golgi). Arrows indicate colocalized APP and Golgi in dendrites. **(b-c)**

Quantification of the proportion of colocalized APP and Golgi complex in soma and dendrites, co-localization as a

percentage of Golgi complex (**b**) or APP (**c**). **(d)** Examples of the compartmental organization of dendritic Golgi

outposts (GOPs) in wild-type and APP neurons. GOPs are labeled by ManII-GFP (green) and GalT-TagRFPt (red),

simultaneously. Arrows point to multi-compartment GOPs (mcGOPs). **(e-h)** Quantification of GOPs in wild-type and

APP neurons. **(e)** Density /100 μm, **(f)** the percentage of mcGOPs and scGOPs, **(g-h)** the distribution patterns of **(g)**

mcGOPs, and **(h)** scGOPs in dendrites. Scale bar: 2 μm.

a

WT | APP
GalT | GalT-alone | GalT-APP

Soma ←   → Distal

b

% of dynamic GOPs

- GalT-WT
- GalT-alone
- GalT-APP

c

WT   ManII/GalT   APP

Soma ←   → Distal   Soma ←   → Distal

d

WT   31.9%   APP   47.1%

n=49   n=49

-40   -20   0   20   40   -40   -20   0   20   40
Retrograde ←   → Anterograde   Retrograde ←   → Anterograde   (µm)

e   **f**   **g**   **h**

% of dynamic GOPs   Total GOPs   mcGOPs   scGOPs

***   ***   ***   **   **   **   **

42   29   Antero Retro   Antero Retro   Antero Retro

i   **j**

Displacement   Velocity

(17) (23) (20) (24)   (34) (28) (57) (34)

(33) (20) (35) (31)   (46) (28) (89) (30)
     **   **

mcGOPs scGOPs   mcGOPs scGOPs

- WT
- APP

Fig. 3. APP disrupts the equilibrium of anterograde/retrograde movements of GOPs.

(a) Examples of dynamic GOPs with/without APP**.** Kymograph of time-lapse imaging of GOPs showing GOPs (labeled by GalT-TagRFPt) dynamic without (red) or with APP-GFP (yellow, GalT-APP). **(b)** Bar charts showing the proportion of dynamic GOPs with/without APP. **(c)** Examples of the trajectories of GOPs in wild-type and APP neurons. Top: snapshots of the GOPs in the straightened dendrites. Bottom: kymograph of time-lapse imaging of GOPs. Dual-colors live imaging showing the compartmental organization of GOPs in dendrites, which are labeled by ManII-GFP (green, *medial*-Golgi) and GalT-TagRFPt (red, *trans*-Golgi). Green or red indicates a single-compartment GOP (scGOP) and yellow shows multi-compartment GOP (mcGOP). **(d)** Compound tracks of dynamic GOPs. Red lines are aligned along with displacement 0; puncta numbers and the proportion of anterograde GOPs are indicated within the boxes. **(e-j)** Quantification of GOPs dynamic in wild-type and APP neurons. **(e)** The proportion of dynamic GOPs. **(f-h)** The percentage of anterograde and retrograde movement of GOPs: **(f)** total GOPs, **(g)** mcGOPs, and **(h)** scGOPs. **(i)** Displacement and **(j)** velocity of mcGOPs and scGOPs. Scale bar: 5 μm / 2 min.

Fig. 4. *Dhc* RNAi rescues the alteration of mcGOPs dynamic induced by APP.

(a) Examples of the trajectories of GOPs dynamic in wild-type and APP neurons with RNAi knockdown of dynein

heavy chain (*Dhc*) or kinesin heavy chain (*Khc*). Top: snapshots of the GOPs in the straightened dendrites. Bottom:

kymograph of time-lapse imaging of GOPs in 10 min. Scale bar: 5 μm / 2 min. **(b)** Quantification of the proportion

of dynamic GOPs. **(c)** Compound tracks of dynamic GOPs. **(d-f)** Quantification of the direction index (DI) for the

GOPs dynamic. The direction index (DI) is calculated using the ratio of anterograde to retrograde movements. **(d)**

total GOPs, **(e)** mcGOPs, **(f)** scGOPs. **(g-h)** Quantification of the displacements of **(g)** mcGOPs and **(h)** scGOPs.

a

Soma | GalT | Syd | Merge
Dendrite

b % of colocalized Golgi or Syd

● Soma
● Dendrite

Golgi Syd

c GOPs density/100 µm

● Control
● Syd RNAi
● Syd OE

d

WT | APP
Syd RNAi | APP+Syd RNAi

ManII/GalT

Soma ← → Distal Soma ← → Distal

e % of dynamic GOPs

f

WT 30.6% n=56 | APP 47.7% n=54
Syd RNAi 23.8% n=52 | APP+Syd RNAi 35.0% n=55

-30 -15 0 15 30 -30 -15 0 15 30 (µm)
Retrograde ← → Anterograde Retrograde ← → Anterograde

0
2
4
6 (×10² s)

g Total GOPs

DI

h mcGOPs

DI

i scGOPs

DI

j mcGOPs

(15) (14) (27) (27)

Antero displacement (µm)

Retro

(33) (63) (27) (43)

k scGOPs

(19) (13) (17) (28)

Antero displacement (µm)

Retro

(36) (13) (28) (47)

● WT
● Syd RNAi
● APP
● APP+Syd RNAi

Fig. 5. Syd is required in alteration of mcGOPs dynamic induced by APP.

(a) Co-localization of Sunday driver (green, Syd-GFP) with Golgi complex (red, GalT-TagRFPt) in soma and dendrite.

Scale bar: 2 μm. **(b)** Quantification of the proportion of colocalized Syd and Golgi complex in soma and dendrites,

co-localization as a percentage of Golgi complex (Golgi) or Syd. **(c)** Bar chart showing the density of GOPs following

overexpression or knockdown of Syd. **(d)** Examples of the trajectories of GOPs movements in wild-type and APP

neurons in Syd knockdown. Top: snapshots of the GOPs in the straightened dendrites. Bottom: kymograph of time-

lapse imaging of GOPs in 10 min. Scale bar: 5 μm / 2 min. **(e)** Quantification of the proportion of dynamic GOPs

with loss of Syd in wild-type and APP neurons. **(f)** Compound tracks of GOPs dynamic. **(g-j)** Quantification of the

direction index (DI) for GOPs dynamic: **(g)** total GOPs, **(h)** mcGOPs, and **(i)** scGOPs. **(j-k)** Quantification of the

displacements of **(j)** mcGOPs and **(k)** scGOPs.

Fig. 6. Loss of Syd recovers the dendritic defects induced by APP.

(a) The dendritic morphologies of C3da ddaA neurons in wild-type and APP neurons following the loss of Syd. The

tracing of the microtubule-based dendritic branches (right). Arrowhead (red) points to the soma and arrows (blue) to

the branch points that are at the 4th order and above. Scale bar: 100 μm. **(b-f)** Quantification of the morphological

characteristics of C3da neuron. **(b)** Total dendrite length, **(c)** The number of branch points, **(d)** The number of branch

points in different orders. **(e-f)** Density **(e)** and length **(f)** of dendritic spikes.