Essentiality of c-di-AMP in *Bacillus subtilis*: Bypassing mutations converge in potassium and

glutamate homeostasis

Larissa Krüger [a], Christina Herzberg [a], Hermann Rath [b], Tiago Pedreira [a], Anja Poehlein [c], Jan Gundlach

[a], Rolf Daniel [c], Uwe Völker [b], Ulrike Mäder [b], and Jörg Stülke [a]*

[a] Department of General Microbiology, GZMB, Georg-August-University Göttingen, Germany

[b] Interfaculty Institute for Genetics and Functional Genomics, University Medicine Greifswald,

Greifswald, Germany

[c] Department of Genomic and Applied Microbiology, GZMB, Georg-August-University Göttingen,

Germany

Running head: Bypasses of cyclic di-AMP essentiality in *B. subtilis*

* Corresponding author:

Jörg Stülke, jstuelk@gwdg.de

ABSTRACT

In order to adjust to changing environmental conditions, bacteria use nucleotide second messengers to transduce external signals and translate them into a specific cellular response. Cyclic di-adenosine monophosphate (c-di-AMP) is the only known essential nucleotide second messenger. In addition to the well-established role of this second messenger in the control of potassium homeostasis, we observed that glutamate is as toxic as potassium for a c-di-AMP-free strain of the Gram-positive model bacterium *Bacillus subtilis*. In this work, we isolated suppressor mutants that allow growth of a c-di-AMP-free strain under these toxic conditions. Characterization of glutamate resistant suppressors revealed that they contain pairs of mutations, in most cases affecting glutamate and potassium homeostasis. Among these mutations, several independent mutations affected a novel glutamate transporter, AimA (Amino acid importer A, formerly YbeC). This protein is the major transporter for glutamate and serine in *B. subtilis*. Unexpectedly, some of the isolated suppressor mutants could suppress glutamate toxicity by a combination of mutations that affect phospholipid biosynthesis and a specific gain-of-function mutation of a mechanosensitive channel of small conductance (YfkC) suggesting the acquisition of a device for glutamate export. Cultivation of the c-di-AMP-free strain on complex medium was an even greater challenge because the amounts of potassium, glutamate, and other osmolytes are substantially higher than in minimal medium. Suppressor mutants viable on complex medium could only be isolated under anaerobic conditions if one of the two c-di-AMP receptor proteins, DarA or DarB, was absent. Also on complex medium, potassium and osmolyte toxicity are the major bottlenecks for the growth of *B. subtilis* in the absence of c-di-AMP. Our results indicate that the essentiality of c-di-AMP in *B. subtilis* is caused by the global impact of the second messenger nucleotide on different aspects of cellular physiology.

AUTHOR SUMMARY

Bacteria are exposed to constantly changing environmental conditions. In order to respond to these

changes, they use nucleotide second messengers to transduce external signals and translate them into

a specific cellular response. Among the repertoire of bacterial second messenger nucleotides, cyclic di-

AMP (c-di-AMP) stands out as it is the only second messenger that is essential for the bacteria that

produce it, including the Gram-positive model organism *Bacillus subtilis*. C-di-AMP plays a major role

in the control of potassium homeostasis, and we found that glutamate is toxic to a *B. subtilis* strain

lacking c-di-AMP to the same extent as potassium. These toxic conditions were the starting point for

an extensive suppressor analysis, which led to the identification of a novel glutamate transporter

(AimA). If the *B. subtilis* strain lacking c-di-AMP was cultivated on complex medium, the isolation of

suppressor mutants was only possible under anaerobic conditions and if either of the two c-di-AMP-

binding signal transduction proteins was absent. This suggests that these proteins are a major burden

for the cell on complex medium in their c-di-AMP free state. Our result underline the complexity of c-

di-AMP signaling and propose new directions for research.

KEYWORDS

Glutamate, potassium, *Bacillus subtilis*, cyclic di-AMP, ion homeostasis, osmoprotection, essential

genes

INTRODUCTION

To achieve appropriate responses to changing environmental conditions, all organisms use second messengers which are formed in response to an environmental cue. In bacteria, specific mono- and dinucleotides play a particularly important role as second messengers [1]. The common nucleotide second messengers cyclic AMP, cyclic di-GMP (c-di-GMP), and guanosine tetra/pentaphosphate ((p)ppGpp) are used to prioritize the utilization of carbon sources and to regulate virulence, to control the choice between sessile and motile lifestyles, and to downregulate essential cellular functions upon amino acid starvation, respectively [2 - 7]. The more recently discovered second messenger cyclic di-AMP is unique in several respects. First, this nucleotide is essential in many bacteria that produce it, suggesting that this molecule plays a role in central cellular activities. Essentiality of c-di-AMP has been observed in many Firmicutes, such as *Bacillus subtilis*, *Listeria monocytogenes*, *Streptococcus agalactiae*, and *Staphylococcus aureus*, but also in the genome-reduced bacterium *Mycoplasma pneumoniae* and in *Chlamydia trachomatis* [8 - 14]. Second, c-di-AMP is toxic if it accumulates in the cells, and in many bacteria this accumulation results in strongly increased stress sensititvity [9, 15 – 19]. Third, c-di-AMP can bind to proteins and the mRNA molecules encoding them to control both the expression and the activity of certain proteins, as in the case of the *B. subtilis* potassium transporters KtrAB and KimA [20 - 23]. Last, but not least, c-di-AMP is not intrinsically present in eukaryotic organisms, and the presence of the molecule is an indication for infection with pathogenic bacteria. Indeed, c-di-AMP is recognized by the human immune system. It binds the STING protein thus resulting the induction of an innate immune response [10].

The essentiality and toxicity of c-di-AMP for many bacteria have induced a great interest in understanding the reasons behind it. Classically, genes were regarded to be essential if they were absolutely needed for growth under optimal growth conditions, typically on complex medium at 37°C [24 - 26]. However, many genes are not described as being essential because there may be paralogues that encode very similar functions as has been described for the c-di-AMP producing diadenylate

cyclases in *B. subtilis* [8, 9]. In some cases, one essential function can be carried out by very different

proteins, and such genes are essential only under specific conditions, *i. e*. the absence of amino acids

from the medium makes the corresponding biosynthetic genes essential. These genes have been called

conditional essential genes. Finally, some genes may not be essential but their loss would cause a

substantial decrease of fitness for the cells, resulting either in the disappearance of such mutants or in

the acquisition of suppressor mutations. Such genes or proteins have been designated quasi-essential,

and the effect of their loss has been studied for the *B. subtilis* DNA topoisomerase TopA, the

endoribonuclease RNase Y, and the modulator of lipid biosynthesis, YqhY [27 - 29]. Very recently,

essential processes have been ranked on a global scale for the first time by the determination of the

speed of death of mutants [30].

Two lines of research have provided important insights into the reasons of c-di-AMP

essentiality. Studies to identify target molecules that interact with c-di-AMP demonstrated that many

proteins involved in potassium and osmolyte homeostasis – both transporters and their regulators –

are controlled by c-di-AMP to limit the intracellular concentrations of potassium and osmotically

compatible compounds to physiologically reasonable levels [11, 13, 23, 31 - 38]. In a complementary

approach, suppressor mutants of bacterial strains that either lack or accumulate c-di-AMP have been

studied. These analyses revealed that c-di-AMP is conditionally essential in *L. monocytogenes*, *S.*

aureus and *B. subtilis*. While the nucleotide is required for growth on complex media, it is dispensable

on minimal media [21, 39, 40]. For a *B. subtilis* mutant lacking c-di-AMP, growth was only possible on

minimal medium as long as the potassium concentration in the medium is low and glutamate is not

present [21, 41]. On complex medium, the lack of c-di-AMP leads to an accumulation of (p)ppGpp [39],

the second messenger that results in a global reprogramming of transcription, translation, and

metabolism upon amino acid starvation. Inactivation of the enzymes that produce (p)ppGpp restores

the viability of the strain lacking c-di-AMP on complex medium. The accumulation of (p)ppGpp leads

to the inability of the transcription factor CodY to bind to its targets and thus in deregulation of a large

regulon that includes many genes for osmolyte uptake and amino acid metabolism, and this

uncontrolled expression of the CodY regulon is responsible for the essentiality of c-di-AMP for *L.*

monocytogenes on complex media [39]. In *S. aureus*, c-di-AMP is dispensable for growth on complex

medium under anaerobic conditions or when the bacteria acquire suppressor mutations that affect

osmolyte or glutamate uptake [40, 42]. Both types of studies support the idea that the control of

potassium and osmolyte homeostasis is the central essential function of c-di-AMP in many bacteria

that produce it [43 - 45].

We are interested in c-di-AMP signaling in the model organism *B. subtilis*. In contrast to *S.*

aureus or *L. monocytogenes*, *B. subtilis* encodes three diadenylate cyclases that produce c-di-AMP, the

constitutively expressed enzymes CdaA and DisA, and the sporulation-specific enzyme CdaS [9, 46].

Mutants lacking both the *cdaA* and *disA* genes that encode the constitutive enzymes are not viable on

complex medium and tolerate only very low potassium concentrations and no glutamate in minimal

medium. In *B. subtilis*, c-di-AMP binds to two signal transduction proteins, DarA and DarB, to several

potassium, osmolyte and magnesium transporters, as well as to the riboswitch that controls the

expression of the high affinity potassium transporters KtrAB and KimA [20, 23, 47].

To get a better understanding of the global role of c-di-AMP for the physiology of *B. subtilis*,

we performed extensive suppressor analysis of strains lacking c-di-AMP that had adapted to tolerate

the presence of glutamate or that are viable on complex medium. These suppressor analyses were

complemented by a transcriptome analysis to understand the roles of glutamate and potassium as well

as of c-di-AMP for global gene expression in *B. subtilis*. Our results indicate that the adaptation to

either glutamate or complex medium usually requires the simultaneous presence of multiple

suppressor mutations that often affect potassium uptake, osmolyte uptake, or glutamate metabolism.

Moreover, a mechanosensitive channel and phospholipid biosynthesis play an important role in the

sensitivity of a strain lacking c-di-AMP to glutamate. Strikingly, this study identified the amino acid

transporter YbeC (renamed AimA, <u>a</u>mino acid <u>im</u>porter <u>A</u>) as the major glutamate transporter in *B.*

subtilis. On complex medium, the c-di-AMP-binding signal transduction proteins DarA and DarB seem

to be toxic in the absence of c-di-AMP, and only in their absence we could isolate suppressor mutants.

The obtained results shed light on the complexicity of c-di-AMP essentiality and open new directions

of research.

RESULTS

Growth of a strain lacking c-di-AMP in the presence of glutamate requires distinct combinations of

suppressor mutations

A *B. subtilis* strain lacking all three diadenylate cyclases is unable to grow at potassium concentrations

of 5 mM or higher or in the presence of glutamate (even at low potassium concentrations) [21, 41]. In

order to get more comprehensive insights into the toxicity of glutamate for the c-di-AMP-free strain

GP2222 (designated Δ*dac* from here on), we decided to isolate and study a larger set of suppressor

mutants that tolerate the presence of glutamate. For this purpose, we plated the Δ*dac* strain [21] on

MSSM medium containing glutamate (1% w/v) and a low concentration of potassium (0.1 mM KCl).

While suppressor mutants appear readily after overnight cultivation when potassium toxicity was the

growth-limiting problem [21], the toxicity of glutamate could only be overcome after several days.

Moreover, the number of suppressor mutants was strongly reduced as compared to selection in the

presence of potassium and ammonium as the nitrogen source. After four days at 37°C, 14 colonies

could be isolated (see Fig. 1, Δ*dac*). These strains were subjected to whole genome sequencing to

identify the responsible mutations (Table 1). The long time for suppressor mutations to appear can be

explained by each mutant harboring multiple mutations. In addition, this explains why the overall

number of suppressors was rather low.

Table 1. Characterization of suppressor mutants derived from the Δ*dac* strain GP2222 in MSSM

minimal medium with glutamate and 0.1 mM KCl. [1]

Strain	Genotype	Ammonium, 0.1 mM KCl	Ammonium, 5 mM KCl	Glutamate, 0.1 mM KCl	Glutamate, 5 mM KCl	LB
GP2840[2]	*ktrC* (fs) *plsC*-A61V *ndhF* (fs) *sigD*-G88D	+++	+++	+++	+++	-
GP3079	Δ(*abh-kinC-ykqA-ktrC*) *plsC*-L48P *yfkC*-R42W *fliP* (fs) *hbs*-I32T *ndoA*-G6S	++	+++	+++	+++	+++
GP3077	*ktrC* (fs) *aimA/glpQ* (trunc) *ybcC* (fs)	+++	+++	+++	+++	-
GP2841	*gudB*+ *accC*-H298Y *fliG* (fs)	+	+	++	+	-
GP2842	*nhaK*-S187F *aimA* (fs)	+++	+++	++	+	-
GP3078	*plsC*-A61V *aimA*-T374M *ktrB*-G313D *sigD*-G88D	+++	+	+++	-	-
GP2849	*ybcC*-K722L, I724F *aimA* (fs)	+++	+++	+++	+++	-
GP2837	*plsC*-L48P *yfkC*-R42W *fliP* (fs)	+++	+++	+++	+++	+++
GP3099	*plsC*-L48P *yfkC*-R42W *fliP* (fs) *cotO*-I34F	+++	+	+++	-	-
GP3464	*plsC*-P110S *flhA* (fs)	++	++	++	-	++
GP2224	*yqeG* (fs) *aimA* (fs) *yqeV* (fs)	+++	+++	+++	+++	-
GP2838	*ndhF*-V237E, M241L, H242L, A243D, G244C, L245F ΔICEBs1 (*int-yddM*) *yoaI* (fs)	+++	++	++	++	++
GP3473	*ndhF* (fs) *aimA* (fs) *gudB*+ *fliH* (fs) *yuiE*-G37V	++	++	++	-	-
GP3474	*ndhF* (fs) *aimA* (fs) *ykpA* (fs) *ktrC* (fs) *fliK* (fs)	++	+++	+++	+++	-

GP2839	fliK (fs) rpoB-Q109P rbsB-R175L	++	++	++	-	-

As observed previously [41], three strains were affected in *ktrC*. In one of these strains

(GP3079), the genomic region from *abh* to *ktrC* was deleted resulting in reduced uptake of potassium

via KtrCD. Similarly, strain GP3078 has a mutation affecting the high-affinity potassium transporter

KtrB, and GP2842 also carries a mutation (*nhaK*) that affects potassium homeostasis. Interestingly, the

same mutation (NhaK-S187F) had been isolated in a suppressor screen of the Δ*dac* strain in the

presence of 5 mM potassium [21] indicating that the mutant protein has an increased activity in

potassium export.

As mentioned above, the suppressor strains usually had multiple mutations. Since we never

experienced the systematic acquisition of multiple mutations in prior suppressor screens [15, 27, 48 -

50], it seems likely that more than one mutation is required to achieve the suppression. Since the

selective pressure in our experiment was directed towards tolerating otherwise toxic glutamate, we

expected that the suppressor mutants would contain mutations affecting glutamate metabolism.

Indeed, the á priori cryptic glutamate dehydrogenase was decryptified by the deletion of nine base

pairs in two mutants (GP2841 and GP3473), thus allowing glutamate degradation [51, 52]. Moreover,

seven independently isolated mutants carried mutations affecting an amino acid transporter, YbeC.

This protein has recently been identified as the major serine transporter of *B. subtilis* [53], and it is

tempting to speculate that YbeC is the major glutamate transporter as well (see below). However,

seven of the isolated mutants did not carry any mutation that is easily linked to glutamate homeostasis.

The previously reported suppressor mutant GP2840 had mutations affecting PlsC, NdhF, and SigD (acyl-

ACP:1-acylglycerolphosphate acyltransferase, NADH dehydrogenase, Sigma factor σ^D, respectively) in

addition to the *ktrC* mutation [41]. Strikingly, *plsC* and *ndhF* are affected in six and four different

suppressor mutants respectively. Several mutants carried mutations affecting motility: In addition to

two *sigD* mutants, seven additional mutants carried mutations in flagellar genes that on one hand

depend on a functional SigD for their expression, and on the other hand are required for SigD activity

[54]. In addition, three of the 14 different independent suppressor mutants carry the same point

mutation in the *yfkC* gene, encoding a mechanosensitive channel that releases ions from the cell [55,

56]. This mutation results in a substitution of Arg-42 by a Trp residue.

We then tested the ability of the suppressor mutants to grow on minimal medium at low and

high potassium concentrations with ammonium or glutamate as the nitrogen source. Moreover, the

ability to grow on a complex medium was tested (see Table 1, S1 Fig.). Since the mutants were isolated

in the presence of glutamate and 0.1 mM potassium, it is not surprising that all strains grew on the

corresponding medium. Moreover, most of the strains were viable with ammonium as the single

nitrogen source irrespective of the potassium concentration. Most of the mutants were also capable

of growing at the elevated potassium concentration in the presence of glutamate. However, only four

of the 15 tested strains were viable on complex medium. Two of them grew well on LB and these

strains had the identical combination of mutations in *yfkC* and *plsC* (see Table 1, see below). This

suggests that the adaptation to complex medium requires more than resistance to potassium and

glutamate.

Stepwise adaptation of a strain lacking c-di-AMP to growth on complex medium

Most of the suppressor mutants that tolerate the presence of glutamate and potassium were unable

to grow on complex medium under standard laboratory conditions. The standard definition of

essentiality refers to optimal growth conditions, *i. e*. to a complex medium and growth at 37°C in the

presence of oxygen for *B. subtilis* [25, 26]. This prompted us to investigate the growth requirements of

the Δ*dac* mutant GP2222 more rigorously. First, we attempted to isolate suppressor mutants derived

from GP2222 under standard conditions as defined above. However, we never succeeded in isolating

a single suppressor mutant. A recent study on c-di-AMP essentiality in *S. aureus* revealed that the

nucleotide is dispensable under anaerobic conditions [40]. Therefore, we tried to isolate suppressor

mutants of GP2222 in the absence of oxygen. While the wild type strain *B. subtilis* 168 grew under

these conditions, again no suppressor mutants of the Δ*dac* strain could be isolated (see Fig. 2).

In *B. subtilis*, c-di-AMP binds two signal transduction proteins, DarA and DarB [23, 47]. In a

suppressor screen using a *L. monocytogenes* strain lacking c-di-AMP, several mutations in the

corresponding genes were observed [39]. We therefore constructed strains that lack both c-di-AMP

and either the *darA* or the *darB* gene, and used these strains to isolate suppressors on complex

medium under aerobic and anaerobic conditions. Again, no suppressors could be obtained under

aerobic conditions; however, we detected one and two suppressors for the Δ*dac* Δ*darA* and Δ*dac*

Δ*darB* mutants, respectively, under anaerobic conditions (see Fig. 2). Whole genome sequencing

revealed that the mutation present in the Δ*dac* Δ*darA* mutant changed Gly-11 of KtrC to a cysteine

residue. The suppressor mutants derived from the Δ*dac* Δ*darB* mutant had a frameshift in *ktrC*

resulting in the expression of a truncated KtrC protein or a mutation affecting the permease domain

OpuAB of the OpuA glycine betaine transporter [57], respectively (Fig. 2). These results strongly

support the idea that the accumulation of potassium or of osmoprotective compatible solutes is the

reason for the essentiality of c-di-AMP [39, 40, 43].

An initial characterization of these suppressor mutants revealed that they were viable under

anaerobic conditions, but not in the presence of oxygen. Therefore, we used the three strains for a

further round of suppressor isolation. Indeed, we were able to obtain suppressors that tolerate the

presence of oxygen for all three mutants (Fig. 2, S3 Fig.). Whole-genome sequencing revealed that the

initial suppressors with defective KtrC proteins had all acquired very specific mutations in the *kimA*

gene, encoding the major high-affinity potassium transporter KimA. In the mutant isolated from the

darA ktrC mutant, the absolutely conserved glycine in transmembrane helix 3 of KimA was deleted.

Helix 3 is essential for potassium binding of KimA [58]. All four independent suppressors isolated from

the Δ*dac* Δ*darB ktrC* mutant had a deletion of 11 amino acids in the transmembrane helix 9. Based on

the high conservation of the deleted residues, we conclude that the mutant KimA proteins were

inactive or less active than the wild type protein. Again, the *kimA* suppressor mutations were

accompanied by other mutations (see Fig. 2). In the suppressor mutant GP3010 derived from the Δ*dac*

Δ*darA ktrC* mutant, we found a 155 bp deletion that results in the truncation the global transcription

regulator for catabolic genes and operons, CcpA, and in the conversion of the start codon of *motP*,

encoding the Na$^+$-coupled MotP flagellar stator protein to a stop codon. For the Δ*dac* Δ*darB ktrC*

suppressor mutant, two aerobic suppressors were subjected to whole genome sequencing. In one

strain (GP2845), we found a point mutation in the *darA* gene that results in a substitution of the

conserved Val-100 residue by an alanine in the DarA protein. The second suppressor (GP3094) carried

mutations affecting the CodY transcription factor (G146V substitution) and the unknown SPβ phage

protein YonD (D255H). In two additional suppressor mutants, we found identical frameshift mutations

in *darA* that result in a truncation of the DarA protein after amino acid 64. Finally, we also characterized

an aerobic suppressor (GP2847) derived from the anaerobic Δ*dac* Δ*darB opuAB* suppressor mutant. In

this case, again a potassium channel subunit (KtrB) was inactivated as the result of a frameshift

mutation. Moreover, this strain carried a frameshift mutation in the essential *accA* gene encoding a

subunit of acetyl-CoA carboxylase that resulted in the formation of a truncation after amino acid 108

and mutations in the genes encoding the putative methylenomycin A exporter Mmr and the tRNA

methylthiotransferase YmcB. These results again highlight the importance of potassium homeostasis

for the adaptation of strains lacking c-di-AMP.

Taken together, a *B. subtilis* strain lacking c-di-AMP is only able to grow under standard

laboratory conditions if it acquires several suppressing mutations. The results of the two suppressor

screens in the presence of glutamate and directly on complex medium demonstrate a distinct selective

pressure under both conditions resulting in different sets of mutations that allow aerobic growth on

complex medium. The importance of the initial loss of the c-di-AMP receptor proteins DarA and DarB

for the direct adaptation to complex medium demonstrates that these proteins interfere with the

growth of *B. subtilis* if they are present in the ligand-free apo-state. Indeed, apo-DarB was recently

shown to trigger growth-inhibitory (p)ppGpp synthesis by the Rel protein [59, 60]. For DarA, this is the

first time that a phenotype for the corresponding mutant is reported. The repeated isolation of

mutations affecting potassium uptake even in iterative cycles of suppressor screens highlights the

severe toxicity of potassium for *B. subtilis* in the absence of c-di-AMP.

A global transcription study of the regulatory effects of potassium, glutamate, and c-di-AMP

In order to obtain deeper insights into the physiological role of c-di-AMP, we decided to study the

impact of a lack of this second messenger on global gene expression by a transcriptome analysis as a

complementary approach to the suppressor analysis. Since c-di-AMP has been implicated in the

interplay of potassium and glutamate homeostasis, we analysed the global gene expression for the

wild type strain *B. subtilis* 168 at low (0.1 mM) and high (5 mM) potassium concentrations and in the

presence of ammonium and glutamate as the nitrogen source (S1 Supporting Information). Moreover,

we used the c-di-AMP free strain (Δ*dac*) GP2222 as well as the isogenic suppressor mutant GP2223

that is able to grow at 5 mM potassium (see Fig. 3 for the regulatory network and S1 Table). Strikingly,

expression of several genes involved in fermentation and respiration was strongly reduced in the

absence of c-di-AMP. The *ldh-lctP* and *cydABCD* operons encoding lactate dehydrogenase and the

lactate exporter as well as the terminal quinole oxidase, are repressed by the NADH-responsive

transcription factor Rex [61, 62]. The about 100-fold repression of these genes in the absence of c-di-

AMP suggests that the NADH levels are reduced in the mutant. This conclusion is also supported by

the increased expression (seven-fold) of the *ndhF-ybcC* operon (see S1 Table) which codes for a NADH

dehydrogenase and thus also contributes to the oxidation of $NADH_2$ to NAD [63]. Importantly, the *ndhF*

gene and its downstream gene *ybcC* were affected in several of the suppressor mutants suggesting

that these mutations counteract the increased expression of the conserved *ndhF-ybcC* operon. In

addition to genes of the Rex regulon, genes of the SigO regulon that are involved in the response to

acid stress [64] and several competence genes were severely reduced in the Δ*dac* mutant GP2222. The

strong reduction of competence gene expression in response to the absence of c-di-AMP suggests that

this second messenger might be required for genetic competence and transformation in *B. subtilis*. To

test this hypothesis, we attempted transformation of strain GP2222 using chromosomal DNA. No

transformants were obtained, whereas the control strain 168 was efficiently transformable. Thus, the

lack of c-di-AMP results in a loss of genetic competence as a result of reduced competence gene

expression.

It has been shown before that the expression of the high affinity potassium uptake systems

KtrAB and KimA is repressed in the presence of potassium via a c-di-AMP sensitive riboswitch [20 - 22].

In agreement with those results, these genes were most strongly repressed by potassium in our

analysis in the wild type strain 168. The expression of the *kimA* transporter gene was 112- and 17-fold

repressed at 5 mM potassium in the presence of glutamate or ammonium, respectively; whereas the

expression of the *ktrAB* channel operon was reduced 29- and 4-fold, respectively (see S2 Table). As

expected, the expression of both *kimA* and *ktrAB* is increased in the absence of c-di-AMP; however,

both genes are still repressed by potassium (eight- and five-fold, respectively), indicating a second, c-

di-AMP independent mechanism of transcriptional regulation of these potassium transport systems.

The particular importance of potassium for the cell and for KimA as a major high-affinity transporter is

highlighted by the fact that the *kimA* gene belongs to the most strongly expressed genes at 0.1 mM

potassium (position #50 in the presence of glutamate), and that it is even the most strongly expressed

transporter in *B. subtilis* under these conditions (see

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE156738 for the full data set).

Taken together, the global transcription analysis supported the idea that c-di-AMP is involved

in multiple cellular functions. Moreover, the analysis confirmed a particular role of genes involved in

controlling NADH homeostasis, as already suggested by the suppressor analysis and by the differential

effects of aerobic and anaerobic conditions to allow suppressor formation of the strains lacking c-di-

AMP.

The mechanosensitive channel YfkC contributes to the glutamate sensitivity of a c-di-AMP free strain

Three of the suppressor mutants isolated in the presence of glutamate carry mutations affecting the

mechanosensitive channel of small conductance, YfkC (see Table 1). Strikingly, all three independently

isolated mutants carry precisely the same mutation that results in a R42W substitution in the protein

(see S4 Fig.). This residue is located at the end of the first trans-membrane helix [65] and the mutation

may affect the relative positioning of the trans-membrane helices 1 and 2. Another striking feature is

that this mutation is in all three cases accompanied by point mutations in the *plsC* gene (see below).

The presence of the same amino acid substitution in YfkC in all three strains suggested that this

mutation resulted in a gain of function rather than in a loss of YfkC activity (from here on, we designate

the mutant allele *yfkC**, the protein YfkC*). To address this question, we constructed Δ*dac* Δ*yfkC* and

Δ*dac yfkC** strains lacking all diadenylate cyclases and assayed the growth of these strains on minimal

medium containing glutamate or ammonium as the nitrogen source at high and low potassium

concentrations (see Fig. 4). Both strains were viable when cultivated on plates containing ammonium

and 0.1 mM potassium whereas they were unable to grow in the presence of glutamate or at the high

potassium concentration. These observations indicate that neither the deletion of the *yfkC* gene nor

*yfkC** mutation alone were sufficient to allow suppression of the glutamate and potassium toxicity of

the Δ*dac* mutant. To address this issue further, we attempted the isolation of suppressor mutants from

the Δ*dac* strain either lacking *yfkC* or carrying the *yfkC** allele in the presence of glutamate. No

suppressor mutants could be isolated from the Δ*dac* Δ*yfkC* mutant, whereas two suppressor mutants

were obtained from the Δ*dac yfkC** mutant (see Fig. 1). This observation supports the idea that the

specific R42W substitution of YfkC* is a prerequisite for the adaptation of the c-di-AMP-free strain to

growth in the presence of glutamate. Whole genome sequencing of the two glutamate-resistant

suppressor mutants identified a mutation in *accA* in strain GP2850 (see below). This mutation supports

the requirement of paired mutations affecting *yfkC* and lipid biosynthesis. The second suppressor

strain, GP3105, carries mutations affecting the amino acid transporter YbeC, the CarB subunit of the

carbamoyl-phosphate synthetase, the endonuclease MutL and the polyketide synthase PksJ. Among

these mutations, the frameshift in the *ybeC* gene is likely the most relevant (see below). These

suppressor mutants were also tested for their growth in the presence of glutamate or increased

potassium concentrations (see Fig. 4). Both were viable in the presence of glutamate irrespective of

the potassium concentration. However, the growth in the presence of glutamate was much better in

the case of the *yfkC* plsC* (L48P) mutant GP2837 (Fig. 4). Taken together, these results suggest that

the *yfkC** mutation in combination with the lipid biosynthetic or *ybeC* mutations is rather specific for

the protection of the Δ*dac* mutant from glutamate toxicity. In the growth assay, we also included the

Δ*dac nhaK* (S187F) mutant that has been isolated for resistance to potassium [21]. As expected, this

strain grew at 5 mM potassium but was as sensitive to glutamate as the original Δ*dac* mutant, however

we observed the rapid formation of suppressor mutants in the presence of glutamate (see Fig. 4, see

below). As the NhaK allele used in this experiment is thought to exhibit increased potassium export

activity [21], it is tempting to speculate the mutant mechanosensitive channel YfkC* is able to export

glutamate to overcome the toxicity of this amino acid.

A reduction of phospholipid biosynthesis helps to overcome the lack of c-di-AMP

The analysis of suppressor mutants that were viable in the presence of glutamate or on complex

medium identified several mutations that affect proteins involved in phospholipid biosynthesis, *i. e.*

AccA, AccC, and PlsC (see Fig. 1, Fig. 2, Table 1). The *accA* and *accC* genes encode subunits of the

essential acetyl-CoA carboxylase. Unexpectedly, a frameshift had occurred in the *accA* gene of strain

GP2847. Since malonyl-CoA synthesis is the essential first step in fatty acid and phospholipid

biosynthesis, it is likely that this mutation can be intrinsically suppressed and thus does not result in a

complete loss of AccA synthesis. The *accA* mutation isolated in the Δ*dac yfkC** mutant results in an

A229V substitution in the active site of the enzyme [66, 67] (see S5 Fig.). For AccC, the mutation

(H298Y) is located in the highly conserved active site motif of the enzyme (S5 Fig). The localization of

these mutations strongly suggests that the initial step of lipid biosynthesis, the formation of malonyl-

CoA, is severely impaired in the suppressor mutants.

PlsC is the acyl-ACP:1-acylglycerolphosphate acyltransferase that catalyzes the formation of

phosphatidic acid in the synthesis of phospholipids. Interestingly, the *plsC* gene is essential for the

growth of *B. subtilis*. In agreement with this essentiality, none of the three different mutations in the

six independent mutants results in a frame shift or a deletion. Instead, residue L48 is replaced by a

proline in three independently isolated mutants, A61 by a valine in two mutants, and P110 by serine

in one mutant. All three residues are located in functionally important regions of the protein. L48 is

close to the HXXXXD motif, which is part the active site of the enzyme. Similarly, A61 and P110 are in

very close proximity to the active center of the enzyme [68] (S6 Fig). The location of all three mutations

in the immediate vicinity of the active center as well as the fact that the mutations affect conserved

residues suggest that they result in a reduced PlsC activity, and thus in reduced phospholipid

biosynthesis.

The requirement of the Δ*dac yfkC** mutant for an additional mutation reducing lipid

biosynthesis to allow growth in the presence of glutamate suggests that the membrane properties are

important for controlling cellular glutamate homeostasis. This conclusion is supported by the well-

established functional interaction between mechanosensitive channels and lipid molecules [69, 70]. It

is tempting to speculate that the combination of altered lipid composition and the mutation in the

trans-membrane helix of YfkC results in a change of channel properties that facilitates the export of

glutamate by YfkC.

AimA (YbeC) is a novel major glutamate transporter

The acquisition of suppressor mutations that provide resistance to glutamate to the Δ*dac* mutant was

a rare event that took several days to occur (see above). However, we observed the rapid appearance

of glutamate-resistant suppressor mutants if the strain already carried a *nhaK* mutation that facilitates potassium export (see Fig. 4). To determine the requirements for glutamate resistance in more detail, we isolated a set of 16 new suppressor mutants (see Fig. 1). Since the originally isolated suppressor mutant GP2842 carried a combination of mutations in *nhaK* and the *ybeC* gene encoding an amino acid transporter, we sequenced the *ybeC* allele in all glutamate-resistant suppressor mutants isolated from the Δ*dac nhaK*(S187F) (from now on referred to as *nhaK** and NhaK* for the gene and protein, respectively). Strikingly, all 16 strains carried mutations in *ybeC*. Among these mutations are frameshifts in ten strains, an insertion of 2 amino acids after residue 226 in one strain, and amino acid substitutions in the remaining five strains (for details, see Table 2). The high number of mutants with inactivated *ybeC* genes strongly suggests that the combination of NhaK* and loss of YbeC allows growth of the Δ*dac* mutant in the presence of glutamate. To test this hypothesis, we deleted the *ybeC* gene in the Δ*dac* mutant. The resulting strain, GP3054, was unable to grow in the presence of glutamate, indicating that only the combination of the *nhaK** and the *ybeC* mutations conferred resistance to glutamate to the Δ*dac* mutant.

Three lines of evidence suggest that YbeC is the major glutamate transporter of *B. subtilis*: (i) We isolated seven independent mutants affected in *ybeC* in the initial screen for mutations conferring glutamate resistance to the Δ*dac* strain, (ii) we reproducibly isolated a large number of *ybeC* mutants based on the Δ*dac nhaK** mutant, and (iii) YbeC is an amino acid transporter of the amino acid-polyamine-organocation (APC) superfamily. We have recently demonstrated that YbeC is the major serine transporter of *B. subtilis* [53]. It is well established that many amino acid transporters can take up multiple different amino acids. To test the possible role of YbeC in glutamate uptake, we first attempted functional complementation in *E. coli* that is unable to grow with glutamate as the single source of carbon and nitrogen due to the low activity of intrinsic glutamate transporters [71]. For this purpose, we transformed *E. coli* JM109 with plasmid pGP2987 that carries the *ybeC* gene or with the empty vector pWH844 [72]. The resulting transformants were tested for growth on minimal medium with glucose and ammonium or with glutamate as the only source of carbon and nitrogen (see Fig. 5A).

Whereas both strains grew well with glucose and ammonium, the strain carrying the empty vector was

unable to utilize glutamate. In contrast, the expression of the *ybeC* gene allowed *E. coli* to use

glutamate as source of carbon and energy. These findings confirm that YbeC indeed is a glutamate

transporter. The protein and the gene were therefore renamed AimA and *aimA*, respectively (amino

acid importer A).

It is very striking that our suppressor screens for glutamate-resistant mutants often identified

AimA but never GltT or GltP, the two described glutamate transporters of *B. subtilis*. Therefore, we

analyzed the contributions of these proteins to glutamate uptake by comparing the growth of strains

expressing only one of the three proteins and a strain lacking all of them (see Fig. 5B). All tested strains

grew well in the presence of ammonium as the nitrogen source, reflecting their ability to synthesize

glutamate. The strain GP3071 lacking all three presumptive glutamate transporters was only viable at

high glutamate concentrations but not at 1 mM or 5 mM. This demonstrates that the bacteria need to

transport glutamate if no ammonium is available for its synthesis. Moreover, it demonstrates that

efficient glutamate transport is not possible in the absence of the three proteins. GP2818 expressing

GltT grew at the three tested glutamate concentrations confirming its role as glutamate transporter.

The strain GP2825 that expressed AimA as the only glutamate transporter did not grow at the lowest

tested glutamate concentration (1 mM), but this strain was able to grow at 5 mM and 20 mM

glutamate suggesting that AimA might be a low-affinity glutamate transporter. In contrast, the strain

expressing GltP was only viable at highest glutamate concentration as observed for strain GP3071

lacking all three proteins. This suggests that GltP plays only a very minor, if any role, in glutamate

uptake, and that yet another protein can transport glutamate at high concentrations.

To study the activities of AimA and GltT in glutamate transport in more detail, we assayed the

growth of strains expressing only one of these transporters at different glutamate concentrations (see

Fig. 5C, 5D). Moreover, we performed this experiment in the presence of 0.1 mM and 5 mM KCl in the

medium to see whether potassium has an effect on glutamate uptake as it has been observed with an

activating effect of glutamate on potassium uptake by KtrCD [41]. As control, we used strain GP3071

that lacks AimA, GltT, and GltP. The results were used to calculate apparent K_S values that reflect the

affinities of the transporters for glutamate. For GltT, we observed K_S values of 0.3 mM and 0.06 mM in

the presence of 0.1 and 5 mM potassium, respectively. These results are in good agreement with the

description of GltT as a high-affinity glutamate transporter [73]. Moreover, they show that the affinity

of GltT for glutamate is five-fold increased in the presence of high potassium concentrations. For AimA,

we determined apparent K_S values of 3.2 and 2.3 mM in the presence of 0.1 and 5 mM potassium,

respectively. In good agreement with the drop dilution assay, this suggests that AimA is a low-affinity

glutamate transporter, and that the affinity of AimA is not affected by the external potassium

concentration. The strain GP3071 lacking all known and suspected glutamate transporters grew only

at high glutamate concentrations. This suggests the presence of an additional transporter with very

low affinity for glutamate (apparent K_S values of 12.3 and 13.8 mM in the presence of 0.1 and 5 mM

potassium, respectively). Thus, AimA is the major glutamate transporter under standard growth

conditions when glutamate is not limiting in the medium.

The role of AimA as the major glutamate transporter is in excellent agreement with the

common isolation of glutamate-resistant Δ*dac aimA* mutants whereas no *gltT* mutants could be

isolated. Indeed, the Δ*dac* Δ*gltT* mutant was unable to grow in the presence of glutamate, but again a

suppressor mutant, GP3093, could be isolated (see Fig. 1). Not surprisingly, this strain carries a

mutation in *aimA* (Leu-419 Pro). Moreover, this strain carries frame-shift mutations in *ndhF* and *fliK*.

The paired occurrence of *aimA* and *ndhF* suppressor mutations has been observed several times

independently (see Table 1).

To get further insights into *gltP*, *gltT* and *aimA*, we determined the expression of these genes

based on the transcriptome analysis under the relevant conditions (see Fig. 5E). The expression of *gltP*

was very low under all tested conditions, which is in agreement with the observation that GltP does

not play a significant role in glutamate uptake. The expression of both GltT and AimA was significant.

In the presence of glutamate in the medium, the expression was reduced five- and two-fold for AimA

and GltT, respectively, at the high potassium concentration (5 mM). In the absence of glutamate, *aimA*

expression did not respond to the supply of potassium whereas the expression of *gltT* was about two-

fold increased at the increased potassium concentration. Thus, if glutamate is limiting, both the

expression of *gltT* and the affinity of the GltT protein for glutamate are increased by potassium.

Adaptation of a Δ*dac* Δ*aimA* mutant to growth in the presence of glutamate

To make the picture of the requirements for a Δ*dac* mutant to tolerate glutamate complete, we also

adapted a Δ*dac* Δ*aimA* mutant to the presence of glutamate. As described above, this mutant was not

viable in the presence of glutamate; however, we could isolate three glutamate-resistant suppressor

mutants (see Fig. 1). The mutations in these strains were identified by whole genome sequencing.

Interestingly, all three suppressor mutants had mutations affecting the essential translation elongation

factor G, in two strains (GP3101 and GP3103) the highly conserved Ala-579 was replaced by a Val (see

S7 Fig.). Ala-579 is located in immediate vicinity of a loop (Loop II) that is required for tRNA

translocation in the ribosome [74, 75], suggesting that the mutation interferes with translation. In the

third strain (GP3102) we found an A21T substitution. This residue is the immediate neighbour of Asp-

20, which binds the Mg^{2+} ion in the active site of the protein [76]. The presence of substitutions in

highly conserved and functionally important regions of FusA in all three suppressors suggests that

reduced translation efficiency helps the Δ*dac* Δ*aimA* mutant to adapt to the presence of glutamate. In

addition to the mutations in *fusA* each strain had a distinct additional mutation. Strain GP3101 had a

frameshift mutation in the *yqeG* gene encoding a putative HAD superfamily phosphatase [77], whereas

the two other strains carried point mutations affecting potassium homeostasis (*nhaK* and *ktrC* for

GP3102 and GP3103, respectively) again highlighting the toxicity of potassium in the presence of

glutamate. Moreover, GP3102 carried amino acid substitutions affecting SigA, the housekeeping sigma

factor (F281V) and MrpA (P397L), a subunit of the sodium exporter of *B. subtilis*, and GP3103 had a

point mutation affecting the quasi-essential DNA topoisomerase 1 TopA (K13N). The analysis of these

strains for growth in the presence of glutamate with 0.1 or 5 mM potassium revealed that all

suppressors were viable at both potassium concentrations, however, the suppression was most

efficient if the *yqeG* or *ktrC* genes were affected in addition to *aimA* (see S8 Fig.).

DISCUSSION

Cyclic di-AMP is essential for many bacteria that produce this second messenger; however, the reasons

behind this essentiality have not been fully understood. In this study, we used the Gram-positive model

organism *B. subtilis* to gain further insights into the critical functions of c-di-AMP. Based on a variety

of distinct suppressor screens to adapt a *B. subtilis* strain lacking c-di-AMP to either the presence of

glutamate or to growth on complex medium, we can conclude that the control of potassium

homeostasis is the major bottleneck that limits growth of the mutant under both conditions. This is in

good agreement with (i) the large variety of c-di-AMP target proteins and RNA molecules that are

involved in the uptake and export of potassium in *B. subtilis* and other bacteria [11, 13, 20, 21, 23, 31,

34, 45, 78], (ii) the fact that the intracellular c-di-AMP levels seem to report the extracellular potassium

concentrations [18, 21, 37], and the isolation of suppressor mutants affecting potassium homeostasis

in response to altered cellular c-di-AMP levels also in other bacteria [18, 37]. However, our study also

clearly demonstrates that potassium is not the only problem for growth of *B. subtilis* in the absence of

c-di-AMP since the presence of mutations that reduce potassium uptake or facilitate its export is not

sufficient to overcome the toxicity of glutamate or complex medium. Suppressor studies with *L.*

monocytogenes and *S. aureus* mutants lacking c-di-AMP revealed that those mutants often had

acquired mutations in transporters for osmoprotectants such as glycine betaine indicating that

intoxication by compatible solutes is the main reason for essentiality of c-di-AMP in these latter

bacteria. The idea of a major role for c-di-AMP in osmoprotection rather than in potassium

homeostasis in these pathogenic bacteria is also supported by the observation that potassium has only

a minor impact on a *L. monocytogenes* strain lacking the only diadenylate cyclase [78]. Thus, while

general features of c-di-AMP signaling are conserved in different bacteria, the specific roles and

mechanisms may differ even between closely related bacteria.

Potassium and glutamate are the most abundant cation and anion, respectively, in every living

cell, and the intracellular concentrations of the two ions need to be balanced [79]. In *B. subtilis*, c-di-

AMP seems to play an important role in achieving this balance. The concentrations of the nucleotide

respond both to the potassium and the nitrogen source availability, they are high at high potassium

concentrations and in the presence of glutamate but not ammonium or glutamine [15, 21,40]. Our

work supports the idea of a close physiological integration of potassium and glutamate homeostasis:

Both potassium and glutamate inhibit growth of the Δ*dac* mutant independently of each other. The

accumulation of amino acids has also been shown to be a problem for the Δ*dac* mutants of *L.*

monocytogenes and *S. aureus*. As presented in this study, in both organisms amino acid transporters

were often inactivated in suppressor mutants that were capable of growing on complex medium

despite the lack of c-di-AMP: In *L. monocytogenes*, several suppressor mutants carried mutations

inactivating the OppABCD oligopeptide transporter [39] whereas the characterization of *S. aureus*

suppressor mutants resulted in the identification of the AlsT glutamine transporter, thus indicating

that glutamine rather than glutamate is toxic for the *S. aureus* Δ*dac* mutant [39, 42]. Again, these

differences between the related organisms support the idea that c-di-AMP has global overarching

functions in the different bacteria, but these can be put into practice differently.

It had been established previously that the accumulation of glutamate can be toxic for *B.*

subtilis [52, 80]. In the case of the Δ*dac* mutant, our work has unraveled three ways to overcome

glutamate toxicity: (i) Glutamate can be efficiently degraded upon decryptification of the otherwise

inactive *gudB* gene encoding the major constitutive glutamate dehydrogenase of *B. subtilis*. Such a

well established *gudB*+ mutation [51] was found in the suppressor mutants GP2841 and GP3473. (ii)

The uptake of glutamate can be prevented, and this was indeed observed in many suppressor mutants.

Surprisingly, the recently characterized serine transporter AimA (YbeC) was identified as the main

glutamate transporter rather than the previsously studied GltT [73]. Even the Δ*dac* mutant lacking *gltT*

was unable to grow in the presence of glutamate unless AimA had been inactivated (see Fig. 1). This

observation as well as our detailed kinetic analyses revealed that AimA is the major low affinity

transporter for glutamate. Moreover, in the absence of c-di-AMP, the *aimA* gene is the most strongly

expressed gene encoding a glutamate transporter (see Fig. 5E). It is interesting to note that the

expression of *aimA* responds to the availability of c-di-AMP (see Fig. 5B), however, the mechanism

responsible for this control has yet to be uncovered. We have previously demonstrated that a

potassium channel (KtrCD) is directly activated and thus converted from a low- to a high-affinity uptake

system in the presence of glutamate [41]. Here, we show that similarly the affinity of the high affinity

glutamate transporter GltT is controlled by the presence of potassium: At high potassium

concentrations, the affinity of GltT for glutamate is increased fivefold. Thus, the uptake systems for

the two most abundant ions in the cell can respond to the presence of the other ion to keep their

intracellular concentrations balanced. (iii) In addition to mutations activating glutamate degradation

or preventing glutamate uptake, we found several mutants with paired mutations affecting lipid

biosynthesis and the mechanosensitive channel of small conductivity YfkC. Moreover, the Δ*dac yfkC**

mutant expressing the altered channel protein had to acquire additional mutations affecting either

glutamate uptake (*aimA*) or fatty acid biosynthesis (*accA*) to tolerate the otherwise toxic glutamate.

We suggest that the altered YfkC protein in conjunction with altered membrane properties facilitates

the export of glutamate. This idea is supported by the observation that YfkC needs a gain-of-function

mutation to bypass glutamate toxicity. Moreover, MscS-like mechanosensitive channels have been

shown to export glutamate in *Corynebacterium glutamicum* [81, 82]. In this bacterium, a gain of

function mutation affecting this channel results in constitutive glutamate secretion [83]. Finally,

specific membrane properties in *C. glutamicum* such as reduced lipid amounts facilitate glutamate

export by the MscS-like channel (see [84] for review). Our results and the parallels in *C. glutamicum*

strongly support the idea that the combination of mutations reducing lipid biosynthesis and that affect

the mechanosensitive channel YfkC allow glutamate export in *B. subtilis*, thus overcoming the toxic

effect of this amino acid.

Three lines of evidence suggest that respiration is a major bottleneck for *B. subtilis* in the

absence of c-di-AMP. First, many suppressors isolated in the presence of glutamate had mutations that

inactivate the *ndhF* gene encoding a subunit of a putative NADH dehydrogenase or the downstream

ybcC gene of unknown function. It is tempting to speculate that the two proteins form a complex that

catalyzes the oxidation of $NADH_2$. Second, when we used complex medium to isolate suppressor

mutants, this was initially only possible in the absence of oxygen if respiration activity was low. This is

paralleled by the observation that the *S. aureus* Δ*dac* is viable in the absence but not in the presence

of oxygen [40]. Finally, the lack of c-di-AMP results in an enhanced expression of the *ndhF-ybcC* operon

and reduced intracellular $NADH_2$ levels as indicated by the strong repression of the genes of the Rex

regulon in the Δ*dac* mutant. Although it is not yet understood how precisely respiration interferes with

the growth of the Δ*dac* mutant, it is tempting to speculate that the proton motive force generated by

respiration plays a key role: the major potassium transporter KimA is actually a proton:potassium

symporter [85],and potassium uptake by the KtrAB and KtrCD channels is directly driven by the proton

motive force [86]. Thus, a reduced respiration lowers the proton motive force and may thus limit the

uptake of potassium, the major growth-limiting factor for the *B. subtilis* Δ*dac* mutant.

The adaptation of the Δ*dac* mutant to complex medium did not only confirm the toxicity of

potassium, but it did also highlight the particular importance of the two c-di-AMP-binding signal

transduction proteins DarA and DarB. The acquisition of suppressor mutations that allow growth under

anaerobic conditions on complex medium was only possible when one of these proteins was absent

(see Fig. 1). This indicates that these proteins inhibit growth in their c-di-AMP-free apo form. For DarB

it has very recently been shown that the apo protein stimulates the synthesis of (p)ppGpp by the

alarmone synthetase Rel both in *B. subtilis* and *L. monocytogenes* [59, 60], and excessive (p)ppGpp

synthesis limits growth of both organisms [39, 59, 60]. For DarA, no function has been identified so far.

It is interesting to note, however, that DarA seems to be even more inhibitory than DarB since the Δ*dac*

Δ*darB* mutants tend to acquire additional mutations affecting DarA when adapting to the presence of

oxygen.

Taken together, our study supports the idea that c-di-AMP plays a central role in the physiology

of *B. subtilis*. Moreover, by comparing the results obtained with suppressor screens in closely related

but distinct bacteria, it becomes clear that the overall themes of c-di-AMP signaling are similar in

different bacteria whereas the precise mechanisms are not. While some of the functions of c-di-AMP

in *B. subtilis* such as the control of potassium homeostasis are already well understood, there remain

many open questions for future research such as the link between c-di-AMP and lipid biosynthesis and

the function of the PII-like signal transduction protein DarA. The work described here provides a solid

ground for further investigation.

MATERIALS AND METHODS

Strains, media and growth conditions. *E. coli* DH5α [87] was used for cloning and for the expression

of recombinant proteins. *E. coli* JM109 (Promega) was used for glutamate transporter

complementation assays. All *B. subtilis* strains used in this study are derivatives of the laboratory strain

168. They are listed in Table 2. *B. subtilis* was grown in Luria-Bertani (LB) or in sporulation (SP) medium

[87, 88]. For the assay of potassium and glutamate toxicity, *B. subtilis* was cultivated in MSSM medium

[21]. In this modified SM medium KH_2PO_4 was replaced by NaH_2PO_4 and KCl was added as indicated.

The media were supplemented with ampicillin (100 μg/ ml), kanamycin (10 and 50 μg/ ml for *B. subtilis*

and *E. coli*, respectively), chloramphenicol (5 μg/ ml), tetracyclin (12.5 μg/ ml), spectinomycin (150 μg/

ml) or erythromycin and lincomycin (2 and 25 μg/ ml, respectively) if required.

Table 2. Bacterial strains used in this study.

Strain	Genotype	Reference
168	*trpC2*	Laboratory collection
BKE14130	*trpC2 ΔdarB::erm*	[89]

GP94	*trpC2 ΔcdaA::spc*	[21]
GP342	*trpC2 amyE*::(*gltA-lacZ kan*)	[90]
GP987	*trpC2 ΔdisA::tet*	[9]
GP991	*trpC2 ΔcdaS::ermC ΔdisA::tet*	[9]
GP997	*trpC2 ΔcdaA::cat*	[9]
GP1360	*trpC2 ΔcdaS::spc*	This study
GP1361	*trpC2 ΔcdaS::spc disA::tet*	GP1360 → GP987
GP1712	*trpC2 ΔdarA::cat*	[47]
GP2181	*trpC2 amyE*::(P-*kimA*-RS-*lacZ cat*)	pGP2912 → 168
GP2222	*trpC2 ΔcdaA::cat ΔcdaS::ermC ΔdisA::tet*	[21]
GP2223	*trpC2 ΔcdaA::cat ΔcdaS::ermC ΔdisA::tet nhaK$_{S187F}$*	[21]
GP2224	*trpC2 ΔcdaA::cat ΔcdaS::ermC ΔdisA::tet aimA $_{frameshift}$* *yqeG$_{frameshift}$ yqeV$_{frameshift}$*	This study
GP2245	*trpC2 ΔcdaS::ermC ΔdisA::tet ΔgltT::kan*	GP2248 → GP991
GP2248	*trpC2 ΔgltT::kan*	This study
GP2259	*trpC2 ΔcdaA::cat ΔcdaS::ermC ΔgltT::kan ΔdisA::tet*	GP997 → GP2245
GP2413	*trpC2 ΔcdaS::ermC ΔdisA::tet ΔdarA::cat*	GP1712 → GP991
GP2420	*trpC2 ΔcdaS::ermC ΔdisA::tet ΔdarA::cat ΔcdaA::spc*	GP94 → GP2413
GP2777	*trpC2 ΔcdaS::spc ΔdisA::tet ΔdarB::ermC*	BKE14130 → GP1361
GP2779	*trpC2 ΔcdaS::spc ΔdisA::tet ΔdarB::ermC ΔcdaA::cat*	GP997 → GP2777
GP2785	*trpC2 ΔyfkC::kan*	This study
GP2786	*trpC2 ΔaimA::kan*	This study
GP2796	*trpC2 ΔcdaS::spc disA::tet ΔyfkC::kan*	GP2785 → GP1361
GP2797	*trpC2 ΔcdaS::spc disA::tet ΔyfkC::kan ΔcdaA::cat*	GP997 → GP2796

GP2798	*trpC2 yfkC$_{R42W}$ kan*	This study
GP2813	*trpC2 ΔcdaS::spc disA::tet yfkC$_{R42W}$ kan*	GP2798 → GP1361
GP2814	*trpC2 ΔcdaS::spc disA::tet yfkC$_{R42W}$ kan ΔcdaA::cat*	GP997 → GP2813
GP2818	*trpC2 ΔgltP::cat ΔaimA::kan*	GP2786 → GP2824
GP2824	*trpC2 ΔgltP::cat*	This study
GP2825	*trpC2 ΔgltT::kan ΔgltP::cat*	GP2824 → GP2248
GP2831	*trpC2 ΔgltT::kan ΔaimA::phleo*	LFH → GP2248
GP2837	*trpC2 ΔcdaA::cat ΔcdaS::ermC ΔdisA::tet yfkC$_{R42W}$ plsC$_{L48P}$ fliP$_{frameshift}$*	This study
GP2838	*trpC2 ΔcdaA::cat ΔcdaS::ermC ΔdisA::tet ndhF$_{V237E, M241L, H242L, A243D, G244C, L245F}$ ΔydcL-yddM yoaI$_{frameshift}$*	This study
GP2839	*trpC2 ΔcdaA::cat ΔcdaS::ermC ΔdisA::tet fliK$_{truncated}$ rpoB$_{Q109P}$ rbsB$_{R175L}$*	This study
GP2840	*trpC2 ΔcdaA::cat ΔcdaS::ermC ΔdisA::tet plsC$_{A61V}$ ndhF$_{frameshift}$ ktrC$_{frameshift}$ sigD$_{G88D}$*	[41]
GP2841	*trpC2 ΔcdaA::cat ΔcdaS::ermC ΔdisA::tet gudB$_{decrypt.}$ accC$_{H298Y}$*	This study
GP2842	*trpC2 ΔcdaA::cat ΔcdaS::ermC ΔdisA::tet nhaK$_{S187F}$ aimA$_{frameshift}$*	This study
GP2845	*trpC2 ΔcdaS::spc ΔdisA::tet ΔdarB::ermC ΔcdaA::cat ktrC* (fs) *kimA$_{Δ354-364}$ darA$_{V100A}$*	This study
GP2849	*trpC2 ΔcdaA::cat ΔcdaS::ermC ΔdisA::tet aimA$_{frameshift}$ ybcC$_{K722L/I724F}$*	This study
GP2850	*trpC2 ΔcdaS::spc disA::tet yfkC$_{R42W}$ kan ΔcdaA::cat accA$_{A229V}$ yerC$_{T58K}$ ypoP$_{KTEYF\ 111\ I}$*	This study

GP3010	*trpC2 ΔcdaS::ermC ΔdisA::tet ΔdarA::cat ΔcdaA::spc*	This study
	ykoX$_{K218STOP}$ ktrC$_{G11C}$ yfkF$_{G294S}$ nusG$_{T82K}$ kimA$_{ΔG112}$ ccpA$_{Δbp850-1005}$	
	(incl 24bp downstream)	
GP3053	*trpC2 ΔcdaS::ermC ΔdisA::tet ΔaimA::phleo*	GP2831 → GP991
GP3054	*trpC2 ΔcdaS::ermC ΔdisA::tet ΔaimA::phleo ΔcdaA::cat*	GP997 → GP3053
GP3071	*trpC2 ΔgltT::kan ΔaimA::phleo ΔgltP::cat*	GP2824 → GP2831
GP3077	*trpC2 ΔcdaA::cat ΔcdaS::ermC ΔdisA::tet ktrC$_{frameshift}$*	This study
	aimA/glpQ$_{truncation}$ ybcC$_{frameshift}$	
GP3078	*trpC2 ΔcdaA::cat ΔcdaS::ermC ΔdisA::tet plsC$_{A61V}$ aimA$_{T374M}$*	This study
	ktrB$_{G313D}$ sigD$_{G88D}$	
GP3079	*trpC2 ΔcdaA::cat ΔcdaS::ermC ΔdisA::tet yfkC$_{R42W}$ plsC$_{L48P}$*	This study
	Δ(abh-kinC-ykqA-ktrC) hbs$_{I32T}$ ndoA$_{G6S}$ fliP$_{frameshift}$	
GP3093	*trpC2 ΔcdaA::cat ΔcdaS::ermC ΔgltT::kan ΔdisA::tet aimA$_{L419P}$*	This study
	ndhF$_{frameshift}$ fliK$_{frameshift}$	
GP3094	*trpC2 ΔcdaS::spc ΔdisA::tet ΔdarB::ermC ΔcdaA::cat*	This study
	ktrC$_{frameshift}$ kimA$_{Δ354-364}$ codY$_{G146V}$ yonD$_{D255H}$	
GP3099	*trpC2 ΔcdaA::cat ΔcdaS::ermC ΔdisA::tet plsC$_{L48P}$ yfkC$_{R42W}$*	This study
	fliP$_{frameshift}$ cotO$_{I34F}$	
GP3101	*trpC2 ΔcdaS::ermC ΔdisA::tet ΔaimA::phleo ΔcdaA::cat*	This study
	fusA$_{A579V}$ yqeG$_{frameshift}$	
GP3102	*trpC2 ΔcdaS::ermC ΔdisA::tet ΔaimA::phleo ΔcdaA::cat*	This study
	nhaK$_{R486C}$ sigA$_{F281V}$ mprA$_{P397L}$ fusA$_{A579V}$	
GP3103	*trpC2 ΔcdaS::ermC ΔdisA::tet ΔaimA::phleo ΔcdaA::cat*	This study
	ktrC$_{G14D}$ topA$_{K13N}$ fusA$_{A21T}$	

GP3105	$trpC2$ $\Delta cdaS{::}spc$ $disA{::}tet$ $yfkC_{R42W}$ kan $\Delta cdaA{::}cat$ $aimA_{frameshift}$ $carB_{T257R}$ $mutL_{frameshift}$ $pksJ_{L4386F}$	This study
GP3464	$trpC2$ $\Delta cdaA{::}cat$ $\Delta cdaS{::}ermC$ $\Delta disA{::}tet$ $plsC_{P110S}$ $flhA_{frameshift}$	This study
GP3473	$trpC2$ $\Delta cdaA{::}cat$ $\Delta cdaS{::}ermC$ $\Delta disA{::}tet$ $ndhF_{frameshift}$ $aimA_{frameshift}$ $gudB^+$ $fliH_{frameshift}$ $yuiE_{G37V}$	This study
GP3474	$trpC2$ $\Delta cdaA{::}cat$ $\Delta cdaS{::}ermC$ $\Delta disA{::}tet$ $ndhF$ (fs) $aimA_{frameshift}$ $ykpA_{frameshift}$ $ktrC_{frameshift}$ $fliK_{frameshift}$ $fliH_{silent}$	This study
GP3486	$trpC2$ $\Delta cdaA{::}cat$ $\Delta cdaS{::}ermC$ $\Delta disA{::}tet$ $nhaK_{S187F}$ $aimA_{frameshift}$	This study
GP3487	$trpC2$ $\Delta cdaA{::}cat$ $\Delta cdaS{::}ermC$ $\Delta disA{::}tet$ $nhaK_{S187F}$ $aimA_{frameshift}$	This study
GP3488	$trpC2$ $\Delta cdaA{::}cat$ $\Delta cdaS{::}ermC$ $\Delta disA{::}tet$ $nhaK_{S187F}$ $aimA_{frameshift}$	This study
GP3489	$trpC2$ $\Delta cdaA{::}cat$ $\Delta cdaS{::}ermC$ $\Delta disA{::}tet$ $nhaK_{S187F}$ $aimA_{P279H}$	This study
GP3490	$trpC2$ $\Delta cdaA{::}cat$ $\Delta cdaS{::}ermC$ $\Delta disA{::}tet$ $nhaK_{S187F}$ $aimA_{I164N}$	This study
GP3491	$trpC2$ $\Delta cdaA{::}cat$ $\Delta cdaS{::}ermC$ $\Delta disA{::}tet$ $nhaK_{S187F}$ $aimA_{frameshift}$	This study
GP3492	$trpC2$ $\Delta cdaA{::}cat$ $\Delta cdaS{::}ermC$ $\Delta disA{::}tet$ $nhaK_{S187F}$ $aimA_{frameshift}$	This study
GP3493	$trpC2$ $\Delta cdaA{::}cat$ $\Delta cdaS{::}ermC$ $\Delta disA{::}tet$ $nhaK_{S187F}$ $aimA_{frameshift}$	This study
GP3494	$trpC2$ $\Delta cdaA{::}cat$ $\Delta cdaS{::}ermC$ $\Delta disA{::}tet$ $nhaK_{S187F}$ $aimA_{S106P}$	This study
GP3495	$trpC2$ $\Delta cdaA{::}cat$ $\Delta cdaS{::}ermC$ $\Delta disA{::}tet$ $nhaK_{S187F}$ $aimA_{N349K}$	This study
GP3496	$trpC2$ $\Delta cdaA{::}cat$ $\Delta cdaS{::}ermC$ $\Delta disA{::}tet$ $nhaK_{S187F}$ $aimA_{frameshift}$	This study
GP3497	$trpC2$ $\Delta cdaA{::}cat$ $\Delta cdaS{::}ermC$ $\Delta disA{::}tet$ $nhaK_{S187F}$ $aimA_{frameshift}$	This study
GP3498	$trpC2$ $\Delta cdaA{::}cat$ $\Delta cdaS{::}ermC$ $\Delta disA{::}tet$ $nhaK_{S187F}$ $aimA_{G70R}$	This study
GP3499	$trpC2$ $\Delta cdaA{::}cat$ $\Delta cdaS{::}ermC$ $\Delta disA{::}tet$ $nhaK_{S187F}$ $aimA_{Insertion}$ $_{after\ aa\ 226\ (+VA)}$	This study
GP3500	$trpC2$ $\Delta cdaA{::}cat$ $\Delta cdaS{::}ermC$ $\Delta disA{::}tet$ $nhaK_{S187F}$ $aimA_{frameshift}$	This study
GP3701	$trpC2$ $\Delta cdaA{::}cat$ $\Delta cdaS{::}ermC$ $\Delta disA{::}tet$ $nhaK_{S187F}$ $aimA_{frameshift}$	This study

GP3702	*trpC2 ΔcdaS::spc ΔdisA::tet ΔdarB::ermC ΔcdaA::cat*	This study
	ktrC$_{frameshift}$ kimA$_{Δ354-364}$ darA$_{frameshift}$	
GP3703	*trpC2 ΔcdaS::spc ΔdisA::tet ΔdarB::ermC ΔcdaA::cat*	This study
	ktrC$_{frameshift}$ kimA$_{Δ354-364}$ darA$_{frameshift}$	

Suppressor screen. For identification of suppressor mutations that rescue the growth of the c-di-AMP

lacking strain GP2222, a suppressor screen was set up. The same set-up was performed for the

suppressor screens with GP2223, GP2814, GP3054, and GP2259 (see Fig. 1). The respective strain was

plated on MSSM minimal plates in the presence of ammonium and 0.1 mM KCl. After incubation for

2-3 days, cell material was scraped from the plates and washed three times in MSSM basal salts

solution. These cells were either plated on MSSM minimal plates (Na-Glutamate, 0.1 mM KCl) or used

for inoculation of 10 ml MSSM minimal medium (Na-Glutamate, 0.1 mM KCl). The bacteria were then

incubated at 42°C for 3-4 days. As soon as suppressors appeared on the plates, they were picked and

re-isolated three times. For liquid cultures, as far as growth could be observed, they were plated.

Suppressors were picked and re-isolated again. For the suppressor screen on complex medium,

GP2222, GP2420, and GP2779 were plated on LB plates. The plates were incubated at 37°C either

under standard conditions (aerobic) or in an anaerobic jar to create an anaerobic environment. As soon

as suppressors appeared, they were picked and re-isolated three times. For the two step adaptation

to complex medium, suppressor mutants previously isolated in the anaerobic screen were plated on

LB plates and incubated at 37°C under aerobic conditions. As soon as new suppressors appeared, they

were picked and re-isolated three times. All isolated mutants were analyzed by whole-genome

sequencing (see below).

Phenotypic characterization. Amylase activity was detected on plates containing nutrient broth (7.5

g/liter), 17 g of Bacto agar per liter (Difco), and 5 g of hydrolyzed starch per liter (Connaught). The

ability to degrade starch was detected by sublimating iodine onto the plates. For quantitative

determination of *lacZ* expression in *B. subtilis,* cells were grown in MSSM medium with the nitrogen

source and potassium concentration as indicated and harvested in the exponential phase. β-

Galactosidase activity was determined with cell extracts obtained by lysozyme treatment as described

previously [88]. One unit of β-galactosidase is defined as the amount of enzyme that produces 1 nmol

of o-nitrophenol per minute at 28°C.

To assay growth of *B. subtilis* mutants at different potassium concentrations, the bacteria were

inoculated in LB medium and precultured in MSSM medium with 0.1 mM KCl. The cultures were grown

until exponential phase, harvested, washed three times in MSSM basal salts solution before the OD_{600}

was adjusted to 1.0. Dilution series were then pipetted onto MSSM plates with ammonium or

glutamate and 0.1 or 5 mM potassium.

Complementation in *E. coli* was assayed by transformation of strain JM109 that is unable to

grow with glutamate as the single source of carbon and nitrogen. For this purpose, the strain was

transformed with the complementation plasmid pGP2987 (*aimA*) and the empty vector control

pWH844. Growth of the transformants was assessed on M9 minimal medium plates containing either

ammonium and glucose or glutamate as carbon and nitrogen source [71].

DNA manipulation and genome sequencing. Transformation of *E. coli* and plasmid DNA extraction

were performed using standard procedures [87]. All commercially available plasmids, restriction

enzymes, T4 DNA ligase and DNA polymerases were used as recommended by the manufacturers.

Chromosomal DNA of *B. subtilis* was isolated as described [88]. *B. subtilis* was transformed with

plasmid and genomic DNA according to the two-step protocol [88].

To identify the mutations in the suppressor mutants, their genomic DNA was subjected to

whole-genome sequencing [91]. Briefly, the reads were mapped on the reference genome of *B. subtilis*

168 (GenBank accession number: NC_000964) [92] using the Geneious software package (Biomatters

Ltd., New Zealand) [93]. Single nucleotide polymorphisms were considered as significant when the

total coverage depth exceeded 25 reads with a variant frequency of ≥90%. All identified mutations

were verified by PCR amplification and Sanger sequencing.

Construction of mutant strains by allelic replacement. Gene deletion was achieved by transformation

of *B. subtilis* 168 with a PCR product constructed using oligonucleotides to amplify DNA fragments

flanking the target genes and an appropriate intervening resistance cassette as described previously

[94]. The mutation *yfkC*-R42W was introduced by amplification of regions containing and flanking the

mutation from strain GP2837, respectively, and a kanamycin resistance cassette was introduced

downstream of *yfkC* in the non-coding region between *yfkC* and *yfkD* (terminator region of both genes

present). The integrity of the regions flanking the integrated resistance cassette was verified by

sequencing PCR products of about 1,000 bp amplified from chromosomal DNA of the resulting mutant

strains.

Plasmids. Construction of a reporter fusion of the *kimA* promoter region with the promoterless *lacZ*

gene was done using the plasmid pAC6 [95]. The promoter fragment was cloned between the EcoRI

and BamHI sites of pAC6. The resulting plasmid was pGP2912. The plasmid was linearized with ScaI

and transformed into *B. subtilis* 168 for chromosomal integration into the *amyE* locus. For *E. coli*

complementation assays, the *aimA* gene was amplified using the primer pair LK167 (5'

AAAGGATCCATGAATCAATTGCATCGAAGAATGGGAACGTTTT)/   AK324   (5'

TTTGTCGACTTATTCTTTTCCGGCAGCAGCTTCTG) and cloned between the BamHI and SalI sites of the

expression vector pWH844 [72]. The resulting plasmid was pGP2987.

Determination of apparent glutamate transporter kinetics. The growth characteristics of *B. subtilis*

glutamate transporter mutants were used to determine the single glutamate transporter kinetics. The

mutants were either positive for *gltT* (*ΔgltP::cat ΔaimA::kan*) or *aimA* (*ΔgltT::kan ΔgltP::cat*) or lacked

both (*ΔgltT::kan ΔaimA::phleo ΔgltP::cat*). The nitrogen source was used as the limiting factor, so that

transport of glutamate was the essential factor for growth. The bacteria were inoculated in LB medium

and precultured in MSSM medium (ammonium) with 0.1/5 mM KCl. The cultures were grown until

exponential phase and the cells were washed three times in MSSM basal salts solution. Afterwards,

the cells were adjusted to OD_{600} 1.0 and used to inoculate a 96 well plate (Microtest Plate 96 Well, F,

Sarstedt) containing the MSSM medium with the respective glutamate and potassium concentrations.

The growth was tracked in an Epoch 2 Microplate Spectrophotometer (BioTek Instruments) at 37°C

with linear shaking at 237 cpm (4 mm) for 20 h, and optical density at 600 nm was measured in 10 min

intervals. The exponential growth phase was used to determine the growth rate μ [h^{-1}]. μ was

calculated from t_1 (early exponential phase) and t_2 (late exponential phase) and the corresponding OD_1

and OD_2 values. The resulting growth rates were then plotted against the glutamate concentrations.

This allowed fitting to the Monod equation and calculation of V_{max} [h^{-1}] and the apparent K_S [mM

glutamate] using the solver tool of Excel 2012 (Microsoft). Experiments were repeated with three

biological replicates.

Transcriptome analysis. Wild type and the c-di-AMP-deficient strains GP2222 Δ*dac* and GP2223 Δ*dac*

nhaK-S187F were grown in MSSM medium with ammonium or glutamate and low (0.1 mM) or high (5

mM) potassium concentrations. The strains were harvested by centrifugation (10.397 × g, 1 min, 4°C)

at mid exponential phase (OD_{600} of 0.4-0.6). A total of two independent biological replicates were

included. The pellets were frozen immediately in liquid nitrogen and stored at −80°C. Isolation of the

total RNA was done as described previously [96, 97], and the quality of the RNA preparations was

assessed by means of an Agilent 2100 Bioanalyzer according to the manufacturer's instructions. 5 µg

of total RNA were used for cDNA synthesis. The synthesis and fluorescence labeling of cDNA followed

a strand-specific method using the FairPlay III Microarray Labeling Kit (Agilent Technologies, Santa

Clara, CA, USA) and actinomycin D (Calbiochem) [98]. 200 ng of Cy3-labeled cDNA were hybridized to

the microarray following Agilent's hybridization, washing and scanning protocol (One-Color

Microarray-based Gene Expression Analysis, version 5.5). Data were extracted and processed using the

Feature Extraction software (version 11.5.1.1). An aggregated expression value was computed for each

annotated coding sequence and previously identified RNA feature [97]. Gene-level intensities were

scaled based on the intensity values of ten different *in vitro*-synthesized transcripts contained in the

One Color RNA Spike-In kit (Agilent Technologies) in order to account for technical variation associated

with sample processing. Genes with at least 2.5-fold difference in expression levels of the wild type at

the different conditions or between wild type and GP2222 mutant were considered significantly

affected.

To analyze the regulatory network, we compared the expression levels of the wild type strain

and the GP2222 mutant at ammonium 0.1 mM KCl. The genes were clustered in operons according to

their genetic localization, as described in *Subti*Wiki [99]. Additionally, gathered operons were grouped

by their corresponding documented regulator/s and metabolism. The visual representation of the

regulatory network was generated with Cytoscape and GIMP softwares [100, 101].

Data availability statement. The transcriptomic data reported in this study have been submitted to

the NCBI Gene Expression Omnibus (GEO; http://www.ncbi.nlm.nih.gov/geo/) under accession

number GSE156738.

ACKNOWLEDGEMENTS

We are grateful to Anika Klewing and Fabian Commichau for helpful discussion. We like to thank Felix

Mehne and Martin Weiss for the help with strain constructions and the initial characterization of a

suppressor mutant, respectively.

REFERENCES

1. Gomelsky M. cAMP, c-di-GMP, c-di-AMP, and now cGMP: bacteria use them all! Mol Microbiol. 2011; 79: 562-565.

2. Görke B, Stülke J. Carbon catabolite repression in bacteria: many ways to make the most out of nutrients. Nat Rev Microbiol. 2008, 6: 613-624.

3. McDonough KA, Rodriguez A. The myriad roles of cyclic AMP in microbial pathogens: from signal to sword. Nat Rev Microbiol. 2012; 10: 27-38.

4. Hengge R, Häussler S, Pruteanu M, Stülke J, Tschowri N, Turgay K. Recent advances and current trends in nucleotide second messenger signaling in bacteria. J Mol Biol. 2019; 431: 908-927.

5. Jenal U, Reinders A, Lori C. Cyclic di-GMP: a second messenger extraordinaire. Nat Rev Microbiol. 2017; 15: 271-284.

6. Liu K, Bittner AN, Wang JD. Diversity in (p)ppGpp metabolism and effectors. Curr Opin Microbiol. 2015; 24: 72-79.

7. Hauryliuk V, Atkinson GC, Murakami KS, Tenson T, Gerdes K. Recent functional insights into the role of (p)ppGpp in bacterial physiology. Nat Rev Microbiol. 2015; 13: 298-309.

8. Luo Y, Helmann JD. Analysis of the role of *Bacillus subtilis* σ^M in β-lactam resistance reveals an essential role for c-di-AMP in peptidoglycan homeostasis. Mol Microbiol. 2012; 83: 623-639.

9. Mehne FM, Gunka K, Eilers H, Herzberg C, Kaever V, Stülke J. Cyclic di-AMP homeostasis in *Bacillus subtilis*: both lack and high level accumulation of the nucleotide are detrimental for cell growth. J Biol Chem. 2013; 288: 2004-2017.

10. Woodward JJ, Lavarone AT, Portnoy DA. c-di-AMP secreted by intracellular *Listeria monocytogenes* activates a host type I interferon response. Science 2010; 328: 1703-1705.

11. Devaux L, Sleiman D, Mazzuoli MV, Gominet M, Lanotte P, Trieu-Cuot P, et al. Cyclic di-AMP regulation of osmotic homeostasis is essential in group B *Streptococcus*. PLoS Genet. 2018; 14: e1007342.

12. Corrigan RM, Bowman L, Willis AR, Kaever V, Gründling A. Cross-talk between two nucleotide-

signaling pathways in *Staphylococcus aureus*. J Biol Chem. 2015; 290: 5826-5839.

13. Blötz C, Treffon K, Kaever V, Schwede F, Hammer E, Stülke J. Identification of the components

involved in cyclic di-AMP signaling in *Mycoplasma pneumoniae*. Front Microbiol. 2017; 8: 1328.

14. Barker JR, Koestler BJ, Carpenter VK, Burdette DL, Waters CM, et al. STING-dependent recognition

of cyclic di-AMP mediates type I interferon responses during *Chlamydia trachomatis* infection.

mBio 2013; 4: e00018-13.

15. Gundlach J, Mehne FMP, Herzberg C, Kampf J, Valerius O, Kaever V, Stülke J. An essential poison:

synthesis and degradation of cyclic di-AMP in *Bacillus subtilis*. J Bacteriol. 2015; 197: 3265-3274.

16. Smith WM, Pham TH, Lei L, Dou J, Soomro AH, et al. Heat resistance and salt hypersensitivity in

Lactococcus lactis due to spontaneous mutation in llmg1816 (*gdpP*) induced by high-temperature

growth. Appl Environ Microbiol. 2012; 78: 7753-7759.

17. Huynh TN, Luo S, Pensinger D, Sauer JD, Tong L, Woodward JJ. An HD-domain phosphodiesterase

mediates cooperative hydrolysis of c-di-AMP to affect bacterial growth and virulence. Proc Natl

Acad Sci USA. 2015; 112: E747-E756.

18. Zarrella TM, Metzger DW, Bai G. Stress suppressor screening leads to detecting regulation of cyclic

di-AMP homeostasis by a Trk-family effector protein in *Streptococcus pneumoniae*. 2018; J

Bacteriol. 200: e00045-18.

19. Zhu Y, Pham TH, Nhiep TH, Vu NM, Marcellin E, et al. Cyclic-di-AMP synthesis by the diadenylate

cyclase CdaA is modulated by the peptidoglycan biosynthesis enzyme GlmM in *Lactococcus lactis*.

Mol Microbiol. 2016; 99: 1015-1027.

20. Nelson JW, Sudarsan N, Furukawa K, Weingerg Z, Wang JX, Breaker RR. Riboswitches in eubacteria

sense the second messenger cyclic di-AMP. Nat Chem Biol. 2013; 9: 834-839.

21. Gundlach J, Herzberg C, Kaever V, Gunka K, Hoffmann T, Weiß M, et al. Control of potassium

homeostasis is an essential function of the second messenger cyclic di-AMP in *Bacillus subtilis*. Sci

Signal. 2017; 10: eaal3011.

22. Gundlach J, Herzberg C, Hertel D, Thürmer A, Daniel R, Link H, Stülke J. Adaptation of *Bacillus*

subtilis to life at extreme potassium limitation. mBio 2017; 8: e00861-17.

23. Gundlach J, Krüger L, Herzberg C, Turdiev A, Poehlein A, Tascón I, et al. Sustained sensing in

potassium homeostasis: Cyclic di-AMP controls potassium uptake by KimA at the levels of

expression and activity. J Biol Chem. 2019; 294: 9605-9614.

24. Hutchison CA, Peterson SN, Gill SR, Cline RT, White O, Fraser CM, et al. Global transposon

mutagenesis and a minimal Mycoplasma genome. Science 1999; 286: 2165-2169.

25. Commichau FM, Pietack N, Stülke J. Essential genes in *Bacillus subtilis*: A re-evaluation after ten

years. Mol Biosyst. 2013; 9: 1068-1075.

26. Rancati G, Moffat J, Typas A, Pvelka N. Emerging and evolving concepts in gene essentiality. Nat

Rev Genet. 2018; 19: 34-49.

27. Reuß DR, Faßhauer P, Mroch PJ, Ul-Haq I, Koo BM, Pöhlein A, et al. Topoisomerase IV can

functionally replace all type 1A topoisomerases in *Bacillus subtilis.* Nucleic Acids Res. 2019; 47:

5231-5242.

28. Benda M, Woelfel S, Gunka K, Klumpp S, Poehlein A, Kálalová D, et al. Quasi-essentiality of RNase

Y in *Bacillus subtilis* is caused by its critical role in the control of mRNA homeostasis. doi:

https://doi.org/10.1101/2020.05.20.106237.

29. Tödter D, Gunka K, Stülke J. The highly conserved Asp23 family protein YqhY plays a role in lipid

biosynthesis in *Bacillus subtilis*. Front Microbiol. 2017; 8: 883.

30. Gallagher LA, Bailey J, Manoil C. Ranking essential bacterial processes by speed of mutant death.

Proc Natl Acad Sci USA. 2020; 117: 18010-18017.

31. Corrigan RM, Campeotto I, Jeganathan T, Roelofs KG, Lee VT, Gründling A. Systematic

identification of conserved bacterial c-di-AMP receptor proteins. Proc Natl Acad Sci USA*.* 2013;

110: 9084-9089.

32. Huynh TN, Choi PH, Sureka K, Ledvina HE, Campillo J, et al. Cyclic di-AMP targets the cystathione

beta-synthase domain of the osmolyte transporter OpuC. Mol Microbiol. 2016; 102: 233-243.

33. Schuster CF, Bellows LE, Tosi T, Campeotto I, Corrigan RM, et al. The second messenger c-di-AMP

inhibits the osmolyte uptake system OpuC in *Staphylococcus aureus*. Sci Signal. 2016; 9: ra81.

34. Quintana IM, Gibhardt J, Turdiev A, Hammer E, Commichau FM, et al. The KupA and KupB proteins

of *Lactococcus lactis* IL1403 are novel c-di-AMP receptor proteins responsible for potassium

uptake. J Bacteriol. 2019; 201: e00028-19.

35. Latoscha A, Drexler DJ, Al-Bassam MM, Kaever V, Findlay KC, et al. 2020. C-di-AMP hydrolysis by

a novel type of phosphodiesterase promotes differentiation of multicellular bacteria. Proc Natl

Acad Sci USA. 2020; 117: 7392-7400.

36. Moscoso JA, Schramke H, Zhang Y, Tosi T, Dehbi A, et al. Binding of cyclic di-AMP to the

Staphylococcus aureus sensor kinase KdpD occurs via the Universal Stress Protein domain and

downregulates the expression of the Kdp potassium transporter. J Bacteriol. 2015; 198: 98-110.

37. Pham HT, Nhiep NTH, Vu TNM, Huynh TN, Zhu Y, et al. Enhanced uptake of potassium or glycine

betaine or export of cyclic di-AMP restores osmoresistance in a high cyclic-di-AMP *Lactococcus*

lactis mutant. PLoS Genet. 2018; 14: e1007574.

38. Rubin BE, Huynh TN, Welkie DG, Diamond S, Simkovsky R, et al. High-throughput interaction

screens illuminate the role of c-di-AMP in cyanobacterial nighttime survival. PLoS Genet. 2018;

14: e1007301.

39. Whiteley AT, Pollock AJ, Portnoy DA. The PAMP c-di-AMP is essential for *Listeria monocytogenes*

growth in macrophages and rich but not minimal medium due to a toxic increase in (p)ppGpp. Cell

Host Microbe 2015; 17: 788-798.

40. Zeden MS, Schuster CF, Bowman L, Zhong Q, Williams HD, Gründling A. Cyclic di-adenosine

monophosphate (c-di-AMP) is required for osmotic regulation in *Staphylococcus aureus* but

dispensable for viability in anaerobic conditions. J. Biol. Chem. 2018; 293: 3180-3200.

41. Krüger L, Herzberg C, Warneke R, Poehlein A, Stautz J, Weiß M, et al. Two ways to convert a low-

affinity potassium channel to high affinity: Control of *Bacillus subtilis* KtrCD by glutamate. J.

Bacteriol. 2020; 202: e00138-20.

42. Zeden MS, Kviatkovski I, Schuster CF, Thomas VC, Fey PD, Gründling A. Identification of the main glutamine and glutamate transporters in *Staphylococcus aureus* and their impact on c-di-AMP production. Mol Microbiol. 2020; 113: 1085-1100.

43. Commichau FM, Gibhardt J, Halbedel S, Gundlach J, Stülke J. 2018. A delicate connection: c-di-AMP affects cell integrity by controlling osmolyte transport. Trends Microbiol. 26:175-185.

44. He J, Yin W, Galperin MY, Chou SH. Cyclic di-AMP, a second messenger of primary importance: tertiary structures and binding mechanisms. Nucleic Acids Res. 2020; 48: 2807-2829.

45. Stülke J, Krüger L. Cyclic di-AMP signaling in bacteria. Annu Rev Microbiol. 2020; 74: 159-179.

46. Mehne FMP, Schröder-Tittmann K, Eijlander RT, Herzberg C, Hewitt L, et al. Control of the diadenylate cyclase CdaS in *Bacillus subtilis*: An autoinhibitory domain limits c-di-AMP production. J Biol Chem. 2014; 289: 21098-21107.

47. Gundlach J, Dickmanns A, Schröder-Tittmann K, Neumann P, Kaesler J, Kampf J, et al. Identification, characterization and structure analysis of the c-di-AMP binding P_{II}-like signal transduction protein DarA. J Biol Chem. 2015; 290: 3069-3080.

48. Rosenberg J, Müller P, Lentes S, Thiele MJ, Zeigler DR, Tödter D, et al. ThrR, a DNA-binding transcription factor involved in controlling threonine biosynthesis in *Bacillus subtilis*. Mol. Microbiol. 2016; 101: 879-893.

49. Reuß DR, Rath H, Thürmer A, Benda M, Daniel R, Völker U, et al. Changes of DNA topology affect the global transcription landscape and allow rapid growth of a *Bacillus subtilis* mutant lacking carbon catabolite repression. Metab. Eng. 2018; 45: 171-179.

50. Kampf J, Gerwig J, Kruse K, Cleverley R, Dormeyer M, Grünberger A, et al. Selective pressure for biofilm formation in *Bacillus subtilis*: differential effect of mutations in the master regulator SinR on bistability. mBio 2018; 9: e00166-18.

51. Belitsky BR, Sonenshein AL. Role and regulation of *Bacillus subtilis* glutamate dehydrogenase genes. J Bacteriol. 1998; 180: 6298-6305.

52. Commichau FM, Gunka K, Landmann JJ, Stülke J. Glutamate metabolism in *Bacillus subtilis*: Gene

expression and enzyme activities evolved to avoid futile cycles and to allow rapid responses to

perturbations in the system. J Bacteriol. 2008; 190:3557-3564.

53. Klewing A, Koo BM, Krüger L, Poehlein A, Reuß D, Daniel R, et al. Resistance to serine in *Bacillus*

subtilis: identification of the serine transporter YbeC and of a metabolic network that links serine

and threonine metabolism. Environ. Microbiol. 2020; in press. doi: https://doi.org/10.1111/1462-

2920.15179.

54. Calvo RA, Kearns DB. FlgM is secreted by the flagellar export apparatus in *Bacillus subtilis*. J

Bacteriol. 2015; 197:8-91.

55. Wahome PG, Setlow P. Growth, osmotic downshock resistance and differentiation of *Bacillus*

subtilis lacking mechanosensitive channels. Arch Microbiol. 2008; 189: 49-58.

56. Hoffmann T, Boiangiu C, Moses S, Bremer E. Responses of *Bacillus subtilis* to hypotonic challenges:

physiological contributions of mechanosensitive channels to cellular survival. Appl Environ

Microbiol. 2008; 74: 2454-2460.

57. Kempf B, Bremer E. OpuA, an osmolytically regulated binding protein-dependent transport

system for the osmoprotectant glycine betaine in *Bacillus subtilis*. J Biol Chem. 1995; 270: 16701-

16713.

58. Tascón I, Sousa JS, Corey RA, Mills DJ, Griwatz D, Aumüller N, et al. Structural basis of proton-

coupled potassium transport in the KUP family. Nat Commun. 2020; 11: 626.

59. Krüger L, Herzberg C, Wicke D, Bähre H, Heidemann JL, Dickmanns, et al. A rendezvous of two

second messengers: The c-di-AMP receptor protein DarB controls (p)ppGpp synthesis in *Bacillus*

subtilis. https://doi.org/10.1101/2020.08.27.268672.

60. Peterson BN, Young MKM, Lou S, Wang J, Whiteley AT, Woodward JJ, et al. (p)ppGpp and c-di-

AMP homeostasis is controlled by CbpB in *Listeria monocytogenes*. mBio 2020; 11: e01625-20.

61. Larsson JT, Rogstam A, von Wachenfeldt C. Coordinated patterns of cytochrome bd and lactate

dehydrogenase expression in *Bacillus subtilis*. Microbiology 2005; 151: 3323-3335.

62. Gyan S, Shiohira Y, Sato I, Takeuchi M, Sato T. Regulatory loop between redox sensing of the

NADH/NAD$^+$ ratio by Rex (YdiH) and oxidation of NADH by NADH dehydrogenase Ndh in *Bacillus*

subtilis. J Bacteriol. 2006; 188: 7062-7071.

63. Qi Y, He Y, She Q, Larese-Casanova P, Li P, Chai Y. Heterogeneity in respiratory electron transfer

and adaptive iron utilization in a bacterial biofilm. Nat Commun. 2019; 10: 3702.

64. MacLellan SR, Wecke T, Helmann JD. A previously unidentified σ factor and two accessory proteins

regulate oxalate decarboxylase expression in *Bacillus subtilis*. Mol Microbiol. 2008; 69: 954-967.

65. Zhang X, Wang J, Feng Y, Ge J, Li W, Sun W, et al. Structure and molecular mechanism of an anion-

selective mechanosensitive channel of small conductance. Proc Natl Acad Sci USA. 2012; 109:

18180-18185.

66. Bilder P, Lightle S, Bainbridge G, Ohren J, Finzel B, Sun F, et al. The structure of the

carboxyltransferase component of acetyl-CoA carboxylase reveals a zinc-binding motif unique to

the bacterial enzyme. Biochemistry 2006; 45: 1712-1722.

67. Silvers MA, Pakhomova S, Neau DB, Silvers WC, Anzalone N, Taylor CM, Waldrop GL. Crystal

structure of carboxyltransferase from *Staphylococcus aureus* bound to the antibacterial agent

moiramide B. Biochemistry 2016; 55: 4666-4674.

68. Robertson RM, Yao J, Gajewski S, Kumar G, Martin EW, Rock CO, White SW. A two-helix motif

positions the lysophatidic acid acyltransferase active site for catalysis within the membrane

bilayer. Nat Struct Mol Biol 2017; 24: 666-671.

69. Pliotas C, Dahl ACE, Rasmussen T, Mahendran KR, Smith TK, Marius P, et al. The role of lipids in

mechanosensation. Nat Struct Mol Biol. 2015; 22: 991-998.

70. Rasmussen T, Rasmussen A, Yang L, Kaul C, Black S, Galbiati, et al. Interaction of the

mechanosensitive channel, MscS, with the membrane bilayer through lipid intercalation into

grooves and pockets. J Mol Biol. 2019; 431: 3339-3352.

71. Halpern YS, Lupo M. Glutamate transport in wild-type and mutant strains of *Escherichia coli*. J

Bacteriol. 1965; 90: 1289-1295.

72. Schirmer F, Ehrt S, Hillen W. Expression, inducer spectrum, domain structure, and function of

MopR, the regulator of phenol degradation in *Acinetobacter calcoaceticus* NCIB8250. J Bacteriol.

1997; 179: 1329-1336.

73. Zaprasis A, Bleisteiner M, Kerres A, Hoffmann T, Bremer E. Uptake of amino acids and their

metabolic conversion into the compatible solute proline confers osmoprotection to *Bacillus*

subtilis. Appl Environ Microbiol. 2015; 81: 250-259.

74. Liu G, Song G, Zhang D, Zhang D, Li Z, Lyu Z, et al. EF-G catalyzes tRNA translocation by disrupting

interactions between decoding center and codon-anticodon duplex. Nat Struct Mol Biol. 2014; 21:

817-824.

75. Peng BZ, Bock LV, Belardinelli, Peske F, Grubmüller H, Rodnina MV. Active role of elongation factor

G in maintaining the mRNA reading frame during translation. Sci Adv. 2019; 5: eaax8030.

76. Maracci C, Peske F, Dannies E, Pohl C, Rodnina MV. Ribosome-induced tuning of GTP hydrolysis

by a translational GTPase. Proc Natl Acad Sci USA. 2014; 111: 14418-14423.

77. Terakawa A, Natsume A, Okada A, Nishihata S, Kuse J, Tanaka K, et al. *Bacillus subtilis* 5'-

nucleotidases with various functions and substrate specificities. BMC Microbiol. 2016; 16: 249.

78. Gibhardt J, Hoffmann G, Turdiev A, Wang M, Lee VT, Commichau FM. C-di-AMP assists

osmoadaptation by regulating the *Listeria monocytogenes* potassium transporters KimA and

KtrCD. J Biol Chem. 2019; 294: 16020-16033.

79. Gundlach J, Commichau FM, Stülke J. Of ions and messengers: an intricate link between potassium,

glutamate and cyclic di-AMP. Curr Genet. 2018; 64:191-195.

80. Gunka K, Tholen S, Gerwig J, Herzberg C, Stülke J, Commichau FM. A high-frequency mutation in

Bacillus subtilis: requirements for the decryptification of the *gudB* glutamate dehydrogenase gene.

J Bacteriol. 2012; 194: 1036-1044.

81. Becker M, Börngen K, Nomura T, Battle AR, Marin K, et al. Glutamate efflux mediated by

Corynebacterium glutamicum MscCG, *Escherichia coli* MscS, and their derivatives. Biochim Biohys

Acta. 2013; 1828:1230-1240.

82. Wang Y, Cao G, Xu D, Fan L, Wu X, Ni X, et al. A novel *Corynebacterium glutamicum* L-glutamate

exporter. Appl Environ Microbiol. 2018; 84:e02691-17.

83. Nakayama Y, Yoshimura K, Iida H. A gain-of-function mutation in gating of *Corynebacterium*

glutamicum NCgl1221 causes constitutive glutamate secretion. Appl Environ Microbiol. 2012;

78:5432-5434.

84. Nakayama Y,Hashimoto KI, Kawasaki H, Martinec B. "Force-From-Lipids" mechanosensation in

Corynebacterium glutamicum. Biophys Rev. 2019; 11:327-333.

85. Tascon I, Sousa JS, Corey RA, Mills DJ, Griwatz D, et al. Structural basis of proton coupled

potassium transport in the KUP family. Nat Commun. 2020; 11:626.

86. Diskowski M, Mikusevic V, Stock C, Hänelt I. 2015. Functional diversity of the superfamily of K^+

transporters to meet various requirements. Biol. Chem. 396:1003-1014.

87. Sambrook J, Russell D. 2001. Molecular Cloning: A Laboratory Manual, Cold Spring Harbor

Laboratory, Cold Spring Harbor.

88. Kunst F, Rapoport G. Salt stress is an environmental signal affecting degradative enzyme synthesis

in *Bacillus subtilis*. J Bacteriol. 1995; 177:2403-2407.

89. Koo BM, Kritikos G, Farelli JD, Todor H, Tong K, Kimsey H, et al. Construction and analysis of two

genome-scale deletion libraries for *Bacillus subtilis*. Cell Syst. 2017; 4: 291-305.

90. Wacker I, Ludwig H, Reif I, Blencke HM, Detsch C, Stülke J. The regulatory link between carbon

and nitrogen metabolism in *Bacillus subtilis*: regulation of the *gltAB* operon by the catabolite

control protein CcpA. Microbiology. 2003; 149: 3001-3009.

91. Reuß DR, Altenbuchner J, Mäder U, Rath H, Ischebeck T, Sappa PK, et al. Large-scale reduction of

the *Bacillus subtilis* genome: consequences for the transcriptional network, resource allocation,

and metabolism. Genome Res. 2017; 27:289-299.

92. Barbe V, Cruveiller S, Kunst F, Lenoble P, Meurice G, Sekowska A, et al. From a consortium

sequence to a unified sequence: the *Bacillus subtilis* 168 reference genome a decade later.

Microbiology 2009; 155:1758-1775.

93. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, et al. Geneious basic: an

integrated and extendable desktop software platform for the organization and analysis of

sequence data. Bioinformatics 2012; 28:1647–1649.

94. Diethmaier C, Pietack N, Gunka K, Wrede C, Lehnik-Habrink M, Herzberg C, et al. A novel factor

controlling bistability in *Bacillus subtilis*: the YmdB protein affects flagellin expression and biofilm

formation. J Bacteriol. 2011; 193: 5997-6007.

95. Stülke J, Martin-Verstraete I, Zagorec M, Rose M, Klier A, Rapoport G. Induction of the *Bacillus*

subtilis ptsGHI operon by glucose is controlled by a novel antiterminator, GlcT. Mol Microbiol.

1997; 25: 65-78.

96. Eymann C, Homuth G, Scharf C, Hecker M. *Bacillus subtilis* functional genomics: global

characterization of the stringent response by proteome and transcriptome analysis. J Bacteriol.

2002; 184: 2500-2520.

97. Nicolas P, Mäder U, Dervyn E, Rochat T, Leduc A, Pigeonneau N, et al. Condition-dependent

transcriptome reveals high-level regulatory architecture in *Bacillus subtilis*. Science 2012; 335:

1103-1106.

98. Mäder U, Nicolas P. Array-based approaches to bacterial transcriptome analysis. In Methods in

Microbiology, Vol. 39, eds. C. Harwood, A. Wipat (Cambridge Academic Press), 151-182.

99. Zhu B, Stülke J. *Subti*Wiki in 2018: from genes and proteins to functional network annotation of

the model organism *Bacillus subtilis*. Nucleic Acids Res. 2018; 46: D743-D748.

100. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: a software

environment for integrated models of biomolecular interaction networks. Genome Res. 2003; 13:

2498-504.

101. The GIMP Development Team. GIMP. 2019; Retrieved from https://www.gimp.org.

1011

 SUPPORTING INFORMATION CAPTIONS

 S1 Supporting Information

 Global gene expression in the *B. subtilis* wild type strain in response to the potassium concentration

 and the nitrogen source.

 S1 Table

 Transcriptomic data for the genes that are most strongly affected by the absence of c-di-AMP (Ammonium

 0.1 mM KCl).

 S2 Table.

 Transcriptomic data of genes involved in c-di-AMP, potassium, and glutamate homeostasis.

 S1 Fig. Acquisition of beneficial mutations allows growth of a c-di-AMP deficient strain in the

 presence of glutamate.

 Growth assay of *B. subtilis* wild type, GP2222 (Δ*dac*), and the isolated glutamate suppressor mutants.

 B. subtilis strains were cultivated in MSSM minimal medium with 0.1 mM KCl and ammonium. The cells

 were harvested, washed, and the OD_{600} was adjusted to 1.0. Serial dilutions were dropped onto MSSM

 minimal plates with the indicated potassium concentration and ammonium or glutamate, or on LB

 plates.

 S2 Fig. Potassium and glutamate influenced promoters.

 The expression of the high-affinity potassium transporter *kimA* (A) and the glutamate synthase *gltA* (B)

 was assessed by fusion of the promoter to the reporter gene *lacZ*. The cells harboring the promoter-

 fusion were cultivated with ammonium or glutamate at 0.1 mM KCl (*kimA*) or ammonium and different

 potassium concentrations (0.1, 5, 20 mM; *gltA*). Promoter activity was analyzed by quantification of β-

1038   galactosidase activity. Statistical analysis was performed using a one-way ANOVA, followed by Tukey's

1039   multiple comparisons test (**** P < 0.0001).

1040

1041   **S3 Fig. Two-step adaptation to complex medium.**

1042   Growth assay of *B. subtilis* wild type, GP2779 (Δ*dac* Δ*darB*), GP2420 (Δ*dac* Δ*darA*) and the isolated

1043   glutamate suppressor mutants. *B. subtilis* strains were precultivated in LB medium. The cells were

1044   harvested, washed, and the OD_{600} was adjusted to 1.0. Serial dilutions were dropped onto MSSM

1045   minimal plates with 0.1 mM KCl and ammonium or on LB plates.

1046

1047   **S4 Fig. Mutations in the mechanosensitive channel YfkC.** Mutations in the *yfkC* gene were obtained

1048   in the suppressor screen with glutamate (Table 1). The amino acid substitution Arg42 to Trp was

1049   observed in three independently isolated clones. In *E. coli* localization of S58 was observed to be highly

1050   dependent on the closing state of the channel (Miller *et al.*, 2003), thus, it appears likely that amino

1051   acid residue 42 in *B. subtilis* is located close to the gate.

1052

1053   **S5 Fig. Mutations in the acetyl-CoA carboxylase subunits.**

1054   Mutations obtained in the suppressor screen with glutamate (Table 1) and LB (Table 4). All mutations

1055   were single amino acid substitutions from independently isolated clones. Both mutations are located

1056   in highly conserved regions of AccA and AccC (hatched boxes) close to the active center.

1057

1058   **S6 Fig. Mutations in the acyl-ACP:1-acylglycerolphosphate acyltransferase.** Mutations in the *plsC*

1059   gene were obtained in the suppressor screen with glutamate (Table 1). All mutations were single amino

1060   acid substitutions from independently isolated clones. All mutations are located in functionally

1061   important regions of the enzyme.

1062

1063

S7 Fig. Mutations in the elongation factor G (FusA). Mutations in the *fusA* gene were obtained in the

suppressor screen of GP3054 (Δ*dac* Δ*aimA*) with glutamate (Table 3). The amino acid substitution

Ala21 to Thr was observed in one suppressor. Ala21 is adjacent to Asp20, an amino acid residue that

binds the Mg ion and is therefore crucial for GTPase activity of the protein (Maracci *et al.*, 2014). In the

two other suppressors Ala579 was mutated to Val. This amino acid residue is located in the conserved

loop II that is required for proper translocation (Peng *et al.*, 2019).

1070

S8 Fig. Effect of the deletion of the glutamate transporters in a c-di-AMP deficient strain.

Growth assay of *B. subtilis* wild type, GP2222 (Δ*dac*), GP3054 (Δ*dac* Δ*aimA*), GP2248 (Δ*dac* Δ*gltT*) and

the isolated glutamate suppressor mutants. *B. subtilis* strains were cultivated in MSSM minimal

medium with 0.1 mM KCl and ammonium. The cells were harvested, washed, and the OD_{600} was

adjusted to 1.0. Serial dilutions were dropped onto MSSM minimal plates with the indicated potassium

concentration and ammonium or glutamate, or on LB plates.

1077

 Figure legends

 Fig. 1. A suppressor screen with a strain lacking c-di-AMP reveals the complexity of glutamate

 toxicity.

 GP2222 (Δ*dac*), GP2814 (Δ*dac yfkC**), GP2223 (Δ*dac nhaK**), GP2259 (Δ*dac* Δ*gltT*), and GP3054 (Δ*dac*

 Δ*aimA*) were cultivated in MSSM medium with glutamate and 0.1 mM KCl and suppressors were

 isolated and analyzed by whole genome sequencing. Note that for GP2222 (Δ*dac*), 14 mutants were

 isolated in this study and one mutant was isolated previously [41] but studied in depth here, thus

 resulting in a total number of 15 mutants.

 Fig. 2. A suppressor screen with a strain lacking c-di-AMP on complex medium.

 GP2222 (Δ*dac*), GP2420 (Δ*dac* Δ*darA*), or GP2779 (Δ*dac* Δ*darB*) were cultivated on LB agar plates and

 incubated under aerobic or anaerobic conditions as indicated. The table shows how the parental strain

 evolved, and under which conditions suppressors could be isolated. The suppressor mutants were

 analyzed by whole genome sequencing.

 Fig. 3. Regulatory network of the most strongly up- and downregulated genes as a result of the lack

 of c-di-AMP.

 Expression ratios of *B. subtilis* wild type and GP2222 (Δ*dac*) with ammonium and 0.1 mM KCl were

 calculated and the most up- or downregulated genes were clustered into operons, regulons, and

 grouped depending on their physiological function. Regulators are depicted as large blue nodes, and

 regulated operons or genes as yellow and orange nodes for up- and downregulated genes,

 respectively). Arrows: green: activation; gray: sigma-factor; red: repression.

 Fig. 4. The influence of the mechanosensitive channel YfkC on growth of a *B. subtilis* strain lacking c-

 di-AMP in the presence of glutamate.

1104   *B. subtilis* strains were cultivated in MSSM minimal medium with 0.1 mM KCl and glutamate. Cells were

1105   harvested, washed, and the OD_{600} was adjusted to 1.0. Serial dilutions were dropped onto MSSM

1106   minimal plates with the indicated potassium concentration and ammonium or glutamate, or on LB

1107   plates.

1108

Fig. 5. Identification of AimA as a novel low-affinity glutamate transporter.

(A) *E. coli* JM109 was transformed with pGP2987 (AimA) and pWH844 (empty vector) and growth of *E.*

coli JM109 was assessed on MSSM minimal medium with glucose and ammonium or glutamate as

carbon and nitrogen source. (B) Growth assay of *B. subtilis* glutamate transporter mutants. *B. subtilis*

strains were cultivated in MSSM minimal medium with 0.1 mM KCl and ammonium. Cells were

harvested, washed, and the OD_{600} was adjusted to 1.0. Serial dilutions were dropped onto MSSM

minimal plates with the indicated potassium concentration and ammonium or glutamate (1, 5, or 20

mM). (C) Determination of glutamate transporter kinetics. The growth rates of *B. subtilis* strains

expressing either AimA (upper panel) or GltT (lower panel) as the only glutamate transporter were

used to determine the apparent glutamate transporter kinetics of AimA and GltT. A *B. subtilis* strain

lacking all glutamate transporters (Δ*gltT* Δ*aimA* Δ*gltP*; light grey) served as a control. The strains were

grown in MSSM minimal medium with 0.1 mM or 5 mM KCl and various glutamate concentrations. The

growth rate was plotted against the glutamate concentration. (D) Kinetic parameters (K_S and v_{max}

values) for AimA and GltT. (E) Expression of *gltT*, *aimA*, and *gltP*. The expression levels of the genes

encoding the glutamate transporters GltT, AimA, and GltP were extracted from the transcriptomic

analysis.

1125

1126

Figure 1, Krüger et al., 2020

Figure 2, Krüger et al., 2020

Figure 3, Krüger et al., 2020

Figure 4, Krüger et al., 2020

Figure 5, Krüger et al., 2020