Title: A single nuclei transcriptomic analysis of the Atlantic salmon gill through smoltification

and seawater transfer

Authors: Alexander C. West[1], Yasutaka Mizoro[2], Shona H. Wood[1], Louise M. Ince[3], Marianne

Iversen[1], Even H. Jørgensen[1], Torfinn Nome[4], Simen Rød Sandve[4], Andrew S. I. Loudon[2] and

David G. Hazlerigg[1]

[1]Arctic Chronobiology and Physiology, University of Tromsø, Framstredet 42, 9019 Tromsø,

Norway

[2]Division of Diabetes, Endocrinology & Gastroenterology, School of Medical Sciences, Faculty

of Biology, Medicine and Health, University of Manchester, UK

[3]University of Geneva, Centre Médical Universitaire (CMU), Department of Pathology and

Immunology, Switzerland

[4]Centre for Integrative Genetics, Department of Animal and Aquaculture Sciences,

Norwegian University of Life Sciences, 1432 Ås, Norway

*Correspondence: david.hazlerigg@uit.no

Abstract

17 Anadromous salmonids begin life adapted to the freshwater environments of their natal

18 streams before a developmental transition, known as smoltification, transforms them into

19 marine-adapted fish. In the wild, the extending photoperiods of spring stimulates

20 smoltification, typified by radical reprogramming of the gill from an ion-absorbing organ to

21 ion-excreting organ. Prior work has highlighted the role of specialized "mitochondrion-rich"

cells in delivering this phenotype. However, transcriptomic studies identify thousands of smoltification-driven differentially regulated genes, indicating that smoltification causes a multifaceted, multicellular change; but direct evidence of this is lacking.

Here, we use single-nuclei RNAseq to characterize the Atlantic salmon gill during smoltification and seawater transfer. We identify 20 distinct clusters of nuclei, including known, but also novel gill cell types. These data allow us to isolate cluster-specific, smoltification-induced changes in gene expression. We also show how cellular make-up of the gill changes through smoltification. As expected, we noted an increase in the proportion of seawater mitochondrion-rich cells, however, we also identify a reduction of several immune-related cells. Overall, our results provide unrivaled detail of the cellular complexity in the gill and suggest that smoltification triggers unexpected immune reprogramming directly preceding seawater entry.

Keywords: Atlantic salmon, smoltification, photoperiod, seasonal, gill

Introduction

The Atlantic salmon migrates between fresh and seawater environments [1]. Atlantic salmon eggs hatch in freshwater streams where they develop for 1-4 years. On reaching a critical size threshold, young "parr" animals are sensitized by several weeks of winter photoperiod (day-lengths), after which long, summer-like photoperiods stimulates the parr to transform into a marine-adapted "smolt" fish [2]. This process, known as smoltification, drives divergent expression of endocrine factors that collectively deliver phenotypic remodeling, of length, weight, silvering, and in particular: gill physiology [1].

The salmonid gill is a complex multifunctional organ, essential for gas exchange,

nitrogenous waste excretion, pH balance and osmoregulation [3]. It is also a major mucosal

immune barrier harboring a dedicated lymphoid tissue [4]. Structurally, the gills are arranged in

symmetrical arches, each of which are populated by numerous filament structures, which are

themselves densely flanked with lamellae. The gill is composed of seven major cell types [5].

Pavement cells (PVCs) have an enlarged surface area on the apical membrane, and form the

majority of the epithelium [6]. Pillar cells (PCs), which are structural cells, define the blood

spaces within the lamellae [7]. Goblet cells (GCs) reside in the filament epithelium and excrete

mucus [8]. Non-differentiated progenitor cells (NDCs) colonize basal and intermediate layers of

the gill epithelium [9]. Chemosensory neuroepithelial cells (NECs) lie along the length of the

efferent edge of the gills and are innervated by the central nervous system [10]. Mitochondrion-

rich cells (MRCs) and their adjacent accessory cells (ACs), finally, are located at the trough

between two lamellae where they abundantly express the channels and pumps required to

maintain the osmotic gradients between blood plasma and both fresh- and seawater [11–13].

Under freshwater, Na^+ ions are directly or indirectly exchanged for protons across the

apical membrane then transported into the blood *via* the sodium potassium ATPase (NKA) on

the basolateral membrane [14–16]. Cl^- ions, meanwhile, are exchanged or channeled across the

apical membrane then enter the blood through an undefined channel [17–20]. Under saltwater,

basolateral NKA generates a chemical and electrical gradient, motivating both loss of Cl^- ions

via apical CFTR channels and paracellular escape of Na^+ ions [13,21] (reviewed in [22]).

Smoltification converts the Atlantic salmon gill from a freshwater-adapted organ to a

seawater-adapted organ. Rising cortisol and growth hormone along with falling prolactin

propels smoltification. This change in endocrinology coincides with a switch in anatomical and

molecular phenotypes of MRCs and ACs, cell types which to date comprise the major focus of

smoltification of gill physiology [1,12,23]. Smoltification is, however, a complex developmental

transition and the smolt gill phenotype likely extends far beyond changing MRC and AC cell

phenotype. Therefore, to realize the complexities of smoltification-driven changes in gill

physiology we adopt a new strategy, single-nuclei RNAseq, which provides transcriptional

responses to smoltification and seawater transfer at individual nuclei-level resolution.

Results & Discussion

A single-nuclei survey of Atlantic salmon gill cells

We profiled 18,844 individual nuclei from eight Atlantic salmon gill samples from four distinct

physiological states (Figure 1A). To define shared correlation structure across datasets we

pooled replicate samples and integrated all four states using diagonalized canonical

correlation analysis followed by L2 normalization. We next identified pairs of mutual nearest

neighbors (MNNs) to identify anchors: cells that represent shared biological states across

datasets. Anchors were then used to calculate "correction" vectors allowing all fours states to

be jointly analyzed as an integrated reference [24]. Unsupervised graph clustering partitioned

the nuclei into 20 clusters, which we visualized using a uniform manifold approximation and

projection (UMAP) dimension reduction technique (Figure 1B).

Lists of co-expressed marker genes define individual clusters. We categorized

individual clusters using gene ontology analysis of marker gene lists and unique expression of

known marker genes. This approach allowed us to infer cell types including fresh- and

seawater MRCs, ACs, neuroendothelial cells, goblet cells, non-differentiated cells, pillar cells,

lymphatic endothelial cells and several types of blood cell. We also identified a novel

population of fibrocyte-like cells, and several types of vascular- and endothelial-like cells that

partitioned across several clusters, together suggesting greater complexity in gill cytology that

previously appreciated (Figure 1B).

We then defined expression signatures for each cell cluster. Our analysis re-identified

several known marker genes, but also identifying several novel cell-type markers (Figure 1C).

For example, the accessory cell signature included highly restricted expression of Slc26a6, an

apical membrane Cl^-/HCO_3 exchanger, associated with gill function but until now

misattributed to expression within MRCs [25,26]. We were interested to note that the markers

defining the erythrocyte population, including beta-globin, were expressed widely among all

cell types. It is unclear exactly what role extra-erythroid haemoglobin plays in the gill,

however, mammalian studies suggest that haemoglobin, in addition to its oxygen carrying

capacity, may play an antimicrobial role [27]. As a major mucosal immune barrier, this capacity

may be pertinent to the gill [28].

Major changes in cell composition during smoltification

To understand how gene expression and cellular complexity changes within the gill during

smoltification and seawater transfer we compared the snRNAseq profiles at different

developmental points (Fig 2A, for confirmation of smolt status see[29]). The abundance of six

nuclei clusters changed dramatically (>3 fold change in percentage abundance) during

smoltification (Figure 2B). SW MRCs increased in proportion steadily from T1-T4, consistent

with previous descriptions of Atlantic salmon gill physiology. We also observed a marked

increase in vascular cell number, with the major differences occurring between T2 and T3,

suggesting that this vascular cell cluster proliferate in line with growth rates (Figure 2C).

Interestingly, four immune-related nuclei clusters representing T cells, myeloid cells, dendritic

cells and lymphatic endothelial cells fell dramatically during smoltification (Figure 2D).

Changes in cell abundance occurred with a similar profile in all immune-associated cell

clusters, with consistent decline observed between T1-T3. In contrast, 24h SW transfer does

not appear to affect immune-cell abundance directly. These results highlight the complex and

dynamic changes in cellular composition that occur in the gill during smoltification.

Cell cluster-specific expression of smoltification-associated factors

Next, we wanted to identify cluster types where smoltification drives changes in local gene

regulation. For statistical power, we cross-referenced our snRNAseq analysis with whole gill

RNAseq analysis of T1-T3, identifying 9746 genes differentially regulated by smoltification

(FDR <0.01). Pearson clustering of these genes resolved five major clusters that were

associated with immune response, structural morphogenesis, autophagy, catabolism and

mitochondrial respiration (Figure 3A). Within our analysis we identified a number of "classical"

smoltification-related genes. As expected, CFTR was highest under constant light (LL), and was

highly localized in expression to MRCs (Figure 3B). We also identified the reciprocal regulation

of sodium-potassium ATPase subunits, specifically, suppression of NKAa1a and increase in

NKAa1b [12]. Inspection of cellular localization within our snRNAseq dataset showed that

expression of these genes were, as anticipated, highest within the MRCs and ACs (Figure 3B).

Our previous work identified genes whose expression are predicated on exposure to

several weeks of short-photoperiod exposure[29]. In Atlantic salmon, these "winter-dependent"

genes are analogous to vernalization dependent genes in *Arabidopsis*[30], where a dosage of

exposure to a winter-like stimulus (in *Arabidopsis*, cold; in Atlantic salmon, short photoperiod)

controls the presentation of a seasonal phenotype under summer-like stimulus (in

Arabidopsis, warmth and long days; in Atlantic salmon, long photoperiod). Winter-dependent

genes are therefore intrinsically linked to unidirectional smolt development, and may play a

mechanistic role in pre-adaptation of the gill for seawater migration. Surprisingly, canonical

markers of smolt status, including the reciprocal expression of NKA subunits, are not winter

dependent, meaning that their expression is passive to photoperiod rather than life history

progression[29].

Using our RNAseq dataset we identified novel, winter dependent genes. Next, we

mined our snRNAseq dataset to identify the cell clusters that express these factors (Figure S1).

Of particular interest was Cuzd1, a gene associated with carcinogenesis, whose expression

was tightly localized to non-differentiated cells [31]. We also identified Rhag, an ammonium

transporter thought to be erythrocyte specific in mammals, but here predominantly expressed

in the vascular cell (VC 3) cluster [32]; and Hg2a (CD74) a ubiquitously expression multi-

functional protein linked to immune defense [33]. Taken together our data show that

smoltification engages all gill cells in diverse regulatory phenotypes.

Cell cluster-specific expression of seawater transfer-associated factors

Smoltification manifests when the Atlantic salmon smolts migrate downstream and arrive in

the marine environment, thereby committing to an oceanic life [1]. To gain insight into this

critical step we compared RNA profiles of gill samples between smolts in freshwater and 24h

in seawater using whole gill RNAseq, identifying 144 induced and 107 suppressed genes

(FDR<0.01). Gene ontologies showed that the induced gene cohort was significantly

associated with keratinization, whereas the suppressed gene cohort related to immune

function (Figure 4A). We cross-referenced our whole gill RNAseq data against our snRNAseq

data to isolate cell-type specificity of gene expression. These data highlight the restricted

expression of key up-regulated genes (Figure 4B). For example, we localize the expression of

an enzyme involved in both ionic and acid/base balance, carbonic anhydrase, to MRCs [34,35].

We also show that ATP-binding cassette sub-family A member 12 (Abca12), a gene important

in epidermal lipid barrier formation [36], is broadly expressed, but particularly concentrated in

MRCs (SW), pavement, vascular, and non-differentiated cells. Interestingly, we show that a

protein chaperone that helps regulate chromatin state, nucleoplasmin [37,38], is expressed

specifically in non-differentiated, vascular and pavement cells groups, suggesting that these

cell types undergo a change in chromatin status under seawater exposure.

Conclusions

Our results bring insightful cellular resolution to the complexity of the Atlantic salmon gill and

the compositional changes that occur during smoltification. Of particular interest was the

suppression of immune cell types, which correlates with reduction in immune-related genes

and suppression of immune function during smoltification [39–41]. These data are a puzzle. The

marine environment is awash with parasites, bacteria and viruses to which the salmon is

potentially vulnerable, so loss of immune function would make little sense. Future work

should focus on why and how the immune system is affected in aquaculture. Conceivably

these data point towards an adaptive immunological reprogramming that helps to avoid

immune shock when the salmon transition between the distinctive pathogen complements of

fresh- and seawater habitats [42,43]. Alternatively, artificial smolt production may drive abnormal

immunosuppression. The constant light routinely used to stimulate smolts would profoundly

undermine the immune defenses of mammals *via* disruption of the circadian clock [44].

Our data also shows that smoltification-driven transcriptional regulation occurs not

only in MRCs and ACs, but also in other distinctive cell types including pavement cells, vascular

cells and non-differentiated cells. We anticipate that novel gene function within the context

of cell function will be a priority for future investigation, and will be assisted by the novel suite

of marker genes which we present here.

<u>References</u>

1.   Stefansson, S. O., Björnsson, B. T., Ebbesson, L. O. E. & McCormick, S. D.

Smoltification. in *Fish Larval Physiology* 639–681 (2008).

2.   Strand, J. E. T., Hazlerigg, D. & Jørgensen, E. H. Photoperiod revisited: is there a critical

day length for triggering a complete parr–smolt transformation in Atlantic salmon

Salmo salar? *J. Fish Biol.* **93**, 440–448 (2018).

3.   Evans, D. H., Piermarini, P. M. & Choe, K. P. The multifunctional fish gill: Dominant site

of gas exchange, osmoregulation, acid-base regulation, and excretion of nitrogenous

waste. *Physiol. Rev.* **85**, 97–177 (2005).

4.   Koppang, E. O. *et al.* Salmonid T cells assemble in the thymus, spleen and in novel

interbranchial lymphoid tissue. *J. Anat.* **217**, 728–739 (2010).

5.   Wilson, J. M. & Laurent, P. Fish gill morphology: Inside out. *J. Exp. Zool.* **293**, 192–213

(2002).

6.   Laurent, P. & Dunel-Erb, S. *The Pseudobranch: Morphology and Function*. *Fish*

Physiology vol. 10 (1984).

7.   Newstead, J. D. Fine structure of the respiratory lamellae of teleostean gills. *Zeitschrift*

für Zellforsch. und Mikroskopische Anat. **79**, 396–428 (1967).

8.   Rojo, M. C., Blánquez, M. J. & González, M. E. A histochemical study of the distribution

of lectin binding sites in the developing branchial area of the trout Salmo trutta. *J.*

Anat. **189**, 609–621 (1996).

9.   Laurent, P., Dunel-Erb, S., Chevalier, C. & Lignon, J. Gill epithelial cells kinetics in a

freshwater teleost, Oncorhynchus mykiss during adaptation to ion-poor water and

hormonal treatments. *Fish Physiol. Biochem.* **13**, 353–370 (1994).

10.   Dunel-Erb, S., Chevalier, C. & Laurent, P. Distribution of neuroepithelial cells and

neurons in the trout gill filament: comparison in spring and winter. *Can. J. Zool.* **72**,

1794–1799 (1994).

11.   Hootman, S. R. & Philpott, C. W. Accessory cells in teleost branchial epithelium. *Am. J.*

Physiol. - Regul. Integr. Comp. Physiol. **7**, (1980).

12.   Mccormick, S. D., Regish, A. M., Christensen, A. K. & Björnsson, B. T. Differential

regulation of sodium – potassium pump isoforms during smolt development and

seawater exposure of Atlantic salmon. 1142–1151 (2013) doi:10.1242/jeb.080440.

13.   Sardet, C., Pisam, M. & Maetz, J. The surface epithelium of teleostean fish gills. *J. Cell*

Biol. **80**, 96–117 (1979).

14.   Edwards, S. L., Tse, C. M. & Toop, T. Immunolocalisation of NHE3-like

immunoreactivity in the gills of the rainbow trout (Oncorhynchus mykiss) and the

blue-throated wrasse (Pseudolabrus tetrious). *J. Anat.* **195**, 465–469 (1999).

15.   Yan, J. J., Chou, M. Y., Kaneko, T. & Hwang, P. P. Gene expression of Na+/H+

exchanger in zebrafish H+-ATPase-rich cells during acclimation to low-Na+ and acidic

environments. *Am. J. Physiol. - Cell Physiol.* **293**, 1814–1823 (2007).

16.   Hirata, T. *et al.* Mechanism of acid adaptation of a fish living in a ph 3.5 lake. *Am. J.*

Physiol. - Regul. Integr. Comp. Physiol. **284**, (2003).

17.   Hiroi, J., McCormick, S. D., Ohtani-Kaneko, R. & Kaneko, T. Functional classification of

226   mitochondrion-rich cells in euryhaline Mozambique tilapia (Oreochromis

227   mossambicus) embryos, by means of triple immunofluorescence staining for Na+/K+-

228   ATPase, Na +/K+/2Cl- cotransporter and CFTR anion channel. *J. Exp. Biol.* **208**, 2023–

229   2036 (2005).

230   18.   Hiroi, J., Yasumasu, S., McCormick, S. D., Hwang, P. P. & Kaneko, T. Evidence for an

231   apical Na-Cl cotransporter involved in ion uptake in a teleost fish. *J. Exp. Biol.* **211**,

232   2584–2599 (2008).

233   19.   Tang, C. H., Hwang, L. Y. & Lee, T. H. Chloride channel CLC-3 in gills of the euryhaline

234   teleost, tetraodon nigroviridis: Expression, localization and the possible role of

235   chloride absorption. *J. Exp. Biol.* **213**, 683–693 (2010).

236   20.   Preest, M. R., Gonzalez, R. J. & Wilson, R. W. A pharmacological examination of Na+

237   and Cl- transport in two species of freshwater fish. *Physiol. Biochem. Zool.* **78**, 259–

238   272 (2005).

239   21.   Marshall, W. S., Lynch, E. M. & Cozzi, R. R. F. Redistribution of immunofluorescence of

240   CFTR anion channel and NKCC cotransporter in chloride cells during adaptation of the

241   killifish Fundulus heteroclitus to sea water. *J. Exp. Biol.* **205**, 1265–1273 (2002).

242   22.   Evans, D. H. Freshwater fish gill ion transport: August Krogh to morpholinos and

243   microprobes. *Acta Physiol. (Oxf).* **202**, 349–359 (2011).

244   23.   Lorgen, M. *et al.* Functional divergence of type 2 deiodinase paralogs in the Atlantic

245   salmon. *Curr. Biol.* **25**, 936–941 (2015).

24.   Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-

1902.e21 (2019).

25.   Leguen, I., Le Cam, A., Montfort, J., Peron, S. & Fautrel, A. Transcriptomic analysis of

trout gill ionocytes in fresh water and sea water using laser capture microdissection

combined with microarray analysis. *PLoS One* **10**, 1–22 (2015).

26.   Boyle, D., Clifford, A. M., Orr, E., Chamot, D. & Goss, G. G. Mechanisms of Cl- uptake in

rainbow trout: Cloning and expression of slc26a6, a prospective Cl-/HCO3- exchanger.

Comp. Biochem. Physiol. -Part A Mol. Integr. Physiol. **180**, 43–50 (2015).

27.   Saha, D. *et al.* Hemoglobin Expression in Nonerythroid Cells: Novel or Ubiquitous? *Int.*

J. Inflam. **2014**, (2014).

28.   Koppang, E. O., Kvellestad, A. & Fischer, U. *Mucosal Health in Aquaculture Fish*

mucosal immunity: gill. *Mucosal Health in Aquaculture* (Elsevier Inc., 2015).

doi:10.1016/B978-0-12-417186-2/00005-4.

29.   Iversen, M. *et al.* RNA profiling identifies novel, photoperiodhistory dependent

markers associated with enhanced saltwater performance in juvenile Atlantic salmon.

PLoS One **15**, 1–21 (2020).

30.   Song, J., Angel, A., Howard, M. & Dean, C. Vernalization - a cold-induced epigenetic

switch. *J. Cell Sci.* **125**, 3723–3731 (2012).

31.   Liaskos, C., Rigopoulou, E. I., Orfanidou, T., Bogdanos, D. P. & Papandreou, C. N.

CUZD1 and Anti-CUZD1 Antibodies as Markers of Cancer and Inflammatory Bowel

Diseases. **2013**, (2013).

32.   Tilley, L. *et al.* A new blood group system , RHAG : three antigens resulting from amino

acid substitutions in the Rh-associated glycoprotein. 151–159 (2010)

doi:10.1111/j.1423-0410.2009.01243.x.

33.   Su, H., Na, N., Zhang, X. & Zhao, Y. The biological function and significance of CD74 in

immune diseases. *Inflamm. Res.* **66**, 209–216 (2017).

34.   Houde, A. L. S. *et al.* Salmonid gene expression biomarkers indicative of physiological

responses to changes in salinity and temperature, but not dissolved oxygen. *J. Exp.*

Biol. **222**, (2019).

35.   Zbanyszek, R. & Smith, L. S. Changes in carbonic anhydrase activity in coho salmon

smolts resulting from physical training and transfer into seawater. *Comp. Biochem.*

Physiol. -- Part A Physiol. **79**, 229–233 (1984).

36.   Akiyama, M. The roles of ABCA12 in epidermal lipid barrier formation and

keratinocyte differentiation. *Biochim. Biophys. Acta - Mol. Cell Biol. Lipids* **1841**, 435–

440 (2014).

37.   Bouleau, A. *et al.* Maternally inherited npm2 mRNA is crucial for egg developmental

competence in zebrafish. *Biol. Reprod.* **91**, 1–9 (2014).

38.   Chen, P. *et al.* Nucleoplasmin is a limiting component in the scaling of nuclear size

with cytoplasmic volume. *J. Cell Biol.* **218**, 4063–4078 (2019).

39.   Jensen, I., Overrein, M. C., Fredriksen, B. N., Strandskog, G. & Seternes, T. Differences

in smolt status affect the resistance of Atlantic salmon (Salmo salar L.) against

infectious pancreatic necrosis, while vaccine-mediated protection is unaffected. *J. Fish*

Dis. **42**, 1271–1282 (2019).

40.   Johansson, L. H., Timmerhaus, G., Afanasyev, S., Jørgensen, S. M. & Krasnov, A.

Smoltification and seawater transfer of Atlantic salmon (Salmo salar L.) is associated

with systemic repression of the immune transcriptome. *Fish Shellfish Immunol.* **58**,

33–41 (2016).

41.   Nuñez-Ortiz, N. *et al.* Atlantic salmon post-smolts adapted for a longer time to

seawater develop an effective humoral and cellular immune response against

Salmonid alphavirus. *Fish Shellfish Immunol.* **82**, 579–590 (2018).

42.   Lee, S.-Y. & Eom, Y.-B. Analysis of Microbial Composition Associated with Freshwater

and Seawater. *Biomed. Sci. Lett.* **22**, 150–159 (2016).

43.   Wang, Y. *et al.* Comparison of the levels of bacterial diversity in freshwater, intertidal

wetland, and marine sediments by using millions of illumina tags. *Appl. Environ.*

Microbiol. **78**, 8264–8271 (2012).

44.   Scheiermann, C., Gibbs, J., Ince, L. & Loudon, A. Clocking in to immunity. *Nature*

Reviews Immunology vol. 18 423–437 (2018).

45.   Matson, K. J. E. *et al.* Isolation of adult spinal cord nuclei for massively parallel single-

nucleus RNA sequencing. *J. Vis. Exp.* **2018**, 1–12 (2018).

Material and Methods

Animal welfare statement

The Atlantic salmon smoltification experiment was conducted as part of the routine, smolt

production at Kårvik havbruksstasjonen, approved by the Norwegian Animal Research

Authority (NARA) for the maintenance of stock animals for experiments on salmonids. This is

in accordance with Norwegian and European legislation on animal research.

Experimental Design

Atlantic salmon (*Salmo salar*, Aquagene commercial stain) were raised from hatching in freshwater, under continuous light (LL, > 200 lux at water surface) at ambient temperature (~10°C). Juvenile salmon were housed in 500 L circular tanks and fed continuously with pelleted salmon feed (Skretting, Stavanger, Norway). At seven months of age parr (mean weight 49.5g) were sampled for T1 (experiment start). Two days later remaining parr were equally distributed between two 100L circular tanks, and over the next seven days the photoperiod was incrementally reduced to a short photoperiod (SP, 8h light:16h darkness). T2 sampling occurred on experimental day 53 (44 days on SP), remaining parr were transferred back to LL on experimental day 60. T3 sampling occurred on experimental day 110 (50 days after return to LL), then a sub-cohort of fish were netted out and transferred to full strength seawater for 24h before the final T4 collection.

RNAseq Analysis

Gill samples were collected, RNA extracted and libraries prepared, sequenced and mapped as Iversen et al (2020). Raw counts were analysed using EdgeR (ver. 3.30.0), using R (ver. 4.0.2) and RStudio (ver. 1.1.456). An ANOVA-like test was used to identify differential expressed genes between T1-T3 samples. Clustering analysis was performed using Pearson correlation, and heapmaps rendered using the R package pheatmap. An exact test was performed to identify differential expressed genes between T3 and T4.

Single nuclei RNAseq Analysis

Gills for single nuclei analysis were collected on dry ice and stored at -80°C. Duplicate samples were processed for T1-T4. Nuclei were released by detergent mechanical lysis, then samples were homogenized (30s) and nuclei isolated by sucrose gradient [45]. Libraries were created

using Chromium Single Cell 3' GEM, Library & Gel Bead Kit v3 (10x technologies) by University

of Manchester genomic technology core facility (UK). Raw data was converted to counts per

cell using Cell Ranger (10x Technologies, ver. 3.1.0) and processed using NCBI annotations.

The R package Seurat (ver. 3.1.5) was used to perform an integrated analysis using all

snRNAseq data [24], further details in results and discussion. Raw data will be available following

peer-reviewed publication.

Acknowledgements

The authors thank all of the animal staff at Kårvik havbruksstasjonen for their expert care of

the research animals, and the University of Manchester Genomics Technology core facility

(UK) for performing chromium 10x library preparation for snRNAseq. ACW is supported by the

Tromsø forskningsstiftelse (TFS) grant awarded to DGH (TFS2016DH). The Sentinel North

Transdisciplinary Research Program Université Laval and UiT awarded to DGH supports this

work. SHW is supported a grant from the Tromsø forskningsstiftelse (TFS) starter grant

TFS2016SW. Experimental costs were covered by HFSP grant "Evolution of seasonal timers"

RGP0030 /2015 awarded to ASIL and DGH.

Author Contributions

Conceptualization, A.C.W., Y.M., E.H.J., A.S.I.L and D.G.H; Resources, A.C.W, Y.M and M.I.;

Investigation, A.C.W., Y.M., M.I. E.H.J and D.G.H; Formal Analysis, A.C.W, Y.M., L.M.I., T.N and

S.R.S; Visualization, A.C.W and S.H.W; Writing – Original Draft, A.C.W and S.H.W; Writing –

Review & Editing, All; Supervision, A.S.I.L and D.G.H; Project Administration, A.S.I.L and D.G.H;

Funding Acquisition, A.S.I.L and D.G.H.

Figure Legends

Figure 1. Single nuclei RNAseq analysis of Atlantic salmon gill tissue. A) Gill tissue processing.

B) UMAP plot of pooled cell data from 18844 cells representing eight samples from four

collection states. The plot indicates 20 separate cell clusters. C) Expression of marker genes in

20 cell clusters. From left to right: hierarchical relatedness of difference cell clusters; total cells

in each cluster; UMI number in each cell cluster; gene features in each cell cluster; violin plots

showing expression pattern of marker genes for each cluster. Abbreviations: ACs - accessory

cells, DCs - dendritic cells, ECs - epithelial cells, fib - fibrocytes, GCs - goblet cells, LECs -

lymphatic endothelial cells, MCs - myeloid cells, MRC - mitochondrion-rich cells, NDCs - non-

differentiated cells, PVCs, pavement cells, RBCs - red blood cells (erythrocytes), TCs - T cells,

VCs - vascular cells.

Figure 2. Comparative abundance of cell clusters at different sampling points. A) Experimental

design. Fish were kept in constant light (LL) from hatching then transferred to short

photoperiod (SP; 8L:16D) for 8 weeks before being returned to constant light (LL) for 8 weeks.

Finally the fish were transferred to sea water for 24h. Sample points are indicated T1-T4. B)

Subset of cell clusters from T2, T3 and T4 (green and red dots) overlaid on T1 cells (grey dots).

C) Increasing abundance of sea-water mitochondrion-rich cells (MRCs SW) and vascular cells

(VC 3) during smoltification C) Decreasing abundance of leukocytes and immune-associated

cells during smoltification.

Figure 3. Photoperiodic changes in gill gene expression and localized cell cluster expression.

A) Heat map representing 9746 genes differentially regulated (FDR <0.01) from T1-T3.

Regulatory patterns for 5 major cluster are shown as amplitude index and 95% confidence

limits. Major gene ontology terms for each cluster are shown. B) RNAseq data for "classical"

smoltification-related genes and violin plots showing their cluster specific expression.

Figure 4. Sea-water transfer-associated changes in gill gene expression and localized cell

cluster expression. A) Genes differentially regulated (FDR <0.01) by 24h seawater transfer.

Major gene ontology terms for each cluster are shown. B) RNAseq data for sea-water transfer-

related genes and violin plots showing their cluster specific expression.

Supplemental Figure 1. RNAseq data for winter-dependent genes and violin plots showing

their cluster specific expression.

Figure 1.

Figure 2.

Figure 3.

Figure 4.