A vapor inhalation method for delivering heroin alters nociception, body temperature and spontaneous activity in female and male rats.

Arnold Gutierrez, Kevin M. Creehan, and Michael A. Taffe

Department of Psychiatry, University of California, San Diego; La Jolla, CA, USA

Running Title: Vapor inhalation of heroin

Address Correspondence to: Dr. Michael A. Taffe, Department of Psychiatry, 9500 Gilman Drive; University of California, San Diego, La Jolla, CA 92093; USA; Email: mtaffe@ucsd.edu

Abstract

Background: The ongoing crisis related to non-medical use of opioids makes it of continued importance to understand the risk factors for opioid addiction, the behavioral and neurobiological consequences of opioid exposure and to seek potential avenues for therapy. Pre-clinical rodent models have been critical to advancing understanding for decades but have been mostly limited to drug delivery by injection or by oral dosing, whereas inhalation, a significant route for some human users, has not been as well-established.

Method: We adapted an e-cigarette based inhalation system, previously shown efficacious for delivery of other drugs to rats, to deliver heroin vapor. Effects *in vivo* were assessed in male and female Sprague-Dawley rats using a warm-water assay for anti-nociception and an implanted radiotelemetry system for evaluating changes in body temperature and spontaneous activity rate.

Results: Inhalation of vapor created by heroin 100 mg/mL in the propylene glycol (PG) vehicle significantly slowed tail-withdrawal from a 52°C water bath, bi-phasically altered activity, and increased temperature in male and female rats. Inhalation of heroin 50 mg/mL for 15 minutes produced significant effects, as the lower bound on efficacy, whereas inhalation of heroin 100 mg/mL for 30 minutes produced robust effects across all endpoints and groups.

Conclusions: This work shows that e-cigarette devices deliver psychoactive doses of heroin to rats, using concentrations of ~50-100 mg/mL and inhalation durations of 15-30 minutes. This technique may be useful to assess the effects of inhaled heroin and other opioid drugs.

1. Introduction

The worldwide crisis related to the non-medical use of opioids continues, as reflected in high rates of deaths that are associated with prescription opioids, recreational drugs such as heroin and increasingly illicit supplies of fentanyl and analogs (CDC/NCHS, 2016; Wilson et al., 2020). The number of individuals seeking treatment for opioid dependence is high (SAMHSA, 2019). Improvement in understanding the risk factors for opioid addiction, as well as the behavioral and neurobiological consequences of opioid exposure, may enhance the discovery of new or improved avenues for therapy. Pre-clinical models in, e.g., laboratory rats, have been critical to advancing understanding but have been mostly limited to the examination of opioids delivered by parenteral injection, the insertion of subcutaneous pellets or pumps, or by oral dosing. Many humans use opioids by the inhalation route and indeed, one of the original public health crises related to non-medical use of opioids in the modern era was associated with the inhalation of vapor created by heating opium (Kane, 1881; Kritikos, 1960). The inhalation route might induce differences in the speed of brain entry, first pass metabolism, sequestration and release from non-brain tissues, etc. compared with these laboratory approaches. There is even some indication that inhalation use of heroin may come before a switch to injection use (Clatts et al., 2011; Cui et al., 2015; Rahimi-Movaghar et al., 2015; Syvertsen et al., 2017; Tyree et al., 2020), perhaps due to perceptions of safety or cultural factors. This may be true with respect to, e.g., disease transmission associated with intravenous injection practices but may involve other risks. For example, inhalation heroin users who develop leukoencephalopathy may be at increased risk for mortality compared with those who inject heroin (Alambyan et al., 2018; Cheng et al., 2019).

Prior studies have shown that the inhalation of heroin produces effects in animal models. Lichtman and colleagues reported anti-nociceptive effects of volatilized heroin and other opioids in mice using a heated glass pipe approach and a tail-withdrawal assay (Lichtman et al., 1996) and Carrol and colleagues demonstrated that monkeys would self-administer volatilized heroin

(Mattox and Carroll, 1996; Mattox et al., 1997). There has, however, been no subsequent broad adoption of either of these techniques, possibly due to the difficulty of creating and maintaining what appear to be one-off apparatus/devices constructed by the respective lab groups. The recent advent of e-cigarette style devices (aka Electronic Nicotine Delivery Systems or ENDS) presents the possibility of delivering a wide range of drugs other than nicotine, including opioids, via vapor inhalation. These devices are widely available (Esterl, 2013; Farnham, 2013), offer significant translational appeal and can be easily adapted to use with laboratory rodents using commonly available sealed-top housing chambers, standard laboratory house vacuum, and simple plumbing. Human use of these devices for ingestion of cannabis extracts or cannabidiol (CBD) is becoming so popular (Jones et al., 2016; Kenne et al., 2017; Mammen et al., 2016; Morean et al., 2015; Morean et al., 2017; Poklis et al., 2019) that terming these devices e-cigarettes or ENDS is becoming a misnomer; Electronic Drug Delivery Systems (EDDS) is more accurate.

The primary goal of this study was to establish methods for delivering physiologically and behaviorally significant amounts of heroin to rats via vapor inhalation. To this end we adapted an e-cigarette based inhalation system we have previously shown is effective for the delivery of nicotine (Javadi-Paydar et al., 2019b), THC (Javadi-Paydar et al., 2018a; Nguyen et al., 2016b), cannabidiol (Javadi-Paydar et al., 2019a), methamphetamine (Nguyen et al., 2016a; Nguyen et al., 2017), 3,4-methylenedioxypyrovalerone (MDPV) (Nguyen et al., 2016a) and oxycodone (Nguyen et al., 2019) to rats. Other work has demonstrated anti-nociceptive effects of inhaled oxycodone in male rats (Nguyen et al., 2019) and a preliminary work demonstrates antinociceptive efficacy of inhaled heroin and methadone (Gutierrez et al., 2020) in male and female rats, and reinforcing effects of inhaled heroin in female rats. Nearly identical systems have been shown to deliver nicotine (Montanari et al., 2020), THC from cannabis extracts (Freels et al., 2020) and the potent synthetic opioids sufentanil and fentanyl (Moussawi et al., 2020; Vendruscolo et al., 2018), to laboratory rodents.

To validate the delivery of heroin by vapor inhalation, we used a warm water tail-withdrawal assay for assessing effects on nociception, and a radiotelemetry system capable of reporting body temperature and activity responses. The telemetry measures were selected based on prior evidence that parenteral injection of opioids causes hyperthermia and increased locomotor activity in both rats and mice. For example, intravenous morphine (6 mg/kg) increased the rectal temperature of anesthetized rats (El Bitar et al., 2016), as did intrathecal morphine (1-15 ug) in freely moving male Wistar rats measured with telemetry (Safrany-Fark et al., 2015). Both morphine and oxycodone produce hyperthermia when injected subcutaneously in rats (Bhalla et al., 2011). Solis and colleagues (2017) showed that intravenous fentanyl (40 ug/kg) initially decreased body temperature of rats in the first hour after administration, (heroin (400 ug/kg) did not), but each drug induced *hyper*thermia in the interval approximately 60-120 minutes after injection (Solis et al., 2017). Heroin has also been shown to increase the locomotor activity of rats when injected subcutaneously or intraperitoneally in doses of ~0.25-1.0 mg/kg, although in some early studies a contrast of this described effect with vehicle injection was not made clear (Hoffman and Wise, 1993; Lamarque et al., 2001; Swerdlow and Koob, 1985; Swerdlow et al., 1985). Nevertheless, in other studies it was shown that heroin injection ~0.25-0.5 mg/kg, s.c., significantly increases the locomotor activity of rats (Raleigh et al., 2013; Singh et al., 2005; Sorge and Stewart, 2006) in comparison with a vehicle injection.

A secondary goal of this study was to determine the parameters necessary for dose control / manipulation within an effective range. As we've shown in prior studies (Javadi-Paydar et al., 2019b; Javadi-Paydar et al., 2018a; Nguyen et al., 2016a; Nguyen et al., 2016b), this can be effected with changes in drug concentration in the vapor vehicle and the duration of inhalation exposure, thus these parameters were under investigation. The final goal was to determine if there are any substantive sex differences in the response to heroin inhalation, see Craft (Craft, 2008) for review. While male and female rats acquire intravenous oxycodone self-administration at similar rates with 1-h access sessions (Mavrikaki et al., 2017), our recent study

with 8-h access found that female rats self-administered more oxycodone in acquisition, but did not reach higher breakpoints in a PR dose-substitution (Nguyen et al., 2020). A prior study reported no sex difference in the anti-nociceptive effect of heroin or oxycodone injected subcutaneously in Sprague-Dawley rats (Peckham and Traynor, 2006), thus no sex difference was predicted for this study.

2. Methods

2.1. Subjects:

Female (N=7) and male (N=7) Sprague-Dawley rats (Harlan, Livermore, CA) rats were housed in humidity- and temperature-controlled (23±2 °C) vivaria on 12:12 hour light:dark cycles. Rats had *ad libitum* access to food and water in their home cages and all experiments were performed in the rats' scotophase. All procedures were conducted under protocols approved by the Institutional Care and Use Committee of the University of California, San Diego.

2.2. Drugs

Heroin (diamorphine HCl) was administered by vapor inhalation with doses described by the concentration in the propylene glycol (PG) vehicle (1, 50, 100 mg/mL) and duration of inhalation (15, 30 minutes) as in prior studies (Javadi-Paydar et al., 2018a; Nguyen et al., 2016b). The heroin was provided by the U.S. National Institute on Drug Abuse Drug Supply Program.

2.3. Inhalation Apparatus

Vapor was delivered into sealed vapor exposure chambers (152 mm W X 178 mm H X 330 mm L; La Jolla Alcohol Research, Inc, La Jolla, CA, USA) through the use of e-vape

controllers (Model SSV-3; 58 watts; La Jolla Alcohol Research, Inc, La Jolla, CA, USA) to trigger

Smok Baby Beast Brother TFV8 sub-ohm tanks. Tanks were equipped with V8 X-Baby M2 0.25

ohm coils. Timed activation of e-vape controllers to deliver the scheduled drug vapor was done

using MedPC IV software (Med Associates, St. Albans, VT USA). The chamber air was

vacuum-controlled by a chamber exhaust valve (i.e., a "pull" system) to flow room ambient air

through an intake valve at ~1 L per minute. This also functioned to ensure that vapor entered

the chamber on each device triggering event. The vapor stream was integrated with the ambient

air stream once triggered.

2.4. Radiotelemetry

The rats were implanted with sterile radiotelemetry transmitters (TA11TA-F20; Data

Sciences International, St Paul, MN) in the abdominal cavity as previously described (Taffe et

al., 2015; Wright et al., 2012) on Post-Natal Day 31. For studies, animals were evaluated in

clean standard plastic homecages (thin layer of bedding) in a dark testing room, separate from

the vivarium, during the (vivarium) dark cycle. Radiotelemetry transmissions were collected via

telemetry receiver plates (Data Sciences International, St Paul, MN; RPC-1 or RMC-1) placed

under the cages as described in prior investigations (Javadi-Paydar et al., 2018b; Nguyen et al.,

2017). Test sessions for inhalation studies started with a 15-minute interval to ensure data

collection, then a 15-minute interval for baseline temperature and activity values. The rats were

then moved to a separate room for the vapor exposure sessions and then returned to the

recording chambers for up to 300 minutes after the start of vapor inhalation. These experiments

were conducted over the adult age interval of PND 204-235. The subjects had participated in

similar telemetry recording experiments following vapor and injected exposure to doses of

nicotine and THC from PND 50 to PND 190 with active dosing conducted no more frequently

than two times per week, in studies not previously reported.

2.5 Experiments:

2.5.1 Experiment 1:

The rats were recorded for a baseline interval and were then exposed to vapor (Heroin 0, 1, 50 mg/mL in the PG) inhalation for the designated interval (15 or 30 minutes) in exposure subgroups of 2-3 rats of the same sex per chamber. The initial dose range was determined by pilot studies and adaptation from self-administration studies published at present only in pre-print form (Gutierrez et al., 2020). The vapor conditions were evaluated in a counterbalanced order by subgroup, with inhalation duration alternating each session by sex. Sessions were conducted up to twice per week, i.e., no more frequently than every 3-4 days. Following this, the effects of Heroin 100 mg/mL for 15 vs 30 minutes, alternating by sex, were assessed on two different days, again at a 3-4 day interval.

2.5.2 Experiment 2:

A tail-withdrawal assay for nociception was conducted immediately after the vapor sessions, i.e. during Experiment 1, before the rats were returned to their individual recording chambers for telemetry assessment. The latency to withdraw the tail from insertion (3-4 cm) into a 52°C water bath was recorded by stopwatch, using methods as previously described (Gutierrez et al., 2020; Javadi-Paydar et al., 2018a; Nguyen et al., 2019). A 15 sec cutoff was used as the maximal latency for this assay to avoid any potential for tissue damage.

2.6 Data Analysis

The telemeterized body temperature and activity rate (counts per minute) were collected on a 5-minute schedule in telemetry studies but are expressed as 30 -minute averages for primary analysis. The time courses for data collection are expressed relative to the initiation of vapor inhalation and times in the figures refer to the end of the interval (e.g. "90 minutes" reflects the data collected from 65 to 90 minutes, inclusively. Due to transfer time after vapor

sessions, the "60 minutes" interval reflects the average of collections from ~40-60 minutes). Any missing temperature values were interpolated from the values before and after the lost time point. Activity rate values were not interpolated because 5-minute to 5-minute values can change dramatically, thus there is no justification for interpolating. Data for the first time-point after inhalation was inadvertently missing for two female rats in the 30-minute PG condition. The values for the second timepoint were used in place of the missing values. These ended up being very similar to the respective values recorded for this timepoint for the 15-minute PG condition. The telemetry data (temperature, activity rate) were generally analyzed with two-way Analysis of Variance (ANOVA) including repeated measures factors for the Vapor inhalation condition and the Time after vapor initiation. A mixed-effects model was used instead of ANOVA wherever there were missing datapoints. The nociception data were analyzed by three-way ANOVA including repeated-measures factors for Duration of inhalation and Vapor inhalation condition and a between-groups factor of Sex. Any significant main effects were followed with post-hoc analysis using Tukey (multi-level factors) or Sidak (two-level factors) procedures. All analysis used Prism 8 for Windows (v. 8.4.3; GraphPad Software, Inc, San Diego CA).

3. Results:

3.1 Experiment 1 (Telemetry):

3.1.1 15-minute inhalation:

Female Rats: The body temperature of the female rats (**Figure 1A**) was significantly elevated by the inhalation of Heroin 50 or 100 mg/mL as reflected in a significant effect of Time after vapor initiation [$F_{(9, 54)} = 13.97$; $P<0.0001$] and of the interaction of Time with Vape Condition [$F_{(27, 162)} = 5.91$; $P<0.0001$] on temperature. The post-hoc test confirmed that temperature was significantly higher after inhalation of Heroin 50 mg/mL compared with the PG condition (60-150 minutes after the start of inhalation) and Heroin 1 mg/mL (60-120 minutes) and likewise higher after inhalation of Heroin 100 mg/mL from 90-150 minutes compared with both the PG

and Heroin 1 mg/mL inhalation conditions. Relatedly, temperature was lower than the baseline after inhalation of PG (120-210 minutes after the start of inhalation) or Heroin 1 mg/mL (90-180; 300 minutes) inhalation conditions, but was not significantly changed from baseline in the Heroin 50 mg/mL condition until 180 and 270 minutes after the start of inhalation and from 180-300 minutes after the start of Heroin 100 mg/mL inhalation.

Figure 1: Mean (N=7 per group; ±SEM) body temperature (A, C) and activity rate (B, D) for female (A, B) and male (C,D) Sprague-Dawley rats after inhalation of vapor from the propylene glycol vehicle (PG), or Heroin (1, 50, 100 mg/mL in the PG) for 15 minutes. Shaded symbols indicate a significant difference from the baseline (Base) value, open symbols indicate a significant difference from the baseline and the PG condition at the respective time-point. A significant difference from the PG condition at a given time is indicated with *, a difference from PG and Heroin 1 mg/mL with &, and a difference from all other treatment conditions with #.

There was also a significant effect of Time [$F_{(9, 54)}$ = 7.42; P<0.0001] and of the interaction of Time with Vape Condition [$F_{(27, 162)}$ = 2.89; P<0.0001] on the activity of the female rats (**Figure 1B**). The post-hoc test confirmed that female rat activity was lower 60 minutes after the start of inhalation of Heroin 100 mg/mL compared with all other conditions and

thereafter higher compared with PG (90-150 minutes) or Heroin 1 mg/mL (90-120 minutes) inhalation.

Male rats: The body temperature of male rats (**Figure 1C**) was significantly altered by Time after vapor initiation [$F_{(9, 54)} = 26.60$; $P<0.0001$], by Vapor Condition [$F_{(3, 18)} = 7.07$; $P<0.005$] and of the interaction of Time with Vapor Condition [$F_{(27, 162)} = 6.41$; $P<0.0001$]. The post-hoc test confirmed that temperature was higher after inhalation of Heroin 100 mg/mL compared with all three other inhalation conditions from 60-180 minutes after the start of inhalation and lower compared with PG inhalation at the 300-minute time-point. Body

Figure 2: *Mean (N=7 per group; ±SEM) body temperature (A, C) and activity rate (B, D) for female (A, B) and male (C,D) Sprague-Dawley rats after inhalation of vapor from the propylene glycol vehicle (PG), or Heroin (1, 50, 100 mg/mL in the PG) for 30 minutes. Shaded symbols indicate a significant difference from the baseline (Base) value, open symbols indicate a significant difference from the baseline and the PG condition at the respective time-point. A significant difference from the PG condition at a given time is indicated with *, a difference from Heroin 1 mg/mL with §, a difference from PG and Heroin 1 mg/mL with &, and a difference from all other treatment conditions with #.*

temperature differed significantly from the baseline after inhalation of Heroin 100 mg/mL (60-120 minutes) but not in the other inhalation conditions. There was a significant effect of Time [F (9, 54) = 13.86; P<0.0001] and of the interaction of Time with Vape Condition [F (27, 162) = 2.37; P<0.001] on the activity of the male rats (**Figure 1D**). The post-hoc test confirmed that activity was greater after inhalation of Heroin 100 mg/mL compared with inhalation of PG (90-180 minutes after the start of inhalation) and compared with the other two heroin conditions (120-150 minutes).

3.1.2 30-minute inhalation:

Female Rats: The body temperature of the female rats (**Figure 2A**) was affected by Heroin inhalation for 30 minutes, as confirmed by a significant effect of Time after vapor initiation [F (9, 54) = 6.98; P<0.0001] and of the interaction of Time with Vape Condition [F (27, 162) = 3.19; P<0.0001] on temperature in the ANOVA. The post-hoc test confirmed that temperature was lower 60-90 minutes after the start of inhalation of Heroin 100 mg/mL compared with the PG (and 60 and 90 minutes after the start of the Heroin 50 mg/mL and Heroin 1 mg/mL conditions, respectively). Temperature was also lower than the baseline 60 minutes after initiation of Heroin 1 or 100 mg/mL. Temperature was then *higher* than in the PG condition after inhalation of Heroin 50 or 100 mg/mL, 120-150 minutes after the start of inhalation. The biphasic nature of the temperature response in the female rats is better illustrated by the 5-minute sample resolution for individuals presented in **Figure 3**.

The activity of female rats (**Figure 2B**) was also altered by Heroin inhalation and the ANOVA confirmed significant effects of Time after vapor initiation [F (9, 54) = 3.24; P<0.005] and of the interaction of Time with Vape Condition [F (27, 162) = 6.09; P<0.0001] on activity rate. Activity was suppressed after Heroin 50 mg/mL (60 minutes after the initiation of vapor) or 100 mg/mL (60-90 minutes) vapor exposure compared with the PG and Heroin 1 mg/mL conditions. Activity was also significantly lower than the baseline 60-90 minutes after the start of

Figure 3: *Individual body temperature responses in the female group after inhalation of Heroin 50 and 100 mg/mL for A) 15 minutes and B) 30 minutes illustrates the consistently biphasic nature of the temperature response.*

100 mg/mL inhalation, and lower compared with the 50 mg/mL condition in the 90-minute time-point. Activity rates were thereafter significantly higher compared with all other conditions after the inhalation of Heroin 50 mg/mL (120 minutes), and higher compared with the PG or Heroin 1 mg/mL after Heroin 50 mg/mL (150 minutes) or 100 mg/mL (150-180 minutes) conditions. Activity differed significantly between Heroin 50 mg/mL and Heroin 1 mg/mL inhalation 180 minutes after the start of inhalation as well.

Male Rats: The body temperature of the male rats (**Figure 2C**) was significantly affected by Heroin inhalation for 30 minutes, and the ANOVA confirmed significant effects of Time after vapor initiation [F (9, 54) = 43.43; P<0.0001], of Vapor Condition [F (3, 18) = 5.16; P<0.01], and of the interaction of Time with Vapor Condition [F (27, 162) = 11.06; P<0.0001] on temperature. The post-hoc test confirmed that body temperature of the male rats was elevated, compared with the respective time after PG or Heroin 1 mg/mL inhalation, after the Heroin 50 mg/mL (60-150 minutes after the start of inhalation) or Heroin 100 mg/mL (60-180 minutes) inhalation (**Figure 2B**). Similarly, temperature was *higher* than the baseline in the Heroin 50 mg/mL (60-120 minutes after the start of

inhalation) and Heroin 100 mg/mL (60-150 minutes) whereas it was *lower* than the baseline for each of the PG (90-180 minutes) and Heroin 1 mg/mL (120-300 minutes) inhalation conditions.

The activity of male rats (**Figure 2D**) was much lower than that of the female rats, but was also altered by Heroin inhalation and the ANOVA confirmed significant effects of Time after vapor initiation [$F_{(9, 54)}$ = 5.69; P<0.0001] and of the interaction of Time with Vape Condition [$F_{(27, 162)}$ = 4.06; P<0.0001] on activity rate. As with the female rats, the activity of the males was suppressed after Heroin 50 or 100 mg/mL (60 minutes after the initiation of vapor) vapor inhalation compared with the PG and Heroin 1 mg/mL conditions. Activity rates were thereafter significantly higher compared with PG inhalation in the Heroin 50 mg/mL (120-150 minutes) condition, and higher compared with the Heroin 1 mg/mL condition following Heroin 50 or 100 mg/mL (120-150 minutes) inhalation.

3.2 Experiment 2 (Anti-nociception):

When assessed after the inhalation sessions, tail withdrawal latency in the male and female rats was significantly increased by heroin vapor inhalation compared with both the baseline assessment and after the PG inhalation condition (**Figure 4**). The effect was robust since the Heroin 50 and 100 mg/mL for 30-minute conditions resulted in maximal effects in different subsets of 5/7 male rats and 5/7 female rats each. The 15-minute, Heroin 50 mg/mL condition resulted in a maximum effect in 1/7, and the 100 mg/mL for 15 minutes in 2/7, female rats.

The three way ANOVA (excluding baseline) confirmed a significant effect of Vapor condition [$F_{(1.857, 22.28)}$ = 142.4; P<0.0001], of Inhalation Duration [$F_{(1.000, 12.00)}$ = 43.88; P<0.0001], of the interaction of Vapor Condition with Duration [$F_{(2.157, 25.88)}$ = 12.66; P<0.0005] and of the interaction of Sex, Vapor Condition and Duration [$F_{(3, 36)}$ = 3.33; P=0.0301], on tail withdrawal latency. The post-hoc test confirmed significant elevations relative

to the respective PG and Heroin 1 mg/mL condition for the 50 and 100 mg/mL for 30-minute

conditions in female rats and for the 50 mg/mL for 30 minutes conditions in male rats. There

was also a significant difference in withdrawal latency in the male group following inhalation of

Heroin 50 mg/mL for 30 minutes compared with the same concentration for 15 minutes. Tail

withdrawal was also significantly slower in male rats after the inhalation of 100 mg/mL for 15

minutes compared with all other 15-minute conditions and slower after the inhalation of Heroin

100 mg/mL for 30 minutes compared with the PG for 30 minutes.

Figure 4: Mean (±SEM) tail withdrawal latency for male and female (N=7 per group) rats under baseline
and post-vapor conditions. The baseline was conducted before any of the counter-balanced inhalation
conditions and was not included in the statistical analysis. Within inhalation duration, a significant
difference from the PG condition is indicated with *, a difference from PG and Heroin 1 mg/mL with &, and
a difference from all other inhalation conditions with #. A difference between inhalation intervals within
drug concentration is indicated with §. Base, baseline. PG, propylene glycol. Her 1, Heroin (1 mg/mL);
Her 50, Heroin (50 mg/mL); Her 100, Heroin (100 mg/mL).

Follow-up analysis that was limited to the 15-minute conditions to further parse the

three-way interaction failed to confirm any sex differences. Collapsed across sex, however, the

Heroin 50 mg/mL condition significantly elevated tail-withdrawal latency compared with either

the PG or 1 mg/mL condition. There was also a significant difference in withdrawal latency

between the Heroin 50 mg/mL and each of the PG and Heroin 1 mg/mL conditions for the female rats.

4. Discussion:

4.1 Main Findings

This study confirms that this method of inhalation of heroin, based on vapor created by an Electronic Drug Delivery System (EDDS) device, produces significant effects *in vivo* in both male and female rats. The effects are dose-dependent, as they vary with both inhalation duration (at a fixed concentration) and with drug concentration in the e-cigarette vehicle (at a given inhalation duration). We have previously shown that heroin vapor exposure (100 mg/mL, 30 minutes) produces moderate anti-nociceptive effects in male and female Wistar rats (Gutierrez et al., 2020) and that is extended to Sprague-Dawley rats, and to a wider range of inhalation conditions, in the present study. As we've previously shown, differences between common laboratory rat strains can result in quantitative or even qualitative differences in response to vapor inhalation of drugs, effects which may vary depending on the outcome measure (Taffe et al., 2020). The present results obtained near-maximal effects in more animals in the 100 mg/mL condition, compared with that prior study, however there were equipment differences between the two studies which may have resulted in more effective drug delivery. For example that prior study reported no anti-nociceptive effect of oxycodone vapor (100 mg/mL, 30 minutes) inhalation using first-generation Protank 3 canisters, whereas a study using the second-generation Herakles Sub Ohm canisters, more similar to the present second-generation SMOK canisters, did report anti-nociception subsequent to 100 mg/mL oxycodone vapor inhalation (Nguyen et al., 2019). In this study, the threshold for observable effects was the 50 mg/mL inhalation for 15-minute condition. These exposure parameters led to significant anti-nociception (both collapsed across sex and within the female group) and produced significant effects on body temperature in the female rats. This study therefore establishes 50-100 mg/mL

as an effective concentration range, and 1 mg/mL as an ineffective concentration, for non-contingent inhalation exposure to heroin in rats.

This is a nontrivial advance from the prior demonstrations that rats and mice, respectively, will self-administer the potent opioids sufentanil (Vendruscolo et al., 2018) and fentanyl (Moussawi et al., 2020). First, the doses needed to produce robust changes in nociception, thermoregulation and locomotor behavior are often far in excess of the doses that rodents will self-administer. Second, we found previously that cocaine and some amphetamine or cathinone class psychomotor stimulants which exhibit both low potency and lower solubility in PG may not readily be delivered in active doses with this approach (Nguyen et al., 2016a). Thus, it was critical to show that a less potent opioid such as heroin can be delivered with this method.

There were biphasic dose-dependent effects of heroin on body temperature which was expressed as lower exposure conditions (doses) producing reliable increases in temperature, and higher doses / exposures initially reducing body temperature, at least in the female rats. A prior study found similar biphasic effects after intravenous fentanyl, which initially decreased body temperature of male Long-Evans rats in the first hour after administration, but induced *hyper*thermia thereafter (Solis et al., 2017). This may reflect potency, or brain penetration speed, as potential differences between the two opioids that permitted this initial hypothermia to be observed. It could similarly be the case in the present study that the inhaled route of administration speeds the brain entry of heroin versus the more common s.c. or i.p. or even i.v. routes of administration. If so, this may be a key area in which the development of this inhalation model allows the investigation of effects of heroin which are unique to this route of administration. It may also be the case that effects of inhalation are more similar to the effects of the intravenous route, in this case the method offers several practical advantages over implanting and maintaining intravenous catheters in rodents. Such advantages include overall

cost, surgical expertise and avoiding subject loss due to occluded catheters or health complications related to the catheter.

The apparent sex difference (males did not express the initial hypothermia under any of the conditions tested) is most likely a difference in effective dose, since pilot work with a group of male Sprague-Dawley rats illustrated a consistent *hypo*thermia associated with a high exposure condition and hyperthermia with a lower exposure condition (not shown). Also, female rats in this study were affected in terms of body temperature and anti-nociception after 15-minute inhalation of Heroin 50 mg/mL, whereas male rats were only affected on the anti-nociception assay and only to an extent that failed to reach statistical significance. Overall, these patterns are more consistent with a dosing difference across sex rather than a sex difference per se. A previous study found no difference in the ED50 for heroin (s.c.) in a warm water tail-withdrawal test between male and female Sprague-Dawley rats (Peckham and Traynor, 2006) and where sex differences were found (morphine, hydrocodone, hydromorphone) the males were *more* sensitive. Nevertheless, female mice develop greater hypothermia than do males after a mg/kg equivalent injection of morphine (Kest et al., 2000).

The initial suppression of activity, followed by a rebound of increased activity, is similar to that reported for injected morphine (Craft et al., 2006) but differs somewhat from prior results for injected heroin, which appeared to show a monotonic effect of time where activity is highest at the first time-points observed after dosing and then declined steadily with time. One possible difference is that some of those studies were conducted with rats in the light part of the cycle (Hoffman and Wise, 1993; Swerdlow and Koob, 1985; Swerdlow et al., 1985), however others reporting similar monotonic time-courses were on a reverse cycle and were tested in the dark (Singh et al., 2005; Sorge and Stewart, 2006). A more consistent difference is that many studies of the effects of injected heroin used a non-housing, specialized photo cell cage (Hoffman and Wise, 1993; Singh et al., 2005; Sorge and Stewart, 2006; Swerdlow and Koob, 1985; Swerdlow et al., 1985) with mesh or rod floors, unlike the plastic floored normal housing cages with a thin

layer of bedding used here. It may be the case that the more familiar, housing-type environment facilitated expression of a more naturalistic response.

4.2 Alternate Methods:

This study reports a set of methods we have evolved from our ongoing experience using EDDS-based vapor exposure. Through the developmental arc of this approach, it is clear that this is but one specific set of methods and that many variations are likely to also work in the hands of other interested laboratories. First, we have used several different iterations of e-cigarette canister/atomizer and found that most differences are subtle in terms of drug delivery. There is a constantly changing and highly varied array of products available online and in local stores and most of them are likely to function for laboratory purposes. Effective vapor delivery can be achieved with many different types of cannister / atomizer product, with only modest methodological adjustments of puffing schedule, drug concentration and inhalation time. Second, the volume of the sealed chamber used to expose animals and the number per chamber can be varied, depending on the goals to be attained and the available equipment. The initial validation step is simply to determine that a vapor cloud to fill the available chamber volume can be produced. Third, while we have adopted a puffing schedule of every 5 minutes, more-, or less-frequent puffing could be used. The most important consideration for developing these methods is to have the ability to contrast and validate drug exposure parameters. Validation should be done with a desired target endpoint (whether that be anti-nociceptive effects, effects on spontaneous behavior, on physiological endpoints such as thermoregulation, or other measure of interest) that is reliably produced in the relevant laboratory species/strain/genotype by injection or other route of administration. Once the target endpoint is determined, systematic manipulation of equipment, dosing protocols or drug concentrations can be used to achieve dose control.

5. Conclusions

In conclusion, the e-cigarette based inhalation approach delivers physiologically and behaviorally relevant amounts of heroin to rats. Given the growing opioid crisis, and the widespread use of e-cigarette technologies, it will be of increasing utility to have a rodent pre-clinical model for the assessment of the effects of inhaled heroin and other opioids.

Acknowledgements

Literature Cited

Alambyan, V., Pace, J., Miller, B., Cohen, M.L., Gokhale, S., Singh, G., Shun, M.C., Hammond, A., Ramos-Estebanez, C., 2018. The Emerging Role of Inhaled Heroin in the Opioid Epidemic: A Review. JAMA Neurol 75(11), 1423-1434.

Bhalla, S., Andurkar, S.V., Gulati, A., 2011. Study of adrenergic, imidazoline, and endothelin receptors in clonidine-, morphine-, and oxycodone-induced changes in rat body temperature. Pharmacology 87(3-4), 169-179.

CDC/NCHS, 2016. National Vital Statistics System, Mortality. U.S. Department of Health and Human Services, CDC, Atlanta, GA.

Cheng, M.Y., Chin, S.C., Chang, Y.C., Wu, T., Lim, S.N., Hsieh, H.Y., Hsu, J.L., Chang, C.W., Tseng, W.J., Li, H.T., Chiang, H.I., Chang, B.L., Tsai, M.H., Ro, L.S., 2019. Different routes of heroin intake cause various heroin-induced leukoencephalopathies. J Neurol 266(2), 316-329.

Clatts, M.C., Goldsamt, L.A., Giang le, M., Colon-Lopez, V., 2011. Accelerated transition to injection among male heroin initiates in Hanoi, Vietnam: implications for early harm reduction interventions. J Community Health 36(6), 999-1003.

Craft, R.M., 2008. Sex differences in analgesic, reinforcing, discriminative, and motoric effects of opioids. Exp Clin Psychopharmacol 16(5), 376-385.

Craft, R.M., Clark, J.L., Hart, S.P., Pinckney, M.K., 2006. Sex differences in locomotor effects of morphine in the rat. Pharmacol Biochem Behav 85(4), 850-858.

Cui, S., Li, Y., Xiang, Y.T., Yan, K.J., Fan, C., Chen, Q., Du, W., Zhang, G., Xiao, L., Li, Y., Yuan, X., Xie, K., Luo, X., 2015. Demographic and Clinical Features of Chinese Heroin Users Who Switch from Non-injection to Injection. Subst Use Misuse 50(14), 1739-1746.

El Bitar, N., Pollin, B., Karroum, E., Pincede, I., Le Bars, D., 2016. Entanglement between thermoregulation and nociception in the rat: the case of morphine. J Neurophysiol 116(6), 2473-2496.

Esterl, M., 2013. E-Cigarettes Fire Up Investors, Regulators, The Wall Street Journal. Forbes, New York.

Farnham, A., 2013. E-Cigarette Sales to Hit $1 Billion, ABC News. ABC News Internet Ventures, New York, NY.

Freels, T.G., Baxter-Potter, L.N., Lugo, J.M., Glodosky, N.C., Wright, H.R., Baglot, S.L., Petrie, G.N., Yu, Z., Clowers, B.H., Cuttler, C., Fuchs, R.A., Hill, M.N., McLaughlin, R.J., 2020. Vaporized Cannabis Extracts Have Reinforcing Properties and Support Conditioned Drug-Seeking Behavior in Rats. J Neurosci 40(9), 1897-1908.

Gutierrez, A., Nguyen, J.D., Creehan, K.M., Taffe, M.A., 2020. Self-administration of heroin by vapor inhalation in female Wistar rats. bioRxiv, 2020.2003.2030.016725.

Hoffman, D.C., Wise, R.A., 1993. Lack of cross-sensitization between the locomotor-activating effects of bromocriptine and those of cocaine or heroin. Psychopharmacology (Berl) 110(4), 402-408.

Javadi-Paydar, M., Creehan, K.M., Kerr, T.M., Taffe, M.A., 2019a. Vapor inhalation of cannabidiol (CBD) in rats. Pharmacol Biochem Behav 184, 172741.

Javadi-Paydar, M., Kerr, T.M., Harvey, E.L., Cole, M., Taffe, M.A., 2019b. Effects of nicotine and THC vapor inhalation administered by an electronic nicotine delivery system (ENDS) in male rats. Drug Alcohol Depend 198, 54-62.

Javadi-Paydar, M., Nguyen, J.D., Kerr, T.M., Grant, Y., Vandewater, S.A., Cole, M., Taffe, M.A., 2018a. Effects of Delta9-THC and cannabidiol vapor inhalation in male and female rats. Psychopharmacology (Berl) 235(9), 2541-2557.

Javadi-Paydar, M., Nguyen, J.D., Vandewater, S.A., Dickerson, T.J., Taffe, M.A., 2018b. Locomotor and reinforcing effects of pentedrone, pentylone and methylone in rats. Neuropharmacology 134(Pt A), 57-64.

Jones, C.B., Hill, M.L., Pardini, D.A., Meier, M.H., 2016. Prevalence and Correlates of Vaping Cannabis in a Sample of Young Adults. Psychol Addict Behav.

Kane, H.H., 1881. American Opium Smokers. Harper's Weekly, 646.

Kenne, D.R., Fischbein, R.L., Tan, A.S., Banks, M., 2017. The Use of Substances Other Than Nicotine in Electronic Cigarettes Among College Students. Subst Abuse 11, 1178221817733736.

Kest, B., Adler, M., Hopkins, E., 2000. Sex differences in thermoregulation after acute and chronic morphine administration in mice. Neurosci Lett 291(2), 126-128.

Kritikos, P.G., 1960. Der Mohn, das Opium u. ihr Gebrauch im Spaetminoicum III. Bemerkungen zu dem gefundenen Idol der Minoischen Gottheit des Mohns. Archives of the Academy of Athens, 21.

Lamarque, S., Taghzouti, K., Simon, H., 2001. Chronic treatment with Delta(9)-tetrahydrocannabinol enhances the locomotor response to amphetamine and heroin. Implications for vulnerability to drug addiction. Neuropharmacology 41(1), 118-129.

Lichtman, A.H., Meng, Y., Martin, B.R., 1996. Inhalation exposure to volatilized opioids produces antinociception in mice. J Pharmacol Exp Ther 279(1), 69-76.

Mammen, G., Rehm, J., Rueda, S., 2016. Vaporizing cannabis through e-cigarettes: Prevalence and socio-demographic correlates among Ontario high school students. Can J Public Health 107(3), e337-e338.

Mattox, A.J., Carroll, M.E., 1996. Smoked heroin self-administration in rhesus monkeys. Psychopharmacology (Berl) 125(3), 195-201.

Mattox, A.J., Thompson, S.S., Carroll, M.E., 1997. Smoked heroin and cocaine base (speedball) combinations in rhesus monkeys. Exp Clin Psychopharmacol 5(2), 113-118.

Mavrikaki, M., Pravetoni, M., Page, S., Potter, D., Chartoff, E., 2017. Oxycodone self-administration in male and female rats. Psychopharmacology (Berl) 234(6), 977-987.

Montanari, C., Kelley, L.K., Kerr, T.M., Cole, M., Gilpin, N.W., 2020. Nicotine e-cigarette vapor inhalation effects on nicotine & cotinine plasma levels and somatic withdrawal signs in adult male Wistar rats. Psychopharmacology (Berl) 237(3), 613-625.

Morean, M.E., Kong, G., Camenga, D.R., Cavallo, D.A., Krishnan-Sarin, S., 2015. High School Students' Use of Electronic Cigarettes to Vaporize Cannabis. Pediatrics 136(4), 611-616.

Morean, M.E., Lipshie, N., Josephson, M., Foster, D., 2017. Predictors of Adult E-Cigarette Users Vaporizing Cannabis Using E-Cigarettes and Vape-Pens. Subst Use Misuse 52(8), 974-981.

Moussawi, K., Ortiz, M.M., Gantz, S.C., Tunstall, B.J., Marchette, R.C.N., Bonci, A., Koob, G.F., Vendruscolo, L.F., 2020. Fentanyl vapor self-administration model in mice to study opioid addiction. Science Advances 6(32), eabc0413.

Nguyen, J.D., Aarde, S.M., Cole, M., Vandewater, S.A., Grant, Y., Taffe, M.A., 2016a. Locomotor Stimulant and Rewarding Effects of Inhaling Methamphetamine, MDPV, and Mephedrone via Electronic Cigarette-Type Technology. Neuropsychopharmacology 41(11), 2759-2771.

Nguyen, J.D., Aarde, S.M., Vandewater, S.A., Grant, Y., Stouffer, D.G., Parsons, L.H., Cole, M., Taffe, M.A., 2016b. Inhaled delivery of Delta(9)-tetrahydrocannabinol (THC) to rats by e-cigarette vapor technology. Neuropharmacology 109, 112-120.

Nguyen, J.D., Bremer, P.T., Hwang, C.S., Vandewater, S.A., Collins, K.C., Creehan, K.M., Janda, K.D., Taffe, M.A., 2017. Effective active vaccination against methamphetamine in female rats. Drug Alcohol Depend 175, 179-186.

Nguyen, J.D., Creehan, K.M., Kerr, T.M., Taffe, M.A., 2020. Lasting effects of repeated (9) - tetrahydrocannabinol vapour inhalation during adolescence in male and female rats. Br J Pharmacol 177(1), 188-203.

Nguyen, J.D., Grant, Y., Creehan, K.M., Hwang, C.S., Vandewater, S.A., Janda, K.D., Cole, M., Taffe, M.A., 2019. Delta(9)-tetrahydrocannabinol attenuates oxycodone self-administration under extended access conditions. Neuropharmacology 151, 127-135.

Peckham, E.M., Traynor, J.R., 2006. Comparison of the antinociceptive response to morphine and morphine-like compounds in male and female Sprague-Dawley rats. J Pharmacol Exp Ther 316(3), 1195-1201.

Poklis, J.L., Mulder, H.A., Peace, M.R., 2019. The unexpected identification of the cannabimimetic, 5F-ADB, and dextromethorphan in commercially available cannabidiol e-liquids. Forensic Sci Int 294, e25-e27.

Rahimi-Movaghar, A., Amin-Esmaeili, M., Shadloo, B., Noroozi, A., Malekinejad, M., 2015. Transition to injecting drug use in Iran: A systematic review of qualitative and quantitative evidence. Int J Drug Policy 26(9), 808-819.

Raleigh, M.D., Pravetoni, M., Harris, A.C., Birnbaum, A.K., Pentel, P.R., 2013. Selective effects of a morphine conjugate vaccine on heroin and metabolite distribution and heroin-induced behaviors in rats. J Pharmacol Exp Ther 344(2), 397-406.

Safrany-Fark, A., Petrovszki, Z., Kekesi, G., Keresztes, C., Benedek, G., Horvath, G., 2015. Telemetry monitoring for non-invasive assessment of changes in core temperature after spinal drug administration in freely moving rats. J Pharmacol Toxicol Methods 72, 19-25.

SAMHSA, 2019. Treatment Episode Data Set (TEDS). 2017. Admissions to and Discharges From Publicly-FundedSubstance Abuse Treatment. Substance Abuse and Mental Health Services Administration, Office of Applied Studies, Rockville, MD.

Singh, M.E., McGregor, I.S., Mallet, P.E., 2005. Repeated exposure to Delta(9)-tetrahydrocannabinol alters heroin-induced locomotor sensitisation and Fos-immunoreactivity. Neuropharmacology 49(8), 1189-1200.

Solis, E., Jr., Cameron-Burr, K.T., Kiyatkin, E.A., 2017. Heroin Contaminated with Fentanyl Dramatically Enhances Brain Hypoxia and Induces Brain Hypothermia. eNeuro 4(5).

Sorge, R.E., Stewart, J., 2006. The effects of long-term chronic buprenorphine treatment on the locomotor and nucleus accumbens dopamine response to acute heroin and cocaine in rats. Pharmacol Biochem Behav 84(2), 300-305.

Swerdlow, N.R., Koob, G.F., 1985. Separate neural substrates of the locomotor-activating properties of amphetamine, heroin, caffeine and corticotropin releasing factor (CRF) in the rat. Pharmacol Biochem Behav 23(2), 303-307.

Swerdlow, N.R., Vaccarino, F.J., Koob, G.F., 1985. Effects of naloxone on heroin-, amphetamine- and caffeine-stimulated locomotor activity in the rat. Pharmacol Biochem Behav 23(3), 499-501.

Syvertsen, J.L., Paquette, C.E., Pollini, R.A., 2017. Down in the valley: Trajectories of injection initiation among young injectors in California's Central Valley. Int J Drug Policy 44, 41-49.

Taffe, M.A., Creehan, K.M., Vandewater, S.A., 2015. Cannabidiol fails to reverse hypothermia or locomotor suppression induced by Delta(9) -tetrahydrocannabinol in Sprague-Dawley rats. Br J Pharmacol 172(7), 1783-1791.

Taffe, M.A., Creehan, K.M., Vandewater, S.A., Kerr, T.M., Cole, M., 2020. Effects of Delta(9)-tetrahydrocannabinol (THC) vapor inhalation in Sprague-Dawley and Wistar rats. Exp Clin Psychopharmacol.

Tyree, G.A., Mosery, N., Closson, E.F., Mabude, Z., du Toit, C., Bangsberg, D.R., Safren, S.A., Mayer, K.H., Smit, J.A., Mimiaga, M.J., Grelotti, D.J., 2020. Trajectories of initiation for the heroin-based drug whoonga - qualitative evidence from South Africa. Int J Drug Policy 82, 102799.

Vendruscolo, J.C.M., Tunstall, B.J., Carmack, S.A., Schmeichel, B.E., Lowery-Gionta, E.G., Cole, M., George, O., Vandewater, S.A., Taffe, M.A., Koob, G.F., Vendruscolo, L.F., 2018. Compulsive-Like Sufentanil Vapor Self-Administration in Rats. Neuropsychopharmacology 43(4), 801-809.

Wilson, N., Kariisa, M., Seth, P., Smith, H.t., Davis, N.L., 2020. Drug and Opioid-Involved Overdose Deaths - United States, 2017-2018. MMWR Morb Mortal Wkly Rep 69(11), 290-297.

Wright, M.J., Jr., Angrish, D., Aarde, S.M., Barlow, D.J., Buczynski, M.W., Creehan, K.M., Vandewater, S.A., Parsons, L.H., Houseknecht, K.L., Dickerson, T.J., Taffe, M.A., 2012. Effect of ambient temperature on the thermoregulatory and locomotor stimulant effects of 4-methylmethcathinone in Wistar and Sprague-Dawley rats. PLoS One 7(8), e44652.