1

Large Mammals Have More Powerful Antibacterial Defenses Than Expected from Their Metabolic Rates

Cynthia J. Downs[1], Laura A. Schoenle[2,3], Eric W. Goolsby[4], Samantha J. Oakey[2], Ray Ball[5], Rays H.Y. Jiang[2], Kirk C. Klasing[6], Lynn B. Martin[2]

[1] Department of Environmental Biology, State University of New York College of Environmental Science and Forestry, Syracuse, NY 13210 USA

[2] Global and Planetary Health, University of South Florida, Tampa, FL 33620 USA

[3] Current address: Office of Undergraduate Biology, Cornell University, Ithaca, NY 14853 USA

[4] Department of Biology, University of Central Florida, Orlando, FL 32816 USA

[5] Department of Biology, Eckerd College, Saint Petersburg, FL 33711 USA

[6] Department of Animal Science, University of California, Davis, CA 95616 USA

Short title: Scaling of mammalian antibacterial defenses

Keywords: allometry, antibacterial capacity, constitutive immunity, immune defenses, safety factory hypothesis, scaling

Article type: Major Article

Elements of the manuscript: main text, supplemental materials document (including Supplementary Methods, Figs. S1 to S7, Tables S1 to S6, Captions for Movies S1), and Movie S1

Abstract

Terrestrial mammals span 7 orders of magnitude in body size, ranging from the < 2 g pygmy shrew (*Suncus etruscus*) to the > 3900 kg African elephant (*Loxodonta africana*). Although body size has profound effects on the behavior, physiology, ecology, and evolution of animals, how investment in immune defenses changes with body size is unknown. Here, we develop a novel 12-point dilution-curve approach to describe and compare antibacterial capacity against 3 bacterial species among >160 terrestrial species of mammals. We show that antibacterial activity in serum across mammals exhibits isometry, but the serum of large mammals is less hospitable to bacteria than would be predicted by their metabolic rates. Specifically, hypometric metabolic rates would predict that a large species should have disproportionately lower antibacterial capacity than small species, but body size is unrelated to killing capacity across species. Scaling of antibacterial immune defenses provides novel perspectives on the ecology of host-pathogen interactions, and on their co-evolutionary dynamics. These results have direct implications for effectively modeling the evolution of immune defenses and identifying potential reservoir hosts of zoonotic pathogens.

Introduction

Hosts can be thought of as islands whose size impacts how many parasites, including microbes, can colonize them (Kuris et al. 1980). Bigger hosts represent bigger islands, leading parasite abundance (Kieft and Simmons 2015), biomass (Poulin and George-Nascimento 2007), diversity (Ezenwa 2004), size (Harrison 1915), size diversity (Poulin 2007), and prevalence (Ishtiaq et al. 2017) to increase with host body size. Large-bodied species also appear more susceptible to some parasite infections (Filion et al. 2020). Large body size is also intimately connected to the relative value of self-defense versus reproduction (Calder 1984), as most hosts reach large size via a long lifespan (Harrison 2017; Downs et al. 2020). Altogether, natural selection should favor the evolution of robust immune systems in large hosts (Sheldon and Verhulst 1996; Downs et al. 2014), and despite a few theoretical (Langman and Cohn 1987; Wiegel and Perelson 2004; Banerjee et al. 2017) and empirical (Lee 2006; Schoenle et al. 2018) calls to investigate scaling of immune defenses, very little works has yet considered whether and how immune defenses scale across the full range of terrestrial mammalian body sizes.

Generally, body size scaling relationships are modeled as power laws, $Y = a\,M^{b}$, with linear transformations, $\log_{10} Y = \log_{10} a + b \log_{10} M$, where Y is the focal trait, M is body mass, a is the intercept, and b is the scaling factor. Four hypotheses make predictions about the values of the scaling factors describing the relationship between immune defenses and body mass. To date, the Protecton and the Complexity of Immunity Hypotheses have received the most attention. Both hypotheses predict that immunity should scale in direct proportion to body size (i.e., isometrically), which results in a mass-invariant relationship ($b = 0$) for concentrations (Langman and Cohn 1987; Wiegel and Perelson 2004). The fractal nature of the circulatory and lymphatic systems is argued to predict isometry because rates of parasite detection and delivery

of defenses should be driven by transit of cells and proteins through the vasculature (e.g., Wiegel

and Perelson 2004; Banerjee et al. 2017; Banerjee 2018). Although this framework was

developed for predicting the number of a single clone of B cells necessary to detect an antigen in

a host and circulation time of a B cell (Wiegel and Perelson 2004), it has been extended to make

predictions about other types of immune defenses but particularly adaptive immunity (Banerjee

et al. 2017; Banerjee 2018). By contrast, the Rate of Metabolism Hypothesis predicts a

hypometric coefficient for immunity ($b < 0$) because all cellular activities are linked to basal

metabolic rates (BMR) (Brown et al. 2004). As mass-specific BMR scales at $b = -0.25$, immune

defenses involving cell turnover and metabolism should scale similarly (Dingli and Pacheco

2006).

Finally, the Safety Factor Hypothesis invokes performance-safety relationships from

biophysics (Harrison 2017) and physiology (Diamond 2002), and posits that large species should

evolve disproportionately stronger (i.e., hypermetric, $b > 0$) damage-mitigation mechanisms than

small ones (Harrison 2017; Downs et al. 2020). Immunologically, the Safety Factor Hypothesis

predicts that large hosts should evolve exceptionally robust constitutive defenses to protect

against the greater infection risks they will experience. Variants of the Safety Factor Hypothesis

exist in the literature already, including Peto's paradox for cancer (Peto 1977) and optimal

defense theory (Shudo and Iwasa 2001). Only recently, however, has direct empirical evidence

been provided for immune defenses, namely hypermetric scaling for both neutrophil

concentrations in mammals ($b = 0.11$) (Downs et al. 2020) and heterophil concentrations in birds

($b = 0.19$) (Ruhs et al. 2020).

Despite evidence for some immune allometries, it remains obscure whether the functional

capacity of immune defenses scale similarly. Does hypermetric scaling of granulocyte

concentrations represent greater immune protection in large animals or simply evidence that

large animals must compensate for lower per-granulocyte effectiveness by circulating more

cells? Cell size scales isometrically with body mass, but per-gram metabolism scales

hypometrically (Schmidt-Nielsen 1984), so the average large-animal leukocyte might be

comparatively ineffective compared to a leukocyte from a small species. Moreover, as selection

tends to act on integrated traits (Bennett and Huey 1990), scaling inference based on processes

that affect host fitness (i.e., control of bacterial infections) will inherently be more insightful than

studies solely based on cell concentrations. Few scaling studies of immunity to date have relied

on integrated immune functions, and others either involved few species, narrow taxonomic

groups, or found body mass effects but did not estimate allometries (e.g., Nunn 2002; Blount et

al. 2003; Nunn et al. 2003; Lee et al. 2008; Schneeberger et al. 2013; Tian et al. 2015;

Fassbinder-Orth et al. 2019; reviewed in Downs et al. 2020).

Our goals here were i) to determine how blood-based antibacterial defenses scaled with

body size among terrestrial mammals, and ii) to test which hypothesis best-predicts the scaling

exponent (b) for this type of immunity. We focused on antibacterial capacity of sera (French and

Neuman-Lee 2012), as this defense is functional and integrative (Demas et al. 2011), is an

important determinant of host health and virulence for certain bacterial pathogens (Taylor 1983),

and is measurable in a standard, reagent-independent way. Briefly, antibacterial capacity

measures bacteriostasis and bacteria killing capacity of sera. To describe interspecific variation

in this immune function in the most confound-free manner possible, we compared the efficacy of

sera of several, independent replicates from adults of mammalian species housed in zoos. To

provide generality to any scaling we observed, we quantified the antibacterial capacity of sera of

160+ mammalian species against three distinct bacteria, *Escherichia coli* (ATCC #8739),

Salmonella enterica (ATCC #13311), and *Micrococcus luteus* (ATCC # 4698). These bacteria

are common pathogens of many vertebrates (Mastroeni et al. 2001), some portion of their natural

infection timeline occurs in blood (Mittrücker and Kaufmann 2000), and they are controlled

effectively by blood-borne immune defenses (Taylor 1983). As the last common evolutionary

ancestor of our focal microbes was ancient, any consistency in allometries should be more

attributable to host immunity than parasite evolutionary history.

To enable meaningful comparisons across species, we characterized antibacterial dilution

series, instead of antibacterial capacity at single dilutions (as is typically used, French and

Neuman-Lee 2012), as the shapes of resultant *ex vivo* antibacterial functions capture salient

aspects of defense (Fig. 1A). For instance, if a very dilute serum sample was effective against a

standard amount of bacteria, a given individual of that host species would be relatively adept at

controlling a large initial dose of bacteria in its bloodstream. Other meaningful antibacterial

parameters included i) the maximum amount of bacteria that serum could protect against, ii) the

serum dilution at the point of most rapid change from protected to vulnerable, and iii) the slope

at this steepest part of the antibacterial function (Fig. 1A). This last parameter could be a proxy

for either how rapidly serum defenses could be recruited to a bacterial replication event (Murphy

et al. 2007) or capture the pattern by which proteins and other molecules were recruited

combinatorically to control bacteria. The Safety Factor Hypothesis, our focal hypothesis, predicts

that large mammals will have disproportionately higher maximal antibacterial capacity, higher

antibacterial capacity at a specific dilution, and a more rapid shift from vulnerable to complete

protection than small species (Fig. 1B). By contrast, the Rate of Metabolism Hypothesis predicts

the opposite for large mammals relative to small ones (Fig 1C), and the Protecton and

Complexity Hypotheses predict that large mammals will have the same antibacterial capacity as

small species (Fig. 1D).

Materials and Methods

Samples

We used the simulations performed by Dingemanse and Dochtermann (Fig. 1 of 2013) to

guide our sample sizes. We obtained serum from healthy, zoo- and lab-housed animals ranging

in body size from 16 g – 3,600 kg, which covers the range of body sizes of extant terrestrial

mammals (Fig. S1) (Smith and Lyons 2011). We supplemented these samples, which comprised

the vast majority of our sample, with those from lab-housed animals to augment the small-bodied

species (< 2 kg) included in our analysis, as these body sizes are rare for mammals in most zoos.

Most zoo samples were collected as part of routine wellness checkups whereas lab samples were

collected at the termination of experiments; all animals from which samples were taken were

outwardly healthy. We could not run all assays against all bacterial species for all samples

because of sample volume constraints. Final sample sizes of mammal species were 199, 186, and

164 for antibacterial capacity against *E. coli, S. enterica,* and *M. luteus*, respectively. We assayed

a mean of ≥ 4 samples per species for each microbe (n = 1-41 replicates per species, Fig. S2). We

also obtained >15 samples from a subset of species representing the full range of body masses to

ensure that a single individual from a large or small species was not biasing a species mean (Fig.

S3). Samples were stored at the collection location at -80 to -20 °C until shipped on dry ice to

our research labs where they were stored at 80 °C until assays. Samples were used in assays

within 24h of thawing. We restricted our analysis to samples collected in the years 2005-2019

because preliminary work indicated that older samples had reduced antibacterial ability. Use of

these samples was approved by the Institutional Animal Care and Use Committees at home

institutions. Samples were collected as part of routine veterinary exams under auspices of zoo

IACUCs and were shared after approval by the appropriate committee at each zoo.

Antibacterial capacity

We measured antibacterial capacity against *E. coli* (ATCC #8739), *S. enterica* (ATCC

#13311), and *M. luteus* (ATCC # 4698) using an adaptation of the microbiocidal assay (a.k.a.

bacteria killing assay, BKA, microbiocidal activity) developed by French and Neuman-Lee

(2012). Briefly, we plated a 12-point dilution curve of each serum sample in triplicate on a

round-bottomed 96-well plate. *E. coli* had two serial dilutions: 6 points from undiluted serum to

1:64, and 6 points from 3:4 to 3:256. *S. enterica* also had two serial dilutions: 6 points from

undiluted to 1:64, and 6 points from 3:4 to 3:128. *M. luteus* had a single serial dilution ranging

from undiluted serum to 1:2048. Super antibacterial capacity curves—curves for species with

100% antibacterial capacity at the lowest dilution of the normal curves—for *E. coli* were 6 points

from 1:4 to 1:256 and 6 points from 3:16 to 3:512 and for were *S. enterica*: 6 points from 1:16 to

1:512 and 6 points from 3:4 to 3:128. We also plated a 4-point dilution curve in Dulbecco's

Phosphate Buffered Saline (PBS, Sigma-Aldrich #D8537) of commercially available cow serum

(Innovative Research Novi, MI 48377, #IBV-Compl) as our inter-assay control. Cow curve

dilutions were 1:32, 1:64, 1:128, 1:256 for *E. coli*; 1:16, 1:32, 1:64, 1:128 for *S. enterica*; and

1;20, 1:80, 1:320, 1:1280 for *M. luteus*. Our final volume of each serum dilution was 18 μl. We

plated three replicates of a 20 μl negative control of PBS and 9 replicates of an 18 μl positive

control. We added 2 μl of a standard concentration (10^4 CFU ml^{-1}) of bacteria to all wells except

the negative controls. We incubated plates for 30 min at 37 °C. We next added 125 μl of tryptic

soy broth (BD #211825) to all wells, and then shook plates for 1 min at 300 rpm. We then

measured baseline absorbance of all wells at 300 nm (Biotek Synergy HTX Multi-Mode Reader)

to serve as an internal control before incubating covered plates for 12 h for *E. coli*, 10 h for *S.*

enterica, and 48 h for *M. luteus* at an incubation temperature of 37 °C. We then measured final

absorbance of all wells again at 300 nm and calculated antibacterial capacity for each well as

$\left(1 - \frac{mean(sample\ end - sample\ baseline)}{mean(positive\ control\ start - positive\ control\ end)}\right) * 100\%$. Inter-assay (9.02%) and intra-assay

variation (0.02%) were both low. Key parameters of the antibacterial capacity assay for each

microbe are summarized in Table S1 and full details about the procedures used to develop and

validate our assays are available in the Supplementary Methods. Assay protocols are deposited

on FigShare (Schoenle et al. 2020).

Data processing and analysis

All data processing and analyses were performed in the statistical programming language

R (R Core Team 2019). We compiled data using a workflow developed using R (all code files

are available on Figshare (Keith et al. 2020; Downs et al. 2021)). We visually checked each

sample dilution curve for outliers (i.e., replicates that were distinct from the other two replicates

and/or fell off the expected sigmoidal curve). We determined whether positive controls, negative

controls, and inter-assay controls (i.e., cow sera) were within the expected range, and discarded a

plate if data fell outside expected ranges and reran samples.

To identify any allometries, we first fit non-linear antibacterial functions for each

individual sample separately. Properly estimating allometries required models that allowed us to

perform phylogenetically multivariate regressions while accounting for the sampling of

numerous individuals within a species because we know that species are not independent

samples (Felsenstein 1985) and a failure to account for within-species variation can produce

inaccurate regression estimates (Garamszegi and Møller 2010; Garamszegi 2014). This modeling

framework is under-served by the available statistical packages, so we modeled data using two

main approaches. For our first approach, we used multivariate phylogenetic comparative

analyses of our data using the R package `Rphylopars` (Goolsby et al. 2021). This approach

allowed us to calculate phylogenetic covariances of traits among species while accounting for

numerous observations within a species (i.e., among-individual variation) (Goolsby et al. 2017).

Although this approach did not estimate scaling coefficients as they would be estimated with a

regression/linear model-based approach, it allowed us to estimate phylogenetic signal of

antibacterial capacity and correlations among the antibacterial curve parameters and between

curve parameters and body mass in the same model. For our second approach, we built

multivariate general linear mixed models using the R package `MCMCglmm` (de Villemereuil and

Nakagawa 2014). Although these models assume each species is an independent point, they

allowed us to estimate b for all curve parameters simultaneously while accounting for the

variance within species by including multiple observations per species variation (Dingemanse

and Dochtermann 2013). This approach better parallels the traditional approach to estimating

scaling coefficients using regression models (Sieg et al. 2009). To test for the strength of

phylogenetic signals in a framework based on general linear models, we built phylogenetic

univariate general linear mixed models for each antibacterial curve parameter individually using

`MCMCglmm` (supplementary methods) (de Villemereuil and Nakagawa 2014). We extracted

species relatedness from a tree build using PhyloT (Letunic 2015, Fig S4)

<u>Fitting non-linear curves for each individual.</u> For the first step, we fit 5-parameter logistic

regression growth curves to each individual antibacterial curve (i.e., the shape of dilution curves

across all 12-serum concentrations) using package `nplr` (Commo and Bot 2016) (Fig. 1A). To

aid in curve-fitting, we $\log_{10}$-transformed serum concentrations (i.e., the dilutions) and converted

the antibacterial ability from a percent to a proportion. Curves could only be fit to values

between 0 and 1. So percent inhibited values > 100 were forced to a random value between 99

and 100 and percent inhibited values < 0 were forced to a random value between 0 and 1 before

conversion to proportions. This approach restricted values to a range near maximal and minimal

antibacterial activity for all other curves. We extracted the curve parameters (inflection$_{\text{dilution,}}$

inflection$_{\text{antibacterial capacity,}}$ bottom asymptote, top asymptote, slope, asymptote coefficient) to use in

univariate and multivariate general linear models (Fig. 1A). Slope and inflection$_{\text{dilution}}$ were

already on a $\log_{10}$-scale. We added 1 to inflection$_{\text{antibacterial capacity,}}$ bottom asymptote, top asymptote

and then $\log_{10}$-transfromed them. We also $\log_{10}$-transformed body mass. All analyses were

performed using transformed data.

<u>Phylogenetic covariance.</u> We performed a multivariate phylogenetic comparative analysis

for antibacterial capacity against each microbe using the R package `Rphylopars` (Goolsby et

al. 2021). All antibacterial capacity curve parameters and body mass were included as response

variables in these models, with one exception. Top asymptote was not included in the model for

antibacterial capacity against *S. enterica* because it was highly correlated (r = 0.998) with

Inflection$_{\text{antibacterial capacity}}$; both could not be included in the same model. We included

phylogenetic effects from a tree we produced by pruning the time-rooted phylogenetic tree

created by Uyeda et al. (2017) to our species list; we scaled tree with height scaled to 1. Models

were fit assuming a Brownian motion model of evolution and we estimated the percent of

variance explained by phylogeny. We tested for phylogenetic signal for each trait using a

likelihood ratio test (df = 1) to compare Pagel's lambda against a null model with a star

phylogeny (lambda = 0) to our model that included phylogeny (Pagel 1997, 1999).

247   <u>Multivariate mixed models.</u> We generated separate multivariate mixed models to query

body mass effects on antibacterial capacity, one for each bacterial species. All curve parameters

were included as response variables, and we allowed each parameter to have different slopes and

intercepts. Body mass was included as a fixed effect, and species was incorporated as a random

effect. Species had independent intercepts and slopes. These models were fit using the

`MCMCglmm` package (Hadfield 2010; Hadfield and Nakagawa 2010; de Villemereuil and

Nakagawa 2014). All mixed models were fit using a weak inverse-Gamma prior with shape and

scale parameters set to 1.002 for the random effect. Default priors for all other fixed effects were

used. Model chains were run for 1.82×10^6 iterations, a 420,000-iteration burn-in, and a 1400

iteration thinning interval. Results were robust across alternative priors, and chain length was

sufficient to yield negligible autocorrelation. We extracted the slopes describing the relationship

between each parameter and body mass across all species.

We included bacterial capacity at the bottom asymptote and asymmetric coefficient in our

models because they helped describe antibacterial capacity curves, although we did not make

predictions about their biological functions. In our study, we did not always collect data at the

lower asymptote because we limited our sampling to 12-dilutions. As a result, the bottom

asymptote estimated by the general linear model was based on incomplete data and not

biologically relevant to our scaling hypotheses. The dilution at the inflection point was also

included in our models because it informed our understanding of the antibacterial capacity at the

inflection point, but it was not obviously biologically meaningful for understanding immune

scaling.

Results

Overall results from all three modeling approaches were consistent and supported

isometric (i.e., mass-invariant) scaling (b=0) for almost all curve parameters for antibacterial

capacity parameters against all three microbes. The one exception to this pattern was the slope

for antibacterial capacity against *M. luteus*; in this one instance, all three modeling approaches

indicated a negative relationship between slope at the intercept with body mass such that slopes

become less steeply positive with body mass.

<u>Phylogenetic covariance.</u> Models built using `Rphylopars` estimated that the covariance

between body mass and any of the curve parameters was very close to zero and had 95% CI

intervals that overlapped zero, with one exception (Table 1): the correlation between slope and

body mass for antibacterial capacity against *M. leuteus* had a negative correlation of –11.0 (95%

CI: -22.8 : -2.6). Lambda values provided evidence of phylogenetic signal in body mass and at

least one curve parameter for antibacterial capacity against all three microbes (Table 2).

Phylogeny explained >97% for variation in body mass, but only $\leq 30\%$ of the variation in most

of the antibacterial capacity curve parameters (Table 2). An exception was that phylogeny

explained 54% if of the variation in Inflection$_{\text{dilution}}$ for antibacterial capacity against *S. enterica*.

Phylogenetic correlations among curve parameters and phylotypic correlations can be found in

Tables S1 and S2, respectively. These models had slightly smaller sample sizes because not all

species were in the time-rooted tree. Final sample sizes of species for these models were 191,

178, and 157 for antibacterial capacity against *E. coli, S. enterica,* and *M. luteus*, respectively

<u>Phylogenetic univariate mixed effects models.</u> Phylogenetic univariate models supported

scaling coefficients of zero. The 95% CI for all curve parameters for antibacterial capacity

against *E. coli, S. enterica, M. luteus* overlapped zero except for slope of antibacterial curves

against *M. luteus* (supplemental materials, Table S3). This parameter scaled hypometerically (-

4.30, 95% CI: -8.34, -0.30). Phylogeny explained 1-18% of the variation in antibacterial curve

parameters (supplemental materials, Table S4).

<u>Multivariate mixed effects models.</u> Multivariate models supported scaling coefficients of

zero for all curve parameters for antibacterial capacity against *E. coli, S. enterica, M. luteus* with

two exceptions (Table 3, Fig. 2) The slope of antibacterial curves against *M. luteus* had a

hypermetric slope of -4.3 (95% CI: -7.7 : -0.79) and the asymmetric coefficient scaled with a

slope of -3.78 (95% CI: -5.604 : -0.91).

Discussion

Our data provide consistent and robust evidence for isometric (i.e., mass-invariant)

scaling of serum antibacterial activity in terrestrial mammals. Additionally, results indicate that

phylogenic signal was surprisingly low for all three microbes. Superficially, isometric scaling

supports the prediction of the Protection and Complexity Hypotheses. These results were

somewhat surprising, given that large species face a greater risk of infection and face greater lost

fitness costs from infections (Downs et al. 2020). Below we argue that the interpretation of these

results might support a more complex interpretation once integrated with our understanding of

how host metabolic rates and other immunological defenses scale.

Mass-invariance of antibacterial capacity was consistent regardless of the bacterial

species studied. This discovery is interesting in light of the ecology and life history differences

among bacterial species. *M. luteus* grows slowly and causes modest disease in most hosts relative

to the other two bacteria; both *S. enterica* and *E. coli* can cause sepsis (van der Poll and Opal

2008), infect via the diet, and have similar replication rates in culture. Our findings thus support

the idea that broadly effective, fast-acting immune responses are generally favored, rather than

by a specific pathogenetic target, across mammalian species.

<u>Isometric scaling and the Protecton/Complexity Hypotheses</u>

Both of the Protecton and Complexity Hypotheses hypotheses argue that large and small

animals need equal defenses against parasites, resulting in immune responses having a scaling

exponent of 0 (Langman and Cohn 1987; Wiegel and Perelson 2004; Banerjee and Moses 2010).

Our results are consistant with this prediction. However, by definition, isometric scaling is

difficult to test because it corresponds with a null model of a mass-invariant or intercept-only

model. Statistically, we cannot distinguish between the null and the model prediction.

Perhaps more importantly, both the Protecton and Complexity Hypotheses were

developed to predict how the number of lymphocyte clones should scale with body size to

provide protection against foreign antigens (Langman and Cohn 1987; Wiegel and Perelson

2004; Banerjee and Moses 2010), not to explain constitutive, innate immunity as studied here.

Assumptions about the distribution and timing of adaptive immune responses are unlikely to

hold for the early-acting, constitutive innate immune responses. The adaptive and innate immune

systems operate on different time scales (Murphy et al. 2007), and the life cycles of lymphocytes

and neutrophils are quite different. Circulating peripheral neutrophils have a lifespan of 1-7 days

(Hidalgo et al. 2018) whereas peripheral, naïve lymphocytes are long lived—T cells live on the

order of years and B cells have a half-life of 4 weeks to several months—and later phases of

adaptive immunity (i.e., memory) are controlled by even longer-lived lymphocytes (Tough and

Sprent 1995). Even "short-lived" lymphocytes involved in early stages of memory have life-

spans of weeks, which is much longer than neutrophils (Tough and Sprent 1995). Although the

Complexity Hypothesis has been extended to make predictions about the size and number of

lymph nodes in mammals (Banerjee 2018) and host capacity to replicate West Nile virus

(Banerjee and Moses 2009, 2010; Banerjee et al. 2017), these extensions are still primarily based

on the dynamics of adaptive immunity. It follows that the underpinning logic of the Complexity

Hypothesis would need to be rederived from first principles to account for dynamics of innate

immune defense and to make predictions for early, broadly active defenses that are part of the

innate immune system, such as antibacterial capacity.

<u>Isometric scaling is steeper than expected by metabolic rates</u>

The mass-invariance of antibacterial capacity is intriguing when considered with the

well-documented hypometric scaling of mass-specific metabolic rates (Brown et al. 2004;

Savage et al. 2004) and the hypermetric scaling of concentrations of some immune cell types

(Downs et al. 2020). The Metabolic Theory of Ecology posits that mass-specific metabolic rate

can be interpreted as the ratio of the average metabolic rate of a cell to the average cell volume,

and this ratio also has a scaling exponent of -0.25 (Savage et al. 2007):

$$\frac{metabolic\ rate}{body\ mass} = \frac{mean\ cellular\ metabolic\ rate}{mean\ cell\ volume} \propto body\ mass^{-0.25}.$$

If cellular metabolic rate increases, cell volume must decrease (Savage et al. 2007). Thus, for cell

types with mass-invariant volumes (b = 0), cellular metabolic rate will scale with the same slope

as mass-specific metabolic rate (b = -0.25). It follows that if basal metabolic rate is the

pacemaker for cellular activity, protein synthesis for these cell types would also scale with a

slope of -0.25 (Brown et al. 2004).

Our assay measured the functional capacity of complement and other circulating proteins

(i.e., lysozyme, mannan-binding protein, beta-defensins) to opsonize and lyse bacteria (Demas et

al. 2011; French and Neuman-Lee 2012). Complement proteins are predominantly produced by

hepatocytes (Lubbers et al. 2017), and hepatocyte volume is mass-invariant (Savage et al. 2007).

As such, production of complement protein production per hepatocyte should scale at -0.25.

Liver mass scales with a slope of 0.895 (Prothero 2015), so. if each hepatocyte produces

complement proteins at the same rate, antibacterial capacity should scale at 0.645 (= 0.895-0.25).

Within this framework, a scaling coefficient of 0 for antibacterial capacity, as we described, is

shockingly high. In other words, large mammals have much higher antibacterial capacity than

expected by their metabolic rate alone. An outstanding question is how large mammalian species

obtain the same antibacterial capacity as small ones if they are constrained by their metabolic

rates.

Why isometric scaling?

Antibacterial capacity and neutrophil concentrations, the other type of first, constitutive

lines of host defense thus far studied in an allometric context (b = 0.11; Downs et al. 2020), both

scaling more steeply than expected by scaling of metabolic rates. These scaling repationships

may have evolved because, relative to small mammals, large mammals (1) have a

disproportionately higher exposure to parasites and (2) have a disproportionately larger gap in

replication rates relative to invading microbes (Downs et al. 2020). We will walk through each in

turn.

First, large mammals have different exposure risks to parasites because of their ecology.

Relative to small mammals, large mammals consume more food (Nagy 2001; Nunn et al. 2003),

move farther and thus are exposed to more of habitat with each unit of movement (e.g., step,

wing flap) (Schmidt-Nielsen 1984), and they have larger home ranges which exposes them to

more types of habitats (Lindstedt et al. 1986; Kelt and Van Vuren 2001). Although a given gram

of tissue in a large host is less exposed to parasites because of a low ratio of surface area to

volume, absolute exposure to parasites is probably much higher in large than small species

(Dobson and Hudson 1986; Downs et al. 2019). Large mammals also have a higher absolute

exposure to parasites over their lifetime because they have longer lifespans than small mammals

(Peters 1983; Calder 1984).

Second, parasites have an evolutionary advantage over hosts because of differences in

replication rates. When parasites grow rapidly, as bacterial (and viral) infections often do,

defenses with short time-delays are favored (Shudo and Iwasa 2001). Barriers against all-comers

are the best option when adversaries can replicate faster than defenses can be honed and

mobilized. The downside to these forms of defense is that they impose comparatively high costs,

especially when adversaries are absent (Sears et al. 2006). Recently, a meta-analysis revealed

that small animal species that are relatively long-lived incur the highest costs of *induced* immune

defenses (Brace et al. 2017). Given the tendency of small species to breed prolifically and

develop rapidly, isometric allometries might manifest as much because small species favor

rapidly-mobilizable defenses that are activated only after exposure to a parasite as they do

because of mandates of large size.

Our understanding of how activation of adaptive immune responses occurs would also

support the need for large animals to have higher early constitutive immune defenses than

expected by metabolic rates. B and T cell receptor diversity is more constrained by over-

responsiveness to self- than under-responsiveness to non-self (De Boer and Perelson 1993).

Indeed, the probability that a parasite will evade the antigen space covered by B and T cell

repertoires is 4×10^{-44} (Perelson and Oster 1979; Banerjee et al. 2017). Adaptive immune systems

will, therefore, eventually detect any antigen that ever evolves. The challenge of relying solely

on these defenses, though, is that they involve major time lags (i.e., minimally 4-10 days but

perhaps longer if the immune system is still developing (Ricklefs 1992). Lags for inducible

defenses generally leave hosts vulnerable as microbial doubling time proceeds over a much

shorter period. This incompatibility might explain why among 25 avian species, the production

rate of infectious virions and the burst size of cells (i.e., the number of infectious virions released

by an infected cell over its infected lifespan) scaled hypometrically, yet the rate of antibody-

mediated neutralization of virus was isometric (Banerjee et al. 2017). Perhaps isometrically-

scaled constitutive immune defenses produced the surprisingly low viral outcomes observed in

that study.

If considered more broadly, our results might in fact be consistent with the Safety Factory

Hypothesis. Perhaps instead of predicting hypermetric scaling of all forms of constitutive innate

immunity, the Safety Factory Hypothesis should instead posit that large and small animals need

equal *functional* protection against pathogens resulting in hypermetric scaling of some defenses

and isometric (or even hypometric) scaling of others. Large mammals (Downs et al. 2020) and

large birds (Ruhs et al. 2020) have disproportionally higher concentrations of early responding

granulocytes relative to their small counterparts. However, this hypermetric scaling may not

translate into hypermetric scaling of functionally because the synthesis of immunological

molecules and cellular rates of reaction might be constrained by metabolic rate (Brown et al.

2004; Savage et al. 2007). In other words, the disproportionately high concentrations of

granulocytes in large species might be necessary to have similar functionality as small species

(Downs et al. 2020; Ruhs et al. 2020). Large animals may use a strategy of many cells with

reduced activity, whereas small animals may use a strategy of fewer cells with higher activity.

To test further test this idea, we would need to determine the scaling coefficient for the

functionality of neutrophils and the scaling of the number of hepatocytes that produce

complement and their functionality.

<u>Phylogenetic signal and other influences on antibacterial capacity</u>

Our results also demonstrated that mammals exhibit a high diversity of antibacterial

curves (Movie S1) and that little of this variation is explained by phylogeny or body size. The

weak phylogenetic signal and a small percentage of variation in antibacterial curves explained by

phylogeny in general were surprising. Previous analyses of scaling of leucocyte concentrations

and in mammals (Downs et al. 2020) and birds (Ruhs et al. 2020) and leucocyte ratios in birds

(Minias 2019) demonstrated that phylogeny explained the majority of interspecific variation.

Although phylogenetic signal was weak for many aspects of antibacterial capacity we measured,

we caution against equating a lack of signal with a lack of phylogenetic effect (Blomberg and

Garland 2002),. Our statistics were not designed to test the coevolution of these traits over time,

rather they look for phylogenetic covariances in the extant traits

Life history, diet, and other differences may also explain some variation in antibacterial

capacity curves among mammalian species. The pace-of-life hypothesis posits that reproductive

rate and longevity, among other life-history traits, should shape investment in immune defenses

(Lee 2006). Although life histories are closely tied to body mass in mammals (Peters 1983;

Calder 1984), species of the same mass, but with different life histories, will have different costs

and benefit structures for immune defense. For example, small species that are relatively long-

lived incur the higher costs of activating immune responses (Brace et al. 2017), natural antibody

titers increased with incubation period in 70 species of tropical birds after accounting for body

mass statistically (Lee et al. 2008), and faster-lived birds produced more of an acute-phase

protein after inoculation with *E. coli* after accounting for body mass (Robinson et al. 2010).

Determining if life history explains the residual variation after accounting for body mass and

phylogeny is a next step towards understanding the variation in interspecific variation in

antibacterial capacity, but are beyond the scope of theis current study

Conclusion

Our findings suggest that large and small species might use different strategies to obtain

the same level of protection against parasites. They have ramifications for human medicine and

management of disease cycles, especially zoonoses, and a greater understanding of immune

scaling could also help improve epidemiological models. A previously developed allometric

framework for host-species competence for multi-host parasites found that data on interspecific

allometries of immune defenses was lacking in the literature (Downs et al. 2019). Because "the

rate that pathogens spread through populations is influenced by the rate of spread through

individual hosts" (De Leo and Dobson 1996), we must understand better how host traits affect

their propensity to maintain and transmit viable parasites (Bolzoni et al. 2008; De Leo et al.

2016).

We expect that many more important immune allometries await description. In particular,

we encourage future scaling work on Order *Chiroptera*, as bats are implicated in the emergence

of many zoonoses including SARS-COV2 (Han et al. 2016; Olival et al. 2017). Unfortunately,

bats are rare in zoos, so few bat species were in the present analysis (but see Ruhs et al. In press).

We also encourage efforts to investigate immune allometries in ectotherms, and especially

vectors, as their immune defenses are more temperature-dependent than mammals (Brown et al.

2004), an important condition for a planet undergoing extensive climate change. We also hope

creative approaches can be applied to scaling studies of more pathogenic microbes, ideally in

more organismal contexts. Such work is challenging, as pathogenicity differs among species as

well as route of infection (Taylor 1983). By continuing to merge cellular and molecular efforts

with the behavioral, ecological, and evolutionary concepts, we can produce a more integrative

immunology and thus more effective options to predict and manage infectious disease risk.

Author Contributions: C.J.D., L.B.M, and K.C.K. conceptualized the study; L.A.S., S.J.O. and

C.J.D., and R.B. procured samples; L.A.S., S.J.O. and C.J.D. established and validated the

laboratory methodology; L.A.S., S.J.O., C.J.D., and L.B.M collected and curated the data; C.J.D.

carried out the formal analysis with input from L.B.M and E.W.G; C.J.D and L.B.M drafted the

manuscript with input from K.C.K; all authors contributed to writing and revision; C.J.D and

L.B.M. administered and supervised the project; C.J.D., L.B.M., and R.H.Y.J acquired funding.

Acknowledgements: We thank Birmingham Zoo, Busch Gardens Tampa Bay (Cara Martel),

Cleveland Metroparks Zoo, Columbus Zoo, the Denver Zoological Foundation, Dickerson Park

Zoo, Indianapolis Zoo, Lincoln Park Zoo, Louisville Zoological Garden, Maryland Zoo in

Baltimore, Milwaukee County Zoo, Naples Zoo, Nashville Zoo at Grassmere, Oregon Zoo

(Mitch Finegan & David Shepherdson), Sedgwick County Zoo, Smithsonian's National Zoo and

Conservation Biological Institute, Topeka Zoo & Conservation Center, Zoo Atlanta (Hayley

Murphy, Stephanie Earhardt), ZooTampa at Lowry Park, and the Demas (Indiana State

University), Hoekstra (Harvard University), Ophir (Cornell University), Place (Cornell

University), and Monteith (University of Wyoming) labs for donating serum samples. We thank

M. Amy, B. Butler, K. Carlson, E. Chinchilli, K. Clausen, H. Droke, M. Espino, A. Fernandez,

M. Galan, M. Henning, N. Huizenga, K. Koller, E. Lee, O. Ogunsina, V. Pappademetriou, A.

Roberts, C. Stanell, S. Travis, A. Vorrath, R. Winner, C. Yeung for help with laboratory work.

N. Keith helped develop the Program R workflows required to process our data. We thank N.A.

Dochtermann and J.C. Uyeda for statistical guidance, and the Martin Lab, HA Woods, J.F.

Harrison, ME Sobolewski, and V Savage for discussion about an early version of the manuscript.

This work was supported by State University of New York College of Environmental Science

and Forestry's Provost Office, the Levitt Center, Hamilton College's Dean of Faculty Office,

University of South Florida's College of Public Health, and the National Science Foundation

(award numbers IOS 1656551 to C.J.D. and IOS 1656618 to L.B.M).

Data Accessibility: The datasets generated during and/or analyzed during the current study and

will be deposited in Figshare after a 3-year embargo. We will consider written requests for the

data prior to the end of the embargo. Lab protocols (Schoenle et al. 2020) and R code (Keith et

al. 2020; Downs et al. 2021) are available on FigShare.

References

Banerjee, S. 2018. A mathematical framework for understanding how lymph node architecture scales with host body
size to produce an efficient immune response 8.

Banerjee, S., and M. Moses. 2010. Scale Invariance of Immune System Response Rates and Times: Perspectives on
Immune System Architecture and Implications for Artificial Immune Systems. arXiv:1008.1380 [cs, math, q-bio].

Banerjee, S., and M. E. Moses. 2009. A Hybrid Agent Based and Differential Equation Model of Body Size Effects
on Pathogen Replication and Immune System Response. Pages 14–18 *in* P. S. Andrews, J. Timmis, N. D. L. Owens,

U. Aickelin, E. Hart, A. Hone, and A. M. Tyrrell, eds. Artificial Immune Systems (Vol. 5666). Springer Berlin
Heidelberg, Berlin, Heidelberg.

Banerjee, S., A. S. Perelson, and M. Moses. 2017. Modelling the effects of phylogeny and body size on within-host
pathogen replication and immune response. Journal of The Royal Society Interface 14:20170479.

Bennett, A. F., and R. B. Huey. 1990. Studying the evolution of physiological performance. Oxford surveys in
evolutionary biology 7:251–284.

Blomberg, S. P., and T. Garland. 2002. Tempo and mode in evolution: phylogenetic inertia, adaptation and
comparative methods: Phylogenetic inertia. Journal of Evolutionary Biology 15:899–910.

Blount, J. D., D. C. Houston, A. P. Moller, and J. Wright. 2003. Do individual branches of immune defence
correlate? A comparative case study of scavenging and non-scavenging birds. Oikos 102:340–350.

Bolzoni, L., G. A. De Leo, M. Gatto, and A. P. Dobson. 2008. Body-size scaling in an SEI model of wildlife
diseases. Theoretical Population Biology 73:374–382.

Brace, A. J., M. J. Lajeunesse, D. R. Ardia, D. M. Hawley, J. S. Adelman, K. L. Buchanan, J. M. Fair, et al. 2017.
Costs of immune responses are related to host body size and lifespan. Journal of Experimental Zoology Part A:
Ecological and Integrative Physiology 327:254–261.

Brown, J. H., J. F. Gillooly, A. P. Allen, V. M. Savage, and G. B. West. 2004. Toward a Metabolic Theory of
Ecology. Ecology 85:1771–1789.

Calder, W. A., III. 1984. Size, function, and life history. Dover Publications, Mineola, NY.

Commo, F., and B. M. Bot. 2016. R package nplr n-parameter logistic regressions 11.

De Boer, R. J., and A. S. Perelson. 1993. How diverse should the immune system be? Proceedings of the Royal
Society of London. Series B: Biological Sciences 252:171–175.

De Leo, G. A., and A. P. Dobson. 1996. Allometry and simple epidemic models for microparaites. Nature 379:720–
722.

De Leo, G. A., A. P. Dobson, and M. Gatto. 2016. Body size and meta-community structure: the allometric scaling
of parasitic worm communities in their mammalian hosts. Parasitology 143:880–893.

de Villemereuil, P., and S. Nakagawa. 2014. General Quantitative Genetic Methods for Comparative Biology. Pages
287–303 *in* L. Z. Garamszegi, ed. Modern Phylogenetic Comparative Methods and Their Application in
Evolutionary Biology. Springer Berlin Heidelberg, Berlin, Heidelberg.

Demas, G. E., D. A. Zysling, B. R. Beechler, M. P. Muehlenbein, and S. S. French. 2011. Beyond
phytohaemagglutinin: assessing vertebrate immune function across ecological contexts: Assessing vertebrate
immune function across ecological contexts. Journal of Animal Ecology 80:710–730.

Diamond, J. 2002. Quantitative evolutionary design. The Journal of Physiology 542:337–345.

Dingemanse, N. J., and N. A. Dochtermann. 2013. Quantifying individual variation in behaviour: mixed-effect
modelling approaches. (M. van de Pol, ed.)Journal of Animal Ecology 82:39–54.

Dingli, D., and J. M. Pacheco. 2006. Allometric Scaling of the Active Hematopoietic Stem Cell Pool across
Mammals. (C. Arendt, ed.)PLoS ONE 1:e2.

Dobson, A. P., and P. J. Hudson. 1986. Parasites, disease and the structure of ecological communities. Trends in
Ecology & Evolution 1:11–15.

Downs, C. J., J. S. Adelman, and G. E. Demas. 2014. Mechanisms and Methods in Ecoimmunology: Integrating
Within-Organism and Between-Organism Processes. Integrative and Comparative Biology 54:340–352.

Downs, C. J., N. A. Dochtermann, R. Ball, K. C. Klasing, and L. B. Martin. 2020. The Effects of Body Mass on
Immune Cell Concentrations of Mammals. American Naturalist 195:107–114.

Downs, C. J., E. W. Goolsby, and L. B. Martin. 2021. Rphylopars code for comparative antimicrobial analysis.
FigShare. DOI:10.6084/m9.figshare.14806962

Downs, C. J., L. A. Schoenle, B. A. Han, J. F. Harrison, and L. B. Martin. 2019. Scaling of Host Competence.
Trends in Parasitology 35:182–192.

Ezenwa, V. O. 2004. Interactions among host diet, nutritional status and gastrointestinal parasite infection in wild
bovids. International Journal for Parasitology 34:535–542.

Fassbinder-Orth, C. A., L. D. Igl, D. C. Hahn, K. M. Watts, T. E. Wilcoxen, and K. R. Ramos-Álvarez. 2019. Do
Life History Traits Influence Patterns of Maternal Immune Elements in New World Blackbirds (Icteridae)?
Integrative Organismal Biology 1.

Felsenstein, J. 1985. Phylogenies and the Comparative Method. The American Naturalist 125:1–15.

Filion, A., A. Eriksson, F. Jorge, C. N. Niebuhr, and R. Poulin. 2020. Large-scale disease patterns explained by
climatic seasonality and host traits. Oecologia.

French, S. S., and L. A. Neuman-Lee. 2012. Improved ex vivo method for microbiocidal activity across vertebrate
species. Biology Open 1:482–487.

Garamszegi, L. Z. 2014. Uncertainties Due to Within-Species Variation in Comparative Studies: Measurement
Errors and Statistical Weights. Pages 157–199 in L. Z. Garamszegi, ed. Modern Phylogenetic Comparative Methods
and Their Application in Evolutionary Biology: Concepts and Practice. Springer Berlin Heidelberg, Berlin,
Heidelberg.

Garamszegi, L. Z., and A. P. Møller. 2010. Effects of sample size and intraspecific variation in phylogenetic
comparative studies: a meta-analytic review. Biological Reviews no-no.

Goolsby, E. W., J. Bruggeman, and C. Ané. 2017. Rphylopars : fast multivariate phylogenetic comparative methods
for missing data and within-species variation. (R. Fitzjohn, ed.)Methods in Ecology and Evolution 8:22–27.

Goolsby, E. W., J. Bruggeman, and C. Ane. 2021. Rphylopars: Phylogenetic Comparative Tools for Missing Data
and Within-Species Variation.

Hadfield, J. D. 2010. MCMC Methods for Multi-Response Generalized Linear Mixed Models: The **MCMCglmm** *R*
Package. Journal of Statistical Software 33.

Hadfield, J. D., and S. Nakagawa. 2010. General quantitative genetic methods for comparative biology: phylogenies,
taxonomies and multi-trait models for continuous and categorical characters. Journal of Evolutionary Biology
23:494–508.

Han, B. A., A. M. Kramer, and J. M. Drake. 2016. Global Patterns of Zoonotic Disease in Mammals. Trends in
Parasitology 32:565–577.

Harrison, J. F. 2017. Do Performance–Safety Tradeoffs Cause Hypometric Metabolic Scaling in Animals? Trends in
Ecology & Evolution 32:653–664.

Harrison, L. 1915. Mallophaga from *Apteryx* , and their Significance; with a Note on the Genus *Rallicola*.
Parasitology 8:88–100.

Ishtiaq, F., C. G. R. Bowden, and Y. V. Jhala. 2017. Seasonal dynamics in mosquito abundance and temperature do
not influence avian malaria prevalence in the Himalayan foothills. Ecology and Evolution 7:8040–8057.

Keith, N. I., L. B. Martin, and C. J. Downs. 2020. Code to compile data and analyzing data for a comparative
antimicrobial analysis. figshare. DOI: 10.6084/m9.figshare.12430103

Kelt, D. A., and D. H. Van Vuren. 2001. The Ecology and Macroecology of Mammalian Home Range Area. The
American Naturalist 157:637–645.

Kieft, T. L., and K. A. Simmons. 2015. Allometry of animal–microbe interactions and global census of animal-
associated microbes. Proceedings of the Royal Society B: Biological Sciences 282:20150702.

Kuris, A. M., A. R. Blaustein, and J. J. Alio. 1980. Hosts as islands. American Naturlists 116:570–586.

Langman, R. E., and M. Cohn. 1987. The E-T (Elephant-Tadpole) paradox necessitates the concept of a unity of B-
cell function: The protecton. Molecular Immunology 24:675–697.

Lee, K. A. 2006. Linking immune defenses and life history at the levels of the individual and the species. Integrative
and Comparative Biology 46:1000–1015.

Lee, K. A., M. Wikelski, W. D. Robinson, T. R. Robinson, and K. C. Klasing. 2008. Constitutive Immune Defences
Correlate with Life-History Variables in Tropical Birds. Journal of Animal Ecology 77:356–363.

Letunic, I. 2015. phyloT: phylogenetic tree generator

Lindstedt, S. L., B. J. Miller, and S. W. Buskirk. 1986. Home Range, Time, and Body Size in Mammals. Ecology
67:413–418.

Lubbers, R., M. F. van Essen, C. van Kooten, and L. A. Trouw. 2017. Production of complement components by
cells of the immune system: Production of complement components. Clinical & Experimental Immunology
188:183–194.

Mastroeni, P., J. A. Chabalgoity, S. J. Dunstan, D. J. Maskell, and G. Dougan. 2001. Salmonella: Immune
Responses and Vaccines. The Veterinary Journal 161:132–164.

Minias, P. 2019. Evolution of heterophil/lymphocyte ratios in response to ecological and life-history traits: A
comparative analysis across the avian tree of life. (D. Ardia, ed.)Journal of Animal Ecology 88:554–565.

Mittrücker, H.-W., and S. H. E. Kaufmann. 2000. Immune response to infection with *Salmonella typhimurium* in
mice. Journal of Leukocyte Biology 67:457–463.

Murphy, K. P., P. Travers, and M. Walport. 2007. Janeway's immunobiology. (7th ed.). Garland Science, New
York.

Nagy, K. A. 2001. Food requirements of wild animals: predictive equations for free-living mammals, reptiles, and
birds. Nutrition Abstracts and Reviews, Series B 71:21R-31R.

Nunn, C. L. 2002. A COMPARATIVE STUDY OF LEUKOCYTE COUNTS AND DISEASE RISK IN
PRIMATES. Evolution 56:177–190.

Nunn, C. L., J. L. Gittleman, and J. Antonovics. 2003. A comparative study of white blood cell counts and disease
risk in carnivores. Proceedings of the Royal Society of London. Series B: Biological Sciences 270:347–356.

Olival, K. J., P. R. Hosseini, C. Zambrana-Torrelio, N. Ross, T. L. Bogich, and P. Daszak. 2017. Host and viral traits
predict zoonotic spillover from mammals. Nature 546:646–650.

Pagel, M. 1997. Inferring evolutionary processes from phylogenies. Zoologica Scripta 26:331–348.

———. 1999. Inferring the historical patterns of biological evolution. Nature 401:877–884.

Perelson, A. S., and G. Oster. 1979. Theoretical Studies of Clonal Selection: Minimal Antibody Repertoire Size and
Reliability of Self-Non-self Discriminationt. Journal of Theoretical Biology 81:645–670.

Peters, R. H. 1983. The ecological implications of body size. Cambridge University Press, New York.

Peto, R. 1977. Epidemiology, multistage models, and short-term mutagenicity tests. Pages 1403–1428 *in* J. D.
Watson, H. H. Hiatt, and J. A. Winsten, eds. Origins of human cancer. Cold Spring Harbor Laboratory, New York,
NY.

Poulin, R. 2007. Evolutionary ecology of parasites. Princeton University Press, Princeton, NJ.

Poulin, R., and M. George-Nascimento. 2007. The scaling of total parasite biomass with host body mass.
International Journal for Parasitology 37:359–364.

Prothero, J. W. 2015. The Design of Mammals: A Scaling Approach. Cambridge University Press, Cambridge.

R Core Team. 2019. R: A language and environment for statistical computing. R Foundation for Statistical
Computing, Vienna, Austria.

Ricklefs, R. E. 1992. Embryonic development period and the prevalence of avian blood parasites. Proceedings of the
National Academy of Sciences 89:4722–4725.

Robinson, W. D., M. Hau, K. C. Klasing, M. Wikelski, J. D. Brawn, S. H. Austin, C. Tarwater, et al. 2010.
Diversification of Life Histories in New World Birds. The Auk 127:253–262.

Ruhs, E. C., D. J. Becker, S. J. Oakey, O. Ogunsina, M. B. Fenton, Simmons, Nancy B., L. B. Martin, et al. In press.
Body size affects immune cell proportions in birds and non-volant mammals, but not bats. Journal of Experimental
Biology.

Ruhs, E. C., L. B. Martin, and C. J. Downs. 2020. The impact of body mass on immune cell concentrations in birds.
Proceedings of the Royal Society B: Biological Sciences 287:20200655.

Savage, V. M., A. P. Allen, J. H. Brown, J. F. Gillooly, A. B. Herman, W. H. Woodruff, and G. B. West. 2007.
Scaling of number, size, and metabolic rate of cells with body size in mammals. Proceedings of the National
Academy of Sciences 104:4718–4723.

Savage, V. M., J. F. Gillooly, W. H. Woodruff, G. B. West, A. P. Allen, B. J. Enquist, and J. H. Brown. 2004. The
predominance of quarter-power scaling in biology. Functional Ecology 18:257–282.

Schmidt-Nielsen, K. 1984. Scaling: why is animal size so important? Cambridge University Press, Cambridge.

Schneeberger, K., G. Á. Czirják, and C. C. Voigt. 2013. Measures of the Constitutive Immune System Are Linked to
Diet and Roosting Habits of Neotropical Bats. (J. D. Brown, ed.)PLoS ONE 8:e54023.

Schoenle, L. A., C. J. Downs, and L. B. Martin. 2018. An Introduction to Ecoimmunology. Pages 901–932 *in* E. L.
Cooper, ed. Advances in Comparative Immunology. Springer International Publishing, Cham.

Schoenle, L. A., L. B. Martin, and C. J. Downs. 2020. Protocols for 12-dilution antibacterial capacity curves for
interspecific comparisons. Figshare. DOI: 10.6084/m9.figshare.12501149

Sears, M. W., J. P. Hayes, C. S. O'Connor, K. Geluso, and J. S. Sedinger. 2006. Individual variation in thermogenic
capacity affects above-ground activity of high-altitude Deer Mice. Functional Ecology 20:97–104.

Sheldon, B. C., and S. Verhulst. 1996. Ecological immunology: costly parasite defences and trade-offs in
evolutionary ecology. Trends in Ecology & Evolution 11:317–321.

Shudo, E., and Y. Iwasa. 2001. Inducible Defense against Pathogens and Parasites: Optimal Choice among Multiple
Options. Journal of Theoretical Biology 209:233–247.

Sieg, A. E., M. P. O'Connor, J. N. McNair, B. W. Grant, S. J. Agosta, and A. E. Dunham. 2009. Mammalian
Metabolic Allometry: Do Intraspecific Variation, Phylogeny, and Regression Models Matter? The American
Naturalist 174:720–733.

Smith, F. A., and S. K. Lyons. 2011. How big should a mammal be? A macroecological look at mammalian body
size over space and time. Philosophical Transactions of the Royal Society B: Biological Sciences 366:2364–2378.

Taylor, P. W. 1983. Bactericidal and bacteriolytic activity of serum against gram-negative bacteria. Microbiological
Reviews 47:46–83.

Tian, J., A. Courtiol, K. Schneeberger, A. D. Greenwood, and G. Á. Czirják. 2015. Circulating white blood cell
counts in captive and wild rodents are influenced by body mass rather than testes mass, a correlate of mating
promiscuity. (T. Piersma, ed.)Functional Ecology 29:823–829.

Tough, D. F., and J. Sprent. 1995. Lifespan of lymphocytes. Immunologic Research 14:1–12.

Uyeda, J. C., M. W. Pennell, E. T. Miller, R. Maia, and C. R. McClain. 2017. The Evolution of Energetic Scaling
across the Vertebrate Tree of Life. The American Naturalist 190:185–199.

van der Poll, T., and S. M. Opal. 2008. Host–pathogen interactions in sepsis. The Lancet Infectious Diseases 8:32–
43.

Wiegel, F. W., and A. S. Perelson. 2004. Some Scaling Principles for the Immune System. Immunology and Cell
Biology 82:127–131.

Table 1. Covariance (Mean ± 95% CI) between body mass and antibacterial capacity curve

parameters from phylogenetic covariance models estimated using the `Rphylopars` in Program

R.

Antibacterial capacity curve parameter	Estimate of covariance by dataset		
	E. coli	*S. enterica*	*M. luteus*
Top asymptote[#]	-0.002 (-0.036 : 0.035)	N/A*	-0.024 (-0.113 : 0.064)
Slope	9.50 (-4.3 : 26.3)	-0.394 (-11.9 : 12.6)	-11.0 (-22.8 : -2.6)
Inflection$_{dilution}$	-0.262 (-0.723 : 0.360)	-0.126 (-0.747 : 0.465)	-0.103 (-0.954 : 0.947)
Asymptote coefficient	-0.022 (-0.345 : 0.178)	-0.107 (-0.341 : 0.033)	-0.022 (-0.390 : 0.156)
Inflection$_{antibacterial\ capacity}$	-0.006 (-0.284 : 0.036)	0.009 (-0.146 : 0.445)	-0.023 (-0.087 : 4.47)
Bottom asymptote	-0.002 (-1.3×10^{78} : 2.1×10^{95})	-0.009 (-0.409 : 0.046)	-0.041 (-2.0×10^{33} : 3.61)

697 [#] Inflection$_{antibacterial\ capacity}$, top asymptote, and bottom asymptote were +1, $\log_{10}$-transformed. Body mass and slope

698 were $\log_{10}$-transformed prior to beta estimation.

699 * Top asymptote was not included in the model for antibacterial capacity against *S. enterica* because it was highly

700 correlated (r = 0.998) with Inflection$_{antibacterial\ capacity}$. Both variables could not be included in the same model

Table 3. Estimates of the percent of variation in antibacterial curves against *E. coil*, *S. enterica,* and *M. luteus* explained by phylogeny from phylogenetic covariance models estimated using the `Rphylopars` in Program R.

Antibacterial capacity curve parameter	Lambda (P-value)			Variance explained (%)		
	E. coli	*S. enterica*	*M. luteus*	*E. coli*	*S. enterica*	*M. luteus*
Body mass*	**1** (**<0.001**)	**1** (**<0.001**)	**1** (**<0.001**)	98.0	97.5	97.5
Top asymptote	0.008 (1)	N/A	0.179 (0.449)	11.6	N/A	24.6
Slope	**0.971** (**<0.001**)	0.996 (0.137)	0.445 (0.016)	28.1	5.47	17.0
Inflection$_{\text{antibacterial capacity}}$	0.075 (0.984)	0.474 (0.363)	0.221 (0.249)	10.7	30.0	23.4
Inflection$_{\text{dilution}}$	**0.996** (**0.002**)	**0.999** (**<0.001**)	**0.998** (**0.004**)	20.2	54.2	16.4
Asymptote coefficient	0.753 (0.378)	0.358 (0.492)	0.198 (0.872)	3.8	3.7	7.1
Bottom asymptote	**0.976** (**<0.001**)	0.431 (0.162)	0.003 (1)	31.1	18.8	10.6

* Inflection$_{\text{antibacterial capacity}}$, top asymptote, and bottom asymptote were +1, log$_{10}$-transformed. Body mass and slope were log$_{10}$-transformed prior to beta estimation.

Table 3. Estimated intercepts and scaling coefficients (mode ± 95% CI) for curve parameters of

antibacterial capacity against *E. coli, S. enterica, M. luteus* from phylogenetic univariate mixed

effects models. Estimates in **bold** have 95% CI intervals that do not overlap 0.

	Fixed effect: $\log_{10}$ body mass					
	Intercept			Slope		
Antibacterial capacity curve parameter	**Beta estimate**	**Effective sample**	**pMCMC**	**Beta estimate**	**Effective sample**	**pMCMC**
E. coli						
Inflection$_{\text{antibacterial capacity}}$*	**-0.876** **(-1.331 : -0.521)**	**1000**	**<0.001**	-0.033 (-0.144 : 0.044)	1000	0.278
Slope	**42.4** **(26.8 : 57.9)**	**1000**	**<0.001**	0.984 (-3.257 : 3.89)	1000	0.828
Top asymptote	**0.266** **(0.199 : 0.343)**	**1102**	**<0.001**	0.008 (-0.007 : 0.026)	1000	0.26
Inflection$_{\text{dilution}}$	**0.179** **(0.107 : 0.253)**	**1000**	**<0.001**	0.006 (-0.011 : 0.021)	1124	0.514
Bottom asymptote	**0.062** **(0.009 : 0.113)**	**1000**	**0.044**	-0.004 (-0.018 : 0.007)	1093	0.362
Asymmetric coefficient	**16.5** **(6.7 : 25.1)**	**1000**	**<0.001**	-0.675 (-2.541 : 1.495)	1000	0.62
S. enterica						
Inflection$_{\text{antibacterial capacity}}$	**-0.905** **(-1.296 : -0.557)**	**886.9**	**<0.001**	-0.027 (-0.102 : 0.061)	887.4	0.754
Slope	**34.6** **(15.4 : 49.8)**	**901.8**	**<0.001**	-0.376 (-4.42 : 3.479)	898.3	0.648

Top asymptote	**0.311** **(0.205 : 0.41)**	**1000**	**<0.001**	0.008 (-0.019 : 0.028)	1000	0.72
Inflection$_{dilution}$	**0.218** **(0.134 : 0.324)**	**1000**	**<0.001**	-0.004 (-0.019 : 0.022)	1000	0.982
Bottom asymptote	0.02 (-0.047 : 0.085)	1225.3	0.51	-0.007 (-0.019 : 0.01)	1220.6	0.502
Asymmetric coefficient	**24.5** **(13.1 : 34.3)**	**1000**	**<0.001**	**-3.778** **(-5.604 : -0.907)**	**1000**	**<0.001**

M. luteus

Inflection$_{antibacterial\ capacity}$	-0.727 (-2.029 : 0.565)	1000	0.302	-0.112 (-0.401 : 0.177)	1000	0.412
Slope	**33.2** **(14.8 : 46.8)**	**1000**	**<0.001**	**-4.309** **(-7.652 : -0.788)**	**1000**	**0.018**
Top asymptote	**0.26** **(0.097 : 0.414)**	**1000**	**<0.001**	0.016 (-0.018 : 0.052)	1000	0.306
Inflection$_{dilution}$	**0.178** **(0.062 : 0.327)**	**1000**	**0.004**	0.006 (-0.017 : 0.042)	1000	0.382
Bottom asymptote	0.074 (-0.025 : 0.282)	1000	0.108	-0.029 (-0.067 : 0.004)	1112	0.112
Asymmetric coefficient	4.8 (-13.2 : 17.7)	1097	0.778	1.967 (-1.652 : 4.864)	1117	0.37

712 * Inflection$_{antibacterial\ capacity}$, top asymptote, and bottom asymptote were +1, $\log_{10}$-transformed. Body mass and slope

713 were $\log_{10}$-transformed prior to beta estimation.

Figures

Figure 1. Antibacterial capacity dilution curves support the Safety Factor Hypothesis. Hypothetical antibacterial dilution curve showing the parameters we estimated and biological interpretation (A) This curve shows, names (underlined), and described the curve parameters and the biological interpretation of the curve parameter (in bold). Hypothetical antibacterial curves for an elephant and mouse showing predictions for the Safety Factory Hypothesis (B), the Rate of Metabolism Hypothesis (C), and the Protecton and Complexity Hypotheses (D).

Figure 2. Parameters of antibacterial capacity curves—instantaneous slope at the inflection point (A-C) maximal
antibiotic capacity (D-F), and antibiotic capacity at the inflection point (G-I)—against *Escherichia coli* (A, D, G,),
Salmonella enterica (B, E, H), *Micrococcus luteus* and (C, F, I) against body mass. Each point represents a species
mean, but analyses were performed on individual-level data.