SARS-CoV-2 infection paralyzes cytotoxic and metabolic functions of the immune cells

Yogesh Singh[1,2,6]*,#, Christoph Trautwein[3]*, Rolf Fendel[4]*, Naomi Krickeberg[4], Georgy Berezhnoy[3], Rosi Bissinger[5], Stephan Ossowski[1,2], Madhuri S. Salker[6], Nicolas Casadei[1,2], Olaf Riess[1,2#] and the Deutsche COVID-19 OMICS Initiative (DeCOI)

[1]Institute of Medical Genetics and Applied Genomics, University of Tübingen, Calwerstrasse 7, 72076, Tübingen, Germany
[2]NGS Competence Center Tübingen (NCCT), University of Tübingen, Calwerstrasse 7, 72076 Tübingen, Germany
[3]Werner Siemens Imaging Center, University of Tübingen, Röntgenweg 13, 72076, Tübingen, Germany
[4]Institute of Tropical Medicine, University Hospital of Tübingen, Tübingen, Wilhelmstrasse 27, 72076, Tübingen, Germany
[5]Department of Internal Medicine, Division of Endocrinology, Diabetology and Nephrology, University Hospital of Tübingen, Germany
[6]Research Institute of Women's Health, University of Tübingen, Calwerstrasse 7/6, 72076, Tübingen, Germany
*Equal contributions
Correspondence
Dr Yogesh Singh or Prof Olaf Riess
Institute of Medical Genetics and Applied Genomics, Tübingen University
Calwerstraße 7, 72076, Tübingen, Germany
Phone: 0049 7071 29 72257/78264 Fax: 0049 7071 29 25355
Email: yogesh.singh@med.uni-tuebingen.de or olaf.riess@med.uni.tuebingen.de

Abstract:

The SARS-CoV-2 virus is the causative agent of the global COVID-19 infectious disease outbreak, which can lead to acute respiratory distress syndrome (ARDS). However, it is still unclear how the virus interferes with immune cell and metabolic functions in the human body. In this study, we investigated the immune response in acute or convalescent COVID19 patients. We characterized the peripheral blood mononuclear cells (PBMCs) using flow cytometry and found that CD8+ T cells were significantly subsided in moderate COVID-19 and convalescent patients. Furthermore, characterization of CD8+ T cells suggested that patients with a mild and moderate course of the COVID-19 disease and convalescent patients have significantly diminished expression of both perforin and granzyme A in CD8+ T cells. Using ^{1}H-NMR spectroscopy, we characterized the metabolic status of their autologous PBMCs. We found that fructose, lactate and taurine levels were elevated in infected (mild and moderate) patients compared with control and convalescent patients. Glucose, glutamate, formate and acetate levels were attenuated in COVID-19 (mild and moderate) patients. In summary, our report suggests that SARS-CoV-2 infection leads to disrupted CD8+ T cytotoxic functions and changes the overall metabolic functions of immune cells.

Key words: COVID-19, CD8+ T cells, Granzyme A, Perforin, Metabolites, ^{1}H-NMR, Flow cytometry

Short title: Defective immune-metabolic functions in COVID-19 patient

51
Introduction:
The first cases of severe acute respiratory coronavirus-2 (SARS-CoV-2) infection appeared in
December 2019, in Wuhan, China[1]. This zoonotic virus has infected by now more than 127.8
million people (30.03.2021) and has resulted more than 2.78 million[2,3] death worldwide. The
containment of the pandemic is challenging and is still growing with roughly 200,000 or more
new infections being reported daily since July 2020[2,3]. There is an urgent need for a better
understanding of immunopathology, as SARS-CoV-2 has become the leading cause of
morbidity (long COVID syndrome) and mortality in many countries.

Coronaviruses (CoV) are a large family of viruses that can cause illnesses such as the
common cold and seasonal influenza[4]. Pathologically, SARS-CoV-2 typically infects via
angiotensin-converting enzyme 2 (ACE2)-expressing nasal epithelial cells in the upper
respiratory tract and type II alveolar epithelial cells in patients exhibiting pneumonitis[1,5]. The
most severe disease courses led frequently to death but, not exclusively in older patients with
and without risk conditions. The primary symptoms of SARS-CoV-2 infections are fatigue,
fever, sore throat, dry cough, loss of smell and taste within 5-21 days of incubation of the
virus[6-9]. COVID-19 symptoms are heterogeneous and range from asymptomatic to mild,
moderate, and severe pathological symptoms, presenting with or without pneumonia[10,11].
However, most infected people develop mild to moderate illness and recover without
hospitalization[12,13]. High serum levels of IL-6, IL-8, IL-10, TNF-α cytokines and an immune
hyper-responsiveness referred to as a 'cytokine storm' is connected with poor clinical
outcome[14,15]. Predominantly, older COVID-19 patients can develop acute severe respiratory
distress syndrome (ARDS) due to a cytokine storm which is a life-threatening situation,
requiring ventilation and intensive care support[16-20].

Several break-through discoveries have extended our understanding as to how the virus takes
advantage of the host and modulates immunity[12,19,21-25]. Recovered COVID-19 patients have
an increased number of antibody-secreting cells and activated CD4[+] and CD8[+] T cells. Further,
Immunoglobulin M (IgM) and SARS-CoV-2 reactive IgG antibodies were also detected in blood
before full symptomatic recovery[26-28]. Most severely affected COVID-19 patients had a lower
T cell but elevated B cell counts when compared with healthy controls[13,14,29,30]. Interestingly,
patients with mild symptoms were also shown to have increased T and B cells compared with
severely affected patients[26,29-31]. There could be several reasons for different disease
outcomes including an over-activated innate or hyper-activated adaptive immune response
leading to cytokine storms and resulting in severe injury to the lungs[10,13,25,32]. Despite several
ongoing efforts, the immunological mechanisms of the host-pathogen interaction are not well
understood[33].

There is an intricate balance between the metabolic state of immune cells and generation of
a robust immune response[19,34-37]. CD8[+] T cells require energy to proliferate and accomplish
their effective functions[38]. Most propagating cells such as lymphocytes utilize the most
abundant energy substrates including, glucose, lipids, and amino acids[39]. In response to
SARS-CoV-2 and other virus infections, CD8[+] T cells play a pivotal role in profound growth
and proliferation to generate their effective functional cells which can produce copious
amounts of effector molecules such as cytokines and cytotoxic granules[30,38-40]. An activated
immune system is coupled with a change in metabolic reprogramming to produce enough
energy needed during (viral) infection[38,39]. Proliferating T cells ferment glucose to lactate even

in the presence of oxygen to meet high energy demands[34,37-39]. Furthermore, glucose and
glutamine are involved in the hexosamine biosynthetic pathway, which regulates the
production of uridine diphosphate N-acetyl glucosamine necessary for T cell clonal expansion
and function[41]. The synthesis of lactate intracellularly is crucial for T cells to have an increased
glycolytic flux[38].

Peripheral blood mononuclear cells (PBMCs) can be analyzed to measure the physiological
dysfunctionality of an individual and can serve as a biomarker[42]. Consequently, the metabolic
status of lymphocytes could help to predict disease severity or to select the optimal therapeutic
intervention to boost the immune function during infection. Generally, most of the metabolism-
related functions in PBMCs during SARS-CoV-2 infections were inferred based on
transcriptomics analysis[34,43] and no functional data (biochemical level) have been presented.
Therefore, understanding the kinetics of the adaptive immune response as well as the
metabolic functions during SARS-CoV-2 infection will help to elucidate the host immune
response to SARS-CoV-2 infection. In this study, using flow cytometry and proton nuclear
magnetic resonance (^{1}H-NMR) spectroscopy, we characterized the PBMCs from SARS-CoV-
2 infected and convalescent patients for their immunophenotypic and metabolic functions.

Results:

Characteristics of study participants

PBMCs were isolated and cryopreserved from blood samples obtained from COVID-19
patients suffering from mild ('Mild (outpatient)') or moderate/severe ('Moderate (inpatient)')
disease or were already recovered ('Convalescent') and from healthy controls ('HC').
Classification of disease severity for this analysis was based on the requirement of
hospitalization. Patients with mild COVID-19 were recruited within three days after
confirmation of infection by RT-qPCR. From moderate to severe COVID-19 patient blood
samples were collected one week after their hospital admittance. The moderate patients were
admitted to the hospital requiring medical care; however, they did not need ventilation or O_2
supply. Recovered patients were included based on a positive SARS-CoV-2 antibody testing.
Study participant characteristics are described in Table 1.

Table 1: Patient demographics

No	COVID-19 status	Blood sampling	COVID-19 severity	Sex	Age
1	Outpatient (mild)	Day1	mild	F	21
2	Outpatient (mild)	Day1	mild	M	59
3	Outpatient (mild)	Day1	mild	F	40
4	Inpatient (moderate)	Day7	Moderate	M	57
5	Inpatient (moderate)	Day7	Moderate	M	47
6	Inpatient (moderate)	Day7	Moderate	F	78
7	Convalescent (Sero +ve)	Convalescent	Recovered, healthy	F	50
8	Convalescent	Convalescent	Recovered, healthy	F	24

	(Sero +ve)				
9	Convalescent (Sero +ve)	Convalescent	Recovered, healthy	M	50
10	Convalescent (Sero +ve)	Convalescent	Recovered, healthy	F	51
11	HC1	-	None	F	36
12	HC2	-	None	M	60
13	HC3	-	None	M	40
14	HC4	-	None	M	37
15	HC5	-	None	M	47

Immunophenotyping of COVID-19 mild, moderate and convalescent COVID-19 patients

To compare the number of lymphocytes and monocytes amongst the four study groups, PBMCs were stained and analysed by flow cytometry. Based on live-cell percentage count (gating strategy; Suppl. Fig. 1a), both, lymphocytes and monocytes were not significantly different among mild, moderate and convalescent COVID-19 patients compared with HC (Suppl. Fig. 1b-d).

Increased inflammatory monocytes and reduced NK cells in moderate COVID-19 patients

Monocytes were further classified into classical, non-classical and intermediate based on the expression of CD16 and/or CD14 and were gated as described earlier[44] (Suppl. Fig. 1a). We found that $CD16^+CD14^-$ patrolling (non-classical) monocytes were significantly increased (p=0.0004) in percentage in moderate patients compared to HC, whereas this number is decreased again significantly compared with convalescent patients (p=0.001) (Fig. 1a). The percentage of $CD16^+CD14^-$ (non-classical) monocytes was also significantly increased (p=0.006) in moderate patients compared with mild patients (Fig. 1a Panel I; left). Interestingly, $CD16^+CD14^+$ pro-inflammatory monocytes (intermediate) were again significantly increased in mild (p=0.03) and moderate (p=0.002) compared with HC as well as between moderate and convalescent (p=0.005) (Fig. 1a panel II; middle). Furthermore, we observed a significantly reduced percentage of $CD14^+CD16^-$ phagocytic monocytes (classical) in moderate compared with mild (p=0.004), HC (p<0.0002) and convalescent (p=0.0007) patients (Fig. 1a Panel III; right). Based on the results above using classical 2D flow cytometry analysis, we observed a major difference between HC *vs* mild or with moderate COVID-19 patient samples, therefore, we concatenated three groups together (equal number of cells from each sample; HC, mild and moderate) to identify the visual clustering of monocytes using Uniform Manifold Approximation and Projection (UMAP) for dimension reduction algorithm[45] (Suppl. Fig. 2). Based on UMAP dimension reduction analysis (data-driven and gated cells), we observed a clear difference in the different subsets of monocytes between all three groups (Suppl. Fig. 2c, d & Fig. 1b). We observed that moderate COVID-19 patients mostly cluster in non-classical and partially in intermediate monocyte regions compared to HC, whilst mild COVID-19 patients cluster in all non-classical, intermediate and classical monocytes regions (Fig. 1b; right). We then explored the lymphoid cells compartment for NK cells ($CD56^+CD8^+CD3^-CD19^-$). We found that moderate patients were significantly different from HC (p=0.04). There was also a

tendency of a decrease in NK cells in mild patients compared with HC (p=0.07), although, not
reaching statistical significance (Fig. 1c).

Dynamics of B and T cells in mild, moderate and convalescent patients

Both T and B cells are indispensable for the immune response against viral infections including
SARS-CoV-2. Firstly, we compared the number of B cells amongst the study groups, which
give rise to virus-specific antibodies (see the gating strategy in Suppl. Fig. 1a). The CD19$^+$CD3$^-$
cells (B cells) were significantly increased in mild (p=0.008; fold change x1.7) and moderate
(p=0.0008; fold change x1.9) patients compared with HC (Fig. 2a). Whilst, B cells were
significantly decreased in moderate compared to convalescent (p=0.04) patients (Fig. 2a).
Comparing CD3$^+$CD19$^-$ lymphocytes among the different patient groups we observed no
significant difference. Moreover, CD3$^+$ cells were analysed for the CD4$^+$ and CD8$^+$ T cell
compartment. There was a tendency of increased CD4$^+$ T cells for mild, moderate, and
convalescent patients compared to HC, but no significant difference was observed among any
of the groups. CD8$^+$ T cells were significantly different between HC compared to moderate
(p=0.04) or convalescent (p=0.04) patients (Fig. 2b). Finally, we characterized
CD4$^+$FOXP3$^+$CD45R$^-$ regulatory T cells (Tregs), however, no significant difference was
observed among the different groups (Suppl. Fig. 3).

Impaired activation and defective cytotoxic functions of CD8$^+$ T cells

We found that the percentage of CD8$^+$ T cells were decreased in mild and convalescent
patients compared to HC. Thus, we explored the activation status of CD8$^+$ T cells based on
HLA-DR expression. We found that CD8$^+$ T cell activation status in all three groups of infected
patients was significantly different from HC (mild p=0.01, moderate p=0.01, and convalescent
p=0.01, Fig. 3a). We characterized the cytotoxic potential of CD8$^+$ T cells based on granzyme
A and perforin levels and found that there was a tendency of decreased granzyme A
expression in mild, moderate and convalescent patients compared with HC (Fig. 3b), however,
it did not reach significance. Perforin was significantly decreased in convalescent (p=0.02)
patients compared with HC (Fig. 3b), although mild patients also had reduced levels (p=0.07),
it did not reach statistical significance. Furthermore, we studied the expression of CD38, a
marker of immune cell activation, which was significantly upregulated in convalescent patients
compared with HC (p=0.04) (Fig. 4a). Similarly, convalescent patients had significantly
increased numbers of PD-1$^+$ CD38$^+$ cytotoxic CD8$^+$ T cells compared with HC (p=0.006),
moderate (p=0.005) and mild (p=0.002), which reflects the exhaustion and non-
responsiveness (anergy) of CD8$^+$ T cells (Fig. 4b). Overall, our data suggested that CD8$^+$ T
cells have reduced activation, diminished expression of cytotoxic molecules such as perforin
and granzyme A and have a severely exhausted phenotype.

Dynamics of metabolites production in the mild, moderate and convalescent patient

To establish a putative link between the metabolic state of immune cells and the impaired
immune response, PBMCs from all patient groups were subjected to ^{1}H-NMR spectroscopy
analysis. We identified and quantified a total of 18 metabolites (Fig. 5a). Hereby, unsupervised
Principal Component Analysis (PCA) showed that spectral data from mild and moderate
patients formed overlapping clusters. However, HC and convalescent patients clustered
together (Fig. 5b), indicating a strong difference in metabolite levels between an infectious

state compared to healthy or recovered groups. Statistical analysis of the four different groups revealed that 15 metabolites showed p-values < 0.05, with the highest significance for metabolites related to energy metabolism (Fig. 5c, Suppl. Fig. 3 & Table 2). The data indicate that during infection, there is a strong consumption (reflecting reduced levels) of glucose, acetate and formate, whilst lactate levels are increased. Furthermore, we also found very high levels of fructose in PBMCs from mild patients, medium concentrations in moderate and, low levels in HC and convalescent patients (Fig. 5c). Furthermore, glutamate was almost abolished in mild and moderate patients, potentially as a consequence of enhanced production of α-ketoglutarate in the TCA cycle in PBMCs *via* glutamate dehydrogenase (Fig. 5c). Levels of other amino acids such as glycine and isoleucine were low in mild and moderate patients compared with HC, while creatine and alanine were high in mild patient and moderate patient respectively (Fig. 5c).

Table 2: Summary of metabolites dysregulated in PBMCs

No	Metabolites	HC	Mild	Moderate	Convalescent
1	Glucose	↑↑	↓↓	↓↓	↑↑
2	Formate	↑↑	↓↓↓	↓↓	↑↑
3	Acetate	↑	↓↓	↓	↑
4	Lactate	↓↓	↑↑	↑↑	↓↓
5	Fructose	↓	↑↑	↑	↓
6	Glutamate	↑	↓	↓	↑
7	Citrate	↓	↑↑	↑	↓
8	Taurine	↓↓	-	↑↑	↓
9	Creatine	↓	↑	-	-
10	Alanine	↓	-	↑↑	↓
11	Glycine	↑	↓	↓	-
12	Isoleucine	-	↓↓	↓	↑

To find an association between different metabolites, we applied the variable importance of projection (VIP) score. We found that formate and glucose had the highest score compared to other metabolites (Fig. 6A). To determine if additional metabolites are positively associated with changes in glucose, lactate and fructose, we performed a pattern hunter analysis for all metabolites. We found that high glucose levels correlated with high formate, acetate and glutamate and low lactate and fructose (Fig. 6b), indicating enhanced glycolysis and TCA cycle in PBMCs. Similarly, fructose, which is entered *via* fructose-1-phosphate and dihydroxyacetone phosphate (DAP) into the glycolysis, is correlated positively with lactate and citrate and a decrease in acetate and formate, respectively (Fig. 6b). Interestingly, levels of the ROS scavenger taurine are only positively correlated with lactate and fructose, but not glucose (Fig. 6).

Discussion:

SARS-CoV-2 infections are an intense and rapidly evolving area of research due to the ongoing global pandemic[14,25]. In this study, we used flow cytometry and [1]H-NMR to decipher the cell proportions and functional state of immune cells (PBMCs) in mild, moderate and convalescent COVID-19 patients compared to HC. Recent reports from COVID-19 patients

suggested that mild and severe patient had lymphopenia[11,46-48]. Here, we found that the
myeloid cell compartment in PBMCs based on CD16 and CD14 markers suggested that the
percentage of non-classical and intermediate monocytes were increased during an active mild
or moderate SARS-CoV-2 infection, once infections are cleared the monocyte percentage
numbers return to normal. These results are in accordance with some of the recently published
studies[49-51].

In our cohort, specifically CD56$^+$CD8$^+$NK cells were decreased during active SARS-CoV-2
viral infections (moderate), while during recovery the numbers were comparable to HC as
reported by others[52]. Similarly, another recent study supports our finding by showing the
decrease in the number of NK cell subsets in COVID-19 patients, with no change in CD56^{bright}
or CD56^{dim} cells[53]. Furthermore, based on single-cell RNA-sequencing data, a reduced
number of NK cells was reported in the recovered COVID-19 patients[48]. Thus, altogether, it
appears that CD56$^+$NK cells and their subsets could have an important function during the
ongoing and clearance of SARS-CoV-2 infections. However, further validation studies are
warranted using different *in vivo* model systems with appropriate control groups to pinpoint
the exact role of different NK cell subsets in SARS-CoV-2 infection. CD19$^+$ B lymphocytes
were increased during infection and remain slightly higher than HC, thus reflecting the
antibody response against the SARS-CoV-2 virus. Thus, our data suggest that these patients
were able to generate SARS-CoV-2 specific B cells, this needs further scientific validation.

A major difference was found in the T lymphocytes compartment. CD8$^+$ T cells were
significantly decreased in moderate and convalescent patients as reported earlier[52]. Thus, it
appears that during viral infection non-virus specific CD8$^+$ T cells undergo apoptosis, whilst
the viral-specific surviving CD8$^+$ T cells are clonally expanded but appeared to have lost their
effector functions[54]. To confirm this, we first measured the activation status of CD8$^+$ T cells
and found that CD8$^+$ T cells appeared to be less activated based on their HLA-DR activation
marker[26]. Further, CD8$^+$ T cells were examined for additional activation marker CD38, which
is involved in cell adhesion, signal transduction and calcium signalling[55], was found to be
upregulated in convalescent patients but not during active infection. These CD38$^+$CD8$^+$ T cells
also expressed higher levels of PD-1, which is an immune checkpoint and marker of
exhaustion[24,30,56-58]. PD-1 further guards against autoimmunity promote apoptosis of antigen-
specific T cells and promotes self-tolerance by suppressing T cell inflammatory activity[59].
Thus, viral infection leaves convalescent patients with exhausted phenotypes. We uncovered
that although there was not a significant change in the numbers of Tregs in COVID-19 patients,
there was a trend towards elevated levels of Tregs in COVID-19 patients and rescued Tregs
in convalescent patients, in agreement with the previous studies[56].

A key finding of our study was the surprising observation that granzyme A and perforin
secreting CD8$^+$ T cells were significantly reduced in convalescent patients. These results are
in agreement with the recent single-cell transcriptomics analysis which suggests that
granzyme B and perforin-1 transcripts were also decreased in convalescent COVID-19
patients compared with moderate or severe illness[60]. In contrast to our results, in young
patients, granzyme A or B and perforin levels were increased in mild and moderate cases.
Conversely, in elderly COVID-19 patients, there was a reduced expression of granzyme A and
perforin[61]. Another study suggested that decreased perforin and granzyme A levels in CD4$^+$ T
cells, CD8$^+$ T cell and NK cells is associated with severely afflicted COVID-19 patients[62].
Furthermore, single-cell transcriptomics analysis of SARS-CoV-2 reactive CD8$^+$ T cells in

exhausted and non-exhausted subsets were analyzed[63]. These exhausted CD8[+] T cells were
increased in frequency and displayed lesser cytotoxicity and inflammation features in mild
COVID-19 patients compared to severe patients[63]. A genetic study performed on two cases
highlights the importance of the perforin gene variant A91V which results in the rapid demise
in young COVID-19 patients[64]. The possible implication of our finding is that convalescent
patients, specifically including cancer patients under treatment, could be susceptible to future
opportunistic infections with other viruses including different variants of SARS-CoV-2.
However, further, a large cohort study is warranted to understand the potential functions of
these molecules (granzyme or perforin) in protection or susceptibility against the COVID-19
infection at gene and function levels.
309
To date, the general metabolic physiology of PBMCs is not well defined in the literature.
However, it is clear now that PBMCs are dependent on circulating nutrients and hormones in
the blood[65]. A defective immune response in COVID-19 patients prompted us to investigate
the metabolic functions of these immune cells. Our metabolomics data indeed shows that
PBMCs from actively infected patients have a distinct metabolic profile from convalescent or
healthy individuals. The most notable difference we observed was for metabolites from the
glycolysis and oxidative phosphorylation (TCA cycle) pathway, which is per recently published
transcriptome data for PBMCs[39,43]. Metabolites such as glucose, formate, acetate and choline
were also reduced in PBMCs in infected patients whereas, HC and convalescent patients had
a normal profile. Accordingly, the glycolytic pathway end products such as lactate were higher
in active mild and moderate COVID-19 patients compared with HC and convalescent
individuals. Therefore, our data suggest that PBMCs (which constitute a major fraction of T
lymphoid cells: 70- 80%) may have modulated their metabolic functions, particularly favouring
the oxidative phosphorylation pathway over the glycolytic pathway, to meet the high demands
of energy needed to combat the ongoing viral infection.
325
A recent report suggested that elevated glucose levels enhance SARS-CoV-2 replication and
cytokine expression in monocytes and glycolysis sustains the viral-induced monocyte
response[66]. Recently, it was emphasized that glucose consumption in PBMCs during COVID-
19 disease could be also a read-out of cytokine storms[34]. Further, a higher abundance of
citrate in PBMCs suggested that perhaps T cells could use the oxidative phosphorylation
pathway for energy consumption to endure the infection, as recent transcriptomic data also
suggested that higher expression of genes related to oxidative phosphorylation both in
peripheral mononuclear leukocytes and bronchoalveolar lavage fluid (BALF) could play a
crucial role in increased mitochondrial activity during SARS-CoV-2 infection[34].
335
Another remarkable finding of our study was the increase of fructose levels in PBMCs during
the course of infection. Previous findings suggested that fructose is involved in the
inflammatory pathways for the production of IL-1β and IL-6 production[67]. Thus, it is
conceivable that the immune cells (most probably monocytes) could be triggered by higher
fructose and simultaneously induce inflammation and IFN-γ production by T cells[67]. These
findings correlate with recent transcriptomic studies on the BALF from infected COVID-19
patients and plasma of COVID-19 patients that also identified changes in fructose
metabolism[34,68].
344
345 Several amino acids are involved in anti-inflammatory effects, especially arginine, glutamine
346 and glycine appeared to improve lung damage induced by infections[69]. Administration of

glutamine reduced the inflammatory cytokines, whereas arginine or glycine reduced IL-6 and CXCL-1 expression in the alveolar epithelium[70]. Additionally, glutamine is an important signalling molecule involved in activating mammalian target of rapamycin (mTOR) signalling which is critical for immune cell activity and inhibiting catabolic functions such as protein degradation and apoptosis[71]. In our study, we found a reduced abundance of glutamate and glycine in PBMCs during the mild and moderate COVID-19 patients. Reduced glucose and glutamine are involved in the hexosamine biosynthetic pathway and could be responsible for poor T cell clonal expansion and function[41]. The exact molecular mechanism of individual amino acids is a complex phenomenon and currently unclear. However, at least in theory, supplementation of glutamine and glycine could be beneficial for COVID-19 patients.

We finally observed a reduction of granzyme A and perforin levels in CD8[+] T cells and detected the ambient level of antioxidant amino acid taurine in the convalescent patients, which could be involved in the modulation of cytotoxic functions of CD8[+] T cells. Both granzyme A and perforin are involved in ROS production and taurine serves as ROS scavenger[72,73]. Thus, decreased granzyme A and perforin could be implicated in reduced ROS production for the impaired effectiveness of CD8[+] T cells in either convalescent patients or COVID-19 patients. This should be the case, as taurine levels are generally increased during active infection in mild/moderate patients compared to healthy controls and are not specifically decreasing due to granzyme A and perforin lacking ROS activity in COVID-19 patients. However, this finding needs further investigation to validate this hypothesis as it is unclear how ROS and taurine act together to affect the cytotoxic functions of immune CD8[+] T cells. In summary, the metabolomics data generated in this study provides first and crucial insights into the complex metabolic changes of PBMCs during SARS-CoV-2 infections, warranting further future in-depth investigation.

Conclusions:
Using immunophenotyping and metabolomics approaches we detected significant changes in PBMC samples of mildly and moderately affected COVID-19 as well as convalescent patients compared with healthy controls. The reduced percentage of NK cells in both mild and moderate patient groups corresponded with the clustering of PBMCs metabolite levels in the principal component analysis distinct from the cluster formed by healthy and convalescent individuals. The dramatically changed metabolic activity and pathways, such as glycolysis and TCA cycle, might not only lead to a vulnerability of COVID-19 patients to subsequent infections but can also offer insights into how PBMCs could be manipulated towards a better survival and personalized treatment of moderate and severe COVID-19 patients.

Limitation of the study:
The current study is performed from the samples obtained during the first wave of infections and suffers from the limited number of patients used in the study. The use of appropriate controls such as influenza viral infection would have been useful for a more generalized conclusion. Therefore, results obtained in future studies might differ from subsequent waves of infections in new patient cohorts. Further, the novel SARS-CoV-2 variants or vaccine candidates might promote alternative host immune evasion strategies to infect the host, thus, eliciting a different immune response. Nonetheless, this study certainly highlights the importance of distinct cell types as well as the crucial function of cell metabolism during SARS-CoV-2 viral infection and even after recovery.

Materials and Methods:

Ethics statement

The study protocols were approved by the University of Tübingen, Germany Human Research Ethics Committee (TÜCOV: 256/2020BO2 (convalescent study), COMIHY: (225/2020AMG1) (outpatient study)-COMIHY, EUDRA-CT: 2020-001512-26, ClinicalTrials.gov ID: NCT04340544, and COV-HCQ: (190//2020AMG1) (inpatient study)-COV-HCQ, EUDRA-CT: 2020-001224-33, ClinicalTrials.gov ID: NCT04342221, 556/2018BO2) and all associated procedures were carried out in accordance with approval guidelines. All participants provided written informed consent in accordance with the Declaration of Helsinki.

Study participants

SARS-CoV-2 positive patients were used for this study and no other virus species were analysed in this study (COMIHY and COV-HCQ). Blood was collected from COVID-19 patients enrolled into two different prospective randomized, placebo-controlled, double-blind clinical trials evaluating the safety and efficacy of hydroxychloroquine in COVID-19 outpatients (COMIHIY) and hospitalized patients (COV-HCQ). We analysed subsets of these study cohort and used outpatient (n=3; COMIHY) which came to a specified outpatient ward at the Institute of Tropical Medicine with mild symptoms and blood was taken and usually defined as D1 outpatients. Inpatients (n=3; COV-HCQ), blood was taken after 7-9 days after study inclusion defined as D7. These patients had moderate symptoms needing hospital care, however not being transferred to the intensive care unit in the hospital. Furthermore, convalescent COVID-19 patients (n=4) were defined as positive for serum antibody reactive to SARS-CoV-2 and blood was taken when they visited the Institute of Tropical Medicine for testing of antibody levels. Amongst this cohort, 3 persons reported mild fever for 10-11 days and 1 individual reported no fever but found positive for SARS-CoV-2 antibodies. Blood from healthy controls (n=5) was obtained from the hospital blood bank.

Flow cytometry and UMAP data analysis

PBMCs were isolated by the standard Ficoll method[74]. A total of 1-2 x10^6 PBMCs per participants were used for three FACS panels (Table 2). In brief, firstly, to distinguish between live from dead, the cells were incubated with LIVE/DEAD Fixable Infra-Red Dead stain (Thermofisher) for 15 minutes at room temperature (RT) into 1:40 diluted dye in DPBS. Subsequently, after LIVE/DEAD staining, cells were stained with surface markers in DPBS (Sigma) with Super Bright stain Buffer (Thermofisher) for 30 minutes at RT. After surface staining cells were also stained for intracellular (IC) markers. Before IC staining, cells were fixed for 30-45 minutes and permeabilized for 5 minutes followed by IC antibody incubation for additional 30 minutes at RT. Cells were washed and resuspended in DPBS containing 2%FBS. Fixing of cells was performed irrespective of whether the panel was used for IC staining or not to prevent the possible contamination during the acquisition of the samples. All the staining procedures were performed in the dark to avoid the photo-bleaching of dyes. For each sample, 200,000 cells were acquired using BD LSRFortessa (core facility) equipped with 4 lasers (violet, blue and yellow-green and Red). Data were analysed using Flow Jo (Tree Star) and fluorescence minus one controls (FMO) were used for setting up the arbitrary gates

for the major cell markers. Furthermore, UMAP dimensional reduction analysis was performed using UMAP plug-in[45] in Flowjo software using default setting except for minimum distance (0.2 instead of 0.5) and population (n=15) as advised by a plug-in. First, dead/debris was removed by gating (FSC-A vs SSC-A; linear scale) then using FSC-A vs FSC-W, we focussed on singlets. The singlets were again gated for live and dead discrimination. An equal number of live cells from each sample (HC, mild and moderate) were concatenated and exported as a single FCS file. This FSC file was subjected to UMAP analysis, each cell population was either monocytes or lymphocytes were again subject to subsequent UMAP analysis for clustering of specific sub-populations. Based on gating or antibodies-stained cells (data driven) analysis was performed and summarized in Suppl. Fig. 2.

Table 2: Antibodies and other reagents used for Flow cytometry

No.	Product Name	Clone	Fluorochrome	Product ID	Company
		NK cells and Monocytes (Panel 1)			
1	CD3	UCHT1	eFluor 450	48-0038-42	Thermofisher
2	CD4	SK3	SuperBright 600	63-0047-42	Thermofisher
3	CD8a	SK1	PerCP-eFluor 710	46-0087-42	Thermofisher
4	CD19	HIB19	eFluor 506	69-0199-42	Thermofisher
5	CD45-RA	HI100	PE-Cy7	25-0458-42	Thermofisher
6	HLA-DR	L243	Alexa Fluor647	A51010	Thermofisher
7	CD38	HIT2	PE-eFluor610	61-0389-42	Thermofisher
8	CD56	MEM188	PE	MA119638	Thermofisher
9	CD16	3G8	Super Bright702	67-0166-42	Thermofisher
10	CD14	61D3	Alexa Fluor700	56-0149-42	Thermofisher
11	Foxp3 (IC)	PCH101	FITC	11-4776-42	Thermofisher
		CD8 exhaustion, T helper follicular cells (Tfh) and antibody-secreting cell (ASC) (Panel 2)			
1	CD3	UCHT1	eFluor 450	48-0038-42	Thermofisher
2	CD19	HIB19	eFluor 506	69-0199-42	Thermofisher
3	CD4	SK3	Super Bright 600	63-0047-42	Thermofisher
4	CD8a	SK1	PerCP-eFluor 710	46-0087-42	Thermofisher
5	CD38	HIT2	PE-eFluor 610	61-0389-42	Thermofisher
6	CD27	O323	Alexa Fluor700	56-0279-42	Thermofisher
7	CXCR5 (CD185)	MU5UBEE	FITC	11-9185-42	Thermofisher
8	ICOS (CD278)	C398.4A	PE	12-9949-81	Thermofisher
9	PD-1 (CD279)	eBioJ105 (J105)	PE-Cy7	25-2799-42	Thermofisher

10	HLA-DR	L243	Alexa Fluor647	A51010	Thermofisher
		Cytotoxic potential			
1	CD4	SK3	SuperBright600	63-0047-42	Thermofisher
2	CD8	SK1	PerCP-eFluor710	46-0087-42	Thermofisher
3	CD19	HIB19	eFluor 506	69-0199-42	Thermofisher
4	CD38	HIT2	PE-eFluor 610	61-0389-42	Thermofisher
5	HLA-DR	L243	Alexa Fluor 647	A51010	Thermofisher
6	GZMA (IC)	CB9	PE	12-9177-42	Thermofisher
7	GZMB (IC)	GB11	Alexa Fluor 488	MA5-23639	Thermofisher
8	Perforin (IC)	dG9	PE-Cy7	12-9177-42	Thermofisher
		Other Flow reagents			
1	Ultracompensation bead			01-2222-42	Thermofisher
2	Live/Dead Fixable Near IR staining kit			L10119	Thermofisher
2	FOXP3/TRN FACTOR STAIN BUFFER SET			00-5523-00	Thermofisher
3	FLOW STAIN BUFFER SOLN			00-4222-57	Thermofisher
4	SB COMPLETE STAINING BUFFER			SB-4401-42	Thermofisher
5	DPBS			D8537	Sigma
6	Pancoll human			P04-601000	Pan Biotech

[1]H-NMR metabolomics

To obtain PBMCs metabolites, PBMCs were suspended in an optimized solvent extraction mixture of 9:1 (methanol: chloroform) as described elsewhere in detail[75] and extracted with a focused ultrasound system (Covaris E220, Woburn, USA). The extraction solutions were evaporated to dryness for 4 hours in a vacuum concentrator and afterwards pellets resuspended with 45 µL in a 1 mM TSP containing deuterated phosphate buffer. After centrifugation at 20,000 x g for 10 min to remove residual macromolecules, 40 µL of the clear supernatant were transferred to 1.7 mm NMR tubes. Spectra were recorded on ultrashielded 600 MHz spectrometer (Bruker AVANCE III HD, Karlsruhe, Germany) with a triple resonance 1.7 mm room temperature probe. Spectra used for analysis were acquired with a 2h 55min lasting CPMG pulse program. Metabolite annotation and quantification were done with ChenomX NMR Suite 8.3.

Statistical analysis

Bar diagrams were prepared using GraphPad Prism 6.0. Data shown are means ± SD. FACS
data were analysed using one-way analysis of variance (ANOVA) for multiple group
comparisons (mild, moderate, convalescent and HC) in GraphPad Prism software. No
matching or pairing was used. Assumed Gaussian distribution with equal standard deviations
(SDs) for experimental design. The mean of each group was compared with the mean of every
other group and Tukey's post-hoc tests for multiple comparisons were employed. P-value
considered significantly less than 0.05 or equal. Metabolite concentrations from [1]H-NMR
analysis were exported as comma-separated value spreadsheet file to MetaboAnalyst 4.0
software, normalized with probabilistic quantile normalization (PQN) and range scaled.
Unsupervised principal component analysis (PCA), a dimensionality-reduction method was
used for clustering of all metabolites (PC1 and PC2 only). Multiple groups comparison was
performed using one way ANOVA. Multivariate data analysis techniques such as partial least
squares analysis (PLSDA) and variable's importance in the PLSDA (VIP) score analysis were
used to find the correlation among different metabolites and various groups. Hierarchical
clustering of all the metabolites was measured and distance measurement was performed
with Pearson r correlation coefficient.

Figure legends:

Fig. 1: Comparison of monocytes and NK cell percentage amongst study groups.

A. The stained PBMCs were gated on the monocyte population and $CD3^+CD19^+$ cells
were excluded. Cell populations are displayed for CD16 and CD14 expression (upper
FACS panel). One exemplary dot plot is shown per study group. The bar diagrams
(lower panel) show the non-classical ($CD16^+CD14^-$), intermediate ($CD16^+CD14^+$) and
classical ($CD16^-CD14^+$) monocytes. *P-value ≤0.05, **P-value ≤0.01 and ***P-value
≤0.001.
B. UMAP plot for different subsets of monocytes (non-classical, intermediate and
classical monocytes; left UMAP plot) from all the samples. Group comparisons among
HC, mild and moderate (right UMAP plot). Colour coded information is provided for
either different subsets of monocyte populations or patient groups.
C. The stained PBMCs were gated on lymphocyte population and further excluded the
$CD3^+CD19^+$ cells and examined for the CD56 and CD8 expression in HC, mild,
moderate and convalescent (upper FACS panel). One exemplary dot plot is shown per
study group. The bar diagram shows the $CD56^+CD8^+CD3^-CD19^-$ NK cells. *P-value
≤0.05.

Fig. 2: Increased B cells in mild and moderate patients and reduced $CD8^+$ cytotoxic T cells in
mild and convalescent patients

A. The stained PBMCs were gated on lymphocyte population and examined for the CD19
and CD3 expression in HC, mild, moderate and convalescent (upper FACS panel).
One exemplary dot plot is shown per study group. The bar diagram shows $CD3^-CD19^+$
B cells. *P-value ≤0.05, **P-value ≤0.01 and ***P-value ≤0.001.
B. The $CD19^-CD3^+$ lymphocytes were examined for $CD4^+$ and $CD8^+$ T marker expression.
One exemplary dot plot is shown per study group. There was statistically significant
difference among HC, mild, moderate and convalescent (upper FACS panel).

However, CD8$^+$ T cells were significantly reduced in outpatient and convalescent
patients. *P-value ≤0.05.
523
524 **Fig. 3:** Decreased activation and cytotoxic functional protein expression of CD8$^+$ T cells in
525 convalescent patients
526
A. CD8$^+$ T cells were examined for the expression of activation marker HLA-DR (upper
FACS panel). One exemplary dot plot is shown per study group. The bar diagram
(lower panel) shows that HLA-DR was a significantly lower on CD8$^+$ T cells in mild,
moderate and convalescent COVID-19$^+$ patients compared with HC. *P-value ≤0.05.
B. CD8$^+$ T cells were examined for the expression of their cytotoxic potential using
granzyme A and perforin expression using IC staining (upper FACS panel). One
exemplary dot plot is shown per study group. There was a statistically significant
difference among HC, mild, moderate and convalescent (upper FACS panel) for
granzyme A. The bar diagram (lower panel) shows that perforin expression was
significantly lower on CD8$^+$ T cells in convalescent COVID-19$^+$ patients compared with
HC, though mild and moderate represent a lower expression of perforin, but it did not
to a significant level. *P-value ≤0.05.
539
540 **Fig. 4:** Increased exhausted CD8$^+$ T cells in convalescent patients
541
542   A. Expression of activation marker CD38 on CD8$^+$ T cells (upper FACS panel). One
543   exemplary dot plot is shown per study group. The bar diagram (lower panel) shows
544   that CD38 expression was significantly higher on CD8$^+$ T cells in convalescent COVID-
545   19$^+$ patients compared with HC. *P-value ≤0.05.
546
547   B. Expression of activation marker CD38 and PD-1 on CD8$^+$ T cells (upper FACS panel).
548   One exemplary dot plot is shown per study group. The bar diagram (lower panel)
549   shows that PD-1$^+$CD38$^+$ expression on was a significantly higher on CD8$^+$ T cells in
550   convalescent COVID-19$^+$ patients compared with HC. **P-value ≤0.01.
551
Fig. 5: ^{1}H-NMR spectroscopy of PBMC extracts
A. Heatmap of featured metabolites' concentrations plotted with SARS-CoV-2
progression group clustering.
B. Principle component analysis (PCA) was performed to identify the clustering of two
different groups. HC and convalescent COVID-19 patient samples cluster together
while SARS-Co-2 infected mild and moderate patients cluster in a separate cluster with
PC1: 90.7% and PC2: 2.6%.
C. Box plots for differentially abundantly present metabolites in the different group
including HC, mild, moderate, and convalescent COVID-19 patient. *P-value ≤0.05,
**P-value ≤0.01 and ***P-value ≤0.001.
562
Fig. 6: Pattern hunter plots provide an insight of close correlations with other metabolites
during COVID-19 infection.
A. Variable Importance in Projection (VIP) scores for all metabolites in the four studied
groups.
B. Pattern hunter plot for glucose.
C. Pattern hunter plot for lactate and fructose.

Suppl. Fig. 1: Total % counts of monocytes and lymphocytes from PBMCs of COVID-19 patients.

    A. Fixed PBMCs samples (1×10^6 cells) were acquired on flow cytometry on 2-3 different days for the entire experiments. Total 200,000 cells were acquired by flow cytometry and gating was performed based on FSC and SSC parameters for lymphocytes, monocytes and dead cells as described earlier[76-78]. Gating strategy for T lymphocytes (CD3, CD4 and CD8) monocytes (CD14 and CD16)[44], NK cells (CD56) using FMO controls.

    B. The bar graphs represent the % of lymphocytes and monocytes.

Suppl. Fig. 2: UMAP analysis of lymphocytes and monocytes cell populations.

    A. UMAP pseudo-colour plot was derived from all the samples. All the FCS files (used only live-cell gated population; minimum cells per samples were 25,000) were concatenated from HC, mild and moderate patients (upper panel). Representation of lymphocytes and monocytes populations in UMAP plot (middle panel). Presentation of different lymphocytes subpopulations like $CD4^+$ T cells, $CD8^+$ T cells, B cells, NK cells, NKT cells and regulatory T cells (lower panel).

    B. UMAP plot for overlaying of different groups.

    C. UMAP plot for different subsets of monocytes.

    D. UMAP plots for individual antibody (protein) expression on monocyte populations. Scale bar shows the expression level of protein expression on the cell clusters (blue - lowest and red highest expression; relative expression).

Suppl. Fig. 3: Kinetics of regulatory T cells is not affected significantly in mild, moderate and convalescent patients.

$Foxp3^+$ expression on $CD19^-CD3^+CD4^+CD45RA^-$ T cells to identify the regulatory T cells in HC, outpatient, outpatient and convalescent (upper FACS panel). There was a statistically significant difference among HC, mild, moderate and convalescent (upper FACS panel).

Suppl. Fig. 4: Metabolite analysis in COVID-19 patients.

    A. Analysis of Variance (ANOVA) for multi-group comparisons.

    B. Partial Least Squares Discriminant Analysis (PLS-DA) scores plot.

    C. Hierarchical clustering of metabolites (distance measured with Pearson r correlation coefficient).

    D. Boxplots for branched-chain amino acids valine and leucine.

Author's contributions:

YS: Overall study design and project coordination, flow cytometry experiments, data analysis and interpretation and initial metabolic sample preparation, funding generation, and writing the manuscript.

CT: Metabolites sample preparation, Processing of the sample on [1]H-NMR, data analysis, data interpretation and writing

MO: Metabolites sample preparation, Processing of the sample on [1]H-NMR

RF, NK: Provided patient materials, Isolation of PBMCs, Performed the experiments for flowcytometry in BSL-2 facility, flow data interpretations.

RB: Isolation of PBMCs from HC and sample preparation for flow cytometry.

SO, MS, NC, OR: Study design, providing research tools, funding generation, data interpretation and writing and amending the manuscript.

All the authors have seen the manuscript, substantially contributed and agreed to be co-author.

Acknowledgements:

We thank you all the patients who participated in this study and FACS core facility (Klinikum-Berg) for accessing the Flow cytometry for the experiments. We also thanks Jana Held and Andrea Kreidenweiss from Institute of Tropical Medicine, Tübingen university for their help with coordination with clinical Trials and proof-reading of the manuscript. We acknowledge support by Deutsche Forschungsgemeinschaft (DFG) and Open Access Publishing Fund of University of Tübingen.

Funding:

RB is supported by Deutsche Forschungsgemeinschaft (DFG Project no. 426724658). EKFS (MS), DFG (OR), Ferring Pharmaceutical (YS and MSS). To MSS the Margarete von Wrangell (MvW 31-7635.41/118/3) habilitation scholarship co-funded by the Ministry of Science, Research and the arts (MWK) of the state of Baden-Württemberg and by the European Social Funds. Funders have no role in the study design and data analysis. To RF the German Federal Ministry of Education and Research (BMBF) (BMBF-01KI2052) and the German Federal Ministry of Health (BMG) (BMG-ZMVI1-1520COR801).

Competing interests:

Authors declare no financial competing interests.

DECOI Consortia
The members of the Deutsche COVID-19 Omics Initiative (DeCOI) are Janine Altmüller, Angel Angelov, Robert Bals, Alexander Bartholomäus, Anke Becker, Daniela Bezdan, Ezio Bonifacio, Peer Bork, Nicolas Casadei, Thomas Clavel, Maria Colome-Tatche, Andreas Diefenbach, Alexander Dilthey, Nicole Fischer, Konrad Förstner, Sören Franzenburg, Julia-Stefanie Frick, Gisela Gabernet, Julien Gagneur, Tina Ganzenmüller, Marie Gauder, Alexander Goesmann, Siri Göpel, Adam Grundhoff, Torsten Hain, Andrè Heimbach, Michael Hummel, Thomas Iftner, Angelika Iftner, Stefan Janssen, Jörn Kalinowski, René Kallies, Birte Kehr, Andreas Keller, Sarah Kim-Hellmuth, Christoph Klein, Oliver Kohlbacher, Karl Köhrer, Michael Knop, Jan O. Korbel, Peter G Kremsner, Denise Kühnert, Ingo Kurth, Markus Landthaler, Yang Li, Kerstin Ludwig, Oliwia Makarewicz, Manja Marz, Alice McHardy, Christian Mertes, Sven Nahsen, Markus Nöthen, Francine Ntoumi, Peter Nürnberg, Uwe Ohler, Stephan Ossowski, Jörg Overmann, Silke Peter, Klaus Pfeffer, Anna R. Poetsch, Alfred Pühler, Nikolaus Rajewsky, Markus Ralser, Olaf Rieß, Stephan Ripke, Ulisses Nunes da Rocha, Philip Rosenstiel, Antoine-Emmanuel Saliba, Leif Erik Sander, Birgit Sawitzki, Philipp Schiffer, Wulf Scheider, Eva-Christina Schulte, Joachim L. Schultze, Alexander Sczyrba, Yogesh Singh, Michael Sonnabend, Oliver Stegle, Jens Stoye, Fabian Theis, Janne Vehreschild, Thirumalaisamy P Velavan, Jörg Vogel, Max von Kleist, Andreas Walker, Jörn

Walter, Dagmar Wieczorek, Sylke Winkler, John Ziebuhr, Simone Scheithauer, Hajo Grundmann, Jonathan Schmid-Burgk, Ulrike Protzer and Inti Alberto De La Rosa Velázquez.

References:

Zhu, N. *et al.* A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med* **382**, 727-733, doi:10.1056/NEJMoa2001017 (2020).

Dashboard, J. C.-L. https://coronavirus.jhu.edu/map.html. (2020).

Dashboard, W. L. https://covid19.who.int. (2020).

Chang, L., Yan, Y. & Wang, L. Coronavirus Disease 2019: Coronaviruses and Blood Safety. *Transfus Med Rev*, doi:10.1016/j.tmrv.2020.02.003 (2020).

Lehner, N. J. M. P. J. How does SARS-CoV-2 cause COVID-19? *Science*, 510-511 (2020).

Liu, J. *et al.* Community Transmission of Severe Acute Respiratory Syndrome Coronavirus 2, Shenzhen, China, 2020. *Emerg Infect Dis* **26**, doi:10.3201/eid2606.200239 (2020).

Singhal, T. A Review of Coronavirus Disease-2019 (COVID-19). *Indian J Pediatr* **87**, 281-286, doi:10.1007/s12098-020-03263-6 (2020).

WHO. <WHO-2019-nCoV-Sci_Brief-Transmission_modes-2020.3-eng.pdf>. *WHO Report* (2020).

Rothe, C. *et al.* Transmission of 2019-nCoV Infection from an Asymptomatic Contact in Germany. *N Engl J Med* **382**, 970-971, doi:10.1056/NEJMc2001468 (2020).

Xu, Z. *et al.* Pathological findings of COVID-19 associated with acute respiratory distress syndrome. *The Lancet Respiratory Medicine* **8**, 420-422, doi:10.1016/s2213-2600(20)30076-x (2020).

Wei Huang, J. B., Michelle McNamara, Suraj Saksena, Marsha Hartman, Tariq Arshad, Scott J. Bornheimerand Maurice O'Gorman. Lymphocyte Subset Counts in COVID-19 Patients: A Meta-Analysis. doi:10.1002/cytoa.24172 (2020).

Chen, G. *et al.* Clinical and immunological features of severe and moderate coronavirus disease 2019. *J Clin Invest* **130**, 2620-2629, doi:10.1172/JCI137244 (2020).

Shi, Y. *et al.* COVID-19 infection: the perspectives on immune responses. *Cell Death Differ*, doi:10.1038/s41418-020-0530-3 (2020).

Daniela Weiskopf1, K. S. S., Matthijs P. Raadsen, Alba Grifoni, Nisreen M.A. Okba, Henrik Endeman, Johannes P.C. van den Akker, Richard Molenkamp, Marion P.G. Koopmans, Eric C.M. van Gorp, Bart L. Haagmans, Rik L. de Swart, Alessandro Sette, Rory D. de Vries. Phenotype and kinetics of SARS-CoV-2-specific T cells in COVID-19 patients with acute respiratory distress syndrome. *Science Immunology*, doi:10.1126/sciimmunol.abd2071 (2020). (2020).

Del Valle, D. M. *et al.* An inflammatory cytokine signature predicts COVID-19 severity and survival. *Nature Medicine*, doi:10.1038/s41591-020-1051-9 (2020).

NcA OANA dOcEA, A. T., DANA ALBULEScU, OANA CRISTEA, & OVIdIU ZLATIAN, M. V., STERGHIOS A. MOScHOS, dIMITRIS TSOUKALAS, MARINA GOUMENOU, NIKOLAOS dRAKOULIS, JOSEF M. dUMANOV, VIcTOR A. TUTELYAN, GENNAdII G. ONISchENKO, MIcHAEL AScHNER, dEMETRIOS A. SPANdIdOS and dANIELA cALINA. A new threat from an old enemy: Re-emergence of coronavirus (Review). *INTERNATIONAL JOURNAL OF MOLEcULAR MEdIcINE* **45**, 1631-1643 (2020).

Ong, E. Z. *et al.* A Dynamic Immune Response Shapes COVID-19 Progression. *Cell Host Microbe* **27**, 879-882 e872, doi:10.1016/j.chom.2020.03.021 (2020).

18   Picchianti Diamanti, A., Rosado, M. M., Pioli, C., Sesti, G. & Lagana, B. Cytokine Release
Syndrome in COVID-19 Patients, A New Scenario for an Old Concern: The Fragile
Balance between Infections and Autoimmunity. *Int J Mol Sci* **21**,
doi:10.3390/ijms21093330 (2020).
19   Vabret, N. *et al.* Immunology of COVID-19: Current State of the Science. *Immunity* **52**,
910-941, doi:10.1016/j.immuni.2020.05.002 (2020).
20   Velavan, T. P. & Meyer, C. G. Mild versus severe COVID-19: Laboratory markers. *Int J*
Infect Dis **95**, 304-307, doi:10.1016/j.ijid.2020.04.061 (2020).
21   Wu, H. *et al.* Clinical and Immune Features of Hospitalized Pediatric Patients With
Coronavirus Disease 2019 (COVID-19) in Wuhan, China. *JAMA Netw Open* **3**,
e2010895, doi:10.1001/jamanetworkopen.2020.10895 (2020).
22   Du, S. Q. & Yuan, W. Mathematical modeling of interaction between innate and
adaptive immune responses in COVID-19 and implications for viral pathogenesis. *J*
Med Virol, doi:10.1002/jmv.25866 (2020).
23   Giamarellos-Bourboulis, E. J. *et al.* Complex Immune Dysregulation in COVID-19
Patients with Severe Respiratory Failure. *Cell Host Microbe* **27**, 992-1000 e1003,
doi:10.1016/j.chom.2020.04.009 (2020).
24   Braciale, T. J. & Hahn, Y. S. Immunity to viruses. *Immunol Rev* **255**, 5-12,
doi:10.1111/imr.12109 (2013).
25   McKechnie, J. L. & Blish, C. A. The Innate Immune System: Fighting on the Front Lines
or Fanning the Flames of COVID-19? *Cell Host Microbe* **27**, 863-869,
doi:10.1016/j.chom.2020.05.009 (2020).
26   Thevarajan, I. *et al.* Breadth of concomitant immune responses prior to patient
recovery: a case report of non-severe COVID-19. *Nature Medicine*,
doi:10.1038/s41591-020-0819-2 (2020).
27   Assis, R. R. d. *et al.* Analysis of SARS-CoV-2 Antibodies in COVID-19 Convalescent
Plasma using a Coronavirus Antigen Microarray
.   *bioRxiv preprint doi:* https://doi.org/10.1101/2020.04.15.043364*.,*
doi:10.1101/2020.04.15.043364 (2020).
28   Cao, Y. *et al.* Potent Neutralizing Antibodies against SARS-CoV-2 Identified by High-
Throughput Single-Cell Sequencing of Convalescent Patients' B Cells. *Cell* **182**, 73-84
e16, doi:10.1016/j.cell.2020.05.025 (2020).
29   Qin, C. *et al.* Dysregulation of immune response in patients with COVID-19 in Wuhan,
China. *Clin Infect Dis*, doi:10.1093/cid/ciaa248 (2020).
30   Song, J. W. *et al.* Immunological and inflammatory profiles in mild and severe cases of
COVID-19. *Nat Commun* **11**, 3410, doi:10.1038/s41467-020-17240-2 (2020).
31   Channappanavar, R., Fett, C., Zhao, J., Meyerholz, D. K. & Perlman, S. Virus-specific
memory CD8 T cells provide substantial protection from lethal severe acute
respiratory syndrome coronavirus infection. *J Virol* **88**, 11034-11044,
doi:10.1128/JVI.01505-14 (2014).
32   Janice Oh, H. L., Ken-En Gan, S., Bertoletti, A. & Tan, Y. J. Understanding the T cell
immune response in SARS coronavirus infection. *Emerg Microbes Infect* **1**, e23,
doi:10.1038/emi.2012.26 (2012).
33   Bost, P. *et al.* Host-Viral Infection Maps Reveal Signatures of Severe COVID-19
Patients. *Cell*, doi:10.1016/j.cell.2020.05.006 (2020).

34   Gardinassi, L. G., Souza, C. O. S., Sales-Campos, H. & Fonseca, S. G. Immune and
Metabolic Signatures of COVID-19 Revealed by Transcriptomics Data Reuse. *Frontiers*
in Immunology **11**, doi:10.3389/fimmu.2020.01636 (2020).

35   Ayres, J. S. Immunometabolism of infections. *Nat Rev Immunol* **20**, 79-80,
doi:10.1038/s41577-019-0266-9 (2020).

36   Thaker, S. K., Ch'ng, J. & Christofk, H. R. Viral hijacking of cellular metabolism. *BMC*
Biol **17**, 59, doi:10.1186/s12915-019-0678-9 (2019).

37   Moreno-Altamirano, M. M. B., Kolstoe, S. E. & Sanchez-Garcia, F. J. Virus Control of
Cell Metabolism for Replication and Evasion of Host Immune Responses. *Front Cell*
Infect Microbiol **9**, 95, doi:10.3389/fcimb.2019.00095 (2019).

38   Gupta, S. S., Wang, J. & Chen, M. Metabolic Reprogramming in CD8+ T Cells During
Acute Viral Infections. *Frontiers in Immunology* **11**, doi:10.3389/fimmu.2020.01013
(2020).

39   Shehata, H. M. *et al.* Sugar or Fat?-Metabolic Requirements for Immunity to Viral
Infections. *Front Immunol* **8**, 1311, doi:10.3389/fimmu.2017.01311 (2017).

40   Pallett, L. J., Schmidt, N. & Schurich, A. T cell metabolism in chronic viral infection. *Clin*
Exp Immunol **197**, 143-152, doi:10.1111/cei.13308 (2019).

41   Swamy, M. *et al.* Glucose and glutamine fuel protein O-GlcNAcylation to control T cell
self-renewal and malignancy. *Nat Immunol* **17**, 712-720, doi:10.1038/ni.3439 (2016).

42   Liew, C. C., Ma, J., Tang, H. C., Zheng, R. & Dempsey, A. A. The peripheral blood
transcriptome dynamically reflects system wide biology: a potential diagnostic tool. *J*
Lab Clin Med **147**, 126-132, doi:10.1016/j.lab.2005.10.005 (2006).

43   Xiong, Y. *et al.* Transcriptomic characteristics of bronchoalveolar lavage fluid and
peripheral blood mononuclear cells in COVID-19 patients. *Emerg Microbes Infect* **9**,
761-770, doi:10.1080/22221751.2020.1747363 (2020).

44   Wong, K. L. *et al.* The three human monocyte subsets: implications for health and
disease. *Immunol Res* **53**, 41-57, doi:10.1007/s12026-012-8297-3 (2012).

45   McInnes, L., Healy, J., Saul, N. & Großberger, L. UMAP: Uniform Manifold
Approximation and Projection. *Journal of Open Source Software* **3**,
doi:10.21105/joss.00861 (2018).

46   Liu, J. *et al.* Longitudinal characteristics of lymphocyte responses and cytokine profiles
in the peripheral blood of SARS-CoV-2 infected patients. *EBioMedicine* **55**, 102763,
doi:10.1016/j.ebiom.2020.102763 (2020).

47   Wang, F. *et al.* Characteristics of peripheral lymphocyte subset alteration in COVID-19
pneumonia. *J Infect Dis*, doi:10.1093/infdis/jiaa150 (2020).

48   Wen, W. *et al.* Immune cell profiling of COVID-19 patients in the recovery stage by
single-cell sequencing. *Cell Discov* **6**, 31, doi:10.1038/s41421-020-0168-9 (2020).

49   Zhang, D. *et al.* COVID-19 infection induces readily detectable morphological and
inflammation-related phenotypic changes in peripheral blood monocytes, the severity
of which correlate with patient outcome. *medRxiv preprint doi:*
https://doi.org/10.1101/2020.03.24.20042655. , doi:10.1101/2020.03.24.20042655
(2020).

50   Silvin, A. *et al.* Elevated Calprotectin and Abnormal Myeloid Cell Subsets Discriminate
Severe from Mild COVID-19. *Cell*, doi:10.1016/j.cell.2020.08.002 (2020).

51   Takahashi, T. *et al.* Sex differences in immune responses that underlie COVID-19
disease outcomes. *Nature*, doi:10.1038/s41586-020-2700-3 (2020).

52   Jiang, Y. *et al.* COVID-19 pneumonia: CD8(+) T and NK cells are decreased in number
but compensatory increased in cytotoxic potential. *Clin Immunol* **218**, 108516,
doi:10.1016/j.clim.2020.108516 (2020).
53   Christopher Maucourant, I. F., Andrea Ponzetta, Soo Aleman, Martin Cornillet, Laura
Hertwig Benedikt Strunz, Antonio Lentini, Björn Reinius, Demi Brownlie, Angelica
Cuapio Gomez, Eivind Heggernes Ask, Ryan M. Hull, Alvaro Haroun-Izquierdo, Marie
Schaffer, Jonas Klingström, Elin Folkesson, Marcus Buggert, Johan K. Sandberg, Lars I.
Eriksson, Olav Rooyackers, Hans-Gustaf Ljunggren, Karl-Johan Malmberg, Jakob
Michaëlsson, Nicole Marquardt, Quirin Hammer, Kristoffer Strålin, Niklas K.
Björkström; and the Karolinska COVID-19 Study Group. Natural killer cell
immunotypes related to COVID-19 disease severity. *Science Immunology*,
doi:10.1126/sciimmunol.abd6832 (2020). (2020).
54   Diao, B. *et al.* Reduction and Functional Exhaustion of T Cells in Patients With
Coronavirus Disease 2019 (COVID-19). *Front Immunol* **11**, 827,
doi:10.3389/fimmu.2020.00827 (2020).
55   Katsuyama, E. *et al.* The CD38/NAD/SIRTUIN1/EZH2 Axis Mitigates Cytotoxic CD8 T Cell
Function and Identifies Patients with SLE Prone to Infections. *Cell Rep* **30**, 112-123
e114, doi:10.1016/j.celrep.2019.12.014 (2020).
56   Zheng, H. Y. *et al.* Elevated exhaustion levels and reduced functional diversity of T cells
in peripheral blood may predict severe progression in COVID-19 patients. *Cell Mol
Immunol*, doi:10.1038/s41423-020-0401-3 (2020).
57   Zhou, R. *et al.* Acute SARS-CoV-2 Infection Impairs Dendritic Cell and T Cell Responses.
Immunity, doi:10.1016/j.immuni.2020.07.026 (2020).
58   De Biasi, S. *et al.* Marked T cell activation, senescence, exhaustion and skewing
towards TH17 in patients with COVID-19 pneumonia. *Nat Commun* **11**, 3434,
doi:10.1038/s41467-020-17292-4 (2020).
59   Francisco, L. M., Sage, P. T. & Sharpe, A. H. The PD-1 pathway in tolerance and
autoimmunity. *Immunol Rev* **236**, 219-242, doi:10.1111/j.1600-065X.2010.00923.x
(2010).
60   Zhang, J. Y. *et al.* Single-cell landscape of immunological responses in patients with
COVID-19. *Nat Immunol* **21**, 1107-1118, doi:10.1038/s41590-020-0762-x (2020).
61   Westmeier, J. *et al.* Impaired Cytotoxic CD8(+) T Cell Response in Elderly COVID-19
Patients. *mBio* **11**, doi:10.1128/mBio.02243-20 (2020).
62   Mazzoni, A. *et al.* Impaired immune cell cytotoxicity in severe COVID-19 is IL-6
dependent. *J Clin Invest*, doi:10.1172/JCI138554 (2020).
63   Kusnadi, A. *et al.* Severely ill COVID-19 patients display impaired exhaustion features
in SARS-CoV-2-reactive CD8(+) T cells. *Sci Immunol* **6**,
doi:10.1126/sciimmunol.abe4782 (2021).
64   Cabrera-Marante, O. *et al.* Perforin gene variant A91V in young patients with severe
COVID-19. *Haematologica* **105**, 2844-2846, doi:10.3324/haematol.2020.260307
(2020).
65   Zeng, Y. *et al.* Peripheral Blood Mononuclear Cell Metabolism Acutely Adapted to
Postprandial Transition and Mainly Reflected Metabolic Adipose Tissue Adaptations
to a High-Fat Diet in Minipigs. *Nutrients* **10**, doi:10.3390/nu10111816 (2018).
66   Codo, A. C. *et al.* Elevated Glucose Levels Favor SARS-CoV-2 Infection and Monocyte
Response through a HIF-1alpha/Glycolysis-Dependent Axis. *Cell Metab*,
doi:10.1016/j.cmet.2020.07.007 (2020).

67 Jaiswal, N., Agrawal, S. & Agrawal, A. High fructose-induced metabolic changes
enhance inflammation in human dendritic cells. *Clin Exp Immunol* **197**, 237-249,
doi:10.1111/cei.13299 (2019).
68 Di Wu, T. S., Xiaobo Yang, Jian-Xin Song Mingliang Zhang Chengye Yao Wen Liu, Muhan
Huang, Yuan Yu, Qingyu Yang Tingju Zhu, Jiqian Xu, Jingfang Mu, Yaxin Wang, Hong
Wang, Tang Tang, Yujie Ren Yongran Wu, Shu-Hai Lin, Yang Qiu, Ding-Yu Zhang, You
Shang, Xi Zhou. Plasma Metabolomic and Lipidomic Alterations Associated with
COVID-19. *Natl Sci Rev*, doi:10.1093/nsr/nwaa086 (2020).
69 Ferrara, F., De Rosa, F. & Vitiello, A. The Central Role of Clinical Nutrition in COVID-19
Patients During and After Hospitalization in Intensive Care Unit. *SN Compr Clin Med*,
1-5, doi:10.1007/s42399-020-00410-0 (2020).
70 Ferreira, C. *et al.* Non-obese diabetic mice select a low-diversity repertoire of natural
regulatory T cells. *Proc Natl Acad Sci U S A* **106**, 8320-8325,
doi:10.1073/pnas.0808493106 (2009).
71 Watford, M. Glutamine and glutamate: Nonessential or essential amino acids? *Anim
Nutr* **1**, 119-122, doi:10.1016/j.aninu.2015.08.008 (2015).
72 Max W. S. Oliveira, J. B. M., Marcos R. de Oliveira, Alfeu Zanotto-Filho, Guilherme A.
Behr, Ricardo F. Rocha, José C. F. Moreira, Fábio Klamt. Scavenging and antioxidant
potential of physiological taurine concentrations against different reactive
oxygen/nitrogen species
. *Pharmaceutical Reports* **62**, 185-193 (2010).
73 Cunningham, L., Simmonds, P., Kimber, I., Basketter, D. A. & McFadden, J. P. Perforin
and resistance to SARS coronavirus 2. *J Allergy Clin Immunol* **146**, 52-53,
doi:10.1016/j.jaci.2020.05.007 (2020).
74 Maskus, D. J. *et al.* Characterization of a novel inhibitory human monoclonal antibody
directed against Plasmodium falciparum Apical Membrane Antigen 1. *Sci Rep* **6**, 39462,
doi:10.1038/srep39462 (2016).
75 Lorenz, M. A., Burant, C. F. & Kennedy, R. T. Reducing time and increasing sensitivity
in sample preparation for adherent mammalian cell metabolomics. *Anal Chem* **83**,
3406-3414, doi:10.1021/ac103313x (2011).
76 Marimuthu, R. *et al.* Characterization of Human Monocyte Subsets by Whole Blood
Flow Cytometry Analysis. *J Vis Exp*, doi:10.3791/57941 (2018).
77 Autissier, P., Soulas, C., Burdo, T. H. & Williams, K. C. Evaluation of a 12-color flow
cytometry panel to study lymphocyte, monocyte, and dendritic cell subsets in humans.
Cytometry A **77**, 410-419, doi:10.1002/cyto.a.20859 (2010).
78 Davies, R., Vogelsang, P., Jonsson, R. & Appel, S. An optimized multiplex flow
cytometry protocol for the analysis of intracellular signaling in peripheral blood
mononuclear cells. *J Immunol Methods* **436**, 58-63, doi:10.1016/j.jim.2016.06.007
(2016).
889

Fig. 1

Fig. 2

a

CD8⁺HLA-DR⁺ T cells

| HC | SARS-CoV-2⁺ | | |

Fig. 3

Fig. 4

Fig. 5

a

VIP

b

Compounds correlated with the Glucose

c

Compounds correlated with the Lactate

Compounds correlated with the Fructose

Fig. 6