A mathematical framework for dissecting the normative foundations of conservation decisions

Guillaume Latombe[1*], Bernd Lenzner[1], Anna Schertler[1], Stefan Dullinger[2], Michael Glaser[1], Ivan Jarić[3,4], Aníbal Pauchard[5,6], John R. U. Wilson[7,8], Franz Essl[1,8]

[1]BioInvasions, Global Change, Macroecology – Group, Department of Botany and Biodiversity Research, University Vienna, Rennweg 14, 1030 Vienna, Austria

[2]Division of Conservation Biology, Vegetation and Landscape Ecology, Department of Botany and Biodiversity Research, University of Vienna, Rennweg 14, 1030 Vienna, Austria

[3]Biology Centre of the Czech Academy of Sciences, Institute of Hydrobiology, Na Sádkách 702/7, 370 05 České Budějovice, Czech Republic

[4]University of South Bohemia, Faculty of Science, Department of Ecosystem Biology, Branišovska 1645/31a, 370 05 České Budějovice, Czech Republic

[5]Laboratorio de Invasiones Biológicas (LIB), Facultad de Ciencias Forestales, University of Concepcion Victoria 631, Concepción, Chile

[6]Institute of Ecology and Biodiversity (IEB), Santiago, Chile

[7]South African National Biodiversity Institute, Kirstenbosch Research Centre, Claremont 7735, South Africa

[8]Centre for Invasion Biology, Department of Botany and Zoology, Stellenbosch University, Private Bag X1, Matieland 7602, South Africa

*Corresponding author: latombe.guillaume@gmail.com

Abstract

Perspectives in conservation are based on a variety of value systems and normative postulates. Such differences in how people value nature and its components lead to different perceptions of the morality of conservation goals and approaches, and often underlie disagreements in the formulation and implementation of environmental management policies. Specifically, whether a specific conservation decision (e.g. killing feral cats to save birds threatened with extinction) is viewed as appropriate or not can vary among people with different value systems. Here, we present a conceptual framework that mathematically formalises the interplay of value systems. This framework is intended to serve as a heuristic tool to clarify normative postulates in conservation approaches, and it highlights how different value systems can lead to different prioritizations of available management options. We compare how management decisions would likely be viewed under three different idealised value systems (ecocentric conservation, new conservation, and sentientist conservation). We illustrate the utility of the framework by applying it to case studies involving invasive alien species, rewilding, and trophy hunting. By making value systems and their consequences in practice explicit, the framework facilitates debates on contested conservation issues. Finally, we believe dissecting the normative postulates of conservation decisions will facilitate understanding and addressing conservation conflicts.

Keywords: environmental management; impact; invasive alien species; moral values; rewilding; speciesism; trophy hunting

INTRODUCTION

The consideration of the moral relationship between humans and nature and the consequent ethical obligations for conservation is relatively recent in Western culture. Environmental ethics only emerged as an academic discipline in the 1970s (Brennan and Lo, 2016) and the concepts of values, duty, and animal welfare, are now increasingly appreciated in applied ecology and conservation (e.g. Díaz et al., 2018; Dubois et al., 2017). These concepts are complex and the formulation and implementation of environmental management policies is often associated with conflicts between different groups of stakeholders and between people with different values and interests, for example for the management of charismatic alien species (Crowley et al., 2017; Jarić et al., 2020; Redpath et al., 2013). An examination of how value systems could be explicitly accounted for in decision making could offer opportunities for better identifying conflicts, potentially helping to resolve them, and overall improve environmental management.

Value systems consider more or less inclusive moral communities (the elements with intrinsic or inherent value), ranging from humans (anthropocentrism), to further incorporate sentient beings (sentientism), living beings (biocentrism), and collectives (such as species and ecosystems; ecocentrism) (Table 1, Figure 1). The definition of moral communities can also be influenced by additional elements (such as spatial elements in the case of nativism), and, at the assessor level, by personal experience. These value systems underlie different sets of explicit or implicit normative postulates that define different conservation approaches. Because these postulates are considered as truths, we have to consider them as non-negotiable. Therefore, if the truths of different value systems are incompatible, conflicts can arise between different groups of stakeholders whose members share common moral values (e.g. Crowley et al., 2017). In particular, conservationists who value biodiversity *per se* [as defined initially by Soulé (1985), called hereafter 'traditional conservation' (Table 1)] can be at odds with those who value biodiversity based on human welfare and economic aspects ['new conservation' (Kareiva and Marvier, 2012)] (Doak et al., 2015; Kareiva, 2014; Soulé, 2014), or with those based on animal welfare ['conservation welfare' (Beausoleil et al., 2018), or, to a certain extent, 'compassionate conservation' (Driscoll and Watson, 2019; Hayward et al., 2019; Wallach et al., 2018)].

In the following, our aim is to conceptualize and decompose value systems and to explore their repercussions for the perception of conservation management actions. First, we recapitulate four archetypal value systems in environmental affairs and relate them to different conservation philosophies. Since identifying commonalities in the perspectives of different parties is key in conflict management (Redpath et al., 2013), we then introduce a formal framework to conceptualise these value systems, and examine how it can be applied to clarify different perspectives. Finally, we discuss opportunities for identifying commonalities between different value systems that may enable identifying widely acceptable solutions to otherwise polarising issues.

Here, we will focus on a Western perspective of value systems that have been internationally considered for environmental policies and the management of nature (e.g. Mace 2014). The archetypes of value systems and of conservation approaches were chosen for their importance in the past and present literature and their clear differences, to illustrate our framework. We acknowledge this is a small part of the global diversity of value systems. It would be interesting to see if our framework could be applied to other contexts, to identify its limitations.

ENVIRONMENTAL ETHICS AND CONSERVATION

From the valuation of humans to that of ecosystems: a complex spectrum of perspectives.

The Western perspective of moral valuation encompasses a diverse set of value systems with respect to the components of nature. Traditionally, one can distinguish at least four archetypal value systems: anthropocentrism, sentientism, biocentrism, and ecocentrism (Palmer et al., 2014; Rolston III, 2003) (Table 1; Figure 1).

Anthropocentrism values nature by the benefits it brings to people through ecosystem services, which encompasses economic, biological, and cultural benefits humans can derive from nature (Díaz et al., 2018). One justification for anthropocentrism is that humans are (arguably) the only self-reflective moral beings, and people are therefore both the subject and object of ethics

(Rolston III, 2003), therefore constituting the moral community (Table 1). In an anthropocentric

system, individuals from non-human species only have value based on their benefits or

disservices for humans (instrumental or non-instrumental).

Sentientism considers that humans and all sentient animals value their life, and experience

pleasure, pain, and suffering. All sentient individuals should therefore also be part of the moral

community (i.e. have an intrinsic value). In this view, it is the sentience (e.g. measured through

cognitive ability, Singer 2009), rather than species themselves, that has intrinsic value.

Biocentrism considers that life has intrinsic value. Although different perspectives on why life

has value exist (see e.g. Taylor 2011), all living organisms are valued equally for being alive, and

not differently based on any other trait.

Some ecocentric, or holistic, value systems consider that ecological collectives, such as species

or ecosystems, have intrinsic value, independently from the individuals that comprise them.

Species can have different values, i.e. speciesism (Table 1), and these values can be influenced

by a multitude of factors, discussed in more details below.

In practice, the separation between these different archetypes is blurry, and values given to

different species may vary under the same general approach. For example, biocentrism can range

from complete egalitarianism between organisms, i.e. universalism (Table 1), to a gradual

valuation resembling sentientism. These four value systems can also interact with other systems

that use other criteria than the intrinsic characteristics of individuals to define the moral

community. For example, nativism is a system that values organisms indigenous to a spatial

location or an ecosystem over those that have been anthropogenically introduced. Nativism can

therefore interact with any of the four systems presented above to alter the value attributed to a

species in a given context. Finally, the attribution of values to individuals from different species

can be deeply embedded in the individual psychologies of the assessor (Palmer et al., 2014;

Waytz et al., 2019). Values and personal interests interact in making and expressing moral

judgements (Essl et al., 2017). Thus, the archetypes of value systems presented above are rarely

expressed in a clear and obvious fashion.

To account for such complexity of the concept of value, in the following, we distinguish between

'intrinsic', 'inherent', and 'utilitarian', value (our definitions). Intrinsic value is the value

possessed by an individual or collective as defined by one of the systems above, and is therefore

fixed in time. Inherent value accounts for influences of many factors to distort the intrinsic value

(note that other scholars have used 'inherent' differently, e.g. Regan, 2004; Taylor, 1987). These

factors include charisma (Courchamp et al., 2018; Jarić et al., 2020), anthropomorphism (Tam et

al., 2013), organismic complexity (Proença et al., 2008), neoteny (Stokes, 2007), cultural

importance (Garibaldi and Turner, 2004), religion (Bhagwat et al., 2011), or parochialism (Waytz

et al., 2019) (Table 1). For example, some alien species that did not have any inherent value prior

to their introduction have been incorporated in local cultures, therefore providing them a novel

and higher inherent value such as horses being linked to a strong local cultural identity in some

parts of the USA (Rikoon, 2006). Inherent value can often be considered to be fixed at the time

scale of a management action, but can nonetheless vary over short time scales in some situations

(see the example of the Oostvaardersplassen nature reserve below). Utilitarian value is

determined only from an anthropocentric perspective. It is context-dependent and can change

rapidly, for example in the case of commercial exploitation.

Moral valuation and the management of nature.

Conservation practices can historically be divided into three main categories, closely related to

specific systems of moral valuation. At one extreme, a 'nature for itself' (Table 1) view mostly

excludes humans from the assessment of the efficacy of conservation management actions. This

ecocentric perspective is the foundation of traditional conservation as defined by Soulé (1985),

and relies on the four following normative postulates: "diversity of organisms is good,"

"ecological complexity is good," "evolution is good," and "biotic diversity has intrinsic value"

(Soulé 1985). It historically underlies widely-used conservation tools, like the IUCN Red List of

Threatened Species (IUCN, 2019). Ecocentrism is often not limited to the valuation of species,

but can encompass wider collectives, i.e. assemblages of species and functions, or ecosystems.

This other perspective is captured, for example, by the IUCN Red List of Ecosystems (IUCN-

CEM, 2016), and it is strongly reflected in international conservation-legislation such as the

175 Convention on Biological Diversity (UNEP CBD, 2010). In the following we refer to traditional

176 conservation as an ecocentric value system where species are intrinsically valuable (nature for

177 itself; Figure 1) and humans are mostly excluded from management. We acknowledge that this is

178 an archetypal view of traditional conservation, and is used here simply for illustrative purposes.

180 By contrast the anthropocentric 'nature for people' perspective (Mace, 2014) values species and

181 ecosystems only to the extent that they contribute to the wellbeing of humans. These values

182 encompass ecosystem services that help sustain human life (Bolund and Hunhammar, 1999) or

183 economic assets (Fisher et al., 2008), and can rely on the assessment of species and ecosystem

184 services in terms of their economic value (Costanza et al., 1997). The 'nature for people'

185 perspective is, in practice, exemplified by 'new conservation', also termed 'social conservation'

186 (Kareiva, 2014; Miller et al., 2011) (Table 1). It has been argued that such an anthropocentric

187 perspective will, by extension, help and even be necessary to maximize the conservation of

188 nature. The exact set of normative postulates proposed by the proponents of new conservation is

189 not clearly defined (Miller et al., 2011), leading to differences of interpretation and heated

190 debates in recent years (Doak et al., 2015; Kareiva, 2014; Kareiva and Marvier, 2012; Soulé,

191 2014).

193 More recently, the necessity to account for the interdependence between the health of nature and

194 human wellbeing (i.e. 'people and nature' (Mace, 2014)) has been advocated in the United

195 Nations Sustainable Development Goals (Weitz et al., 2018). Similarly, "nature-based solutions"

196 is an approach endorsed by the IUCN, which aims at protecting, sustainably managing, and

197 restoring, natural or modified ecosystems, to simultaneously provide human wellbeing and

198 biodiversity benefits (Cohen-Shacham et al., 2016). The 'One Health' approach, endorsed by the

199 Food and Agriculture Organization, the World Health Organization, and the World Organisation

200 for Animal Health also acknowledges the interdependence between the state of ecosystems and

201 human health and zoonoses (Gibbs, 2014) The difference between people and nature and new

202 conservation approaches therefore lies in the fact that it merges anthropocentric and ecocentric

203 systems, rather than considering that the latter will be addressed by focusing on the former (see

204 Section "Nature despite/for/and people" below for details).

206   Finally, recent approaches advocate for the consideration of animal welfare in conservation,

207   therefore incorporating sentientist perspectives ['conservation welfare' (Beausoleil et al., 2018);

208   'compassionate conservation' (Ramp and Bekoff, 2015; Wallach et al., 2018)].

FRAMING MORAL VALUES FOR CONSERVATION

Formulation of a mathematical framework.

Many of the conflicts in conservation are grounded in the failure to identify and formalize

differences in world views, which contain elements of the four archetypes presented above,

influenced by cultural norms, economic incentives etc. (Essl et al., 2017). Here, we propose a

mathematical formulation as a method to clarify moral discourses in conservation. We argue that

defining concepts as mathematical terms can make differences in value systems and their

normative postulates more transparent to all participants of the discourse. Doing so can help to

identify and facilitate the discussion of shared values and incompatibilities between different

environmental policies and management options (Miller et al., 2011), and contribute to manage

conflicts (Redpath et al., 2013). In a similar vein, Parker et al. (1999) proposed a mathematical

framework for assessing the environmental impacts of alien species. This work was highly

influential in the conceptualisation of biological invasions (being cited nearly 2,000 times until

October 2020 according to Google Scholar), rather than by its direct quantitative application. We

also acknowledge that this approach has specific limitations, which are discussed below.

Our mathematical formalisation conceptualises the consequences of environmental management

actions. We argue that it can account for different value systems, including the anthropocentric,

sentientist, biocentric and species-based ecocentric systems (see Appendix S1 for an extension to

ecocentrism beyond species and considering wider collectives, i.e. ecosystems), while also

accounting for cultural and personal contexts. These consequences C can be conceptualised as a

combination of the impact of an action on the different species or individuals involved and the

value given to said species and individuals under different value systems as follows:

236                          $$C = \sum_{species\ s} \bar{I}_s \times V_s \times N_s{}^a$$ Eq.1

where $\bar{I}_s$ is a function (e.g. mean, maximum, etc.) of the impact (direct and indirect) resulting from the management action on all individuals of species s, V_s is the inherent value attributed to an individual of species s (as described above), N_s is the abundance of species s, and a determines the importance given to a species based on its abundance or rarity (and enables to account for the importance of a species rather than an individual, see below). The unit of C depends on how other parameters are defined, which themselves depend on the value system considered. In summary, the higher the impact on species with high values, the higher the consequences.

The parameter a can take both positive and negative values. A value of 1 would mean that the same importance is given to all individuals in the moral community (Table 1) and be typical of individual-centred value systems, i.e. anthropocentrism, sentientism, and biocentrism. As a result, impacts on larger populations would weigh more on the consequences. As a decreases towards 0, the correlation between the value of a species and its abundance decreases. For $a = 0$, the consequence of a management action becomes abundance-independent. For $a < 0$, rare species would be valued higher than common species (or the same impact would be considered to be higher for rare species), for example due to the higher risk of extinction. $a > 1$ would give a disproportionate weight to abundant species, which are often important for providing ecosystem services (Gaston, 2010).

The impact I_s can be limited to the death of individuals or the risk of extinction of species, but can also include animal welfare, biophysical states, etc. I_s encompasses both the direct impact of a management action, and its indirect impact resulting from biotic interactions. One would therefore need to define a baseline corresponding to either i) the lowest possible measurable level of impact (e.g. being alive if death is the only measure of impact, or no sign of disease and starvation for biophysical states; this would obviously be more complicated for welfare), so that I would only be positive; or ii) the current state of the system, in which case impacts could be positive or negative for different species. The duration over which to measure such impact should also be determined. The exact quantification of impact will be influenced by different value systems and personal subjectivity, and some impacts may be considered incommensurable

(Essl et al., 2017). The average impact $\bar{I}_s$ could be used as a measure at the species-level, as
different individuals may experience different impacts, if the management action targets only
part of a given population. Using the average impact is not without shortcomings though, and
other measures such as the maximum impact experienced by individuals, or more complex
functions accounting for the variability of impacts and values across individuals of a same
species, and of different types of impact, may be used. Under anthropocentric perspectives,
impacts are influenced by the utilitarian values of species.
275
276 **Application of the mathematical framework.**
277
Note that Equation 1 produces relative rather than absolute values. The values of consequences C
of a management action under different value systems cannot be directly compared with each
other, because the unit and range of values of C can vary between value systems, as they may
consider different measures of impact. Instead, Equation 1 can be used to rank a set of
management actions for each value system based on their assessed consequences, to identify
management actions representing consensus, compromises or conflicts amongst value systems.
284
285 If different types of impacts considered under a value system occur simultaneously but are not
286 directly comparable, Equation 1 should be applied to each one separately. The overall
287 consequences of management actions can then be ranked under a given value system using
288 approaches such as multi-criteria decision analyses (MCDA, Huang, Keisler, & Linkov, 2011).
289
Similarly, some actions might not follow moral norms compared to others despite having more
desirable consequences. For example, killing individuals may be considered less moral, but more
efficient to preserve biodiversity or ecosystem services than using landscape management.
Solving these moral dilemmas is complex, and how to do so is beyond the scope of this
publication. It is nonetheless important to account for the morality of actions to address
environmental conflicts. Different approaches for resolving environmental conflicts, including
MCDA (Wittmer et al., 2006) and operational research (Kunsch et al., 2009), may offer avenues
to do so.
298

In the following, we show how, despite the difficulty to quantify the variables described above,

this framework can be used as a heuristic tool to capture the implications of considering different

value systems for determining the appropriateness of a conservation action, and to better

understand conservation disputes.

NATURE DESPITE/FOR/AND PEOPLE

Over the past decade there has been some debate between proponents of traditional conservation,

and those of new conservation (Table 1), as each group assumes different relationships between

nature and people. Here, we show how the formal conceptualisation of Equation 1 could help

clarifying the position of the new conservation approach in response to its criticisms (Kareiva,

2014).

Nature despite people and traditional conservation

Traditional conservation is based on an ecocentric value system and seeks to maximize diversity

of organisms, ecological complexity, and to enable evolution (Soulé, 1985). For simplification,

we will consider an extreme perspective of traditional conservation, championed by 'fortress

conservation' (Büscher, 2016; Siurua, 2006), i.e. excluding humans from the moral community.

To capture these aspects, consequences C in Equation 1 can be more specifically expressed as

follows:

$$C = \sum_{\text{species s (excluding humans)}} \bar{I}_s \times V_s \times N_s^{a<0} \qquad \text{Eq. 2}$$

Assigning a stronger weight to rare species ($a < 0$) accounts for the fact that rare species are

more likely to go extinct, decreasing the diversity of organisms. Evolution is not explicitly

accounted for in Equation 2, but emphasising the importance of rare species will lead to fewer

extinctions and can increase the chance of species with high evolutionary potential to remain

within the species pool. Ecological complexity can be accounted for by attributing higher values

to species with rare traits and specific functional roles.

Nature for people and new conservation

New conservation considers that many stakeholders ("resource users", Kareiva, 2014) tend to have an anthropocentric value system, and that conservation approaches that do not incorporate such a perspective will likely not succeed at maximizing diversity of organisms (Kareiva, 2014; Kareiva and Marvier, 2012). Under anthropocentrism, species are only conserved due to their utilitarian value, i.e. their effect on *I* for humans, rather than based on an inherent value *V*. Different groups of stakeholders are likely to be impacted differently (e.g. different monetary benefits / losses), and we propose the following extension of Equation 1 to account for this variability:

$$C = \sum_{\text{stakeholders } t} \bar{I}_t \times V_t \times N_t \qquad\qquad \text{Eq.3}$$

where $\bar{I}_t$ is the average impact of management on the group of stakeholders *t*, including indirect impacts through the effect of management of non-human species, V_t is the value of the group of stakeholders *t*, and N_t is its abundance. Note that including inherent values V_t in Equation 3 does not imply that we consider that different humans should be valued differently, but that is a view that some people have, and this needs to appear here to capture the full spectrum of perceived consequences of a management action.

New conservation holds an ambiguous perspective, stating that it should make "sure people benefit from conservation", while at the same time does not "want to replace biological-diversity based conservation with a humanitarian movement" (Kareiva, 2014). Using our framework, we interpret this as considering that one can design management actions that minimize consequences *C* under both Equations 2 and 3. The link between biodiversity and ecosystem services has been demonstrated, even if many unknowns remain (Cardinale et al., 2012; Chivian and Bernstein, 2008), implying that high biodiversity can indeed support the provision of ecosystem services to humans. Such an approach will necessarily distinguish between "useful" species and others, and impacts will be perceived differently by different groups of stakeholders. Considering multiple types of impacts (economic benefits/losses, access to nature, health, etc.) while accounting for cultural differences, would increase the pool of useful species (comparing the resulting equation

outputs using, for example, MCDA). The outcome of the two approaches would then potentially
be more aligned with each other. This broad utilitarian perspective is captured in the most recent
developments of new conservation approaches, which consider a wide range of nature
contributions to people, rather than just ecosystem services (Díaz et al., 2018).

People and nature

People and nature views seek to simultaneously benefit human wellbeing and biodiversity. Under
this perspective, Equations 2 and 3 should therefore be combined in a single approach, for
example using MCDA (Huang et al., 2011), to capture a more diverse set of value systems than
Equations 2 and 3 alone, even if the two approaches generate divergent results.

Equations 2 and 3 clearly show how failing to explicitly define normative postulates for
conservation approaches can lead to extreme interpretations (excluding humans or considering
specific utilities of species) and conflicts and to overlook nuances between perspectives. In
particular, it has been argued that the normative postulates of new conservation need to be more
explicitly defined (Miller et al., 2011). Our framework could help doing so, by being more
explicit about how new conservation would be defined relative to the traditional conservation
and the people and nature perspective.

THE CASE OF ANIMAL WELFARE

The question of if and how animal welfare should be integrated into conservation practice is
increasingly debated (e.g. Hampton and Hyndman, 2018). Recently, conservation welfare (Table
1) has proposed to consider both the "fitness" (physical states) and "feelings" (mental
experiences) of non-human individuals in conservation practice (Beausoleil et al., 2018). Based
on virtue ethics rather than consequentialism, compassionate conservation (Table 1) also
emphasises animal welfare and is based on the "growing recognition of the intrinsic value of
conscious and sentient animals" (Wallach et al., 2018). It opposes the killing of sentient invasive
alien species; the killing of sentient native predators threatening endangered species; or the
killing of sentient individuals from a given population to fund broader conservation goals.

Despite the near-universal support of conservation practitioners and scientists for compassion

towards wildlife and ensuring animal welfare (e.g. Russell et al. 2016, Oommen et al. 2019,

Hayward et al. 2019), compassionate conservation has sparked vigorous responses (Driscoll and

Watson, 2019; Griffin et al., 2020; Hampton et al., 2018; Hayward et al., 2019; Oommen et al.,

2019). Amongst the main criticisms of compassionate conservation is that the absence of action

can result in unintended detrimental effects and increased suffering for individuals of other or the

same species (including humans), as a result of altered biotic interactions across multiple trophic

levels, i.e. "not doing anything" is an active choice that has consequences (Table 2). However,

since compassionate conservation is not based on consequentialism, it uses different criteria to

assess the appropriateness of conservations actions (but see Wallach et al., 2020 for responses to

some criticisms). Our purpose here is not to discuss the relevance or irrelevance of virtue ethics

for conservation (see Griffin et al., 2020 for such criticism). Instead, we propose discussing

animal welfare from the perspective of consequentialism (Hampton et al., 2018), i.e. more

aligned with the approach of conservation welfare (Beausoleil et al., 2018), and to show how it

may be aligned with or oppose the traditional and new conservation approaches.

A mathematical conceptualisation of animal welfare

A consequentialist, sentientist perspective aims at maximizing happiness, or conversely

minimising suffering, for all sentient beings, an approach also termed 'utilitarianism' (Singer,

1980; Varner, 2008). Suffering is therefore considered as a measure of impact (or, in

mathematical terms, impact is a function of suffering, which can be expressed as $I(S_s)$ in

Equation 1).

It has become widely accepted that animals experience emotions (de Waal, 2011). Emotions have

been shown to be linked to cognitive processes (Boissy and Lee, 2014), which differ greatly

among species (MacLean et al., 2012), and behavioural approaches have been used to evaluate

and grade emotional responses (e.g. Désiré et al. 2002; but see Shriver 2006 and Bermond et al.

2008 for different conclusions about the capacity of animals to experience suffering). We

therefore postulate that the quantification of suffering is conceptually feasible in the context of

the heuristic tool presented here. In a utilitarian approach, the inherent value of a species would

therefore be a function of its capacity to experience emotions and suffering E_s, which can be

expressed as $V(E_s)$ instead of V_s in Equation 1.

Under these considerations for defining impact and value of species, the consequences of a

conservation action can be computed as:

$$C = \sum_{\text{species s}} \overline{I(S_s)} \times V(E_s) \times N_s{}^{a=1}$$ Eq.4

Although $V(E_s)$ should be measured in an objective fashion, many factors may influence the

relationship between the inherent value and the emotional capacity of a species. For example,

high empathy (Table 1) from the observer will tend to make the distribution uniform, whereas

anthropomorphism and parochialism (Table 1) may lead to higher rating of the emotional

capacities of species phylogenetically close to humans or with which humans are more often in

contact, such as pets. Finally, we assumed that $a = 1$, to give equal importance to any individual

regardless of the abundance of its species, as suffering and wellbeing are usually considered at

the individual level (Beausoleil et al., 2018).

Assessing suffering in the presence and absence of conservation management actions

The short-term suffering resulting from pain and directly caused by lethal management actions,

such as the use of poison to control invasive alien species (Twigg and Parker, 2010) or the use of

firearms and traps to cull native species threatening other native species (e.g. Proulx et al. 2016)

or humans (e.g. Gibbs and Warren 2015), is the most straightforward type of suffering that can

be assessed, and is usually sought to be minimised in all conservation approaches. Suffering can

have many other causes, and suffering of an individual may be assessed through a wide variety

of proxies, including access to food and water, death, number of dead kin for social animals,

physiological measurements of stress hormones, etc. Suffering can take various forms, and

commensurability can be an issue (Table 2), making the distinction between the morality of

lethal actions and non-lethal suffering complex. Non-lethal suffering can result from

unfavourable environmental conditions (e.g. leading to food deprivation) and occur over long

periods, while lethal actions could be carried out in a quick, non-painful fashion (e.g. Shao et al.
2018), and even lead to improved animal welfare (Wilson and Edwards, 2019), but may be
deemed immoral.
457
The concept of animal welfare and how to measure it is extremely complex (Beausoleil et al.,
2018), and defining it precisely is beyond the scope of this study. We nonetheless advocate a
conceptual approach that takes into account indirect consequences of management actions within
a certain timeframe and consider uncertainty (Table 2). Direct and indirect biotic interactions
may be explicitly modelled to quantify the impact on animals and their suffering. Simulation
models can also make projections on how populations may change in time, accounting for future
suffering.
465

Are traditional conservation and animal welfare compatible?

It has been argued that sentientism and ecocentrism are not fully incompatible (Varner, 2011).
The relationship between biodiversity and animal suffering can be formalised more clearly using
the traditional conservation and the sentientist Equations 2 and 4, to explore if the same
management action can minimize the consequences evaluated using the two equations (see also
Appendix S2 for the application of the framework to theoretical cases). The main difference with
the traditional vs new conservation debate here is that Equations 2 and 4 share a number of
species, whereas the new conservation Equation 3 only contains humans, which are excluded
from Equation 2. Even though the variables of Equation 4 differ from those of Equation 2 (V and
I are computed differently, and the value of a is different), it is possible that these equations will
vary in similar way for different management actions due to their similar structure, although this
would depend on the variety of impacts on humans that are considered in Equation 3. Finally, as
for the people and nature approach, the consequences of sentientist and ecocentric approaches
can be evaluated in combination, as suggested by conservation welfare (Beausoleil et al., 2018),
using tools such as MCDA (Huang et al., 2011; Wittmer et al., 2006).
482
One issue that may be irreconcilable between ecocentric approaches such as traditional
conservation and approaches based on sentientism is the fate of rare and endangered species with

limited or no sentience. Under utilitarian sentientism, the conservation of non-sentient species

ranks lower than the conservation of sentient species, and consequently they are not included in

Equation 4. For example, endangered plant species that are not a resource for the maintenance of

sentient populations would receive no attention, as there would be few arguments for their

conservation. Traditional conservation would focus on their conservation, as they would have a

disproportionate impact in Equation 2, due to low abundance leading to a high value for $N^{a<0}$.

Finally, it is important to note that the current body of knowledge shows that the link between

biodiversity and animal welfare mentioned above especially applies to the increase of native

biodiversity. Local increase of biodiversity due to the introduction of alien species may only be

temporary due to extinction debt (Kuussaari et al. 2009) and often results in a reduction of

ecosystem functioning (Cardinale et al., 2012). Therefore, it is important to distinguish between

nativism (Table 1) and the detrimental effects of *invasive* alien species on biodiversity and

ecosystem functioning and services (Bellard et al., 2016). Nativism would result in increasing the

inherent value V_s of native species (Figure 1), whereas in the second case, insights from science

on the impact of invasive alien species would modify the distribution $I(S)$ rather than the

distribution V_s. This can also apply to native species whose impacts on other species, such as

predation, are increased through environmental changes (Carey et al., 2012).

UNRESOLVED QUESTIONS AND LIMITATIONS

This framework shares similarities with mathematical approaches used in conservation triage

(Bottrill et al., 2008), but has two crucial differences. First, conservation triage is an ecocentric

perspective with variables that are comparatively easy to quantify. Bottrill et al. (2008) provided

an example using phylogenetic diversity as a measure of value V, and a binomial value b to

quantify biodiversity benefit that can be interpreted as the presence or absence of a species (i.e. I

$= 1 / b$). Because it is ecocentric, local species abundance is not considered, which corresponds

to setting $a = 0$. In this example, consequences (C) in the general Equation 1 are therefore

defined simply by V / b.

In contrast, our framework allows much more flexibility to encompass a range of value systems,

as shown above. Given that the data needed for quantifying parameters of Equations 1 to 4

related to value, impact, emotional capacity and suffering are scarce and often very difficult to

measure, this framework in its current form would nonetheless be difficult to use as a

quantitative decision tool to evaluate alternative management actions, contrary to triage

equations. Rather, our equations decompose the question underlying many controversies around

management decisions in conservation: what or who is valued, how, and by how much?

There are nonetheless a number of approaches that may be used to develop quantification

schemes for the different parameters of the framework. Grading systems may be developed to

assess impact and suffering based on various indicators, including appearance, physiology, and

behaviour (Beausoleil et al., 2018; Broom, 1988). For assessing the value of different species,

questionnaires may be used to assess how different species are valued by people, and influenced

by their social and cultural background, similar to what has been done to assess species charisma

(e.g. Colléony et al. 2017, Albert et al. 2018). It will nonetheless be important to acknowledge

the corresponding uncertainties in the assessment of impact and value, differences in perception

among societal groups for different taxa and potential shifts in perception over time (Table 2).

The second difference from conservation triage is that the latter considers additional criteria that

were not addressed here, including feasibility, cost, and efficiency (including related

uncertainties). The combination of these different perspectives calls for appropriate methods to

include them all in decision making, which can be done using MCDA (Huang et al., 2011). Here,

good communication and transparency of the decision process is key to achieve the highest

possible acceptance across stakeholders, and to avoid biases in public perception (see case

studies below for examples).

The issue of spatial and temporal scale also warrants consideration (Table 2). In the case of a

species that may be detrimental to others in a given location but in decline globally, the spatial

scale and the population considered for evaluating the terms of Equations 1 to 4 is crucial to

determine appropriate management actions. Similarly, management actions may also result in a

temporary decrease in welfare conditions for animals, which may increase later on (Ohl and Van

der Staay, 2012), or the impacts may be manifested with a temporal lag. In that case, determining

the appropriate time period over which to evaluate the terms of Equations 1 to 4 will not be

straightforward. Impacts will be of different importance depending on whether they occur in the

short- or long-term, especially since long-term impacts are harder to predict and involve higher

uncertainty. Discount rates (Table 2) may therefore be applied, in a similar way they are applied

to the future effects of climate change and carbon emissions (Essl et al., 2018), or to assess the

impact of alien species (Essl et al., 2017).

Equations 1 to 4 assume that all individuals from a given species have the same value or

emotional capacities (or use the average of the value across individuals). However, intraspecific

differences in value may be important for conservation. For example, reproductively active

individuals contributing to population growth/recovery may be given a higher value in an

ecocentric perspective. Trophy hunters might prefer to hunt adult male deer with large antlers.

Intraspecific value may also vary spatially, for example between individuals in nature reserves or

in highly disturbed ecosystems. Equation 1 may therefore theoretically be adapted to use custom

groups of individuals with specific values within species, similar to Equation 3.

Finally, it is crucial to account for biotic interactions in our framework to comprehensively

assess the indirect impacts of management actions on different species (Table 2). Some species

with low values V_s in a certain value system may be crucial for assessing the impact I_s on other

species. These biotic interactions will therefore determine the time frame over which the

framework should be applied, as impacts on one species at a given time may have important

repercussions in the future. These biotic interactions can be complex, and several tools, such as

simulation models and ecological network analyses can be used to address them. Concepts such

as keystone species (Mills et al., 1993) can also offer a convenient way to overcome such

complexity by modifying V_s rather than $\bar{I}_s$. Let us assume that a management action will have a

direct impact on a keystone species, which will result in indirect impacts on multiple other

species with inherent values. Increasing the value of the keystone species can result in the same

assessment of C as to explicitly model the biotic interactions and compute the resulting indirect

impacts $\bar{I}_s$.

CASE STUDIES ILLUSTRATING ETHICAL CONFLICTS IN CONSERVATION DECISIONS

In the following, we present three case studies where conservation actions have either failed, had adverse effects, or were controversial, and we explore how our framework can help to identify normative postulates underlying these situations.

Invasive alien species management: the case of the alien grey squirrel in Italy

The grey squirrel (*Sciurus carolinensis*) is native to North America and was introduced in various locations in Europe during the late nineteenth and the twentieth century (Bertolino, 2008). It threatens native European red squirrel (*Sciurus vulgaris*) populations through competitive exclusion, and is also a vector of transmission of squirrel poxvirus in Great Britain (Schuchert et al., 2014). Furthermore, it has wider impacts on woodlands and plantations, reducing value of tree crops, and potentially affects bird populations through nest predation (Bertolino, 2008).

Based on the impacts of the grey squirrel, an eradication campaign was implemented in 1997 in Italy, with encouraging preliminary results (Genovesi and Bertolino, 2001). However, this eradication campaign was halted by public pressure from animal rights movements. The strategy of the animal rights activists consisted in (i) humanising the grey squirrel and using emotive messages (referring to grey squirrels as "Cip and Ciop", the Italian names of the Walt Disney "Chip and Dale" characters) and (ii) minimising or denying the effect of grey squirrel on native taxa, especially the red squirrel (Genovesi and Bertolino, 2001). In addition, the activists did not mention (iii) the difference in abundance between a small founding population of grey squirrels that could be eradicated by managers, and a large population of native red squirrels that would be extirpated or severely impacted by grey squirrels if control was not implemented.

Genovesi & Bertolino (2001) explain that the main reason for the failure of the species management is a different perspective on primary values: the eradication approach was underlain

by species valuation, following traditional conservation, whereas the animal right activists and

the public were more sensitive to animal welfare. However, the application of our framework

reveals some inconsistencies in the animal right activists' arguments that could have been used to

advocate for the eradication approach. Translating this situation in our framework indicates that

(i) the humanisation of the grey squirrel consists of increasing the perception of its emotional

capacity $E_{gs} > E_{rs}$ (and therefore $V(E_{gs}) > V(E_{rs})$), (ii) minimising the impact of the grey squirrel

is equal to restricting the time scale to a short one and to likely minimising the amount of

suffering S caused by grey squirrels on other species, i.e. $S_{gs} = S_{rs}$ (and therefore $I(S_{gs}) = I(S_{rs})$)

without management and $S_{gs} > S_{rs}$ (and therefore $I(S_{gs}) > I(S_{rs})$) under management, and (iii) not

mentioning differences in species abundance corresponds to setting $a = 0$. Following these three

points, the consequences under management $C_m = I(S_{gs}) \times V(E_{gs}) + I(S_{rs}) \times V(E_{rs})$ are

higher than without management, due to the increase in $V(E_{gs})$ and $I(S_{gs})$.

The framework can thus be used to provide recommendations for what the advocates for the

eradication campaign would have needed to have done: i) increase the value E_{rs} of red squirrels

in a similar way as what was done for grey squirrels, so that their relative values compared to

grey squirrels would remain the same as before the communication campaign by the animal right

activists; ii) better explain the differences in animal suffering caused by the long-term presence

of the grey squirrel compared to the short-term, carefully designed euthanasia protocol, which

would avoid a subjective perception of the distribution of S; and iii) highlight the differences in

the number of individuals affected. The consequences would then be computed as $C = V(E_{gs}) \times$

$I(S_{gs}) \times N_{gs} + V(E_{rs}) \times I(S_{rs}) \times N_{rs}$. In that case, assuming for simplification the same

suffering through euthanasia for grey squirrels as red squirrels suffer from the grey sqirrels, and

the same value to individuals of each species (i.e. avoiding nativism), the mere differences $N_{rs} >$

N_{gs} in abundance would lead to a higher value of C without management. This would further

increase by extending the impacts of grey squirrels to other species, as mentioned above.

A more fundamental issue, however, is that in some value systems it would not be acceptable to

actively kill individuals, even if that meant letting grey squirrels eliminate red squirrels over long

periods of time (e.g. Wallach et al., 2018). The reluctance to support indirectly positive

conservation programs is a common issue (Courchamp et al., 2017). Whether an acceptable

threshold on consequences over which killing individuals could be determined through

discussion would depend, in part, on the willingness of the affected parties to compromise.

De-domestication: the case of Oostvaardersplassen nature reserve

De-domestication, the intentional reintroduction of domesticated species to the wild is a recent

practice in conservation that raises new ethical questions related to the unique status of these

species (Gamborg et al., 2010). Oostvaardersplassen is a Dutch nature reserve where two

domesticated species of large herbivores (Heck cattle, *Bos primigenius*, and konik horses, *Equus*

ferus caballus) have been 'rewilded' in addition to the reintroduction of the red deer (*Cervus*

elaphus) to act as landscape engineers by grazing (ICMO2, 2010). The populations increased

rapidly, as natural predators are missing and population regulation was not conducted, as a result

of a 'non-intervention-strategy'. The project was widely criticized when a considerable number

of individuals died from starvation during a harsh winter, resulting in the introduction of

population reduction by culling weak animals in order to prevent starvation (other approaches,

such as the reintroduction of large predators were discarded due to lack of experience and too

many uncertainties in efficiency, ICMO2 2010).

From a traditional conservation perspective, disregarding animal welfare and focusing on species

diversity and ecological restoration, the project was a success. The introduction of the three

herbivore species led to sustainable populations (despite high winter mortality events), and

ensured stability of bird populations without the need for further interventions (ICMO2 2010),

i.e. the conditions of many species were improved (the impact was lowered), leading to lower

consequences C overall (Equation 2). However, the welfare of individuals from the three

charismatic large herbivorous species became a point of conflict. Interestingly, it appears that the

conflict was driven by a shift in attitude, from considering the herbivore species as a natural way

to manage the grasslands to being part of the ecosystem changed the value V_s, or by the

importance given to their emotional capacity E_s (Ohl and Van der Staay, 2012), therefore leading

to increasing the consequences $C = V(E_s) \times I(S_s) \times N_s^{\,1}$ under sentientism, with S_s and N_s

constant. Temporal changes in the distributions of the V and E variables should therefore be

taken into account when implementing conservation management actions, and even monitored

through time in a way similar to adaptive management approaches. Another possible explanation

for this shift in attitude is that the notion of responsibility (Table 2) affected the moral aspect of

culling that should then be considered in parallel to its consequences. If culling animals can be

considered acceptable in some cases, it may not be the case if these individuals were

purposefully introduced.

As a result, the reserve management has examined a number of sustainable measures to improve

the welfare of individuals from the three species (therefore decreasing S_s to compensate the

increase in V_s). Among those were recommendations to increase access to natural shelter in

neighbouring areas of woodland or forestry, to create shelter ridges to increase survival in winter

as an ethical and sustainable solution, and to use early culling to regulate populations and avoid

suffering from starvation in winter (ICMO2 2010). This example shows how a combination of

two complementary management actions (the rewilding of the OVP and the provision of shelter)

led to minimised consequences under both the traditional conservation and the sentientist

Equations 2 and 4, whereas only rewilding would increase consequences under Equation 4.

Culling may still face opposition based on moral arguments though.

Trophy hunting

Trophy hunting, the use of charismatic species for hunting activities, has been argued to be good

for conservation when revenues are reinvested properly into nature protection and redistributed

across local communities, but faces criticisms for moral reasons (Di Minin et al., 2016; Lindsey

et al., 2007a). The action of killing some individuals to save others might be incompatible with a

deontological perspective, but, assuming a consequentialist perspective, the framework can be

applied to formalise the assessment of different management options (we are not considering

canned hunting here, the practice of farming animals for the specific purpose of being hunted).

In traditional conservation, trophy hunting is desirable if it directly contributes to the

maintenance of species diversity. The potential of trophy hunting to contribute to the

maintenance of biodiversity is via creating economic revenues, i.e. an anthropocentric

perspective, and it therefore falls under the umbrella of new conservation. In theory, trophy

hunting should lead to lower consequences than doing nothing for both the traditional and new

conservation (Equations 2 and 3), and therefore for the 'people and nature' approach, as they are

in this case not independent from each other (Lindsey et al., 2007b). Many social and biological

factors currently affect the efficacy of trophy hunting as a conservation tool. Corruption and

privatisation of the benefits have sometimes prevented the revenues to be reinvested into

conservation, but also to be redistributed across local communities, whereas doing so has been

shown to increase their participation in conservation actions with proven benefits for local

biodiversity (Di Minin et al., 2016). In other words, a decrease in the anthropocentric Equation 2

leads to a decrease in the ecocentric Equation 3. In addition, trophy hunting can lead to

unexpected evolutionary consequences (Coltman et al., 2003), overharvesting of young males

(Lindsey et al., 2007a), and disproportionate pressure on threatened species (Palazy et al., 2013,

2012, 2011) and therefore to population declines and potential detrimental effects on

biodiversity. Despite these issues, it has been argued that banning trophy hunting may create

replacement activities that would be more detrimental to biodiversity (Di Minin et al., 2016).

From an animal welfare perspective, trophy hunting appears to be in direct contradiction with a

decrease in animal suffering, and has been criticised by proponents of compassionate

conservation (Wallach et al., 2018). However, as for the culling of invasive alien species, we

suspect the story is more complex. First, there may be direct benefits for animal welfare, if

money from trophy hunting is reinvested in protection measures against poaching (which is

likely more detrimental to the hunted individuals). Second, to our knowledge, only few studies

have compared the welfare of individual animals to quantify the elements of the sentientist

Equation 4 (for example assessed through access to resources) in areas where trophy hunting is

practiced and where it is not. Given the links between biodiversity and animal welfare described

above, it seems plausible that good practice in trophy hunting may benefit the welfare of

individuals from other and from the same species.

CONCLUSIONS

A variety of value systems exist in conservation, which are based on different underlying

normative postulates and can differ between stakeholders, resulting in differing preferences for

conservation practices among people. Here, we have proposed a framework with a formal set of
equations to conceptualize and decompose these different perspectives. In this framework, the
different value systems supported by different conservation approaches follow the same
structure, but can differ in the variables that are used, and in the values they are taking. While
such a formalisation by necessity does not capture the full range of complex and nuanced real-
world situations in environmental decision-making, it provides a method to make their
underlying value systems and the resulting conflicts explicit and transparent, which is essential
for the planning and implementation of pro-active management. The search for consensus in
conservation can be counter-productive and favour status-quo against pro-active management
(Peterson et al., 2005), however our framework may help identify hidden commonalities between
seemingly antagonistic stances. We hope that this framework can foster debates on contested
conservation issues, and will ultimately contribute to a broader appreciation of different
viewpoints. In an increasingly complex world shaped by human activities, this is becoming ever
more important.
746
747

748 ## Acknowledgements
749

We thank Franck Courchamp, Vincent Devictor, Jordan Hampton, Jonathan Jeschke, and Thomas
Potthast for extremely useful comments on this manuscript. GL, BL, AS, FE, and SD were
supported by the Austrian Science Foundation FWF (BiodivERsA-Belmont Forum Project 'Alien
Scenarios', grant no. I 4011-B32; grant no. I3757-B29). AP was funded by Conicyt PIA CCTE
AFB170008. IJ acknowledges support by the J. E. Purkyně Fellowship of the Czech Academy of
Sciences. JRUW thanks the South African Department of Forestry, Fisheries, and the
Environment (DFFtE) for funding noting that this publication does not necessarily represent the
views or opinions of DFFtE or its employees.

REFERENCES

Albert, C., Luque, G.M., Courchamp, F., 2018. The twenty most charismatic species. PLoS One 13, e0199149. https://doi.org/10.1371/journal.pone.0199149

Beausoleil, N.J., Mellor, D.J., Baker, L., Baker, S.E., Bellio, M., Clarke, A.S., Dale, A., Garlick, S., Jones, B., Harvey, A., 2018. "Feelings and Fitness" not "Feelings or Fitness" – the raison d'être of conservation welfare, which aligns conservation and animal welfare objectives. Front. Vet. Sci. 5, 296.

Bellard, C., Cassey, P., Blackburn, T.M., 2016. Alien species as a driver of recent extinctions. Biol. Lett. 12, 20150623. https://doi.org/10.1098/rsbl.2015.0623

Bermond, B., Armstrong, S.J., Botzler, R.G., 2008. A neuropsychological and evolutionary approach to animal consciousness and animal suffering. Anim. ethics Read. 2, 99–112.

Bertolino, S., 2008. Introduction of the American grey squirrel (*Sciurus carolinensis*) in Europe: a case study in biological invasion. Curr. Sci. 95, 903–906.

Bhagwat, S.A., Dudley, N., Harrop, S.R., 2011. Religious following in biodiversity hotspots: challenges and opportunities for conservation and development. Conserv. Lett. 4, 234–240. https://doi.org/10.1111/j.1755-263X.2011.00169.x

Boissy, A., Lee, C., 2014. How assessing relationships between emotions and cognition can improve farm animal welfare. Rev. Sci. Tech. 33, 103–110.

Bolund, P., Hunhammar, S., 1999. Ecosystem services in urban areas. Ecol. Econ. 29, 293–301. https://doi.org/10.1016/S0921-8009(99)00013-0

Bottrill, M.C., Joseph, L.N., Carwardine, J., Bode, M., Cook, C., Game, E.T., Grantham, H., Kark, S., Linke, S., McDonald-Madden, E., 2008. Is conservation triage just smart decision making? Trends Ecol. Evol. 23, 649–654. https://doi.org/10.1016/j.tree.2008.07.007

Brennan, A., Lo, Y.-S., 2016. Environmental Ethics. Stanford Encycl. Philos.

Broom, D.M., 1988. The scientific assessment of animal welfare. Appl. Anim. Behav. Sci. 20, 5–19. https://doi.org/10.1016/0168-1591(88)90122-0

Büscher, B., 2016. Reassessing fortress conservation? New media and the politics of distinction in Kruger National Park. Ann. Am. Assoc. Geogr. 106, 114–129. https://doi.org/10.1080/00045608.2015.1095061

Cardinale, B.J., Duffy, J.E., Gonzalez, A., Hooper, D.U., Perrings, C., Venail, P., Narwani, A., Mace, G.M., Tilman, D., Wardle, D.A., Kinzig, A., Daily, G.C., Loreau, M., Grace, J.B.,

Larigauderie, A., Srivastava, D.S., Naeem, S., 2012. Biodiversity loss and its impact on

humanity. Nature 486, 59. https://doi.org/10.1038/nature11148

Carey, M.P., Sanderson, B.L., Barnas, K.A., Olden, J.D., 2012. Native invaders–challenges for

science, management, policy, and society. Front. Ecol. Environ. 10, 373–381.

https://doi.org/10.1890/110060

Chivian, E., Bernstein, A., 2008. Sustaining life: how human health depends on biodiversity.

Oxford University Press.

Cohen-Shacham, E., Walters, G., Janzen, C., Maginnis, S., 2016. Nature-based solutions to

address global societal challenges. IUCN, Gland. Switz. 97.

Colléony, A., Clayton, S., Couvet, D., Saint Jalme, M., Prévot, A.-C., 2017. Human preferences

for species conservation: Animal charisma trumps endangered status. Biol. Conserv. 206,

263–269. https://doi.org/10.1016/j.biocon.2016.11.035

Coltman, D.W., O'Donoghue, P., Jorgenson, J.T., Hogg, J.T., Strobeck, C., Festa-Bianchet, M.,

2003. Undesirable evolutionary consequences of trophy hunting. Nature 426, 655.

https://doi.org/10.1038/nature02177

Costanza, R., D'Arge, R., De Groot, R., Farber, S., Grasso, M., Hannon, B., Limburg, K.,

Naeem, S., O'neill, R. V, Paruelo, J., Raskin, R.G., Sutton, P., van den Belt, M., 1997. The

value of the world's ecosystem services and natural capital. Nature 387, 253–260.

https://doi.org/10.1016/S0921-8009(98)00020-2

Courchamp, F., Fournier, A., Bellard, C., Bertelsmeier, C., Bonnaud, E., Jeschke, J.M., Russell,

J.C., 2017. Invasion biology: specific problems and possible solutions. Trends Ecol. Evol.

32, 13–22. https://doi.org/10.1016/j.tree.2016.11.001

Courchamp, F., Jaric, I., Albert, C., Meinard, Y., Ripple, W.J., Chapron, G., 2018. The

paradoxical extinction of the most charismatic animals. PLoS Biol. 16, e2003997.

https://doi.org/10.1371/journal.pbio.2003997

Crowley, S.L., Hinchliffe, S., McDonald, R.A., 2017. Conflict in invasive species management.

Front. Ecol. Environ. 15, 133–141. https://doi.org/10.1002/fee.1471

de Waal, F.B.M., 2011. What is an animal emotion? Ann. N. Y. Acad. Sci. 1224, 191–206.

https://doi.org/10.1111/j.1749-6632.2010.05912.x

Désiré, L., Boissy, A., Veissier, I., 2002. Emotions in farm animals:: a new approach to animal

welfare in applied ethology. Behav. Processes 60, 165–180. https://doi.org/10.1016/S0376-

6357(02)00081-5

Di Minin, E., Leader-Williams, N., Bradshaw, C.J.A., 2016. Banning trophy hunting will

exacerbate biodiversity loss. Trends Ecol. Evol. 31, 99–102.

https://doi.org/10.1016/j.tree.2015.12.006

Díaz, S., Pascual, U., Stenseke, M., Martín-López, B., Watson, R.T., Molnár, Z., Hill, R., Chan,

K.M.A., Baste, I.A., Brauman, K.A., 2018. Assessing nature's contributions to people.

Science (80-.). 359, 270–272. https://doi.org/10.1126/science.aap8826

Doak, D.F., Bakker, V.J., Goldstein, B.E., Hale, B., 2015. What is the future of conservation?, in:

Protecting the Wild. Springer, pp. 27–35. https://doi.org/10.5822/978-1-61091-551-9_4

Driscoll, D.A., Watson, M.J., 2019. Science denialism and compassionate conservation: response

to Wallach et al. 2018. Conserv. Biol. 33, 777–780. https://doi.org/10.1111/cobi.13273

Dubois, S., Fenwick, N., Ryan, E.A., Baker, L., Baker, S.E., Beausoleil, N.J., Carter, S.,

Cartwright, B., Costa, F., Draper, C., 2017. International consensus principles for ethical

wildlife control. Conserv. Biol. 31, 753–760. https://doi.org/10.1111/cobi.12896

Essl, F., Erb, K., Glatzel, S., Pauchard, A., 2018. Climate change, carbon market instruments, and

biodiversity: focusing on synergies and avoiding pitfalls. Wiley Interdiscip. Rev. Clim.

Chang. 9, e486. https://doi.org/10.1002/wcc.486

Essl, F., Hulme, P.E., Jeschke, J.M., Keller, R., Pyšek, P., Richardson, D.M., Saul, W.-C., Bacher,

S., Dullinger, S., Estévez, R.A., Kueffer, C., Roy, H.E., Seebens, H., Rabitsch, W., 2017.

Scientific and normative foundations for the valuation of alien-species impacts: thirteen

core principles. Bioscience 67, 166–178. https://doi.org/10.1093/biosci/biw160

Fisher, B., Turner, K., Zylstra, M., Brouwer, R., De Groot, R., Farber, S., Ferraro, P., Green, R.,

Hadley, D., Harlow, J., 2008. Ecosystem services and economic theory: integration for

policy-relevant research. Ecol. Appl. 18, 2050–2067. https://doi.org/10.1890/07-1537.1

Gamborg, C., Gremmen, B., Christiansen, S.B., Sandoe, P., 2010. De-domestication: ethics at the

intersection of landscape restoration and animal welfare. Environ. Values 19, 57–78.

Garibaldi, A., Turner, N., 2004. Cultural keystone species: implications for ecological

conservation and restoration. Ecol. Soc. 9.

Gaston, K.J., 2010. Valuing common species. Science (80-.). 327, 154–155.

https://doi.org/10.1126/science.1182818

Genovesi, P., Bertolino, S., 2001. Human dimension aspects in invasive alien species issues: the

case of the failure of the grey squirrel eradication project in Italy. Gt. reshuffling Hum.

Dimens. invasive alien species. IUCN, Gland 113–119.

Gibbs, E.P.J., 2014. The evolution of One Health: a decade of progress and challenges for the

future. Vet. Rec. 174, 85–91.

Gibbs, L., Warren, A., 2015. Transforming shark hazard policy: Learning from ocean-users and

shark encounter in Western Australia. Mar. Policy 58, 116–124.

https://doi.org/10.1016/j.marpol.2015.04.014

Griffin, A.S., Callen, A., Klop-Toker, K., Scanlon, R.J., Hayward, M.W., 2020. Compassionate

conservation clashes with conservation biology: should empathy, compassion, and

deontological moral principles drive conservation practice? Front. Psychol. 11, 1139.

Hampton, J.O., Hyndman, T.H., 2018. Underaddressed animal-welfare issues in conservation.

Conserv. Biol.

Hampton, J.O., Warburton, B., Sandøe, P., 2018. Compassionate versus consequentialist

conservation. Conserv. Biol.

Hayward, M.W., Callen, A., Allen, B.L., Ballard, G., Broekhuis, F., Bugir, C., Clarke, R.H.,

Clulow, J., Clulow, S., Daltry, J.C., Davies-Mostert, H.T., Fleming, P.J.S., Griffin, A.S.,

Howell, L.G., Kerley, G.I.H., Klop-Toker, K., Legge, S., Major, T., Meyer, N., Montgomery,

R.A., Moseby, K., Parker, D.M., Périquet, S., Read, J., Scanlon, R., Seeto, R., Shuttleworth,

C., Somers, M.J., Tamessar, C.T., Tuft, K., Upton, R., Valenzuela-Molina, M., Wayne, A.,

Witt, R.R., Wüster, W., 2019. Deconstructing compassionate conservation. Conserv. Biol.

33, 760–768. https://doi.org/10.1111/cobi.13366

Huang, I.B., Keisler, J., Linkov, I., 2011. Multi-criteria decision analysis in environmental

sciences: ten years of applications and trends. Sci. Total Environ. 409, 3578–3594.

https://doi.org/10.1016/j.scitotenv.2011.06.022

ICMO2, 2010. Natural processes, animal welfare, moral aspects and management of the

Oostvaardersplassen, Report of the second International Commission on Management of the

Oostvaardersplassen (ICMO2). The Hague/Wageningen, Netherlands. Wing rapport 039.

November 2010.

IUCN-CEM, 2016. The IUCN Red List of Ecosystems. Version 2016-1. <http://iucnrle.org>.

IUCN, 2019. The IUCN Red List of Threatened Species. Version 2019-1.

<http://www.iucnredlist.org>.

Jarić, I., Courchamp, F., Correia, R.A., Crowley, S.L., Essl, F., Fischer, A., González-Moreno, P.,

Kalinkat, G., Lambin, X., Lenzner, B., 2020. The role of species charisma in biological

invasions. Front. Ecol. Environ. https://doi.org/10.1002/fee.2195

Kareiva, P., 2014. New conservation: setting the record straight and finding common ground.

Conserv. Biol. 28, 634–636. https://doi.org/10.1111/cobi.12295

Kareiva, P., Marvier, M., 2012. What is conservation science? Bioscience 62, 962–969.

https://doi.org/10.1525/bio.2012.62.11.5

Kunsch, P.L., Kavathatzopoulos, I., Rauschmayer, F., 2009. Modelling complex ethical decision

problems with operations research. Omega 37, 1100–1108.

Kuussaari, M., Bommarco, R., Heikkinen, R.K., Helm, A., Krauss, J., Lindborg, R., Öckinger,

E., Pärtel, M., Pino, J., Rodà, F., 2009. Extinction debt: a challenge for biodiversity

conservation. Trends Ecol. Evol. 24, 564–571. https://doi.org/10.1016/j.tree.2009.04.011

Lindsey, P.A., Frank, L.G., Alexander, R., Mathieson, A., Romanach, S.S., 2007a. Trophy

hunting and conservation in Africa: problems and one potential solution. Conserv. Biol. 21,

880–883. https://doi.org/10.1111/j.1523-1739.2006.00594.x

Lindsey, P.A., Roulet, P.A., Romanach, S.S., 2007b. Economic and conservation significance of

the trophy hunting industry in sub-Saharan Africa. Biol. Conserv. 134, 455–469.

https://doi.org/10.1016/j.biocon.2006.09.005

Mace, G.M., 2014. Whose conservation? Science (80-.). 345, 1558–1560.

https://doi.org/10.1126/science.1254704

MacLean, E.L., Matthews, L.J., Hare, B.A., Nunn, C.L., Anderson, R.C., Aureli, F., Brannon,

E.M., Call, J., Drea, C.M., Emery, N.J., 2012. How does cognition evolve? Phylogenetic

comparative psychology. Anim. Cogn. 15, 223–238. https://doi.org/10.1007/s10071-011-

0448-8

Miller, T.R., Minteer, B.A., Malan, L.-C., 2011. The new conservation debate: the view from

practical ethics. Biol. Conserv. 144, 948–957. https://doi.org/10.1016/j.biocon.2010.04.001

Mills, L.S., Soulé, M.E., Doak, D.F., 1993. The keystone-species concept in ecology and

conservation. Bioscience 43, 219–224. https://doi.org/10.2307/1312122

Ohl, F., Van der Staay, F.J., 2012. Animal welfare: At the interface between science and society.

Vet. J. 192, 13–19. https://doi.org/10.1016/j.tvjl.2011.05.019

Oommen, M.A., Cooney, R., Ramesh, M., Archer, M., Brockington, D., Buscher, B., Fletcher,

R., Natusch, D.J.D., Vanak, A.T., Webb, G., 2019. The fatal flaws of compassionate conservation. Conserv. Biol. 33, 784–787. https://doi.org/10.1111/cobi.13329

Palazy, L., Bonenfant, C., Gaillard, J.-M., Courchamp, F., 2013. On the use of the IUCN status for the management of trophy hunting. Wildl. Res. 39, 711–720. https://doi.org/10.1071/WR12121

Palazy, L., Bonenfant, C., Gaillard, J.-M., Courchamp, F., 2011. Cat dilemma: too protected to escape trophy hunting? PLoS One 6, e22424. https://doi.org/10.1371/journal.pone.0022424

Palazy, L., Bonenfant, C., Gaillard, J.M., Courchamp, F., 2012. Rarity, trophy hunting and ungulates. Anim. Conserv. 15, 4–11. https://doi.org/10.1111/j.1469-1795.2011.00476.x

Palmer, C., McShane, K., Sandler, R., 2014. Environmental ethics. Annu. Rev. Environ. Resour. 39, 419–442. https://doi.org/10.1146/annurev-environ-121112-094434

Parker, I.M., Simberloff, D., Lonsdale, W.M., Goodell, K., Wonham, M., Kareiva, P.M., Williamson, M.H., Von Holle, B., Moyle, P.B., Byers, J.E., Goldwasser, L., 1999. Impact: toward a framework for understanding the ecological effects of invaders. Biol. Invasions 1, 3–19. https://doi.org/10.1023/A:1010034312781

Peterson, M.N., Peterson, M.J., Peterson, T.R., 2005. Conservation and the myth of consensus. Conserv. Biol. 19, 762–767. https://doi.org/10.1111/j.1523-1739.2005.00518.x

Proença, V.M., Pereira, H.M., Vicente, L., 2008. Organismal complexity is an indicator of species existence value. Front. Ecol. Environ. 6, 298–299. https://doi.org/10.1890/1540-9295(2008)6[298:OCIAIO]2.0.CO;2

Proulx, G., Brook, R.K., Cattet, M., Darimont, C., Paquet, P.C., 2016. Poisoning wolves with strychnine is unacceptable in experimental studies and conservation programmes. Environ. Conserv. 43, 1–2. https://doi.org/10.1017/S0376892915000211

Ramp, D., Bekoff, M., 2015. Compassion as a practical and evolved ethic for conservation. Bioscience 65, 323–327. https://doi.org/10.1093/biosci/biu223

Redpath, S.M., Young, J., Evely, A., Adams, W.M., Sutherland, W.J., Whitehouse, A., Amar, A., Lambert, R.A., Linnell, J.D.C., Watt, A., 2013. Understanding and managing conservation conflicts. Trends Ecol. Evol. 28, 100–109. https://doi.org/10.1016/j.tree.2012.08.021

Regan, T., 2004. The case for animal rights. Univ of California Press.

Rikoon, J.S., 2006. Wild horses and the political ecology of nature restoration in the Missouri Ozarks. Geoforum 37, 200–211. https://doi.org/10.1016/j.geoforum.2005.01.010

Rolston III, H., 2003. Environmental ethics, in: Bunnin, N., Tsui-James, E.P. (Eds.), The
Blackwell Companion to Philosophy, 2nd Edition. Blackwell Publishing, Oxford.

Russell, J.C., Jones, H.P., Armstrong, D.P., Courchamp, F., Kappes, P.J., Seddon, P.J., Oppel, S.,
Rauzon, M.J., Cowan, P.E., Rocamora, G., 2016. Importance of lethal control of invasive
predators for island conservation. Conserv. Biol. 30, 670–672.
https://doi.org/10.1111/cobi.12666

Schuchert, P., Shuttleworth, C.M., McInnes, C.J., Everest, D.J., Rushton, S.P., 2014. Landscape
scale impacts of culling upon a European grey squirrel population: can trapping reduce
population size and decrease the threat of squirrelpox virus infection for the native red
squirrel? Biol. Invasions 16, 2381–2391. https://doi.org/10.1007/s10530-014-0671-8

Shao, Y., Li, M., Zhang, W., Ji, Y., Hayes, D.J., 2018. World's Largest Pork Producer in Crisis:
China's African Swine Fever Outbreak. Agric. Policy Rev. 2018, 1.

Shriver, A., 2006. Minding mammals. Philos. Psychol. 19, 433–442.
https://doi.org/10.1080/09515080600726385

Singer, P., 2009. Speciesism and moral status. Metaphilosophy 40, 567–581.
https://doi.org/10.1111/j.1467-9973.2009.01608.x

Singer, P., 1980. Utilitarianism and vegetarianism. Philos. Public Aff. 325–337.

Siurua, H., 2006. Nature above people: Rolston and " fortress" conservation in the south. Ethics
Environ. 71–96.

Soulé, M., 2014. The "New Conservation." Conserv. Biol. 27, 895–897.
https://doi.org/10.1111/cobi.12147

Soulé, M.E., 1985. What is conservation biology? Bioscience 35, 727–734.
https://doi.org/10.2307/1310054

Stokes, D.L., 2007. Things we like: human preferences among similar organisms and
implications for conservation. Hum. Ecol. 35, 361–369. https://doi.org/10.1007/s10745-
006-9056-7

Tam, K.-P., Lee, S.-L., Chao, M.M., 2013. Saving Mr. Nature: Anthropomorphism enhances
connectedness to and protectiveness toward nature. J. Exp. Soc. Psychol. 49, 514–521.
https://doi.org/10.1016/j.jesp.2013.02.001

Taylor, P.W., 2011. Respect for nature: A theory of environmental ethics. Princeton University
Press.

Taylor, P.W., 1987. Inherent value and moral rights. Monist 70, 15–30.

Twigg, L.E., Parker, R.W., 2010. Is sodium fluoroacetate (1080) a humane poison? The influence of mode of action, physiological effects, and target specificity. Anim. Welf. 19, 249–263.

UNEP CBD, 2010. Strategic plan for biodiversity 2011–2020 and the Aichi targets, in: Report of the Tenth Meeting of the Conference of the Parties to the Convention on Biological Diversity.

Varner, G., 2011. Environmental ethics, hunting, and the place of animals, in: The Oxford Handbook of Animal Ethics. https://doi.org/10.1093/oxfordhb/9780195371963.013.0032

Varner, G., 2008. Utilitarianism and the evolution of ecological ethics. Sci. Eng. Ethics 14, 551–573. https://doi.org/10.1007/s11948-008-9102-5

Wallach, A.D., Batavia, C., Bekoff, M., Alexander, S., Baker, L., Ben-Ami, D., Boronyak, L., Cardilin, A.P.A., Carmel, Y., Celermajer, D., Coghlan, S., Dahdal, Y., Gomez, J.J., Kaplan, G., Keynan, O., Khalilieh, A., Kopnina, H., Lynn, W.S., Narayanan, S.R., Santiago-Ávila, F.J., Yanco, E., Zemanova, M.A., Ramp, D., 2020. Recognizing animal personhood in compassionate conservation. Conserv. Biol. https://doi.org/10.1111/cobi.13494

Wallach, A.D., Bekoff, M., Batavia, C., Nelson, M.P., Ramp, D., 2018. Summoning compassion to address the challenges of conservation. Conserv. Biol. 32, 1255–1265. https://doi.org/10.1111/cobi.13126

Waytz, A., Iyer, R., Young, L., Haidt, J., Graham, J., 2019. Ideological differences in the expanse of the moral circle. Nat. Commun. 10, 1–12. https://doi.org/10.1038/s41467-019-12227-0

Weitz, N., Carlsen, H., Nilsson, M., Skånberg, K., 2018. Towards systemic and contextual priority setting for implementing the 2030 Agenda. Sustain. Sci. 13, 531–548. https://doi.org/10.1007/s11625-017-0470-0

Wilson, G.R., Edwards, M., 2019. Professional kangaroo population control leads to better animal welfare, conservation outcomes and avoids waste. Aust. Zool. 40, 181–202.

Wittmer, H., Rauschmayer, F., Klauer, B., 2006. How to select instruments for the resolution of environmental conflicts? Land use policy 23, 1–9.

Tables

1005

1006   **Table1.** Glossary of terms as they are used for the purposes of this paper.

Term	Definition
Anthropocentrism	Value system that considers humans to be the sole, or primary, holder of moral standing, and therefore the concern of direct moral obligations. Non-human species are considered only to the extent that they affect humans (Rolston III 2003; Palmer *et al.* 2014).
Anthropomorphism	"The attribution of human personality or characteristics to something non-human, like an animal, object, etc." (Oxford English Dictionary).
Biocentrism	Value system considering all living beings as the concern of direct moral obligations (Rolston III 2003; Palmer *et al.* 2014).
Collectivism	Value system in which a group or collective has a higher value than the individuals that compose it (Wallach *et al.* 2018).
Compassionate conservation	Conservation approach inspired by virtue ethics based on four tenets: i) do no harm; ii) individuals matter; iii) inclusivity (the value of an individual is independent from the context of the population, e.g. nativity, rarity, etc.); and iv) peaceful coexistence (Ramp & Bekoff 2015; Wallach *et al.* 2018).
Conservation welfare	Conservation approach aiming at minimizing animal suffering (Beausoleil et al., 2018).
Consequentialism	"An ethical doctrine which holds that the morality of an action is to be judged solely by its consequences" (Oxford English Dictionary).
Deontology	A normative ethical theory considering that "choices are morally required, forbidden, or permitted" (Alexander & Moore 2016).
Ecocentrism	Value system considering that species, their assemblages and their functions, as well as more broadly ecosystems, rather than individuals, are the concern of direct moral obligations (Rolston III 2003; Palmer *et al.* 2014).

Empathy	"The quality or power of projecting one's personality into or mentally identifying oneself with an object of contemplation, and so fully understanding or appreciating it." (Oxford English Dictionary). Empathy will influence the inherent value given to individuals from other species.
Impact (for the purposes of the framework, Eq.1)	Impact refers to any effect that modifies the wellbeing, health or resilience (for non-sentient beings) of an individual, from physical pain to emotional suffering and death (these notions being interrelated, but not equivalent).
Inherent value (our definition)	Value possessed by an individual or collective, accounting for the effects of multiple context-dependent factors to modify intrinsic value.
Intrinsic value	Value possessed by an individual or collective as defined by a system of moral valuation, such as anthropocentrism, sentientism, biocentrism or ecocentrism. Intrinsic value is fixed in time.
Invasive alien species	"Plants, animals, pathogens and other organisms that are non-native to an ecosystem, and which may cause economic or environmental harm or adversely affect human health" (*Regulation (EU) No 1143/2014 of the European Parliament and of the Council of 22 October 2014 on the prevention and management of the introduction and spread of invasive alien species*).
Moral community	The group of beings considered to have intrinsic moral value (Shoemaker 2010). The size of the group depends on the value system. For example, the moral community is restricted to humans in case of Anthropocentrism.
Nativism	Value system considering that species that have evolved in a given location have a higher value in this location than species that have evolved somewhere else. In nativism, value varies spatially (Wallach *et al.* 2018).
Nature despite people	Management conceptual approach aiming at conserving biological diversity (focusing on species and habitats) specifically in response to human impacts on the environment, e.g. sustainable use (Mace 2014).

Nature for itself	Management conceptual approach aiming at conserving biological diversity (focusing on wilderness and natural habitats) through human exclusion, for example through the creation of parks and protected areas (Mace 2014).
Nature for people	Management conceptual approach aiming at conserving the components of nature beneficial to humans (focusing on ecosystems and their services) (Mace 2014).
Neoteny	"The retention of juvenile characteristics in a (sexually) mature organism" (Oxford English Dictionary).
New conservation	Discipline aiming at preserving biological diversity through the conservation of natural elements providing services and contribution to human wellbeing (Kareiva & Marvier 2012; Kareiva 2014).
Normative postulate	Value statements that make up the basis of an ethic of appropriate attitudes toward other forms of life (Soulé 1985).
Parochialism	Ideology in which moral regard is directed "towards socially closer and structurally tighter targets, relative to socially more distant and structurally looser targets", and, by extension, to species phylogenetically, cognitively, or in appearance closer to humans (Waytz *et al.* 2019).
People and nature	Management conceptual approach considering that humans and nature are interdependent and therefore aiming at achieving compromises in the conservation of nature and human wellbeing. (Mace 2014).
Sentience	The ability to experience phenomenal consciousness, i.e. the qualitative, subjective, experiential, or phenomenological aspects of conscious experience, rather than just the experience of pain and pleasure (Allen & Trestman 2017).
Sentientism	Value system considering sentient beings as the concern of direct moral obligations (Rolston III 2003; Palmer *et al.* 2014).
Speciesism	Value system in which some species are considered to have a higher value than others, for various possible reasons (Singer 2009). Speciesism

	is often used to refer to the superiority of humans, which is a specific expression of speciesism as considered in this paper.
Suffering	Negative emotion, sometimes called emotional distress, experienced by sentient beings, and which can result from different causes, including but not limited to physical pain (Dawkins 2008; Farah 2008).
Traditional conservation	Discipline aiming at preserving biological diversity through the management of nature, and based on four value-driven normative postulates: "diversity of organisms is good," "ecological complexity is good," "evolution is good," and "biotic diversity has intrinsic value" (Soulé 1985). Traditional conservation is rooted in ecocentrism.
Utilitarian value	Value given to an individual or collective by humans, based on its utility.
Virtue ethics	Ethical doctrine that emphasizes the virtues, or moral character as the reason for action (Hursthouse & Pettigrove 2018).

1007

1008 **Table 2.** List of factors to consider regarding the effects of environmental management actions

1009 from an environmental ethics perspective.

Factor	Influence on variables and outputs in Equations 1 to 5
Biotic interactions	The impact or suffering of individuals from one species can be caused by individuals from another species, either through direct or indirect interactions. Management actions can therefore also have non-trivial indirect impacts on some species.
Capacity to provide ecosystem services	The presence of a specific species may increase the fitness/welfare of other species through the ecosystem services it provides. Since these effects can be difficult to quantify explicitly, the value of such species may be increased in Equations 1 to 4 to account for them.
Discounting rate	Rate at which impacts that occur in the future lose importance.
Impact quantification and commensurability	How the impacts of management actions are quantified is also dependent on value systems, as some impacts (such as death) may be considered incommensurable to others (such as suffering).
Responsibility from previous actions	Previous human actions on certain species, such as reintroduction of domesticated species or the introduction of alien species can change the perception of the public and therefore change the inherent value attributed to these species, or change the morality of an action, in addition to obviously having an impact on these species.
Spatial scale	The spatial scale will change the abundance N and the number of species considered. As a result, a management action that is more beneficial than another at small scale may not be such at a larger scale, and reciprocally. Additionally, the spatial scale can change the inherent value of species, for example under nativism, or because of the range of cultures that are considered.
Temporal scale	The time frame over which the impact or the suffering of individuals is computed can change their values. Management actions may decrease welfare of individuals on the short term, but

	be beneficial on the long term once the ecosystem has stabilised. Similarly, not culling some population may cause less suffering on the short term, but increase it in the future by disrupting ecosystem services, leading to population collapse due to lack of resources, etc.
Uncertainty of impact	The complexity of an ecological system can make the impact of management actions on different species difficult to assess precisely, therefore creating potential errors, especially in the presence of multiple biotic interactions. This may lead to an incorrect estimation of the consequences C.
Uncertainty of value expressions and preferences	Quantifying the value given by a person or a group of people to an individual is difficult, context-dependent, and highly subjective. Sensitivity analyses on the distribution of values can be used to account for such uncertainty.

1010

1011

1012

Figure

Figure 1. Differences between value systems influenced by a) anthropocentrism, b) sentientism, c) biocentrism and d) ecocentrism. Anthropocentrism, sentientism and biocentrism all value individuals inherently, but consider different moral communities, i.e. their values depend on the category of species they belong to, with {humans} ∈ {sentient beings} ∈ {all living organisms}. Ecocentrism, in contrast, is not based on individuals, but on ecological collectives, i.e. on species or on species assemblages and ecosystems. Species outside of the moral community may have a utilitarian value for species in the moral community (represented by the arrows), which will be reflected by changes in the impact variable. Note that species can have both an inherent and a utilitarian value. Within the moral community, species may have equal inherent values, but subjective perceptions and different value systems can assign different values to different species. The skewness of the value distribution then indicates the degree or strength of

1027   speciesism with respect to the species of reference, assumed here to be the human species, and is

1028   influenced by many factors, including charisma, cultural context, etc.