A minimal yet flexible likelihood framework to assess correlated evolution

Abdelkader Behdenna[1,2,*]   Patrice Petot[1,2]   Joël Pothier[1]

Camille Nous[3]   Amaury Lambert[2,4]   Guillaume Achaz[5,2,1]

[1]Institut de Systématique, Évolution, Biodiversité (ISYEB), Muséum National d'Histoire Naturelle, CNRS UMR 7205, Sorbonne Université, École Pratique des Hautes Études, Université des Antilles, Paris, France

[2]SMILE group, Center for Interdisciplinary Research in Biology (CIRB), Collège de France, CNRS UMR 7241, INSERM U 1050, Paris, France

[3]Laboratoire Cogitamus, Sorbonne Université, Paris, France

[4]Laboratoire de Probabilités, Statistique et Modélisation (LPSM), Sorbonne Université, CNRS UMR 8001, Université de Paris, Paris, France

[5]Éco-anthropologie, Muséum National d'Histoire Naturelle, CNRS UMR 7206, Université de Paris, Paris, France

[*]Current affiliation: Epigene Labs, 7 square Gabriel Fauré, 75017 Paris, France

Corresponding author: `guillaume.achaz@mnhn.fr`

1   Abstract

An evolutionary process is reflected in the sequence of changes through time of any trait (e.g. morphological, molecular). Yet, a better understanding of evolution would be procured by characterizing correlated evolution, or when two or more evolutionary processes interact. A wide range of parametric methods have previously been proposed to detect correlated evolution but they often require significant computing time as they rely on the estimation of many parameters. Here we propose a minimal likelihood framework modelling the joint evolution of two traits on a known phylogenetic tree. The type and strength of correlated evolution is characterized by few parameters tuning mutation rates of each trait and interdependencies between these rates. The framework can be applied to study any discrete trait or character ranging from nucleotide substitution to gain or loss of a biological function. More specifically, it can be used to 1) test for independence between two evolutionary processes, 2) identify the type of interaction between them and 3) estimate parameter values of the most likely model of interaction. In its current implementation, the method takes as input a phylogenetic tree together with mapped discrete evolutionary events on it and then maximizes the likelihood for one or several chosen scenarios. The strengths and limits of the method, as well as its relative power when compared to a few other methods, are assessed using both simulations and data from 16S rRNA sequences in a sample of 54 γ-enterobacteria. We show that even with datasets of less than 100 species, the method performs well in parameter estimation and in the selection of evolutionary scenario.

2   Introduction

Evolutionary processes often turn out to be often interdependent at all levels of organization, from molecules (Shindyalov et al., 1994) to ecosystems (Van Valen, 1973). The work described in this article focuses on what we term *correlated evolution*, a lack of independence in the evolution of two or more traits of the same living entity whether it is an individual, a sequence or a species (Achaz and Dutheil, 2020). For the sake of simplicity, we will use the generic term *trait* to describe any morphological trait, life history trait or any molecular character. In many cases, one mutation in one trait impacts the evolution of other traits. We naturally acknowledge patterns of correlated evolution between morphological traits (e.g. size and body mass) or between molecular traits (e.g. residues of the same proteins) but functional constraints can also arise in metabolic or regulation networks (Fraser et al., 2004) without any physical contact between the

partners.

Patterns of apparent correlated evolution can have different causes. First, and this would be the central focus of this work, they can be caused by relevant biological interactions. Depending on the exact nature of the interaction, patterns can differ. For example, synergistic effects result in positive correlation whereas antagonistic ones cause negative correlations. Genetic interactions at the molecular level are named epistasis (Phillips, 2008). A mutation may for example be deleterious in certain contexts, while beneficial in others (Weinreich et al., 2005). More generally, a combination of two mutations can be more or less fitter than the sum of the two individual effects. The catalog of fitness values of all possible combinations of mutations constitutes a fitness landscape (Wright, 1932), an object that still fascinates evolutionary biologist today (Achaz et al., 2014; Visser and Krug, 2014; Yi and Dean, 2019). The structure of the fitness landscape is highly predictive of the expected pattern of correlated evolution (Achaz and Dutheil, 2020).

In this study, we restrict ourselves to the simple case of only two evolutionary processes. Compensated Pathogenic Deviations (CPDs) constitute an illuminating example of correlated evolution. CPDs are instances of the more general Bateson-Dobzhansky-Muller incompatibilities (Bateson, 1909; Dobzhansky, 1934; Muller, 1942; Orr, 1996; Welch, 2004). They are alleles that are highly deleterious in a focal species but wild type in one or several other species. The deleteriousness of the mutated allele must then be balanced by one or more *compensatory* mutations, assuming that the effect is not simply due to different non-genic environments. The more deleterious the mutation, the stronger the epistatic compensation. CPDs have been reported for not only for humans (Kondrashov et al., 2002) but also for insects (Kulathinal et al., 2004) and they can be as common as 10% of the deleterious amino acid substitutions. Depending on the exact nature of the epistasis between the two mutations, different mutational histories will have different likelihoods. If the first mutation to occur is the deleterious one, then the occurrence of the compensatory mutation must follow very quickly or even co-occur with the first one. When the compensatory mutation occurs first, the second one, that is no more deleterious, can occur with some time lag. Both possible scenarios can be re-expressed in terms of induction of one event onto the other; the induction represents the intensity in which the triggering event favors the other. The inferred strength of the induction will be however different for both scenarios: for example, a strong induction in the first case and only a mild induction in the second one.

Although direct interactions cause patterns of correlated evolution, the reciprocal is not true, for at least two reasons.

First, an observed pattern of correlation can result from indirect interactions between the two focal traits. As always, correlation does not imply causation. For example, two morphological traits can jointly respond to changes of a hidden environmental variable without having direct interaction. Similarly, two residues in proteins can show patterns of correlated evolution because they both interact with a third residue. To overcome this issue, an interesting proposal is to fit a model where only pairwise interactions are allowed, preventing indirect coupling in the model. A popular and successful approach is the DCA (Direct Coupling Analysis) method (Weigt et al., 2009; Morcos et al., 2011). It models the abundance of sequences in nature by probabilities of presence in a Potts model. In such a model, the distribution depends not only on marginal frequencies at each site (local abundance) but also on joint frequencies for all pairs of sites (pairwise epistasis). The underlying assumption is that deleterious combinations of residues should be rare enough so that we do no see them in the wild. The main challenge of DCA is to infer the parameters of a highly parameterized model (one value per residue per site plus one value per pair of residues per pair of sites), which typically requires thousands of pseudo-independent sequences. Several methods have been developed to infer correctly and efficiently the model parameters (Weigt et al., 2009; Morcos et al., 2011; Baldassi et al., 2014; Ekeberg et al., 2013) but they are obviously computationally demanding. Importantly, the co-evolving sites detected by DCA have been shown to be excellent predictors of amino acid contacts in the 3D structures (Marks et al., 2011), much better than the ones detected by simple correlation metrics such as Mutual Information (MI) (Chiu and Kolodziejczak, 1991; Martin et al., 2005). MI methods have however been shown to be decent indicators of interactions (Bitbol, 2018) and are so fast to compute that they can be measured for all pairs of sites on alignments of complete bacterial genomes (Pensar et al., 2019).

Second, patterns of apparent correlated evolution can be due to "phylogenetic inertia" (Harvey and Pagel, 1991). More specifically, species, sequences or individuals cannot be regarded as statistically independent samples because they partially share a history. This shared history is often represented by a phylogenetic tree. Species closer in the tree have traits that are closer in values and this extends to pairs of traits. Hence, the variable phylogenetic proximity between species create patterns of correlation for traits that are not biologically interacting. In this regard, phylogeny can be seen as a hidden variable. After the initial proposal of Felsenstein (1985) to use of statistically independent contrasts (differences of traits values for sister species), several methods have been developed to specifically take phylogenetic inertia into account. Among them, we can cite a generalized version of least squares, entitled PGLS, used to compute correlated

evolution among continuous traits (Grafen, 1989; Martins, 1994; Martins and Hansen, 1997).

In this article, we are interested in quantifying correlated evolution between discrete traits. We will restrict the term, *evolutionary process*, to describe any process that results in *discrete evolutionary events* on a phylogenetic tree. Discrete events abstract mutations at the molecular level, but more generally can be applied to any change, gain, or loss of any biological function, of any morphological trait or of any life history trait. These events can be mapped on the phylogenetic tree using ancestral character reconstruction (Shindyalov et al., 1994). Once the events are mapped, one can then test for the independence between the evolutionary paths of the two traits and whether both traits tend to show co-variation in the same branches of the tree (Tufféry and Darlu, 2000; Dutheil et al., 2005; Dutheil and Galtier, 2007) or whether one type of event tends to precede the other (Kryazhimskiy et al., 2011; Behdenna et al., 2016). For cases where events are placed (ignoring reconstruction uncertainties), p-values can be computed analytically for any type of correlated evolution (one event precedes the other, both events co-occur, one event prohibits the other, etc.) using matrix formalism (Behdenna et al., 2016).

The null model of independence used to assign a p-value assumes that evolutionary events follow a Poisson process, so that they occur uniformly on the tree. More generally, evolutionary processes can be modeled by time-continuous Markov processes (Felsenstein, 1981). It is then possible to generalize the model for two evolutionary processes that explicitly depend on one onto the other (Pagel, 1994; Milligan, 1994). The model of independence corresponding to a sub-space of parameter values of the general model makes it is possible to test whether the two evolutionary processes are independent (or not) using likelihood ratio tests (LRT). Based on this idea, powerful likelihood-based approaches were proposed (Pagel, 1994; Milligan, 1994; Schöniger and von Haeseler, 1994; Tillier and Collins, 1995; Pollock et al., 1999; Baum and Donoghue, 2001; Pagel and Meade, 2006; Yeang et al., 2007; Dib et al., 2014) but they are generally computationally demanding. The methods involve exploring or maximizing over a likelihood surface that has as many dimensions as the number of parameters. For two binary traits, there are 4 possible states for the two traits (00, 01, 10, 11). When the two processes are independent, there are four transition rates (two for each trait in both directions) but there is a 4-by-4 rate matrix describing the complete process. The rate matrix is defined by 12 parameters, or 8 when double mutations are forbidden. More complex models, like 2 sites with 20 amino acids each, depend on more parameters (on the order of few hundreds). Consequently, these methods need larger datasets and

expensive computation time.

Here, we propose a likelihood framework that is based on a minimal model of correlated evolution. The framework generates a series of nested models with 2 to 8 parameters that correspond to the mutation rates of each process and interactions between them. We show that these parameters are core values that characterize patterns of correlated evolution. The current implementation of the method assumes that the tree is correctly inferred and that the events are correctly placed on the tree. Using maximum likelihood, the method can (i) test for independence between two evolutionary processes, (ii) find the most likely type of correlated evolution between them (obligate or preferential sequential order, reciprocal synergy, incompatibility, etc.) and (iii) estimate the strength of the interaction.

3   Model

3.1   A minimal model of correlated evolution

The model detailed here describes the joint distribution of two processes describing sequences of evolutionary events E_1 and E_2 respectively, on a given tree T.

We assume that events E_i, $i \in \{1, 2\}$ occur on the branches of tree T according to a Poisson process with intensity m_i hereafter called *occurrence rate*. The originality of our model is that the value of m_i can change as a function of the *realization of events E_i and E_j* on the tree, as we now describe.

For each event E_i, its occurrence rate m_i can take 4 values μ_i, μ_i^*, ν_i or ν_i^*. The idea is that the trait modeled by process i can take two values, say A/a for process 1, B/b for process 2, and be in a "basal" state or in an "excited" state, resulting in four states, e.g. for process 1: A, A^*, a, a^*, as depicted in Figure 1A:

- The two basal rates μ_i and ν_i are meant to represent "natural" rates of occurrence of the event E_i, when trait i is in its basal state (non-starred), e.g. for process 1: A mutates to a at rate μ_1 and a mutates to A at rate ν_1.

- The starred rates μ_i^* and ν_i^* are meant to model "excited" rates of occurrence of the event E_i, when trait i is in its excited state (starred), e.g. for process 1: A^* mutates to a at rate μ_1^* and a^* mutates to A at rate ν_1^*. Note that a trait in its excited state falls back into a basal state upon mutation.

- Last, a trait i can jump from the basal state to the excited state, e.g. for process 1, from A to A^* or from a to a^*. These events are induced by occurrences of the

other type of event, here E_j with $j \neq i$.

Alternatively, the model can be fully specified by characterizing states by the *rates at which the next event will occur* and thanks to the following two rules (see again Figure 1):

1. When an E_j event occurs at a non-starred rate (the process j was either at μ_j or ν_j), the m_i rate of process i simultaneously switches to (or remains in) its corresponding starred rate: from μ_i (or μ_i^*) to μ_i^*, and similarly, from ν_i (or ν_i^*) to ν_i^*.

2. When an E_i event occurs, m_i switches from μ_i (or μ_i^*) to ν_i, and conversely, from ν_i (or ν_i^*) to μ_i, simultaneously losing its star if any.

The ratio $\lambda_i = \mu_i^*/\mu_i$ (potentially also $\lambda_i' = \nu_i^*/\nu_i$) can be thought of as an *induction* factor which measures the influence of an event on another. There is no order requirement between the rates: if $\lambda_i > 1$, the induction is positive (*i.e.* the intensity of the process i is increased after an event E_j) whereas it is negative when $\lambda_i < 1$. We assume that the interaction only concerns the next "induced" event, as it models the impact of an E_j event on the rate of occurrence of the next E_i event (and *vice-versa*). Framed in terms of compensatory events, the second event can be seen as compensating for the need created by the first one. In this sense, rule 2 also "consumes" the induction, as starred rates switch back to non-starred ones. It is noteworthy to mention that, in this model, a second subsequent occurrence of E_i will not set back the rate of j to a basal one (the "need" remains); only an E_j event will do it. Finally, rule 2 also sets the back and forth transitions between the μ and ν rates.

This framework can be used to evaluate many scenarios, some of which are particularly relevant from the biological standpoint and are detailed in the next section.

3.2   A selection of scenarios

Assuming a single occurrence rate for each trait (*i.e.* $\mu = \nu$) results in a model where the rate of E_i events are only modified by the occurrences of the other E_j events (and vice-versa). Having two rates (μ and ν) for the same process allows us to cover a wider range of scenarios. A possible example is a scenario where the first event is an adaptive change triggered by an environmental change whereas the second one is neutral (thus $\mu \gg \nu$), assuming that subsequent ones are not observed. An interesting usage of ν is the case of scenarios where events can occur only once, like irrevocable losses of genes or

functions. Events first occur at their standard basal rate ($\mu > 0$) and are then prohibited ($\nu = 0$).

Independence (H_0). A first basic null model is when events occur independently of one another and have a single rate. When the two processes are independent, their intensities are constant, regardless of the position of the occurrences of each type of event on the tree. The model therefore reduces to 2 parameters μ_1 and μ_2 (assuming that the ν rates equal the μ rates):

$$\mu_1 = \mu_1^* \; ; \; \mu_2 = \mu_2^*$$

The intensities of both processes are constant in the whole tree. When the two processes are independent but have more than one rate, a more relevant approach is to consider a 4-parameter model, where starred rates are equal to non-starred ones but where rates μ differ from rates ν.

$$\mu_1 = \mu_1^* \; ; \; \mu_2 = \mu_2^*$$
$$\nu_1 = \nu_1^* \; ; \; \nu_2 = \nu_2^*$$

Prohibition. This scenario models the case where E_2 events are not allowed unless they occur after an E_1 event (E_1 triggers the appearance of E_2). This would be the case of a strongly deleterious mutation that cannot occur without being compensated upstream. If we assume a single rate ($\mu = \nu$), there are 3 parameters

$$\mu_1 = \mu_1^* \; ; \; \mu_2 = 0 \; ; \; \mu_2^* > 0$$

Again, by letting ν rates differ from μ rates, one can also expand the model to include two rates for each process.

Asymmetric induction. E_1 events have a constant rate while the rate of E_2 is increased after an E_1 event has occurred. This scenario models, for example, a compensatory mutation that follows the fixation of a slightly deleterious mutation. More generally, this 3-parameter model describes the process by which direct compensatory events occur.

$$\mu_1 = \mu_1^* \; ; \; \mu_2 \; ; \; \mu_2^* = \lambda_\mu \mu_2$$

If one wants to include a second rate for both processes, it is possible to add 2 or 3 parameters (e.g. $\nu_1 = \nu_1^* \; ; \; \nu_2 \; ; \; \nu_2^* = \lambda_\nu \nu_2$), depending on whether one wants to have

$\lambda_\mu = \lambda_\nu$ or not. Interestingly, unless one is specifically interested in positive induction, there is no need to restrict $\lambda_{(\mu,\nu)} > 1$, as $\lambda_{(\mu,\nu)} < 1$ would result in a scenario where occurrences of E_1 would slow down the second evolutionary process.

Reciprocal induction. Here, we consider a symmetric interaction between both processes. E_1 events enhance the occurrence rate of E_2 events and, reciprocally, E_2 events enhance the rate of occurrence of E_1 events. This scenario mimics for example reciprocal sign epistasis, where two mutations are beneficial only in the double mutant (Weinreich et al., 2005; Poelwijk et al., 2007).

$$\mu_1 \; ; \; \mu_1^* = \lambda_1 \mu_1 \; ; \; \mu_2 \; ; \; \mu_2^* = \lambda_2 \mu_2$$

Several variants of this scenario also exist depending on whether $\lambda_1 = \lambda_2$ or not, where the λ are restricted to be larger than 1 or not, and finally whether the μ rates differ from the ν rates.

4 The likelihood framework

Let E_1 and E_2 be two types of events whose occurrences are distributed on a tree T, whose topology and branch lengths are known. We compute the likelihood function as the probability of the positions of the occurrences of both events on T, under the model previously described, conditioned on a set of parameters $(\mu_1, \mu_1^*, \nu_1, \nu_1^*, \mu_2, \mu_2^*, \nu_2, \nu_2^*)$. As described below, we can also estimate the rates of the process at the root as well as the order of all events in the tree.

In our current implementation, we assume that no more than one event of each type can occur on a single branch. In the limit of small occurrence rates, this is an excellent approximation. The derivation of the likelihood for more than one event of each type on a branch becomes cumbersome otherwise.

4.1 The likelihood on a single branch

The likelihood of a branch $\mathcal{B}$ of the tree depends on (i) the branch length, (ii) the occurrence rates at the beginning of the branch and (iii) which event(s) occur on the branch. Considering an initial state of rates (m_1, m_2) and the possible occurrences of E_1 and/or E_2 on the branch, we can determine the final state of rates of the current branch, which will be used as the initial rate for the daughter branches, when they exist. All transitions are deterministic and provided in the Appendix. It is noteworthy to mention

that, in the model described here, both processes can never be at the same time in a starred rate.

The likelihood of the branch $\mathcal{B}$ is calculated through the following cases (formula and derivations are detailed in the Appendix): (1) no event occurred on the branch, (2) one event of type i occurred on the branch, for $i \in [1, 2]$ and finally (3) one event of each type (i followed by j or j followed by i) occurred on the branch. In the later case, where E_1 and E_2 occurred on $\mathcal{B}$, we cannot assume any order between them, as both orders ($E_1 \rightarrow E_2$) and ($E_2 \rightarrow E_1$) are possible *a priori*; furthermore, the end rate of both processes after the branch depends on this order. We solve this issue by computing the likelihood of the branch for both possible orders (see below).

4.2 The likelihood on the whole tree

The initial rate state at the root is either (μ_1, μ_2), (μ_1^*, μ_2), or (μ_1, μ_2^*). All other pairs are either impossible (e.g. (μ_1^*, μ_2^*)) or equal up to a swap between μ with ν. Indeed μ and ν can be swapped without any loss of generality (e.g. (μ_1, ν_2^*) is the same as (μ_1, μ_2^*) in the model). This remark is useful to limit the computing time, and has no consequence, since the μ rates and the ν rates play a completely symmetric role in our model. We calculate separately the likelihoods considering the 3 possible initial states, using the following algorithm:

- We know the initial rates of both processes on a branch (inherited from ancestor branch or given by initialization if root edge).

- We check the occurrence of zero, one or both events on the concerned branch and calculate its likelihood and the rates of both processes at the end of the branch. If the branch has both types of events, we calculate separately the likelihood for the two sub-cases and propagate them in parallel with their two respective final rates.

- If the branch is not terminal, we apply this algorithm to its daughter branches, transmitting the final rate. In the case of both events in the branch, we transmit both possible final rates separately to two different paths of the recursion.

The likelihood of the whole tree is the product of the likelihoods of its branches. The number of full-tree likelihoods that must be computed scales with 2^{b_2}, where b_2 is the number of branches with both types of events. These two-event branches are the main limitation of the method, in terms of computation time.

4.3 Estimating the maximum likelihood (ML)

The set of 8 parameters $(\mu_1, \mu_1^*, \nu_1, \nu_1^*, \mu_2, \mu_2^*, \nu_2, \nu_2^*)$ maximizing the likelihood describes the most likely scenario leading to the observed joint distribution of the occurrences of E_1 and E_2 on the tree T under our model of correlated evolution. The 8-uple maximizing the likelihood is searched by a dedicated implementation based on a mix of (a) sequential unidimensional gradient ascent (each dimension is optimized successively), (b) multidimensional gradient ascent, both using golden section search and (c) a Newton-Raphson algorithm when possible (when the curvature is negative). The implementation has been optimized for the full model but also for the nested submodels.

Each possible sequence of ordered events from the root to the leaves of the tree is evaluated and the most likely is returned. As an option, we also estimate which one of the three possible initial rates is more likely (maximizing the likelihood for each of them independently); by default we assume that it is (μ_1, μ_2) and we disregard the others. As a result, the method returns the ordered events, the initial rates (optionally) and the set of parameters maximizing the likelihood.

4.3.1 Scenario selection

The likelihood can also be maximized for submodels by constraining some of the parameters, reducing the parameter space and shortening computation time : for example, setting $\mu_1 = \mu_1^*$ and $\nu_1 = \nu_1^*$ is equivalent to assuming that the occurrence rate of E_1 is not influenced by the occurrences of E_2. Constraining one or more parameters to a fixed value, or to be equal, can define a nested sequence of models, each being a particular case of a more general one (the most general having 8 free parameters). We compare nested models through Likelihood-Ratio Tests (LRT) (Neyman and Pearson, 1933). For example, when considering only two nested models, we can use the fact that twice the difference of their maximum log-likelihoods approximately follows a χ^2 distribution with degrees of freedom equal to the difference in numbers of parameters: $2\ln(\mathcal{L}_a^{\max}/\mathcal{L}_b^{\max}) \sim \chi^2(a - b)$. Then by comparing $2\ln(\mathcal{L}_a^{\max}/\mathcal{L}_b^{\max})$ to the quantiles of $\chi^2(a - b)$, we can decide to reject or not the model with more parameters. Within the specifications of the model described in this article, we have specified different, simple scenarios, some of them being nested into each other.

Cases of particular interest involves submodels without correlated evolution. There are at least two such models in our framework. The most simplistic one has only 2 parameters: μ_1 and μ_2 (or even a single parameter $\mu = \mu_1 = \mu_2$). A more flexible version of independence allows for intrinsic rate changes and is modeled by 4 parameters: μ_1,

μ_2, ν_1 and ν_2. Interestingly, one can also test the relevance of intrinsic rate variations by comparing these two models of independence. Then, one can test for correlated evolution by measuring (and statistically testing) the likelihood improvement with extra parameters, letting the starred rates being different from the basal ones. Although we have not come up with a generalist decision tree using a standard series LRT with a growing number of parameters, we recommend to compute the ML for several scenarios and check which parameters significantly improve the likelihood (as computation time is short enough). This exploration of the different scenarios that are embedded in the general framework (that has at most 8 parameters) will provide an excellent guide to select which scenario is the most parsimonious for the two evolutionary processes under study.

Implementation   The inference program, *epocs*, has been implement in C language and is available at `http://bioinfo.mnhn.fr/abi/public/epocs/`. Simulations were performed using a home made simulator written in Ocaml language, available upon request.

5 Results

5.1 Simulated data

We first assessed the power of the method on simulated data. Simulations give an overview of the theoretical power and the limitations of the method. Through different controlled scenarios, we evaluate (a) to what extent the method correctly estimates the occurrence rates (b) its power to reject independence and (c) its ability to find the strength and nature of the interactions between the two evolutionary processes.

5.1.1 Estimating the occurrence rates

We first tested the accuracy of the method to retrieve occurrence rates that have been arbitrarily selected for a series of simulations. We simulated $10,000$ replicates of a scenario where $\mu_1 = 5$, $\mu_2 = 4$, $\mu_1^* = 20$ and $\mu_2^* = 50$ (all ν were equal to the μ). The input tree is perfectly symmetric (balanced) with 6 synchronized series of bifurcations, resulting in 64 leaves; all branches have the same length. We also used different types of trees (a perfectly imbalanced tree, called caterpillar tree, with 64 leaves and the phylogeny of *enterobacteria* used below) but as results were similar (data not shown), we decided to only report results for the symmetric tree for the sake of conciseness. As rates are rescaled by total tree length, $\mu_1 = 5$ implies that there are on average 5 occurrences of E_1 when the process is at its basal rate (*i.e.* not counting the ones that are triggered by occurrences of E_2). We applied the maximum likelihood method to each simulated replicate generating a mapping of events E_1 and E_2 on the tree, in order to estimate the four rates (the ν were equal to the μ in these simulations). When more than one event of a kind occurred in the branch, only one was kept.

Figure 2 shows the distribution of estimated rates. We observe that the distribution of all four rates shows a peak typically at the true values, demonstrating that our framework is indeed able to correctly estimate the rates. The dispersion of estimates around the true values mostly comes from the stochasticity of the simulations: since the events are distributed according to Poisson processes, in some cases we can get more or less occurrences than the average corresponding to the true rates. This also explains why we observe peaks at low rates that correspond to 1 or 2 occurrences. As the current implementation does not estimate rates larger than 1000, the last bin (1000+) encompasses all large rates.

The means ($\pm$ stdev) of the estimated values are $\bar{\mu}_1 = 4.8$ (± 2.4) and $\bar{\mu}_2 = 3.9$ (± 2.1) while for the higher rates, $\bar{\mu}_1^* = 28.4$ (± 58.2) and $\bar{\mu}_2^* = 80.1$ (± 74.0). This indicates that

estimations of basal rates are unbiased whereas estimations of induced rates with high values may be biased upward, although their distributions show a clear mode centered on the true values.

5.1.2 Power to detect induction

We further assessed the power of likelihood-ratio tests (LRTs) to detect an induction from E_1 on E_2 and compared them to the power of our previously published method based on counts in the phylogeny (Behdenna et al., 2016). The power is estimated by the fraction of replicates that reject significantly the null model of independence.

First, we simulated a scenario of asymmetric induction of E_1 on E_2 using 3 distinct parameters: μ_1, μ_2 and μ_2^*, with an induction rate ($\lambda = \mu_2^*/\mu_2$) varying from 1 (no induction) to 1000 (very strong induction); all ν values are equal to the corresponding μ. Two LRTs were computed, comparing an independence model (H_0, 2 parameters: μ_1 and μ_2) to the simulated model (H_1, 3 parameters: μ_1, μ_2 and μ_2^*) or to an over-parameterized model with inductions in both directions (H_2, 4 parameters: μ_1, μ_2, μ_1^* and μ_2^*). P-values were computed assuming a χ^2 distribution for twice the log of maximum likelihood ratios, with either 1 (H_1 vs H_0) or 2 (H_2 vs H_0) degrees of freedom. We also report the power of two other tests that assess the probability of independence using either a) counts of pairs of E_1 and E_2 events that co-occur in the same branch (referred to as "epics Id" in reference to Behdenna et al. 2016) or b) counts of pairs of events that co-occur plus counts of E_2 events following E_1 events in the tree ("epics S+Id"). Results (Figure 3a) show that both LRTs outperform our previous tests based solely on counts and that the LRT based on the over-parameterized model is slightly less powerful. For inductions such that $\lambda > 10$, LRTs show a very good power ($> 70\%$) that reaches $\sim 100\%$ for inductions $\lambda > 100$.

Second, we simulated a scenario of symmetric (reciprocal) induction between E_1 and E_2 using 3 distinct parameters: μ_1, μ_2 and $\lambda = \mu_1^*/\mu_1 = \mu_2^*/\mu_2$, with $\lambda \in [1, 1000]$. Two other LRTs were computed, comparing an independence model (H_0, 2 parameters: μ_1 and μ_2) to the simulated model (H_1, 3 parameters: μ_1, μ_2 and λ) or to an over-parameterized model with inductions in both directions but of different strengths (H_2, 4 parameters: μ_1, μ_2, μ_1^* and μ_2^*). Results (Figure 3b) show that again both LRTs outperform our previous tests based solely on counts, with an even stronger power that reaches $\sim 100\%$ for inductions $\lambda > 10$.

5.1.3 Power to detect induction and/or irreversible loss

We next assessed the power of LRT to discriminate between the effect of *interactions between the events* (modeled by the induced rates μ^*) and *intrinsic rate changes* (modeled by the ν rates). More specifically, we designed three different scenarios where E_1 events tend to precede E_2 events:

1. **Induction**: E_1 events favors E_2 events ($\lambda_2 > 1$), while E_2 events slow down E_1 events ($\lambda_1 < 1$). This is a special case of asymmetric induction of E_1 on E_2.

2. **Loss**: E_1 events have a higher rate of occurrence ($\mu_1 > \mu_2$) and both events can occur only once ($\nu_1 = \nu_2 = 0$). In this scenario, there is no interaction between the two processes and the sequential pattern (E_1 precedes E_2) is only due to their respective intrinsic rates. A biological example would be the irreversible loss of two functions, one being lost faster.

3. **Induction+Loss**: irreversible loss ($\nu = 0$) plus asymmetric induction ($\lambda_1 < 1$, $\lambda_2 > 1$).

The exact rates that we used in the simulations are indicated in Table 1.

Scenario	μ_1	μ_1^*	μ_2	μ_2^*	ν
Induction	10	5	10	$[5, 5000]$	μ
Loss	$[5, 5000]$	μ_1	5	5	0
Induction+Loss	25	5	25	$[5, 5000]$	0

Table 1: *Rates used for the simulations*

To assess the presence of induction and/or loss, we computed 4 LRTs, that compared 4 pairs of nested models illustrated in Figure 4a. LRT_1 and LRT_4 test whether adding a single parameter $\lambda_2 = \mu_2^*/\mu_2$ improves significantly the likelihood, demonstrating an induction of E_1 on E_2. LRT_2 and LRT_3 test whether the addition of 2 parameters (ν_1 and ν_2) significantly improves the likelihood, demonstrating that rate changes occur regardless of the interactions. When none of the LRTs is significant, a scenario of no induction+no loss (referred to as H_0) is inferred. When LRT_1 and LRT_4 are the only two significant ones, a scenario of induction+no loss is inferred. When LRT_2 and LRT_3 are the only two significant ones, a scenario of loss+no-induction is inferred. When 3 LRTs are significant, a scenario of induction+loss is said "likely". Finally when all LRTs are significant, a scenario of induction+loss is inferred.

We first simulated a scenario of asymmetric induction. Results (Figure 4B) show that a scenario of induction+no loss is correctly preferred, provided that λ_2 is large enough ($\lambda_2 > 10$). Second, we simulated a scenario of loss+no induction: the correct scenario is inferred (Figure 4C), provided that μ_1 is large enough, *i.e.*, E_1 events must occur rapidly and only once (followed by E_2 events, that occur also once but later). Finally, we simulated a case with irreversible loss+induction. Evidence supporting both rate changes (intrinsic plus interaction) is higher as the induction of E_1 on E_2 gets larger (Figure 4D). When the induction is not very strong, the method only perceives the effect of loss.

From this set of simulations, we conclude that model selection based on LRT can be used to assess the presence of interactions between the two types of events as well as intrinsic rate changes that are independent of any interaction.

Analysis of rRNA 16S nucleotides

The ribosomal ribonucleic acid (rRNA) is a major component of the ribosome. rRNA sequences, already in the genome of the last universal common ancestor, are very well conserved and therefore have been used to show the diversity of living organisms, in particular the existence of the three kingdoms of life (Fox and Woese, 1975; Woese and Fox, 1977). The 3D structure of RNA has been resolved experimentally (Doty et al., 1959; Leontis and Westhof, 2001), and its folding reveals physical interactions between nucleotides. In particular, the single-stranded structure of rRNA is in a large part determined by Watson-Crick pairs (hereafter WC pairs) that form stems. As the RNA structure is strongly stabilized by stems of WC pairs, the mutation of one nucleotide involved in a WC pair is very often deleterious and is typically only observed together

with a second compensatory mutation on the paired nucleotide. WC pairs constitute a classical case of correlated evolution at the molecular level (Chiu and Kolodziejczak, 1991; Leontis and Westhof, 1998; Moore, 1999). As the interactions between both nucleotides of a pair are very strong, we expect a very short time lag, if any, between the two substitutions at paired bases.

We report all pairs of nucleotides that show significant correlated evolution in the RNA 16S gene in a sample of 54 gamma-enterobacteria (plus one beta-enterobacterium as outgroup). We retrieved the sequences from the SILVA rRNA database (Pruesse et al., 2007; Quast et al., 2013), aligned them using Muscle v3.6 (Edgar, 2004), and trimmed the poorly aligned positions (*i.e.*, too divergent regions) of the alignment using Gblocks v0.91b with default parameter values (Castresana, 2000; Talavera and Castresana, 2007). From the resulting 1,233 nucleotide alignment, we inferred a phylogenetic tree, using Phyml (Guindon et al., 2010) with a generalized time reversible (GTR) substitution model; finally, we inferred the ancestral states by maximum likelihood with Bayestraits Multistate (Pagel et al., 2004). At each site, for each branch, we assimilated any inferred substitutions to an evolutionary event, disregarding the exact nature of the substitution. In the *E. coli* rRNA 16S structure, that we assume is conserved for all enterobacterial rRNAs of our datasets, there is a total of 477 WC pairs.

The list of the 55 species, the filtered alignment and the input tree (with or without events) are provided as Supplementary Material.

WC pairs show evidence of correlated evolution

We used rRNA WC pairs as a positive control to evaluate the performance of two tests for independence: LRT_{23} compares the maximum likelihood estimate of a 2-parameter model with no induction (μ_1 and μ_2) to a 3-parameter model with reciprocal induction (μ_1, μ_2 and $\lambda = \mu_1^*/\mu_1 = \mu_2^*/\mu_2$). LRT_{45} compares the maximum likelihood estimate of a 4-parameter model with no induction (μ_1, μ_2, ν_1 and ν_2) to a 5-parameter model with reciprocal induction (same plus $\lambda = \mu_1^*/\mu_1 = \mu_2^*/\mu_2 = \nu_1^*/\nu_1 = \nu_2^*/\nu_2$). The rationale is to check whether including switches of intrinsic rates improves the performance of LRTs. We also report the performance of our previous method based on counts of co-occurring events (epics-Id). The comparison between the pairs that are significantly coevolving and the documented WC pairs shows an overall good overlap between them (Table 2). The stronger the evidence of correlated evolution (*i.e.*, the smaller the p-values), the higher the enrichment of WC pairs.

We however note that a large fraction of the 477 WC pairs show no evidence of co-evolution, regardless of the method. At least two reasons can be put forward. First,

	epics-Id		LRT$_{23}$		LRT$_{45}$	
pval	N	WC	N	WC	N	WC
0.05	3,006	74	9,464	89	6,850	86
10^{-2}	698	57	2,940	74	2,477	75
10^{-3}	101	40	752	61	426	59
10^{-4}	39	29	266	50	118	45
10^{-5}	22	20	68	39	49	37
10^{-6}	12	11	37	34	30	28

Table 2: *For each method, N indicates the number of pairs with evidence of correlated evolution, for the risk given in the first column; WC is the number of Watson-Crick pairs among them.*

we have filtered the alignment with Gblocks. Out of the complete set of 477 WC pairs, only 349 WC pairs (0.05% of the 759,528 unmasked possible nucleotide pairs) have both their positions unmasked after Gblocks. Furthermore, any site that is monomorphic on the alignment cannot show a pattern of correlated evolution in the tree. The alignment contains 422 polymorphic sites. This reduces the dataset of WC pairs that could be detected by the method to a maximum of 106 WC pairs, among a total of 88,831 polymorphic possible nucleotide pairs. Most of these 106 WC pairs show evidence of correlated evolution for a risk of 5%, but only ~ 30 very strong evidence of correlated evolution (p-value $< 10^{-6}$). Finally, adding variable intrinsic rates (allowing ν rates to differ from μ rates) improves the specificity of LRTs (at the cost of only a marginal loss of sensitivity), so we only used this LRT$_{45}$ in the following and simply referred to it as LRT.

To better understand what type of WC pairs show a significant pattern of correlated evolution, we computed the induction rate λ of all 106 WC pairs, regardless of the significance of the LRT. Results (Figure 5) clearly demonstrate that the 75 WC pairs with decent support for correlated evolution (p-value $< 1\%$) have stronger induction rate than the others. Any WC pair with an induction rate of $\lambda > 100$ has significant support for correlated evolution at this risk.

What other nucleotide pairs show evidence for correlated evolution?

We assessed the presence of correlated evolution using a selection of four different methods:

- the LRT$_{45}$ described above: *LRT*,

- our previous method based on counts of co-occurring events: *epics-Id* (Behdenna et al., 2016),

- another method based on co-occurrences: *CoMap* (Dutheil et al., 2005),

- a method based on Mutual Information "corrected" for the phylogeny: *MIp* (Gloor et al., 2005).

We ranked the 75 "best" nucleotide pairs, that show the strongest statistical evidence for correlated evolution (disregarding the significance of p-values). We then used the 3D positions of the nucleotides in the crystal structure of the 16S ribosome (Korostelev et al., 2006) to compute the distance (in Å) between each pair of nucleotides, taking the barycenter of the base as its position in space. Each pair was then tagged depending on whether it was a WC pair, a non-WC pair with both nucleotides close in space (< 10Å) or a distant pair. Results (Figure 6) show that the LRT method is marginally more specific at detecting WC pairs, while epics-Id was also able to detect significant correlated evolution among close nucleotides that are not WC pairs.

To further characterize the non-WC nucleotide pairs that show patterns of correlated evolution, we remapped them on the secondary structure: stems of WC pairs and associated loops. Results (Figure 6b) show that several non-WC pairs close in space belong to the same stem and are either consecutive (positions k and $k+1$, that are stacked nucleotides) or shifted by one when compared to WC pairs. This suggests that alternative non-WC interactions also constrain the evolution of the rRNA 16S.

6   Discussion

In this study, we have developed a likelihood framework based on a minimal model of correlated evolution between two evolutionary processes that generate discrete events on a given phylogenetic tree. This model depends on at most 8 parameters, 4 for each process, that represent *interactions* between the two processes (the starred rates) and intrinsic rate switches ($\nu \neq \mu$). For a given tree with evolutionary events mapped on it, we estimate the parameters by maximum likelihood for any (sub)model defined with (a subset of) the 8 parameters. Maximizing this probability is equivalent to searching for an optimal set of parameters describing the processes that led to the observed distribution of the events on the tree. Nested models also permit the direct comparison of scenarios

using likelihood ratio tests (LRTs), in particular to assess the support for interaction and/or intrinsic rate switches.

Using simulated controlled scenarios, we have shown that our method is accurate for estimating the occurrence rates and has good power to assess the presence of interactions despite the use of limited data (symmetric trees with only 64 leaves). Moreover, we show that the method can assess correctly which process can lead to ordered pairs of events: differential rates with irreversible loss or interaction between the two processes. Finally, based on an analysis of nucleotide pairs in an alignment of 16S sampled in only 52 enterobacteria, we show that the method detects correlated evolution in many polymorphic WC pairs, especially among the ones that exhibit strong reciprocal induction. In addition we show that several pairs that are not WC but in the same stem also show evidence of correlated evolution. Comparison of performance with a selection of three previously existing methods (our previous one, and two among the popular ones, that are representative of the different approaches to detect correlated evolution) demonstrates that the LRT based on the current framework is more specific at detecting WC pairs but, more importantly, can also estimate the strength of their reciprocal induction.

As the model has a small number of parameters, the computation time of likelihood maximization is fast enough to analyze correlated evolution in a very large number of pairs. In the 16S analysis, we had to maximize 2,304,091 likelihood functions (finding all possible orders of events for all pairs). The total computation time on a 4Ghz laptop, without any parallelization, is 15 min for the 2-parameter model of independence (μ_1 and μ_2), 110 min ($\approx 2h$) for the 4-parameter model of independence, and at most 306 min ($\approx 5h$) for the full 8-parameter model (that was not used in the analysis above). This suggests that MCMC Bayesian surface exploration, that typically requires millions of likelihood estimations, could be run to analyze cases on a dataset of a "reasonable" size. Computation time remains, however, orders of magnitude larger than in the methods based on counts we have developed before (Behdenna et al., 2016), which takes only 5 sec to compute correlated evolution from counts of co-occurrence.

The framework described here relates to the one proposed by Pagel (1994), as they both rely on likelihood computation of an explicit continuous time Markov chain. In both frameworks, the independence model is nested in a subspace of the more general model of correlated evolution and statistical tests such as LRT can be done to assess the statistical support for independence. However, our minimal framework differs from the one of Pagel (1994) in several important aspects. First, and this is a major difference, our model has at most 8 parameters, whereas Pagel (1994) computes the transition rates between all possible pairs of states for the two traits. In the case of two binary traits,

Pagel (1994) has 8 parameters but it has hundreds for sites with 20 amino acids. In the case of two binary traits, both frameworks are identical if we restrict our minimal framework to the 4-parameter model of independence that is the null model of Pagel (1994) and the largest 8-parameter model that corresponds to the full model of Pagel (1994). However, our minimal model is flexible enough that one can test for likelihood improvement due to the addition of intrinsic rates changes ($\nu \neq \mu$) and/or addition of interaction between the processes ($\mu^* \neq \mu$). More generally, our minimal model constitutes a powerful tool to assess the statistical support for any scenario that is defined within the framework, ignoring extra unnecessary parameters. An obvious and direct drawback is that processes which occurrence rates are tuned by more than 2 states (e.g. for a given site, many amino acid transitions occur in the input tree with "very" different rates). In this regard, our minimal framework is most relevant when only a few *different* types of transitions occur in the tree. Deep trees saturated with evolutionary events should be avoided. Finally, the comparison of the various rates of our minimal model is straightforward as they have an explicit biological interpretation (e.g., computing the λ immediately tells whether the interaction is an induction of a repression).

The framework developed here takes as input a phylogenetic tree and a set of events placed on the tree. Like any method based on phylogenies, it will suffer from uncertainties from the phylogenetic reconstruction. Typically, trees estimated on very close sequences or on very divergent ones should be handled with care. As the method relies on branch lengths to estimate the rates of the process, one has to be careful while using a tree as input, especially on deep branches that are specifically subject to uncertainties. Furthermore, as our method assumes that evolutionary events are correctly placed on the tree, we also recommend great care in the reconstruction of the ancestral state. When possible, likelihood based methods (such as ML or Bayesian methods) should be preferred over parsimony; we nonetheless note that when there are only few events on the tree, all methods will likely perform similarly. One possible extension of our framework would be to integrate over all possible ancestral reconstructions. Finally, we also stress that we only consider at most one event of each kind on each branch. Altogether, we suggest again that the method presented here is best suited for phylogenies that are not too deep and where the evolutionary events of interest are sparsely located.

One interesting feature of the framework presented here is its inherent flexibility that allows the analysis of diverse types of data. Regardless of its nature, any discrete evolutionary event that is mapped on a phylogenetic tree can be treated, whether it is a molecular (*e.g.*, substitution) or a non-molecular (*e.g.*, gain or loss of a character or of

a biological function) event.

As a concluding remark, we would like to mention that GWAS (Genome Wide Association) studies can be seen as computing correlated evolution between genomic variants and a phenotypic trait of interest (Achaz and Dutheil, 2020). As any method of correlated evolution, GWA studies also suffer from phylogenetic inertia, via what is called stratification of the population into subpopulations (Price et al., 2006). For bacterial GWAS, it has been suggested to fully include the tree for the detection of a positive correlation of genomic variants with the phenotype of interest (Collins and Didelot, 2018). This suggests that the framework presented here, that can easily assess and quantify correlated evolution between genetic variants and discrete changes of a trait, can also be used to perform GWA-like studies.

7   Ackowledgments

The authors want to thank EPC Rocha and R Kulathinal for their critical reading and helpful comments on the manuscript as well as O Tenaillon for fruitful discussions on the work. This work was supported by a state grant from the *Agence Nationale de la Recherche* through the *Investissements d'Avenir* program ANR-16-CONV-0005. AB, PP, AL and GA thank the Center for Interdisciplinary Research in Biology (CIRB, Collège de France) for funding.

References

Achaz, G. and J. Dutheil. 2020. Correlated evolution: models and methods. xx xx vol. xx. ISTE.

Achaz, G., A. Rodriguez-Verdugo, B. S. Gaut, and O. Tenaillon. 2014. The reproducibility of adaptation in the light of experimental evolution with whole genome sequencing. Adv Exp Med Biol 781:211–31.

Baldassi, C., M. Zamparo, C. Feinauer, A. Procaccini, R. Zecchina, M. Weigt, and A. Pagnani. 2014. Fast and accurate multivariate Gaussian modeling of protein families: predicting residue contacts and protein-interaction partners. PLoS ONE 9:e92721.

Bateson, W. 1909. Heredity and variation in modern lights. chap. III, Pages 87–110 *in* Danvinand Modmz Science (A. Seward, ed.) gutenberg projet http://www.gutenberg.org/files/22430/22430-h/22430-h.htm ed. Cambridge University Press.

Baum, D. A. and M. J. Donoghue. 2001. A likelihood framework for the phylogenetic analysis of adaptation. Pages 24–44 *in* Adaptation and Optimality (S. Orzack and E. Sober, eds.). Cambridge University Press, New York.

Behdenna, A., J. Pothier, S. S. Abby, A. Lambert, and G. Achaz. 2016. Testing for Independence between Evolutionary Processes. Syst. Biol. 65:812–823.

Bitbol, A.-F. 2018. Inferring interaction partners from protein sequences using mutual information. PLOS Computational Biology 14:e1006401 publisher: Public Library of Science.

Castresana, J. 2000. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. Mol Biol Evol 17:540–52.

Chiu, D. K. and T. Kolodziejczak. 1991. Inferring consensus structure from nucleic acid sequences. Comput. Appl. Biosci. 7:347–352.

Collins, C. and X. Didelot. 2018. A phylogenetic method to perform genome-wide association studies in microbes that accounts for population structure and recombination. PLOS Computational Biology 14:e1005958 publisher: Public Library of Science.

Dib, L., D. Silvestro, and N. Salamin. 2014. Evolutionary footprint of coevolving positions in genes. Bioinformatics 30:1241–9.

Dobzhansky, T. 1934. Studies on hybrid sterility. Zeitschrift für Zellforschung und Mikroskopische Anatomie 21:169–223.

Doty, P., H. Boedtker, J. R. Fresco, R. Haselkorn, and M. Litt. 1959. Secondary structure in ribonucleic acids. Proc Natl Acad Sci U S A 45:482–99.

Dutheil, J. and N. Galtier. 2007. Detecting groups of coevolving positions in a molecule: a clustering approach. BMC Evol Biol 7:242.

Dutheil, J., T. Pupko, A. Jean-Marie, and N. Galtier. 2005. A model-based approach for detecting coevolving positions in a molecule. Mol Biol Evol 22:1919–28.

Edgar, R. C. 2004. Muscle: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Res 32:1792–7.

Ekeberg, M., C. Lövkvist, Y. Lan, M. Weigt, and E. Aurell. 2013. Improved contact prediction in proteins: using pseudolikelihoods to infer Potts models. Phys Rev E Stat Nonlin Soft Matter Phys 87:012707.

Felsenstein, J. 1981. Evolutionary trees from dna sequences: a maximum likelihood approach. J Mol Evol 17:368–76.

Felsenstein, J. 1985. Phylogenies and the comparative method. The American Naturalist 125:1–15.

Fox, G. E. and C. R. Woese. 1975. 5s rna secondary structure. Nature 256:505–7.

Fraser, H. B., A. E. Hirsh, D. P. Wall, and M. B. Eisen. 2004. Coevolution of gene expression among interacting proteins. Proceedings of the National Academy of Sciences of the United States of America 101:9033–9038.

Gloor, G. B., L. C. Martin, L. M. Wahl, and S. D. Dunn. 2005. Mutual information in protein multiple sequence alignments reveals two classes of coevolving positions. Biochemistry 44:7156–65.

Grafen, A. 1989. The phylogenetic regression. Philos. Trans. R. Soc. Lond., B, Biol. Sci. 326:119–157.

Guindon, S., J.-F. Dufayard, V. Lefort, M. Anisimova, W. Hordijk, and O. Gascuel. 2010. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of phyml 3.0. Syst Biol 59:307–21.

Harvey, P. H. and M. D. Pagel. 1991. The Comparative Method In Evolutionary Biology. Oxford University Press, U.S.A., Oxford ; New York.

Kondrashov, A. S., S. Sunyaev, and F. A. Kondrashov. 2002. Dobzhansky-muller incompatibilities in protein evolution. Proc Natl Acad Sci U S A 99:14878–83.

Korostelev, A., S. Trakhanov, M. Laurberg, and H. F. Noller. 2006. Crystal structure of a 70s ribosome-trna complex reveals functional interactions and rearrangements. Cell 126:1065–1077.

Kryazhimskiy, S., J. Dushoff, G. A. Bazykin, and J. B. Plotkin. 2011. Prevalence of epistasis in the evolution of influenza A surface proteins. PLoS Genet. 7:e1001301.

Kulathinal, R. J., B. R. Bettencourt, and D. L. Hartl. 2004. Compensated deleterious mutations in insect genomes. Science 306:1553–4.

Leontis, N. B. and E. Westhof. 1998. A common motif organizes the structure of multi-helix loops in 16 s and 23 s ribosomal rnas. Journal of molecular biology 283:571–583.

Leontis, N. B. and E. Westhof. 2001. Geometric nomenclature and classification of rna base pairs. RNA 7:499–512.

Marks, D. S., L. J. Colwell, R. Sheridan, T. A. Hopf, A. Pagnani, R. Zecchina, and C. Sander. 2011. Protein 3D Structure Computed from Evolutionary Sequence Variation. PLOS ONE 6:e28766 publisher: Public Library of Science.

Martin, L. C., G. B. Gloor, S. D. Dunn, and L. M. Wahl. 2005. Using information theory to search for co-evolving residues in proteins. Bioinformatics 21:4116–24.

Martins, E. P. 1994. Estimating the Rate of Phenotypic Evolution from Comparative Data. The American Naturalist 144:193–209 publisher: [University of Chicago Press, American Society of Naturalists].

Martins, E. P. and T. F. Hansen. 1997. Phylogenies and the Comparative Method: A General Approach to Incorporating Phylogenetic Information into the Analysis of Interspecific Data. The American Naturalist 149:646–667 publisher: [University of Chicago Press, American Society of Naturalists].

Milligan, B. 1994. Estimating evolutionary rates for discrete characters. Pages 299–311 *in* Models in phylogeny reconstruction (R. W. Scotland, D. J. Siebert, and D. M. Williams, eds.) vol. Systematics Association Special Volume Number 52. Clarendon, Oxford, UK.

Moore, P. B. 1999. Structural motifs in rna. Annual review of biochemistry 68:287–300.

Morcos, F., A. Pagnani, B. Lunt, A. Bertolino, D. S. Marks, C. Sander, R. Zecchina, J. N. Onuchic, T. Hwa, and M. Weigt. 2011. Direct-coupling analysis of residue coevolution captures native contacts across many protein families. Proc. Natl. Acad. Sci. U.S.A. 108:E1293–1301.

Muller, H. 1942. Isolating mechanisms, evolution, and temperature. Biology Symposium 6:71–125.

Neyman, J. and E. Pearson. 1933. On the problems of the most efficient tests of statistical hypotheses. Philosophical Transactions of the Royal Society of London .

Orr, H. A. 1996. Dobzhansky, bateson, and the genetics of speciation. Genetics 144:1331–5.

Pagel, M. 1994. Detecting correlated evolution on phylogenies: a general method for the comparative analysis of discrete characters. Proceedings of the Royal Society of London B: Biological Sciences 255:37–45.

Pagel, M. and A. Meade. 2006. Bayesian analysis of correlated evolution of discrete characters by reversible-jump markov chain monte carlo. Am Nat 167:808–25.

Pagel, M., A. Meade, and D. Barker. 2004. Bayesian estimation of ancestral character states on phylogenies. Syst Biol 53:673–84.

Pensar, J., S. Puranen, B. Arnold, N. MacAlasdair, J. Kuronen, G. Tonkin-Hill, M. Pesonen, Y. Xu, A. Sipola, L. Sánchez-Busó, J. A. Lees, C. Chewapreecha, S. D. Bentley, S. R. Harris, J. Parkhill, N. J. Croucher, and J. Corander. 2019. Genome-wide epistasis and co-selection study using mutual information. Nucleic Acids Res 47:e112–e112 publisher: Oxford Academic.

Phillips, P. C. 2008. Epistasis–the essential role of gene interactions in the structure and evolution of genetic systems. Nat Rev Genet 9:855–67.

Poelwijk, F. J., D. J. Kiviet, D. M. Weinreich, and S. J. Tans. 2007. Empirical fitness landscapes reveal accessible evolutionary paths. Nature 445:383–6.

Pollock, D. D., W. R. Taylor, and N. Goldman. 1999. Coevolving protein residues: maximum likelihood identification and relationship to structure. J Mol Biol 287:187–98.

Price, A. L., N. J. Patterson, R. M. Plenge, M. E. Weinblatt, N. A. Shadick, and D. Reich. 2006. Principal components analysis corrects for stratification in genome-wide association studies. Nature Genetics 38:904–909 number: 8 Publisher: Nature Publishing Group.

Pruesse, E., C. Quast, K. Knittel, B. M. Fuchs, W. Ludwig, J. Peplies, and F. O. Glöckner. 2007. Silva: a comprehensive online resource for quality checked and aligned ribosomal rna sequence data compatible with arb. Nucleic Acids Res 35:7188–96.

Quast, C., E. Pruesse, P. Yilmaz, J. Gerken, T. Schweer, P. Yarza, J. Peplies, and F. O. Glöckner. 2013. The silva ribosomal rna gene database project: improved data processing and web-based tools. Nucleic Acids Res 41:D590–6.

Schöniger, M. and A. von Haeseler. 1994. A stochastic model for the evolution of auto-correlated dna sequences. Mol Phylogenet Evol 3:240–7.

Shindyalov, I. N., N. A. Kolchanov, and C. Sander. 1994. Can three-dimensional contacts in protein structures be predicted by analysis of correlated mutations? Protein Engineering, Design and Selection 7:349–358.

Talavera, G. and J. Castresana. 2007. Improvement of phylogenies after removing divergent and ambiguously aligned blocks from protein sequence alignments. Syst Biol 56:564–77.

Tillier, E. R. M. and R. A. Collins. 1995. Neighbor joining and maximum likelihood with rna sequences: Adressing the interdependence of sites. Mol Biol Evol 12:7–15.

Tufféry, P. and P. Darlu. 2000. Exploring a phylogenetic approach for the detection of correlated substitutions in proteins. Mol. Biol. Evol. 17:1753–1759.

Van Valen, L. 1973. A New Evolutionary Law. Evol. Theory 1:1–30.

Visser, J. A. G. M. d. and J. Krug. 2014. Empirical fitness landscapes and the predictability of evolution. Nat Rev Genet 15:480–490 number: 7 Publisher: Nature Publishing Group.

Weigt, M., R. A. White, H. Szurmant, J. A. Hoch, and T. Hwa. 2009. Identification of direct residue contacts in protein–protein interaction by message passing. PNAS 106:67–72 publisher: National Academy of Sciences Section: Physical Sciences.

Weinreich, D. M., R. A. Watson, and L. Chao. 2005. Perspective: Sign epistasis and genetic constraint on evolutionary trajectories. Evolution 59:1165–74.

Welch, J. J. 2004. Accumulating dobzhansky-muller incompatibilities: reconciling theory and data. Evolution 58:1145–56.

Woese, C. R. and G. E. Fox. 1977. Phylogenetic structure of the prokaryotic domain: the primary kingdoms. Proc Natl Acad Sci U S A 74:5088–90.

Wright, S. 1932. The roles of mutation, inbreeding, crossbreeding and selection in evolution. Pages 356–366 *in* Proceedings of the Sixth International Congress on Genetics vol. 1.

Yeang, C.-H., J. Darot, H. Noller, and D. Haussler. 2007. Detecting the coevolution of biosequences an example of rna interaction prediction. Molecular biology and evolution 24:2119–31.

Yi, X. and A. M. Dean. 2019. Adaptive Landscapes in the Age of Synthetic Biology. Mol Biol Evol 36:890–907 publisher: Oxford Academic.

Appendix

A   Initial and final states

The following tables sum up the final rate state corresponding to each initial rate state, for each possible case. Note that the states where both events have a starred occurrence rate are forbidden.

No event on the branch

In this particular case, the final state is always the same as the initial state.

One occurrence of E_i only on the branch

E_1		E_2	
Initial state	Final state	Initial state	Final state
(a_1, a_2)	(b_1, b_2)	(a_1, a_2)	(b_1, b_2)
(μ_1, μ_2)	(ν_1, μ_2^*)	(μ_1, μ_2)	(μ_1^*, ν_2)
(μ_1, μ_2^*)	(ν_1, μ_2^*)	(μ_1, μ_2^*)	(μ_1, ν_2)
(μ_1, ν_2)	(ν_1, ν_2^*)	(μ_1, ν_2)	(μ_1^*, μ_2)
(μ_1, ν_2^*)	(ν_1, ν_2^*)	(μ_1, ν_2^*)	(μ_1, μ_2)
(μ_1^*, μ_2)	(ν_1, μ_2)	(μ_1^*, μ_2)	(μ_1^*, ν_2)
(μ_1^*, ν_2)	(ν_1, ν_2)	(μ_1^*, ν_2)	(μ_1^*, μ_2)
(ν_1, μ_2)	(μ_1, μ_2^*)	(ν_1, μ_2)	(ν_1^*, ν_2)
(ν_1, μ_2^*)	(μ_1, μ_2^*)	(ν_1, μ_2^*)	(ν_1, ν_2)
(ν_1, ν_2)	(μ_1, ν_2^*)	(ν_1, ν_2)	(ν_1^*, μ_2)
(ν_1, ν_2^*)	(μ_1, ν_2^*)	(ν_1, ν_2^*)	(ν_1, μ_2)
(ν_1^*, μ_2)	(μ_1, μ_2)	(ν_1^*, μ_2)	(ν_1^*, ν_2)
(ν_1^*, ν_2)	(μ_1, ν_2)	(ν_1^*, ν_2)	(ν_1^*, μ_2)

Both events are present on the branch

$E_1 \rightarrow E_2$			$E_2 \rightarrow E_1$		
Initial state	Intermediate state	Final state	Initial state	Intermediate state	Final state
(a_1, a_2)	(b_1, b_2)	(c_1, c_2)	(a_1, a_2)	(b_1, b_2)	(c_1, c_2)
(μ_1, μ_2)	(ν_1, μ_2^*)	(ν_1, ν_2)	(μ_1, μ_2)	(μ_1^*, ν_2)	(ν_1, ν_2)
(μ_1, μ_2^*)	(ν_1, μ_2^*)	(ν_1, ν_2)	(μ_1, μ_2^*)	(μ_1, ν_2)	(ν_1, ν_2^*)
(μ_1, ν_2)	(ν_1, ν_2^*)	(ν_1, μ_2)	(μ_1, ν_2)	(μ_1^*, μ_2)	(ν_1, μ_2)
(μ_1, ν_2^*)	(ν_1, ν_2^*)	(ν_1, μ_2)	(μ_1, ν_2^*)	(μ_1, μ_2)	(ν_1, μ_2^*)
(μ_1^*, μ_2)	(ν_1, μ_2)	(ν_1^*, ν_2)	(μ_1^*, μ_2)	(μ_1^*, ν_2)	(ν_1, ν_2)
(μ_1^*, ν_2)	(ν_1, ν_2)	(ν_1^*, μ_2)	(μ_1^*, ν_2)	(μ_1, μ_2)	(ν_1, μ_2)
(ν_1, μ_2)	(μ_1, μ_2^*)	(μ_1, ν_2)	(ν_1, μ_2)	(ν_1^*, ν_2)	(μ_1, ν_2)
(ν_1, μ_2^*)	(μ_1, μ_2^*)	(μ_1, ν_2)	(ν_1, μ_2^*)	(ν_1, ν_2)	(μ_1, ν_2^*)
(ν_1, ν_2)	(μ_1, ν_2^*)	(μ_1, μ_2)	(ν_1, ν_2)	(ν_1^*, μ_2)	(μ_1, μ_2)
(ν_1, ν_2^*)	(μ_1, ν_2^*)	(μ_1, μ_2)	(ν_1, ν_2^*)	(ν_1, μ_2)	(μ_1, μ_2^*)
(ν_1^*, μ_2)	(μ_1, μ_2)	(μ_1^*, ν_2)	(ν_1^*, μ_2)	(ν_1^*, ν_2)	(μ_1, ν_2)
(ν_1^*, ν_2)	(μ_1, ν_2)	(μ_1^*, μ_2)	(ν_1^*, ν_2)	(ν_1, μ_2)	(μ_1, μ_2)

B   Likelihood function for a single branch

Let T be a tree, and $\mathcal{B}$ a branch of this tree, of length l. We consider two events E_i, $i \in \{1, 2\}$, of respective occurrence rates $(\mu_i, \mu_i^*, \nu_i, \nu_i^*)$. Each event is assumed to occur at most once on $\mathcal{B}$.

We detail here the likelihood function for this single branch, the three possible cases being treated separately. Let $f_\mathcal{B}$ be the likelihood for the branch $\mathcal{B}$, given the parameter values.

No event on the branch

Let a_1 (resp. a_2) be the occurrence rate for E_1 (resp. E_2) on the branch.

$$f_\mathcal{B} \quad = e^{-l(a_1 + a_2)}$$

One occurrence of E_i, $i \in \{1, 2\}$ on the branch

In this case, we define:

- a_1 (resp. a_2) as the occurrence rate for E_1 (resp. E_2) before the occurrence of E_i on the branch

- b_1 (resp. b_2) as the occurrence rate for E_1 (resp. E_2) after the occurrence of E_i on the branch.

Those parameters take different values depending on the initial states on the branch, and according to the model. We easily get

$$f_{\mathcal{B}} = \int_0^l a_i e^{-a_1 t} e^{-a_2 t} e^{-b_1(l-t)} e^{-b_2(l-t)} dt$$

$$= -\frac{a_i}{a_1+a_2-b_1-b_2}(e^{-l(a_1+a_2)} - e^{-l(b_1+b_2)})$$

When $(a1 + a2 - b_1 - b_2) \to 0$:

$$f_{\mathcal{B}} \to a_i l e^{-l(b_1+b_2)}$$

One occurrence of each event on the branch

In this subsection, we assume that the occurrence of E_i precedes the occurrence of E_j on branch $\mathcal{B}$ ($\{i,j\} \in \{1,2\}, i \neq j$).

By analogy with the previous case, we define:

- a_1 (resp. a_2) as the occurrence rate for E_1 (resp. E_2) before the occurrence of E_i on the branch

- b_1 (resp. b_2) as the occurrence rate for E_1 (resp. E_2) after the occurrence of E_i and before the occurrence of E_j on the branch.

- c_1 (resp. c_2) as the occurrence rate for E_1 (resp. E_2) after the occurrence of E_j on the branch.

Those parameters take different values depending on the initial states on the branch, and according to the model. We easily get

$$f_{\mathcal{B}} = \int_0^l a_i e^{-a_1 t} e^{-a_2 t}(\int_0^{l-t} b_j e^{-b_1 x} e^{-b_2 x} e^{-c_1(l-t-x)} e^{-c_2(l-t-x)} dx) dt$$

$$= \frac{a_i b_j}{c_1+c_2-b_1-b_2}\left(\frac{e^{-l(a_1+a_2)}-e^{-l(b_1+b_2)}}{b_1+b_2-a_1-a_2} - \frac{e^{-l(a_1+a_2)}-e^{-l(c_1+c_2)}}{c_1+c_2-a_1-a_2}\right)$$

If all three denominators tend to 0,

$$f_{\mathcal{B}} \to \frac{1}{2}a_i b_j l^2 e^{-l(b_1+b_2)}$$

When $c_1 + c_2 - b_1 - b_2 \to 0$,

$$f_{\mathcal{B}} \quad \rightarrow \frac{a_i b_j}{b_1+b_2-a_1-a_2}\left(\frac{e^{-l(a_1+a_2)}-e^{-l(b_1+b_2)}}{b_1+b_2-a_1-a_2} - le^{-l(b_1+b_2)}\right)$$

When $b_1 + b_2 - a_1 - a_2 \rightarrow 0$,

$$f_{\mathcal{B}} \quad \rightarrow \frac{a_i b_j}{c_1+c_2-b_1-b_2}\left(le^{-l(b_1+b_2)} - \frac{e^{-l(b_1+b_2)}-e^{-l(c_1+c_2)}}{c_1+c_2-b_1-b_2}\right)$$

When $a_1 + a_2 - c_1 - c_2 \rightarrow 0$,

$$f_{\mathcal{B}} \quad \rightarrow \frac{a_i b_j}{c_1+c_2-b_1-b_2}\left(\frac{e^{-l(b_1+b_2)}-e^{-l(c_1+c_2)}}{c_1+c_2-b_1-b_2} - le^{-l(c_1+c_2)}\right)$$

A) State changes for process 1

B) Few state changes for both processes

C) A phylogenetic tree with events and rates
(not showing the hidden states)

Figure 1: A minimal model of correlated evolution between two processes. (A) An illustration showing the hidden states of process 1 (here noted with As) and their corresponding occurrence rates (indicated within the disks). Lines with black disks represent switches due to the occurrence of E_1 events, whereas lines with grey disks represent the effect of occurrences of E_2 events. (B) A subset of hidden state and rate switches for the two processes. Notation is the same as in (A). (C) An illustration of the occurrence rate switches on a mock phylogeny where the initial rates at the root are μ_1 and μ_2 (the hidden state would be A, B). Black disks represent E_1 events whereas grey ones represent E_2 events.

Figure 2: Empirical distribution of 10,000 replicates of rate estimation by maximum likelihood. For each replicate, we simulated on a perfectly symmetric tree with 64 leaves, occurrences of events with the following rates: $\mu_1 = 5$, $\mu_2 = 4$, $\mu_1^* = 20$ and $\mu_2^* = 50$. The different peaks at low rates correspond to cases where the numbers of realized events are small (1 or 2).

Figure 3: **Power of the likelihood ratio tests (LRT).** The power of LRTs (LRT curves) has been estimated and compared to the power of our previous methods (epics curves). (A) In the left panel, the simulated scenario has 3 parameters: $\mu_1 = 5$, $\mu_2 = 5$ and $\lambda = \mu_2^*/\mu_2$; increasing λ values represents greater induction of E_1 events on E_2. Two LRTs are computed : LRT_1df and LRT_2df (see text). Thresholds of the LRT are computed assuming a χ^2 distribution with 1 or 2 degrees of freedom. Our previous method (epics) computes exact p-values using either counts of co-occurring events (epics_Id) or counts of co-occurring plus sequentially ordered events (epics_S+Id). (B) In the right panel, the simulated scenario has a symmetric induction that is tuned by $\lambda = \mu_1^*/\mu_1 = \mu_2^*/\mu_2$.

Figure 4: **Power to assess correctly whether there is an induction, a loss or both**. **(A) Model selection and color code.** All LRTs compare nested models with at least 2 rates (μ_1 and μ_2) and optional induction ($\lambda_2 = \mu_2^*/\mu_2$) and/or loss ($\nu_1 = \nu_2 = 0$). Induction is "inferred" (blue area) when LRT_1 and LRT_4 are the only significant tests. Loss is "inferred" (red area) when LRT_2 and LRT_3 are the only significant tests. Both induction and loss are "likely" (dark grey area) when LRT_1 and LRT_2 are significant as well as either LRT_3 or LRT_4. Both are "inferred" (black areas) when all four tests are significant. White areas mark cases where no LRT is significant. All other cases are represented by light grey areas. **(B-D) Results.** For each value of the x-axis, we simulated 1,000 replicates of one of three scenarios: (B) Induction of E_1 on E_2, with $\mu_1 = \mu_2 = 10$ and for increasing values of μ_2^*; (C) Loss scenario, where $\nu_1 = \nu_2 = 0$, for increasing values of μ_1; (D) Induction+loss scenario, with $\mu_1 = \mu_2 = 25$ and for increasing values of μ_2^*. The y-axis indicates the fraction of each scenario that is inferred/likely.

Figure 5: **Analysis of Watson-Crick pairs of RNA 16S, in a tree of 54 enterobacteria.** Assuming a model with 5 parameters (μ_1, μ_2, ν_1, ν_2 and $\lambda = \mu_i^*/\mu_i = \nu_i^*/\nu_i$), we estimated the reciprocal induction rate of mutation of one site onto the other. The reported rates are colored in black or white, depending on whether there is statistical support for correlated evolution (at a risk of 1%).

A) Ranking the 75 best pairs

Rank	1	10	20	30	40	50	60	70

epics-Id: Log10(p-val) -11.8 -7.0 -5.1 -4.4 -3.9 -3.7 -3.4 -3.3

LRT: Log10(p-val) -23.4 -10.8 -8.0 -6.2 -5.2 -4.9 -4.6 -4.3

CoMap: Log10(p-val) -5.9 -5.7 -5.3 -5.2 -5.1 -4.6 -4.6 -4.3

MIp: Log10(p-val) -5.3 -4.9 -4.6 -3.8 -3.6 -3.0 -2.8 -2.6

■ WC   ▨ <10 Å   ▢ >10 Å

B) Structural positioning of the 75 best pairs

											>10Å
epics-Id	39	7	6	2	1	1	1	-	-	-	18
LRT	40	2	3	1	1	-	1	-	-	-	27
CoMap	39	1	1	-	1	-	1	1	1	-	30
MIp	39	5	3	2	2	-	1	1	1	1	20

Figure 6: **Analysis of the 75 best nucleotide pairs of 16S showing correlated evolution.** (A) We report the nature of the 75 "best" pairs (the ones with the lowest probabilities of independence) among all nucleotide pairs, according to 4 different methods: our previous method with co-occurrences (epics-Id), an LRT based method (see legend of Figure 5) using our likelihood framework, the CoMap method based on co-occurrences on the tree and the MIp method based on Mutual Information, corrected for phylogeny. Pairs are colored in black when they are Watson-Crick, on dark grey when they are less than 10Å apart in the 3D structure and in light grey otherwise. Below their rank, we also report their associated p-value. (B) Relative structural positions on the stems-loops for the nucleotides of the 75 "best" pairs, according to the 4 methods.