Genome-wide approaches for the identification of markers and genes associated with sugarcane yellow leaf virus resistance

Ricardo José Gonzaga Pimenta[1], Alexandre Hild Aono[1], Roberto Carlos Burbano Villavicencio[2], Alisson Esdras Coutinho[3], Carla Cristina da Silva[1], Ivan Antônio dos Anjos[4], Dilermando Perecin[3], Marcos Guimarães de Andrade Landell[4], Marcos Cesar Gonçalves[5], Luciana Rossini Pinto[4], Anete Pereira de Souza[1,6*]

[1]Centre of Molecular Biology and Genetic Engineering (CBMEG), University of Campinas (UNICAMP), Campinas, Brazil

[2]Gustavo Galindo Velasco Campus, Littoral Polytechnic Superior School (ESPOL), Guayaquil, Ecuador

[3]São Paulo State University (UNESP), Jaboticabal, Brazil

[4]Advanced Centre for Technological Research in Sugarcane Agribusiness, Agronomic Institute of Campinas (IAC/APTA), Ribeirão Preto, Brazil

[5]Plant Protection Research Centre, Biological Institute (IB/APTA), São Paulo, Brazil

[6]Department of Plant Biology, Institute of Biology, University of Campinas (UNICAMP), Campinas, Brazil

* Correspondence
Anete Pereira de Souza
anete@unicamp.br

Keywords: *Saccharum,* **SCYLV, RT-qPCR, AFLP, SSR, GBS, GWAS, Machine learning.**

Abstract

The breeding of sugarcane, a leading sugar and energy crop, is particularly hampered by the extremely complex sugarcane genome. One of the main diseases that affect this crop is sugarcane yellow leaf (SCYL), caused by the sugarcane yellow leaf virus (SCYLV). Regardless of the manifestation of SCYL symptoms, infection by SCYLV can lead to substantial yield losses, making resistance to this virus highly relevant to sugarcane breeding. However, the genetic basis of this trait has not been unraveled or widely explored. In this context, several methodologies have arisen as promising tools for the identification of molecular markers associated with SCYLV resistance, which are crucial for marker-assisted selection and can contribute to the understanding of the mechanisms involved in the response to this virus. Here, we investigated the genetic basis of SCYLV resistance in sugarcane using dominant and codominant markers and genotypes of interest for breeding. A sugarcane panel inoculated with SCYLV was analyzed for SCYL symptom severity, and viral titer was estimated by reverse transcription quantitative PCR. This panel was genotyped with 662 dominant markers, as well as by genotyping-by-sequencing, which enabled the identification of 70,888 SNPs and indels with information on allele proportion and position on the *Saccharum spontaneum* genome. We then performed polyploid-adapted genome-wide association analyses and used machine-learning algorithms coupled with feature selection methods to establish marker-trait associations. While each approach identified a unique set of markers associated with the phenotypes, we also observed convergences between them, demonstrating their complementarity. Finally, by aligning the sequences neighboring these markers to the genomic coding sequences of closely related

plant species, we annotated genes harboring 176 of these markers. With this procedure, we identified
genes encoding emblematic participants in virus resistance mechanisms, as well as other candidates
whose involvement in the responses to viruses has not been reported before. Our approach has the
potential to accelerate sugarcane breeding targeting SCYLV resistance and is a powerful tool for
studies on the biological processes leading to this trait.

1   Introduction

Sugarcane is one of the world's most important crops, ranking first in production quantity and sixth
in net production value in 2016 (FAO, 2020). It is by far the most relevant sugar crop, accounting for
approximately 80% of the world's sugar production (FAO, 2020; ISO, 2020) and is also a prominent
energy crop. It is estimated that by 2022, sugarcane will account for 40% of the world's first-
generation biofuels (Lalman et al., 2016), and biomass-rich energy cane is a promising source of
second-generation ethanol (Carvalho-Netto et al., 2014). However, this crop has an extremely
complex genome; modern cultivars are the product of a few crosses between two autopolyploid
species conducted at the end of the 19th century (Bremer, 1961). *Saccharum spontaneum* ($2n = 5x =$
40 to $16x = 128$; $x = 8$) (Panje and Babu, 1960), a wild stress-resistant but low-sugar species, was
hybridized with *Saccharum officinarum* ($2n = 8x = 80$, $x = 10$) (D'Hont et al., 1998), which has a
high sugar content but is sensitive to drought and susceptible to diseases. This was followed by
several backcrosses with *S. officinarum* to restore sugar productivity in a process known as
nobilization (Roach, 1972). These procedures gave origin to plants with very large (ca. 10 Gb)
(D'Hont et al., 1998), highly polyploid (D'Hont and Glaszmann, 2001), aneuploid (Sforça et al.,
2019) and remarkably repetitive (Mancini et al., 2018) genomes. This complexity directly affects
sugarcane research and breeding; for instance, until recently, it prevented the use of codominance
information in marker-assisted breeding strategies for this crop, leading to loss of information during
the performance of these approaches and therefore limiting their use (Premachandran et al., 2011;
Bourke et al., 2018).

One of the diseases that affect this crop is sugarcane yellow leaf (SCYL), which is caused by
sugarcane yellow leaf virus (SCYLV), a positive-sense ssRNA belonging to the *Polerovirus* genus
(Moonan et al., 2000; Smith et al., 2000). The expression of SCYL symptoms is complex and usually
occurs in late stages of plant development. The main symptom of these infections is the intense
yellowing of midribs in the abaxial surface of leaves, and other visible manifestations are necrosis of
leaf tips, leaf chlorosis and plant stunting (Scagliusi and Lockhart, 2000; ElSayed et al., 2015). The
virus is known to induce alterations in the metabolism and transport of sucrose and in photosynthetic
efficiency (Gonçalves et al., 2005; Lehrer et al., 2010); this impairs plant development and eventually
reflects losses in sugarcane productivity (Vega et al., 1997; Grisham et al., 2001; Vasconcelos et al.,
2009; Viswanathan et al., 2014). Many symptoms of SCYL may, however, be caused by other biotic
and abiotic stresses or plant senescence (Vega et al., 1997; Aljanabi et al., 2001; ElSayed et al.,
2015), making SCYL identification troublesome. Therefore, the diagnosis of infection by molecular
methods is of great assistance. Immunological assays were the first methods developed to detect and
quantify this virus (Scagliusi and Lockhart., 2000), but later assessments by reverse transcription
followed by quantitative polymerase chain reaction (RT-qPCR) are much more sensitive and precise
in estimating the levels of viral infection (Gonçalves et al., 2002; Korimbocus et al., 2002; Zhu et al.,
2010).

Due to SCYL's elusive symptomatology, the spread of its causal agent is silent; SCYLV is
considered endemic in all sugarcane-producing regions of the world (Gonçalves, 2007). The virus is
disseminated mostly during sugarcane vegetative propagation by setts but is also transmitted by

aphids, mainly the white sugarcane aphid *Melanaphis sacchari* (Zehntner, 1897) (Scagliusi and
Lockhart, 2000). Unlike other pathogens, SCYLV is not efficiently eliminated by thermal treatments
(Delage et al., 1999); the only way to thoroughly remove the virus from plants is by meristem
micropropagation (Chatenet et al., 2001; Fitch et al., 2001), which is time-consuming and requires
specialized infrastructure and personnel. These features make varietal resistance to SCYLV the
resource that can be most efficiently exploited to prevent damage and losses caused by this virus.
Sugarcane resistance to SCYLV has been explored in breeding programs, but research on SCYL
genetics is not exempt from the difficulties generated by the complexity of this crop's genome
(Gonçalves et al., 2012).

The identification of loci linked to resistance to SCYLV has been investigated in a few studies,
which involved either quantitative trait locus (QTL) mapping in biparental progenies (Costet et al.,
2012; Islam et al., 2018) or genome-wide association studies (GWASs) (Débibakas et al., 2014;
Gouy et al., 2015; Yang et al., 2019a). Due to sugarcane's polyploid nature, most of these studies
employed molecular markers that were treated as dominant, which implies a great loss of genetic
information (Garcia et al., 2013). Additionally, these works employed immunological methods to
phenotype SCYLV resistance. The usage of dominant markers and the poor reliability of
phenotyping, allied with flaws in the assembly of association panels, were listed as factors that
limited the power of the detection of QTLs and markers associated with disease resistance
(Débibakas et al., 2014; Gouy et al., 2015).

In this study, we evaluated the efficacy of several genome-wide approaches to identify markers and
genes associated with SCYLV resistance. We analyzed a panel of 97 *Saccharum* accessions
inoculated with SCYLV, which were graded for the severity of SCYL symptoms, and their viral
titration was estimated by relative and absolute RT-qPCR. This panel was genotyped with amplified
fragment length polymorphisms (AFLPs) and simple sequence repeats (SSRs), which were treated as
dominant. We also constructed a genotyping-by-sequencing (GBS) library for 94 genotypes of this
population, which was used for the discovery of single nucleotide polymorphisms (SNPs) and
insertions and deletions (indels) using a monoploid set of chromosomes of *S. spontaneum* (Zhang et
al., 2018) as a genomic reference. We then employed three distinct methodologies to detect marker-
trait associations: the fixed and random model circulating probability unification (FarmCPU) method
using dominant markers; mixed linear modeling using SNPs and indels, in which allele proportions
(APs) in each locus were employed to establish genotypic classes and estimate additive and dominant
effects; and several machine learning (ML) methods coupled with feature selection (FS) techniques,
using all markers to predict genotype attribution to phenotypic clusters identified by clustering
analyses. Finally, by aligning neighboring sequences of the genomes of plant species closely related
to sugarcane, we annotated genes containing markers associated with phenotypes. We discuss the
putative participation of these genes in the mechanisms underlying resistance to SCYLV.

2   Material and Methods

2.1   Plant Material and Inoculation

The plant material and inoculation methods employed in the present study are described by
Villavicencio et al. (in press). The experimental population consisted of a panel of 97 sugarcane
genotypes that comprised wild germplasm accessions of *S. officinarum*, *S. spontaneum* and
Saccharum robustum; traditional sugarcane and energy cane clones; and commercial cultivars
originating from Brazilian breeding programs. The accession names and pedigree information are
available in Supplementary Table 1. To ensure plant infection with SCYLV, a field nursery was

established in March 2016 at the Advanced Centre for Technological Research in Sugarcane
Agribusiness located in Ribeirão Preto, São Paulo, Brazil (4°52'34" W, 21°12'50" S). Seedlings
from sprouted setts of each genotype were planted in 1-meter plots with an interplot spacing of 1.5
meters. The cultivar SP71-6163, which is highly susceptible to SCYLV (Vega et al., 1997), was
interspersed with the panel genotypes. *M. sacchari* aphids were reared on detached SP71-6163 leaves
under laboratory conditions and released weekly in the field nursery in July 2016. Setts obtained
from this nursery were used to install a field experiment following a randomized complete block
design with three blocks in May 2017. Plants were grown in 1-meter-long three-row plots with row-
to-row and interplot spacings of 1.5 and 2 meters, respectively. Each plot contained two plants,
totaling six plants of each genotype per plot. To further assist infection by SCYLV, the cultivar
SP71-6163 was planted in the borders and between blocks, and *M. sacchari* aphids were released in
the field weekly for five months, starting from November 2017.

142   **2.2   Phenotyping**

Plants were phenotyped in two crop seasons: plant cane in June 2018 (13 months after planting) and
ratoon cane in July 2019 (9 months after the first harvest). The severity of SCYL symptoms was
assessed by three independent evaluators, who classified the top visible dewlap leaves (TVDLs) of
each plot using a diagrammatic scale established by Villavicencio et al. (in press). The scale
consisted of four levels, as follows: (1) green leaves with no symptoms; (2) slight yellowing of the
midrib abaxial surface, with discrete or no progression to the leaf blade; (3) intense yellowing of the
midrib abaxial surface and a more pronounced yellowing of the leaf blade; and (4) intense yellowing
of both the midrib abaxial surface and leaf blade (Supplementary Figure 1).

In the same week as symptom evaluation was performed, fragments from the median region of at
least one TVDL per plot were collected and stored at -80°C until processing. Total RNA was
extracted from this tissue using TRIzol (Invitrogen, Carlsbad, USA). Samples were subjected to an
additional purification process consisting of three steps: (i) mixing equal volumes of RNA extract
and chloroform, (ii) precipitating the RNA overnight with 2.5 volumes of 100% ethanol and (iii) a
conventional cleaning step with 70% ethanol. RNA was then quantified on a NanoDrop 2000
spectrophotometer (Thermo Scientific, Waltham, USA) and subjected to electrophoresis on a 1%
agarose gel stained with ethidium bromide for integrity checks. The samples were next diluted and
treated with RNase-Free RQ1 DNase (Promega, Madison, USA); after this step, the RNA was
quantified and diluted again for standardization. The RNA was then converted to cDNA using the
ImProm-II Reverse Transcription System kit (Promega, Madison, USA).

The SCYLV titer in each sample was determined by qPCR using GoTaq qPCR Master Mix
(Promega, Madison, USA) on a Bio-Rad CFX384 Touch detection system (Bio-Rad, Philadelphia,
USA). Two viral quantification methodologies were employed – one relative and one absolute –
using primers and conditions as described by Chinnaraja and Viswanathan (2015). For both methods,
a set of primers was used to amplify a 181-bp fragment from SCYLV ORF3 (YLSRT). For the
relative quantification, an additional set of primers was used to amplify a 156-bp fragment of the 25S
subunit of sugarcane ribosomal RNA (25SrRNA), which was used as an internal control. The $2^{-\Delta\Delta CT}$
method (Livak and Schmittgen, 2001) was used to correct cycle threshold (CT) values; the sample
with the highest CT and a melting temperature of 82.5 ± 0.5°C for the YLSRT primers was used as a
control for phenotyping in each year. The absolute quantification followed the methodology
described by Chinnaraja et al. (2014). A pGEM-T Easy vector (Promega, Madison, USA) cloned
with a 450-bp fragment from SCYLV ORF3 previously amplified by RT-PCR was used to construct

a serial dilution curve with six points and tenfold dilutions between points; this curve was amplified
on all qPCR plates. All reactions were performed using three technical replicates.

2.3 Phenotypic Data Analyses

The normality of phenotypic data was assessed by Shapiro-Wilk tests, and normalization was carried
out using the bestNormalize package (Peterson, 2017) in R software (R Core Team, 2011). The best
linear unbiased predictors (BLUPs) were estimated for each trait with the breedR R package (Muñoz
and Sanchez, 2015) using a mixed model as follows:

$$Y_{ijm} = \mu + B_j + Y_m + BY_{jm} + G_{i(jm)} + e_{ijm}$$

where $Yijm$ is the phenotype of the ith genotype considering the jth block and the mth year of
phenotyping. The trait mean is represented by μ; fixed effects were modeled to estimate the
contributions of the jth block (B_j), the mth year (Y_m) and the interaction between block and year (B_{Yjm}).
Random effects included the genotype (G) and the residual error (e), representing nongenetic effects.

Pearson's correlation tests were performed using the BLUPs to check the correlation between the
three traits (symptom severity, relative quantification and absolute quantification), and correlation
distributions were plotted using the GGally R package (Schloerke et al., 2011). To investigate the
separation of genotypes according to phenotypes, we performed two hierarchical clustering on
principal components (HCPC) analyses with the factoMineR package (Lê et al., 2008) – first using
only viral quantification and then employing the three analyzed traits. The factoextra R package
(Kassambara and Mundt, 2017) was used to plot graphs associated with these analyses. Statistical
differences between the phenotypes of the clusters identified in each HCPC were assessed by
Kruskal-Wallis tests or analyses of variance (ANOVAs), depending on the distribution of the data.
Post hoc Dunn's tests using the Bonferroni correction were performed with the R package dunn.test
(Dinno, 2017) to verify pairwise differences between clusters.

2.4 Genotyping

2.4.1 Dominant Markers

Total DNA was extracted from leaves of each genotype following the method described by Aljanabi
et al. (1999). AFLPs were developed using *Eco*RI and *Msp*I restriction enzymes (New England
BioLabs). Digestion reactions were prepared in a final volume of 20 µL containing 300 ng DNA, 2.5
U of each restriction enzyme in 1X RL Buffer (New England BioLabs) and incubated in a
thermocycler for 3 hours at 37°C and for 5 min at 70°C. Adapter ligation was conducted in a final
volume of 40 µL containing 20 µL of the digestion reaction, 5 X buffer (40 mM Tris pH 8.4, 100 mM
KCl), 0.5 µM *Eco*RI adaptor, 5 µM *Msp*I adaptor, 1 mM ATP and 0.85 U of T4 DNA ligase (67
U/µL) (BioLabs). Ligation was performed at 37°C for 2 hours and 16°C for 16 hours in a
thermocycler. Preamplification was conducted with primers complementary to restriction enzyme
adaptors and devoid of selective nucleotides at the 3' end (*Eco*RI+0 and *Msp*I+0 primers) and using a
6X dilution of the digestion/ligation product. This reaction was performed in a final volume of 15 µL
containing 2 µL of the 6X dilution digestion/ligation product, 1X PCR buffer (20 mM Tris pH 8.4, 50
mM KCl), 3.3 µM *Eco*RI+0 and *Msp*I+0 primers, 0.17 mM dNTPs, 2 mM MgCl$_2$ and 0.07 U Taq
DNA polymerase. The cycling conditions were as follows: 29 cycles at 94°C for 30 seconds, 56°C
for 1 minute and 72°C for 1 minute. Preamplification reactions were diluted 10X and used for
selective amplification reactions using combinations of *Eco*RI/*Msp*I primers with three selective
nucleotides at the 3`end and the *Eco*RI primer labeled with fluorophores IRDye700 or IRDye800.

Thirty-five selective primer combinations were used (Supplementary Table 2). The reaction was
performed in a final volume of 10 μL containing 2.5 μL of the 10X diluted preamplification, 1X PCR
buffer (20 mM Tris pH 8.4, 50 mM KCl), 0.05 μM of selective Eco700 labeled primer (or 0.07 μM
Eco800 primer), 0.25 μM for Msp selective primer, 0.25 μM dNTPs, 2 mM MgCl$_2$, 0.5 U of Taq
DNA polymerase. The cycling conditions were as follows: one cycle at 94°C for 30 seconds, 65°C
for 30 seconds and 72°C for 1 minute followed by 12 cycles at 94°C for 30 seconds, 65°C for 30
seconds (decreasing 0.7°C/cycle) and 72°C for 1 minute, followed by 23 cycles of 94°C for 30
seconds, 56°C for 30 seconds and 72°C for 1 minute. Final amplicons were separated on a 6%
denaturing polyacrylamide gel and visualized with a LI-COR 4300 DNA Analyzer (LI-COR,
Lincoln, NE, USA).

Twelve SSR loci previously isolated from the sugarcane expressed sequence tag database
(Maccheroni et al., 2007; Oliveira et al., 2007, 2009; Marconi et al., 2011) were used for SSR
genotyping; details on each SSR locus can be found in Supplementary Table 3. PCR mixes were
prepared, and amplifications were conducted in a Bio-Rad MyCycler thermal cycler (Bio-Rad,
Philadelphia, USA) following the conditions previously established by Oliveira et al. (2009) and
Marconi et al. (2011); primers were labeled with fluorescent dyes IRDye700 and IRDye800 to allow
band visualization. Amplicons were separated on a 6% denaturing polyacrylamide gel and visualized
with a LI-COR 4300 DNA Analyzer. Due to sugarcane's polyploidy, both AFLPs and SSRs were
treated as dominant and scored based on the presence (1) or absence (0) of bands. After genotyping,
genotypes and markers with over 10% missing data were removed, as well as markers with a minor
allele frequency (MAF) below 10%.

2.4.2   Genotyping-by-sequencing

Genomic DNA was extracted from young leaves using the GenElute Plant Genomic DNA Miniprep
Kit (Sigma-Aldrich, St. Louis, USA). The integrity of the DNA was verified by electrophoresis on a
1% agarose gel stained with ethidium bromide, and its concentration was determined using a Qubit
3.0 fluorometer (Thermo Scientific, Wilmington, USA). The construction of the GBS library was
based on a protocol by Poland et al. (2012) and used a combination of *Pst*I and *Mse*I restriction
enzymes. For operational reasons, 94 out of the 97 genotypes of the panel were included in the
library, which did not include genotypes 87, 88 and 95 (see Supplementary Table 1). The library was
subjected to a purification step using polyethylene glycol as described by Lundin et al. (2010) with
slight modifications. It was then validated with a Fragment Analyzer (Agilent Technologies, Santa
Clara, USA) and quantified by RT-qPCR in a Bio-Rad CFX384 Touch detection system using the
KAPPA KK4824 kit (Kapa Biosystems, Wilmington, USA). Two 150-bp single-end sequencing
libraries were prepared using the NextSeq 500/550 High Output Kit (Illumina, San Diego, USA) and
sequenced on a NextSeq 500 (Illumina, San Diego, USA).

After checking sequencing quality with FastQC (Andrews, 2010), we used Stacks software version
1.42 (Catchen et al., 2013) for demultiplexing and checking the amount of data generated for each
sample. The TASSEL4-POLY pipeline (Pereira et al., 2018), developed from TASSEL-GBS
(Glaubitz et al., 2014), was used for variant calling. Most parameters were set at their standard
values; exceptions were the use of the "inclGaps" argument in the "DiscoverySNPCaller" plugin, the
"misMat" argument with a value of 0.3 and the "callHets" argument in the "MergeDuplicateSNPs"
plugin. Rather than aligning raw reads to a reference genome, the TASSEL-GBS pipeline first
generates "tags" – unique sequences representing redundant reads – to reduce computation time
(Glaubitz et al., 2014). We tested the mapping of tags against nine genomic references using two
aligners: BWA version 0.7.2 (Li and Durbin, 2009) and Bowtie2 version 2.2.5 (Langmead and

Salzberg, 2012). The genomic references used were as follows: the *Sorghum bicolor* genome
(Paterson et al., 2009), the methyl-filtered genome of the sugarcane cultivar SP70-1143 (Grativol et
al., 2014), a sugarcane RNA-Seq assembly (Cardoso-Silva et al., 2014), a de novo assembly
generated from GBS data following the GBS-SNP-CROP pipeline (Melo et al., 2016), a draft
genome of the sugarcane cultivar SP80-3280 (Riaño-Pachón and Mattiello, 2017), a sugarcane
transcriptome generated by Iso-Seq (Hoang et al., 2017), the mosaic monoploid genome of the
sugarcane cultivar R570 (Garsmeur et al., 2018), the *S. spontaneum* genome (Zhang et al., 2018) and
a monoploid chromosomic set obtained from this same reference that included the "A" haplotype and
unassembled scaffolds. After variant calling, VCFtools version 0.1.13 (Danecek et al., 2011) was
used to retain biallelic markers with a MAF of 0.1, no missing data and a minimum sequencing depth
of 50 reads. The most appropriate reference was chosen, and adopting the method proposed by Yang
et al. (2019b), the ratio between alleles (allele proportions, APs) of each variant was transformed into
genotypes with a fixed ploidy of 12; this conversion was performed using the vcfR R package (Knaus
and Grünwald, 2017).

2.5   Population Structure and Linkage Disequilibrium Analyses

Three procedures were used to evaluate genetic structuring in the panel, employing dominant and
codominant markers separately; for all analyses, the maximum number of clusters in the panel was
set to 10. The first method was a discriminant analysis on principal components (DAPC), performed
in the adegenet R package (Jombart et al., 2008). The second was a principal component analysis
(PCA) followed by K-means, for which missing data were imputed with the pcaMethods package
(Stacklies et al., 2007) and for which the optimal number of clusters was evaluated using the elbow,
silhouette and gap statistic methods in the factoextra package. The last was a Bayesian clustering of
genotypes into predetermined numbers of clusters (K) performed on STRUCTURE software
(Pritchard et al., 2000), assuming an admixture model with correlated allelic frequencies between
populations. Ten independent runs were implemented for each K, and for dominant markers,
estimates of probabilities of values of K in each run were taken following 100,000 generations as
burn-in and 200,000 generations sampled in a Monte Carlo Markov Chain (MCMC). For Bayesian
clustering using SNPs and indels, we used a subset of 7,000 markers randomly sampled from the
total dataset, parallelized STRUCTURE with StrAuto software (Chhatre and Emerson, 2016) and
sampled 100,000 generations in the MCMC. In both cases, the most likely number of genetic clusters
was determined by the ad hoc statistics ΔK (Evanno et al., 2005) and the LnP(D) probability
logarithm; the output was interpreted in STRUCTURE HARVESTER software version 0.6.94 (Earl
et al., 2012). Clumpak software (Kopelman et al., 2015) was used to average the admixture
proportions of runs and to estimate cluster membership coefficients for genotypes.

For SNPs and indels, we also measured linkage disequilibrium (LD) on the ldsep R package (Gerard,
2020) by calculating the squared correlation coefficient (r^2) between pairs of markers on the same
chromosome. The decay of LD over physical distance was investigated by pooling all chromosomes,
plotting pairwise r^2 values against the distance between markers and fitting a curve using the
equation proposed by Hill and Weir (1988). The critical r^2 for LD decay was set to 0.1, the most
commonly used threshold for determining the existence of LD (Vos et al., 2017). Only comparisons
with $p < 0.05$ were used in this analysis.

2.6   Association Analyses

2.6.1   FarmCPU

Association analyses with dominant markers were performed with the FarmCPU method in R (Liu et
al., 2016). For these analyses, markers were recoded to indicate the presence (0) and absence (2) of
bands. We tested FarmCPU using no covariates and including matrices obtained from the three
genetic structure analyses described in the previous section as such. In each case, a "quantile-
quantile" (Q-Q) plot of the $-\log_{10}(p)$ values of markers was generated, and the genomic inflation
factor λ (Devlin and Roeder, 1999) was calculated. The average λ from the analyses of the three traits
employing each covariate matrix was calculated and used to select the model that best controlled
inflation. The Bonferroni correction with α = 0.05 was used to establish the significance threshold for
associations. The phenotypic variance explained by each marker was estimated for significant
marker-trait associations using a linear model in R software.

2.6.2   Mixed Modeling in GWASpoly

Association analyses using SNPs and indels were performed using mixed linear model approaches in
the GWASpoly R package (Rosyara et al., 2016). The output of the three genetic structure analyses
previously described was used to build Q matrices, which were included in the models as fixed
effects. Similarly, three different genetic kinship matrices (K) of the panel were computed and
included as random effects: (I) a realized relationship model (MM^T) matrix (VanRaden, 2008), built
on GWASpoly; (II) a complete autopolyploid matrix based on Slater et al. (2016), built with the
AGHmatrix R package (Amadeu et al., 2016); and (III) a pseudodiploid matrix based on Slater et al.
(2016), also built with AGHmatrix. We tested twelve Q + K combinations and, for each of them, six
marker-effect models were used: general, additive, simplex dominant reference, simplex dominant
alternative, diploidized general and diploidized additive. For each model, a Q-Q plot of the $-\log_{10}(p)$
values of markers was generated, and λ was calculated. The average λ of all traits and models
employing each Q + K combination was calculated and used to select the best set of matrices. Once
this combination was chosen, Manhattan plots were generated for all models and traits. The
Bonferroni and false discovery rate (FDR) correction methods with α = 0.05 were assessed to
establish the significance threshold for associations.

2.6.3   Machine Learning Coupled with Feature Selection

Finally, we assessed the capacity of ML strategies to predict the attribution of genotypes to the
phenotypic groups identified in the HCPC analyses based on all markers, following the genomic
prediction approach proposed by Aono et al. (2020). For this approach, we selected accessions
successfully genotyped with both SNPs/indels and AFLPs/SSRs; missing data in dominant markers
were imputed as the means. We evaluated the accuracy of eight ML algorithms: adaptive boosting
(AB) (Freund and Schapire, 1997), decision tree (DT) (Quinlan, 1986), Gaussian naive Bayes (GNB)
(Friedman et al., 1997), Gaussian process (GP) (Rasmussem, 2003), K-nearest neighbor (KNN)
(Cover and Hart, 1967), multilayer perceptron (MLP) neural network (Popescu et al., 2009), random
forest (RF) (Breiman, 2001) and support vector machine (SVM) (Cristianini and Shawe-Taylor,
2000), all implemented in the scikit-learn Python 3 module (Pedregosa et al., 2011). As a cross-
validation strategy, we used a stratified K-fold (k=5) repeated 100 times for different data
configurations.

We then tested five FS techniques to obtain feature importance and create subsets of marker data:
gradient tree boosting (FS1) (Chen and Guestrin, 2016), L1-based FS through a linear support vector

classification system (FS2) (Cristianini and Shawe-Taylor, 2000), extremely randomized trees (FS3)
(Geurts et al., 2006), univariate FS using ANOVA (FS4) and RF (FS5) (Breiman, 2001); all FS
approaches were implemented in the scikit-learn Python 3 module. We tested the differences in the
accuracy between the selected FS methods using ANOVAs and multiple comparisons by Tukey's
tests implemented in the agricolae R package (De Mendiburu and de Mendiburu, 2020). We also
evaluated the intersection of these datasets. Finally, the area under the receiver operating
characteristic (ROC) curve (AUC) was calculated for the best ML-FS combination and plotted using
the Matplotlib library89 with Python 3.

2.6   Marker Mapping and Annotation

The distribution of markers identified by all analyses along *S. spontaneum* "A" chromosomes was
visualized using MapChart (Voorips, 2002). Markers previously associated with SCYLV resistance
by QTL mapping (Costet et al., 2012; Islam et al., 2018) and GWAS (Débibakas et al., 2014; Yang et
al., 2019a) were also retrieved and included in the map. Finally, the sequences of associated markers
were annotated by aligning SSR flanking sequences or the 2000-bp window adjacent to SNPs and
indels against the coding sequences (CDSs) of the genomes of 14 Poaceae species (*Brachypodium
distachyon*, *Brachypodium hybridum*, *Brachypodium silvaticum*, *Hordeum vulgare*, *Oryza sativa,
Oropetium thomaeum*, *Panicum hallii*, *Panicum virgatum*, *S. bicolor*, *Setaria italica*, *Setaria viridis*,
Triticum aestivum, *Thinopyrum intermedium* and *Zea mays*) and *Arabidopsis thaliana* retrieved from
the Phytozome database (https://phytozome.jgi.doe.gov/pz/portal.html). For this, BLASTn (Altschul
et al., 1990) was used with an E-value of 1e-30, and the best alignment of each sequence was kept for
analysis.

3   Results

3.1   Phenotypic Data Analyses

A total of 97 sugarcane accessions inoculated with SCYLV were evaluated for the severity of SCYL
symptoms and for viral titer by relative and absolute RT-qPCR quantification of individual samples
in two consecutive years. We observed a skew in the distribution of symptom severity data towards
the absence of symptoms, and despite normalization procedures, these data did not follow a normal
distribution, as indicated by a Shapiro-Wilk test (p = 2.2e-16). This was not the case for the SCYLV
quantification data; both the relative (p = 0.999) and absolute (p = 0.930) quantification data were
successfully normalized. Plots of the distribution of the BLUPs calculated for the three traits
analyzed and of the correlations between them are depicted in Figure 1. Pearson's tests indicated that
symptom severity was not significantly correlated with the viral titer determined by the relative (p =
0.117) or absolute (p = 0.296) quantification. We found, however, a significant (p < 2.2e-16) and
strong (r^2 = 0.772) correlation between the values obtained by the two quantification methods,
indicating their reliability.

We performed two HCPC analyses to investigate the classification of genotypes according to the
phenotypes associated with SCYLV resistance. The first one, using BLUP values of SCYLV titer
determined by RT-qPCR, indicated the division of the panel into three clusters (Supplementary
Figure 2); a factor map wherein these groups are plotted onto the first two dimensions of this HCPC
is shown in Figure 2A, and the attribution of genotypes to each cluster is available in Supplementary
Table 4. Boxplots of the distribution of data among these clusters (Supplementary Figure 3) indicated
that each of these groups presented distinct SCYLV titers, as indicated by the two quantification
methods: cluster Q1 was composed of accessions that showed the lowest titers of SCYLV; cluster Q2
was formed by accessions that presented intermediate virus titers; and cluster Q3 included genotypes

with the highest virus titers. ANOVAs indicated that these differences were significant ($p < 2e-16$),
and Dunn tests showed that the virus titer determined by both quantification methods was
significantly different between all pairs of clusters (Supplementary Table 5).

The second HCPC, performed with the BLUPs of the three traits analyzed, also indicated the division
of the panel into three clusters (Supplementary Figure 4); a factor map wherein these groups are
plotted onto the first two dimensions of this HCPC is shown in Figure 2B, and the attribution of
genotypes to each cluster is available in Supplementary Table 4. Boxplots of the distribution of
phenotypic data among the three clusters (Supplementary Figure 5) clearly show that this division
resulted in a separation of groups with contrasting phenotypes. Cluster SQ1 was composed of
accessions that showed the least severe SCYL symptoms and presented low titers of SCYLV, as
indicated by the two quantification methods; interestingly, all genotypes attributed to Q1 were
included in SQ1. Cluster SQ2 was formed by accessions that also displayed mild disease symptoms
but presented high viral titers. Finally, cluster SQ3 included accessions with the most severe disease
symptoms and high virus titers. These differences were statistically significant; a Kruskal-Wallis test
indicated the existence of significant ($p = 4.078e-08$) differences in symptom severity between all
clusters, and ANOVAs showed that the same significance was present for the values of relative and
absolute viral quantifications ($p < 2e-16$). Dunn tests showed that the severity of SCLV symptoms in
each cluster was significantly different from the others and that SQ1 showed significantly lower
SCYLV titers than the other two clusters as estimated by both quantification methods
(Supplementary Table 5).

3.2   Genotyping

The genotyping of the 97 accessions of the panel using 12 SSR loci and 35 AFLP selective primer
combinations generated 981 polymorphic markers (167 SRR fragments and 814 AFLP fragments).
Upon filtering, four genotypes (74, 93, 96 and 97) and 319 markers were removed as they presented
call rate below 90% or MAF below 10%. Therefore, 93 genotypes and 662 dominant markers were
retained for further analyses that involved them.

The sequencing of the GBS library including 94 sugarcane genotypes generated 119,117,758 raw
reads, 87.88% of which presented a Q-value above 30. Genotypes 31 and 77 presented very low
amounts of generated data (0.19 and 1.5 KB, respectively) in contrast with the library average (598
KB) and were therefore excluded from subsequent analyses to avoid biasing. The results from tag
mapping against each genomic reference using BWA and Bowtie2 aligners can be found in
Supplementary Table 6. To avoid sampling of duplicated regions – which are abundant in the
sugarcane genome and could bias the calculation of APs – we did not include tags with multiple
alignments in the ensuing analyses. BWA presented slightly superior results in terms of uniquely
mapped tags and was therefore chosen as the aligning tool. Regarding the nine genomic references
tested for variant calling, the methyl-filtered genome of the cultivar SP70-1143 resulted in the largest
number of isolated markers among all references, yielding 60,173 SNPs and 43,054 indels after
rigorous filtering (Supplementary Table 7). However, knowing that a high-quality genome assembly
can greatly improve the results of GWASs (Benevenuto et al., 2019) and also allow the estimation of
long-distance LD, we opted to perform ensuing analyses with markers obtained from a reference that
not only yielded a large number of markers but also provided reliable information on their position.
Thus, we chose to use the monoploid chromosome set isolated from the *S. spontaneum* genome as a
reference; despite not yielding the highest number of markers (38,710 SNPs and 32,178 indels), this
genome had a performance superior to most of the other references tested here and, unlike many
others, provided markers with information of position at chromosome level.

3.3   Panel Genetic Structure and Linkage Disequilibrium

The genetic structure of the panel was investigated using two separate sets of markers – AFLPs and
SSRs scored as dominant and codominant SNPs and indels with AP information – and three different
approaches – a DAPC, a PCA followed by k-means and a Bayesian clustering implemented in
STRUCTURE software. Supplementary Table 8 summarizes the allocation of genotypes to the
clusters identified in each analysis. First, the three methods were performed with dominant markers.
The DAPC was performed retaining 70 principal components (PCs) that together explained ~90% of
the observed variance (Supplementary Figure 6A); the Bayesian information criterion (BIC) curve
indicated k = 3 as the most likely number of clusters in the panel (Supplementary Figure 6B).
Therefore, genotypes were separated into three clusters (A1-3); the plotting of the panel elements
onto the first two linear discriminants from this analysis is shown in Figure 3A. Despite resulting in
the subdivision of the panel into four clusters (B1-4), as indicated by two out of three statistical
methods performed to assess the optimal number of clusters in the panel (Supplementary Figures 7A-
C), the clustering by PCA followed by k-means yielded results that shared similarities with the
DAPC. Genotypes forming B1 and B4, for instance, were entirely represented by A3 and A1,
respectively. Figure 3B depicts a projection of groups identified by k-means onto the PCA's first two
PCs, which accounted for ~18% of the variation in the genotypic data. Last, based on the Bayesian
clustering, both ΔK and LnP(D) methods clearly indicated K = 2 as the most likely number of
clusters in the panel (Supplementary Figure 8A and B, respectively). Therefore, we present here a
plot of the estimated association coefficients of individuals for these two clusters (Figure 3C), named
C1 and C2. Clustering by this method was highly consistent with clustering by k-means, as C1
corresponded to the union of B1 and B2, and C2 corresponded to the union of B3 and B4.

These same three analyses were next performed with codominant markers. The DAPC was
performed retaining 60 PCs explaining ~90% of the observed variance (Supplementary Figure 9A);
the BIC curve indicated k = 3 as the most likely number of clusters in the panel (Supplementary
Figure 9B). Thus, the panel was separated into three groups (D1-3); the plotting of the panel elements
onto the first two linear discriminants from this analysis is shown in Figure 4A. The PCA followed
by k-means was executed assuming the existence of two clusters (E1 and E2), as clearly indicated by
all three testing methods employed (Supplementary Figure 10A-C). Despite indicating the
subdivision of the panel into two clusters, the attribution of genotypes to these groups shared
similarities with the DAPC clustering. E1 corresponded to D2, and E2 was equivalent to D1 and D3
combined. Figure 4B depicts a projection of these groups onto the PCA's first two PCs, which
accounted for ~33% of the variation in the genotypic data. Finally, based on Bayesian clustering,
both ΔK and LnP(D) methods indicated K = 2 as the most likely number of clusters in the panel
(Supplementary Figure 11A and B, respectively). Thus, we present here a plot of the estimated
association coefficients of individuals for these two clusters, named F1 and F2 (Figure 4C).
Clustering by this procedure was also consistent with clustering by the other two methods, as F1
corresponded to the union of D2 and D3, and F2 corresponded to D1.

Despite observing extensive correspondences in clustering results between methods when the same
genotypic data were used, we found that using dominant or codominant markers yielded noticeably
different outcomes. Some overlap was observed between clusters identified by the analyses using
each set of markers (e.g., D2 corresponded roughly to the union of A1 and A2), but overall, groups
identified by these analyses shared little resemblance. Additionally, the results from these methods
did not present correspondences with the phenotype-based HCPCs. Although most accessions in the
HCPC group SQ3 were attributed to the same genotypic cluster in each population structure analysis,
such a pattern was not observed for the other phenotypic groups.

We also estimated pairwise LD between codominant markers and analyzed its decay over distance.
We observed high r^2 values (~0.4) between closely distanced markers, which dropped to 0.1 at
approximately 2 Mb (Figure 5).

3.4   Association Analyses

3.4.1   FarmCPU

For FarmCPU analyses, we tested matrices obtained from each genetic structure analysis as
covariates and ran the models with no covariates. Supplementary Figure 12 depicts a boxplot of the
distribution of λ in each case; these data followed a normal distribution ($p = 0.975$), and an ANOVA
indicated that no significant differences ($p = 0.084$) existed between models. Thus, we chose to
conduct association analyses using no covariates, as this resulted in the median value of λ closest to
its theoretical value under the null hypothesis ($λ = 1$). Appropriate profiles of inflation of p-values
were obtained for the three traits, as shown in the Q-Q plots generated by FarmCPU (Supplementary
Figure 13). Using a Bonferroni-corrected threshold of 0.05, one marker-trait association was detected
for symptom severity, and five associations were detected for the viral titer estimated by each
quantification method – with one marker (MSP_TTG-ECO_ACA-700-0133) being mutually
associated with both (Supplementary Table 9); this could be an indication of the consistency of the
association or simply reflect the correlation between the two measurements. All markers detected to
be trait-associated were AFLP fragments, and the percentage of phenotypic variance explained by
each marker ranged from 9 to 30%.

3.4.2   Mixed Modeling

Twelve combinations of population structure (Q) and kinship (K) matrices were tested as effects in
the association models of the three traits analyzed. A boxplot of the distribution of λ in each Q + K
model is shown in Supplementary Figure 14. A Shapiro-Wilk test indicated that the data did not
follow a normal distribution ($p = 3.253e-06$), and a Kruskal-Wallis test indicated that there were no
significant differences ($p = 0.869$) between the inflation of p-values of the models. Thus, we chose to
conduct the following analyses with a Q + K combination that resulted in the median value of λ
closest to 1, which was obtained with the combination of the first three PCs from a PCA (Q_{PCA}) with
both the realized relationship (K_{MM}) and pseudodiploid (K_{Pseudo}) kinship matrices. Taking into
account that the MM^T matrix can be computed by the GWASpoly package, we considered the use of
the $Q_{PCA} + K_{MM}$ combination to be the most straightforward approach. Q-Q plots of the association
analyses for SCYL symptom severity and SCYLV relative and absolute quantifications can be found
in Supplementary Figure 15. In general, all models showed an appropriate profile of inflation of p-
values.

A stringent significance threshold ($p < 0.05$ corrected by the Bonferroni method) was used to identify
35 nonredundant markers (19 SNPs and 16 indels) that were significantly associated with SCYL
symptom severity (Figure 6). Using this correction, no markers were significantly associated with
SCYLV titer. In an attempt to establish a less conservative threshold for association analyses of these
two traits, we employed the FDR for the correction of p-values, which resulted in very low
significance thresholds and the identification of thousands of associations as significant (data not
shown). Therefore, we ultimately opted to use an arbitrary threshold of $p < 0.0001$ ($-\log_{10}(p) > 4$) to
determine markers strongly associated with the two quantification traits. This threshold resulted in 13
markers (4 SNPs and 9 indels) associated with SCYLV titers determined by relative quantification
and 9 markers (6 SNPs and 3 indels) associated with SCYLV titers determined by absolute
quantification (Figure 6); one marker (S3_5029059) was common to both viral quantification

methods. Supplementary Table 10 supplies information on all marker-trait associations identified by
this approach. For each trait analyzed, we observed a redundancy between markers identified as
significant by the six different marker-effect models employed; this observation was particularly
common between the simplex dominant alternative and the two diploidized models. No markers were
identified as significantly associated with phenotypes using the simplex dominant reference model.

3.4.3   Machine Learning Coupled with Feature Selection

As a last marker-trait association method, we tested eight ML algorithms for predicting the
attribution of genotypes to the phenotypic clusters identified in the two HCPC analyses. When
assessing their potential in this task using the full marker dataset, the predictive accuracies varied
greatly depending on the method and the phenotypic groups under analysis. They were lower for the
prediction of clusters associated with viral titer (Q), ranging from 39.2% with DT to 49.6% with
SVM, with an average of 44.5% (Supplementary Figure 16A). For the clusters identified including
symptom severity data (SQ), accuracies were overall higher, albeit varying even more and being still
unsatisfactory; they then ranged from 7.9% with GNB to 73.9% with SVM (Supplementary Figure
16B) and had an average of 58%. Therefore, we tested applying five different FS methods to reduce
the marker dataset, keeping markers that were more explanatory of the phenotypic clustering results.
Furthermore, we constructed three additional reduced marker datasets consisting of intersections
between FS methods: markers selected by at least two of the five methods (Inter1); markers selected
by at least two of the three best methods (Inter2); and markers selected by all three best methods
(Inter3). We then reassessed the predictive accuracies of all ML algorithms using these new marker
datasets.

After these procedures, we could achieve considerably higher accuracies in predicting Q and SQ
clusters. An ANOVA followed by Tukey's test showed that three of the FS methods tested (FS1, FS2
and FS4) presented notably superior effects in both cases, while the other two (FS3 and FS5) did not
result in substantial increases in accuracy (Supplementary Figure 17). In the two scenarios, the most
accurate model-FS combination was MLP coupled with FS2, which was composed of 232 markers
for Q and 170 markers for SQ. This combination resulted in average accuracies of 97.6% and 96.5%
for the prediction of Q and SQ clusters, respectively (Supplementary Tables 11 and 12). However, in
both scenarios, MLP also achieved the second-best results when using Inter2 datasets, which were
composed of 190 markers for Q and 120 markers for SQ. With this strategy, we could achieve
average accuracies of 95.7% for Q and 95.4% for SQ. As the usage of Inter2 represented an even
further reduction in the number of markers with minimal decreases in the predictive accuracies, we
considered the MLP-Inter2 combination as the most efficient approach to predict clusters associated
with SCYLV resistance. To further evaluate the performance of MLP, we produced ROC curves and
calculated their respective AUCs (Supplementary Figures 18 and 19). Prior to FS, MLP did not
present satisfactory results, with ROC curves very close to the chance level and AUCs of 0.45-0.61
for Q and 0.40-0.56 for SQ (Supplementary Figures 18A and 19A). When Inter2 was used, the ROC
curves showed a much better performance of the model, with AUCs of 1.00 for Q and of 0.98-1.00
for SQ (Supplementary Figures 18B and 19B). These results confirm that the markers present in
Inter2 are in fact associated with SCYLV resistance and that MLP is an appropriate model to predict
clustering based on this marker dataset.

The markers representing the reduced datasets associated with Q and SQ clusters can be found in
Supplementary Tables 13 and 14, respectively. We observed twelve marker overlaps between the two
datasets, i.e., two AFLP fragments, seven SNPs and three indels were present in both of them.
Interestingly, several of these markers were also identified as associated with phenotypes in the

FarmCPU and mixed modeling analyses. This was the case, for instance, of MSP_TTG-ECO_ACA-
700-0133, an AFLP fragment associated with SCYLV titer estimated by relative and absolute
quantification and with Q and SQ clustering, and of S3_5029059, an indel associated with relative
and absolute SCYLV quantification by GWAS that was also present in both the Q and SQ Inter2
datasets.

3.5   Marker Mapping and Annotation

For a better visualization of the physical location of all markers associated with SCYLV resistance,
we constructed a map of their distribution along *S. spontaneum*'s "A" chromosomes (Figure 7), in
which we also included markers identified as associated with SCYLV resistance in previous mapping
studies. Overall, markers were considerably spread along chromosomes; however, we observed
regions of dense concentration of markers identified by various methods, such as the long arms of
chromosomes 1 and 3. We also verified the proximity between several markers identified in the
present work and by other authors, indicating their convergence and the reliability of the methods
employed here.

Out of the 362 nonredundant markers associated with all phenotypes, 176 were located in genic
regions and could be annotated by aligning their 2000-bp neighboring regions with the CDSs of 14
Poaceae species and *A. thaliana* genomes; Supplementary Table 15 contains data on the alignment
with the highest percentage of identity for each marker. In some cases, where two or more markers
were closely located, coincident alignments and annotations were obtained; consequently, 148 genes
were representative of all the best alignments. The large majority of top-scoring alignments (117)
occurred with CDSs of *S. bicolor*, the phylogenetically closest species among those used for
alignment; fewer alignments also occurred with the CDSs of *Z. mays* (19), *P. hallii* (4), *H. vulgare*
(2)*, Oropetium thomaeum* (2), *P. virgatum* (2), *S. viridis* (2), *B. sylvaticum* (1), *S. italica* (1) and *T.*
intermedium (1). All of the annotated markers but one SSR fragment (SCB381.09) consisted of SNPs
or indels. Several of the annotated genes could be associated with plant resistance to viruses, as
detailed in the discussion.

4   Discussion

In the present study, we evaluated the severity of SCYL symptoms and SCYLV titer in a panel of 97
sugarcane accessions. These two traits are of great concern to breeding, as the exhibition of SCYL
symptoms and high viral titers have both been associated with higher yield losses in SCYLV-infected
sugarcane plants (Lehrer et al., 2007; Zhu et al., 2010; Viswanathan et al., 2014). Prior to
phenotyping, plants were subjected to high SCYLV inoculum pressure, an innovation over all
previous mapping studies (Costet et al., 2012; Débibakas et al., 2014; Gouy et al., 2015; Islam et al.,
2018; Yang et al., 2019a), which relied on natural infection under field conditions. Using RT-qPCR,
currently regarded as the most precise method for SCYLV quantification (Zhu et al., 2010), we
assessed the viral titer in these genotypes; together with the study by Villavicencio et al. (in press),
this is also the first time RT-qPCR was used for this purpose in a large number of sugarcane
genotypes. We found a strong and positive correlation between the BLUPs calculated for the SCYLV
titers obtained by the two quantification methods employed, showing the consistency of the data. The
absence of a perfect correlation might have arisen from intrinsic differences between the two
methods, which have been responsible for disparities in viral quantification by RT-qPCR in other
plant-virus interactions (Jarošová et al., 2009).

However, we observed no significant correlation between the BLUPs calculated for the severity of
disease symptoms and viral titers. This finding corroborates a growing body of evidence suggesting

that these traits are not strongly or necessarily correlated, i.e., high SCYLV titers are not a guarantee
of more severe yellowing or of its development at all (Comstock et al., 1998; Lehrer and Komor,
2008; Chinnaraja et al., 2014). Although Zhu et al. (2010) claim to have observed a correlation
between these two parameters, this was not statistically shown. Our results, on the other hand,
demonstrate that no quantitative correlation between the severity of SCYL symptoms and SCYLV
titers exists across the sugarcane genotypes analyzed. This reinforces the importance of SCYLV
molecular screening of sugarcane clones by breeding programs; this should be done to avoid the
employment of genotypes that accumulate high viral loads asymptomatically but may
inconspicuously suffer yield losses, in addition to serving as a virus reservoir for vector transmission
to other susceptible genotypes.

To further explore this issue, we performed two HCPC analyses to discriminate accessions based on
their response to SCYLV infection, which led to the separation of clusters with considerable
phenotypic differences. In the first HCPC, using only data from the two quantification methods, we
could discern groups with significant variation in viral titers; in the second analysis, which also
included symptom severity data, clusters with even more contrasting responses to SCYLV could be
discriminated. In 1983, Cooper and Jones proposed a new terminology addressing plant responses to
viral infections that is still employed by several authors in this area (Beoni et al., 2016; Foresman et
al., 2016; Mansilla-Córdova et al., 2018). According to their proposal, once infected, plants present
differences in the ability to restrict viral replication and invasion; at the extremes of a spectrum of
behaviors are plants termed susceptible (in which the virus rapidly replicates and disperses) and
resistant (in which viral replication and dispersal is slow). In addition, plants may also respond
differently to the infection: another spectrum exists, at the extremes of which are sensitive plants
(which exhibit severe symptoms) and tolerant plants (which exhibit little or no symptoms). In view of
this nomenclature, we propose that the clusters identified in this second HCPC be described by the
three following designations: (SQ1) resistant, for sugarcane genotypes distinguished by low SCYLV
titer and mild or no SCYL symptoms; (SQ2) tolerant, for genotypes that, despite exhibiting higher
viral titers, presented fewer or no disease symptoms; and (SQ3) susceptible, for genotypes with the
most severe symptoms and presenting high viral titers. This classification per se is of great use in
sugarcane breeding, as it distinguishes not only sources of tolerance to SCYLV but also an
exceptionally promising group of genotypes that are truly resistant to this virus.

The main objective of this work was, however, to identify markers associated with SCYLV
resistance in a broader sense. With this aim, we genotyped the plants in the panel with a combination
of dominant and codominant markers, which has never been described for sugarcane to date; this is
also the first study employing a two-enzyme GBS protocol in this crop. We evaluated the impact of
using genomic references from various backgrounds in variant calling. In previous sugarcane
GWASs, variant calling was performed using the genome of *S. bicolor* (Fickett et al., 2019; Yang et
al., 2019a, 2019b, 2020), and until recently, this was sugarcane's closest relative with a well-
assembled and annotated genome; however, in our analyses, this reference yielded a number of
markers considerably inferior to other references, which discouraged its use. The reference that
yielded the largest number of markers was the methyl-filtered genome of the SP70-1143 cultivar, in
agreement with the results of a previous study employing GBS (Balsalobre et al., 2017). This is a
plausible outcome, as the protocol used for library preparation reduces genome complexity by
avoiding sampling of methylated regions (Elshire et al., 2011), which were also filtered out for
assembly of the SP70-1143 genome (Grativol et al., 2014). However, to choose the best reference for
further analyses, we also considered the quality of the assembly, which has been shown to greatly
affect the results of GWASs in polyploids (Benevenuto et al., 2019). The best-assembled sugarcane
genome available to date is the allele-defined genome of a haploid *S. spontaneum* accession (Zhang

et al., 2018). Despite presenting one of the highest total tag alignment rates, this reference also gave a
very high rate of multiple alignments, which led to the identification of a relatively low number of
markers. This was probably due to the alignment of tags to homeologous regions of different alleles
rather than simply to the duplicated regions that we intended to avoid. To circumvent this situation,
we opted to conduct our analyses with markers isolated using a monoploid chromosome set obtained
from this genome, which provided a large number of markers (38,710 SNPs and 32,178 indels) with
reliable position information.

Using these codominant markers, we analyzed the decay of LD over distance. LD has long been
hypothesized to be high in sugarcane due to the short breeding history and narrow genetic basis of
this crop; many studies using dominant markers have estimated it to be especially high at 5-10 cM
(Jannoo et al., 1999; Raboin et al., 2008, Wei et al., 2010; Singh et al., 2016; Barreto et al., 2019).
The first study to use SNPs for this task and estimate LD decay in bp (Yang et al., 2019c) indicated
that LD was extremely long lasting, with the average r^2 decaying to 0.2 at 3.5 Mb in hybrids. Our
results further confirm the persistence of LD at long distances in sugarcane, albeit indicating that it
decayed more quickly – with r^2 dropping to 0.2 at less than 1 Mb and to 0.1 at 2 Mb. These results
impact mapping studies, as a high LD implies that a low density of markers might be needed for
accurate mapping of quantitative traits.

With the two distinct marker datasets generated – including dominant and codominant markers with
AP information – we tested several approaches to evaluate population structure in the panel; these
assessments yielded remarkably different results. Some of the few existing studies contrasting the
usage of dominant and codominant markers in plants have also shown discrepancies in measures of
genetic structure and diversity (Gaudeul et al., 2004; Fang et al., 2006; Roncallo et al., 2019).
However, this sort of comparison has never been performed including markers with AD or AP
information in polyploids, let alone in sugarcane, which has one the most complex genomes known.
In this crop, the most relatable findings available are those reported by Creste et al. (2010), who
compared the effect of using dominant AFLP, SSR and target region amplification polymorphism
markers in the estimation of genetic relationships in sugarcane. Alike we demonstrate now, these
authors showed that using different types of markers can bias genetic analyses in this crop and that
thus the choice of marker to be used must be guided by the specific goal of each study. For genome-
wide mapping of quantitative traits – for which a high density of markers is necessary – codominant
SNPs and indels are currently more cost-effective, as they can be easily identified in much larger
numbers. Additionally, in the case of polyploids, these markers offer the possibility of estimating
allele dosages or APs, which are highly informative for genetic studies (de Bem Oliveira, 2019;
Matias et al., 2019; Aono et al., 2020). Hence, we believe the results we obtained with codominant
SNPs and indels are more reliable, as they rely on much larger amounts of genetic information.

In contrast with the differences arisen from the type of marker used, we observed little divergence
between the results of the methods used for each marker dataset; the separation of genotypes into
groups was overall similar, with consistent correspondences between groups identified in each
analysis. Additionally, eventual discrepancies between methods did not result in significant
differences in the inflation of the models used for both FarmCPU and codominant mixed modeling
analyses; the inflation patterns observed in Q-Q plots were similar to those of previous studies
applying similar genotyping and marker-trait association methodologies (Barreto et al., 2019; Yang
et al., 2019a, 2019b, 2020). For both association mapping strategies, we opted to perform ensuing
association analyses using the covariates that resulted in the value of λ closest to 1. For FarmCPU,
this corresponded to the "naive" model with no covariates; for codominant mixed modeling analysis,
this was the $Q_{PCA} + K_{MM}$ combination. K_{MM} is the usual choice of relationship matrix in polyploid

association mapping (Rosyara et al., 2016; Berdugo-Cely et al., 2017; Byrne et al., 2020), as Q
matrices obtained from PCA are commonly used to control population structure in GWASs
(Nimmakayala et al., 2016; Su et al., 2016; Ferrão et al., 2018).

FarmCPU analyses using dominant markers identified one AFLP fragment significantly associated
with symptom severity, which explained a small part of the phenotypic variation ($r^2 = 0.116$). Eight
out of the nine markers found to be associated with viral titer explained larger parts of the variation in
the phenotypes (21-30%). These results are more promising than those obtained in a previous GWAS
employing dominant markers in the investigation of SCYLV resistance, which found markers
explaining smaller percentages (9-14%) of the variation observed (Débibakas et al., 2014). Albeit
low, values in this range are, in fact, the rule rather than the exception in sugarcane association
studies. Evidence indicates that almost all of this crop's traits are highly quantitative, with the notable
exception of brown rust resistance, for which two major QTLs are known (Daugrois et al., 1996;
Raboin et al., 2006). For other relevant traits, such as yield, sugar or fiber content and resistance to
other diseases, it is common to find the most associated markers explaining $\leq 10\%$ of the phenotypic
variation (Gouy et al., 2013; Barreto et al., 2019; Fickett et al., 2019).

A few authors have suggested that these suboptimal results could be improved with the usage of
markers with dosage; this was also performed here, using SNPs and indels with AP information for
codominant mixed modeling analyses. Although we successfully identified markers associated with
SCYL symptom severity using the Bonferroni correction, the same was not observed for SCYLV
quantification data obtained by RT-qPCR. This was probably influenced by the limited size of the
panel, a factor that restricts the power of GWASs (Wang et al., 2012; Yang et al., 2014). As pointed
out by Racedo et al. (2016), assembling and phenotyping large sugarcane association panels is a
challenging task, and this was especially burdensome in our study, in which laborious inoculation
and quantification techniques were employed. Furthermore, the Bonferroni method is notorious for
its conservative nature, which contributes to a low capacity for controlling false negatives (Gao et al.,
2010; Johnson et al., 2010; Fadista et al., 2016). This led us to establish an arbitrary threshold ($p <$
0.0001) to select markers strongly associated with SCYLV titer for further investigation. Using this
methodology, we identified 57 nonredundant markers associated with the three phenotypes.

Finally, as a last approach to identify marker-trait associations, we tested several ML algorithms
coupled with FS methods to predict genotype attribution to phenotypic clusters identified by HCPC
analyses. We could achieve very high accuracies of prediction (95%) with considerably reduced
marker datasets comprising 120-190 SNPs and indels. These results are very similar to what was
obtained for predicting sugarcane brown rust resistance groups using the same methods, where an
accuracy of 95% was obtained using 131 SNPs (Aono et al., 2020). Marker datasets selected by ML
have rarely been employed in genetic association studies in plants, but the few existing examples
show their power to identify genes associated with phenotypes of interest (Steinfath et al., 2009; Heer
et al., 2018; Zhou and Bellis, 2019).

Here, we annotated 176 markers associated with SCYLV resistance to 148 genes. Many of these
candidates do not allow extensive discussion on their involvement in resistance to this viral disease,
as they either have very generic descriptions (e.g., "expressed protein") or have not been previously
linked to plant virus resistance. Other proteins, such as helix-loop-helix proteins or
glycosyltransferases, have occasionally been associated with responses to viruses but are members of
very large gene families with extremely diverse biological roles. Thus, we will focus on discussing
genes whose involvement in disease resistance is better established.

Remarkably, a few candidates harboring markers associated with SCYL resistance encode proteins
previously associated with the response to SCYLV infection. This was the case for
SbRio.10G317500.1, encoding a peroxidase precursor and associated with SQ clustering.
Peroxidases are long known to be activated in response to pathogens, but most notably, there is
evidence that a guaiacol peroxidase is more active in sugarcane plants exhibiting SCYL symptoms
than in uninfected or asymptomatic plants (Scagliusi et al., 2009). Our results provide further
evidence that these enzymes are in fact involved in the response to SCYLV and in the development
of yellowing symptoms. Three other candidates harboring markers associated with SCYL symptom
severity and SQ clustering encode proteins with motifs that, despite not being commonly linked to
defense against viruses, were also found to be associated with SCYLV resistance in a previous
GWAS (Yang et al., 2019a). These are Sobic.001G023900, encoding a GATA zinc finger protein,
and Sobic.001G200200 and Zm00001d037864_T030, which encode proteins containing
tetratricopeptide repeats.

Other annotations included classic participants in more general disease resistance mechanisms, such
as several genes encoding proteins with leucine-rich repeat (LRR) motifs. These structures are an
important part of nucleotide-binding LRR (NBS-LRR) proteins, receptors that detect pathogen-
associated proteins and elicit effector-triggered immunity in plants (DeYoung and Innes, 2006).
NBS-LRR proteins represent over 80% of all plant disease resistance (R) genes known (Dubey and
Singh, 2018) and have been shown to determine resistance to viruses in several crops (Dinesh-Kumar
et al., 2000; Seo et al., 2006; Xun et al., 2019). In our analyses, we found one LRR protein
(Sobic.008G156600.1) associated with SCYLV titer determined by relative quantification, one
disease resistance NBS-LRR (Sobic.007G085400.1) and one N-terminal leucine zipper NBS-LRR
resistance gene analog (Sobic.005G203500.1) associated with Q clustering, and an additional LRR
protein (Sobic.001G452600.1) associated with SQ clustering. Furthermore, we annotated one gene
(Sobic.009G204800.1) associated with Q clustering that encodes a precursor of a receptor-like
serine/threonine-protein kinase, i.e., the family to which LRR proteins belong. This is similar to a
finding by Yang et al. (2019), who identified a serine/threonine-protein kinase associated with
SCYLV resistance. We consider these proteins to be highly promising candidates to be involved in
the recognition of infection by SCYLV, which could trigger response mechanisms leading to the
restriction of the virus. Further virus-host interaction studies involving these proteins might help
confirm this hypothesis, which would represent a major breakthrough in understanding resistance to
SCYLV.

Two other annotated genes associated with Q and SQ clustering were readily identified as involved
in plant disease resistance mechanisms. Sobic.010G131300.2 contains a Bric-a-Brac, Tramtrack,
Broad Complex/Pox virus and Zinc finger (BTB/POZ) domain, while Sobic.007G198400.1 contains
two BTB domains, as well as ankyrin repeat regions. These domains are present in and are essential
for the function of NONEXPRESSOR OF PATHOGENESIS-RELATED GENES 1 (NPR1), a
central player in plant disease responses (Cao et al., 1997; Rochon et al., 2006). This family of
transcription factors is involved in establishing both systemic acquired resistance and induced
systemic resistance (Pieterse et al., 1998), mediating the crosstalk between salicylic acid and
jasmonic acid/ethylene responses (Spoel et al., 2003). Correspondingly, NPR1 has been widely
shown to be involved in resistance to viruses (Liu et al., 2002; Lin et al., 2004); therefore, it is
reasonable to suggest its participation in the response to infection by SCYLV.

We also found a few candidates with putative roles in the RNA interference mechanism, one of the
most prominent processes that contribute to resistance against viruses in plants. This is the case for
Sobic.001G214000.1, associated with SQ clustering, which encodes a Dicer. Dicers are part of a

mechanism known as RNA silencing, where they are responsible for the recognition and cleavage of
long double-stranded RNA molecules into mature small RNAs, which in turn guide the site-directed
cleavage of viral mRNAs that disrupt virus replication (Llave, 2010); accordingly, these nucleases
have been linked to resistance to viruses in several plant species (Deleris, 2006; Alam et al., 2019).
Another gene possibly involved in RNA interference is Sobic.009G121100, which harbors a marker
associated with SCYLV titer determined by absolute quantification. This gene encodes a protein
related to calmodulin binding – a calcium transducer that regulates the activity of various proteins
with diverse functions (Bouché et al., 2005) and has been widely implicated in viral resistance in
plants, often playing roles in RNA interference (Heo et al., 1999; Nakahara et al., 2012; Li et al.,
2014). Therefore, we consider these genes promising candidates in the regulation of SCYLV
replication and spreading *in planta*, as well as in the development of SCYL symptoms.

Two additional annotations linked to the mechanism of RNA interference are those of genes
encoding proteins with F-box domains, SbRio.03G158900 and Sobic.002G019750.1. These are
motifs with ca. 50 amino acids that occur in large numbers in plant genomes (Correa et al., 2013). F-
box proteins are involved in virus resistance in several plant species (Cao et al., 2008; Thiel et al.,
2012); a particularly interesting case is FBW2 from *A. thaliana*, which regulates AGO1, an
Argonaute protein with a central role in RNA silencing (Earley et al., 2010) and posttranscriptional
repression of target viral RNAs through a sequence homology-dependent mechanism (Morel et al.,
2002; Chen et al., 2015; Yang and Li, 2018). Even more intriguing is the fact that one of the proteins
encoded by the SCYLV genome, P0, contains an F-box-like domain and mediates the destabilization
of AGO1, leading to the suppression of host posttranscriptional gene silencing (Mangwende et al.,
2009). Whether the F-box proteins identified here play active roles in the mechanisms involved in
posttranscriptional gene silencing of SCYLV replication in sugarcane and in the manifestation of
symptoms remains a question to be investigated by further studies.

Several other annotated genes may represent host factors involved in various steps of plant-virus
interactions. For instance, Sobic.010G160500.4, associated with Q clustering, encodes an RNA
helicase with a DEAD-box domain; these host proteins are often coopted by viruses to promote viral
translation or replication, playing important roles in regulating infection (Huang et al., 2010; Kovalev
et al., 2012; Li et al., 2016). Similarly, soluble N-ethylmaleimide-sensitive-factor attachment protein
receptor (SNARE) proteins, characterized by the motif present in Q clustering-associated
Sobic.001G528000.1, are essential in the biogenesis and fusion of vesicles involved in intercellular
replication and movement of several plant viruses (Wei et al., 2013; Cabanillas et al., 2018; Sasvari
et al., 2018; Ibrahim et al., 2020). We also found one gene encoding a myosin (Sobic.002G108000.1)
associated with Q clustering and two genes related to kinesin (Sobic.001G346600.1 and
Sobic.001G399200.2) associated with SQ clustering. These are two classes of filament-associated
motor proteins that influence numerous biological processes due to their role in the transport of
organelles (Heinlein, 2015). In a few cases, both myosins (Harries et al., 2009; Amari et al., 2011,
2014) and kinesins (Abdelkhalek et al., 2019) have been shown to be involved in viral intercellular
movement, but the mechanisms are currently unclear. One last interesting annotation was
Sobic.003G101500.1, a protein with a DNAJ domain associated with Q clustering. DNAJs have been
shown to interact with proteins of various plant viruses and to be associated with resistance,
sometimes being crucial for virus infection and spread (Hofius et al., 2007; Lu et al., 2009; Shimizu
et al., 2009; Liu et al., 2013). We consider these genes to be promising candidates as host cofactors in
the response to SCYLV infection.

In conclusion, this array of genome-wide analyses allowed us to detect markers significantly
associated with SCYLV resistance in sugarcane. If validated, these markers represent an especially

valuable resource for sugarcane breeding programs, as the results can be directly employed in
marker-assisted strategies for the early selection of clones. The annotation of several genes wherein
these markers are located revealed many candidates with long-established and pivotal roles in viral
disease resistance, further demonstrating the efficiency of the methods employed for this purpose.
Additionally, this annotation provides valuable insights into the unexplored mechanisms possibly
involved in sugarcane's response to infection by SCYLV, introducing new candidates whose role in
this process can be further investigated in future studies.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

Author Contributions

DP, MGAL, MCG, LRP and APS conceived the project and designed the experiments. RJGP,
RCBV, CCS, IAA and LRP performed phenotyping. RJGP, RCBV and AEC performed genotyping.
RJGP and AHA analyzed the data and interpreted the results. RJGP wrote the manuscript. All
authors read and approved the manuscript.

Funding

This work was supported by grants from the São Paulo Research Foundation (FAPESP), the
Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPQ, grant 424050/2016-1), the
Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES, Computational Biology
Program), the Littoral Polytechnic Superior School (ESPOL) and the Secretaría Nacional de Ciencia
y Tecnología (SENESYT). RJGP received an MSc fellowship from CAPES (grant
88887.177386/2018-00) and MSc and PhD fellowships from FAPESP (grants 2018/18588-8 and
2019/21682-9). AHA received a PhD fellowship from FAPESP (grant 2019/03232-6). RCBV
received a PhD fellowship from PAEDEx-AUIP. CCS received a PD fellowship from FAPESP
(grant 2015/24346-9).

Abbreviations

AB: adaptive boosting

AFLP: amplified fragment length polymorphism

ANOVA: analysis of variance

AP: allele proportion

AUC: area under the receiver operating characteristic curve

BIC: Bayesian information criterion

BLUP: best linear unbiased predictor

BTB/POZ: Broad Complex/Pox virus and Zinc finger

CDS: coding sequence

CT: cycle threshold

DAPC: discriminant analysis of principal components

DT: decision tree

FarmCPU: fixed and random model circulating probability unification

FDR: false discovery rate

FS: feature selection

GBS: genotyping-by-sequencing

GNB: Gaussian naive Bayes

GP: Gaussian process

GWAS: genome-wide association study

HCPC: hierarchical clustering on principal components

Indel: insertion and deletion

KNN: K-nearest neighbor

LD: linkage disequilibrium

LRR: leucine-rich repeat

MAF: minimum allele frequency

ML: machine learning

MLP: multilayer perceptron

NBS: nucleotide-binding site

NPR1: NONEXPRESSOR OF PATHOGENESIS-RELATED GENES 1

PC: principal component

PCA: principal component analysis

Q-Q: quantile-quantile

QTL: quantitative trait locus

RF: random forest

ROC: receiver operating characteristic

RT-qPCR: reverse transcription quantitative polymerase chain reaction

SCYL: sugarcane yellow leaf

SCYLV: sugarcane yellow leaf virus

SNARE: soluble N-ethylmaleimide-sensitive-factor attachment protein receptor

SNP: single nucleotide polymorphism

SSR: simple sequence repeat

SVM: support vector machine

TVDL: top visible dewlap leaf

Acknowledgments

We thank Aline C. L. Moraes for assistance in constructing and sequencing the GBS library and
Maicon Volpin for assistance in fieldwork.

References

Abdelkhalek, A., Ismail, I. A., Dessoky, E. S., El-Hallous, E. I., & Hafez, E. (2019). A tomato kinesin-like protein is associated with Tobacco mosaic virus infection. Biotechnology & Biotechnological Equipment, 33(1), 1424-1433. doi: 10.1080/13102818.2019.1673207

Alam, C. M., Jain, G., Kausar, A., Singh, A. K., Mandal, B., Varma, A.... & Chakraborty, S. (2019). Dicer 1 of Candida albicans cleaves plant viral dsRNA in vitro and provides tolerance in plants against virus infection. Virus Disease, 30(2), 237-244. doi: 10.1007/s13337-019-00520-x

Aljanabi, S. M., Forget, L., & Dookun, A. (1999). An improved and rapid protocol for the isolation of polysaccharide- and polyphenol-free sugarcane DNA. Plant Molecular Biology Reporter, 17(3), 281-282. doi: 10.1023/A:1007692929505

Aljanabi, S. M., Parmessur, Y., Moutia, Y., Saumtally, S., and Dookun, A. (2001). Further evidence of the association of a phytoplasma and a virus with yellow leaf syndrome in sugarcane. Plant Pathol. 50, 628-636. doi: 10.1046/j.1365-3059.2001.00604.x

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., and Lipman, D. J. (1990). Basic local alignment search tool. J. Mol. Biol. 215, 403-410. doi: 10.1016/s0022-2836(05)80360-2

Amadeu, R. R., Cellon, C., Olmstead, J. W., Garcia, A. A. F., Resende, M. F. R., and Muñoz, P. R. (2016). AGHmatrix: R package to construct relationship matrices for autotetraploid and diploid species: a blueberry example. Plant Genome 9, 1-10. doi: 10.3835/plantgenome2016.01.0009

Amari, K., Lerich, A., Schmitt-Keichinger, C., Dolja, V. V., & Ritzenthaler, C. (2011). Tubule-guided cell-to-cell movement of a plant virus requires class XI myosin motors. PLoS Pathogens, 7(10), e1002327. doi: 10.1371/journal.ppat.1002327

Amari, K., Di Donato, M., Dolja, V. V., & Heinlein, M. (2014). Myosins VIII and XI play distinct roles in reproduction and transport of tobacco mosaic virus. PLoS Pathogens, 10(10), e1004448. doi: 10.1371/journal.ppat.1004448

Andrews, S. (2010). FastQC: A Quality Control Tool for High Throughput Sequence Data. Babraham: Babraham Bioinformatics.

Aono, A. H., Costa, E. A., Rody, H. V. S., Nagai, J. S., Pimenta, R. J. G., Mancini, M. C., dos Santos, F. R. C., Pinto, L. R., Landell, M. G. de A., de Souza, A. P., & Kuroshu, R. M. (2020). Machine learning approaches reveal genomic regions associated with sugarcane brown rust resistance. Scientific Reports, 10. doi:10.1038/s41598-020-77063-5

Balsalobre, T. W. A., da Silva Pereira, G., Margarido, G. R. A., Gazaffi, R., Barreto, F. Z., Anoni, C. O.... and de Souza, A. P. (2017). GBS-based single dosage markers for linkage and QTL mapping allow gene mining for yield-related traits in sugarcane. BMC Genomics, 18, 72. doi: 10.1186/s12864-016-3383-x

Barreto, F. Z., Rosa, J. R. B. F., Balsalobre, T. W. A., Pastina, M. M., Silva, R. R., Hoffmann, H. P.... & Carneiro, M. S. (2019). A genome-wide association study identified loci for yield

component traits in sugarcane (*Saccharum* spp.). PloS One, 14(7), e0219843. doi:
10.1371/journal.pone.0219843

Benevenuto, J., Ferrão, L. F. V., Amadeu, R. R., and Munoz, P. (2019). How can a high-quality
genome assembly help plant breeders? GigaScience 8, giz068. doi:
10.1093/gigascience/giz068

Beoni, E., Chrpová, J., Jarošová, J., and Kundu, J. K. (2016). Survey of Barley yellow dwarf virus
incidence in winter cereal crops, and assessment of wheat and barley resistance to the virus.
Crop Pasture Sci. 67, 1054-1063. doi: 10.1071/cp16167

Berdugo-Cely, J., Valbuena, R. I., Sánchez-Betancourt, E., Barrero, L. S., and Yockteng, R. (2017).
Genetic diversity and association mapping in the Colombian Central Collection of *Solanum*
tuberosum L. Andigenum group using SNPs markers. PLoS One 12, e0173039. doi:
10.1371/journal.pone.0173039

Bouché, N., Yellin, A., Snedden, W. A., and Fromm, H. (2005). Plant-specific calmodulin-binding
proteins. Annu. Rev. Plant Biol. 56, 435-466. doi:
10.1146/annurev.arplant.56.032604.144224

Bourke, P. M., Voorrips, R. E., Visser, R. G. F., and Maliepaard, C. (2018). Tools for genetic studies
in experimental populations of polyploids. Front. Plant Sci. 9, 513. doi:
10.3389/fpls.2018.00513

Breiman, L. (2001). Random forests. Machine Learning, 45(1), 5-32. doi: 10.1023/A:1010933404324

Bremer, G. (1961). Problems in breeding and cytology of sugar cane. Euphytica 10, 59-78. doi:
10.1007/bf00037206

Byrne, S., Meade, F., Mesiti, F., Griffin, D., Kennedy, C., and Milbourne, D. (2020). Genome-wide
association and genomic prediction for fry color in potato. Agronomy 10, 90. doi:
10.3390/agronomy10010090

Cabanillas, D. G., Jiang, J., Movahed, N., Germain, H., Yamaji, Y., Zheng, H., & Laliberté, J. F.
(2018). Turnip mosaic virus uses the SNARE protein VTI11 in an unconventional route for
replication vesicle trafficking. The Plant Cell, 30(10), 2594-2615. doi: 10.1105/tpc.18.00281

Cao, H., Glazebrook, J., Clarke, J. D., Volko, S., & Dong, X. (1997). The *Arabidopsis* NPR1 gene
that controls systemic acquired resistance encodes a novel protein containing ankyrin repeats.
Cell, 88(1), 57-63. doi: 10.1016/S0092-8674(00)81858-9

Cao, Y., Yang, Y., Zhang, H., Li, D., Zheng, Z., and Song, F. (2008). Overexpression of a rice
defense-related F-box protein geneOsDRF1in tobacco improves disease resistance through
potentiation of defense gene expression. Physiol. Plant. 134, 440-452. doi: 10.1111/j.1399-
3054.2008.01149.x

Cardoso-Silva, C. B., Costa, E. A., Mancini, M. C., Balsalobre, T. W. A., Canesin, L. E. C., Pinto, L.
R., et al. (2014). De novo assembly and transcriptome analysis of contrasting sugarcane
varieties. PLoS One 9, e88462. doi: 10.1371/journal.pone.0088462

Carvalho-Netto, O. V., Bressiani, J. A., Soriano, H. L., Fiori, C. S., Santos, J. M., Barbosa, G. V. S.,
993   et al. (2014). The potential of the energy cane as the main biomass crop for the cellulosic
994   industry. Chem. Biol. Technol. Agric. 1, 20. doi: 10.1186/s40538-014-0020-2

995   Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A., and Cresko, W. A. (2013). Stacks: an
996   analysis tool set for population genomics. Mol. Ecol. 22, 3124-3140. doi: 10.1111/mec.12354

997   Chatenet, M., Delage, C., Ripolles, M., Irey, M., Lockhart, B. E. L., and Rott, P. (2001). Detection of
998   sugarcane yellow leaf virus in quarantine and production of virus-free sugarcane by apical
999   meristem culture. Plant Dis. 85, 1177-1180. doi: 10.1094/pdis.2001.85.11.1177

1000   Chen, H., Zhang, L., Yu, K., and Wang, A. (2015). Pathogenesis of Soybean mosaic virus in soybean
1001   carrying Rsv1 gene is associated with miRNA and siRNA pathways, and breakdown of
1002   AGO1 homeostasis. Virology 476, 395-404. doi: 10.1016/j.virol.2014.12.034

1003   Chen, T., & Guestrin, C. (2016). Xgboost: A scalable tree boosting system. In Proceedings of the
1004   22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining
1005   (pp. 785-794). doi: 10.1145/2939672.2939785

1006   Chhatre, V. E., & Emerson, K. J. (2017). StrAuto: automation and parallelization of STRUCTURE
1007   analysis. BMC Bioinformatics, 18, 1-5. doi: 10.1186/s12859-017-1593-0

1008   Chinnaraja, C., and Viswanathan, R. (2015). Quantification of sugarcane yellow leaf virus in
1009   sugarcane following transmission through aphid vector, *Melanaphis sacchari*. Virus Disease
1010   26, 237-242. doi: 10.1007/s13337-015-0267-7

1011   Chinnaraja, C., Viswanathan, R., Sathyabhama, M., Parameswari, B., Bagyalakshmi, K., Malathi, P.,
1012   et al. (2014). Quantification of sugarcane yellow leaf virus in in vitro plantlets and
1013   asymptomatic plants of sugarcane by RT-qPCR. Curr. Sci. 106, 729-734

1014   Comstock, J. C., Irey, M. S., Lockhart, B. E. L., and Wang, Z. K. (1998). Incidence of yellow leaf
1015   syndrome in CP cultivars based on polymerase chain reaction and serological techniques.
1016   Sugar Cane 4, 21-24

1017   Cooper, J. I., and Jones, A. T. (1983). Responses of plants to viruses: proposals for the use of terms.
1018   Phytopathology 73, 127-128. doi: 10.1094/phyto-73-127

1019   Correa, R. L., Bruckner, F. P., de Souza Cascardo, R., and Alfenas-Zerbini, P. (2013). The role of F-
1020   box proteins during viral infection. International journal of molecular sciences, 14, 4030-
1021   4049. doi: 10.3390/ijms14024030

1022   Costet, L., Raboin, L.-M., Payet, M., D'Hont, A., and Nibouche, S. (2012). A major quantitative trait
1023   allele for resistance to the sugarcane yellow leaf virus (Luteoviridae). Plant Breed. 131, 637-
1024   640. doi: 10.1111/j.1439-0523.2012.02003.x

1025   Cover, T., & Hart, P. (1967). Nearest neighbor pattern classification. IEEE transactions on
1026   information theory, 13(1), 21-27. doi: 10.1109/TIT.1967.1053964

Creste, S., Sansoli, D. M., Tardiani, A. C. S., Silva, D. N., Gonçalves, F. K., Fávero, T. M.... & Pinto,
L. R. (2010). Comparison of AFLP, TRAP and SSRs in the estimation of genetic
relationships in sugarcane. Sugar Tech, 12(2), 150-154. doi: 10.1007/s12355-010-0029-1

Cristianini, N., & Shawe-Taylor, J. (2000). An introduction to support vector machines and other
kernel-based learning methods. Cambridge University Press. doi:
10.1017/CBO9780511801389

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A.... and McVean, G.
(2011). The variant call format and VCFtools. Bioinformatics, 27, 2156-2158. doi:
10.1093/bioinformatics/btr330

Daugrois, J. H., Grivet, L., Roques, D., Hoarau, J. Y., Lombard, H., Glaszmann, J. C., & D'Hont, A.
(1996). A putative major gene for rust resistance linked with a RFLP marker in sugarcane
cultivar 'R570'. Theoretical and Applied Genetics, 92(8), 1059-1064. doi:
10.1007/BF00224049

Deleris, A., Gallego-Bartolome, J., Bao, J., Kasschau, K. D., Carrington, J. C., & Voinnet, O. (2006).
Hierarchical action and inhibition of plant Dicer-like proteins in antiviral defense. Science,
313(5783), 68-71. doi: 10.1126/science.1128214

DeYoung, B. J., & Innes, R. W. (2006). Plant NBS-LRR proteins in pathogen sensing and host
defense. Nature Immunology, 7(12), 1243-1249. doi: 10.1038/ni1410

D'Hont, A., Ison, D., Alix, K., Roux, C., and Glaszmann, J. C. (1998). Determination of basic
chromosome numbers in the genus Saccharum by physical mapping of ribosomal RNA genes.
Genome 41, 221-225. doi: 10.1139/g98-023

D'Hont, A., and Glaszmann, J. C. (2001). Sugarcane genome analysis with molecular markers: a first
decade of research, in Proceedings of the International Society of Sugarcane Technologists
XXVI Congress, ed. D. M. Hogarth (Brisbane, Australia: The Australian Society of Sugar
Cane Technologists), 556-559.

De Bem Oliveira, I., Resende, M. F., Ferrão, L. F. V., Amadeu, R. R., Endelman, J. B., Kirst, M.... &
Munoz, P. R. (2019). Genomic prediction of autotetraploids; influence of relationship
matrices, allele dosage, and continuous genotyping calls in phenotype prediction. G3: Genes,
Genomes, Genetics, 9(4), 1189-1198. doi: 10.1534/g3.119.400059

De Mendiburu, F., & De Mendiburu, M. F. (2020). Package 'agricolae'. R package version, 1-2.

Débibakas, S., Rocher, S., Garsmeur, O., Toubi, L., Roques, D., D'Hont, A., et al. (2014).
Prospecting sugarcane resistance to sugarcane yellow leaf virus by genome-wide association.
Theor. Appl. Genet. 127, 1719-1732. doi: 10.1007/s00122-014-2334-7

Delage, C., Rippoles, M., Chatenet, M., Irey, M., and Rott, P. (1999). Elimination of sugarcane
yellow leaf virus from sugarcane by meristem tip culture, in Proceedings of the XXIII
Congress of the International Society of Sugar Cane Technologists. New Delhi, India: ISSCT
Congress (Vol. 404).

Devlin, B., and Roeder, K. (1999). Genomic control for association studies. Biometrics 55, 997-1004.
1065   doi: 10.1111/j.0006-341x.1999.00997.x

1066   Dinesh-Kumar, S. P., Tham, W. H., & Baker, B. J. (2000). Structure–function analysis of the tobacco
1067   mosaic virus resistance gene N. Proceedings of the National Academy of Sciences, 97(26),
1068   14789-14794. doi: 10.1073/pnas.97.26.14789

1069   Dinno, A. (2017). dunn.test: Dunn's test of multiple comparisons using rank sums. R package
1070   version, 1(4).

1071   Dubey, N., & Singh, K. (2018). Role of NBS-LRR proteins in plant defense. In Molecular Aspects of
1072   Plant-Pathogen Interaction (pp. 115-138). Springer, Singapore. doi: 10.1007/978-981-10-
1073   7371-7_5

1074   Earl, D. A., and vonHoldt, B. M. (2012). Structure harvester: a website and program for visualizing
1075   structure output and implementing the Evanno method. Conserv. Genet. Resour. 4, 359-361.
1076   doi: 10.1007/s12686-011-9548-7

1077   Earley, K., Smith, M., Weber, R., Gregory, B., and Poethig, R. (2010). An endogenous F-box protein
1078   regulates ARGONAUTE1 in *Arabidopsis thaliana*. Silence 1, 15. doi: 10.1186/1758-907X-1-
1079   15

1080   ElSayed, A. I., Komor, E., Boulila, M., Viswanathan, R., and Odero, D. C. (2015). Biology and
1081   management of sugarcane yellow leaf virus: an historical overview. Arch. Virol. 160, 2921-
1082   2934. doi: 10.1007/s00705-015-2618-5

1083   Elshire, R. J., Glaubitz, J. C., Sun, Q., Poland, J. A., Kawamoto, K., Buckler, E. S., and Mitchell, S.
1084   E. (2011). A robust, simple genotyping-by-sequencing (GBS) approach for high diversity
1085   species. PloS one, 6(5), e19379. doi: 10.1371/journal.pone.0019379

1086   Evanno, G., Regnaut, S., and Goudet, J. (2005). Detecting the number of clusters of individuals using
1087   the software structure: a simulation study. Mol. Ecol. 14, 2611-2620. doi: 10.1111/j.1365-
1088   294x.2005.02553.x

1089   Fadista, J., Manning, A. K., Florez, J. C., and Groop, L. (2016). The (in)famous GWAS P-value
1090   threshold revisited and updated for low-frequency variants. Eur. J. Hum. Genet. 24, 1202-
1091   1205. doi: 10.1038/ejhg.2015.269

1092   Fang, J., Twito, T., Zhang, Z., & Chao, C. T. (2006). Genetic relationships among fruiting-mei
1093   (*Prunus mume* Sieb. et Zucc.) cultivars evaluated with AFLP and SNP markers. Genome,
1094   49(10), 1256-1264. doi: 10.1139/g06-097

1095   FAO (2020). FAOSTAT: Production Sheet. Rome: FAO.

1096   Ferrão, L. F. V., Benevenuto, J., Oliveira, I. D. B., Cellon, C., Olmstead, J., Kirst, M., et al. (2018).
1097   Insights into the genetic basis of blueberry fruit-related traits using diploid and polyploid
1098   models in a GWAS context. Front. Ecol. Evol. 6, 107. doi: 10.3389/fevo.2018.00107

1099   Fickett, N., Gutierrez, A., Verma, M., Pontif, M., Hale, A., Kimbeng, C., & Baisakh, N. (2019).
1100   Genome-wide association mapping identifies markers associated with cane yield components

and sucrose traits in the Louisiana sugarcane core collection. Genomics, 111(6), 1794-1801.
doi: 10.1016/j.ygeno.2018.12.002

Fitch, M. M. M., Lehrer, A. T., Komor, E., and Moore, P. H. (2001). Elimination of sugarcane
yellow leaf virus from infected sugarcane plants by meristem tip culture visualized by tissue
blot immunoassay. Plant Pathol. 50, 676-680. doi: 10.1046/j.1365-3059.2001.00639.x

Foresman, B. J., Oliver, R. E., Jackson, E. W., Chao, S., Arruda, M. P., and Kolb, F. L. (2016).
Genome-wide association mapping of barley yellow dwarf virus tolerance in spring oat
(*Avena sativ*a L.). PLoS One 11, e0155376. doi: 10.1371/journal.pone.0155376

Freund, Y., & Schapire, R. E. (1997). A decision-theoretic generalization of on-line learning and an
application to boosting. Journal of computer and system sciences, 55(1), 119-139. doi:
10.1006/jcss.1997.1504

Friedman, N., Geiger, D., & Goldszmidt, M. (1997). Bayesian network classifiers. Machine
Learning, 29(2-3), 131-163. doi: 10.1023/A:1007465528199

Gerard, D. (2020). Pairwise Linkage Disequilibrium Estimation for Polyploids. bioRxiv. doi:
10.1101/2020.08.03.234476.

Gao, X., Becker, L. C., Becker, D. M., Starmer, J. D., and Province, M. A. (2010). Avoiding the high
Bonferroni penalty in genome-wide association studies. Genet. Epidemiol. 34, 100-105. doi:
10.1002/gepi.20430

Garcia, A. A. F., Mollinari, M., Marconi, T. G., Serang, O. R., Silva, R. R., Vieira, M. L. C., et al.
(2013). SNP genotyping allows an in-depth characterisation of the genome of sugarcane and
other complex autopolyploids. Sci. Rep. 3, 3399. doi: 10.1038/srep03399

Garsmeur, O., Droc, G., Antonise, R., Grimwood, J., Potier, B., Aitken, K., et al. (2018). A mosaic
monoploid reference sequence for the highly complex genome of sugarcane. Nat. Commun.
9, 2638. doi: 10.1038/s41467-018-05051-5

Gaudeul, M., Till-Bottraud, I., Barjon, F., & Manel, S. (2004). Genetic diversity and differentiation
in *Eryngium alpinum* L. (Apiaceae): comparison of AFLP and microsatellite markers.
Heredity, 92(6), 508-518. doi: 10.1038/sj.hdy.6800443

Geurts, P., Ernst, D., & Wehenkel, L. (2006). Extremely randomized trees. Machine Learning, 63(1),
3-42. doi: 10.1007/s10994-006-6226-1

Glaubitz, J. C., Casstevens, T. M., Lu, F., Harriman, J., Elshire, R. J., Sun, Q., et al. (2014).
TASSEL-GBS: a high capacity genotyping by sequencing analysis pipeline. PLoS One 9,
e90346. doi: 10.1371/journal.pone.0090346

Gonçalves, M.C.; Klerks, M.M.; Verbeek, M.; Vega, J.; Van Den Heuvel, J.F.J.M. (2002). The use of
molecular beacons combined with NASBA for the sensitive detection of Sugarcane yellow
leaf virus. Eur. J. Plant Pathol. 08, 401-407. doi: 10.1023/a:1016040314260.

Gonçalves, M. C. (2007). Doenças causadas por vírus, in Cana-de-açúcar, eds. L. L. Dinardo-
Miranda, A. C. M. Vasconcelos, and M. G. A. Landell. (Campinas: Instituto Agronômico de
Campinas), 150-165.

Gonçalves, M. C., Pinto, L. R., Souza, S. C., and Landell, M. G. A. (2012). Virus diseases of
sugarcane: a constant challenge to sugarcane breeding in Brazil. Funct. Plant Sci. Biotechnol.
6, 108-116.

Gonçalves, M. C., Vega, J., Oliveira, J. G., and Gomes, M. M. A. (2005). Sugarcane yellow leaf
virus infection leads to alterations in photosynthetic efficiency and carbohydrate
accumulation in sugarcane leaves. Fitopatol. Bras. 30, 10-16. doi: 10.1590/s0100-
41582005000100002

Gouy, M., Rousselle, Y., Chane, A. T., Anglade, A., Royaert, S., Nibouche, S., et al. (2015). Genome
wide association mapping of agro-morphological and disease resistance traits in sugarcane.
Euphytica 202, 269-284. doi: 10.1007/s10681-014-1294-y

Grativol, C., Regulski, M., Bertalan, M., McCombie, W. R., da Silva, F. R., Zerlotini Neto, A., et al.
(2014). Sugarcane genome sequencing by methylation filtration provides tools for genomic
research in the genus *Saccharum*. Plant J. 79, 162-172. doi: 10.1111/tpj.12539

Grisham, M., Pan, Y., Legendre, B., Godshall, M., and Eggleston, G. (2001). Effect of sugarcane
yellow leaf virus on sugarcane yield and juice quality. Proc. Int. Soc. Sugar Cane Technol.
24, 434-438

Harries, P. A., Park, J. W., Sasaki, N., Ballard, K. D., Maule, A. J., & Nelson, R. S. (2009). Differing
requirements for actin and myosin by plant viruses for sustained intercellular movement.
Proceedings of the National Academy of Sciences, 106(41), 17594-17599. doi:
10.1073/pnas.0909239106

Heer, K., Behringer, D., Piermattei, A., Bässler, C., Brandl, R., Fady, B.... & Vendramin, G. G.
(2018). Linking dendroecology and association genetics in natural populations: Stress
responses archived in tree rings associate with SNP genotypes in silver fir (*Abies alba* Mill.).
Molecular Ecology, 27(6), 1428-1438. doi: 10.1111/mec.14538

Heinlein, M. (2015). Plant virus replication and movement. Virology, 479, 657-671. doi:
10.1016/j.virol.2015.01.025

Heo, W. D., Lee, S. H., Kim, M. C., Kim, J. C., Chung, W. S., Chun, H. J., et al. (1999). Involvement
of specific calmodulin isoforms in salicylic acid-independent activation of plant disease
resistance responses. Proc. Natl. Acad. Sci. U. S. A. 96, 766-771. doi: 10.1073/pnas.96.2.766

Hill, W. G., & Weir, B. S. (1988). Variances and covariances of squared linkage disequilibria in
finite populations. Theoretical Population Biology, 33(1), 54-78. doi: 10.1016/0040-
5809(88)90004-4

Hoang, N. V., Furtado, A., Mason, P. J., Marquardt, A., Kasirajan, L., Thirugnanasambandam, P. P.,
et al. (2017). A survey of the complex transcriptome from the highly polyploid sugarcane
genome using full-length isoform sequencing and de novo assembly from short read
sequencing. BMC Genom. 18, 395-395. doi: 10.1186/s12864-017-3757-8

Hofius, D., Maier, A. T., Dietrich, C., Jungkunz, I., Börnke, F., Maiss, E., & Sonnewald, U. (2007).
Capsid protein-mediated recruitment of host DnaJ-like proteins is required for Potato virus Y
infection in tobacco plants. Journal of Virology, 81(21), 11870-11880. doi:
10.1128/JVI.01525-07

Huang, T. S., Wei, T., Laliberté, J. F., and Wang, A. (2010). A host RNA helicase-like protein,
AtRH8, interacts with the potyviral genome-linked protein, VPg, associates with the virus
accumulation complex, and is essential for infection. Plant Physiology, 152(1), 255-266. doi:
10.1104/pp.109.147983

Ibrahim, A., Yang, X., Liu, C., Cooper, K. D., Bishop, B. A., Zhu, M.... & Nelson, R. S. (2020).
Plant SNAREs SYP22 and SYP23 interact with Tobacco mosaic virus 126 kDa protein and
SYP2s are required for normal local virus accumulation and spread. Virology, 547, 57-71.
doi: 10.1016/j.virol.2020.04.002

Islam, M. S., Yang, X., Sood, S., Comstock, J. C., and Wang, J. (2018). Molecular characterization
of genetic basis of sugarcane yellow leaf virus (SCYLV) resistance in *Saccharum* spp. hybrid.
Plant Breed. 137, 598-604. doi: 10.1111/pbr.12614

ISO (2020). International Sugar Organization. London: ISO.

Jannoo, N., Grivet, L., Dookun, A., D'Hont, A., & Glaszmann, J. C. (1999). Linkage disequilibrium
among modern sugarcane cultivars. Theoretical and Applied Genetics, 99(6), 1053-1060. doi:
10.1007/s001220051414

Jarošová, J., Gadiou, S., and Kumar, J. K. (2009). Real-time RT-PCR quantitative analysis of plant
viruses in stone fruit tissues. Julius-Kühn-Archiv, 61

Johnson, R. C., Nelson, G. W., Troyer, J. L., Lautenberger, J. A., Kessing, B. D., Winkler, C. A., et
al. (2010). Accounting for multiple comparisons in a genome-wide association study
(GWAS). BMC Genom. 11, 724. doi: 10.1186/1471-2164-11-724

Jombart, T. (2008). adegenet: a R package for the multivariate analysis of genetic markers.
Bioinformatics 24, 1403-1405. doi: 10.1093/bioinformatics/btn129

Kassambara, A., and Mundt, F. (2017). Factoextra: extract and visualize the results of multivariate
data analyses. R Package Version 1, 2017

Knaus, B. J., and Grünwald, N. J. (2017). vcfr: a package to manipulate and visualize variant call
format data in R. Mol. Ecol. Resour. 17, 44-53. doi: 10.1111/1755-0998.12549

Kopelman, N. M., Mayzel, J., Jakobsson, M., Rosenberg, N. A., and Mayrose, I. (2015). Clumpak: a
program for identifying clustering modes and packaging population structure inferences
across K. Mol. Ecol. Resour. 15, 1179-1191. doi: 10.1111/1755-0998.12387

Korimbocus, J., Coates, D., Barker, I., and Boonham, N. (2002). Improved detection of sugarcane
yellow leaf virus using a real-time fluorescent (TaqMan) RT-PCR assay. J. Virol. Methods
103, 109-120. doi: 10.1016/s0166-0934(01)00406-2

Kovalev, N., Pogany, J., and Nagy, P. D. (2012). A co-opted DEAD-box RNA helicase enhances
tombusvirus plus-strand synthesis. PLoS Pathogens, 8(2), e1002537. doi:
10.1371/journal.ppat.1002537

Lalman, J. A., Shewa, W. A., Gallagher, J., and Ravella, S. (2016). Biofuels production from
renewable feedstocks, in Quality Living Through Chemurgy and Green Chemistry, ed. P. C.
K. Lau. (Berlin, Heidelberg: Springer), 193-220. doi: 10.1007/978-3-662-53704-6_8

Langmead, B., and Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. Nat. Methods
9, 357-359. doi: 10.1038/nmeth.1923

Lê, S., Josse, J., and Husson, F. (2008). FactoMineR: an R package for multivariate analysis. J. Stat.
Softw. 25, 1-18. doi: 10.18637/jss.v025.i01

Lehrer, A. T., Moore, P. H., and Komor, E. (2007). Impact of sugarcane yellow leaf virus (SCYLV)
on the carbohydrate status of sugarcane: comparison of virus-free plants with symptomatic
and asymptomatic virus-infected plants. Physiological and Molecular Plant Pathology, 70(4-
6), 180-188. doi: 10.1016/j.pmpp.2007.09.005

Lehrer, A., Yan, S.-L., Fontaniella, B., ElSayed, A., and Komor, E. (2010). Carbohydrate
composition of sugarcane cultivars that are resistant or susceptible to sugarcane yellow leaf
virus. J. Gen. Plant Pathol. 76, 62-68. doi: 10.1007/s10327-009-0210-0

Lehrer, A. T., and Komor, E. (2008). Symptom expression of yellow leaf disease in sugarcane
cultivars with different degrees of infection by sugarcane yellow leaf virus. Plant Pathol. 57,
178-189. doi: 10.1111/j.1365-3059.2007.01696.x

Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics (Oxford, England) 25, 1754-1760. doi:
10.1093/bioinformatics/btp324

Li, F., Huang, C., Li, Z., and Zhou, X. (2014). Suppression of RNA silencing by a plant DNA virus
satellite requires a host calmodulin-like protein to repress RDR6 expression. PLoS Pathog.
10, e1003921. doi: 10.1371/journal.ppat.1003921

Li, Y., Xiong, R., Bernards, M., and Wang, A. (2016). Recruitment of *Arabidopsis* RNA helicase
AtRH9 to the viral replication complex by viral replicase to promote turnip mosaic virus
replication. Scientific Reports, 6, 30297. doi: 10.1038/srep30297

Lin, W. C., Lu, C. F., Wu, J. W., Cheng, M. L., Lin, Y. M., Yang, N. S.... & Cheng, C. P. (2004).
Transgenic tomato plants expressing the Arabidopsis NPR1 gene display enhanced resistance
to a spectrum of fungal and bacterial diseases. Transgenic Research, 13(6), 567-581. doi:
10.1007/s11248-004-2375-9

Liu, Y., Schiff, M., Marathe, R., & Dinesh-Kumar, S. P. (2002). Tobacco Rar1, EDS1 and
NPR1/NIM1 like genes are required for N-mediated resistance to tobacco mosaic virus. The
Plant Journal, 30(4), 415-429. doi: 10.1046/j.1365-313x.2002.01297.x

Liu, J. Z., & Whitham, S. A. (2013). Overexpression of a soybean nuclear localized type–III DnaJ
domain-containing HSP40 reveals its roles in cell death and disease resistance. The Plant
Journal, 74(1), 110-121. doi: 10.1111/tpj.12108

Liu, X., Huang, M., Fan, B., Buckler, E. S., & Zhang, Z. (2016). Iterative usage of fixed and random
effect models for powerful and efficient genome-wide association studies. PLoS Genetics,
12(2), e1005767. doi: 10.1371/journal.pgen.1005767

Livak, K. J., and Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time
quantitative PCR and the $2^{-\Delta\Delta CT}$ method. Methods 25, 402-408. doi: 10.1006/meth.2001.1262

Llave, C. (2010). Virus-derived small interfering RNAs at the core of plant–virus interactions.
Trends in Plant Science, 15(12), 701-707. doi: 10.1016/j.tplants.2010.09.001

Lu, L., Du, Z., Qin, M., Wang, P., Lan, H., Niu, X.... & Wu, Z. (2009). Pc4, a putative movement
protein of Rice stripe virus, interacts with a type I DnaJ protein and a small Hsp of rice. Virus
Genes, 38(2), 320-327. doi: 10.1007/s11262-008-0324-z

Lundin, S., Stranneheim, H., Pettersson, E., Klevebring, D., and Lundeberg, J. (2010). Increased
throughput by parallelization of library preparation for massive sequencing. PLoS One 5,
e10029. doi: 10.1371/journal.pone.0010029

Maccheroni, W., Jordão, H., Degaspari, R., & Matsuoka, S. (2007). Development of a dependable
microsatellite-based fingerprinting system for sugarcane. Proceedings of the International
Society of Sugar Cane Technologistics, Durban.

Mancini, M. C., Cardoso-Silva, C. B., Sforça, D. A., and Pereira de Souza, A. (2018). "Targeted
sequencing by gene synteny," a new strategy for polyploid species: sequencing and physical
structure of a complex sugarcane region. Front. Plant Sci. 9, 397. doi:
10.3389/fpls.2018.00397

Mangwende, T., Wang, M.-L., Borth, W., Hu, J., Moore, P. H., Mirkov, T. E., et al. (2009). The P0
gene of sugarcane yellow leaf virus encodes an RNA silencing suppressor with unique
activities. Virology 384, 38-50. doi: 10.1016/j.virol.2008.10.034

Mansilla-Córdova, P. J., Bampi, D., Rondinel-Mendoza, N. V., Melo, P. C. T., Lourenção, A. L., and
Rezende, J. A. M. (2018). Screening tomato genotypes for resistance and tolerance to Tomato
chlorosis virus. Plant Pathol. 67, 1231-1237. doi: 10.1111/ppa.12826

Marconi, T. G., Costa, E. A., Miranda, H. R., Mancini, M. C., Cardoso-Silva, C. B., Oliveira, K. M....
& Souza, A. P. (2011). Functional markers for gene mapping and genetic diversity studies in
sugarcane. BMC Research Notes, 4(1), 264. doi: 10.1186/1756-0500-4-264

Matias, F. I., Alves, F. C., Meireles, K. G. X., Barrios, S. C. L., do Valle, C. B., Endelman, J. B., &
Fritsche-Neto, R. (2019). On the accuracy of genomic prediction models considering multi-
trait and allele dosage in *Urochloa* spp. interspecific tetraploid hybrids. Molecular Breeding,
39(7), 100. doi: 10.1007/s11032-019-1002-7

Melo, A. T. O., Bartaula, R., and Hale, I. (2016). GBS-SNP-CROP: a reference-optional pipeline for
SNP discovery and plant germplasm characterization using variable length, paired-end
genotyping-by-sequencing data. BMC Bioinform. 17, 29. doi: 10.1186/s12859-016-0879-y

Moonan, F., Molina, J., and Mirkov, T. E. (2000). Sugarcane yellow leaf virus: an emerging virus
that has evolved by recombination between luteoviral and poleroviral ancestors. Virology
269, 156-171. doi: 10.1006/viro.1999.0162

Morel, J.-B., Godon, C., Mourrain, P., Béclin, C., Boutet, S., Feuerbach, F., et al. (2002). Fertile
hypomorphic ARGONAUTE (ago1) mutants impaired in post-transcriptional gene silencing
and virus resistance. Plant Cell 14, 629-639. doi: 10.1105/tpc.010358

Muñoz, F., and Sanchez, L.S. (2014). breedR: statistical methods for forest genetic resources
analysis. In Trees for the future: plant material in a changing climate.

Nakahara, K. S., Masuta, C., Yamada, S., Shimura, H., Kashihara, Y., Wada, T. S., et al. (2012).
Tobacco calmodulin-like protein provides secondary defense by binding to and directing
degradation of virus RNA silencing suppressors. Proc. Natl. Acad. Sci. U. S. A. 109, 10113-
10118. doi: 10.1073/pnas.1201628109

Nimmakayala, P., Tomason, Y. R., Abburi, V. L., Alvarado, A., Saminathan, T., Vajja, V. G., et al.
(2016). Genome-wide differentiation of various melon horticultural groups for use in GWAS
for fruit firmness and construction of a high resolution genetic map. Front. Plant Sci. 7, 1437.
doi: 10.3389/fpls.2016.01437

Oliveira, K. M., Pinto, L. R., Marconi, T. G., Margarido, G. R., Pastina, M. M., Teixeira, L. H. M....
& Souza, A. P. (2007). Functional integrated genetic linkage map based on EST-markers for a
sugarcane (*Saccharum* spp.) commercial cross. Molecular Breeding, 20(3), 189-208. doi:
10.1007/s11032-007-9082-1

Oliveira, K. M., Pinto, L. R., Marconi, T. G., Mollinari, M., Ulian, E. C., Chabregas, S. M.... &
Souza, A. P. (2009). Characterization of new polymorphic functional markers for sugarcane.
Genome, 52(2), 191-209. doi: 10.1139/g08-105

Panje, R. R., and Babu, C. N. (1960). Studies in Saccharum spontaneum distribution and
geographical association of chromosome numbers. Cytologia 25, 152-172. doi:
10.1508/cytologia.25.152

Paterson, A. H., Bowers, J. E., Bruggmann, R., Dubchak, I., Grimwood, J., Gundlach, H., et al.
(2009). The *Sorghum bicolor* genome and the diversification of grasses. Nature 457, 551-556.
doi: 10.1038/nature07723

Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O.... & Vanderplas, J.
(2011). Scikit-learn: Machine learning in Python. The Journal of Machine Learning Research,
12, 2825-2830.

Pereira, G. S., Garcia, A. A. F., and Margarido, G. R. A. (2018). A fully automated pipeline for
quantitative genotype calling from next generation sequencing data in autopolyploids. BMC
Bioinform. 19, 398. doi: 10.1186/s12859-018-2433-6

Peterson, R. (2017). Estimating normalization transformations with bestNormalize.
https://github.com/petersonR/bestNormalize. [Accessed September 2, 2020].

Pieterse, C. M., Van Wees, S. C., Van Pelt, J. A., Knoester, M., Laan, R., Gerrits, H.... & Van Loon,
L. C. (1998). A novel signaling pathway controlling induced systemic resistance in
Arabidopsis. The Plant Cell, 10(9), 1571-1580. doi: 10.1105/tpc.10.9.1571

Poland, J. A., Brown, P. J., Sorrells, M. E., and Jannink, J.-L. (2012). Development of high-density
genetic maps for barley and wheat using a novel two-enzyme genotyping-by-sequencing
approach. PLoS One 7, e32253. doi: 10.1371/journal.pone.0032253

Popescu, M. C., Balas, V. E., Perescu-Popescu, L., & Mastorakis, N. (2009). Multilayer perceptron
and neural networks. WSEAS Transactions on Circuits and Systems, 8(7), 579-588.

Premachandran, M. N., Prathima, P. T., and Maya, L. (2011). Sugarcane and polyploidy - a review. J.
Sugarcane Res. 1, 1-15

Quinlan, J. R. (1986). Induction of decision trees. Machine Learning, 1(1), 81-106. doi:
10.1007/BF00116251

Pritchard, J. K., Stephens, M., and Donnelly, P. (2000). Inference of population structure using
multilocus genotype data. Genetics 155, 945-959

R Core Team (2011). R: A Language and Environment for Statistical Computing. Vienna, Austria: R
Foundation for Statistical Computing.

Raboin, L. M., Oliveira, K. M., Lecunff, L., Telismart, H., Roques, D., Butterfield, M.... & D'Hont,
A. (2006). Genetic mapping in sugarcane, a high polyploid, using bi-parental progeny:
identification of a gene controlling stalk colour and a new rust resistance gene. Theoretical
and Applied Genetics, 112(7), 1382-1391. doi: 10.1007/s00122-006-0240-3

Raboin, L. M., Pauquet, J., Butterfield, M., D'Hont, A., & Glaszmann, J. C. (2008). Analysis of
genome-wide linkage disequilibrium in the highly polyploid sugarcane. Theoretical and
Applied Genetics, 116(5), 701-714. doi: 10.1371/journal.pbio.0030196

Racedo, J., Gutiérrez, L., Perera, M. F., Ostengo, S., Pardo, E. M., Cuenya, M. I., ... and Castagnaro,
A. P. (2016). Genome-wide association mapping of quantitative traits in a breeding
population of sugarcane. BMC Plant Biology, 16(1), 142. doi: 10.1186/s12870-016-0829-x

Rasmussen, C. E. (2003). Gaussian processes in machine learning. In Summer School on Machine
Learning (pp. 63-71). Springer, Berlin, Heidelberg. doi: 10.1007/978-3-540-28650-9_4

Riaño-Pachón, D. M., and Mattiello, L. (2017). Draft genome sequencing of the sugarcane hybrid
SP80-3280. F1000Research 6, 861. doi: 10.12688/f1000research.11859.2

Roach, B. T. (1972). Nobilisation of sugarcane. Proc. Int. Soc. Sugar Cane Technol. 14, 206-216

Rochon, A., Boyle, P., Wignes, T., Fobert, P. R., & Després, C. (2006). The coactivator function of
Arabidopsis NPR1 requires the core of its BTB/POZ domain and the oxidation of C-terminal
cysteines. The Plant Cell, 18(12), 3670-3685. doi: 10.1105/tpc.106.046953

Roncallo, P. F., Beaufort, V., Larsen, A. O., Dreisigacker, S., & Echenique, V. (2019). Genetic diversity and linkage disequilibrium using SNP (KASP) and AFLP markers in a worldwide durum wheat (*Triticum turgidum* L. var *durum*) collection. PloS One, 14(6), e0218562. doi: 10.1371/journal.pone.0218562

Rosyara, U. R., De Jong, W. S., Douches, D. S., and Endelman, J. B. (2016). Software for genome-wide association studies in autopolyploids and its application to potato. Plant Genome 9, 1-10. doi: 10.3835/plantgenome2015.08.0073

Sasvari, Z., Kovalev, N., Gonzalez, P. A., Xu, K., & Nagy, P. D. (2018). Assembly-hub function of ER-localized SNARE proteins in biogenesis of tombusvirus replication compartment. PLoS Pathogens, 14(5), e1007028. doi: 10.1371/journal.ppat.1007028

Scagliusi, S. M., and Lockhart, B. E. L. (2000). Transmission, characterization, and serology of a luteovirus associated with yellow leaf syndrome of sugarcane. Phytopathology 90, 120-124. doi: 10.1094/phyto.2000.90.2.120

Scagliusi, S. M., Basu, S. K., de Gouvea, J. A., & Vega, J. (2009). Physiological Alterations in Brazilian Sugarcane Varieties Infected by Sugarcane yellow leaf virus (ScYLV). Functional Plant Science and Biotechnology, 3(1), 19-25.

Schloerke, B., Crowley, J., Cook, D., Hofmann, H., Wickham, H., Briatte, F., et al. (2011). Ggally: Extension to ggplot2.

Seo, Y. S., Rojas, M. R., Lee, J. Y., Lee, S. W., Jeon, J. S., Ronald, P.... & Gilbertson, R. L. (2006). A viral resistance gene from common bean functions across plant families and is up-regulated in a non-virus-specific manner. Proceedings of the National Academy of Sciences, 103(32), 11856-11861. doi: 10.1073/pnas.0604815103

Sforça, D. A., Vautrin, S., Cardoso-Silva, C. B., Mancini, M. C., Romero-da Cruz, M. V., Pereira, G. d. S., et al. (2019). Gene duplication in the sugarcane genome: a case study of allele interactions and evolutionary patterns in two genic regions. Front. Plant Sci. 10, 553. doi: 10.3389/fpls.2019.00553

Shimizu, T., Yoshii, A., Sakurai, K., Hamada, K., Yamaji, Y., Suzuki, M.... & Hibi, T. (2009). Identification of a novel tobacco DnaJ-like protein that interacts with the movement protein of tobacco mosaic virus. Archives of Virology, 154(6), 959-967. doi: 10.1007/s00705-009-0397-6

Singh, R. K., Banerjee, N., Khan, M. S., Yadav, S., Kumar, S., Duttamajumder, S. K.... and Paterson, A. H. (2016). Identification of putative candidate genes for red rot resistance in sugarcane (*Saccharum* species hybrid) using LD-based association mapping. Molecular Genetics and Genomics, 291(3), 1363-1377. doi: 10.1007/s00438-016-1190-3

Slater, A. T., Cogan, N. O. I., Forster, J. W., Hayes, B. J., and Daetwyler, H. D. (2016). Improving genetic gain with genomic selection in autotetraploid potato. Plant Genome 9, 1-15. doi: 10.3835/plantgenome2016.02.0021

Smith, G. R., Borg, Z., Braithwaite, K. S., Lockhart, B. E. L., and Gibbs, M. J. (2000). Sugarcane
yellow leaf virus: a novel member of the Luteoviridae that probably arose by inter-species
recombination. J. Gen. Virol. 81, 1865-1869. doi: 10.1099/0022-1317-81-7-1865

Spoel, S. H., Koornneef, A., Claessens, S. M., Korzelius, J. P., Van Pelt, J. A., Mueller, M. J.... &
Van Loon, L. C. (2003). NPR1 modulates cross-talk between salicylate-and jasmonate-
dependent defense pathways through a novel function in the cytosol. The Plant Cell, 15(3),
760-770. doi: 10.1105/tpc.009159

Stacklies, W., Redestig, H., Scholz, M., Walther, D., & Selbig, J. (2007). pcaMethods – a
bioconductor package providing PCA methods for incomplete data. Bioinformatics, 23(9),
1164-1167. doi: 10.1093/bioinformatics/btm069

Steinfath, M., Gärtner, T., Lisec, J., Meyer, R. C., Altmann, T., Willmitzer, L., & Selbig, J. (2010).
Prediction of hybrid biomass in *Arabidopsis thaliana* by selected parental SNP and metabolic
markers. Theoretical and Applied Genetics, 120(2), 239-247. doi: 10.1007/s00122-009-1191-
2

Su, J., Fan, S., Li, L., Wei, H., Wang, C., Wang, H., et al. (2016). Detection of favorable QTL alleles
and candidate genes for lint percentage by GWAS in Chinese Upland cotton. Front. Plant Sci.
7, 1576. doi: 10.3389/fpls.2016.01576

Thiel, H., Hleibieh, K., Gilmer, D., and Varrelmann, M. (2012). The P25 pathogenicity factor of beet
necrotic yellow vein virus targets the sugar beet 26S proteasome involved in the induction of
a hypersensitive resistance response via interaction with an F-box protein. Mol. Plant-
Microbe Interact. 25, 1058-1072. doi: 10.1094/mpmi-03-12-0057-r

VanRaden, P. M. (2008). Efficient methods to compute genomic predictions. J. Dairy Sci. 91, 4414-
4423. doi: 10.3168/jds.2007-0980

Vasconcelos, A., Gonçalves, M. C., Pinto, L. R., Landell, M. G., and Perecin, D. (2009). Effects of
sugarcane yellow leaf virus on sugarcane yield and root system development. Funct. Plant
Sci. Biotechnol. 3, 31-35

Vega, J., Scagliusi, S. M. M., and Ulian, E. C. (1997). Sugarcane yellow leaf disease in Brazil:
evidence of association with a luteovirus. Plant Dis. 81, 21-26. doi:
10.1094/pdis.1997.81.1.21

Villavicencio, R. C. B., Gonçalves, M. C., Nóbile, P. M., Anjos, I. A., Silva, M. F., Perecin, D.,
Anjos, L. S., Fernandes, V. B., Pinto, L. R. R. (in press) Screening of *Saccharum* spp.
genotypes for sugarcane yellow leaf virus resistance by combining symptom phenotyping and
highly precise virus titration. Crop Protection.

Viswanathan, R., Chinnaraja, C., Malathi, P., Gomathi, R., Rakkiyappan, P., Neelamathi, D., et al.
(2014). Impact of sugarcane yellow leaf virus (SCYLV) infection on physiological efficiency
and growth parameters of sugarcane under tropical climatic conditions in India. Acta Physiol.
Plant. 36, 1805-1822. doi: 10.1007/s11738-014-1554-4

Voorrips, R. E. (2002). MapChart: software for the graphical presentation of linkage maps and QTLs.
Journal of Heredity, 93(1), 77-78. doi: 10.1093/jhered/93.1.77

Vos, P. G., Paulo, M. J., Voorrips, R. E., Visser, R. G., van Eck, H. J., & van Eeuwijk, F. A. (2017).
Evaluation of LD decay and various LD-decay estimators in simulated and SNP-array data of
tetraploid potato. Theor. App. Genet. 130, 123-135. doi: 10.1007/s00122-016-2798-8

Wang, H., Smith, K. P., Combs, E., Blake, T., Horsley, R. D., and Muehlbauer, G. J. (2012). Effect
of population size and unbalanced data sets on QTL detection using genome-wide association
mapping in barley breeding germplasm. Theor. Appl. Genet. 124, 111-124. doi:
10.1007/s00122-011-1691-8

Wei, X., Jackson, P. A., Hermann, S., Kilian, A., Heller-Uszynska, K., and Deomano, E. (2010).
Simultaneously accounting for population structure, genotype by environment interaction,
and spatial variation in marker–trait associations in sugarcane. Genome, 53(11), 973-981. doi:
10.1139/G10-050

Wei, T., Zhang, C., Hou, X., Sanfaçon, H., & Wang, A. (2013). The SNARE protein Syp71 is
essential for turnip mosaic virus infection by mediating fusion of virus-induced vesicles with
chloroplasts. PLoS Pathogens, 9(5), e1003378. doi: 10.1371/journal.ppat.1003378

Xun, H., Yang, X., He, H., Wang, M., Guo, P., Wang, Y.... & Liu, B. (2019). Over-expression of
GmKR3, a TIR–NBS–LRR type R gene, confers resistance to multiple viruses in soybean.
Plant Molecular Biology, 99(1-2), 95-111. doi: 10.1007/s11103-018-0804-z

Yang, N., Lu, Y., Yang, X., Huang, J., Zhou, Y., Ali, F., et al. (2014). Genome wide association
studies using a new nonparametric model reveal the genetic architecture of 17 agronomic
traits in an enlarged maize association panel. PLoS Genet. 10, e1004573-e1004573. doi:
10.1371/journal.pgen.1004573

Yang, Z., Li, Y. (2018). Dissection of RNAi-based antiviral immunity in plants. Current Opinion in
Virology, 32, 88-99. doi: 10.1016/j.coviro.2018.08.003

Yang, X., Sood, S., Luo, Z., Todd, J., and Wang, J. (2019a). Genome-wide association studies
identified resistance loci to orange rust and yellow leaf virus diseases in sugarcane
(*Saccharum* spp.). Phytopathology 109, 623-631. doi: 10.1094/phyto-08-18-0282-r

Yang, X., Todd, J., Arundale, R., Binder, J. B., Luo, Z., Islam, M. S., et al. (2019b). Identifying loci
controlling fiber composition in polyploid sugarcane (*Saccharum* spp.) through genome-wide
association study. Ind. Crops Prod. 130, 598-605. doi: 10.1016/j.indcrop.2019.01.023

Yang, X., Song, J., Todd, J., Peng, Z., Paudel, D., Luo, Z.... and Zhao, Y. (2019c). Target enrichment
sequencing of 307 germplasm accessions identified ancestry of ancient and modern hybrids
and signatures of adaptation and selection in sugarcane (*Saccharum* spp.), a 'sweet'crop with
'bitter'genomes. Plant Biotechnology Journal, 17(2), 488-498. doi: 10.1111/pbi.12992

Yang, X., Luo, Z., Todd, J., Sood, S., & Wang, J. (2020). Genome-wide association study of multiple
yield traits in a diversity panel of polyploid sugarcane (*Saccharum* spp.). The Plant Genome,
e20006. doi: 10.1002/tpg2.20006

Zhang, J., Zhang, X., Tang, H., Zhang, Q., Hua, X., Ma, X., et al. (2018). Allele-defined genome of
the autopolyploid sugarcane *Saccharum spontaneum* L. Nat. Genet. 50, 1565-1573. doi:
10.1038/s41588-018-0237-2

Zhou, W., & Bellis, E. S. (2019). Minor QTLs mining through the combination of GWAS and
machine learning feature selection. BioXriv. doi: 10.1101/702761

Zhu, Y. J., Lim, S. T. S., Schenck, S., Arcinas, A., and Komor, E. (2010). RT-PCR and quantitative
real-time RT-PCR detection of sugarcane yellow leaf virus (SCYLV) in symptomatic and
asymptomatic plants of Hawaiian sugarcane cultivars and the correlation of SCYLV titre to
yield. Eur. J. Plant Pathol. 127, 263-273. doi: 10.1007/s10658-010-9591-3

1478

1479

1480

1481

1482

1483

1484

1485

1486

1487

1488

1489

1490

1491

1492

1493

1494

1495

1496

1497

1498

1499

Figure Legends

Figure 1. Distributions of BLUP values and correlations between the three traits analyzed.
Asterisks (***) indicate statistical significance ($p < 0.0005$).

Figure 2. Factorial maps generated in the two hierarchical clustering on principal components
(HPCP) analyses using BLUP values. (A) Factorial map of HCPCs performed using the SCYLV
titer determined by RT-qPCR. A division into three clusters (Q1, Q2 and Q3) was considered. **(B)**
Factorial map of HCPC performed using SCYL symptom severity and SCYLV titer determined by
RT-qPCR. A division into three clusters (SQ1, SQ2 and SQ3) was considered.

Figure 3: Graphical outputs of the three clustering analyses employed to evaluate the genetic
structure for 93 genotypes of the panel using 662 amplified fragment length polymorphisms
(AFLPs) and simple sequence repeats (SSRs). (A) Plotting of the panel onto the first two linear
discriminants from a discriminant analysis of principal components (DAPC). A division into three
clusters (A1, A2 and A3) was considered. **(B)** Projection of the panel onto the first two principal
components (PCs) from a principal component analysis (PCA). Genotypes are divided into four
clusters (B1, B2, B3 and B4) as indicated by a k-means clustering analysis. Ellipses represent a 95%
confidence interval. **(C)** Bayesian clustering performed on STRUCTURE. Each genotype is
represented by a vertical bar split in accordance with the admixture coefficients for two clusters (C1
in dark blue and C2 in red). Genotypes are ordered according to ID number (see Supplementary
Table 1).

Figure 4. Graphical outputs of the three clustering analyses employed to evaluate the genetic
structures for 92 genotypes in the panel using codominant markers. (A) Plotting of the panel
onto the first two linear discriminants from a discriminant analysis of principal components (DAPC)
using 70,888 single-nucleotide polymorphisms (SNPs) and insertions and deletions (indels). A
division into three clusters (D1, D2 and D3) was considered. **(B)** Projection of the panel onto the first
two principal components (PCs) from a principal component analysis (PCA) using 70,888 SNPs and
indels. The genotypes were divided into two clusters (E1 and E2) as indicated by k-means clustering
analysis. Ellipses represent a 95% confidence interval. **(C)** Bayesian clustering performed on
STRUCTURE using 7,000 SNP and indel markers. Each genotype is represented by a vertical bar
that is split in accordance with the admixture coefficients for two clusters (F1 in blue and F2 in
orange). The genotypes are ordered according to ID number (see Supplementary Table 1).

Figure 5. Decay of linkage disequilibrium (r^2) as a function of physical distance (bp) between
pairs of 67,007 single nucleotide polymorphisms (SNPs) and insertions and deletions (indels)
located on *Saccharum spontaneum* chromosomes 1A-8A. Only r^2 values with $P < 0.05$ are
included.

Figure 6. Manhattan plots generated in association analyses using the best linear unbiased
predictor (BLUP) values of the three traits analyzed. Six different models were tested: general,
additive, simplex dominant reference (1-dom-ref), simplex dominant alternative (1-dom-alt),
diploidized general (diplo-general) and diploidized additive (diplo-additive). On the x-axis, S
represents scaffolds not associated with any of the *S. spontaneum* chromosomes.

Figure 7. Distribution of markers associated with SCYLV resistance along *Saccharum*
***spontaneum* "A" chromosomes.** In each chromosome, marker positions are shown on the left, and
marker names are indicated on the right, labeled and colored according to the method employed for
their identification. Markers identified by previous mapping studies are colored in gray.

Figures

1544

1545 **Figure 1**

1546

1547

1548 **Figure 2**

1549

1550

1551   **Figure 3**

1552

1553

1554

1555

1556

1557

1558

1559

1560

1561

1562   **Figure 4**

1563

1564

1565

1566

1567

1568

1569

1570

1571

1572

1573

1574

1575

1576

1577

1578 **Figure 5**

1579

1580

1581

1582

1583 **Figure 6**

1584

1585

1586

1587

1588

1589   **Figure 7**