Shifts in mutation spectra enhance access to beneficial mutations

Mrudula Sane[1], Gaurav D Diwan[1,2,3], Bhoomika A Bhat[1,4], Lindi M Wahl[5]* and Deepa Agashe[1]*

[1] National Centre for Biological Sciences, GKVK Campus, Bellary Road, Bengaluru, India

[2] CellNetworks, Bioquant, University of Heidelberg, 69120 Heidelberg, Germany

[3] Biochemie Zentrum Heidelberg (BZH), University of Heidelberg, 69120 Heidelberg, Germany

[4] Undergraduate Programme, Indian Institute of Science, Bengaluru, India

[5] Western University, London, Ontario, Canada

*Correspondence to:

dagashe@ncbs.res.in

lwahl@uwo.ca

ABSTRACT

Biased mutation spectra are pervasive, with widely varying direction and magnitude of mutational

biases that influence genome evolution and adaptation. Why are unbiased spectra rare, and how

do such diverse biases evolve? Our experiments show that changing the mutation spectrum allows

populations to sample previously under-sampled mutational space, including beneficial mutations.

The resulting shift in the distribution of fitness effects is advantageous: beneficial mutation supply

and beneficial pleiotropy both increase, and deleterious load reduces. More broadly, simulations

indicate that the evolution of a mutational bias in an unbiased ancestor is selectively neutral; but

reversing the direction of a long-term bias is always selectively favoured. Indeed, we predict

frequent spectrum changes in the bacterial phylogeny, typically involving reversals of ancestral

bias. Thus, shifts in mutation spectra may evolve under selection, and can directly alter the

outcome of adaptive evolution by facilitating access to beneficial mutations.

SIGNIFICANCE STATEMENT

Mutations are important because they provide raw material for evolution. Some types of mutations

occur more often than others, but the direction and degree of such mutational biases varies. How

do mutational biases evolve, and do they influence evolution? We experimentally measured the

immediate effects of changing the mutation bias of *E. coli*, and used simulations to test the long-

term effects. Altering mutational bias is beneficial whenever the new bias increases sampling of

mutational classes that were previously under-sampled. We also show that historically, bacteria

have often experienced such beneficial bias switches. Our work demonstrates the importance of

mutational biases in evolution. By allowing exploration of new mutational space, altered mutation

biases could drive rapid adaptation even in constant environments.

INTRODUCTION

The mutation spectrum describes the frequency of various classes of sampled mutations (e.g., transversions vs. transitions (Tv/Ts)), and is often determined by the action of DNA repair enzymes. By determining the pool of genetic variants available for selection, mutation spectra can shape key genome features (e.g., nucleotide (1, 2), codon and amino acid composition (3)), even under selection for thousands of generations (4); determine the genetic basis of adaptation (5–9), driving convergent evolution (10–12); and shape the evolution of resistance to antibiotics (13–15) and anti-cancer drugs (16). Thus, mutation spectra are important for adaptive evolution, but their role is underappreciated (5). The mutation spectrum is instantaneously and relatively easily altered by loss-of-function mutations in DNA repair genes, e.g., as observed in "mutators" that also have higher mutation rates (17). The elevated mutation rate increases the supply of beneficial mutations, facilitating rapid adaptation (18). However, spectrum shifts also occur without associated mutation rate changes, such as with the loss of some bacterial DNA repair enzymes (17), and under nutritional or anaerobic stress (19, 20). Diverse mutation spectra occur even across natural yeast strains and closely related species of *Chlamydomonas*, despite similar mutation rates (21, 22).

In fact, most species analysed so far have a skewed mutation spectrum (i.e., some mutational classes are over-represented relative to the null expectation), with substantial variation in the direction and magnitude of bias (23, 24) (experimental data from microbes summarized in Table S1). This diversity implies major evolutionary shifts in mutation spectra. However, the frequency, underlying evolutionary processes, and consequences of spectrum shifts are unknown. For instance, why are unbiased spectra rare? Do spectrum shifts – for example, a change from an unbiased to a biased state, or altered direction of bias – evolve under selection? A spectrum shift may persist if the new bias favours mutational classes that are inherently advantageous (5, 25), allowing the evolved bias to hitchhike with beneficial mutations. It has also been speculated that stress-induced changes in mutation spectra may enhance sampling of new beneficial mutations by altering the distribution of fitness effects (DFE) (19, 26). However, prior studies have failed to find consistent support for general selective benefits of specific mutational classes. For instance,

transition mutations (Ts) are not consistently more beneficial (or less deleterious) than

transversions (Tv) (27, 28); and the fixation probability of Ts vs. Tv mutations in genomes is clade-

specific (29). Hence, particular mutational classes do not seem to be universally beneficial, and

the observed diversity in mutation bias remains puzzling.

We addressed these gaps using a combination of experiments, simulations, and phylogenetic

analyses. To measure the immediate evolutionary impact of the mutation spectrum, we estimated

the genome-wide DFE of new mutations in *Escherichia coli*. We manipulated the mutation

spectrum by deleting a DNA repair gene (Δ*mutY*, "mutator" strain) from our wild type strain ("WT"),

altering both the mutation spectrum and rate (17). Allowing independent lineages of both strains

to evolve under mutation accumulation (MA), we sequenced whole genomes of evolved isolates

to identify strains carrying a single new mutation each. Thus, we minimized the impact of selection

on the mutation spectrum and the DFE, while also decoupling the effects of the mutator's spectrum

from its high mutation rate. Measuring the effect of each single mutation on population growth rate,

we determined the effects of mutation spectrum on the beneficial mutation supply, genetic load,

and pleiotropic effects. Next, to test the generality and longer-term evolutionary consequences of

shifts in mutation spectra, we performed both adaptive walk and full population simulations. Finally,

we inferred evolutionary transitions in DNA repair enzymes across the bacterial phylogeny to

estimate the frequency of shifts in mutation bias. Our results show that shifts in mutation spectra

can evolve under selection and fuel adaptation via previously under-sampled evolutionary paths.

RESULTS AND DISCUSSION

The mutator DFE has more beneficial mutations that reduce the expected genetic load and

alter pleiotropic effects

From MA experiments conducted in rich media (LB), we obtained 80 evolved WT (30) and 79

mutator strains, each carrying a single distinct mutation with respect to its ancestor (Supplementary

Data). We measured the selective fitness effect (relative growth rate) of each mutation in the MA

environment (LB) as well as 15 other environments with different carbon sources, including some

that are used by *E. coli* in the mouse gut (31). With this large set of fitness effects, we constructed

environment-specific empirical DFEs (WT data for some environments were previously reported

(30)). Since bacterial MA experiments can cause over-sampling of beneficial mutations and under-

sampling of deleterious mutations, we used a correction (32) that retains the measured selective

effect (*s*) of each mutation, but estimates and corrects for selection bias by changing the frequency

of mutations with a given selective effect (Fig. S1A–C, Supplementary Data). These corrected

DFEs thus provide accurate estimates of the fraction of beneficial mutations (f_b) and the average

selective effect (*s*).

Correcting for selection bias during MA results in DFEs with coarsely binned fitness histograms,

we calculated the fraction of neutral (–0.05 < *s* < 0.05, conservatively accounting for a maximum

of ~5% fitness measurement error), beneficial (*s* > 0.05), and deleterious (*s* < –0.05) mutations for

each strain in all 16 environments. The DFEs and f_b values varied across environments (Fig. 1,

Fig. S1), but on average the mutator had a greater fraction of beneficial mutations (f_b = 36% vs.

28%) and fewer deleterious mutations (Fig. 1, Table S2). Thus, compared to WT, the mutator had

more beneficial-shifted DFEs. Mean fitness effects of mutations were also significantly more

beneficial in the mutator (paired Welch's two-sample test, p = 0.043; Fig. S2A), although medians

were not significantly different (p = 0.35, Fig. S2B). Finally, mutations had larger absolute effects

in WT than in the mutator (Wilcoxon's rank-sum test, W = 662920, p = 1.4×10^{-14}); hence, the

mutator's beneficial-shifted DFEs could not have arisen because of overall larger-effect mutations.

These results indicate a broad selective advantage to the mutator, independent of its mutation

rate. In addition, accounting for the skewed DFEs and order-of-magnitude higher mutation rate

(μ_{WT} = 9.3×10^{-11} per bp per generation, $\mu_{mutator}$ = 89×10^{-11}, Table S3), we estimated (following (33))

that on average the mutator should have an ~16-fold greater genome-wide supply of beneficial

mutations, and ~6-fold higher genetic load than the WT (Tables S4, S5). In contrast, if we ignored

the mutator's distinct DFE and accounted only for its mutation rate, both the supply of beneficial

mutations and the deleterious genetic load would be nearly 10-fold greater than WT (Tables S4,

S5; only the load is significantly affected by the correct DFE). Thus, depending on their mutation

rate and DFE, mutators may have a substantially higher supply of beneficial mutations and smaller

genetic load than previously believed.

The high f_b values observed in some environments (Fig. 1) contradict the general expectation that

beneficial mutations should be rare (34). However, many other studies – including with microbes

and plants – have also reported a large number of beneficial mutations (35–41). Such large f_b

values have been partially attributed to an improved ability to measure small-effect beneficial

mutations (38, 42, 43) that are more common than expected (44), especially at low effective

population sizes (40). Most prior studies (34) provide approximate or indirect estimates of the DFE

via random mutagenesis of a single gene, and focus only on the beneficial part of the DFE, or infer

the DFE by fitting a model to time series data from selection or MA lines. Further, they do not

estimate the number of underlying mutations to account for epistasis. Our direct, genome-wide

estimates of the DFE of single, randomly-sampled new mutations support the idea (45, 46) that

new beneficial mutations may be more common than generally believed.

Next, we estimated the pleiotropic effects of mutations (Fig. 2A), which are important because they

can facilitate adaptation (via synergistic pleiotropic benefits) and shape fitness tradeoffs (via

antagonistic pleiotropy) in new environments. The mutator had a distinct distribution of pleiotropic

effects in all 16 environments (compare Fig. S2C–D, chi-square tests: $p < 10^{-5}$ in each case, Table

S6), driven by a higher incidence of beneficial synergistic pleiotropy and lower deleterious

synergistic pleiotropy. However, both strains had low antagonistic pleiotropy (Fig. 2B–C).

Beneficial mutations in the mutator were also beneficial across many more environments

(Kolmogorov-Smirnov test comparing the two distributions in Fig. 2F, D=0.41, $p=3.1 \times 10^{-6}$);

whereas deleterious mutations were deleterious in fewer environments than WT (D=0.56,

$p=2.63 \times 10^{-11}$; Fig. 2F, Fig. S3). Despite this qualitative association between mutational effects,

overall, the magnitude of fitness effects was not correlated across environments (only 38 of 120

correlations were significant in WT and 22 of 120 significant in mutator, Spearman's rank

correlation, $p < 0.05$). Thus, new beneficial mutations in the mutator are more likely to facilitate

adaptation across many environments, but are no more likely to generate tradeoffs. Interestingly,

simulating an increase in the median fitness effect of the WT DFE without changing its shape ("WT-

beneficial shift") mimicked pleiotropic effects observed in the mutator in all but two environments

(Fig. 2D, Fig. S2E, Table S6). Conversely, simply reducing the median fitness effect in the WT

DFE ("WT-deleterious shift") lowered beneficial synergistic pleiotropy in all 16 environments (Fig.

2E, Fig. S2F, Table S6). Ignoring the mutation spectrum can thus cause overestimation of a

mutator's genetic load, and underestimation of the supply and pleiotropic effects of beneficial

mutations.

The mutator's beneficial-shifted DFEs arise from its distinct mutation spectrum

We had hypothesized that the mutator's distinct DFE could arise from its biased spectrum.

However, other global effects of the initial *mutY* gene deletion could also lead to a DFE shift. For

instance, if deleting *mutY* reduced the mutator's fitness, a larger fraction of new mutations might

be beneficial (47). However, the mutator ancestor had lower fitness than WT in only three

environments (Fig. 3A, Table S7). Alternatively, epistatic interactions with *mutY* could increase the

deleterious effect of new mutations in the WT. However, paired WT and mutator strains carrying

the same mutation had similar fitness in all but one environment (Fig. 3B, Table S8). Thus, the

observed shift in the mutator DFE cannot be explained by global effects of the *mutY* deletion. The

difference across strains was also robust across mutational classes: mutations were more

beneficial in the mutator even when comparing only base-pair substitutions, or coding, non-coding,

synonymous, or non-synonymous mutations (Fig. S4).

Next, we asked whether the mutator is biased towards mutational classes that happen to be more

beneficial (or less deleterious). We defined mutation bias as the proportion of mutations belonging

to a specific mutational class (e.g., Tv bias = Number of Tv mutations / (Number of Tv + Number

of Ts mutations)). Note that the null expectation ("unbiased" state) differs across each type of bias.

For Tv bias, the null is 0.67 because 2/3rd of all possible single nucleotide changes are

transversions. For GC→AT bias, the null is 0.5 because for a randomly generated genome half of

all mutations should be GC→AT. For actual genomes with extreme GC content, the null

expectation will change (e.g., in a GC-rich genome, the baseline probability of AT→GC mutations

is less than 0.5 because there are few A or T bases that can mutate). This is not a major problem

for *E. coli* since its genomic GC content is ~50%. Regardless of the type of bias, we could compare

mutation biases across the new mutations sampled by WT and mutator, because barring the

deletion of *mutY* both ancestral genomes were identical. The mutator strongly favours

transversions, GC→AT mutations, base pair substitutions (BPS) and coding mutations (Fig. 3C,

Table S3). Overall, transversions and GC→AT mutations were less deleterious than transitions

and AT→GC mutations respectively (Fig. 3D–E; fitness effects pooled across environments, Table

S9), but the other mutation biases did not show consistent differences (Fig. S4, Table S9).

Additionally, other aspects of the mutations (e.g., genomic location or specific amino acid changes)

also did not differ across strains (Fig. S5, S6; Tables S10, S11). Thus, as predicted, the mutation

spectrum – specifically the strong Tv and GC→AT bias – appeared to drive the distinct DFE of the

mutator, because the mutator over-samples these specific mutational classes. However, it was

unclear why Tv and GC→AT mutations should be more beneficial.

Simulations demonstrate a general benefit of reversing mutation spectra

To test the generality of our empirical results and uncover underlying mechanisms, we simulated

adaptive walks in an NK fitness landscape ((48), see Methods), modelling sequences of N

nucleotides such that mutations could be classified as Ts vs. Tv or GC→AT vs. AT→GC. Starting

from a randomly chosen ancestor sequence and mutation spectrum (Tv bias = fraction of

transversions), we allowed populations to explore successive mutational steps at randomly chosen

loci (bases). A mutation that confers a relative fitness increase of 1+s fixes (a step in the walk is

taken) with probability 2s; that is, we assumed a strong-selection weak-mutation (SSWM) regime.

At various points during the walk, we simulated mutations to generate a DFE, computed fitness

effects given the underlying fitness landscape (affected by K other randomly chosen loci, to

incorporate epistasis), and calculated the fraction of beneficial (f_b) and deleterious (f_d) mutations

and their effect sizes. We initially set N=200 and K=1 (each locus epistatically coupled to

1/199=0.5% loci), and present the average outcomes for 500 adaptive walks on 500 randomly-

generated fitness landscapes (Fig. S7A–B shows variation across walks).

As expected, mean population fitness increased during the adaptive walk, with a concomitant

reduction in f_b and beneficial effect size, an increase in f_d, and a relatively constant deleterious

effect size (Fig. 4A). Setting ancestral Tv bias to 0.45 (mimicking our WT), we compared the DFE

generated by the ancestor at various time points, to the DFE that would be created if the bias were

changed at that time point (mimicking our Δ*mutY* strain). The bias-shifted ("shifted") strain thus

has the same fitness and sequence as the ancestor and an identical mutation rate, differing only

in spectrum. As the ancestor evolved (i.e., f_b decreased), a stronger bias shift (i.e., higher Tv)

allowed the bias-shifted strain to sample proportionally more beneficial mutations than the ancestor

(Fig. 4B). More generally, exploring well-adapted populations (f_b = 0.04) but varying both the

ancestral and shifted mutation spectra, we find that shifted strains that reverse the ancestral bias

(i.e., have a strong bias in the opposite direction) have the greatest advantage (Fig. S8). Our results

hold across a different axis of the mutation spectrum (GC→AT bias, Fig. S9), and in more rugged

fitness landscapes with higher epistasis (Fig. S10A–B). Importantly, adapting populations did not

"exhaust" a small set of beneficial mutations; rather, they fixed distinct mutations as they explored

a large number of diverse pathways that represent a subset of many possible fitness peaks (Fig.

S10C). Our results also hold in full population simulations with multiple segregating mutations

(relaxing the SSWM assumption), where deleterious mutations can fix (Fig. 4C–D, Fig. S11). In

fact, when more mutations are accessible, we observe the same beneficial effects of bias

reversals, but these effects emerge more quickly. Finally**,** these effects are also observed in full

population simulations when the fitness landscape is defined using a codon-based rather than a

nucleotide model (Fig. S12).

Thus, our simulation results can be broadly generalized in the form of three key predictions: (1)

reversing an ancestral mutation bias is always beneficial (2) the magnitude of the benefit increases

with the degree to which the new bias opposes the ancestral bias, and (3) the magnitude of the

benefit increases when the ancestor has evolved with a given bias for longer intervals of time (i.e.,

as the ancestor gets better-adapted over time). Our experimental data with the Δ*mutY* strain

support the first prediction, since we found that overall the mutator (with a reversal of the Tv bias,

Fig. 3D) had higher f_b values than WT. Interestingly, the WT GC→AT bias was reinforced in the

mutator, rendering the results in Fig. 3E puzzling. However, ~88% of all GC→AT mutations in the

mutator are also transversions. Hence, our mutator's observed beneficial-shifted DFE is explained

by its strong Tv bias, which opposes the existing WT bias towards Ts.

Although our data are generally explained by the simulation results, some puzzling observations

remain. The simulations predict that the mutator should have higher f_b in all environments, but we

do not observe this pattern as consistently as expected (Fig. 1). Further, in apparent contradiction

to prediction 3 above, the difference in f_b between WT and mutator in lowest in LB (where ancestral

growth rate is highest and presumably *E. coli* is best-adapted). However, note that our strains

evolved for a very short time in LB under minimal selection, so LB is not an "ancestral" environment

in an evolutionarily meaningful sense. Additionally, absolute growth rate is not a reliable indicator

of the relative degree of or period of adaptation to an environment, because it is also strongly

influenced by nutrient availability. Indeed, f_b was not correlated with ancestral growth rate across

environments in either strain (Spearman's rank correlation test, WT: rho = –0.217, Mutator: rho =

–0.18; p > 0.05 in each case). Hence, to understand the environment-specific differences in DFEs,

we need to explicitly test predictions 2 and 3 (described above) under conditions where the precise

degree of bias change and the length of evolutionary time in an environment can be measured

and/or manipulated.

Our simulation results demonstrate general conditions under which a change in mutational bias

can be selectively advantageous. But how does mutation bias evolve in the first place? Starting

from an unbiased ancestor, we find that introducing any bias is selectively neutral, regardless of

how long the ancestor had evolved in the same fitness landscape (Fig. 4E–F). However, if the

ancestor had previously evolved with a biased spectrum (e.g., our WT), the population would have

already sampled many beneficial mutations within the class favoured by the existing bias. Hence,

a shift in the opposite direction (e.g., in our Δ*mutY* strain) allows the population to explore

mutational space that was previously under-sampled, increasing the probability of finding new

beneficial mutations. Importantly, this means that the benefit of a specific mutation class (e.g., Tv)

is not universal, but is context-dependent. Increased sampling of any mutational class that was

previously poorly explored would therefore be beneficial. Thus, after a period of evolving with a

new spectrum, a change in the fitness landscape (e.g., due to epistasis or environmental change)

may again render a spectrum shift advantageous. Together, these results generate two additional,

broad predictions: (a) starting from an unbiased spectrum, a biased spectrum can only evolve

through drift, and (b) after a period of adaptation with a specific mutation bias (and/or following

environmental change), reversing the bias becomes beneficial. We tested the second prediction

using a broad phylogenetic analysis of bacterial taxa, expecting that evolutionary shifts in mutation

spectra should be enriched in bias reversals.

Evolutionary transitions in DNA repair enzymes indicate mutation spectrum reversals in

most bacterial lineages

Altering DNA repair function is a simple and instantaneous mechanism to generate spectrum shifts.

Hence, to infer long-term mutation spectrum dynamics we mapped evolutionary transitions in 11

bacterial DNA repair enzymes with empirically measured effects on mutation spectra, i.e., whose

loss changes the Tv or GC→AT bias in *E. coli* (Table S12). Identifying gene orthologues in 1093

extant bacterial genomes, we used ancestral reconstruction to infer enzyme gains and losses

across a pruned tree derived from a previously published whole-genome bacterial phylogeny

based on >400 proteins (following (49); Fig. 5A). Broadly consistent with previous work with a few

DNA repair genes (50), we found frequent evolutionary transitions in most repair enzymes with

known spectrum effects (Fig. 5A), as well as for 33 other genes with unknown effects on mutation

spectra (Fig. S13). Using the predicted set of enzymes present in each hypothetical ancestor

(nodes), we estimated the Tv and GC→AT bias at each node relative to WT *E. coli* and traced the

change in relative bias over time (Fig. 5B, Fig. S14A). Consistent with our simulation results,

successive enzyme gain or loss events led to a bias reversal in >80% lineages, and we observed

only a few cases of reinforced bias (Fig. 5C). This is distinct from the pattern expected by chance

(10,000 phylogenetic stochastic simulations using similar enzyme transition rates; Fig. 5C, Fig.

S14B). Notably, bias reversals are more frequent than expected (one-sample t-tests, Tv bias: t = -

72.38, $p<2\times10^{-16}$; GC→AT bias: t = -59.42, $p<2\times10^{-16}$) while reinforcements are less frequent than

expected (Tv bias: t = -72.38, $p<2\times10^{-16}$; GC→AT bias: t = -59.42, $p<2\times10^{-16}$) (Fig. S14C–D).

Hence, our results cannot be attributed to peculiarities of the enzyme set, tree topology, or number

of evolutionary transitions. Instead, we suggest that such frequent bias reversals likely reflect

selection on mutation rate and/or spectrum. Although it is difficult to separate these effects, two

pieces of evidence suggest that the observed dynamics at least partly reflect selection for reversal

of mutation spectra. First, enzymes with a weak effect on mutation rate but a strong impact on

spectrum are also highly dynamic (e.g. *ung* and *mutM*; Table S12). Second, for enzymes with a

strong impact on both mutation rate and spectrum, either bias reversals or reinforcements would

change mutation rates. Hence, selection for mutation rate alone cannot explain the higher-than-

expected frequency of bias reversals that we observe.

We discuss here a few limitations of this analysis. First, reconstruction of ancestral character states

in phylogenetic datasets is not always accurate and does not account for uncertainty in the

underlying phylogeny (51). Second, the estimated biases do not account for other repair enzymes

whose effects on the mutation spectrum are unknown, or for variation in the impact of enzyme loss

or gain. We also likely underestimate the actual number of bias shifts because we could not include

cases where repair function may be altered without gene deletion (e.g., due to loss-of-function

point mutations). However, given the lack of relevant empirical data for most bacteria, the values

we report are our best estimates of the mutation biases of diverse bacteria relative to *E. coli*, and

quantify at least a proportion of the "true" changes in mutation spectra that have occurred across

lineages. Available experimental data also show that the qualitative effects of deleting a specific

DNA repair enzyme are largely consistent across species (Table S13), suggesting that the broad

trends observed here should be robust. Finally, recent work indicates similar dynamics of genes

involved in DNA repair in bacteria (52) and fungi (53). Thus, sustained shifts in mutation spectra

likely occur frequently, and their long-term impacts on the pool of genetic variants available for

selection deserve further investigation.

CONCLUSIONS

We show that pervasive mutational biases can evolve by genetic drift, whereas shifts in the

strength and direction of bias could evolve under selection. Thus, a species' evolutionary history

may be sufficient to explain the evolution of genomic bias (e.g., in base composition), without

invoking specific underlying mechanisms or selection favouring particular mutational classes.

Previous work showed that the transversion-biased mutators Δ*mutT* and Δ*mutY* have distinct DFEs

for antibiotic resistance compared to WT, but the effect was attributed to transversions being more

beneficial for the specific antibiotic target (13). However, by recasting the puzzle of mutational

biases as a problem of searching a vast and dynamic mutational space, our model allows broad

generalization across mutation spectra. We suggest that the rarity of species with unbiased

mutation spectra is not surprising, since shifts from unbiased to biased spectra may occur easily

and frequently via changes in DNA repair function and are unopposed by selection. Hence, a

biased spectrum should be the norm rather than the exception.

While it is well established that access to _more_ mutations (through an increased mutation rate) is

advantageous (18), here we demonstrate that accessing _different_ mutations can likewise provide

an advantage. Our results have important implications for the evolution of mutator genotypes, in

which shifts in mutation spectra and mutation rates are often (though not always) entangled.

Mutator dynamics are governed by the genetic load and supply of beneficial mutations (18, 33,

54). In a new environment (away from fitness optima), these parameters are primarily driven by

mutators' high mutation rate (55) – particularly under strong selection (56) – allowing them to

hitchhike with beneficial mutations (57). We show here that the genome-wide beneficial supply

rate is strongly influenced by the mutation spectrum across environments, and this effect is

strongest in well-adapted populations. Further, we predict that the advantage of a high mutation

rate should be enhanced if accompanied by a reversal of the ancestral spectrum, but diminished

otherwise. A recent simulation study also predicts that differences in the mutation spectrum of

mutators can change the magnitude of deleterious mutations; such differences can facilitate the

loss of mutators under selection (58). Thus, mutation rate and spectrum may jointly govern the

observed rise, persistence, and fall of mutators under selection (13, 33, 59). Such effects may

explain why mutators with relatively small increases in mutation rate are also abundant in natural

bacterial populations (60), and deserve further attention (13, 58) with theoretical and experimental

analyses to dissect the role of mutation rate vs. spectrum.

Together with studies showing that mutation biases are pervasive and influence the genetic basis

of adaptation under diverse conditions (see Introduction), our results demonstrate that mutation

spectra may be important drivers of adaptation and innovation under myriad scenarios. When the

beneficial mutation supply is limited by the existing spectrum, an antiparallel (opposite direction on

the same axis) or orthogonal jump (on a different axis of the spectrum) could enhance sampling of

new beneficial mutations, facilitating rapid adaptation. Our phylogenetic analysis likely

underestimated such evolutionary shifts in mutation spectra, which may occur even more

frequently on shorter timescales via horizontal transfer and recombination (61), and drive

polymorphism in spectra across natural bacterial isolates (60, 62). We predict multiple cascading

effects of shifts in mutation spectra, including a reduction in the waiting time for beneficial

mutations, decreased likelihood of mutational meltdown under genetic drift, and distinct genetic

pathways of adaptation. We hope that future work will test these predictions.

METHODS SUMMARY

Here, we provide a summary of the methods important to understand our study. Details are given

in Supplementary Materials and Methods.

Bacterial strains and mutation accumulation (MA) experiments

We obtained the wild-type (WT) *E. coli* K-12 MG1655 strain and mutator strains (Δ*mutY*, Δ*ung* and

Δ*mutM* in *E. coli* K-12 BW25113) from the Coli Genetic Stock Centre (CGSC, Yale University). We

moved the Δ*mutY*, Δ*ung* and Δ*mutM* loci into the WT strain using P1-phage transduction and

confirmed the knockout using PCR and whole genome sequencing. To test whether genetic

background alters the fitness effects of single mutations observed during mutation accumulation

(MA), we deleted the *mutY* locus from 19 evolved WT MA lines using P1-phage transduction.

We founded WT, Δ*mutY*, Δ*ung* and Δ*mutM* MA lines from single colonies, and propagated them

through daily single-colony bottlenecks on LB agar for 8250, 330, 4400 and 5775 generations

respectively. The MA protocol minimizes the effect of selection, allowing us to sample a wide range

of mutations largely independent of their fitness consequences.

Whole-genome sequencing to identify single mutations and determine mutation spectra

We sequenced individual colonies from MA experiments to identify all clones carrying only a single

mutation relative to the WT or Δ*mutY* ancestor, using the Illumina HiSeq 2500 platform. For each

sample, we aligned quality-filtered reads to the NCBI reference *E. coli* K-12 MG1655 genome

(RefSeq accession ID GCA_000005845.2) and generated a list of base-pair substitutions and short

indels (<10bp). We calculated mutational biases as the fraction of each class of mutations

observed in our evolved strains (Table S3).

Constructing single-mutation DFEs

We measured growth rates of all evolved MA isolates with single mutations and their respective

ancestors in 16 different environments, and estimated maximum growth rate as a proxy of fitness.

For each isolate, we used the average growth rate of three technical replicates to calculate the

fitness effect (*s*) as (growth rate of evolved isolate/growth rate of ancestor) – 1. We used *s* values

of mutations to construct strain- and environment-specific distributions of fitness effects (DFE).

Although selection is minimized in MA experiments, it is not entirely avoided. We corrected both

WT and mutator DFEs for the effect of such selection as described recently (32), for each

environment. Calculations are shown in the Supplementary Data file.

Estimating genetic load, beneficial mutation supply, and pleiotropy

We used the mutation rate and f_b (estimated via the DFE) of WT and mutator in each environment

to estimate the deleterious load and supply of beneficial mutations, as described earlier (33)

(Tables S4 and S5). To predict the impact of the strain-specific DFEs on adaptation in new habitats,

we estimated the incidence of antagonistic and synergistic pleiotropy among new mutations

sampled by each strain. Results for WT *E. coli* in some environments were previously reported

(30). We similarly estimated pleiotropy of the 79 single mutations that arose in the mutator (Δ*mutY*)

background, and compared the incidence of pleiotropy across strains.

Simulations to test the effect of mutation spectrum changes

We used an established evolutionary model to test whether and to what extent changes in mutation

spectra are generally expected to affect the DFE. Following Stoltzfus (63), we simulated adaptive

walks on the NK fitness landscape (48), modelling sequences composed of strings of four bases

(A, C, G, T), allowing mutations to be classified as transitions or transversions, GC→AT or AT→GC,

and so on. In an NK fitness landscape, each locus in the sequence is randomly assigned K

neighbours from among all loci in the model. The fitness contribution of locus i (ω_i) then depends

on the state of that locus and the state of its neighbours. For computational efficiency we initially

simulated adaptive walks (strong-selection weak-mutation regime). At various times during the

adaptive walk, we generated a DFE by randomly sampling new mutations from the set of available

single-step mutations and estimated key parameters such as f_b.

Next, we relaxed a number of assumptions to test the generality of the simulation results. We

investigated higher degrees of epistasis by varying both N and K. We also simulated full

populations, in which large numbers of mutations simultaneously segregate, and deleterious

mutations can fix. Finally, we implemented a codon-based model in which nucleotide sequences

evolved as described above, but fitness was defined by an NK fitness landscape based on amino

acid sequence.

Inferring past evolutionary changes in mutation spectra

We used data from mutation accumulation studies with *E. coli* to obtain the predicted effects of the

loss of 11 genes encoding DNA repair enzymes on the Tv bias and GC→AT bias (Table S12). To

determine the evolutionary history of these genes, we detected orthologues and mapped predicted

gain and loss events for each gene across the bacterial phylogeny, as described previously (49).

We used the inferred state of all enzymes at every node in the tree (including the eubacterial

ancestor and the extant taxa) to estimate the Tv and GC$\rightarrow$AT bias at each node, relative to WT *E.*

coli. For all lineages in the tree, we traced the change in bias over time, up to the extant taxon and

used these data to calculate the observed number and direction of changes in bias in each lineage,

categorising them as "no change", "single change", "reinforcement" and "reversal" (Fig. S14A). To

derive expected changes in bias based on chance, we ran forward simulations on the transition

rate matrices (describing inferred rates of enzyme state transitions) and used the number and

direction of mutation bias changes in each simulation to calculate the average proportion of

lineages showing a given number of reversals (or other changes in mutation bias) expected by

chance (Fig. S14B).

Acknowledgements

We thank Shyamsunder Buddh, Brian Charlesworth, Deborah Charlesworth, Joachim Krug,

Krushnamegh Kunte, Saurabh Mahajan, Christopher Marx, and Mukund Thattai for discussion;

Joshua Miranda for assistance with MA experiments; and the NGS facility at the Bangalore Life

Sciences Cluster. **Funding:** We acknowledge funding and support from the National Centre for

Biological Sciences (NCBS-TIFR), the Council for Scientific and Industrial Research India (Senior

Research Fellowship to MS), the University Grants Commission of India (Senior Research

Fellowship to GDD), the Department of Science and Technology India (KVPY Fellowship to BAB),

the Natural Sciences and Engineering Research Council of Canada (LMW), and the Wellcome

Trust (GDD is supported by grant 210585/B/18/Z to Robert B. Russell; and DBT/Wellcome Trust

India Alliance grant IA/I/17/1/503091 to DA). We also thank the International Centre for Theoretical

Sciences (ICTS) for supporting the Bangalore School on Population Genetics and Evolution (code:

ICTS/popgen2020/01), where this collaboration was initiated. **Competing interests:** Authors

declare no competing interests. **Author contributions:** MS designed and conducted experiments,

analysed data, and drafted the manuscript. GDD designed and conducted phylogenetic analyses.

BAB conducted experiments and analysed data. LMW obtained funding, designed and conducted

simulation analyses, and drafted the manuscript. DA conceived and designed experiments and

phylogenetic analyses, analysed data, obtained funding, and wrote the manuscript. **Data and**

materials availability: All data used for experimental and phylogenetic analysis are available as

Supplementary Information files. Code for simulations and phylogenetic analysis is available on

Github (https://github.com/lmwahl/MutationSpectrum and

https://github.com/gauravdiwan89/dfe_ms_phylo_analysis).

Figure 1. Mutator DFEs have a larger fraction of beneficial mutations than WT. Stacked bar plots show the observed proportion of beneficial (f_b), deleterious and neutral mutations in WT and mutator (Mut). Average values across environments are given in the key; asterisks indicate significant differences between WT and mutator f_b, colored to show the strain with higher f_b (Table S2).

Figure 2. Beneficial-shifted DFE alters pleiotropic effects. (A) Schematic showing possible
pleiotropic effects of mutations in two environments (W: relative fitness). (B–E) Median proportions
of pleiotropic fitness effects (across 15 focal environments, excluding LB; also see Fig. S2C–F) of
mutations in WT and mutator, and WT DFE artificially shifted towards either beneficial or
deleterious mutations. Schematics indicate global DFEs (WT=cyan, mutator=pink, WT-
shifted=light blue); black lines: relative fitness = 1. (F) Pleiotropic effects of mutations across
multiple environments. Frequency distributions show the proportion of mutations with consistent
fitness effects (beneficial or deleterious) in a given number of environments. Black lines indicate
medians.
493

Figure 3. Differences in WT and mutator DFEs are best explained by differences in mutation spectra. (A) Growth rate of WT and mutator ancestors (n=12–24/strain/environment). Asterisks: significant differences between strains (Table S7). (B) Pairwise differences in the fitness effects of 19 mutations across genetic backgrounds (Mutator – WT); asterisks: significant difference from zero (black dotted line; Table S8). (C) Observed mutation biases in WT and mutator MA lines (including those with multiple mutations); the number of mutations of each type is noted. Asterisks: significant differences in proportions across strains; dashed line: no bias (Tv bias = 0.67, GC→AT bias = 0.5) (Table S3). (D–E) Fitness effects of different mutation classes, for MA-evolved isolates carrying single mutations. Boxplots (colored by strain and mutation class) show median, quartiles and range of relative fitness effects of mutations (open diamonds: outliers). Each boxplot was constructed with n x 16 data points (n = number of mutations, shown below each plot; pooled effects from 16 environments). Black horizontal lines: fitness impact = 0. Asterisks: significant differences across mutation classes (ANOVA output, Table S9; for imbalanced sample sizes, permutation tests as described for Fig. S4); nd: not determined.

Figure 4. Simulations show a general benefit of reversing mutation spectra. (A) Change in
mean population fitness, fraction of beneficial and deleterious mutations (f_b and f_d), and mean
magnitude of beneficial and deleterious effects (s_b and s_d) over the course of adaptive walks of a
WT ancestor. (B) Impact of altering the WT mutation bias on f_b at different points along the adaptive
walk (Tv bias = Tv/(Ts+Tv) mutations; grey dashed line: WT bias), as a function of the new mutation
bias (shifted Tv bias). Line color: WT f_b at a given time; steeper lines (lighter colors): later
evolutionary time, better adapted WT. (C–D) Results as in (A–B) but for simulated populations of
100,000 individuals evolving for up to 100,000 generations with a mutation rate of 10^{-4} per genome
per generation; the mean number of segregating genotypes in each generation was 77.8. (E–F)
Impact of introducing a mutation bias in an unbiased ancestor ("anc"), as a function of the new bias
(shifted Tv or GC→AT bias). Panels show the results of simulations identical to those described in
panel B, but starting with an unbiased ancestor.
522

Figure 5. Phylogenetic analysis shows frequent evolutionary transitions in DNA repair genes and reversals in mutation spectra. (A) Patterns of gene presence/absence in extant taxa, used to map evolutionary transitions in DNA repair genes on the bacterial phylogeny (total transitions are shown in the key), and infer the magnitude of Tv and GC→AT biases relative to WT *E. coli* in extant taxa (heatmaps) based on experimental data from *E. coli* (Table S12). (B) Inferred trajectories of relative mutational biases across 1093 lineages (C) Observed number of lineages showing spectrum shifts due to successive evolutionary transitions in repair enzymes (% values calculated as fraction across all lineages; blue=observed; black=expected values, mean of 10,000 stochastic forward simulations, Fig. S14 C–D).

References

1. D. Agashe, N. Shankar, The evolution of bacterial DNA base composition. *Journal of Experimental*
Zoology Part B: Molecular and Developmental Evolution 322, 517–528 (2014).
2. H. Long, *et al.*, Evolutionary determinants of genome-wide nucleotide composition. *Nature Ecology*
& Evolution 48, 582 (2018).
3. J. Bohlin, O. Brynildsrud, T. Vesth, E. Skjerve, D. W. Ussery, Amino acid usage is asymmetrically
biased in AT-and GC-rich microbial genomes. *PLoS ONE* 8, e69878 (2013).
4. A. Couce, *et al.*, Mutator genomes decay, despite sustained fitness gains, in a long-term
experiment with bacteria. *Proceedings of the National Academy of Sciences of the United States*
of America 259, 201705887–10 (2017).
5. A. Stoltzfus, L. Y. Yampolsky, Climbing Mount probable: Mutation as a cause of nonrandomness in
evolution. *Journal of Heredity* 100, 637–647 (2009).
6. M. A. Streisfeld, M. D. Rausher, Population genetics, pleiotropy, and the preferential fixation of
mutations during adaptive evolution. *Evolution* 65, 629–642 (2010).
7. D. R. Rokyta, P. Joyce, S. B. Caudle, H. A. Wichman, An empirical test of the mutational landscape
model of adaptation using a single-stranded DNA virus. *Nature Genetics* 37, 441–444 (2005).
8. K. Gomez, J. Bertram, J. Masel, Mutation bias can shape adaptation in large asexual populations
experiencing clonal interference. *Proc Royal Soc B* 287, 20201503 (2020).
9. A. V. Cano, J. L. Payne, Mutation bias interacts with composition bias to influence adaptive
evolution. *Plos Comput Biol* 16, e1008296 (2020).
10. A. Stoltzfus, D. M. McCandlish, Mutational biases influence parallel adaptation. *Molecular Biology*
and Evolution 34, 2163–2172 (2017).
11. J. F. Storz, *et al.*, The role of mutation bias in adaptive molecular evolution: insights from
convergent changes in protein function. *Philosophical Transactions of the Royal Society B:*
Biological Sciences 374, 20180238–10 (2019).
12. A. V. Cano, H. Rozhoňová, A. Stoltzfus, D. M. McCandlish, J. L. Payne, Mutation bias shapes the
spectrum of adaptive substitutions. *Proc National Acad Sci* 119, e2119720119 (2022).
13. A. Couce, J. R. Guelfo, J. Blázquez, Mutational spectrum drives the rise of mutator bacteria. *PLoS*
Genetics 9, e1003167 (2013).
14. J. L. Payne, *et al.*, Transition bias influences the evolution of antibiotic resistance in
Mycobacterium tuberculosis. *PLoS Biology* 17, e3000265-23 (2019).
15. A. Bachar, *et al.*, Point mutations in topoisomerase I alter the mutation spectrum in E. coli and
impact the emergence of drug resistance genotypes. *Nucleic Acids Research* 48, 761–769 (2019).
16. S. M. Leighow, C. Liu, H. Inam, B. Zhao, J. R. Pritchard, Multi-scale predictions of drug resistance
epidemiology identify design principles for rational drug design. *Cell Reports* 30, 3951-3963.e4
(2020).
17. P. L. Foster, H. Lee, E. Popodi, J. P. Townes, H. Tang, Determinants of spontaneous mutation in
the bacterium Escherichia coli as revealed by whole-genome sequencing. *Proceedings of the*
National Academy of Sciences of the United States of America, 201512136–10 (2015).
18. Y. Raynes, P. D. Sniegowski, Experimental evolution and the dynamics of genomic mutation rate
modifiers. *Heredity* 113, 375–380 (2014).
19. R. P. Maharjan, T. Ferenci, A shifting mutational landscape in 6 nutritional states: Stress-induced
mutagenesis as a series of distinct stress input–mutation output relationships. *PLoS Biology* 15,
e2001477-22 (2017).
20. S. Shewaramani, *et al.*, Anaerobically grown Escherichia coli has an enhanced mutation rate and
distinct mutational spectra. *PLoS Genetics* 13, e1006570-22 (2017).
21. P. Jiang, *et al.*, A modified fluctuation assay reveals a natural mutator phenotype that drives
mutation spectrum variation within Saccharomyces cerevisiae. *Elife* 10, e68285 (2021).
22. E. López-Cortegano, *et al.*, De novo mutation rate variation and its determinants in
Chlamydomonas. *Mol Biol Evol* 38, msab140- (2021).
23. T. Ferenci, Irregularities in genetic variation and mutation rates with environmental stresses.
Environmental Microbiology 21, 3979–3988 (2019).
24. V. Katju, U. Bergthorsson, Old trade, new tricks: Insights into the spontaneous mutation process
from the partnering of classical mutation accumulation experiments with high-throughput genomic
approaches. *Genome Biology and Evolution* 11, 136–165 (2018).
25. L. Y. Yampolsky, A. Stoltzfus, Bias in the introduction of variation as an orienting factor in
evolution. *Evolution and Development* 3, 73–83 (2001).
26. D. Agashe, The road not taken: Could stress-specific mutations lead to different evolutionary
paths? *PLoS Biology* 15, e2002862-5 (2017).

27. A. Stoltzfus, R. W. Norris, On the causes of evolutionary transition:transversion bias. *Molecular*
Biology and Evolution 33, 595–602 (2016).
28. D. M. Lyons, A. S. Lauring, Evidence for the Selective Basis of Transition-to-Transversion
Substitution Bias in Two RNA Viruses. *Mol Biol Evol* 34, 3205–3215 (2017).
29. Z. Zou, J. Zhang, Are nonsynonymous transversions generally more deleterious than
nonsynonymous transitions? *Mol Biol Evol* 38, msaa200- (2020).
30. M. Sane, J. J. Miranda, D. Agashe, Antagonistic pleiotropy for carbon use is rare in new
mutations. *Evolution* 72, 2202–2213 (2018).
31. D.-E. Chang, *et al.*, Carbon nutrition of Escherichia coli in the mouse intestine. *P Natl Acad Sci*
Usa 101, 7427–7432 (2004).
32. L. M. Wahl, D. Agashe, Selection bias in mutation accumulation. *Evolution* 76, 528–540 (2022).
33. S. Wielgoss, *et al.*, Mutation rate dynamics in a bacterial population reflect tension between
adaptation and genetic load. *Proceedings of the National Academy of Sciences* 110, 222–227
(2013).
34. A. Eyre-Walker, P. D. Keightley, The distribution of fitness effects of new mutations. *Nature*
Reviews Genetics 8, 610–618 (2007).
35. M. Weng, *et al.*, Fitness effects of mutation in natural populations of Arabidopsis thaliana reveal a
complex influence of local adaptation. *Evolution* (2020) https:/doi.org/10.1111/evo.14152.
36. C. Zeyl, J. A. DeVisser, Estimates of the rate and distribution of fitness effects of spontaneous
mutation in Saccharomyces cerevisiae. *Genetics* 157, 53–61 (2001).
37. D. W. Hall, S. B. Joseph, A High Frequency of Beneficial Mutations Across Multiple Fitness
Components in Saccharomyces cerevisiae. *Genetics* 185, 1397–1409 (2010).
38. S. B. Joseph, D. W. Hall, Spontaneous Mutations in Diploid Saccharomyces cerevisiae More
Beneficial Than Expected. *Genetics* 168, 1817–1825 (2004).
39. W. J. Dickinson, Synergistic Fitness Interactions and a High Frequency of Beneficial Changes
Among Mutations Accumulated Under Relaxed Selection in Saccharomyces cerevisiae. *Genetics*
178, 1571–1578 (2008).
40. L. Perfeito, L. Fernandes, C. Mota, I. Gordo, Adaptive mutations in bacteria: high rate and small
effects. *Science* 317, 813–815 (2007).
41. C. Kozela, M. O. Johnston, Effect of Salt Stress on Mutation and Genetic Architecture for Fitness
Components in Saccharomyces cerevisiae. *G3 Genes Genomes Genetics* 10, g3.401593.2020
(2020).
42. R. G. Shaw, F. H. Shaw, C. Geyer, What fraction of mutations reduces fitness? A reply to
Keightley and Lynch. *Evolution* 57, 686–689 (2003).
43. J. W. Thatcher, J. M. Shaw, W. J. Dickinson, Marginal fitness contributions of nonessential genes
in yeast. *Proc National Acad Sci* 95, 253–257 (1998).
44. M. Imhof, C. Schlötterer, Fitness effects of advantageous mutations in evolving Escherichia coli
populations. *Proc National Acad Sci* 98, 1113–1117 (2001).
45. F. H. Shaw, C. J. Geyer, R. G. Shaw, A comprehensive model of mutations affecting fitness and
inferences for Arabidopsis thaliana. *Evolution* 56, 453–463 (2002).
46. R. G. Shaw, D. L. Byers, E. Darmo, Spontaneous Mutational Effects on Reproductive Traits of
Arabidopsis thaliana. *Genetics* 155, 369–378 (2000).
47. H. A. Orr, The population genetics of adaptation: The distribution of factors fixed during adaptive
evolution. *Evolution* 52, 935–949 (1998).
48. C. A. Macken, A. S. Perelson, Protein evolution on rugged landscapes. *Proceedings of the*
National Academy of Sciences of the United States of America 86, 6191–6195 (1989).
49. G. D. Diwan, D. Agashe, Wobbling Forth and Drifting Back: The Evolutionary History and Impact
of Bacterial tRNA Modifications. *Mol Biol Evol* 35, 2046–2059 (2018).
50. J. A. Eisen, P. C. Hanawalt, A phylogenomic study of DNA repair genes, proteins, and processes.
Mutation research 435, 171–213 (1999).
51. J. B. Losos, Seeing the Forest for the Trees: The Limitations of Phylogenies in Comparative
Biology: (American Society of Naturalists Address). *Am Nat* 177, 709–727 (2011).
52. M. Sharda, A. Badrinarayanan, A. S. N. Seshasayee, Evolutionary and Comparative analysis of
bacterial Non-Homologous End Joining Repair. *Genome Biol Evol* 12, evaa223- (2020).
53. M. A. Phillips, J. L. Steenwyk, X.-X. Shen, A. Rokas, Examination of gene loss in the DNA
mismatch repair pathway and its mutational consequences in a fungal phylum. *Genome Biol Evol*,
evab219- (2021).
54. K. Jain, A. James, Fixation probability of a nonmutator in a large population of asexual mutators.
Journal of Theoretical Biology 433, 85–93 (2017).

55. J. A. G., *et al.*, Diminishing Returns from Mutation Supply Rate in Asexual Populations. *Science*
283, 404–406 (1999).
56. O. Tenaillon, B. Toupance, H. L. Nagard, F. Taddei, B. Godelle, Mutators, population size,
adaptive landscape and the adaptation of asexual populations of bacteria. *Genetics* 152, 485–493
(1999).
57. C. F. Gentile, S.-C. Yu, S. A. Serrano, P. J. Gerrish, P. D. Sniegowski, Competition between high-
and higher-mutating strains of Escherichia coli. *Biology Letters* 7, 422–424 (2010).
58. A. Couce, O. Tenaillon, Mutation bias and GC content shape antimutator invasions. *Nature*
communications 10, 3114–9 (2019).
59. L. Notley-McRobb, S. Seeto, T. Ferenci, Enrichment and elimination of mutY mutators in
Escherichia coli populations. *Genetics* 162, 1055–1062 (2002).
60. I. Matic, *et al.*, Highly variable mutation rates in commensal and pathogenic Escherichia coli.
Science 277, 1833–1834 (1997).
61. E. Denamur, *et al.*, Evolutionary implications of the frequent horizontal transfer of mismatch repair
genes. *Cell* 103, 711–721 (2000).
62. J. E. LeClerc, B. Li, W. L. Payne, T. A. Cebula, High mutation frequencies among Escherichia coli
and Salmonella pathogens. *Science* 274, 1208–1211 (1996).
63. A. Stoltzfus, Mutation-biased adaptation in a protein NK model. *Molecular Biology and Evolution*
23, 1852–1862 (2006).
676