**Characterization of ethylene-inducible pedicel-fruit abscission zone formation in non-**

**climacteric sweet cherry (*Prunus avium* L.)**

Seanna Hewitt[1,2*], Benjamin Kilian[1,2*], Tyson Koepke[1,2], Jonathan Abarca[1], Matthew Whiting[1,3],

and Amit Dhingra[1,2§]

[1]Department of Horticulture, Washington State University, Pullman WA 99164-6414

[2]Molecular Plant Sciences Graduate Program, Washington State University, Pullman, WA 99164

[3]Irrigated Agriculture Research and Extension Center, Washington State University, Prosser,

WA 99350

*Authors contributed equally to this work

[§]Corresponding author

**Abstract**

Harvesting of sweet cherry (*Prunus avium* L.) fruit is a labor-intensive process. Mechanical harvesting of sweet cherry fruit is feasible; however, it is dependent on the formation of an abscission zone at the fruit-pedicel junction. The natural propensity for pedicel-fruit abscission zone (PFAZ) formation varies by cultivar, and the general molecular basis for PFAZ formation is not well characterized. In this study, ethylene-inducible change in pedicel fruit retention force (PFRF) was recorded in a developmental time course with a concomitant analysis of the PFAZ transcriptome from three sweet cherry cultivars. In 'Skeena', mean PFRF for both control and treatment fruit dropped below the 0.40kg-force (3.92N) threshold for mechanical harvesting and indicating the formation of a discrete PFAZ. In 'Bing', mean PFRF for both control and treatment groups decreased over time. However, a mean PFRF conducive to mechanical harvesting was achieved only in the ethylene-treated fruit. While in 'Chelan' the mean PFRF of the control and treatment groups did not meet the threshold required for efficient mechanical harvesting. Transcriptome analysis of the PFAZ followed by the functional annotation, differential expression analysis, and gene ontology (GO) enrichment analyses of the data facilitated the identification of phytohormone-responsive and abscission-related transcripts as well as processes that exhibited differential expression and enrichment in a cultivar-dependent manner over the developmental time-course. Additionally, read alignment-based variant calling revealed several short variants in differentially expressed genes, associated with enriched gene ontologies and associated metabolic processes, lending potential insight into the genetic basis for different abscission responses between the cultivars. These results provide genetic targets for induction or inhibition of PFAZ formation, depending on the desire to harvest the fruit with or without the stem attached. Understanding the genetic mechanisms underlying the development of

39    the PFAZ will inform future cultivar development while laying a foundation for mechanized

40    sweet cherry harvest.

41    **Key Words**

42    Ethephon, ethylene, auxin, RNA-seq, transcriptome, *Prunus avium*, abscission, RT-qPCR, gene

43    expression, Rosaceae, mechanical harvest

**Introduction**

Sweet cherry (*Prunus avium* L.), a member of the Rosaceae family, is a commercially important tree fruit crop throughout the world, with approximately 2.3 million tons produced annually (Axe and Bush, 2017; Blando and Oomah, 2019). In addition to its worldwide economic value, production, and consumption of sweet cherries has increased in recent years as consumers have become aware of their nutritional benefits (Blando and Oomah, 2019). The industry is evolving to address the growing market, availability of labor during harvest, and has begun adapting harvesting strategies to meet new consumer demand for stemless cherries (Kappel et al., 2012; Quero-García et al., 2017). Traditional harvesting methods focus on separating the fruit from the tree at the pedicel-peduncle junction, leaving the pedicel (stem) attached to the fruit (Wittenbach and Bukovac, 1974). Mechanical harvesting, on the other hand, is best achieved when the fruit abscises easily at the fruit-pedicel junction (Quero-García et al., 2017). Increasing labor costs associated with traditional hand harvesting, in addition to the growing demand for stemless fruit, has made adoption of mechanical harvesting strategies attractive if they can be made uniform across cultivars (Zhao et al., 2013). However, the harvesting of sweet cherries has presented a unique set of challenges. Unlike sour cherries (*Prunus cerasus*), which develop an anatomically and histologically distinct fruit-pedicel abscission zone (PFAZ) and separate with ease, sweet cherry cultivars display phenotypic differences in PFAZ formation and consequent ease of fruit separation—some cultivars require excessive force to separate fruit at the PFAZ, which tends to compromise fruit quality and integrity (Stösser et al., 1969; Zhao et al., 2013).

While sweet cherry and peach belong to the same sub-family, the former bears non-climacteric fruits that do not produce ethylene autocatalytically during ripening and senescence.

This is most likely due to the presence of several stop codon mutations in the ethylene

biosynthesis and perception genes in sweet cherry (Koepke et al., 2013). However, sweet cherry

fruit from some cultivars exhibit a novel developmental response to exogenous ethylene

application (Hiwasa-Tanase and Ezura, 2014).  Exogenous ethylene application can induce or

enhance PFAZ formation, loosening the fruit, and facilitating efficient mechanical harvesting

(Wittenbach and Bukovac, 1974; Smith and Whiting, 2010).

Pedicel-fruit retention force (PFRF) is used as a direct measure of PFAZ formation and

serves as a metric to determine the mechanical harvestability of the fruit. An average PFRF value

of 0.40kg-force (3.92N) is considered the threshold for mechanical harvestability, though this

will depend on the actuation method of the harvester (Bukovac and MJ, 1979; Smith and

Whiting, 2010; Zhao et al., 2013). In sweet cherry, a reduction in PFRF can be induced through

the application of ethephon (2-chloroethylephosphonic acid), a commercially available plant

growth regulator that is rapidly metabolized to ethylene (Smith and Whiting, 2010; Zhao et al.,

2013). In both sweet and sour cherry, ethephon has been demonstrated to be a viable option for

reducing the threshold for mechanical harvest without negatively impacting fruit quality, when

applied in appropriate concentrations; however high rates of ethephon have been associated with

reduction in fruit quality (e.g. gummosis, defoliation, lenticel enlargement) and, thus, cultivars

with naturally low PFRF are desired (Looney and McMechan, 1970; Bukovac and MJ, 1979;

Smith and Whiting, 2010; Zhao et al., 2013). The natural PFRF of sweet cherry varies by

cultivar, with some varieties, like 'Skeena', exhibiting an auto-abscising phenotype and requiring

no exogenous ethylene to induce abscission. Representing an intermediate phenotype, 'Bing' can

be induced to abscise when ethephon is applied approximately 14 days before harvest (Smith and

Whiting, 2010; Zhao et al., 2013). 'Chelan', on the contrary, does not abscise naturally or in the

presence of ethephon (Smith and Whiting, 2010).

Sweet cherry phenotyping studies have shown that PFRF values remain consistent for

these cultivars across multiple years, indicating that the abscission phenotypes are genetically

stable and can perhaps be manipulated at the genetic level (Zhao et al., 2013). Furthermore, these

findings suggest that standardization of PFRF for mechanical harvesting across cultivars is

possible if the ideal ethephon or other treatment regimens are determined for individual sweet

cherry cultivars. Despite the extensive physiological characterization of PFAZ integrity across

sweet cherry varieties, the underlying molecular basis for PFAZ structural differences has not

previously been elucidated.

As understood from studies in model plant systems such as Arabidopsis and tomato,

abscission at the fruit-pedicel junction entails a series of hallmark structural changes: the middle

lamella is dissolved by hydrolytic enzymes, such as polygalacturonase and cellulase (Taylor et

al., 1991); cell walls in the separation layer thicken, and cell wall components become hydrated

as a result (Huberman et al., 1983); primary cell walls break down as abscission progresses,

resulting in the formation of large intercellular cavities (Tabuchi et al., 2001; Grover, 2012); and

lignin deposits accumulate proximally to the abscission zone, forming part of a peridermal

boundary layer that will serve to protect the pedicel scar following fruit separation (Sexton and

Roberts, 1982; Tabuchi et al., 2001; Merelo et al., 2017; Meir et al., 2019). This process is

thought to operate in a similar manner in fruit crops including apple, peach, and olive although

species and cultivar-specific differences, particularly with regards to chemical induction of

abscission in these crops, are not yet well understood (Botton et al., 2011; Zhu et al., 2011; Ali et

al., 2012; Gil-Amado and Gomez-Jimenez, 2013; Goldental-Cohen et al., 2017).

According to the current model of abscission in plants, the genetic events underlying

cellular structure modification at the PFAZ center around an interplay between ethylene and

auxin-associated pathways. Binding of ethylene to corresponding receptors (ETRs) in PFAZ

cells initiates signal transduction pathways leading to the activation of numerous, ethylene-

responsive transcription factors (ERFs), which elicit different modulatory roles. The ethylene

response and signaling network ultimately results in the initiation of cell death, reduction in cell

wall adhesion, and separation of the fruit from the pedicel (Roberts et al., 2002; Meir et al.,

2019). Working antagonistically to ethylene is the phytohormone auxin. Auxin's biologically

active form, free indole-3-acetic acid (IAA), decreases the sensitivity of plant organs to ethylene.

The genetic and metabolic factors governing auxin homeostasis ensure that an appropriate

balance is maintained between free IAA, IAA-conjugates, and auxin degradation during different

developmental stages (Meir et al., 2010, 2015). A presence of higher levels of free IAA

corresponds to inhibited or delayed ethylene-dependent developmental responses like abscission

zone formation (Else et al., 2004). Decreasing polar auxin transport across the abscission zone in

sweet cherries by girdling methods results in increased fruit abscission (Blanusa et al., 2005).

Additionally, in grape, the application of inhibitors of auxin transport has been shown to promote

increased abscission (Kühn et al., 2016).

While there is evidence implicating the involvement of numerous ethylene-associated

transcription factors and auxin-associated genes in the modulation of PFAZ formation, species-

specific modes of action have yet to be resolved (Roberts et al., 2002). Furthermore, mechanisms

involving transmission of hormonal and other signals upstream of AZ formation, as well as

specifics of the enzyme-driven remodeling and redeposition of cell wall constituents in sweet

cherry are in need of further elucidation. The stimulation of PFAZ formation in response to

exogenous ethylene application in the non-climacteric fruit sweet cherry represents a unique

biological system to elucidate the process of inducible abscission. An improved understanding of

the interplay between hormone response, signaling, and activation of abscission-associated genes

and pathways in sweet cherry will facilitate improved strategies for planned induction or

inhibition of PFAZ formation.

To elucidate the molecular bases for differences in abscission phenotypes among sweet

cherry cultivars, and to correlate this information with fruit development, time course

physiological measurements of the PFRF along with concomitant transcriptome analysis of the

PFAZ tissue from ethylene-treated and control 'Bing', 'Skeena', and 'Chelan' were conducted.

The hypothesis that cultivar-specific gene expression differences in ethylene- and auxin-

responsive pathways are directly correlated to the differences in abscission phenotypes was

evaluated. Additionally, this work aimed to reveal other genes and enriched gene ontologies

associated with key metabolic processes involved in pedicel-fruit abscission in sweet cherry. The

results of this study provide potential genetic targets for PFAZ formation in sweet cherry, which

are expected to inform strategies for improving PFAZ phenotypes conducive to different

harvesting approaches.

**Methods**

*Plant Material*

The sweet cherry trees used in this study are located at Washington State University's

Roza Farm, 10 km north of Prosser, Washington, USA (46.2°N, 119.7°). Trees were irrigated

weekly from bloom to leaf senescence with low-volume, under-tree, micro-sprinklers, and grown

using standard orchard management practices. Trees had an in-row spacing of 2.44 m (8 ft) and

between row spacing of 4.27 m (14 ft). Rows were planted in a north-south orientation and

trained to a Y-trellis architecture.

*Ethephon application*

Ethephon (formula 240 g/l [2lbs/gal]) was applied via air-blast sprayer at 3.5 L ha−1 (3 pt

A−1) with a 1,871 L ha−1 (200 g A−1) spray volume (Smith and Whiting, 2010). Ethephon

applications and measurements were conducted in three different years (2010, 2013, and 2014).

Each replication was performed in the same orchard block, using distinct trees within the block.

Treatment application was done early in the morning (between 0600 and 0800 hours) to reduce

the effects of ethylene evolution from warm temperatures and wind, as previously described

(Smith and Whiting, 2010).

Optimal ethephon application time for 'Bing' had been established previously as 14 days

before harvest (DBH) (Smith and Whiting, 2010), or 80% fruit maturation. Because 'Bing',

'Chelan', and 'Skeena' have different timelines for the maturation of fruit after bloom, ethephon

was applied at ca. 80% maturation for each of the cultivars in the 2014 growing season. This

percentage coincided with 14 DBH for 'Bing', 12 DBH for 'Chelan' and 16 DBH for 'Skeena'

(Supplementary File 1). Information regarding ethephon treatment, and PFRF results for 2010

and 2013 can be found in Supplementary File 2.

*PFRF measurements, color evaluation, and abscission zone sampling*

In all three years, sampling and measurements were conducted at the following time

points for each sweet cherry cultivar: (1) immediately prior to application of ethephon or $H_2O$;

(2) 6 hours after the application of ethephon or $H_2O$, and (3) at harvest.

At each sampling time, ten fruit were randomly selected for analysis from each of four

trees/cultivar/treatment. PFRF was measured using a modified digital force gauge (Imada). Each

fruit was manually categorized by exocarp color based on 1–7 scale developed specifically for

sweet cherry by CTIFL (Centre technique interprofessionnel des fruit et legumes, France)

(Supplementary File 3) (Zhang and Whiting, 2011).  In addition to the collection of PFRF values,

the abscission zones of 10 fruit, representing ten biological replicates, from each

cultivar/treatment were harvested from corresponding trees at each time point per the following

steps: 1.) Using a new razor blade, the fruit was first cut approximately 0.5 cm below the pedicel,

leaving the pedicel and a thin disc of fruit/skin attached, 2.) two sets of parallel cuts were made

downward on the cherry fruit disc on either side of the stem, effectively making a cubic piece of

fruit 3mm x 3mm x 3mm attached to the pedicel, 3.) the pedicel was cut off directly above the

fruit and the cube of fruit tissue consisting the abscission zone and some fruit and pedicel tissue

was placed in a 15ml falcon tube and flash frozen for subsequent processing (Supplementary File

4).

*Total RNA Extraction*

Excised sweet cherry abscission zone tissue derived from ten fruits representing ten

biological replicates / cultivar / time point was pooled, pulverized and homogenized into a single

sample using a SPEX SamplePrep® FreezerMill 6870 (Metuchen, NJ USA) and transferred to

storage at -80°C. Total RNA was extracted using an acid guanidinium thiocyanate phenol

chloroform extraction method similar to that previously described (Chomczynski and Sacchi,

1987). Briefly, 1mL of 0.8M guanidinium thiocyanate, 0.4M ammonium thiocyanate, 0.1M

sodium acetate pH 5.0, 5% w/v glycerol, and 38% v/v water saturated phenol were added to

approximately 100 mg powdered tissue, shaken to evenly mix the sample and incubated at room

temperature (RT) for 5 minutes. 200μL chloroform was added and shaken vigorously until the

entire sample became homogenously cloudy and then was incubated (RT, 3 minutes). Samples

were centrifuged at 17,000 x g at 4°C for 15 minutes, and the aqueous upper phase was

transferred to a sterile 1.5mL microcentrifuge tube. To this, 600µl 2-propanol was added,

inverted 5-6 times, and incubated at RT for 10 minutes. Samples were centrifuged 17,000 x g at

4°C for 10 minutes, and the supernatant decanted. 1 mL 75% DEPC ethanol was added to the

pellet, vortexed for 10 seconds, and centrifuged 9,500 x g at 4°C for 5 minutes. Pellets were

suspended in RNase free water and incubated at 37°C with RNase free *DNaseI* for 30 minutes,

which was subsequently inactivated (65°C, 10 minutes).

Extracted RNA was quantified, and its quality was checked using the Bio-Rad (Hercules,

CA) Experion system using the Experion RNA High Sensitivity Analysis kit or the Agilent

(Santa Clara, CA) 2100 Bioanalyzer system using the RNA NanoChip and Plant RNA Nano

Assay Class.

*RNA sequencing and assembly*

RNA samples that passed the quality threshold of RIN 8.0 were sequenced at the

Michigan State University Genomics Service Center for library preparation and sequencing. The

Illumina Hi Seq 2000 sequencing platform (San Diego, CA) was used to sequence 2x100 PE

reads from the cDNA libraries generated from the above RNA extractions, representing a single

sample derived from 10 biological replicates of each cultivar, treatment, and time point. cDNA

and final sequencing library molecules were generated with Illumina's TruSeq RNA Sample

Preparation v2 kit (San Diego, CA) and instructions with minor modifications. Modifications to

the published protocol include a decrease in the mRNA fragmentation incubation time from 8

minutes to 30 seconds to create the final library of proper molecule size range. Additionally,

Aline Biosciences' (Woburn, MA) DNA SizeSelector-I bead-based size selection system was

utilized to target final library molecules for mean size of 450 base pairs. All libraries were

quantified on a Life Technologies (Carlsbad, CA) Qubit Fluorometer and qualified on an Agilent

(Santa Clara, CA) 2100 Bioanalyzer.

Read preprocessing and assembly were conducted in CLC Genomics Workbench (8.5.1).

Briefly, RNAseq read datasets were processed with the CLC Create Sequencing QC Report tool

to assess read quality. The CLC Trim Sequence process was used to trim the first 16 bases due to

GC ratio variability and for a Phred score of 30. All read datasets were trimmed of ambiguous

bases. Illumina reads were then processed through the CLC Merge Overlapping Pairs tool, and

all reads were *de novo* assembled to produce contiguous sequences (contigs). A single master

assembly was generated from the combined read data from 'Bing', 'Chelan', and 'Skeena'

cultivars at each time point (Supplementary File 5). Assembled contigs passed the filter criteria

of >200 base length combined with >2x average read coverage. The cultivar-specific, non-

trimmed read sets were mapped back to the master assembly to generate individual read

mappings for each cultivar, treatment, and time point. Read counts were normalized for each

mapping group using the Reads Per Kilobase per Million reads (RPKM) method (Mortazavi et

al. 2008). All the sequence data is available from NCBI Short Read Archive – Bing accession:

SRX2210365; Chelan accession: SRX2210366; Skeena accession: SRX2210367 submitted

under BioProject PRJNA329134.

*Differential expression analysis*

A two-pronged approach, using Kal's Z-test and the NOIseq-sim R package, was

employed to identify contigs with highest likelihood of being differentially expressed (Kal et al.,

1999; Tarazona et al., 2013). Only contigs that passed established threshold filtering for both

methods were considered for further analysis.

Genes with highest probability of being differentially expressed were first identified

using the NOISeq-sim package in OmicsBox, which is designed to infer probability of

differential expression by modeling biological replications in silico for RNAseq experiments

(Tarazona et al., 2011, 2013). This approach has been successfully employed for differential

expression analysis in other crops, including peach and rice (De La Fuente et al., 2015; Altúzar-

Molina et al., 2020). Default parameters were used to simulate five, in silico replications with a

simulated replicate size of 0.2 and a set variability of 0.02 in each replication.

Next, Kal's Z-tests were performed in CLC Genomics Workbench 8.5.1 to add another

level of stringency to the identification of differentially expressed genes. A paired experiment

comparing the read count values for ethephon treatment to control values corresponding to each

cultivar was performed.

The final sequence selection was reduced to 3,190 contigs, which were considered to

have a high probability of being differentially expressed based on conformity to all of the

following criteria for at least one treatment/time: (1) $|log2FC|>1$, (2) NOIseq probability of DE >

0.8, (3) Kal's Z-test FDR corrected p-value <0.05 (Supplementary File 6).

*Functional annotation and GO enrichment analysis*

Assembled sweet cherry contigs were annotated using the Blast2GO feature in OmicsBox

(version 1.2.4). Contigs were blasted for greatest sequence homology against the NCBI

Viridiplantae database and subsequently assigned to their corresponding gene ontology (GO)

terms as described previously (Götz et al., 2008; Hewitt et al., 2020a, 2020b).

GO enrichment analysis using Fisher's Exact Test was also conducted in OmicsBox to

identify cellular components, molecular functions, and biological processes that were over or

under-represented in the ethephon treated fruit at harvest in comparison with the control fruit

(Götz et al., 2008). Based on the differential expression analysis, for each sweet cherry cultivar,

lists representing transcripts with NOIseq probability>0.8, Kal's Z-test FDR corrected p-

value<0.05, and |logFC|>1 at the harvest time point were produced. These lists served as the

treatment datasets for enrichment analyses, and the master annotated transcriptome was used as

the reference dataset (Supplementary File 7). The FDR corrected p-value cutoff for enrichment

was set to 0.05. Following separate enrichment analyses for each cultivar, enriched GO terms

that were shared between cultivars or unique to a single cultivar were identified using the Venn

Diagram application in OmicsBox.

*RT-qPCR Validation*

Targets for RT-qPCR validation were selected from a list of genes known to be involved

in ethylene response and cell wall breakdown (Supplementary File 8). Primers were designed

based on the near full-length transcript sequences to amplify an approximately 100-150 bp

region in the 3' region of target transcripts. A bacterial luciferase gene was used as a spiked

reference, with 50 ng added per reaction.

Library preparation, target amplification, and expression analysis were conducted in

accordance with previously published methods, with minor modifications (Hendrickson et al.

2019). VILO cDNA synthesis kit (Invitrogen™) was used to generate three biological replicates

of cDNA from three independent RNA samples isolated for each time point per manufacturer's

instructions. Replicate cDNAs were then pooled into a single sample (50 ng/ul). At least 4-6

technical replicate RT-qPCR reactions were performed using iTAQ with ROX and SYBR

(BioRad), and 20μL reactions were prepared as per the manufacturer recommendations. A total

of 2μL of cDNA diluted to 50ng/μL RNA equivalents was used per reaction with 5μL $H_2O$, 2μL

of each primer (10μM), and 10μL of iTAQ SYBR® Green Supermix with ROX. The RT-qPCR

reactions were performed on a Stratagene MX3005 using the following parameters: 95°C 5 min;

50 cycles of 95°C 30 sec, 57°C 30 sec, 72°C 30 sec; 72°C 5 min. Fluorescence readings were

taken at the end of each elongation step. A melting step was performed following the cycles at

95°C for 30 sec, 54°C for 30 sec and ramp up to 95°C to produce a dissociation curve.

To capture PCR efficiency in the data, Cq values and efficiencies were calculated for

each reaction using the LinRegPCR tool (Ramakers et al. 2003, Ruijter et al. 2009). Cq values

resulting from efficiencies below 1.80 or above 2.20 were judged unacceptable and were treated

as unsuccessful or undetected amplifications. Cq values with efficiency values that were within

expected parameters but exceeded (or equaled) 40.00 were also deemed unacceptable and

disregarded in downstream analysis. In the same manner, Cq values between (35.00-39.99) were

determined to be of low confidence and were marked for special consideration in downstream

analysis. Fold-change expression was determined from the Cq values of all gene targets (among

all replicates of all samples) among the 'Bing', 'Chelan', and 'Skeena' cultivars using the Pfaffl

method (Pfaffl 2001). Expression values were determined with reference to the luciferase spiked

gene (Supplementary File 8).

*Short variant identification*

The GATK best practices pipeline for short variant discovery was used to identify SNPs

and indels in key, differentially expressed ethylene- and auxin-associated contigs, with minor

modifications (Van der Auwera et al., 2013; Poplin et al., 2018). Briefly, a group of paired,

untrimmed reads from each sweet cherry cultivar was aligned to a designated reference fasta in

CLC Genomics Workbench 8.5.1. The reference file contained only the sequences for the

assembled contigs that had been previously assigned GO annotations in OmicsBox. The resulting

three alignments were exported as BAM files for subsequent use in the GATK pipeline. A

reference fasta index and dictionary were created using Samtools and Picard software programs,

respectively. Within the GATK (v. 4.1.7.0) suite, the HaplotypeCaller tool was used to identify

variants between parental haplotypes for each cultivar; the results from all three cultivars were

then merged into a single GVCF file using the CombineGVCFs tool. Finally, The

GenotypeGVCFs tool was used to perform joint genotyping on the GVCF file containing variant

information for each cherry cultivar. The SelectVariants tool was used to filter out low-quality

and low-confidence calls; minimum depth of coverage (DP) was set at 30x and minimum quality

of depth (QD), which is directly related to Phred score, was set to 30. The results were visualized

and called variants in differentially expressed genes of interest were confirmed, using Integrative

Genomics Viewer (v. 2.8.2).

**Results and Discussion**

*Pedicel-Fruit Retention Force*

Application of ethephon at 80% of fruit development (Smith and Whiting, 2010) ensured

developmental equivalency of PFRF and tissue sampling time points across cultivars. This

percentage coincided with 12 DBH for 'Chelan', 14 DBH 'Bing', and 16 DBH for 'Skeena'. The

PFRF values of control fruits decreased naturally over time, with reductions of 75.5%, 74.0%,

and 37.5% observed for 'Skeena', 'Bing', and 'Chelan,' respectively. Application of ethephon

decreased mean PFRF value in comparison with the respective controls, but whether this

decrease was biologically significant and resulted in the achievement of the threshold for

mechanical harvesting varied across cultivars (Figure 1).

In 'Skeena', mean PFRF for both control and treatment fruit dropped below the 0.40kg-

force (3.92N) threshold for mechanical harvesting, reaching final mean PFRF values of 0.29kg-

force (2.84N) and 0.156kg-force (1.53N), respectively (Figure 1). These findings indicate that

while 'Skeena' is capable of natural abscission, ethephon application does significantly increase

the AZ formation response, causing PFRF values to decrease significantly compared to control

fruit by the harvest date.

Similarly, for 'Bing', mean PFRF for both control and treatment groups decreased over

time; however, a mean PFRF conducive to mechanical harvesting was only achieved in the

ethephon treated fruit, which reached a final PFRF value of 0.215kg-force (2.11N), significantly

lower than the control value of 0.418kg-force (4.1N) (Figure 1). This suggests that the

inducibility of 'Bing' is resultant of a similar, yet less dramatic, natural decrease in PFRF that is

enhanced by ethephon treatment.

'Chelan' exhibited a statistically significant PFRF response at the time of harvest in the

ethephon treated fruit in comparison with the control; however, the final PFRF of 0.832kg-force

(8.16N) of the treatment group was not reduced to the threshold required for efficient mechanical

harvesting (Figure 1). These physiological results support the observations that 'Chelan' forms

neither a developmental nor an ethylene-induced PFAZ (Smith and Whiting 2010). In the

absence of a discrete PFAZ, the pedicel fruit junction region in 'Chelan' corresponding to the

PFAZ in 'Bing' and 'Skeena' was used for subsequent RNAseq analysis.

*Transcriptome assembly and annotation*

The transcriptome assembly resulted in the generation of 82,587 contigs from

1,061,563,488 total trimmed reads. Contigs were subsequently filtered for >200 base length and

>2x coverage, for a final total of 81,852 contigs for downstream processing (Supplementary File

5). Functional annotation conducted using the OmicsBox genomics suite resulted in the

assignation of annotations to a total of 30,946 (37.8%) contigs (Supplementary File 7).

*Gene Ontology Enrichment Analysis of Differentially Expressed Contigs*

Based on our NOIseq/Kal's test approach, 1,274 genes were identified to be differentially

expressed in ethephon-treated 'Bing', 715 in 'Skeena', and 523 in 'Chelan' at least one

treatment/timepoint in comparison with the control. To understand what biological processes,

molecular functions, and cellular components were enriched across the three cultivars, and

thereby determine pathways of the greatest interest to this study, GO enrichment analysis of

differentially expressed contigs was conducted, followed by filtering for the most specific

ontologies using an FDR-corrected p-value cutoff of 0.05. This strategy resulted in identification

of enriched GOs at 6-hours post-ethephon and at harvest for all genotypes. No GOs were

significantly enriched at the 0-hour time point.

<u>Unique, Enriched Gene Ontologies at 6 Hours Post-Ethephon Treatment</u>

From samples collected 6 hours after treatment, 84 unique GOs were identified for

'Bing', 8 for 'Chelan', and 33 for 'Skeena' (Figures 2A and 3A). Among the ontologies unique

to 'Bing' were several terms associated with chitin breakdown and metabolism, including:

"chitinase activity", "chitin binding", "chitin catabolic process", and "response to chitin".

Besides the hallmark role of plant chitinases and associated pathways in breakdown of fungal

pathogens, these enzymes are involved in a number of other important aspects of development,

such as generation of signal molecules that regulate organ morphogenesis (van Loon et al.,

2006), regulation of lignin deposition and cell shape (Zhong et al., 2002; Johnston, 2006; Agustí

et al., 2008), regulation of programmed cell death (Grover, 2012), and formation of abscission

zones (Roberts et al., 2000; Grover, 2012). Chitin-associated processes have also been shown to

be activated in response to jasmonic acid, methyl jasmonate, ethylene, and gibberellic acid

(Graham and Sticklen, 1994; Kasprzewska, 2003; van Loon et al., 2006), as well as auxin and

cytokinin (Shinshi et al., 1987). The unique enrichment of chitin-associated GO terms in 'Bing'

was accompanied by simultaneous overrepresentation of several hormone signaling pathways,

including "methyl jasmonate esterase activity", "methyl salicylate esterase activity", "jasmonic

acid catabolic process", "salicylic acid metabolic process", "gibberellin catabolic process", and

"hormone-mediated signaling process", the latter of which included genes associated with

ethylene, auxin, and abscisic acid (ABA) signaling and response. As jasmonic acid (JA) and

salicylic acid (SA) metabolic pathways have established roles in plant growth regulation and

stress response, the unique co-enrichment of all of these terms, in conjunction with the "ROS

response" enriched term, suggest that the inducible abscission phenotype may result from the

unique jasmonic acid and redox/ROS signaling pathways, which are known to be activated in

response to ethylene response (Müller and Munné-Bosch, 2015). These pathways, in turn, may

lead to activation of chitin-associated processes that lead to changes in lignin deposition, cell

growth, and programmed cell death in a cascade of events leading up to abscission zone

formation following ethephon application in 'Bing'.

In 'Chelan' a relatively small number of GO terms were enriched at 6 hours post-

ethephon. Among the few enriched terms was "negative regulation of endopeptidase activity".

As with plant chitinases, endopeptidases contribute to programmed cell death and cell wall

breakdown (Helm et al., 2008) and have been shown to be activated by exogenous ethylene

application (LIU et al., 2005). Enrichment of this ontology suggests that molecular mechanisms

are at play to stabilize the ethylene response in 'Chelan', whereas such a mechanism does not

appear to play a substantial role in 'Bing'.

'Skeena' displayed a number of enriched ontologies associated with epigenetic

modifications, including "chromatin binding", "double-stranded methylated DNA binding",

"regulation of DNA methylation", "gene silencing by RNA-directed DNA methylation",

"histone acetylation", and "histone binding". Such processes have previously been implicated in

abscission of citrus (Sabbione et al., 2019) and litchi (Peng et al., 2017). In general, methylation

of cell wall structural components, such as pectin, is thought to be critical for maintenance of cell

wall integrity (Zhang and Zhang, 2020). Enrichment for such processes, which may occur in an

ethylene-independent manner, could underlie the natural abscission process in 'Skeena'.

<u>Unique, Enriched GOs at Harvest</u>

At harvest, 22 unique enriched ontologies were identified in 'Bing', 56 in 'Chelan', and 2

in 'Skeena' (Figures 2B and 3B). Among the ontologies unique to 'Bing' were "lignin catabolic

process" and "plant organ morphogenesis", along with several GO terms that were also enriched

at the 6-hour time point— chitin-metabolism and response-associated terms and JA/SA-

associated processes. Lignin, a derivative of phenylpropanoid metabolism, is abundant in

abscission zones, aiding in the development of a protective boundary layer (Sexton and Roberts,

1982), serving as a mechanical brace to localize cell wall breakdown (Lee et al., 2018), and

accumulating specifically in the PFAZ regions during ethylene-promoted abscission (Merelo et

al., 2017). In addition to the instrumental role of lignin, general processes involved in "plant

organ morphogenesis" appear to be at work in inducible abscission of 'Bing'.

In 'Chelan', a comparatively high number of terms was observed at harvest. Among the

enriched GOs were "cell wall-related processes" and "methylation" (Figure 2B, Supplementary

File 9).  These abscission-associated terms are enriched much later than they are for 'Bing' and

'Skeena', both of which showed enrichment of associated processes at the 6-hour time point.

Furthermore, in contrast to the 6-hour time point, at which few GOs were enriched, the elevated

number of enriched terms observed in 'Chelan' at harvest suggest a delayed developmental

response to ethylene, which may contribute to the failure of this cultivar to develop a PFAZ

naturally and in the presence of ethephon.

In 'Skeena', at harvest, only two GO terms were uniquely enriched: "phenylpropanoid

biosynthetic process" and "stress response". The limited number of unique terms at the later time

point indicate that the majority of impactful processes contributing to PFAZ formation likely

take place earlier in development. Among the enriched ontologies, phenylpropanoid metabolism

is involved in production of lignin precursors and has been previously implicated in fruit

abscission (Kostenyuk et al., 2002). Additionally, stress-associated processes appear to be active

at this stage of development and are likely to further accentuate processes associated with

programmed cell death, changes in cell wall integrity, and ultimately PFAZ formation in

'Skeena'.

Shared, Enriched Gene Ontologies

In addition to unique GOs, shared ontologies reveal PFAZ formation-associated

processes that are similar between two or more cultivars following ethephon treatment (Figures

2A, 2B, and 4). At harvest, 'Bing' and 'Chelan' shared the enriched GO term "response to

auxin", while 'Skeena' did not. One possible contributing factor to the auto-abscission of

'Skeena' is that this cultivar may have naturally lower free auxin levels than 'Bing' or 'Chelan'

and, promoting natural formation of the PFAZ. This concept is explored further in a later section.

At the 6-hour time point, all three genotypes shared the enriched GO term, "DNA-

binding transcription factor activity". This GO term, among which a number of ethylene-

responsive transcription factors were identified (Figure 4), remained enriched in 'Bing' and

'Chelan' at harvest. This observation provides evidence that all three cultivars have ethylene-

responsive capacity; however, the previously described negative regulation of ethylene-

associated mechanisms in 'Chelan', along with the enrichment of auxin response at harvest,

suggests that a higher degree of feedback inhibition serves to quell the extent of ethylene-

signaling and response that culminates in strong PFAZ formation in the other two cultivars

(naturally, in 'Skeena,' and following ethephon stimulation, in 'Bing').

The term "integral component of membrane" was shared between 'Chelan' and 'Skeena'

at 6 hours post-ethephon and between all three cultivars at harvest. A large number of

differentially expressed genes were associated with this term (Supplementary File 9); among the

notable genes were cell wall and integrity-associated *patatin-like protein 2* (*PLP2*), which plays

a role in programmed cell death, and protein *walls-are-Thin 1* (*WAT1*), which, in addition to its

role in auxin mobilization, plays a role in secondary cell wall formation and stability (Camera et

al., 2009; Ranocha et al., 2010). As all three cultivars undergo a decrease in PFRF following

application of ethephon, it is logical that biological processes, molecular functions, and cellular

components associated with membrane and cell wall integrity are affected (Figure 2).

Overall, the GO enrichment results provide a global picture of the processes underlying

the natural abscission capacity of 'Skeena', the inducibility-of the abscission process in 'Bing',

and recalcitrance to PFAZ formation in 'Chelan'. Examination of the differentially expressed

genes associated with the enriched ontologies, described in the next section, lends further insight

into the mechanisms underlying these three, unique sweet cherry abscission phenotypes.

*Differential Expression of Genes Associated with Key, Enriched Ontologies*

Results of the GO enrichment analysis provided a basis for distinguishing pathways in

which to observe expression patterns of significant genes, and to compare these patterns of

across cultivars following ethephon or control treatments.

478
479    <u>Differential Expression of Genes at 6 Hours Post-Ethephon Treatment</u>
480

In 'Bing', a number of differentially expressed contigs corresponding to endochitinases,

including several transcript variants of *endochitinase EP3-like*, *endochitinase-like*, and

*endochitinase 2-like* were induced at the 6-hour time point (Figure 3A). At the same time,

hormone-mediated signaling genes showed heightened expression, including *abscisic acid*

*receptor PYR1-like* (*PYR1-like*), *ethylene receptor 2* (*ETR2*), *auxin-responsive protein IAA1-like*

(*IAA1-like*), and JA responsive protein*s TIFY5A-like* and *TIFY 9*. While, genes associated with

chitinase activity, hormone signaling, and ROS response displayed consistently elevated fold-

change expression in 'Bing', in the other two cultivars, the responses varied.  In general, a trend

of low or negative fold change expression in ethephon-treated versus control PFAZ tissues was

observed for this gene set in 'Chelan' and 'Skeena'; notably, however, *PYR1-like* and *IAA1-like*

fold change expression was elevated in both 'Bing' and 'Chelan'. High expression of ABA-

associated *PYR1-like,* which promotes senescence and stress responses leading up to abscission,

in the PFAZ of ethephon-treated 'Bing' and 'Chelan' cherries suggests that, despite ultimately

different abscission phenotypes, both cultivars possess upstream mechanics that precede PFAZ

formation. Furthermore, the low abscission capacity of 'Chelan' may lie in a comparatively high

auxin versus ethylene response, along with negative regulatory mechanisms that have been

previously undescribed. Specifically, several transcripts corresponding to *alpha-humulene 10-*

*hydroxylase-like*, associated with the GO term "negative regulation of endopeptidase activity",

were consistently downregulated in all three cultivars following ethephon treatment, but most

drastically so in 'Chelan'. The specific role of alpha-humulene hydroxylase-like proteins, which

modulate plant secondary metabolites (mainly sesquiterpenes) in response to herbivory, in

abscission remains to be elucidated; however, the unique expression pattern in 'Chelan' points

towards a potentially novel mechanism of negative regulation of abscission zone formation.

Many genes associated with epigenetic modification were differentially expressed in

'Skeena' at the 6-hour time point, including three transcript variants corresponding to *increased*

*DNA methylation 1* and two transcript variants corresponding to *histone acetyltransferase HAC1-*

*like isoform X1* (*HAC1-like*) (Figure 3A). These genes were also induced in 'Bing' at the 6-hour

time point, but they were not significantly differentially expressed like in 'Skeena'. This finding

suggests that methylation plays an important role in abscission, and that methylation and

associated processes may occur via both ethylene-dependent ('Bing') and independent

('Skeena') pathways.

Differential Expression of Genes at Harvest

Among the genes differentially expressed in 'Bing' at the harvest time point was "lignin

catabolism"-associated *laccase 15-like, laccase 7-like, and laccase 9* (Figure 3B). The former,

laccase genes, are involved in lignin synthesis and deposition; these processes are known to

occur during AZ formation, and are critical for lignin-promoted architectural changes at the

cellular level (Agustí et al., 2008; Merelo et al., 2017). The heightened expression of these genes

observed uniquely in 'Bing' at harvest suggests that laccases, and therefore structural

modifications associated with lignin accumulation, are active in the abscission process for this

cultivar.

Also differentially expressed at harvest in 'Bing' was "plant organ morphogenesis"-

associated *lateral boundary domain containing protein 18* (*LOB18*) (Figure 3B). *LOB18* has

previously been implicated in boundary formation during root development. It is possible that

this gene, or additional members of the LOB gene family, plays a similar role in boundary

formation at the PFAZ (Majer and Hochholdinger, 2011; Wang et al., 2013). *LOB8* expression

was high at harvest in both 'Bing' and 'Skeena', but the expression difference between

ethephon-treated and control PFAZ tissues was significant only in the former. In addition to

*LOB18*, a differentially expressed transcript corresponding to receptor-like protein kinase

*HAESA-LIKE 1* (*HSL1*), which is also associated with the "plant organ morphogenesis" ontology

(Figure 3B), was significantly downregulated in both 'Bing' and 'Skeena'. This is consistent

with previous work in Arabidopsis that revealed decreased expression levels of HSL1

immediately prior to abscission; furthermore, loss of  HSL1 function resulted in impeded floral

and leaf abscission (Niederhuth et al., 2013). The differential expression of both *LOB18* and

*HSL1* may underlie the ability to induce abscission in 'Bing' through boundary formation and

promotion of abscission-associated processes.

In addition to the genes that are directly involved in modification of the PFAZ

ultrastructure, elevated expression of several transcript variants of corresponding to *salicylic acid*

*binding protein 2-like*, which is associated with JA and SA metabolism, was observed at the

harvest time point in 'Bing', indicating that JA signaling mechanisms remain active throughout

development in 'Bing' following ethephon treatment. Similarly, expression of JA-induced,

chitinase-associated genes, were also observed at the harvest.

In 'Chelan', several genes associated with cell wall modification, including *beta-*

*galactosidase 8*, *pectinesterase-like,* and two *expansin* genes, displayed significantly heightened

expression at harvest. These genes are involved in cell wall loosening, degradation of linkages

between structural components, and digestion of cell walls and middle lamella in the AZ (Brown,

1997; Cho and Cosgrove, 2000; Roberts et al., 2002; Wu and Burns, 2004). While cell wall

modification genes displayed similar expression patterns in 'Bing' and 'Skeena', the extent of

expression change between treatment and control was not significant for these cultivars.

Expression patterns of several other genes in 'Chelan' indicate that this cultivar may have

reduced lignin biosynthetic capacity in comparison to 'Bing' and 'Skeena'. While the expression

of cell wall modifying associated genes suggests that 'Chelan' is partially competent to develop

an PFAZ, it is possible that this cultivar doesn't possess some of the key molecular equipment to

coordinate all of the associated processes, including lignification. This is evidenced by the

observation of significantly decreased expression in a methylation-associated transcript

corresponding to *caffeic acid 3-O-methyltransferase-like* (*CAOMT*) at harvest; conversely, this

transcript displayed increased abundance in both 'Bing' and 'Skeena' at 6 hours and at harvest.

*CAOMT* lies upstream of the lignin biosynthetic process; decreased expression of this gene has

been shown to lead to reduced lignification in poplar and expression changes in *CAOMT-like*

genes parallels PFAZ formation in citrus (Van Doorsselaere et al., 1995; Jouanin et al., 2000;

Merelo et al., 2017).

The expression patterns of genes corresponding to cell wall modifiers and lignin

biosynthesis precursors that were identified in 'Chelan' lend insight regarding why this non-

abscising cultivar displays reduced PFRF but fails to form a PFAZ that is discrete enough to

facilitate mechanical harvesting; furthermore, low expression of other terminal abscission-

associated genes, like the laccases displaying heightened expression in 'Bing' lend further

credibility to this hypothesis.

Most notable among the differentially expressed genes in 'Skeena', were stress-response

associated *peroxidase 10*, *peroxidase 17-like*, and *16.9 kDa class I heat shock protein 1-like.*

Elevated expression of stress-responsive genes indicates an increased need for ROS scavenging

to prevent oxidative damage while these signal molecules serve the purpose of activating

abscission associated processes. Based on the pronounced decrease of stress-responsive genes in

'Skeena' following ethephon application, and the observation that PFRF of 'Skeena' is reduced

the least out of all the cultivars following treatment, it is plausible that 'Skeena' may only

perceive the exogenous ethylene as environmental stress, whereas 'Bing' and 'Chelan' appear to

host concerted, ethylene-induced signal transduction responses upstream of abscission in

response to ethephon application. In conjunction with increased peroxidase gene expression

reduced expression of pathogen defense-associated transcripts corresponding to *major latex*

*protein* (*MLP-like protein*) *28* and *43* was observed. This is consistent with a study of *MLP-like*

expression patterns in peach abscission zones, wherein it was suggested that *MLP-like* genes play

role in mitigation in defense in early AZ formation, but display reduced expression immediately

prior to abscission and are further downregulated in response to ethylene (Ruperti et al., 2002).

Similar responses were observed for additional plant-defense associated genes, including

*dessication protectant protein Lea14*, *defensin Ec-AMP-D2-like*, in both 'Skeena' and 'Bing'.

 *Differential Expression of Ethylene-associated Contigs*

In addition to observing the expression patterns of differentially expressed contigs

associated with key enriched metabolic pathways for each cultivar, it was of interest to

specifically examine expression of genes involved in ethylene response, to better understand how

these responses may vary at the cultivar level, and thereby differentially contribute to abscission

phenotypes. The basis for the inducibility of PFAZ formation in 'Bing' during fruit development

is further evident at the genetic level with respect to expression of ethylene biosynthesis,

signaling, and responsive elements in the transcriptome (Supplementary File 9). Throughout the

time course in 'Bing', a transcript corresponding to the rate-limiting ethylene biosynthetic gene

*1-aminocyclopropane-1-carboxylate oxidase 1 (ACO1)* was highly elevated in expression, as

were transcripts corresponding to *ethylene receptor 2* (*ETR2*) and *ethylene insensitive 2* (*EIN2*),

two key genes involved in ethylene perception and signaling, respectively. The activation of

ethylene biosynthesis that is observed to occur uniquely in 'Bing' suggests that exogenous

ethephon stimulates the autocatalytic production of endogenous ethylene, which thereby

accentuates the overall degree and impact of the ethylene response in this cultivar. Moreover, the

naturally high expression of *ETR2* and *EIN2*, which are augmented in the presence of exogenous

ethephon lend further testament to the responsiveness of 'Bing' to ethylene.  A negative

regulator of ethylene, *reversion-to-ethylene-sensitivity 1* (*RTE1*) was induced in 'Bing' following

ethephon application, but levels remained lower overall than in 'Chelan' and 'Skeena', an

observation which is consistent with the comparatively high degree of sensitivity of 'Bing' to

ethylene. In contrast to 'Bing', 'Skeena' and 'Chelan' displayed basal levels of expression of

*ACO1* and *ETR2*, and elevated levels of *RTE1*, suggesting minimal ethylene response,

endogenous production, and sensitivity (Figure 5, Supplementary File 9).

Transmission of ethylene-activated signal to the nucleus involves ethylene-insensitive 2

(EIN2) and ethylene insensitive 2-like (EIL) family of proteins. Upon activation of ETRs by

ethylene, the C-terminal domain of EIN2 is cleaved and translocated to the nucleus where it

activates transcription of ethylene-responsive factors (ERFs) involved in the regulation of

downstream ripening responses (Liu et al., 2015; Chen et al., 2018). Numerous transcripts

encoding ERFs were identified, which were differentially expressed in at least one cultivar and

time point (Figure 4). Consistent with the aforementioned results, a majority of these contigs

were highly induced in 'Bing' following ethephon treatment.    In particular, transcripts

corresponding to *ERF1B-like*, *ERF27* and *ERF109-like, ERF113-like*, *ERF109-like,* and *ERF*

*ABR1-like* were highly differentially expressed in 'Bing' 6 hours following ethephon treatment,

and also at harvest.   In all three cultivars *ERF96-like* was highly induced following ethephon

treatment (Figure 4, Supplementary File 9). In addition to ERFs, transcripts corresponding to

members of the of the WRKY and MYB transcription factor families displayed highly elevated

expression in 'Bing' following ethephon treatment. This is consistent with previous work

demonstrating the activation of both of these transcription factor families during abscission, and

eliciting actions downstream of initial ethylene response (Corbacho et al., 2013; Li et al., 2015;

Jiang et al., 2017; Zhang et al., 2019).

*Differential Expression of Auxin Response-associated Contigs*

In addition to the differential ethylene responses, the ethephon-treated 'Chelan', 'Bing',

and 'Skeena' PFAZ tissues displayed a differential reduction in the expression of auxin-

responsive genes in comparison to their respective controls. This suggests that the response to

auxin and/or reduction in auxin transport, was inhibited in the presence of exogenously applied

ethylene.

Auxin-responsive transcription factor genes *IAA11*, *IAA13*, and *IAA27*, auxin transport

facilitator *WAT1*, and auxin efflux carrier component 1c displayed comparatively low expression

in 'Skeena', intermediate expression in 'Bing', and higher expression in 'Chelan' control PFAZ

tissues at 6 hours post-ethephon and at harvest. Ethephon treatment attenuated this expression in

PFAZ tissue for all three cultivars, but the same relative expression trend was maintained.

*IAA13*, *IAA27*, and *WAT1* displayed decreased expression at harvest in all ethephon-treated

cultivars in comparison with their respective controls. Interestingly, 'Chelan' displayed a natural

increase in the expression of these transcripts from the 0-hour time point to harvest, which was

inhibited and reversed by the application of ethephon (Figure 6). For both 'Bing' and 'Skeena',

the same transcripts naturally decreased in abundance over time—a decrease that was accelerated

in the presence of ethephon.

While free auxin was not measured in this study, differences in the abundance of auxin-

responsive and auxin mobilization-associated transcripts in the three sweet cherry cultivars over

time lends to extrapolation of information regarding cultivar-specific, endogenous free IAA

concentrations. If free auxin levels are high, abundance and/or activity of auxin-responsive

protein encoding transcripts (ARFs, IAAs) is expected to be higher to accommodate them.

'Skeena' does not require ethylene for abscission; however, ethephon application appeared to

further offset the auxin to ethylene ratios in favor of ethylene to accelerate abscission in this

cultivar. The ability of 'Skeena' to auto-abscise suggests that endogenous accumulation of free

auxin at the site of abscission may be naturally lower in this cultivar than it is for 'Bing' and

'Chelan'. Conversely, the comparatively higher expression of transcripts associated with auxin

response and movement proteins in 'Chelan' is suggestive of a naturally higher level of

endogenous IAA. While the current industrial standard levels of ethephon application were

insufficient to reduce the PFRF of 'Chelan' fruit to the threshold required for mechanical

harvesting, both PFRF and auxin-associated transcript abundance in 'Chelan' did decrease as a

result of ethephon application. Ethephon application rates as high as 5.8 L ha$^{-1}$ remain

insufficient to induce a reduction of 'Chelan' PFRF values to the threshold for mechanical

harvest (Smith and Whiting, 2010). An early study found that a high application rate of 500 ppm

(7.8 L ha$^{-1}$ [6.7 pt A$^{-1}$]), ethephon begins to deleteriously affect some sweet cherry cultivars by

inducing unwanted leaf abscission and terminal shoot necrosis, although it is not reported to

what extent 'Chelan' is impacted by such a concentration (Bukovac and MJ, 1979; Smith and

Whiting, 2010). Considering the present results alongside the insight gained from previous work,

it is possible that mechanical abscission of 'Chelan' could be achieved through a combination of

a slightly higher ethephon application rate and application of auxin inhibitors, the latter of which

could further shift the ethylene/auxin ratio in a manner favorable to abscission while reducing the

need for excessively high and potentially phytotoxic ethephon application rates.

'Bing' PFAZ tissues displayed an abundance of auxin-associated transcripts intermediate

to that of 'Skeena' and 'Chelan'. Following ethephon treatment, the phenotypic observations of

inducible abscission were supported at the gene expression level; with the responses greatly

attenuated to lower than those of 'Skeena' ethephon treated and control PFAZ tissues.

*Proposed Models for PFAZ Formation*

Together, the GO enrichment and differential expression analysis results inform a new

model for abscission (Figure 7). In 'Bing', ethephon elicits an ethylene-response that stimulates

hormonal regulatory signals and metabolic processes, including JA and SA-mediated signaling.

These processes, in turn, activate downstream processes associated with morphological changes

at the PFAZ, including chitin metabolism and lignin biosynthesis and deposition. The

culmination of these events ultimately facilitates mechanical separation of the fruit at the PFAZ.

The components of this process, which appears complete in 'Bing', seem to be partially lacking

in 'Chelan', perhaps due to inherently high auxin levels that limit the ethylene-response capacity

from the beginning. 'Skeena' appears to orchestrate PFAZ formation in via a different and

ethylene-independent mechanism. While this mechanism will require further elucidation, lack of

enrichment for the "auxin response" ontology and reduced expression of the corresponding

auxin-associated genes in 'Skeena', suggests that low levels of free auxin, and therefore minimal

antagonism of a natural abscission process, may underly the auto-abscission phenotype.

*Short Variant Discovery in Differentially Expressed Genes*

Variant calling using the GATK pipeline revealed several SNPs and indels in the ORFs

of DE genes associated with key, enriched GOs (Table 1).

An INDEL was identified in "DNA-binding transcription factor activity"-associated

*AP2/ERF and B3 domain containing protein RAV1*, wherein deletion of a 6 bp segment results in

loss of two amino acid residues without impacting the reading frame. Interestingly, 'Skeena' was

homozygous for this deletion, 'Bing' was heterozygous, and 'Chelan' did not possess the

deletion. The resulting protein changes likely impact the natural propensity of the cultivars to

transduce signals associated with stress and hormone responses upstream of PFAZ formation and

could ultimately underlie differences in abscission phenotype. Additionally, a SNP, resulting in

an amino acid substitution, in the coding sequence of *auxin efflux carrier component 1c*, was

identified in 'Chelan'. This variant could impact the overall auxin responsiveness, which appears

to be greater in 'Chelan' than 'Bing' or 'Skeena', at the genetic level.

Additional variants identified include a SNP in "ROS response"-associated *17.4 kDA*

*class III heat shock protein*, an INDEL in the epigenetic modification-associated gene *polycomb*

*group EMBRYONIC FLOWER 2 isoform x1,* and a SNP in the 3' UTR of the "chitinase

activity"-associated *endochitinase-like* gene. *A* member of the diverse polycomb group (PcG)

family, *EMF2* is involved in epigenetic silencing of developmental repressors, and was

significantly downregulated in 'Skeena' at the 6-hour time point. While further work is needed to

determine the precise nature of these variants and their impact on PFAZ formation, their

presence in DE genes associated with enriched functions following ethephon application makes

them key targets for understanding activation of abscission in sweet cherry cultivars.

*RT-qPCR validation*

RT-qPCR analysis of 10 ethylene-responsive/abscission-related genes resulted in 70%

correspondence of general expression trends for 'Bing' and 'Chelan' and an 80% correspondence

for 'Skeena'. Validated transcripts whose RT-qPCR expression pattern (fold-change calculated

using the $2^{-(\Delta\Delta Ct)}$ method) was consistent with that of the RNAseq-based expression

(ethephon/control RPKM ratio) included genes associated with ethylene biosynthesis (ACS1 and

ACO1), perception (ETR2), and response (ERF1B-like, ERF027-like, and WRKY1), as well as

cell wall breakdown-associated polygalacturonase (PG) (Supplementary File 8).

**Conclusions**

This study investigated the changes in PFRF, gene expression changes, and enriched biological processes underlying PFAZ formation in sweet cherry. The results provide transcriptomic insight regarding the gene expression-level effects of exogenous ethylene application on the abscission phenotype and PFAZ development, which can be utilized by the industry to customize harvest strategies.

Consistent with previous work, PFAZ formation was observed to occur in a cultivar-specific manner, and the abscission phenotype of each cultivar was affected to a different degree by exogenous application of ethylene. Observation of unique, heightened expression of ethylene biosynthesis, perception, signaling, and response genes in 'Bing' at 6 hours following ethephon application and at harvest parallels the decrease in PFRF and may be partially responsible for the inducibility of abscission in this cultivar. GO enrichment analysis revealed key biochemical pathways containing additional differentially expressed genes potentially underlying the unique, abscission responses of each cultivar. Among these enriched pathways were a number of hormone-signaling pathways, ROS signaling, chitin metabolic processes, and lignin biosynthesis, and DNA-binding transcription factor activity (among which were numerous ERFs).

Overall, the results of this study point towards a potential genetic basis for the inducible abscission response in 'Bing', the auto-abscission of 'Skeena', and the recalcitrance to abscise displayed by 'Chelan'. The natural abscission of 'Skeena', which is enhanced by ethephon, may result from naturally lower levels of free auxin, as evidenced by the low abundance of IAA/ARF-associated transcripts in comparison with 'Bing' and 'Chelan'. Furthermore, 'Skeena' exhibited fewer significant gene expression changes than the other cultivars following ethephon application, which corresponded to fewer enriched ontologies at the time of harvest. In 'Chelan',

comparatively high abundance auxin-associated transcripts in control fruit, that was attenuated

following ethephon treatment, may indicate higher levels of endogenous free auxin. The

increased capacity for auxin mobilization, indicated by IAA/ARF expression, antagonizes the

abscission-promoting effects of exogenous ethylene. This observation, along with the PFRF

results, provides insight regarding the recalcitrance of 'Chelan' fruit to abscise at the pedicel-

fruit junction.

Identification of cultivar-specific, differentially expressed genes and enriched GO terms

involved in abscission, as well as ontologies that are shared across cultivars, provides

information that will inform future efforts to promote controlled, timely abscission of sweet

cherries. This, in turn, will lead to the improvement and standardization of mechanical

harvesting, thereby improving efficiency and increasing the economic profitability of the sweet

cherry industry.  Ultimately, the outcomes of this work may be extended to other crops where

planned abscission can be useful in managing the harvest.

**Acknowledgments**

This study was supported by a USDA Special Crop Research Initiative (SCRI) program grant

(Project No. 2009-02559) to MW and AD. Work in the Dhingra lab was supported in part by

Washington State University Agriculture Center Research Hatch Grant WNP00011. SLH, BK,

and TK acknowledge the support received from ARCS Seattle Chapter and National Institute of

Health/National Institute of General Medical Sciences through an institutional training grant

award T32-GM008336. The contents of this work are solely the responsibility of the authors and

do not necessarily represent the official views of the NIGMS or NIH. The authors thank Dr.

Richard Sharpe for assistance with RT-qPCR assays and analysis. This manuscript has been

released as a pre-print at BioRxiv (Hewitt, Kilian, et al. 2020).

**Author Contributions**

SH – Transcriptome assembly, data analysis, manuscript preparation, and editing

BK – Experimental design, data analysis, manuscript preparation

TK – Experimental design, PFRF, and tissue collection

JA – Assisted with ethephon treatment and collection of PFRF data and tissue collection

MW – Experimental design, guidance on application of ethephon and tissue collection

AD – Conceived the study, experimental design, data analysis, manuscript preparation and

editing

**Competing Interest**

770 The authors declare no competing interests, or other interests that might be perceived to

771 influence the results and/or discussion reported in this paper.

**Figure 1.** Endpoint mean control and ethephon-treatment PFRF values for (A) 'Bing' (B), 'Chelan', and (C) 'Skeena' PFAZ tissues. The dotted line represents the threshold PFRF for mechanical harvest. Asterisks indicate significant difference between ethephon-treated and control fruit at harvest (p<0.05).

**A.** ENRICHED GO TERMS: 6 HOURS POST-ETHEPHON APPLICATION

| BING | CHELAN | SKEENA |
|---|---|---|
| 1.) DNA-binding transcription factor activity  2.) iron ion binding | | |
| 1.) extracellular region  2.) heme binding | | |
| | 1.) integral component of membrane | |
| 1.) circadian rhythm.  2.) quercetin O-glucosyltransferase | | 1.) circadian rhythm  2.) quercetin O-glucosyltransferase |
| 1.) abscisic acid binding | 43.) monophenol monooxygenase activity | 1.) aromatase activity |
| 2.) activation of protein kinase activity | 44.) negative regulation of phosphoprotein phosphatase | 2.) ATP binding |
| 3.) acylglycerol lipase activity | 45.) nucleus | 3.) Cdc73/Paf1 complex |
| 4.) alpha-amino acid metabolic process | 46.) omega-6 fatty acid desaturase activity | 4.) chromatin binding |
| 5.) anthocyanidin 3-O glucosyltransferase activity | 47.) oxidoreductase activity, acting on paired donors | 5.) double-stranded methylated DNA binding |
| 6.) beta-amyrin synthase activity | 48.) pentacyclic triterpenoid biosynthetic process | 6.) gene silencing by RNA-directed DNA methylation |
| 7.) C-19 gibberellin 2-beta dioxygenase activity | 49.) phosphate ion transmembrane transport | 7.) glutathione catabolic process |
| 8.) carbohydrate transmembrane transport | 50.) phosphate ion transmembrane transporter activity | 8.) helicase activity |
| 9.) carbohydrate:proton symporter activity | 51.) phospholipase activity | 9.) histone acetyltransferase activity |
| 10.) carbon-nitrogen lyase activity | 52.) pigment biosynthetic process | 10.) histone binding |
| 11.) carboxylic acid catabolic process | 53.) poly(A)-specific ribonuclease activity | 11.) histone H3-K14 acetylation |
| 12.) catechol oxidase activity | 54.) positive transcriptional regulation in response to heat | 12.) histone H3-K18 acetylation |
| 13.) CCR4-NOT complex | 55.) protein complex oligomerization | 13.) histone H3-K23 acetylation |
| 14.) cell wall macromolecule catabolic process | 56.) protein dimerization activity | 14.) IMP biosynthetic process |
| 15.) cellular response to phosphate starvation | 57.) protein folding chaperone | 15.) mitotic spindle assembly checkpoint |
| 16.) cellular response to unfolded protein | 58.) protein phosphatase inhibitor activity | 16.) monooxygenase activity |
| 17.) chaperone cofactor-dependent protein refolding | 59.) protein refolding | 17.) mRNA splicing, via spliceosome |
| 18.) chitin binding | 60.) protein self-association | 18.) nuclear pore |
| 19.) chitin catabolic process | 61.) pyridoxal phosphate biosynthetic process | 19.) phosphoribosylformyl glycinamidine synthase activity |
| 20.) chitinase activity | 62.) regulation of biological quality | 20.) plasmodesma |
| 21.) cytokinin 7-beta glucosyltransferase activity | 63.) regulation of protein serine/threonine phosphatase | 21.) protein import into nucleus |
| 22.) cytoplasm | 64.) regulation of seed germination | 22.) regulation of DNA methylation |
| 23.) damaged DNA binding | 65.) regulation of transition time from vegetative-reproductive | 23.) regulation of DNA-dependent DNA replication |
| 24.) defense response | 66.) response to abscisic acid | 24.) regulation of transcription, DNA templated |
| 25.) gibberellin catabolic process | 67.) response to biotic stimulus | 25.) RNA binding |
| 26.) glutathione binding | 68.) response to chitin | 26.) signal transduction |
| 27.) glutathione metabolic process | 69.) response to reactive oxygen species | 27.) spliceosomal complex |
| 28.) glutathione transferase activity | 70.) response to salt stress | 28.) constituent of nuclear pore |
| 29.) glycopeptide alpha-N acetylgalactosaminidase | 71.) response to water deprivation | 29.) transcription coregulator activity |
| 30.) glycosyl compound metabolic process | 72.) RNA polymerase II cis-regulatory region DNA binding | 30.) transcription elongation from RNA polymerase II promoter |
| 31.) heat acclimation | 73.) salicylic acid metabolic process | 31.) transcriptionally active chromatin |
| 32.) heat shock protein binding | 74.) secondary active sulfate transmembrane transporter | 32.) transmembrane transport |
| 33.) hormone-mediated signaling pathway | 75.) shikimate O-hydroxycinnamoyltransferase | 33.) transmembrane transporter activity |
| 34.) hydrolase activity, acting on carbon-nitrogen bonds | 76.) signaling receptor activity | |
| 35.) integral component of plasma membrane | 77.) sucrose alpha-glucosidase activity | |
| 36.) jasmonic acid metabolic process | 78.) sucrose catabolic process | |
| 37.) lanosterol synthase activity | 79.) sulfate transmembrane transport | |
| 38.) lipid droplet | 80.) unfolded protein binding | |
| 39.) methyl indole-3-acetate esterase activity | 81.) 2-hydroxyisoflavanone dehydratase activity | |
| 40.) methyl jasmonate esterase activity | 82.) 3-oxo-myristate decarboxylase activity | |
| 41.) methyl salicylate esterase activity | 83.) 3-oxo-palmitate decarboxylase activity | |
| 42.) misfolded protein binding | 84.) 3-oxolaurate decarboxylase activity | |

**B.** ENRICHED GO TERMS: HARVEST

| BING | CHELAN | | SKEENA |
|---|---|---|---|
| 1.) extracellular region  2.) heme binding  3.) integral component of membrane | | | |
| 1.) DNA-binding transcription factor  2.) quercetin O-glucosyltransferase  3.) plasma membrane  4.) auxin response | | | |
| 1.) carbohydrate transmembrane transport | 1.) active ion transmembrane transporter activity | 29.) phosphate ion transport | 1.) phenylpropanoid biosynthetic process |
| 2.) carbohydrate transmembrane transporter activity | 2.) alpha,alpha-trehalose phosphatesynthase activity | 30.) phyllome development | 2.) response to stress |
| 3.) cell wall macromolecule catabolic process | 3.) amino acid transmembrane transport | 31.) plasmodesma | |
| 4.) chitin binding | 4.) ATP binding | 32.) polyprenol biosynthetic process | |
| 5.) chitin catabolic process | 5.) carbohydrate derivative catabolic process | 33.) polyprenyltransferase activity | |
| 6.) chitinase activity | 6.) carboxylic acid transmembrane transporter activity | 34.) polysaccharide binding | |
| 7.) DNA binding | 7.) carboxylic ester hydrolase activity | 35.) protein modification by small protein conjugation | |
| 8.) ferroxidase activity | 8.) cell wall | 36.) protein phosphorylation | |
| 9.) hydroquinone:oxygen oxidoreductase activity | 9.) cellular response to carbohydrate stimulus | 37.) protein serine/threonine kinase activity | |
| 10.) iron ion homeostasis | 10.) ceramide 1-phosphate binding | 38.) regulation of seed germination | |
| 11.) iron ion transport | 11.) ceramide 1-phosphate transfer activity | 39.) regulation of transition time from vegetative - reproductive phase | |
| 12.) jasmonic acid metabolic process | 12.) ceramide 1-phosphate transport | 40.) response to ethylene | |
| 13.) lignin catabolic process | 13.) female meiotic nuclear division | 41.) response to salt stress | |
| 14.) methyl indole-3-acetate esterase activity | 14.) floral meristem determinacy | 42.) response to water deprivation | |
| 15.) methyl jasmonate esterase activity | 15.) glycopeptide alpha-N acetylgalactosaminidase activity | 43.) S-adenosylmethionine-dependent methyltransferase activity | |
| 16.) methyl salicylate esterase activity | 16.) growth factor activity | 44.) secondary active transmembrane transporter activity | |
| 17.) nitrate import | 17.) identical protein binding | 45.) signal transduction | |
| 18.) plant organ morphogenesis | 18.) intermembrane lipid transfer | 46.) sucrose alpha-glucosidase activity | |
| 19.) post-embryonic plant morphogenesis | 19.) intracellular membrane-bounded organelle | 47.) sucrose catabolic process | |
| 20.) regulation of root development | 20.) intracellular sequestering of iron ion | 48.) sulfate transmembrane transport | |
| 21.) regulation of transcription, DNA-templated | 21.) iron ion binding | 49.) sulfate transmembrane transporter activity | |
| 22.) salicylic acid metabolic process | 22.) methylation | 50.) systemic acquired resistance | |
| | 23.) O-methyltransferase activity | 51.) terpene synthase activity | |
| | 24.) oligopeptide transmembrane transport | 52.) transcription regulatory region DNA binding | |
| | 25.) oligopeptide transmembrane transporter activity | 53.) trehalose biosynthetic process | |
| | 26.) oxidoreductase activity, acting on paired donors | 54.) trehalose metabolism in response to stress | |
| | 27.) oxidoreductase activity, acting on single donors | 55.) trehalose-phosphatase activity | |
| | 28.) peptidase activity | | |

Legend:
- Enriched GO terms unique to 'Bing'
- Enriched GO terms unique to 'Chelan'
- Enriched GO terms unique to 'Skeena'
- Enriched GO terms shared between 'Bing' and 'Chelan'
- Enriched GO terms shared between 'Bing' and 'Skeena'
- Enriched GO terms shared between 'Chelan' and 'Skeena'
- Enriched GO terms shared between 'Bing', Chelan' and, 'Skeena'

**Figure 2.** Chart displaying shared and unique enriched GO terms in ethephon-treated 'Bing', 'Chelan', and 'Skeena' at 6 hours post-ethephon treatment (A) and at harvest (B). Enrichment results are based on Fisher's Exact test with an FDR corrected p-value <0.05.

## A.

**6 HOURS POST-ETHEPHON**

Color scale: C/Eth Exp. High — C/Eth Exp. Low — Eth/C Exp. High

**Expression of Genes Associated with Enriched GOs**

| Enriched GO Term | Cultivar | DE Genes Associated with Enriched GOs | Contig | Bing 0 Hr | Bing 6 Hr | Bing Harv | Chelan 0 Hr | Chelan 6 Hr | Chelan Harv | Skeena 0 Hr | Skeena 6 Hr | Skeena Harv |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Chitinase activity/binding/catabolism; Chitin response | B | endochitinase 2-like | 22753 | | * | * | | | | | | |
| | B | endochitinase EP3-like v1 | 18291 | | * | | | | | | | |
| | B | endochitinase EP3-like v2 | 18292 | | * | | | | | | | |
| | B | endochitinase EP3-like v3 | 43552 | | * | | | | | | * | |
| | B | endochitinase-like | 3317 | | * | | | * | | | | |
| Hormone-mediated signaling pathway | B | abscisic acid receptor PYR1-like | 11946 | | * | | | | | | | |
| | B | ethylene receptor ETR2 | 6010 | | * | | | | | | | |
| | B | auxin responsive protein IAA1-like | 2474 | | * | | | * | | | | |
| | B | protein TIFY 5A-like | 44985 | | * | | | | | | * | |
| | B | protein TIFY 9 | 7749 | | * | | | | | | | |
| Jasmonic/salicylic acid metabolic process; Methyl IAA esterase activity | B | allene oxide cyclase, chloroplastic-like | 6411 | | * | | | | | | | |
| | B | 2-hydroxyisoflavanone dehydratase-like v1 | 17848 | | * | | | | * | | | |
| | B | 2-hydroxyisoflavanone dehydratase-like v2 | 17850 | | * | | | | * | | | |
| | B | 2-hydroxyisoflavanone dehydratase-like v3 | 31946 | | * | | | | | | | |
| | B | probable carboxylesterase 15 | 8498 | | * | | | | | | | |
| | B | probable carboxylesterase 18 | 9714 | | * | | | | | | | |
| | B | salicylic acid-binding protein 2-like v1 | 3684 | | * | | | | | | | |
| | B | salicylic acid-binding protein 2-like v2 | 3685 | | * | | | | | | | |
| | B | salicylic acid-binding protein 2-like v3 | 3686 | | * | * | | | | | | |
| | B | salicylic acid-binding protein 2-like v4 | 3687 | | * | * | | | | | | |
| | B | salicylic acid-binding protein 2-like v5 | 3688 | | * | | | | | | | |
| | B | salicylic acid-binding protein 2-like v6 | 3689 | | * | | | | | | | |
| | B | salicylic acid-binding protein 2-like v7 | 3690 | | * | * | | | | | | |
| | B | salicylic acid-binding protein 2-like v8 | 8874 | | * | * | | | | | | |
| | B | salicylic acid-binding protein 2-like v9 | 47678 | | * | * | | | | | | |
| ROS response | B | 16.9 kDa class I heat shock protein 1-like isoform X2 | 41037 | | * | | | | | | | |
| | B | 17.1 kDa class II heat shock protein-like | 14189 | | * | | | | | | | |
| | B | 17.4 kDa class III heat shock protein | 31787 | | * | | | | | | | |
| | B | 17.6 kDa class II heat shock protein-like v1 | 14190 | | * | | | | | | | |
| | B | 17.6 kDa class II heat shock protein-like v2 | 14191 | | * | | | | | | | |
| | B | 18.1 kDa class I heat shock protein-like v1 | 49189 | | * | | | | | | | |
| | B | 18.1 kDa class I heat shock protein-like v2 | 49819 | | * | | | | | | | |
| | B | 18.5 kDa class I heat shock protein v1 | 52677 | | * | | | | | | | |
| | B | 18.5 kDa class I heat shock protein v2 | 52677 | | * | | | | | | | |
| | B | 18.5 kDa class I heat shock protein v3 | 58968 | | * | * | | | | | | |
| | B | heat shock factor protein HSF30 isoform X1 | 7368 | | * | | | | | | | |
| Negative regulation of endopeptidase activity; Endopeptidase inhibitor activity | C | alpha-humulene 10-hydroxylase-like v1 | 11871 | | | | | | * | | | |
| | C | alpha-humulene 10-hydroxylase-like v2 | 11873 | | | | | | * | | | |
| | C | alpha-humulene 10-hydroxylase-like v3 | 16462 | | | | | | * | | | |
| | C | alpha-humulene 10-hydroxylase-like v4 | 16964 | | | | | | * | | | |
| Chromatin binding; Methylated DNA binding; Regulation of DNA methylation; Gene silencing by RNA-directed DNA methylation; Histone acetylation; Histone binding | S | nucleolar complex protein 3 homolog isoform X1 v1 | 3330 | | | | | | | | * | |
| | S | nucleolar complex protein 3 homolog isoform X1 v2 | 11292 | | | | | | | | * | |
| | S | polycomb group protein EMBRYONIC FLOWER 2 isoform X1 | 13007 | | | | | * | | | * | |
| | S | increased DNA methylation 1 v1 | 10315 | | | | | | | | * | |
| | S | increased DNA methylation 1 v2 | 17311 | | | | | | | | * | |
| | S | increased DNA methylation 1 v3 | 20390 | | | | | | | | * | |
| | S | histone acetyltransferase HAC1-like isoform X1 v1 | 944 | | | | | | | | * | |
| | S | histone acetyltransferase HAC1-like isoform X1 v2 | 19137 | | | | | | | | * | |
| | S | chromatin structure-remodeling complex protein SYD isoform X2 | 396 | | | | | | | | * | |

## B.

**HARVEST**

Color scale: C/Eth Exp. High — C/Eth Exp. Low — Eth/C Exp. High

**Expression of Genes Associated with Enriched GOs**

| GO Enriched Term | Cultivar | Description | Contig | Bing 0 Hr | Bing 6 Hr | Bing Harv | Chelan 0 Hr | Chelan 6 Hr | Chelan Harv | Skeena 0 Hr | Skeena 6 Hr | Skeena Harv |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Chiting binding and catabolism; Chitinase activity | B | class V chitinase-like | 72624 | | | * | | | | | | |
| | B | endochitinase 2-like | 22753 | | * | * | | | | | | |
| | B | endochitinase EP3-like v1 | 16426 | | | * | | | | | | |
| | B | endochitinase EP3-like v2 | 16495 | | | * | | | | | | |
| | B | endochitinase EP3-like v3 | 16496 | | | * | | * | | | * | |
| | B | endochitinase EP3-like v4 | 6142 | | | * | | | | | | |
| Lignin catabolic process | B | laccase-15-like | 47488 | | | * | | | | | | |
| | B | laccase-7-like | 48560 | | | * | | | | | | |
| | B | putative laccase-9 v1 | 10596 | | | * | | | | | | |
| | B | putative laccase-9 v2 | 62006 | | | * | | | * | | | |
| Jasmonic acid/salicylic acid metabolic process | B | salicylic acid-binding protein 2-like v4 | 3687 | | * | * | | | | | | |
| | B | salicylic acid-binding protein 2-like v5 | 3688 | | * | * | | | | | | |
| | B | salicylic acid-binding protein 2-like v8 | 8874 | | * | * | | | | | | |
| | B | salicylic acid-binding protein 2-like v9 | 47678 | | * | * | | | | | | |
| Plant organ morphogenesis | B | LOB domain-containing protein 18 | 19670 | | | * | | | | | | |
| | B | NAC transcription factor 29-like | 11020 | | | * | | | | | | |
| | B | receptor-like protein kinase HSL1 | 2398 | | | * | | | | | | * |
| | C | beta-galactosidase 8 | 2206 | | | | | | * | | | |
| | C | expansin-A1-like | 15072 | | | | | | * | | | |
| | C | pectinesterase-like | 21520 | | | | | | * | | | |
| | C | xyloglucan endotransglucosylase/hydrolase protein 32 | 46901 | | | | | | * | | | |
| | C | xyloglucan endotransglucosylase/hydrolase protein B | 26608 | | | | | | * | | | |
| Methylation | C | caffeic acid 3-O-methyltransferase-like v1 | 35033 | | | | | | * | | | |
| | C | caffeic acid 3-O-methyltransferase-like v2 | 36566 | | | | | * | * | | | |
| | C | methyltransferase-like protein 2 | 13169 | | | | | | * | | | |
| | C | methyltransferase At1g27930 | 36027 | | * | | | | * | | | |
| | C | salicylate carboxymethyltransferase-like | 12961 | | | | | | * | | * | |
| Phenylpropanoid biosynthetic process | S | fatty acyl-CoA reductase 3-like | 32869 | | | | | | | | | * |
| | S | phenylalanine ammonia-lyase 1 | 4716 | | * | | | | | | | * |
| Response to stress | S | 16.9 kDa class I heat shock protein 1-like isoform X2 | 41037 | | * | | | | | | | * |
| | S | defensin Ec-AMP-D2-like | 5976 | | | * | | | | | | * |
| | S | desiccation protectant protein Lea14 homolog | 8453 | | | | | | | | | * |
| | S | MLP-like protein 28 | 13433 | | | * | | | * | | | * |
| | S | MLP-like protein 43 | 1079 | | | * | | | | | | * |
| | S | peroxidase 10 | 68269 | | | | | | | | | * |
| | S | peroxidase 17-like | 2517 | | | * | | | | | | * |

**Figure 3.** Heatmap displaying expression of genes associated with significantly enriched unique gene ontologies (GOs) for each genotype at 6 hours post-ethephon treatment (A) and at harvest (B). Columns 1 and 2 display enriched GO terms and the genotype for which they were enriched, respectively. Columns 3 and 4 display the names of the differentially expressed genes associated with each enriched GO and their corresponding contig numbers, respectively. Heatmap represents fold change expression of ethephon treated abscission zones (AZs) versus the control AZs for each cultivar. Asterisk indicates significant differential expression (Kal's Z-test FDR-corrected p-value <0.05, NOISeq DE probability>0.8, |LOG$_2$FC|>1).

**A. 6 HOURS POST-ETHEPHON**

DNA-binding transcription factor activity

| Enriched GO Term | B | C | S | DE Genes Associated with Enriched GOs | Contig # |
|---|---|---|---|---|---|
| DNA-binding transcription factor activity | X | X | X | AP2/ERF and B3 domain-containing transcription factor RAV1 | 14058 |
| | X | X | X | AP2/ERF and B3 domain-containing transcription repressor RAV2-like | 20340 |
| | X | X | X | ethylene-responsive transcription factor 2-like | 37379 |
| | X | X | X | ethylene-responsive transcription factor 4-like v1 | 22307 |
| | X | X | X | ethylene-responsive transcription factor ERF ABR1-like | 17992 |
| | X | X | X | ethylene-responsive transcription factor ERF027 | 39727 |
| | X | X | X | ethylene-responsive transcription factor ERF034-like | 30914 |
| | X | X | X | ethylene-responsive transcription factor ERF096-like | 68474 |
| | X | X | X | ethylene-responsive transcription factor ERF098-like | 64590 |
| | X | X | X | ethylene-responsive transcription factor ERF109-like v1 | 23882 |
| | X | X | X | ethylene-responsive transcription factor ERF109-like v4 | 41515 |
| | X | X | X | ethylene-responsive transcription factor ERF113-like | 1199 |
| | X | X | X | ethylene-responsive transcription factor ERF1B-like v1 | 47562 |
| | X | X | X | ethylene-responsive transcription factor ERF1B-like v2 | 50021 |
| | X | X | X | ethylene-responsive transcription factor ERF2-like | 20167 |
| | X | X | X | ethylene-responsive transcription factor TINY | 44704 |
| | X | X | X | ethylene-responsive transcription factor TINY-like | 54876 |
| | X | X | X | ethylene-responsive transcriptionfactor ERF027-like | 33777 |
| | X | X | X | transcription factor E2FB-like isoform X2 | 15691 |
| | X | X | X | CBF/DREB1-like protein a | 66067 |
| | X | X | X | transcription factor MYB114-like | 13853 |
| | X | X | X | MYB-related protein 308-like v2 | 75886 |
| | X | X | X | probable WRKY transcription factor 26 | 2506 |
| | X | X | X | probable WRKY transcription factor 33 | 7843 |
| | X | X | X | probable WRKY transcription factor 40 | 12849 |
| | X | X | X | probable WRKY transcription factor 69 | 30796 |
| | X | X | X | probable WRKY transcription factor 75 v1 | 22200 |

Expression of Genes Associated with Enriched GOs — Bing (0 Hr, 6 Hr, Harv), Chelan (0 Hr, 6 Hr, Harv), Skeena (0 Hr, 6 Hr, Harv). Cultivar columns: B, C, S.

**B. HARVEST**

| Enriched GO Term | B | C | S | DE Genes Associated with Enriched GOs | Contig # |
|---|---|---|---|---|---|
| Auxin response | X | X | | auxin response factor ARF3 | 9103 |
| | X | X | | auxin-responsive protein IAA27 v1 | 864 |
| | X | X | | auxin-responsive protein IAA27 v2 | 8321 |
| | X | X | | auxin-responsive protein SAUR32 | 22530 |
| | X | X | | auxin-responsive protein SAUR50 | 41960 |
| | X | X | | Probable auxin efflux carrier component 1c | 5865 |
| | X | X | | protein BIG GRAIN 1-like A | 7489 |
| | X | X | | VAN3-binding protein-like isoform X1 | 8523 |
| DNA-binding transcription factor activity | X | X | | AP2/ERF and B3 domain-containing transcription repressor RAV2-like | 20340 |
| | X | X | | ethylene-responsive transcription factor 4-like v1 | 22307 |
| | X | X | | ethylene-responsive transcription factor 4-like v2 | 17802 |
| | X | X | | ethylene-responsive transcription factor ERF017 | 31967 |
| | X | X | | ethylene-responsive transcription factor ERF023 | 31950 |
| | X | X | | ethylene-responsive transcription factor ERF027 | 39727 |
| | X | X | | ethylene-responsive transcription factor ERF034-like | 30914 |
| | X | X | | ethylene-responsive transcription factor ERF096-like | 68474 |
| | X | X | | ethylene-responsive transcription factor ERF109-like v1 | 23882 |
| | X | X | | ethylene-responsive transcription factor ERF109-like v2 | 41438 |
| | X | X | | ethylene-responsive transcription factor ERF109-like v3 | 23883 |
| | X | X | | ethylene-responsive transcription factor ERF109-like v5 | 58830 |
| | X | X | | ethylene-responsive transcription factor ERF1B-like v1 | 47562 |
| | X | X | | ethylene-responsive transcription factor ERF1B-like v2 | 50021 |
| | X | X | | MYB-related protein 308-like v1 | 40430 |
| | X | X | | MYB-related protein Myb4-like | 61117 |
| | X | X | | MYB10 V1-2 | 4548 |
| | X | X | | transcription factor MYB108-like | 24917 |
| | X | X | | transcription factor MYB36-like | 7229 |
| | X | X | | transcription factor MYB41-like | 60269 |
| | X | X | | transcription factor MYB8-like | 26167 |
| | X | X | | probable WRKY transcription factor 53 | 16663 |
| | X | X | | probable WRKY transcription factor 71 | 19579 |
| | X | X | | probable WRKY transcription factor 75 v2 | 31022 |
| | X | X | | transcription repressor MYB6-like | 13058 |

Expression of Genes Associated with Enriched GOs — Bing (0 Hr, 6 Hr, Harv), Chelan (0 Hr, 6 Hr, Harv), Skeena (0 Hr, 6 Hr, Harv). Cultivar columns: B, C, S.

**Figure 4.** Heatmap displaying expression of genes associated with significantly enriched gene ontologies (GOs) shared between two cultivars 6 hours post-ethephon treatment (A) and at harvest (B). Columns 1 and 2 display enriched GO terms and the genotype for which they were enriched, respectively. Columns 3 and 4 display the names of the differentially expressed genes associated with each enriched GO and their corresponding contig numbers, respectively. Heatmap represents fold change expression of ethephon treated abscission zones (AZs) versus the control AZs for each cultivar.

**Figure 5.** Expression patterns of ethylene biosynthesis gene *1-aminocyclopropane-1-carboxylate oxidase 1* (*ACO1*), *ethylene receptor 2* (*ETR2*), and signal transducer *ethylene insensitive 2* (*EIN2*), and ethylene negative regulatory protein *reversion-to-ethylene-sensitivity 1* (*RTE1*) at 0 hours, 6 hours post-ethephon treatment, and harvest. Asterisks indicate significant difference, according to both Kal's Z-test (p<0.05) and NOIseq-sim (probability > 0.8) differential expression probability analysis, and a |logFC|>1.

**Figure 6.** Expression patterns of auxin-associated genes. A shared expression pattern was observed for auxin-associated genes at harvest. Auxin responsive transcription factors *IAA11*, *IAA13*, and *IAA27*, auxin transport facilitator *walls-are-thin 1* (*WAT1*), and auxin efflux carrier component 1c displayed comparatively lower expression in 'Skeena', intermediate expression in 'Bing', and higher expression in 'Chelan' control at harvest. Following ethephon treatment, expression was attenuated in all three cultivars, however the same relative expression trend was maintained, with 'Skeena' ethephon-treated fruit exhibiting lowest expression at harvest and 'Chelan' ethephon-treated fruit displaying the highest expression at harvest. Asterisks indicate significant difference, according to both Kal's Z-test ($p < 0.05$) and NOIseq-sim differential expression probability analysis (probability>0.8), and a |logFC|>1.

**Figure 7.** Proposed models for abscission, based on gene expression and gene ontology (GO) enrichment results following ethephon application of 'Bing', 'Chelan', and 'Skeena' sweet cherry. In 'Bing' (A), ethephon offsets the balance between ethylene and auxin in favor of ethylene. This triggers endogenous ethylene-responses that stimulates JA and SA-mediated signaling, additional hormone mediated signaling pathways, and reactive oxygen species (ROS) responses. Together, these ethylene-induced signaling pathways activate downstream processes associated with morphological changes at the PFAZ, including chitin metabolism, cell wall modification, plant organ morphogenesis, and lignin biosynthesis and deposition. The culmination of these events ultimately facilitates mechanical separation of the fruit at the PFAZ. In 'Chelan' (B), the components required for inducible abscission seem to be partially lacking. While ethephon likely contributes to enhanced ethylene signaling, free auxin levels in 'Chelan' may be high enough initially to overwhelm any ethylene response, thus limiting the cascade of signaling events leading to abscission. The activation of cell wall modifying enzymes may be responsible for the observed reduction in PFRF, however, it does not seem to be enough to facilitate complete PFAZ formation. In 'Skeena', ethephon does little to affect the overall abscission phenotype, as this cultivar is capable of natural abscission. While ethylene signaling is not high, low initial levels of free auxin may naturally shift the hormonal tug-of-war in favor of abscission, leading to moderate activation of JA metabolism and signaling. Near the time of harvest, upregulation in processes associated with epigenetic modification and, ultimately, activation of cell wall modifiers may represent intermediates in an alternate, ethylene-independent pathway to PFAZ formation.

**Table 1.** Variants identified in differentially expressed genes associated with enriched gene ontologies. The GATK RNAseq short variant discovery pipeline was used for characterization of SNPs and indels (Poplin et al., 2018).

| Gene | Associated Enriched GOs | Variant position on contig (bp) | Full Protein Length (residues) | Variant Residue Number | Variant Type | Amino Acid/ ORF change | Genotype | | |
| --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| | | | | | | | Bing | Chelan | Skeena |
| *Endochitinase-like* Contig 3317 (1140 bp) | Chitinase activity/binding/catabolism; Chitin response | 38 | 316 Translation -2 | In 3' UTR | SNP:A↔G | N/A | A/G | A/G | G/G |
| *Polycomb group protein EMBRYONIC FLOWER 2 isoform X1* Contig 13007 (913 bp) | Chromatin binding; Methylated DNA binding; Regulation of DNA methylation; Gene silencing by RNA-directed DNA methylation; Histone acetylation; Histone binding | 724 | 577 Translation -1 | 453 | DEL: ACTTGTTCCAAT GTCATTGG* | Frame Shift | DEL/* | */* | */* |
| *17.4 kDa class III heat shock protein* Contig 31787 (1075 bp) | ROS Response | 747 | 161 Translation +1 | 52 | DEL: CCACTGGAATGC TAGCCGCTCCTT TGTTCTCGTTGT TGGTTTCGTGGG CC* | Frame shift resulting in early STOP | */* | */* | DEL/DEL |
| *Probable auxin efflux carrier component 1c* Contig 5865 (2848 bp) | Auxin response | 1770 | 619 Translation +1 | 341 | SNP: G↔C | Glu↔Asp | G/G | C/C | G/G |
| *AP2/ERF and B3 domain-containing transcription factor RAV1* Contig 14058 (1503 bp) | DNA-binding transcription factor activity | 363 | 422 Translation +2 | 353 | DEL: CGGCAT* | Gly-Ile deletion; reading frame unaffected | */DEL | */* | DEL/DEL |

**Supplementary File 1.** Ethephon treatment and sampling timeline

**Supplementary File 2**. PFAZ data from 2010, 2013, and 2014 seasons

**Supplementary File 3.** Supplemental fruit color data collected at the same time as PFRF and AZ tissue sampling.

**Supplementary File 4.** Picture of fruit section sampled for PFAZ analysis

**Supplementary File 5.** Cherry assembly fasta file

**Supplementary File 6.** Spreadsheet with RPKM values of contigs that passed stringent statistical and probability filters for differential expression based on Kal's Z-test, NOIseq-sim analysis, and Log2FC expression filtering.

**Supplementary File 7.** List of all annotated contigs >2x coverage and >200bp in length

**Supplementary File 8.** RT-qPCR primers; Cq values and fold change expression summary by template

**Supplementary File 9.** Expression heatmaps for differentially expressed (DE) genes associated with shared and unique enriched gene ontology (GO) terms and DE genes associated with ethylene and auxin pathways.

**References**

Ali T, Subramanian J, Peter K P. 2012. Ethephon-Induced Abscission of "Redhaven" Peach. American Journal of Plant Sciences.2012.

Altúzar-Molina A, Lozano L, Ortíz-Berrocal M, Ramírez M, Martínez L, de Lourdes Velázquez-Hernández M, Dhar-Ray S, Silvente S, Mariano N, Shishkova S. 2020. Expression of the Legume-Specific Nod Factor Receptor Proteins Alters Developmental and Immune Responses in Rice. Plant Molecular Biology Reporter.1-20.

Axe SSA, Bush KG. 2017. 1.2 Global Sweet Cherry Production. Cherries: Botany, Production and Uses.76:2.

Blando F, Oomah BD. 2019. Sweet and sour cherries: Origin, distribution, nutritional composition and health benefits. Trends Food Sci Technol.

Blanusa T, Else MA, Atkinson CJ, Davies WJ. 2005. The regulation of sweet cherry fruit abscission by polar auxin transport. Plant growth regulation.45:189-198.

Botton A, Eccher G, Forcato C, Ferrarini A, Begheldo M, Zermiani M, Moscatello S, Battistelli A, Velasco R, Ruperti B. 2011. Signaling pathways mediating the induction of apple fruitlet abscission. Plant Physiol.155:185-208.

Bukovac M, Zucconi F, Wittenbach V, Flore J, Inoue H. 1971. Effects of (2-chloroethyl) phosphonic acid on development and abscission of maturing sweet cherry (Prunus avium L.) fruit. Amer Soc Hort Sci J.

Camera SL, Balagué C, Göbel C, Geoffroy P, Legrand M, Feussner I, Roby D, Heitz T. 2009. The Arabidopsis patatin-like protein 2 (PLP2) plays an essential role in cell death execution and differentially affects biosynthesis of oxylipins and resistance to pathogens. Mol Plant-Microbe Interact.22:469-481.

Chomczynski P, Sacchi N. 1987. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. Anal Biochem.162:156-159.

De La Fuente L, Conesa A, Lloret A, Badenes ML, Ríos G. 2015. Genome-wide changes in histone H3 lysine 27 trimethylation associated with bud dormancy release in peach. Tree Genet Genom.11:45.

Else MA, Stankiewicz-Davies AP, Crisp CM, Atkinson CJ. 2004. The role of polar auxin transport through pedicels of Prunus avium L. in relation to fruit development and retention. J Exp Bot.55:2099-2109.

Gil-Amado JA, Gomez-Jimenez MC. 2013. Transcriptome analysis of mature fruit abscission control in olive. Plant and Cell Physiology.54:244-269.

Götz S, García-Gómez JM, Terol J, Williams TD, Nagaraj SH, Nueda MJ, Robles M, Talón M, Dopazo J, Conesa A. 2008. High-throughput functional annotation and data mining with the Blast2GO suite. Nucleic Acids Res.36:3420-3435.

Han S, Hwang I. 2018. Integration of multiple signaling pathways shapes the auxin response. J Exp Bot.69:189-200.

Hendrickson C, Hewitt S, Swanson ME, Einhorn T, Dhingra A. 2019. Evidence for pre-climacteric activation of AOX transcription during cold-induced conditioning to ripen in European pear (Pyrus communis L.). PloS one.14:e0225886.

Hewitt SL, Ghogare R, Dhingra A. 2020. Glyoxylic acid overcomes 1-MCP-induced blockage of fruit ripening in Pyrus communis L. var.'D'Anjou'. Scientific Reports.10:1-14.

Hewitt SL, Hendrickson CA, Dhingra A. 2020. evidence for the involvement of Vernalization-related Genes in the Regulation of cold-induced Ripening in 'D'Anjou'and 'Bartlett'pear fruit. Scientific reports.10:1-14.

Hewitt SL, Kilian B, Koepke T, Abarca J, Whiting M, Dhingra A. 2020. Characterization of ethylene-inducible pedicel-fruit abscission zone formation in non-climacteric sweet cherry *Prunus avium* L.). bioRxiv.2020.2009.2005.284497.

Hiwasa-Tanase K, Ezura H. 2014. Climacteric and non-climacteric ripening. Fruit Ripening, Physiology, Signalling and Genomics.1-14.

Huberman M, Goren R, Zamski E. 1983. Anatomical aspects of hormonal regulation of abscission in citrus–The shoot☐peduncle abscission zone in the non☐abscising stage. Physiol Plant.59:445-454.

Kal AJ, van Zonneveld AJ, Benes V, van den Berg M, Koerkamp MG, Albermann K, Strack N, Ruijter JM, Richter A, Dujon B. 1999. Dynamics of gene expression revealed by comparison of serial analysis of gene expression transcript profiles from yeast grown on two different carbon sources. Molecular biology of the cell.10:1859-1872.

Kappel F, Granger A, Hrotkó K, Schuster M. 2012. Cherry. In: Fruit Breeding. Springer. p. 459-504.

Koepke T, Schaeffer S, Harper A, Dicenta F, Edwards M, Henry RJ, Møller BL, Meisel L, Oraguzie N, Silva H. 2013. Comparative genomics analysis in Prunoideae to identify biologically relevant polymorphisms. Plant Biotechnol J.11:883-893.

Kühn N, Serrano A, Abello C, Arce A, Espinoza C, Gouthu S, Deluc L, Arce-Johnson P. 2016. Regulation of polar auxin transport in grapevine fruitlets (Vitis vinifera L.) and the proposed role of auxin homeostasis during fruit abscission. BMC Plant Biol.16:234.

Majer C, Hochholdinger F. 2011. Defining the boundaries: structure and function of LOB domain proteins. Trends Plant Sci.16:47-52.

Meir S, Philosoph-Hadas S, Sundaresan S, Selvaraj KSV, Burd S, Ophir R, Kochanek B, Reid MS, Jiang C-Z, Lers A. 2010. Microarray Analysis of the Abscission-Related Transcriptome in the Tomato Flower Abscission Zone in Response to Auxin Depletion1[C][W][OA]. Plant Physiol.154:1929-1956.

Merelo P, Agustí J, Arbona V, Costa ML, Estornell LH, Gómez-Cadenas A, Coimbra S, Gómez MD, Pérez-Amador MA, Domingo C. 2017. Cell wall remodeling in abscission zone cells during ethylene-promoted fruit abscission in citrus. Frontiers in plant science.8:126.

Mortazavi A, Williams BA, Mccue K, Schaeffer L, Wold B. 2008. Mapping and quantifying mammalian transcriptomes by RNA-Seq. Nature Methods.5:621-628.

Niederhuth CE, Patharkar OR, Walker JC. 2013. Transcriptional profiling of the Arabidopsis abscission mutant hae hsl2by RNA-Seq. BMC Genomics.14:37.

Pfaffl MW. 2001. A new mathematical model for relative quantification in real-time RT-PCR. Nucleic Acids Res.29:e45.

Poplin R, Ruano-Rubio V, DePristo MA, Fennell TJ, Carneiro MO, Van der Auwera GA, Kling DE, Gauthier LD, Levy-Moonshine A, Roazen D, et al. 2018. Scaling accurate genetic variant discovery to tens of thousands of samples. bioRxiv.201178.

Quero-García J, Schuster M, López-Ortega G, Charlot G. 2017. Sweet cherry varieties and improvement. Cherries: botany, production and uses CAB International, Boston.60-94.

Ramakers C, Ruijter JM, Lekanne Deprez RH, Moorman AFM. 2003. Assumption-free analysis of quantitative real-time polymerase chain reaction (PCR) data. Neurosci Lett.339:62-66.

Ranocha P, Denancé N, Vanholme R, Freydier A, Martinez Y, Hoffmann L, Köhler L, Pouzet C, Renou JP, Sundberg B. 2010. Walls are thin 1 (WAT1), an Arabidopsis homolog of Medicago

truncatula NODULIN21, is a tonoplast localized protein required for secondary wall formation in fibers. The Plant Journal.63:469-483.

Ranocha P, Dima O, Felten J, Freydier A, Hoffmann L, Ljung K, Lacombe B, Corratgé C, Thibaud J-B, Sundberg B. WAT1 (WALLS ARE THIN1) defines a novel auxin transporter in plants and integrates auxin signaling in secondary wall formation in Arabidopsis fibers. Proceedings of the BMC proceedings; 2011: BioMed Central.

Ranocha P, Dima O, Nagy R, Felten J, Corratgé-Faillie C, Novák O, Morreel K, Lacombe B, Martinez Y, Pfrunder S. 2013. Arabidopsis WAT1 is a vacuolar auxin transport facilitator required for auxin homoeostasis. Nature communications.4:1-9.

Reed JW. 2001. Roles and activities of Aux/IAA proteins in Arabidopsis. Trends Plant Sci.6:420-425.

Roberts Ja, Elliott Ka, Gonzalez-Carranza ZH. 2002. Abscission, dehiscence, and other cell separation processes. Annu Rev Plant Biol.53:131-158.

Ruijter JM, Ramakers C, Hoogaars WMH, Karlen Y, Bakker O, van den hoff MJB, Moorman AFM. 2009. Amplification efficiency: Linking baseline and bias in the analysis of quantitative PCR data. Nucleic Acids Res.37.

Smith E, Whiting M. 2010. Effect of ethephon on sweet cherry pedicel-fruit retention force and quality is cultivar dependent. Plant growth regulation.60:213-223.

Stösser R, Rasmussen H, Bukovac MJ. 1969. Histochemical changes in the developing abscission layer in fruits of Prunus cerasus L. Planta.86:151-164.

Tabuchi T, Ito S, Arai N. 2000. Structural changes in the abscission zones of pedicels at different ripening stages of tomato. Journal of the Japanese Society for Horticultural Science.69:705-707.

Tabuchi T, Wada S, Arai N. Anatomical and cytological study of abscission on processing tomato fruits. Proceedings of the VI International Symposium on Processing Tomato & Workshop on Irrigation & Fertigation of Processing Tomato 487; 1998.

Tarazona S, Furió-Tarı P, Ferrer A, Conesa A. 2013. NOISeq: Differential Expression in RNA-seq.

Tarazona S, García F, Ferrer A, Dopazo J, Conesa A. 2011. NOIseq: a RNA-seq differential expression method robust for sequencing depth biases. EMBnet journal.17:18-19.

Taylor I, Walker JC. 2018. Transcriptomic evidence for distinct mechanisms underlying abscission deficiency in the Arabidopsis mutants haesa/haesa-like 2 and nevershed. BMC research notes.11:1-5.

Taylor JE, Tucker GA, Lasslett Y, Smith CJS, Arnold CM, Watson CF, Schuch W, Grierson D, Roberts JA. 1991. Polygalacturonase expression during leaf abscission of normal and transgenic tomato plants. Planta.183:133-138.

Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, del Angel G, Levy-Moonshine A, Jordan T, Shakir K, Roazen D, Thibault J, et al. 2013. From FastQ Data to High-Confidence Variant Calls: The Genome Analysis Toolkit Best Practices Pipeline. Current Protocols in Bioinformatics. 2013/10/01;43:11.10.11-11.10.33.

Wittenbach VA, Bukovac MJ. 1974a. Cherry Fruit Abscission, Evidence for Time of initiation and the involvement of Ethylene. Plant Physiol.54:494-498.

Wittenbach VA, Bukovac MJ. 1974b. Cherry Fruit Abscission: Evidence for Time of Initiation and the Involvement of Ethylene. Plant physiology.54:494-498.

Zhao Y, Athanson B, Whiting M, Oraguzie N. 2013. Pedicel-fruit retention force in sweet cherry (Prunus avium L.) varies with genotype and year. Scientia horticulturae.150:135-141.

Zhu H, Dardick CD, Beers EP, Callanhan AM, Xia R, Yuan R. 2011. Transcriptomics of shading-induced and NAA-induced abscission in apple (Malus domestica) reveals a shared pathway involving reduced photosynthesis, alterations in carbohydrate transport and signaling and hormone crosstalk. BMC Plant Biol.11:138.

# 'Chelan'

# 'Skeena'

## A. ENRICHED GO TERMS: 6 HOURS POST-ETHEPHON APPLICATION

**Shared between Bing, Chelan and Skeena:** 1.) DNA-binding transcription factor activity  2.) iron ion binding

**Shared between Bing and Chelan:** 1.) extracellular region  2.) heme binding

**Shared between Chelan and Skeena:** 1.) integral component of membrane

**Shared between Bing and Skeena:** 1.) circadian rhythm.  2.) quercetin O-glucosyltransferase

| BING | CHELAN | SKEENA |
|---|---|---|
| 1.) abscisic acid binding | 1.) aromatase activity | 1.) ADP binding |
| 2.) activation of protein kinase activity | 2.) carotene catabolic process | 2.) ATP binding |
| 3.) acylglycerol lipase activity | 3.) carotenoid dioxygenase activity | 3.) Cdc73/Paf1 complex |
| 4.) alpha-amino acid metabolic process | 4.) cysteine-type endopeptidase inhibitor activity | 4.) chromatin binding |
| 5.) anthocyanidin 3-O glucosyltransferase activity | 5.) negative regulation of endopeptidase activity | 5.) double-stranded methylated DNA binding |
| 6.) beta-amyrin synthase activity | 6.) sterol metabolic process | 6.) gene silencing by RNA-directed DNA methylation |
| 7.) C-19 gibberellin 2-beta dioxygenase activity | 7.) transferase activity, transferring hexosyl groups | 7.) glutathione catabolic process |
| 8.) carbohydrate transmembrane transport | 8.) UDP-glucosyltransferase activity | 8.) helicase activity |
| 9.) carbohydrate:proton symporter activity | | 9.) histone acetyltransferase activity |
| 10.) carbon-nitrogen lyase activity | | 10.) histone binding |
| 11.) carboxylic acid catabolic process | | 11.) histone H3-K14 acetylation |
| 12.) catechol oxidase activity | | 12.) histone H3-K18 acetylation |
| 13.) CCR4-NOT complex | | 13.) histone H3-K23 acetylation |
| 14.) cell wall macromolecule catabolic process | | 14.) IMP biosynthetic process |
| 15.) cellular response to phosphate starvation | | 15.) mitotic spindle assembly checkpoint |
| 16.) cellular response to unfolded protein | | 16.) monooxygenase activity |
| 17.) chaperone cofactor-dependent protein refolding | | 17.) mRNA splicing, via spliceosome |
| 18.) chitin binding | | 18.) nuclear pore |
| 19.) chitin catabolic process | | 19.) phosphoribosylformyl glycinamidine synthase activity |
| 20.) chitinase activity | | 20.) plasmodesma |
| 21.) cytokinin 7-beta glucosyltransferase activity | | 21.) protein import into nucleus |
| 22.) cytoplasm | | 22.) regulation of DNA methylation |
| 23.) damaged DNA binding | | 23.) regulation of DNA-dependent DNA replication |
| 24.) defense response | | 24.) regulation of transcription, DNA templated |
| 25.) gibberellin catabolic process | | 25.) RNA binding |
| 26.) glutathione binding | | 26.) signal transduction |
| 27.) glutathione metabolic process | | 27.) spliceosomal complex |
| 28.) glutathione transferase activity | | 28.) constituent of nuclear pore |
| 29.) glycopeptide alpha-N acetylgalactosaminidase | | 29.) transcription coregulator activity |
| 30.) glycosyl compound metabolic process | | 30.) transcription elongation from RNA polymerase II promoter |
| 31.) heat acclimation | | 31.) transcriptionally active chromatin |
| 32.) heat shock protein binding | | 32.) transmembrane transport |
| 33.) hormone-mediated signaling pathway | | 33.) transmembrane transporter activity |
| 34.) hydrolase activity, acting on carbon-nitrogen bonds | | |
| 35.) integral component of plasma membrane | | |
| 36.) jasmonic acid metabolic process | | |
| 37.) lanosterol synthase activity | | |
| 38.) lipid droplet | | |
| 39.) methyl indole-3-acetate esterase activity | | |
| 40.) methyl jasmonate esterase activity | | |
| 41.) methyl salicylate esterase activity | | |
| 42.) misfolded protein binding | | |
| 43.) monophenol monooxygenase activity | | |
| 44.) negative regulation of phosphoprotein phosphatase | | |
| 45.) nucleus | | |
| 46.) omega-6 fatty acid desaturase activity | | |
| 47.) oxidoreductase activity, acting on paired donors | | |
| 48.) pentacyclic triterpenoid biosynthetic process | | |
| 49.) phosphate ion transmembrane transport | | |
| 50.) phosphate ion transmembrane transporter activity | | |
| 51.) phospholipase activity | | |
| 52.) pigment biosynthetic process | | |
| 53.) poly(A)-specific ribonuclease activity | | |
| 54.) positive transcriptional regulation in response to heat | | |
| 55.) protein complex oligomerization | | |
| 56.) protein dimerization activity | | |
| 57.) protein folding chaperone | | |
| 58.) protein phosphatase inhibitor activity | | |
| 59.) protein refolding | | |
| 60.) protein self-association | | |
| 61.) pyridoxal phosphate biosynthetic process | | |
| 62.) regulation of biological quality | | |
| 63.) regulation of protein serine/threonine phosphatase | | |
| 64.) regulation of seed germination | | |
| 65.) regulation of transition time from vegetative-reproductive | | |
| 66.) response to abscisic acid | | |
| 67.) response to biotic stimulus | | |
| 68.) response to chitin | | |
| 69.) response to reactive oxygen species | | |
| 70.) response to salt stress | | |
| 71.) response to water deprivation | | |
| 72.) RNA polymerase II cis-regulatory region DNA binding | | |
| 73.) salicylic acid metabolic process | | |
| 74.) secondary active sulfate transmembrane transporter | | |
| 75.) shikimate O-hydroxycinnamoyltransferase | | |
| 76.) signaling receptor activity | | |
| 77.) sucrose alpha-glucosidase activity | | |
| 78.) sucrose catabolic process | | |
| 79.) sulfate transmembrane transport | | |
| 80.) unfolded protein binding | | |
| 81.) 2-hydroxyisoflavanone dehydratase activity | | |
| 82.) 3-oxo-myristate decarboxylase activity | | |
| 83.) 3-oxo-palmitate decarboxylase activity | | |
| 84.) 3-oxolaurate decarboxylase activity | | |

## B. ENRICHED GO TERMS: HARVEST

**Shared between Bing, Chelan and Skeena:** 1.) extracellular region  2.) heme binding  3.) integral component of membrane

**Shared between Bing and Chelan:** 1.) DNA-binding transcription factor  2.) quercetin O-glucosyltransferase  3.) plasma membrane  4.) auxin response

| BING | CHELAN | | SKEENA |
|---|---|---|---|
| 1.) carbohydrate transmembrane transport | 1.) active ion transmembrane transporter activity | 29.) phosphate ion transport | 1.) phenylpropanoid biosynthetic process |
| 2.) carbohydrate transmembrane transporter activity | 2.) alpha,alpha-trehalose phosphatesynthase activity | 30.) phyllome development | 2.) response to stress |
| 3.) cell wall macromolecule catabolic process | 3.) amino acid transmembrane transport | 31.) plasmodesma | |
| 4.) chitin binding | 4.) ATP binding | 32.) polyprenol biosynthetic process | |
| 5.) chitin catabolic process | 5.) carbohydrate derivative catabolic process | 33.) polyprenyltransferase activity | |
| 6.) chitinase activity | 6.) carboxylic acid transmembrane transporter activity | 34.) polysaccharide binding | |
| 7.) DNA binding | 7.) carboxylic ester hydrolase activity | 35.) protein modification by small protein conjugation | |
| 8.) ferroxidase activity | 8.) cell wall | 36.) protein phosphorylation | |
| 9.) hydroquinone:oxygen oxidoreductase activity | 9.) cellular response to carbohydrate stimulus | 37.) protein serine/threonine kinase activity | |
| 10.) iron ion homeostasis | 10.) ceramide 1-phosphate binding | 38.) regulation of seed germination | |
| 11.) iron ion transport | 11.) ceramide 1-phosphate transfer activity | 39.) regulation of transition time from vegetative - reproductive phase | |
| 12.) jasmonic acid metabolic process | 12.) ceramide 1-phosphate transport | 40.) response to ethylene | |
| 13.) lignin catabolic process | 13.) female meiotic nuclear division | 41.) response to salt stress | |
| 14.) methyl indole-3-acetate esterase activity | 14.) floral meristem determinacy | 42.) response to water deprivation | |
| 15.) methyl jasmonate esterase activity | 15.) glycopeptide alpha-N acetylgalactosaminidase activity | 43.) S-adenosylmethionine-dependent methyltransferase activity | |
| 16.) methyl salicylate esterase activity | 16.) growth factor activity | 44.) secondary active transmembrane transporter activity | |
| 17.) nitrate import | 17.) identical protein binding | 45.) signal transduction | |
| 18.) plant organ morphogenesis | 18.) intermembrane lipid transfer | 46.) sucrose alpha-glucosidase activity | |
| 19.) post-embryonic plant morphogenesis | 19.) intracellular membrane-bounded organelle | 47.) sucrose catabolic process | |
| 20.) regulation of root development | 20.) intracellular sequestering of iron ion | 48.) sulfate transmembrane transport | |
| 21.) regulation of transcription, DNA-templated | 21.) iron ion binding | 49.) sulfate transmembrane transporter activity | |
| 22.) salicylic acid metabolic process | 22.) methylation | 50.) systemic acquired resistance | |
| | 23.) O-methyltransferase activity | 51.) terpene synthase activity | |
| | 24.) oligopeptide transmembrane transport | 52.) transcription regulatory region DNA binding | |
| | 25.) oligopeptide transmembrane transporter activity | 53.) trehalose biosynthetic process | |
| | 26.) oxidoreductase activity, acting on paired donors | 54.) trehalose metabolism in response to stress | |
| | 27.) oxidoreductase activity, acting on single donors | 55.) trehalose-phosphatase activity | |
| | 28.) peptidase activity | | |

**Legend:**

- ▢ Enriched GO terms unique to 'Bing'
- ▢ Enriched GO terms unique to 'Chelan'
- ▢ Enriched GO terms unique to 'Skeena'
- ▢ Enriched GO terms shared between 'Bing' and 'Chelan'
- ▢ Enriched GO terms shared between 'Bing' and 'Skeena'
- ▢ Enriched GO terms shared between 'Chelan' and 'Skeena'
- ▢ Enriched GO terms shared between 'Bing', Chelan and, 'Skeena'

## A. 6 HOURS POST-ETHEPHON

Color scale: C/Eth Exp. High — C/Eth Exp. Low — Eth/C Exp. High

**Expression of Genes Associated with Enriched GOs**

| Enriched GO Term (1) | Cultivar (2) | DE Genes Associated with Enriched GOs (3) | Contig (4) | Bing 0 Hr | Bing 6 Hr | Bing Harv | Chelan 0 Hr | Chelan 6 Hr | Chelan Harv | Skeena 0 Hr | Skeena 6 Hr | Skeena Harv |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Chitinase activity/binding/catabolism; Chitin response | B | endochitinase 2-like | 22753 | | * | * | | | | | | |
| | B | endochitinase EP3-like v1 | 18291 | | * | | | | | | | |
| | B | endochitinase EP3-like v2 | 18292 | | * | | | | | | | |
| | B | endochitinase EP3-like v3 | 43552 | | * | | | | | | * | |
| | B | endochitinase-like | 3317 | | * | | | * | | | | |
| Hormone-mediated signaling pathway | B | abscisic acid receptor PYR1-like | 11946 | | * | | | | | | | |
| | B | ethylene receptor ETR2 | 6010 | | * | | | | | | | |
| | B | auxin responsive protein IAA1-like | 2474 | | * | | * | | | | | |
| | B | protein TIFY 5A-like | 44985 | | * | | | | | | * | |
| | B | protein TIFY 9 | 7749 | | * | | | | | | | |
| Jasmonic/salicylic acid metabolic process; Methyl IAA esterase activity | B | allene oxide cyclase, chloroplastic-like | 6411 | | * | | | | | | | |
| | B | 2-hydroxyisoflavone dehydratase-like v1 | 17848 | | * | | | * | | | | |
| | B | 2-hydroxyisoflavone dehydratase-like v2 | 17850 | | * | | | * | | | | |
| | B | 2-hydroxyisoflavone dehydratase-like v3 | 31946 | | * | | | | | | | |
| | B | probable carboxylesterase 15 | 8498 | | * | | | | | | | |
| | B | probable carboxylesterase 18 | 9714 | | * | | | | | | | |
| | B | salicylic acid-binding protein 2-like v1 | 3684 | | * | | | | | | | |
| | B | salicylic acid-binding protein 2-like v2 | 3685 | | * | | | | | | | |
| | B | salicylic acid-binding protein 2-like v3 | 3686 | | * | * | | | | | | |
| | B | salicylic acid-binding protein 2-like v4 | 3687 | | * | * | | | | | | |
| | B | salicylic acid-binding protein 2-like v5 | 3688 | | * | | | | | | | |
| | B | salicylic acid-binding protein 2-like v6 | 3689 | | * | | | | | | | |
| | B | salicylic acid-binding protein 2-like v7 | 3690 | | * | * | | | | | | |
| | B | salicylic acid-binding protein 2-like v8 | 8874 | | * | * | | | | | | |
| | B | salicylic acid-binding protein 2-like v9 | 47678 | | * | * | | | | | | |
| ROS response | B | 16.9 kDa class I heat shock protein 1-like isoform X2 | 41037 | | * | | | | | | | |
| | B | 17.1 kDa class II heat shock protein-like | 14189 | | * | | | | | | | |
| | B | 17.4 kDa class III heat shock protein | 31787 | | * | | | | | | | |
| | B | 17.6 kDa class II heat shock protein-like v1 | 14190 | | * | | | | | | | |
| | B | 17.6 kDa class II heat shock protein-like v2 | 14191 | | * | | | | | | | |
| | B | 18.1 kDa class I heat shock protein-like v1 | 49189 | | * | | | | | | | |
| | B | 18.1 kDa class I heat shock protein-like v2 | 49819 | | * | | | | | | | |
| | B | 18.5 kDa class I heat shock protein v1 | 52677 | | * | | | | | | | |
| | B | 18.5 kDa class I heat shock protein v2 | 52677 | | * | | | | | | | |
| | B | 18.5 kDa class I heat shock protein v3 | 58968 | | * | * | | | | | | |
| | B | heat shock factor protein HSF30 isoform X1 | 7368 | | * | | | | | | | |
| Negative regulation of endopeptidase activity; Endopeptidase inhibitor activity | C | alpha-humulene 10-hydroxylase-like v1 | 11871 | | | | | * | | | | |
| | C | alpha-humulene 10-hydroxylase-like v2 | 11873 | | | | | * | | | | |
| | C | alpha-humulene 10-hydroxylase-like v3 | 16462 | | | | | * | | | | |
| | C | alpha-humulene 10-hydroxylase-like v4 | 16964 | | | | | * | | | | |
| Chromatin binding; Methylated DNA binding; Regulation of DNA methylation; Gene silencing by RNA-directed DNA methylation; Histone acetylation; Histone binding | S | nucleolar complex protein 3 homolog isoform X1 v1 | 3330 | | | | | | | | * | |
| | S | nucleolar complex protein 3 homolog isoform X1 v2 | 11292 | | | | | | | | * | |
| | S | polycomb group protein EMBRYONIC FLOWER 2 isoform X1 | 13007 | | | | | * | | | * | |
| | S | increased DNA methylation 1 v1 | 10315 | | | | | | | | * | |
| | S | increased DNA methylation 1 v2 | 17311 | | | | | | | | * | |
| | S | increased DNA methylation 1 v3 | 20390 | | | | | | | | * | |
| | S | histone acetyltransferase HAC1-like isoform X1 v1 | 944 | | | | | | | | * | |
| | S | histone acetyltransferase HAC1-like isoform X1 v2 | 19137 | | | | | | | | * | |
| | S | chromatin structure-remodeling complex protein SYD isoform X2 | 396 | | | | | | | | * | |

## B. HARVEST

Color scale: C/Eth Exp. High — C/Eth Exp. Low — Eth/C Exp. High

**Expression of Genes Associated with Enriched GOs**

| GO Enriched Term (1) | Cultivar (2) | Description (3) | Contig (4) | Bing 0 Hr | Bing 6 Hr | Bing Harv | Chelan 0 Hr | Chelan 6 Hr | Chelan Harv | Skeena 0 Hr | Skeena 6 Hr | Skeena Harv |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Chiting binding and catabolism; Chitinase activity | B | class V chitinase-like | 72624 | | | * | | | | | | |
| | B | endochitinase 2-like | 22753 | | * | * | | | | | | |
| | B | endochitinase EP3-like v1 | 16426 | | | * | | | | | | |
| | B | endochitinase EP3-like v2 | 16495 | | | * | | | | | | |
| | B | endochitinase EP3-like v3 | 16496 | | | * | | | * | | * | |
| | B | endochitinase EP3-like v4 | 6142 | | | * | | | | | | |
| Lignin catabolic process | B | laccase-15-like | 47488 | | | * | | | | | | |
| | B | laccase-7-like | 48560 | | | * | | | | | | |
| | B | putative laccase-9 v1 | 10596 | | | * | | | | | | |
| | B | putative laccase-9 v2 | 62006 | | | * | | | * | | | |
| Jasmonic acid/salicylic acid metabolic process | B | salicylic acid-binding protein 2-like v4 | 3687 | | * | * | | | | | | |
| | B | salicylic acid-binding protein 2-like v5 | 3688 | | * | * | | | | | | |
| | B | salicylic acid-binding protein 2-like v8 | 8874 | | * | * | | | | | | |
| | B | salicylic acid-binding protein 2-like v9 | 47678 | | * | * | | | | | | |
| Plant organ morphogenesis | B | LOB domain-containing protein 18 | 19670 | | | * | | | | | | |
| | B | NAC transcription factor 29-like | 11020 | | | * | | | | | | |
| | B | receptor-like protein kinase HSL1 | 2398 | | | * | | | | | | * |
| | C | beta-galactosidase 8 | 2206 | | | | | | * | | | |
| | C | expansin-A1-like | 15072 | | | | | | * | | | |
| | C | pectinesterase-like | 21520 | | | | | | * | | | |
| | C | xyloglucan endotransglucosylase/hydrolase protein 32 | 46901 | | | | | | * | | | |
| | C | xyloglucan endotransglucosylase/hydrolase protein B | 26608 | | | | | | * | | | |
| Methylation | C | caffeic acid 3-O-methyltransferase-like v1 | 35033 | | | | | | * | | | |
| | C | caffeic acid 3-O-methyltransferase-like v2 | 36566 | | | | | * | * | | | |
| | C | methyltransferase-like protein 2 | 13169 | | | | | | * | | | |
| | C | methyltransferase At1g27930 | 36027 | | * | | | | * | | | |
| | C | salicylate carboxymethyltransferase-like | 12961 | | | | | | * | | * | |
| Phenylpropanoid biosynthetic process | S | fatty acyl-CoA reductase 3-like | 32869 | | | | | | | | | * |
| | S | phenylalanine ammonia-lyase 1 | 4716 | | * | | | | | | | |
| Response to stress | S | 16.9 kDa class I heat shock protein 1-like isoform X2 | 41037 | | * | | | | | | | * |
| | S | defensin Ec-AMP-D2-like | 5976 | | | * | | | | | | * |
| | S | desiccation protectant protein Lea14 homolog | 8453 | | | | | | | | | * |
| | S | MLP-like protein 28 | 13433 | | | * | | | * | | | * |
| | S | MLP-like protein 43 | 1079 | | | * | | | * | | | * |
| | S | peroxidase 10 | 68269 | | | | | | | | | * |
| | S | peroxidase 17-like | 2517 | | | * | | | | | | |

# A. 6 HOURS POST-ETHEPHON

Color scale: C/Eth — C/Eth — Eth/C

## Expression of Genes Associated with Enriched GOs

| Enriched GO Term [1] | B | C | S | DE Genes Associated with Enriched GOs | Contig # [4] | Bing 0 Hr | Bing 6 Hr | Bing Harv | Chelan 0 Hr | Chelan 6 Hr | Chelan Harv | Skeena 0 Hr | Skeena 6 Hr | Skeena Harv |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| DNA-binding transcription factor activity | X | X | X | AP2/ERF and B3 domain-containing transcription factor RAV1 | 14058 | | * | | | | | | | |
| | X | X | X | AP2/ERF and B3 domain-containing transcription factor repressor RAV2-like | 20340 | | * | * | | | | | | |
| | X | X | X | ethylene-responsive transcription factor 2-like | 37379 | | * | | | | | | | |
| | X | X | X | ethylene-responsive transcription factor 4-like v1 | 22307 | | | * | | * | | | | |
| | X | X | X | ethylene-responsive transcription factor ERF ABR1-like | 17992 | | * | | | | | | | |
| | X | X | X | ethylene-responsive transcription factor ERF027 | 39727 | | * | | | | * | | * | * |
| | X | X | X | ethylene-responsive transcription factor ERF034-like | 30914 | | | | | * | * | | | |
| | X | X | X | ethylene-responsive transcription factor ERF096-like | 68474 | | * | | | | * | | | |
| | X | X | X | ethylene-responsive transcription factor ERF098-like | 64590 | | * | | | | | | | |
| | X | X | X | ethylene-responsive transcription factor ERF109-like v1 | 23882 | | | | | * | * | | * | |
| | X | X | X | ethylene-responsive transcription factor ERF109-like v4 | 41515 | | * | | | | | | | |
| | X | X | X | ethylene-responsive transcription factor ERF113-like | 1199 | | * | | | | * | | | |
| | X | X | X | ethylene-responsive transcription factor ERF1B-like v1 | 47562 | | * | * | | | | | | |
| | X | X | X | ethylene-responsive transcription factor ERF1B-like v2 | 50021 | | * | * | | | | | | |
| | X | X | X | ethylene-responsive transcription factor ERF2-like | 20167 | | * | * | | | | | | |
| | X | X | X | ethylene-responsive transcription factor TINY | 44704 | | | | | | | | * | |
| | X | X | X | ethylene-responsive transcription factor TINY-like | 54876 | | | | | | | | * | |
| | X | X | X | ethylene-responsive transcriptionfactor ERF027-like | 33777 | | * | | | | | | * | |
| | X | X | X | transcription factor E2FB-like isoform X2 | 15691 | | | | | * | | | | |
| | X | X | X | CBF/DREB1-like protein a | 66067 | | * | | | | | | * | |
| | X | X | X | transcription factor MYB114-like | 13853 | | | | | | | | | |
| | X | X | X | MYB-related protein 308-like v2 | 75886 | | * | | | | | | | |
| | X | X | X | probable WRKY transcription factor 26 | 2506 | | * | | | | | | | |
| | X | X | X | probable WRKY transcription factor 33 | 7843 | | * | | | | * | | * | |
| | X | X | X | probable WRKY transcription factor 40 | 12849 | | * | | | | | | | |
| | X | X | X | probable WRKY transcription factor 69 | 30796 | | * | | | | | | | |
| | X | X | X | probable WRKY transcription factor 75 v1 | 22200 | | * | | | | | | | |

# B. HARVEST

Color scale: C/Eth — C/Eth — Eth/C

## Expression of Genes Associated with Enriched GOs

| Enriched GO Term [1] | B | C | S | DE Genes Associated with Enriched GOs | Contig # [4] | Bing 0 Hr | Bing 6 Hr | Bing Harv | Chelan 0 Hr | Chelan 6 Hr | Chelan Harv | Skeena 0 Hr | Skeena 6 Hr | Skeena Harv |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Auxin response | X | X | | auxin response factor ARF3 | 9103 | | | | | | * | | | |
| | X | X | | auxin-responsive protein IAA27 v1 | 864 | | | * | | | * | | | |
| | X | X | | auxin-responsive protein IAA27 v2 | 8321 | | | | | | * | | | |
| | X | X | | auxin-responsive protein SAUR32 | 22530 | | | * | | | | | | |
| | X | X | | auxin-responsive protein SAUR50 | 41960 | | | | | | * | | | |
| | X | X | | Probable auxin efflux carrier component 1c | 5865 | | | * | | | | | | |
| | X | X | | protein BIG GRAIN 1-like A | 7489 | | | * | | | | | | |
| | X | X | | VAN3-binding protein-like isoform X1 | 8523 | | | * | | | * | | | * |
| DNA-binding transcription factor activity | X | X | | AP2/ERF and B3 domain-containing transcription repressor RAV2-like | 20340 | | * | | | | | | | |
| | X | X | | ethylene-responsive transcription factor 4-like v1 | 22307 | | | * | | * | | | | |
| | X | X | | ethylene-responsive transcription factor 4-like v2 | 17802 | | * | * | | * | | | | |
| | X | X | | ethylene-responsive transcription factor ERF017 | 31967 | | | | | | * | | | |
| | X | X | | ethylene-responsive transcription factor ERF023 | 31950 | | * | | | | | | | |
| | X | X | | ethylene-responsive transcription factor ERF027 | 39727 | | * | | | | * | | * | * |
| | X | X | | ethylene-responsive transcription factor ERF034-like | 30914 | | | | | * | * | | | |
| | X | X | | ethylene-responsive transcription factor ERF096-like | 68474 | | * | * | | | * | | | |
| | X | X | | ethylene-responsive transcription factor ERF109-like v1 | 23882 | | | | | * | * | | * | |
| | X | X | | ethylene-responsive transcription factor ERF109-like v2 | 41438 | | | * | | | | | | |
| | X | X | | ethylene-responsive transcription factor ERF109-like v3 | 23883 | | | | | * | * | | * | |
| | X | X | | ethylene-responsive transcription factor ERF109-like v5 | 58830 | | | | | | * | | | |
| | X | X | | ethylene-responsive transcription factor ERF1B-like v1 | 47562 | | * | * | | | | | | |
| | X | X | | ethylene-responsive transcription factor ERF1B-like v2 | 50021 | | * | * | | | | | | |
| | X | X | | MYB-related protein 308-like v1 | 40430 | | | | | | * | | | |
| | X | X | | MYB-related protein Myb4-like | 61117 | | * | * | | | | | | |
| | X | X | | MYB10 V1-2 | 4548 | | * | | | | | | | |
| | X | X | | transcription factor MYB108-like | 24917 | | * | * | * | | * | | | |
| | X | X | | transcription factor MYB36-like | 7229 | | | | | | * | | | |
| | X | X | | transcription factor MYB41-like | 60269 | | | | | | * | | | |
| | X | X | | transcription factor MYB8-like | 26167 | | | | | | | | | |
| | X | X | | probable WRKY transcription factor 53 | 16663 | | | * | | | * | | | |
| | X | X | | probable WRKY transcription factor 71 | 19579 | | | | | | * | | | |
| | X | X | | probable WRKY transcription factor 75 v2 | 31022 | | | * | | | * | | | |
| | X | X | X | transcription repressor MYB6-like | 13058 | | | * | | | | | | |