TITLE

NLRC5 regulates expression of MHC-I and provides a target for anti-tumor immunity in

transmissible cancers

RUNNING TITLE

NLRC5 upregulates MHC-I on transmissible tumors

AUTHORS

Chrissie E. B. Ong[1*], Amanda L. Patchett[1], Jocelyn M. Darby[1], Jinying Chen[1,2], Guei-Sheung

Liu[1,3], A. Bruce Lyons[4], Gregory M. Woods[1], Andrew S. Flies[1*]

AFFILIATIONS

[1]Menzies Institute for Medical Research, College of Health and Medicine, University of

Tasmania, Hobart, TAS, Australia

[2]Department of Ophthalmology, the First Affiliated Hospital of Jinan University, Guangzhou,

China

[3]Ophthalmology, Department of Surgery, University of Melbourne, East Melbourne,

Australia

[4]Tasmanian School of Medicine, College of Health and Medicine, University of Tasmania,

Hobart, TAS, Australia

CORRESPONDING AUTHORS CONTACT INFORMATION

Andrew S. Flies, PhD

Menzies Institute for Medical Research, College of Health and Medicine

University of Tasmania

Private Bag 23, Hobart TAS 7000

phone: +61 3 6226 4614; email: Andy.Flies@utas.edu.au

Chrissie E. B. Ong, PhD candidate

Menzies Institute for Medical Research, College of Health and Medicine

University of Tasmania

Private Bag 23, Hobart TAS 7000

email: Chrissie.Ong@utas.edu.au

KEYWORDS

transmissible cancer, devil facial tumor, allograft, MHC-I, NLRC5, contagious cancer,

immune evasion, wild immunology

ABSTRACT

Downregulation of major histocompatibility complex I (MHC-I) on tumor cells is a primary

means of immune evasion by many types of cancer. Additionally, MHC-I proteins are a

primary target of humoral and cellular mechanisms associated with transplant rejection.

Transmissible tumors that overcome allograft rejection mechanisms and evade anti-tumor

immunity have killed thousands of wild Tasmanian devils (*Sarcophilus harrisii*). Interferon-

gamma (IFNG) upregulates surface MHC-I expression on devil facial tumor (DFT) cells but is

not sufficient to induce tumor regressions. Transcriptome analysis of IFNG-treated DFT cells

revealed strong upregulation of *NLRC5*, a master regulator of MHC-I in humans and mice. To

explore the role of NLRC5 in transmissible cancers, we developed DFT cells lines that

constitutively overexpress NLRC5. Transcriptomic results suggest that the role of NLRC5 as

a master regulator of MHC-I is conserved in devils. Furthermore, NLRC5 was shown to drive

the expression of many components of the antigen presentation pathway. To determine if

MHC-I is a target of allogeneic immune responses, we tested serum from devils with natural

DFT regressions against DFT cells. Antibody binding occurred with cells treated with IFNG

and overexpressed NLRC5. However, CRISPR/Cas9-mediated knockout of MHC-I subunit

beta-2-microglobulin (*B2M*) eliminated antibody binding to DFT cells. Consequently, MHC-I

could be identified as a target for anti-tumor and allogeneic immunity and provides mechanistic

insight into MHC-I expression and antigen presentation in marsupials. NLRC5 could be a

promising target for immunotherapy and vaccines to protect devils from transmissible cancers

and inform development of transplant and cancer therapies for humans.

INTRODUCTION

In 1996, a wild Tasmanian devil (*Sarcophilus harrisii*) was photographed with a large facial tumor. In subsequent years, similar devil facial tumors (DFTs) were recorded[1], and in 2006, it was confirmed that DFTs are clonally transmissible cancers that spread among devils through social interactions[2,3]. In 2014, a second genetically independent transmissible devil facial tumor (DFT2) was discovered in southern Tasmania[4]. Despite the independent origin of the first devil facial tumor (DFT1) and DFT2, both clonal tumors arose from a Schwann cell lineage[5,6], suggesting devils could be prone to transmissible Schwann cell cancers. These lethal and unique tumors are simultaneously cancers, allografts, and infectious diseases, and have been the primary driver of an average 77% decline in devil populations across the island state of Tasmania[7].

The successful transmission and seeding of DFT cells from one devil to another as an allograft[3] reveals its ability to circumvent both allogeneic and anti-tumor immune responses. DFT1 cells generally express little or no major histocompatibility complex class I (MHC-I) on their surface[8], an immune escape mechanism commonly observed in human cancers[9] that prevents recognition of tumor cells by cytotoxic anti-tumor CD8$^+$ T cells. Beta-2-microglobulin (B2M) is necessary for surface MHC-I expression and the clonal DFT1 cell lineage has a hemizygous mutation in the *B2M* gene[10], suggesting that immune evasion through reduced MHC-I expression has been a target of evolutionary selection pressure. Loss of MHC-I should lead to recognition and cytotoxic responses by natural killer (NK) cells. Devils have demonstrated NK-like activity *in vitro*[11] but the ongoing transmission of DFT1 cells suggests that NK cytotoxic response against DFT1 cells either do not occur or are ineffective. All DFT1 cell lines tested to date can upregulate MHC-I in response to interferon gamma (IFNG) treatment[8]. Rare cases of DFT1 regression have been reported in the wild[12] and serum antibody responses of these devils are generally higher against cell lines treated with IFNG to upregulate MHC-I[12,13]. In contrast to DFT1 cells, DFT2 cells constitutively express MHC-I, but the most highly-expressed alleles appear to be those shared by the DFT2 cells and the host devil[14]. This further suggest a critical role of MHC-I in immune evasion by DFT cells.

Upregulation of MHC-I on DFT1 cells via treatment with IFNG has served as the foundation for a vaccine against devil facial tumor disease (DFTD), which is caused by DFT1 cells. However, there are caveats to using a pleiotropic cytokine such as IFNG. IFNG plays multiple roles in the innate and adaptive immune system and can function to drive either an anti-tumor

or a pro-tumor response depending on the circumstances[15]. While IFNG is well known for

directing the immune response towards anti-tumor immunity, it also causes the upregulation of

programmed death ligand 1 (PDL1)[16] and non-classical, monomorphic MHC-I SAHA-UK on

DFT cells[14]. PDL1 and SAHA-UK molecules can be counterproductive to the cell-mediated

immune response mediated by MHC-I recognition. Additionally, the inhibition of cell

proliferation and increased DFT cell death associated with IFNG[17] constrain large-scale

production of IFNG-treated DFT cells for whole cell vaccines.

NLRC5 (NLR caspase recruitment domain containing protein 5), a member of the NOD-like

receptor (NLR) family, was identified in 2010 as the transcriptional activator of MHC-I

genes[18]. NLRC5 is strongly upregulated by IFNG and is found to be a critical mediator for

IFNG-induced MHC-I expression in humans and mice[18], but little is known about NLRC5 in

other species. NLRC5 acts with high specificity[18], and functions in MHC-I regulation by

interacting with several other transcription factors[19] to form a multi-protein complex called the

enhanceosome[20,21]. The enhanceosome activates the promoters of MHC-I genes and

components of the antigen processing machinery such as B2M, immunoproteasome subunits

PSMB8 (also known as LMP7) and PSMB9 (also known as LMP2), and transporter associated

with antigen processing 1 (TAP1)[9,18]. Aside from MHC-I regulation, NLRC5 has been reported

to be involved in innate immune responses as well as malignancy of certain cancers[22]. Despite

a potential central role of NLRC5 in immune evasion, studies of NLRC5 remain limited and

several hypothesized secondary roles of NLRC5 remain unexplored[22].

In this study, we take advantage of a unique natural experiment in which two independent

clonal tumor cell lines have essentially been passaged through hundreds of free-living animals

to assess the role of NLRC5 and MHC-I in immune evasion. The overexpression of NLRC5 in

DFT1 and DFT2 cells induced the expression of *B2M*, MHC-I heavy chain *SAHAI-01* and other

functionally-related genes. *PDL1* and the non-classical MHC-I *SAHA-UK* which are

upregulated by IFNG were not induced by NLRC5. MHC-I was constitutively expressed on

the surface of DFT cells overexpressing NLRC5, which suggests that modulation of NLRC5

expression could be a potential substitute for IFNG to increase DFT cell immunogenicity.

Additionally, MHC-I molecules on DFT cells were revealed to be an immunogenic target of

allogeneic responses in wild devils.

MATERIALS AND METHODS

Cells and Cell Culture Conditions

DFT1 cell line C5065 strain 3[23] (RRID:CVCL_LB79) and DFT2 cell lines RV (RRID:CVCL_LB80) and JV (RRID not available) were used in this study as indicated. DFT1 C5065 was provided by A-M Pearse and K. Swift of the Department of Primary Industries, Parks, Water and Environment (DPIPWE) (Hobart, TAS, Australia) and was previously established from DFT1 biopsies obtained under the approval of the Animal Ethics Committee of the Tasmanian Parks and Wildlife Service (permit numbers 33/2004-5 and 32/2005-6). DFT2 cell lines RV and JV were established from single cell suspensions obtained from tumor biopsies[4]. Cells were cultured at 35 °C with 5% CO_2 in complete RPMI medium: RPMI 1640 medium with L-glutamine (Thermo Fisher Scientific, Waltham, MA, USA), 10% heat-inactivated fetal bovine serum (Bovogen Biologicals, Melbourne, VIC, Australia), 1% (v/v) Antibiotic-Antimycotic (100X) (Thermo Fisher Scientific), 10 mM HEPES (Thermo Fisher Scientific) and 50 μM 2-mercaptoethanol (Sigma-Aldrich, St. Louis, MO, USA).

RNA Sequencing and Analysis

Initial RNA sequencing was performed using DFT1 C5065 and DFT2 RV cells treated with and without 5 ng/mL recombinant devil IFNG (provided by Walter and Eliza Hall Institute (WEHI), Melbourne, VIC, Australia) for 24 h according to the previously described protocols[6,24]. For the remaining cell lines (**Table 1**, ID # 5–9), total RNA was extracted using the NucleoSpin® RNA plus kit (Macherey Nagel, Bethlehem, USA) per manufacturer's instructions. Two replicates were prepared for each cell line. RNA sequencing was conducted at the Ramaciotti Centre for Genomics (Sydney, NSW, Australia) using the following methods. RNA integrity was assessed using Agilent TapeStation (Agilent Technologies, Santa Clara, CA, USA). All samples had RNA Integrity Number (RIN) scores of 10.0. mRNA libraries were prepared using the TruSeq Stranded mRNA Library Prep (Illumina Inc., San Diego, CA, USA). The libraries were sequenced on an Illumina NovaSeq 6000 platform (Illumina) with 100 base-pair single-end reads. The quality of the sequencing reads were analyzed using FastQC version 0.11.9[25]. Raw FASTQ files have been deposited to the European Nucleotide Archive (ENA) and are available at BioProject # PRJEB39847.

The sequencing reads were mapped to the Tasmanian devil reference genome (GCA_902635505.1 mSarHar1.11) using Subread version 2.0.0[26]. Uniquely mapped reads

were counted and assigned to genes using featureCounts[27]. Differential expression analysis of

gene counts was performed using statistical software R studio[28] on R version 4.0.0[29]. Firstly,

genes with less than 100 aligned reads across all samples were filtered out to exclude lowly

expressed genes. Gene counts were then normalized across samples by upper quartile

normalization using edgeR[30–32] and EDASeq[33,34]. Normalized read counts were scaled by

transcripts per kilobase million (TPM) to account for varied gene lengths. For differential

expression analysis, gene expression of NLRC5-overexpressing cell lines (DFT1.NLRC5,

DFT2.NLRC5) were compared against BFP-control cell lines (DFT1.BFP, DFT2.BFP) while

IFNG-treated cells (DFT1.WT + IFNG, DFT2.WT^{RV} + IFNG) were compared against the

untreated wild-type (DFT1.WT, DFT2.WT^{RV}), according to their respective tumor origin.

Differential gene expression was calculated using the *voom*[35] function in *limma*[36] with linear

modelling and empirical Bayes moderation[37] (**Supplementary Table 1**). Genes were defined

as significantly differentially expressed by applying FDR < 0.05, and log$_2$FC ≥ 2.0 (upregulated)

or ≤ -2.0 (downregulated) thresholds.

A bar plot of fold changes in mRNA expression upon treatment was created from TPM values

in GraphPad Prism version 5.03. Venn diagrams of differentially expressed genes were

developed using Venny version 2.1[38]. Heatmaps were created from log$_2$TPM values using the

ComplexHeatmap[39] package in R studio. For functional enrichment analysis, over-

representation of gene ontology (GO) and Reactome pathways was analyzed on differentially

expressed genes in R studio using functions *enrichGO* in ClusterProfiler[40] and *enrichPathway*

in ReactomePA[41], respectively. Significant GO terms and Reactome pathways were selected

by applying the cut-offs p-value < 0.001, q-value < 0.05 and adjusted p-value < 0.05. P-values

were adjusted for multiple testing using Benjamini–Hochberg method.

Plasmid Construction

The coding sequence for full length devil *NLRC5* (ENSSHAT00000015489.1) was isolated

from cDNA of devil lymph node mononuclear cells stimulated with recombinant devil IFNG[16]

(10 ng/ml, 24 h). Devil *NLRC5* was then cloned into plasmid pAF105 (detailed description of

pAF105 plasmid construction available in **Supplementary Methods 1**). For this study, devil

NLRC5 was amplified from pAF105 with overlapping ends to the 5' and 3' SfiI sites of the

Sleeping Beauty transposon plasmid pSBbi-BH[42] (a gift from Eric Kowarz; Addgene # 60515,

Cambridge, MA, USA) using Q5$^{®}$ Hotstart High-Fidelity 2X Master Mix (New England

Biolabs (NEB), Ipswich, MA, USA) (see **Supplementary Figure 1** for plasmid maps and

Supplementary Table 2 for primers and reaction conditions). The fragment was cloned into
SfiI-digested (NEB) pSBbi-BH using NEBuilder[®] HiFi DNA Assembly Cloning Kit (NEB)
and the assembled plasmid pCO1 was transformed into NEB[®] 5-alpha competent *Escherichia*
coli (High Efficiency) (NEB) according to manufacturer's instructions. Positive clones were
identified by colony PCR and the plasmid was purified using NucleoSpin[®] Plasmid EasyPure
kit (Macherey-Nagel, Düren, Germany). The cloned devil *NLRC5* transcript was verified by
Sanger sequencing using Big Dye[TM] Terminator v3.1 Cycle Sequencing Kit (Applied
Biosystems (ABI), Foster City, CA, USA) and Agencourt[®] CleanSEQ[®] (Beckman Coulter,
Brea, CA, USA) per manufacturer's instructions. The sequences were analyzed on 3500xL
Genetic Analyzer (ABI) (see **Supplementary Table 3** for list of sequencing primers). For
detailed step-by-step protocols for plasmid design and construction, reagent recipes, and
generation of stable cell lines see Bio-protocol # e3986[43].

Transfection and Generation of Stable Cell Lines
Stable cell lines of both DFT1 and DFT2 (C5065 and JV cell lines respectively) overexpressing
NLRC5 were prepared as follows. 5×10^5 cells were seeded into a 6-well plate and incubated
overnight to achieve 50–80% confluency on the day of transfection. As the vector constructed
uses a Sleeping Beauty (SB) transposon system for gene transfer, co-transfection of an
expression vector encoding a SB transposase enzyme pCMV(CAT)T7-SB100[44] (a gift from
from Zsuzsanna Izsvak; Addgene plasmid # 34879) was needed to facilitate this process. Per
2.0 mL of culture volume, 2.0 µg of plasmid DNA (1.5 µg pCO1 + 0.5 µg pCMV(CAT)T7-
SB100) was diluted in phosphate-buffered saline (PBS) to 100 µL and then added to 6.0 µg of
polythymidine (PEI) (1 mg/mL, linear, 25 kDa; Polysciences, Warrington, FL, USA) diluted
in PBS to 100 µL (3:1 ratio of PEI to DNA (w/w)). The DNA:PEI solution was mixed by gentle
pipetting and incubated at room temperature for 15 to 20 min. The media on DFT cells were
replaced with fresh complete RPMI medium and the transfection mix was added dropwise to
the cells. The cells were incubated with the DNA:PEI solution overnight at 35 °C with 5% CO_2.
The next morning, media was replaced with fresh complete RPMI media. 48 h post-transfection,
the cells were observed for fluorescence through expression of reporter gene mTagBFP and
were subjected to seven days of positive selection by adding 1 mg/mL hygromycin B (Sigma-
Aldrich) in complete RPMI medium. Once selection was complete, the cells were maintained
in 200 µg/mL hygromycin B in complete RPMI medium. pSBbi-BH was used as a control to
account for the effects of the transfection and drug selection process.

Flow Cytometric Analysis of B2M Expression

Cells were harvested and plated in a round-bottom 96-well plate (1×10^5 per well) and centrifuged at 500g for 3 min at 4 °C to discard the medium. Cells were blocked with 50 µL of 1% normal goat serum (Thermo Fisher Scientific) in FACS buffer (PBS with 0.5% BSA, 0.02% sodium azide) for 10 min on ice. After blocking, 0.4 µL anti-devil B2M mouse antibody in supernatant (a gift from Hannah Siddle)[8] diluted to a total of 50 µL in FACS buffer was added to the cells for 15 min on ice. The cells were washed with 150 µL FACS buffer and centrifuged at 500g for 3 min at 4 °C. Goat anti-mouse IgG-Alexa Fluor 488 (Thermo Fisher Scientific) was diluted in FACS buffer to 4 µg/mL and 50 µL of the solution was incubated with the target cells in the dark for 30 min on ice. The cells were washed twice with FACS buffer to remove excess secondary antibody. Lastly, the cells were resuspended in 200 µL FACSbuffer with propidium iodide (PI) (500 ng/mL) (Sigma-Aldrich) prior to analysis on BD FACSCanto[TM] II (BD Biosciences, Franklin Lakes, NJ, USA). As a positive control for surface B2M expression, DFT1 C5065 and DFT2 JV cells were stimulated with 5 ng/mL recombinant devil IFNG[16] for 24 h.

Generation of *B2M* CRISPR/Cas9 Knockout Cell Lines (B2M[-/-])

Two single guide RNAs (sgRNAs) targeting the first exon of devil *B2M* gene (ENSSHAG00000017005) were designed using a web-based CRISPR design tool CHOPCHOP[45] (**Supplementary Figure 2**). Complementary oligonucleotides encoding each *B2M* sgRNA sequence were synthesized (Integrated DNA Technologies (IDT), Coralville, IA, USA), phosphorylated and annealed before cloning into lentiCRISPRv2 plasmid[46] (a gift from Feng Zhang; Addgene # 52961) at BsmBI (NEB) restriction sites using T4 DNA ligase (NEB) (see **Supplementary Table 4** for oligonucleotide sequences). The ligated plasmids pAF217 and pAF218 were then transformed into NEB® Stable Competent *Escherichia coli* (High Efficiency) (NEB). Single colonies were selected, and the plasmids were purified using ZymoPURE™ Plasmid Miniprep Kit (Zymo Research, Irvine, CA, USA). The sgRNA sequence in each plasmid was validated by Sanger sequencing according to the method described above (see **Supplementary Table 3** for list of sequencing primers).

B2M targeting vectors pAF217 and pAF218 were each transfected into DFT1.WT and DFT1.NLRC5 cells to generate *B2M* knockout cell lines DFT1.B2M[-/-] and

252   DFT1.NLRC5.B2M$^{-/-}$. Transfection of cells were carried out as described above with the
253   exception that 1.5 μg of plasmid was used instead of 2.0 μg. A day after transfection, the cells
254   were subjected to positive selection by adding 100 μg/mL puromycin (InvivoGen, San Diego,
255   CA, USA) for a week.
256
Post-drug selection, the cells were screened and sorted multiple rounds using a Beckman-
Coulter MoFlo Astrios cell sorter to select DFT1.B2M$^{-/-}$ and DFT1.NLRC5.B2M$^{-/-}$ cells with
negative B2M expression. DFT1.B2M$^{-/-}$ cells were treated with 10 ng/mL devil recombinant
IFNG[16] for 24 h to stimulate surface B2M upregulation prior to analysis. For flow cytometry,
cells were first harvested by centrifugation at 500g for 3 min at 4 °C, and then blocked with
100 μL of 1% normal goat serum (Thermo Fisher Scientific) in complete RPMI medium for
10 min on ice. After blocking, the cells were incubated with 0.8 μL anti-devil B2M mouse
antibody in supernatant[8] diluted in complete RPMI to a total of 100 μL for 15 min on ice. The
cells were washed with 2.0 mL complete RPMI and centrifuged at 500g for 3 min at 4 °C. Next,
the cells were incubated with 100 μL of 2 μg/mL goat anti-mouse IgG-Alexa Fluor 647
(Thermo Fisher Scientific) diluted in complete RPMI in the dark for 15 min on ice. The cells
were washed with 2.0 mL of complete RPMI medium to remove excess secondary antibody.
Lastly, the cells were resuspended to a concentration of 1×10^7 cells/ml in 200 ng/mL DAPI
(Sigma-Aldrich) diluted in complete RPMI medium. B2M negative cells were selected and
bulk-sorted using cell sorter Moflo Astrios EQ (Beckman Coulter).
272
After multiple rounds of sorting to establish a B2M negative population, genomic DNA of the
cells was isolated and screened for mutations in the *B2M* gene by Sanger sequencing (see
Supplementary Table 3 for sequencing primers). Indels (insertions or deletions) in the *B2M*
gene were assessed using Inference of CRISPR Edits (ICE) analysis tool version 2.0 from
Synthego[47] (Menlo Park, CA, USA) (**Supplementary Figure 2**). *B2M* knockout cell lines: (i)
DFT1.B2M$^{-/-}$ derived from DFT1 cells transfected with pAF217, and (ii) DFT1.NLRC5.B2M$^{-/-}$
$^{-/-}$ derived from DFT1.NLRC5 transfected with pAF218 were selected for downstream analysis
(see **Table 1** for full list of cell lines).
281
Flow Cytometric Analysis of Serum Antibody Target
Serum samples from wild Tasmanian devils were collected as described[12,48]. To induce surface
expression of MHC-I, DFT cells were treated with 10 ng/mL devil recombinant IFNG[16] for 24

h prior to analysis. Cells were washed with cold FACS buffer and 1×10^5 cells per well were

plated in a round-bottom 96-well plate. The cells were centrifugated at 500*g* for 3 min at 4 °C

to discard the medium. Serum samples (see **Supplementary Table 5** for serum sample

information) were thawed on ice and diluted 1:50 with FACS buffer. 50 μL of diluted serum

was added to cells and incubated for 1 h on ice. After incubation, the cells were washed twice

with 200 μL FACS buffer. 50 μL of 10 μg/mL monoclonal mouse anti-devil IgG2b antibody

(A4-D1-2-1, provided by WEHI)[49] in FACS buffer was added to the cells and incubated for 30

min on ice. The cells were washed twice with FACS buffer and then incubated with 50 uL of

4 μg/ml goat anti-mouse IgG-Alexa Fluor 488 (Thermo Fisher Scientific) in FACS buffer for

30 min on ice, protected from light. The cells were washed twice with ice-cold PBS (Thermo

Fisher Scientific). After washing, the cells were stained with LIVE/DEAD[TM] Fixable Near-IR

Dead Cell Stain (Thermo Fisher Scientific) per manufacturer's instructions. For B2M surface

expression analysis, the cells were stained as described in the protocol above. However,

LIVE/DEAD[TM] Fixable Near-IR Dead Cell Stain (Thermo Fisher Scientific) was used instead

of PI to determine cell viability. All cells were fixed with FACS fix (0.02% sodium azide, 1.0%

glucose, 0.4% formaldehyde) diluted by 20 times prior to analysis on BD FACSCanto[TM] II

(BD Biosciences).

RESULTS

NLRC5 is upregulated in DFT1 and DFT2 cells treated with IFNG

IFNG has been shown to upregulate MHC-I[8] and PDL1[16] on DFT cells. To probe the

mechanisms driving upregulation of these key immune proteins, we performed RNA-seq using

mRNA extracted from IFNG-treated DFT1 cell line C5065 (DFT1.WT) and an IFNG-treated

DFT2 cell line RV (DFT2.WT[RV]). Markers for Schwann cell differentiation, SRY-box 10

(SOX10) and neuroepithelial marker nestin (NES), that are expressed in both DFT1 and DFT2

cells[6], were selected as internal gene controls. As expected, transcriptome analysis showed that

B2M, MHC-I gene *SAHAI-01*, and *PDL1* were strongly upregulated by IFNG. MHC-I

transactivator *NLRC5* was also upregulated upon IFNG treatment, more than a 100-fold in

DFT1.WT (275-fold) and DFT2.WT[RV] cells (124-fold) relative to untreated cells (**Fig. 1**).

NLRC5 upregulates MHC-I and antigen presentation genes but not PDL1 and non-

classical MHC-I

To assess the role of NLRC5 in antigen processing and presentation, we developed an expression vector that stably upregulates NLRC5 in DFT cells. DFT1 cell line C5065 and DFT2 cell line JV were used for production of NLRC5-overexpressing DFT cells. Following drug selection to create stable cell lines, we performed RNA-seq on DFT1 and DFT2 cells stably transfected with BFP-control and NLRC5 vectors (see **Table 1** for list of cell lines). Changes in the mRNA expression profile of DFT cells overexpressing NLRC5 relative to BFP-control cells were examined in parallel with changes observed in wild-type DFT cells following IFNG treatment (**Fig. 2** and **Supplementary Fig. 3**). The transcriptome for IFNG-treated DFT2 cells was previously generated from the DFT2 RV cell line (DFT2.WT[RV])[6]. Otherwise, all DFT2 results are from DFT2 JV.

Differential expression analysis showed that 159 genes were upregulated by IFNG (DFT1.WT + IFNG) in contrast to 40 genes by NLRC5 (DFT1.NLRC5) in DFT1 cells (**Fig. 2**). In DFT2 cells, 288 genes were upregulated by IFNG (DFT2.WT[RV] + IFNG) and 30 genes by NLRC5 (DFT2.NLRC5) (**Fig. 2**). There were ten genes that were upregulated by both IFNG and NLRC5 in DFT1 and DFT2 cells. These shared genes were predominantly related to MHC-I antigen processing and presentation pathway which suggests a role of NLRC5 in IFNG-induced MHC-I expression.

A heatmap was used to explore the expression profiles of genes associated with MHC-I and MHC-II antigen processing and presentation. SOX10 and NES, which are Schwann cell differentiation markers highly expressed in DFT1 and DFT2 cells[6], and the myelin protein periaxin (PRX), a marker for DFT1 cells[50], were included as internal controls. Overall, NLRC5 upregulated genes involved in MHC-I and MHC-II pathways to a smaller magnitude than IFNG (**Fig. 3**). NLRC5 upregulated a subset of IFNG-induced MHC-I and antigen processing machinery genes, including *B2M*, MHC-I heavy chains *SAHAI-01*, *SAHAI (LOC105750614)* and *SAHAI (LOC100927947)*, *PSMB8*, *PSMB9*, and *TAP1*. In comparison, other IFNG-induced genes such as *PSMB10*, *TAP2* and TAP binding protein (*TAPBP*) were not upregulated by NLRC5 in either DFT1.NLRC5 or DFT2.NLRC5 cells. MHC-I genes that were induced by IFNG but not NLRC5 include non-classical MHC-I genes *SAHA-UK* and *SAHA-MR1*, although the latter was only induced in DFT2 cells treated with IFNG. Additionally, *PDL1* was upregulated by IFNG, but not NLRC5. Examination of the promoter elements immediately upstream of *SAHA-UK* and *PDL1* did not identify the putative MHC-I-conserved SXY module[51] necessary for NLRC5-mediated transcription in the devil genome. A putative

interferon-stimulated response element (ISRE) for devil MHC-I genes was identified 127 bp upstream of the start codon of *SAHA-UK* (**Supplementary Fig. 4**).

NLRC5 did not consistently regulate MHC-II genes. However, the invariant chain associated with assembly of MHC-II complexes, *CD74*, was significantly upregulated in DFT1.NLRC5. Similarly, IFNG treatment on DFT1 cells only upregulated MHC-II transactivator *CIITA*. Strikingly, IFNG treatment on DFT2 cells induced several MHC-II genes such as *HLA-DRA (LOC100923003)*, *HLA-DMA (LOC100925801)*, *HLA-DMB (LOC100925533)*, *CD74* and *CIITA*.

NLRC5 primarily functions in MHC-I antigen processing and presentation but is not limited to immune-related functions

The majority of research into NLRC5 has been devoted to its role as a regulator of MHC-I expression. In addition, some studies have reported possible roles of NLRC5 in antiviral immunity, inflammation and cancer through modulation of various signaling pathways[52–57]. To identify additional biological functions of NLRC5 in DFT cells, over-representation analysis of gene ontology (GO) biological processes and Reactome pathways was performed using the list of differentially expressed genes between NLRC5-overexpressed DFT cells and BFP-controls (FDR < 0.05, $\log_2$FC ≥ 2.0 or ≤ −2.0). Both analyses revealed significant up- and downregulation of genes associated with immune system processes and developmental processes in cells overexpressing NLRC5.

Among the list of genes upregulated in DFT1.NLRC5 and DFT2.NLRC5 cells, the most significantly associated GO biological process was *antigen processing and presentation of exogenous peptide antigen via MHC class I, TAP-dependent* (**Figs. 4A** and **5A**). Several additional immune-related processes were also associated with NLRC5 overexpression, particularly in DFT1 cells. Some of these included *positive regulation of immune response*, *interferon-gamma-mediated signaling pathway*, *immune response-regulating cell surface receptor signaling pathway* (**Fig. 4A**), and *regulation of interleukin-6 biosynthetic process* (**Fig. 4C**). In DFT1.NLRC5 and DFT2.NLRC5, GO terms related to development that were significantly over-represented included *morphogenesis of an epithelium* (**Fig. 4A)** and *negative regulation of epidermis development* (**Fig. 5A**), respectively.

As DFT cells are of neuroendocrine origin, specifically of the Schwann cell lineage[5,6], a number of neural-related genes were targeted by NLRC5. In DFT2 cells, NLRC5 upregulated genes that are involved in *myelination*, which are usually expressed at low levels in DFT2 cells[6] (**Fig. 5A**). These genes include brain enriched myelin associated protein 1 (*BCAS1*), myelin binding protein (*MBP*), myelin protein zero (*MPZ*) and UDP glycosyltransferase 8 (*UGT8*) (**Fig. 5B**). Furthermore, many of the downregulated genes in DFT2.NLRC5 were related to nervous system function, mainly pertaining to *synaptic signaling* and *sensory perception* (**Fig. 5C**).

Reactome pathway analysis revealed an enrichment of pathways that were consistent with those identified by GO analysis. This included enrichment of the *ER-phagosome pathway* and *antigen processing-cross presentation* in DFT1.NLRC5 (**Table 2**) and DFT2.NLRC5 (**Table 3**); *signaling by the B cell receptor (BCR)* in DFT1.NLRC5; and *transmission across chemical synapses* in DFT2.NLRC5 cells. Interestingly, nuclear factor of activated T cells 1 (*NFATC1*), protein kinase C beta (*PRKCB*), *PSMB8* and *PSMB9*, associated with several GO immune-related processes in DFT1 cells overexpressing NLRC5 (**Fig. 4B**), were enriched for the *beta-catenin independent WNT signaling* pathway (**Table 2**). Other enriched pathways included those involved in extracellular matrix organization such as *collagen chain trimerization* (**Table 2**) and *assembly of collagen fibrils and other multimeric structures* (**Table 3**).

NLRC5 induces MHC-I expression on the cell surface

To determine if NLRC5 is capable of regulating MHC-I expression at the protein level, surface MHC-I was analyzed by flow cytometry in DFT cells overexpressing NLRC5 using a monoclonal antibody against B2M[8]. The overexpression of NLRC5 induced upregulation of surface expression of B2M in both DFT1.NLRC5 (**Fig. 6A**) and DFT2.NLRC5 cells (**Fig. 6B**). The level of B2M expression was also comparable to wild-type DFT cells treated with IFNG.

Next, we assessed the stability of NLRC5-induced MHC-I expression by examining the expression of B2M in long-term cultures. One-month post-drug selection, DFT1.NLRC5 cells cultured in the presence or absence of hygromycin B were stained for B2M every four weeks for a total of 12 weeks. As shown in **Fig. 6A**, MHC-I expression was stably maintained in DFT1.NLRC5 cells, with or without ongoing drug selection pressure throughout the 12-week culture thus, demonstrating the relative stability of the human EF1a promoter driving NLRC5 expression in long-term cell cultures. PDL1 was also not upregulated on the cell surface in

NLRC5-overexpressing DFT cells compared to IFNG-treated DFT cells (**Supplementary Fig. 5**).

MHC-I is a predominant target of anti-DFT antibody responses

It was previously reported that the antibodies from devils infected with DFT1 were specific to MHC-I, as determined by incubating serum from these devils with IFNG-treated DFT cells[12]. Considering the diverse roles of IFNG, there could be other IFNG-induced antigens that can serve as targets for the anti-DFT antibody response.

To establish if MHC-I is the target of anti-DFT serum antibodies, surface MHC-I expression was first ablated by knocking out the hemizygous *B2M* allele[10] in wild-type DFT1 cells (DFT1.WT) and NLRC5-overexpressing DFT1 cells (DFT1.NLRC5) using CRISPR/Cas9 technology. Gene disruption of *B2M* was confirmed by genomic DNA sequencing (**Supplementary Fig. 2**), and flow cytometry using a monoclonal anti-B2M antibody (**Fig. 7**). CRISPR/Cas9-mediated *B2M* knockout (B2M$^{-/-}$) in DFT1 cells rendered the cells irreversibly deficient for surface expression of B2M despite IFNG and NLRC5 stimulation (DFT1.B2M$^{-/-}$ + IFNG and DFT1.NLRC5.B2M$^{-/-}$). Due to the pivotal role of B2M in stability of MHC-I complex formation and surface presentation[58-62], absence of surface B2M is indicative of a lack of surface MHC-I expression.

After surface MHC-I ablation was confirmed, serum from six wild devils (TD1–TD6) that demonstrated natural DFT1 regressions[12] was tested against DFT1 cell lines with B2M knocked out (DFT1.B2M$^{-/-}$ and DFT1.NLRC5.B2M$^{-/-}$). Serum from a healthy devil (TD7) and an immunized devil with induced tumor regression (My)[48] were used as negative and positive controls for antibody binding. All six sera from DFTD$^+$ devils (TD1–TD6) showed weak to no binding to DFT1.WT and DFT1.BFP, which are inherently negative for surface MHC-I (**Fig. 7**). With forced expression of MHC-I using IFNG (DFT1.WT + IFNG) and NLRC5 (DFT1.NLRC5), a positive shift in antibody binding was observed. There was no apparent difference in the level of antibody binding between IFNG-treated and NLRC5-overexpressing DFT1 cells, suggesting a similarity between the antibody target(s) induced by IFNG and NLRC5. Following B2M knockout, antibody binding of all six sera was reduced in both IFNG-induced (DFT1.B2M$^{-/-}$ + IFNG) and NLRC5-induced B2M knockout DFT1 cells (DFT1.NLRC5.B2M$^{-/-}$), suggesting that MHC-I is a target of DFT1-specific antibody responses in natural tumor regressions.

DISCUSSION

Overexpression of NLRC5 in DFT cells has revealed a major and evolutionarily conserved role for NLRC5 in MHC-I antigen processing and presentation. Consistent with studies in human and mouse cell lines[9,18,63–65], NLRC5 induced expression of classical MHC-I genes (*SAHAI-01, SAHAI (LOC105750614), SAHAI (LOC100927947)), B2M*, *PSMB8*, *PSMB9* and *TAP1* in both DFT1 and DFT2 cells. Despite the lack of increase in *TAP2* expression, the selective upregulation of MHC-I and other functionally-related genes by NLRC5 was sufficient to restore MHC-I molecules on the cell surface. Although the peptide transport function of TAP proteins typically involves the formation of TAP1 and TAP2 heterodimers, homodimerization of TAP proteins have been described[66,67]. However, the functionality of TAP1 homodimers remains to be verified. The conservation of genes of the MHC-I pathway, regulated by NLRC5 across species, highlights the important role of NLRC5 in MHC-I regulation.

Previous studies have shown that sera from wild devils with natural DFT1 regressions contained high titers of antibody that bound to IFNG-treated DFT1 cells. It was proposed that the primary antibody targets were MHC-I proteins[12]. Additionally, some of these devils experienced tumor regression despite the lack of strong evidence for immune cell infiltration into tumors. The function of NLRC5 that is mainly restricted to MHC-I regulation compared with IFNG provided an opportunity to re-examine antibody target(s) of serum antibodies from wild devils burdened with DFTs. A clear understanding of immunogenic targets of DFTs will provide direction for a more effective vaccine against DFTs.

The MHC-I complex was identified as the predominant target of anti-DFT serum antibodies. The antibody binding intensity against NLRC5-overexpressing DFT cells was similar to IFNG-treated DFT cells, suggesting similar levels of target antigen expression. When MHC-I expression was ablated through *B2M* knockout, antibody binding was reduced to almost background levels despite IFNG and NLRC5 stimulation. This discovery presents an option to exploit NLRC5 for induction of immunity via MHC-I expression to potentiate an anti-tumor immune response in Tasmanian devils. Although cellular immunity is likely a key mechanism for tumor rejection, B cells and antibodies can play eminent roles in transplant rejection[68] and anti-tumor immunity[69]. B cells can promote rejection through antibody-dependent mechanisms that facilitate FcR-mediated phagocytosis by macrophages, antibody-dependent cellular cytotoxicity (ADCC) by NK cells, complement activation and antigen uptake by dendritic cells

(reviewed by Yuen et al.)[70]. Moreover, B cells can enhance immune surveillance and response

through direct antigen presentation to T cells and production of immune-modulating molecules

such as cytokines and chemokines[70].

Caldwell et al. reported that the most highly expressed MHC alleles on DFT2 cells are those

that matched host MHC alleles[14], which suggests that DFT cells may hide from host defenses

or induce immunological tolerance via shared MHC alleles. If MHC-I is the major antibody

target and potentially the overall immune system target, devils having the largest MHC

mismatch with DFT cells will be the most likely to have strong MHC-I specific responses and

reject DFTs, leading to natural selection in the wild. For example, previous studies have shown

that some devils have no functional MHC-I allele at the UA loci and that these individuals can

be homozygous at the UB and UC loci[48]. These individuals present a reduced MHC-peptide

that would have the lowest probability of a match to the DFT MHC alleles that induce host

DFT1 tolerance. However, selection for reduced genetic diversity in MHC alleles would be

unfavorable for long-term conservation. A prophylactic vaccine would ideally be designed to

assist in the preservation of the genetic diversity of wild devils[71].

Although the MHC proteins themselves are likely a primary target of humoral and cellular

immunity, MHC-I alleles generally differ by only a few amino acids[14,72]. Mutations in DFTs

and somatic variation between host and tumor cells provide a rich source of additional antigenic

targets for humoral and cellular immunity[10]. The reduction in antibody binding to B2M

knockout cells suggests that these tumor antigens are unlikely to be the primary antibody

targets, although binding of antibodies to peptide-MHC complexes cannot be excluded.

Knocking out individual MHC alleles in DFT cells or overexpression of MHC alleles in

alternative non-DFT cell lines could be used to disentangle the importance of specific alleles

and investigate the potential for peptide-MHC complexes to be antibody targets.

Our results confirm that IFNG affects more immunoregulatory processes than NLRC5.

However, the functional dichotomy of IFNG in cancer means that NLRC5 modulation could

be an alternative to IFNG treatment for enhancing tumor cell immunogenicity in a range of

species, including human. Importantly, NLRC5 upregulated B2M on the surface of DFT cells

to similar levels as IFNG, but it does not upregulate inhibitory molecules. The restoration of

functional MHC-I molecules without concomitant upregulation of PDL1 and SAHA-UK has

multiple advantages over IFNG for triggering effective cytotoxic responses against DFT cells.

First, cells transfected with NLRC5 constitutively express MHC-I and therefore do not require

culturing in IFNG, which can be problematic as IFNG can also reduce cell viability[17]. Second,

PDL1 negatively regulates T cell responses by inducing T cell anergy[73] and apoptosis[74] while

limiting T cell activity[75]. Moreover, PDL1 promotes tumor growth and survival by stimulating

cell proliferation[76] and resistance to T cell killing[77,78]. Third, the expression of monomorphic

MHC-I SAHA-UK induced by IFNG would allow DFT cells to escape cytotoxic attack from

both NK cells and CD8[+] T cells[79]. Fourth, several other immune checkpoint protein receptor-

ligand interactions were recently shown to be conserved in devils[80,81], but we found no

significant upregulation these genes by NLRC5. The ability to improve tumor immunogenicity

in the absence of inhibitory signals has positive implications for immunization and

immunotherapeutic strategies. NLRC5 could evoke protective anti-tumor immunity against

DFTs, similar to NLRC5-expressing B16-F10 melanoma cells in mice[64].

The absence of a regulatory effect on *SAHA-UK* and *PDL1* by NLRC5 in contrast to IFNG

could be due to the composition of the promoter elements of these genes. The promoter of

MHC class I genes consists of three conserved cis-regulatory elements: a NFκB-binding

Enhancer A region, an interferon-stimulated response element (ISRE) and a SXY module[82,83].

The SXY module is critical for NLRC5-mediated MHC-I transactivation as it serves as the

binding site for the multi-protein complex formed between NLRC5 and various transcription

factors[19,21,84]. An ISRE and SXY module is present within 200 base pairs of the start codon for

all three classical devil MHC-I genes[51]. We identified an ISRE element in the *SAHA-UK*

promoter region but were unable to identify an SXY module in this region. This could explain

the upregulation of *SAHA-UK* upon IFNG stimulation but not in NLRC5-overexpressing DFT

cells. Similarly, the SXY module was not identified in orthologues of SAHA-UK, Modo-UK

in the grey short-tailed opossum[85] and Maeu-UK in the tammar wallaby[86]. The difference in

regulation and therefore, pattern of expression of the UK gene in marsupials[85–87] may reflect a

separate function from classical MHC-I. The marsupial UK gene has been hypothesized to play

a marsupial-specific role in conferring immune protection to vulnerable newborn marsupials

during their pouch life[86]. SXY modules are typically not found in the promoter region of

PDL1[88] therefore, it is not expected for NLRC5 to be a regulator of PDL1. Rather, IFNG-

mediated induction of PDL1 occurs via transcription factor interferon regulatory factor 1 (IRF-

1)[88], which is induced by STAT1[89].

Beyond MHC-I regulation, NLRC5 expression in DFT1 cells displayed other beneficial immune-regulating functions, mainly via the non-canonical β-catenin-independent WNT signaling pathway. One of the downstream effectors that was upregulated by NLRC5 included *PRKCB*, an activator of NFκB in B cells[90]. NFκB is a family of pleiotropic transcription factors known to regulate several immune and inflammatory responses including cellular processes such as cell proliferation and apoptosis[91]. In recent years, aberrations in NFκB signaling have been implicated in cancer development and progression[92,93]. The regulation of NFκB signaling by NLRC5 has been documented in several studies although the findings have been contradictory[54,55,94,95].

In summary, we have demonstrated the role of NLRC5 in MHC-I regulation of DFT cells thereby, displaying the functional conservation of NLRC5 across species. The finding that MHC-I is a major antibody target in wild devils with natural DFT regression can help guide DFTD vaccine development and conservation management strategies. NLRC5-overexpressing DFT cells can be harnessed to elicit both cellular and humoral immunity against future and pre-existing DFT infections in wild devils using MHC-I as a target. Given the prevalence of altered MHC-I expression in cancer as a form of immune escape mechanism[96–98], NLRC5 presents as a new target for providing an insight into the role of MHC-I in cancer as well as transplantation, and its manipulation for human cancer treatment and transplant tolerance.

ACKNOWLEDGEMENTS

The authors would like to thank Patrick Lennard, Peter Murphy, and Candida Wong for assistance in the lab and Terry Pinfold for assistance in flow cytometry. We thank Hannah Siddle for supplying the monoclonal antibody for B2M and for offering her expertise in devil MHC-I immunogenetics. We wish to thank G. Ralph for ongoing care of Tasmanian devils, the Bonorong Wildlife Sanctuary for providing access to Tasmanian devils, and R. Pye for providing care for devils and collecting blood samples. This work was supported by ARC DECRA grant # DE180100484 and ARC Discovery grant # DP180100520, University of Tasmania Foundation Dr. Eric Guiler Tasmanian Devil Research Grant through funds raised by the Save the Tasmanian Devil Appeal (2013, 2015, 2017).

AUTHOR CONTRIBUTIONS

ABL, ALP, ASF, CEBO and GMW designed the study. ALP, ASF, CEBO, GSL, JC, and JMD developed the technology. CEBO and JMD performed the experiments. ALP and CEBO

587 performed bioinformatic analyses. CEBO created the figures. ALP, ASF, and CEBO analyzed

588 the data. CEBO wrote the manuscript, and all authors edited the manuscript.

590 **CONFLICT OF INTEREST**

591 The authors declare that the research was conducted in the absence of any commercial or

592 financial relationships that could be construed as a potential conflict of interest.

REFERENCES

Jones ME, Paetkau D, Geffen E, Moritz C. Genetic diversity and population structure of Tasmanian devils, the largest marsupial carnivore. *Mol Ecol* 2004; **13**: 2197–2209.

Loh R, Hayes D, Mahjoor A, O'Hara A, Pyecroft S, Raidal S. The Immunohistochemical Characterization of Devil Facial Tumor Disease (DFTD) in the Tasmanian Devil (Sarcophilus harrisii). *Vet Pathol* 2006; **43**: 896–903.

Pearse A-M, Swift K. Allograft theory: Transmission of devil facial-tumour disease. *Nature* 2006; **439**: 549.

Pye RJ, Pemberton D, Tovar C, Tubio JMC, Dun KA, Fox S *et al.* A second transmissible cancer in Tasmanian devils. *Proc Natl Acad Sci U S A* 2016; **113**: 374–379.

Murchison EP, Tovar C, Hsu A, Bender HS, Kheradpour P, Rebbeck CA *et al.* The Tasmanian devil transcriptome reveals Schwann cell origins of a clonally transmissible cancer. *Science* 2010; **327**: 84–87.

Patchett AL, Coorens THH, Darby J, Wilson R, McKay MJ, Kamath KS *et al.* Two of a kind: transmissible Schwann cell cancers in the endangered Tasmanian devil (*Sarcophilus harrisii*). *Cell Mol Life Sci* 2020; **77**: 1847–1858.

Lazenby BT, Tobler MW, Brown WE, Hawkins CE, Hocking GJ, Hume F *et al.* Density trends and demographic signals uncover the long-term impact of transmissible cancer in Tasmanian devils. *J Appl Ecol* 2018; **55**: 1368–1379.

Siddle H V, Kreiss A, Tovar C, Yuen CK, Cheng Y, Belov K *et al.* Reversible epigenetic down-regulation of MHC molecules by devil facial tumour disease illustrates immune escape by a contagious cancer. *Proc Natl Acad Sci U S A* 2013; **110**: 5103–5108.

Yoshihama S, Roszik J, Downs I, Meissner TB, Vijayan S, Chapuy B *et al.* NLRC5/MHC class I transactivator is a target for immune evasion in cancer. *Proc Natl Acad Sci U S A* 2016; **113**: 5999–6004.

Stammnitz MR, Coorens THH, Gori KC, Hayes D, Fu B, Wang J *et al.* The Origins and Vulnerabilities of Two Transmissible Cancers in Tasmanian Devils. *Cancer Cell* 2018; **33**: 607-619.e15.

Brown GK, Kreiss A, Lyons AB, Woods GM. Natural killer cell mediated cytotoxic responses in the Tasmanian devil. *PLoS One* 2011; **6**: e24475.

Pye R, Hamede R, Siddle H V., Caldwell A, Knowles GW, Swift K *et al.* Demonstration of immune responses against devil facial tumour disease in wild

Tasmanian devils. *Biol Lett* 2016; **12**: 20160553.

13   Pye R, Patchett A, McLennan E, Thomson R, Carver S, Fox S *et al.* Immunization
Strategies Producing a Humoral IgG Immune Response against Devil Facial Tumor
Disease in the Majority of Tasmanian Devils Destined for Wild Release. *Front*
Immunol 2018; **9**: 259.

14   Caldwell A, Coleby R, Tovar C, Stammnitz MR, Kwon YM, Owen RS *et al.* The
newly-arisen Devil facial tumour disease 2 (DFT2) reveals a mechanism for the
emergence of a contagious cancer. *Elife* 2018; **7**: e35314.

15   Zaidi MR. The Interferon-Gamma Paradox in Cancer. *J Interf Cytokine Res* 2019; **39**:
30–38.

16   Flies AS, Lyons AB, Corcoran LM, Papenfuss AT, Murphy JM, Knowles GW *et al.*
PD-L1 is not constitutively expressed on Tasmanian devil facial tumor cells but is
strongly upregulated in response to IFN-γ and can be expressed in the tumor
microenvironment. *Front Immunol* 2016; **7**: 581.

17   Ong CEB, Lyons AB, Woods GM, Flies AS. Inducible IFN-γ Expression for MHC-I
Upregulation in Devil Facial Tumor Cells. *Front Immunol* 2019; **9**: 3117.

18   Meissner TB, Li A, Biswas A, Lee K-H, Liu Y-J, Bayir E *et al.* NLR family member
NLRC5 is a transcriptional regulator of MHC class I genes. *Proc Natl Acad Sci U S A*
2010; **107**: 13794–13799.

19   Meissner TB, Liu Y-J, Lee K-H, Li A, Biswas A, van Eggermond MCJA *et al.*
NLRC5 cooperates with the RFX transcription factor complex to induce MHC class I
gene expression. *J Immunol* 2012; **188**: 4951–4958.

20   Gobin SJP, van Zutphen M, Westerheide SD, Boss JM, van den Elsen PJ. The MHC-
specific enhanceosome and its role in MHC class I and β2-microglobulin gene
transactivation. *J Immunol* 2001; **167**: 5175–5184.

21   Neerincx A, Rodriguez GM, Steimle V, Kufer TA. NLRC5 controls basal MHC class I
gene expression in an MHC enhanceosome-dependent manner. *J Immunol* 2012; **188**:
4940–4950.

22   Benkő S, Kovács EG, Hezel F, Kufer TA. NLRC5 Functions beyond MHC I
Regulation-What Do We Know So Far? *Front Immunol* 2017; **8**: 150.

23   Pearse A-M, Swift K, Hodson P, Hua B, McCallum H, Pyecroft S *et al.* Evolution in a
transmissible cancer: a study of the chromosomal changes in devil facial tumor (DFT)
as it spreads through the wild Tasmanian devil population. *Cancer Genet* 2012; **205**:
101–112.

24 Patchett AL, Wilson R, Charlesworth JC, Corcoran LM, Papenfuss AT, Lyons BA *et al*. Transcriptome and proteome profiling reveals stress-induced expression signatures of imiquimod-treated Tasmanian devil facial tumor disease (DFTD) cells. *Oncotarget* 2018; **9**: 15895–15914.

25 Andrews S. FastQC: A Quality Control tool for High Throughput Sequence Data. 2010.https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

26 Liao Y, Smyth GK, Shi W. The Subread aligner: fast, accurate and scalable read mapping by seed-and-vote. *Nucleic Acids Res* 2013; **41**: e108–e108.

27 Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* 2014; **30**: 923–930.

28 RStudio Team. RStudio: Integrated Development Environment for R. 2020.http://www.rstudio.com/.

29 R Core Team. R: A Language and Environment for Statistical Computing. 2020.https://www.r-project.org/.

30 Robinson MD, McCarthy DJ, Smyth GK. edgeR: A Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 2009; **26**: 139–140.

31 Robinson MD, Oshlack A. A scaling normalization method for differential expression analysis of RNA-seq data. *Genome Biol* 2010; **11**: R25.

32 Anders S, Huber W. Differential expression analysis for sequence count data. *Genome Biol* 2010; **11**: R106.

33 Bullard JH, Purdom E, Hansen KD, Dudoit S. Evaluation of statistical methods for normalization and differential expression in mRNA-Seq experiments. *BMC Bioinformatics* 2010; **11**: 94.

34 Risso D, Schwartz K, Sherlock G, Dudoit S. GC-Content Normalization for RNA-Seq Data. *BMC Bioinformatics* 2011; **12**: 480.

35 Law CW, Chen Y, Shi W, Smyth GK. voom: Precision weights unlock linear model analysis tools for RNA-seq read counts. *Genome Biol* 2014; **15**: R29.

36 Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W *et al. limma* powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res* 2015; **43**: e47.

37 Phipson B, Lee S, Majewski IJ, Alexander WS, Smyth GK. Robust hyperparameter estimation protects against hypervariable genes and improves power to detect differential expression. *Ann Appl Stat* 2016; **10**: 946–963.

38   Oliveros JC. Venny. An interactive tool for comparing lists with Venn's diagrams.

2015.https://bioinfogp.cnb.csic.es/tools/venny/index.html.

39   Gu Z, Eils R, Schlesner M. Complex heatmaps reveal patterns and correlations in

multidimensional genomic data. *Bioinformatics* 2016; **32**: 2847–2849.

40   Yu G, Wang LG, Han Y, He QY. ClusterProfiler: An R package for comparing

biological themes among gene clusters. *Omi A J Integr Biol* 2012; **16**: 284–287.

41   Yu G, He QY. ReactomePA: An R/Bioconductor package for reactome pathway

analysis and visualization. *Mol Biosyst* 2016; **12**: 477–479.

42   Kowarz E, Löscher D, Marschalek R. Optimized Sleeping Beauty transposons rapidly

generate stable transgenic cell lines. *Biotechnol J* 2015; **10**: 647–653.

43   Flies AS, Darby JM, Murphy PR, Pinfold TL, Patchett AL, Lennard PR. Generation

and Testing of Fluorescent Adaptable Simple Theranostic (FAST) Proteins. *Bio-*

protocol 2020; **10**: e3696.

44   Mátés L, Chuah MKL, Belay E, Jerchow B, Manoj N, Acosta-Sanchez A *et al.*

Molecular evolution of a novel hyperactive Sleeping Beauty transposase enables

robust stable gene transfer in vertebrates. *Nat Genet* 2009; **41**: 753–761.

45   Labun K, Montague TG, Krause M, Torres Cleuren YN, Tjeldnes H, Valen E.

CHOPCHOP v3: Expanding the CRISPR web toolbox beyond genome editing.

Nucleic Acids Res 2019; **47**: W171–W174.

46   Sanjana NE, Shalem O, Zhang F. Improved vectors and genome-wide libraries for

CRISPR screening. *Nat Methods* 2014; **11**: 783–784.

47   Synthego. Synthego Performance Analysis, ICE Analysis.

2019.https://www.synthego.com/products/bioinformatics/crispr-analysis (accessed 31

Jul2020).

48   Tovar C, Pye RJ, Kreiss A, Cheng Y, Brown GK, Darby J *et al.* Regression of devil

facial tumour disease following immunotherapy in immunised Tasmanian devils. *Sci*

Rep 2017; **7**: 43827.

49   Howson LJ, Morris KM, Kobayashi T, Tovar C, Kreiss A, Papenfuss AT *et al.*

Identification of dendritic cells, B cell and T cell subsets in Tasmanian devil lymphoid

tissue; evidence for poor immune cell infiltration into devil facial tumors. *Anat Rec*

(Hoboken) 2014; **297**: 925–938.

50   Tovar C, Obendorf D, Murchison EP, Papenfuss AT, Kreiss A, Woods GM. Tumor-

specific diagnostic marker for transmissible facial tumors of Tasmanian devils:

immunohistochemistry studies. *Vet Pathol* 2011; **48**: 1195–1203.

51   Cheng Y, Stuart A, Morris K, Taylor R, Siddle H, Deakin J *et al.* Antigen-presenting

genes and genomic copy number variations in the Tasmanian devil MHC. *BMC*

Genomics 2012; **13**: 87.

52   Kuenzel S, Till A, Winkler M, Häsler R, Lipinski S, Jung S *et al.* The nucleotide-

binding oligomerization domain-like receptor NLRC5 is involved in IFN-dependent

antiviral immune responses. *J Immunol* 2010; **184**: 1990–2000.

53   Neerincx A, Lautz K, Menning M, Kremmer E, Zigrino P, Hösel M *et al.* A role for

the human nucleotide-binding domain, leucine-rich repeat-containing family member

NLRC5 in antiviral responses. *J Biol Chem* 2010; **285**: 26223–26232.

54   Cui J, Zhu L, Xia X, Wang HY, Legras X, Hong J *et al.* NLRC5 negatively regulates

the NF-kappaB and type I interferon signaling pathways. *Cell* 2010; **141**: 483–496.

55   Benko S, Magalhaes JG, Philpott DJ, Girardin SE. NLRC5 Limits the Activation of

Inflammatory Pathways. *J Immunol* 2010; **185**: 1681–1691.

56   Davis BK, Roberts RA, Huang MT, Willingham SB, Conti BJ, Brickey WJ *et al.*

Cutting edge: NLRC5-dependent activation of the inflammasome. *J Immunol* 2011;

186: 1333–1337.

57   Wang J quan, Liu Y ru, Xia Q, Chen R nan, Liang J, Xia Q rong *et al.* Emerging roles

for NLRC5 in immune diseases. Front. Pharmacol. 2019; **10**: 1352.

58   Boyd LF, Kozlowski S, Margulies DH. Solution binding of an antigenic peptide to a

major histocompatibility complex class I molecule and the role of β_2-microglobulin.

Proc Natl Acad Sci U S A 1992; **89**: 2242–2246.

59   Kozlowski S, Takeshita T, Boehncke WH, Takahashi H, Boyd LF, Germain RN *et al.*

Excess β_2 microglobulin promoting functional peptide association with purified

soluble class I MHC molecules. *Nature* 1991; **349**: 74–77.

60   Vitiello A, Potter TA, Sherman LA. The role of beta 2-microglobulin in peptide

binding by class I molecules. *Science (80-)* 1990; **250**: 1423–1426.

61   Williams DB, Barber BH, Flavell RA, Allen H. Role of beta 2-microglobulin in the

intracellular transport and surface expression of murine class I histocompatibility

molecules.[erratum appears in J Immunol 1989 Jul 15;143(2):761]. *J Immunol* 1989;

142: 2796–2806.

62   Arce-Gomez B, Jones EA, Barnstable CJ, Solomon E, Bodmer WF. The Genetic

Control of HLA-A and B Antigens in Somatic Cell Hybrids: Requirement for β_2

Microglobulin. *Tissue Antigens* 1978; **11**: 96–112.

63   Biswas A, Meissner TB, Kawai T, Kobayashi KS. Cutting edge: impaired MHC class I

expression in mice deficient for NLRC5/class I transactivator. *J Immunol* 2012; **189**:

516–520.

64   Rodriguez GM, Bobbala D, Serrano D, Mayhue M, Champagne A, Saucier C *et al.*

NLRC5 elicits antitumor immunity by enhancing processing and presentation of tumor

antigens to CD8[+] T lymphocytes. *Oncoimmunology* 2016; **5**: e1151593.

65   Yao Y, Wang Y, Chen F, Huang Y, Zhu S, Leng Q *et al.* NLRC5 regulates MHC class

I antigen presentation in host defense against intracellular pathogens. *Cell Res* 2012;

22: 836–847.

66   Lapinski PE, Miller GG, Tampé R, Raghavan M. Pairing of the nucleotide binding

domains of the transporter associated with antigen processing. *J Biol Chem* 2000; **275**:

6831–6840.

67   Antoniou AN, Ford S, Pilley ES, Blake N, Powis SJ. Interactions formed by

individually expressed TAP1 and TAP2 polypeptide subunits. *Immunology* 2002; **106**:

182–189.

68   Schmitz R, Fitch ZW, Schroder PM, Choi AY, Jackson AM, Knechtle SJ *et al.* B cells

in transplant tolerance and rejection: friends or foes? *Transpl Int* 2020; **33**: 30–40.

69   Nelson BH. CD20 + B Cells: The Other Tumor-Infiltrating Lymphocytes . *J Immunol*

2010; **185**: 4977–4982.

70   Yuen GJ, Demissie E, Pillai S. B Lymphocytes and Cancer: A Love–Hate

Relationship. *Trends in Cancer* 2016; **2**: 747–757.

71   Flies AS, Flies EJ, Fox S, Gilbert A, Johnson SR, Liu G-S *et al.* An oral bait

vaccination approach for the Tasmanian devil facial tumor diseases. *Expert Rev*

Vaccines 2020; **19**: 1–10.

72   Gastaldello A, Sh R, Bailey A, Owen R, Turner S, Kontouli A *et al.* Passage of

transmissible cancers in the Tasmanian devil is due to a dominant, shared peptide

motif and a limited repertoire of MHC-I allotypes. *bioRxiv* 2020.

doi:10.1101/2020.07.03.184416.

73   Selenko-Gebauer N, Majdic O, Szekeres A, Höfler G, Guthann E, Korthäuer U *et al.*

B7-H1 (Programmed Death-1 Ligand) on Dendritic Cells Is Involved in the Induction

and Maintenance of T Cell Anergy. *J Immunol* 2003; **170**: 3637–3644.

74   Dong H, Strome SE, Salomao DR, Tamura H, Hirano F, Flies DB *et al.* Tumor-

associated B7-H1 promotes T-cell apoptosis: A potential mechanism of immune

evasion. *Nat Med* 2002; **8**: 793–800.

75   Butte MJ, Keir ME, Phamduy TB, Sharpe AH, Freeman GJ. Programmed death-1

ligand 1 interacts specifically with the B7-1 costimulatory molecule to inhibit T cell
responses. *Immunity* 2007; **27**: 111–122.

76 Ghebeh H, Tulbah A, Mohammed S, ElKum N, Amer SM Bin, Al-Tweigeri T *et al.*
Expression of B7-H1 in breast cancer patients is strongly associated with high
proliferative Ki-67-expressing tumor cells. *Int J Cancer* 2007; **121**: 751–758.

77 Iwai Y, Ishida M, Tanaka Y, Okazaki T, Honjo T, Minato N. Involvement of PD-L1
on tumor cells in the escape from host immune system and tumor immunotherapy by
PD-L1 blockade. *Proc Natl Acad Sci U S A* 2002; **99**: 12293–12297.

78 Azuma T, Yao S, Zhu G, Flies AS, Flies SJ, Chen L. B7-H1 is a ubiquitous
antiapoptotic receptor on cancer cells. *Blood* 2008; **111**: 3635–3643.

79 Kochan G, Escors D, Breckpot K, Guerrero-Setas D. Role of non-classical MHC class
I molecules in cancer immunosuppression. Oncoimmunology. 2013; **2**: e26491.

80 Flies AS, Blackburn NB, Lyons AB, Hayball JD, Woods GM. Comparative Analysis
of Immune Checkpoint Molecules and Their Potential Role in the Transmissible
Tasmanian Devil Facial Tumor Disease. *Front Immunol* 2017; **8**: 513.

81 Flies AS, Darby JM, Lennard PR, Murphy PR, Ong CEB, Pinfold TL *et al.* A novel
system to map protein interactions reveals evolutionarily conserved immune evasion
pathways on transmissible cancers. *Sci Adv* 2020; **6**: eaba5031.

82 Van Den Elsen PJ, Gobin SJP, Van Eggermond MCJA, Peijnenburg A. Regulation of
MHC class I and II gene transcription: Differences and similarities. *Immunogenetics*
1998; **48**: 208–221.

83 Van Den Elsen PJ, Peijnenburg A, Van Eggermond MC j. a., Gobin SJ p. Shared
regulatory elements in the promoters of MHC class I and class II genes. *Immunol*
Today 1998; **19**: 308–312.

84 Ludigs K, Seguín-Estévez Q, Lemeille S, Ferrero I, Rota G, Chelbi S *et al.* NLRC5
exclusively transactivates MHC class I and related genes through a distinctive SXY
module. *PLoS Genet* 2015; **11**: e1005088.

85 Belov K, Deakin JE, Papenfuss AT, Baker ML, Melman SD, Siddle H V *et al.*
Reconstructing an Ancestral Mammalian Immune Supercomplex from a Marsupial
Major Histocompatibility Complex. *PLoS Biol* 2006; **4**: e46.

86 Siddle H V., Deakin JE, Coggill P, Hart E, Cheng Y, Wong ESW *et al.* MHC-linked
and un-linked class I genes in the wallaby. *BMC Genomics* 2009; **10**: 310.

87 Cheng Y, Belov K. Characterisation of non-classical MHC class I genes in the
Tasmanian devil (*Sarcophilus harrisii*). *Immunogenetics* 2014; **66**: 727–735.

88 Lee S-J, Jang B-C, Lee S-W, Yang Y-I, Suh S-I, Park Y-M *et al.* Interferon regulatory
factor-1 is prerequisite to the constitutive expression and IFN-γ-induced upregulation
of B7-H1 (CD274). *FEBS Lett* 2006; **580**: 755–762.

89 Loke P, Allison JP. PD-L1 and PD-L2 are differentially regulated by Th1 and Th2
cells. *Proc Natl Acad Sci USA* 2003; **100**: 5336–5341.

90 Saijo K, Mecklenbräuker I, Santana A, Leitger M, Schmedt C, Tarakhovsky A. Protein
kinase C β controls nuclear factor κB activation in B cells through selective regulation
of the IκB kinase α. *J Exp Med* 2002; **195**: 1647–1652.

91 Hayden MS, Ghosh S. Signaling to NF-κB. *Genes Dev* 2004; **18**: 2195–2224.

92 Xia L, Tan S, Zhou Y, Lin J, Wang H, Oyang L *et al.* Role of the NFκB-signaling
pathway in cancer. *Onco Targets Ther* 2018; **11**: 2063–2073.

93 Verzella D, Pescatore A, Capece D, Vecchiotti D, Ursini MV, Franzoso G *et al.* Life,
death, and autophagy in cancer: NF-κB turns up everywhere. *Cell Death Dis* 2020; **11**:
1–14.

94 Tong Y, Cui J, Li Q, Zou J, Wang HY, Wang R-F. Enhanced TLR-induced NF-κB
signaling and type I interferon responses in NLRC5 deficient mice. *Cell Res* 2012; **22**:
822–35.

95 Liu Y ru, Yan X, Yu H xia, Yao Y, Wang J quan, Li X feng *et al.* NLRC5 promotes
cell proliferation via regulating the NF-κB signaling pathway in Rheumatoid arthritis.
Mol Immunol 2017; **91**: 24–34.

96 Garrido F, Cabrera T, Concha A, Glew S, Ruiz-Cabello F, Stern PL. Natural history of
HLA expression during tumour development. Immunol. Today. 1993; **14**: 491–499.

97 Hicklin DJ, Marincola FM, Ferrone S. HLA class I antigen downregulation in human
cancers: T-cell immunotherapy revives an old story. Mol. Med. Today. 1999; **5**: 178–
186.

98 Campoli M, Ferrone S. HLA antigen changes in malignant cells: Epigenetic
mechanisms and biologic significance. Oncogene. 2008; **27**: 5869–5885.

 TABLES

859 **Table 1.** Devil facial tumor (DFT) cell lines and treatments

ID #	Sample name	Parent cell line	Treatment
1	DFT1.WT*	DFT1 C5065	Untreated
2	DFT1.WT + IFNG	DFT1 C5065	5 ng/ml IFNG, 24h
3	DFT2.WT[RV]*	DFT2 RV	Untreated
4	DFT2.WT[RV] + IFNG	DFT2 RV	5 ng/ml IFNG, 24h
5	DFT1.BFP	DFT1 C5065	Transfected with control vector pBSbi-BH
6	DFT1.NLRC5	DFT1 C5065	Transfected with NLRC5 vector pCO1
7	DFT2.WT	DFT2 JV	Untreated
8	DFT2.BFP	DFT2 JV	Transfected with control vector pBSbi-BH
9	DFT2.NLRC5	DFT2 JV	Transfected with NLRC5 vector pCO1
10	DFT1.B2M$^{-/-}$	DFT1 C5065	Transfected with *B2M* targeting vector pAF217
11	DFT1.B2M$^{-/-}$ + IFNG	DFT1 C5065	Transfected with *B2M* targeting vector pAF217 and treated with 5 ng/ml IFNG for 24h
12	DFT1.NLRC5.B2M$^{-/-}$	DFT1 C5065	Transfected with NLRC5 vector pCO1 and *B2M* targeting vector pAF218

860 *DFT1.WT data from Patchett et al., 2018[24] and DFT2.WT[RV] from Patchett et al., 2020[6] available
861 through European Nucleotide Archive # PRJNA416378 and # PRJEB28680, respectively.

Table 2. Reactome pathways enriched in differentially expressed genes in DFT1.NLRC5

Reactome ID	Pathway	Count	Term size	p-value	p.adjust	Genes
Upregulated						
R-HSA-1236974	ER-Phagosome pathway	4	74	4.69E-05	4.75E-03	*B2M, PSMB8, PSMB9, TAP1*
R-HSA-1168372	Downstream signaling events of B Cell Receptor (BCR)	4	80	6.37E-05	4.75E-03	*NFATC1, PRKCB, PSMB8, PSMB9*
R-HSA-1236975	Antigen processing-Cross presentation	4	81	6.69E-05	4.75E-03	*B2M, PSMB8, PSMB9, TAP1*
R-HSA-983705	Signaling by the B Cell Receptor (BCR)	4	104	1.77E-04	9.44E-03	*NFATC1, PRKCB, PSMB8, PSMB9*
R-HSA-3858494	Beta-catenin independent WNT signaling	4	129	4.06E-04	1.73E-02	*NFATC1, PRKCB, PSMB8, PSMB9*
R-HSA-1169091	Activation of NF-kappaB in B cells	3	64	7.19E-04	2.55E-02	*PRKCB, PSMB8, PSMB9*
Downregulated						
R-HSA-216083	Integrin cell surface interactions	5	62	1.16E-05	4.04E-03	*CDH1, COL18A1, COL6A1, COL6A2, JAM2*
R-HSA-1251985	Nuclear signaling by ERBB4	3	28	3.33E-04	4.93E-02	*EREG, GFAP, S100B*
R-HSA-5173105	O-linked glycosylation	4	73	4.27E-04	4.93E-02	*ADAMTS7, B3GNT7, GALNT13, GALNT17*
R-HSA-913709	O-linked glycosylation of mucins	3	34	5.96E-04	4.93E-02	*B3GNT7, GALNT13, GALNT17*
R-HSA-8948216	Collagen chain trimerization	3	36	7.06E-04	4.93E-02	*COL18A1, COL6A1, COL6A2*

A cut-off p-value < 0.001 and p.adjust < 0.05 were used to display significant pathways. P-values were adjusted (p.adjust) for multiple testing using Benjamini–Hochberg method. See also Supplementary Table 9 for full list of Reactome pathways.

Table 3. Reactome pathways enriched in differentially expressed genes in DFT2.NLRC5

Reactome ID	Pathway	Count	Term size	p-value	p.adjust	Genes
Upregulated						
R-HSA-1236974	ER-Phagosome pathway	4	74	3.43E-06	3.80E-04	*B2M, PSMB8, PSMB9, TAP1*
R-HSA-1236975	Antigen processing-Cross presentation	4	81	4.93E-06	3.80E-04	*B2M, PSMB8, PSMB9, TAP1*
R-HSA-983169	Class I MHC mediated antigen processing & presentation	5	312	5.96E-05	3.06E-03	*B2M, PSMB8, PSMB9, TAP1, TRIM69*
R-HSA-983170	Antigen Presentation: Folding, assembly and peptide loading of class I MHC	2	18	3.31E-04	1.27E-02	*B2M, TAP1*
R-HSA-162909	Host Interactions of HIV factors	3	119	6.91E-04	1.36E-02	*B2M, PSMB8, PSMB9*
Downregulated						
R-HSA-112316	Neuronal System	33	276	1.73E-06	1.28E-03	*see Supplementary Table 10*
R-HSA-1474228	Degradation of the extracellular matrix	16	97	1.82E-05	6.73E-03	*see Supplementary Table 10*
R-HSA-264642	Acetylcholine Neurotransmitter Release Cycle	5	10	5.87E-05	1.45E-02	*see Supplementary Table 10*
R-HSA-181429	Serotonin Neurotransmitter Release Cycle	5	12	1.70E-04	2.27E-02	*see Supplementary Table 10*
R-HSA-181430	Norepinephrine Neurotransmitter Release Cycle	5	12	1.70E-04	2.27E-02	*see Supplementary Table 10*
R-HSA-112315	Transmission across Chemical Synapses	21	179	1.84E-04	2.27E-02	*see Supplementary Table 10*
R-HSA-1474244	Extracellular matrix organization	24	224	2.65E-04	2.80E-02	*see Supplementary Table 10*
R-HSA-166658	Complement cascade	6	21	4.02E-04	3.38E-02	*see Supplementary Table 10*
R-HSA-1296072	Voltage gated Potassium channels	7	29	4.11E-04	3.38E-02	*see Supplementary Table 10*
R-HSA-2022090	Assembly of collagen fibrils and other multimeric structures	9	49	5.62E-04	4.16E-02	*see Supplementary Table 10*
R-HSA-210500	Glutamate Neurotransmitter Release Cycle	5	16	7.96E-04	4.91E-02	*see Supplementary Table 10*
R-HSA-212676	Dopamine Neurotransmitter Release Cycle	5	16	7.96E-04	4.91E-02	*see Supplementary Table 10*

A cut-off p-value < 0.001 and p.adjust < 0.05 were used to display significant pathways. P-values were adjusted (p.adjust) for multiple testing using Benjamini–Hochberg method. See also Supplementary Table 10 for full list of Reactome pathways.

FIGURES

Figure 1. Upregulation of NLRC5 by IFNG in DFT1 and DFT2 cells. Fold change in
mRNA expression (transcripts per kilobase million (TPM)) of *B2M*, MHC class I gene *SAHAI-*
01, *PDL1* and *NLRC5* upon IFNG treatment in DFT1 C5065 cell line (DFT1.WT) and DFT2
RV cell line (DFT2.WT[RV]). *SOX10* and *NES* were included as internal controls. Bars show the
mean of *N*=2 replicates per treatment. Error bars indicate standard deviation.

Figure 2. Venn diagram of genes significantly upregulated upon IFNG treatment and NLRC5 overexpression in DFT1 and DFT2 cells. Genes were defined as significantly upregulated when false discovery rate (FDR) < 0.05 and $\log_2FC \geq 2.0$. Total number of genes upregulated for each treatment is indicated in parentheses under the sample name. The box shows genes upregulated in all four treatments: (i) IFNG-treated DFT1 cells (DFT1.WT + IFNG), (ii) IFNG-treated DFT2 cells (DFT2.WT[RV] + IFNG), (iii) NLRC5-overexpressing DFT1 cells (DFT1.NLRC5), and (iv) NLRC5-overexpressing DFT2 cells (DFT2.NLRC5). See Supplementary Table 1 for a full list of differentially expressed genes and Supplementary Table 6 for description of devil-specific genes.

Figure 3. Heatmap showing expression profiles of genes involved in MHC-I and MHC-II antigen processing and presentation pathways, and PDL1, in IFNG-treated, and NLRC5-overexpressing DFT1 and DFT2 cells. Log$_2$TPM expression values were scaled across each gene (rows) and represented by Z-Score, with red and blue representing high and low relative expression, respectively. Replicates for each treatment (*N*=2) are included in the heatmap. *SAHAI* encodes the Tasmanian devil MHC-I heavy chain gene. See Supplementary Table 6 for corresponding NCBI gene symbols and description of devil-specific genes.

Figure 4. GO biological processes that were enriched in DFT1 cells with NLRC5 overexpression. GO biological process terms associated with genes upregulated (UP) **(A, B)** and downregulated (DN) **(C)** in DFT1.NLRC5. **(B)** Heatplot of genes associated with each positively-regulated GO term. The cut-offs p-value < 0.001 and adjusted p-value (p.adjust) < 0.05 were used to determine significant biological processes. P values were adjusted for multiple testing using Benjamini–Hochberg method. See also Supplementary Table 7 for full list of GO biological processes.

Figure 5. GO biological processes that were enriched in DFT2 cells with NLRC5 overexpression. GO biological process terms associated with genes upregulated (UP) **(A, B)** and downregulated (DN) **(C)** in DFT2.NLRC5. **(B)** Heatplot of genes associated with each positively-regulated GO term. The cut-offs p-value < 0.001 and adjusted p-value (p.adjust) < 0.05 were used to determine significant biological processes. P values were adjusted for multiple testing using Benjamini–Hochberg method. See also Supplementary Table 8 for full list of GO biological processes.

Figure 6. **Upregulation of MHC-I following NLRC5 overexpression.** Surface expression of B2M in DFT1.NLRC5 **(A)** and DFT2.NLRC5 **(B)**. B2M expression in the NLRC5 cell lines were compared to wild-type (DFT.WT), BFP-control (DFT.BFP), and IFNG-treated (DFT.WT + IFNG) DFT cells. **(A)** Stable expression of B2M in DFT1.NLRC5 was assessed every four weeks for 12 weeks post-drug selection in the presence and absence of hygromycin B (hygB) selection pressure. Secondary antibody-only staining (DFT.WT (2° ab only)) was included as a control. The results shown are representative of N = 3 replicates/treatment.

Figure 7. Flow cytometric analysis of serum antibody binding from devils with anti-DFT1 antibody response. Ablation of surface B2M in CRISPR/Cas9-mediated *B2M* knockout cells (B2M$^{-/-}$) was confirmed using a monoclonal anti-B2M antibody (anti-B2M ab). Sera from six devils (TD1-TD6) with seroconversion (immune) following DFTD infection were tested against wild-type DFT1 (DFT1.WT), IFNG-treated DFT1 (DFT1.WT + IFNG), IFNG-treated *B2M*-knockout DFT1 (DFT1.B2M$^{-/-}$ + IFNG), BFP-control (DFT1.BFP), DFT1 overexpressing NLRC5 (DFT1.NLRC5) and *B2M*-knockout NLRC5-overexpressing DFT1 (DFT1.NLRC5.B2M$^{-/-}$) cells. An immunized devil (My) with MHC-I^{+} antibodies was included as a positive control, meanwhile serum from a healthy devil (TD7) was included as a negative control as represented in the shaded grey area. *Ab*, antibody; *AF488*, Alexa Fluor 488.