Seasonal and long-term consequences of esca on grapevine stem xylem

integrity

Running title: Impact of the trunk disease esca on stem hydraulic integrity

Highlight: Our study reveals that esca can critically affect xylem water movement in grapevine

perennial organs, by the presence of plant-derived tyloses.

G. Bortolami[a], E. Farolfi[a], E. Badel[b], R. Burlett[c], H. Cochard[b], N. Ferrer[a], A. King[d], L.J.

Lamarque[c,e], P. Lecomte[a], M. Marchesseau-Marchal[a], J. Pouzoulet[f], J.M. Torres-Ruiz[b], S.

Trueba[c,g], S. Delzon[c], G.A. Gambetta[f], C.E.L. Delmas[a*]

[a]INRAE, BSA, ISVV, SAVE, 33882 Villenave d'Ornon, France

[b]Université Clermont-Auvergne, INRAE, PIAF, 63000 Clermont-Ferrand, France

[c]Univ. Bordeaux, INRAE, BIOGECO, 33615 Pessac, France

[d]Synchrotron SOLEIL, L'Orme des Merisiers, Gif-sur-Yvette, 91192, France

[e]Département des Sciences de l'Environnement, Université du Québec à Trois-Rivières, Trois-

Rivières, Québec, G9A 5H7, Canada

[f]EGFV, Bordeaux-Sciences Agro, INRAE, Université de Bordeaux, ISVV, 210 chemin de

Leysotte, 33882 Villenave d'Ornon, France

[g]School of Forestry and Environmental Studies, Yale University, New Haven, CT 06511, USA

*Author for correspondence

Chloé E. L. DELMAS

chloe.delmas@inrae.fr

ORCID ID: 0000-0003-3568-605X

ABSTRACT

Hydraulic failure has been extensively studied during drought-induced plant dieback, but its role in plant-pathogen interactions is under debate. During esca, a grapevine (*Vitis vinifera*) disease, symptomatic leaves are prone to irreversible hydraulic dysfunctions but little is known about the hydraulic integrity of perennial organs over the short- and long-term. We investigated the effects of esca on stem hydraulic integrity in naturally infected plants within a single season and across season(s). We coupled direct (k_s) and indirect (k_{th}) hydraulic conductivity measurements, and tylose and vascular pathogen detection with in vivo X-ray microtomography visualizations. We found xylem occlusions (tyloses), and subsequent loss of stem k_s, in all of the shoots with severe symptoms (apoplexy) and in more than 60% of the shoots with moderate symptoms (tiger-stripe), and no tyloses in shoots that were currently asymptomatic. In vivo stem observations demonstrated that tyloses were observed only when leaf symptoms appeared, and resulted in more than 50% PLC in 40% of symptomatic stems, unrelated to symptom age. The impact of esca on xylem integrity was only seasonal and no long-term impact of disease history was recorded. Our study demonstrated how and to what extent a vascular disease such as esca, affecting xylem integrity, could amplify plant mortality by hydraulic failure.

Key words: Esca, hydraulic failure, plant dieback, tyloses, vascular pathogens, *Vitis vinifera* L., X-ray microCT, xylem anatomy

INTRODUCTION

In agricultural and forest ecosystems, perennial plant dieback causes decreases in plant productivity and longevity (Aleemullah and Walsh, 1996; Eskalen *et al*., 2013; Urbez-Torres *et al*., 2013; Alvindia and Gallema, 2017). Plant dieback is a complex process where different biotic and/or abiotic stress factors interact and contribute to leaf and crown wilting and ultimately plant death (Desprez-Lostau *et al*., 2006; Anderegg *et al*., 2013, Cailleret *et al*., 2017; Bettenfeld *et al*., 2020). Drought-mediated plant dieback has been extensively studied, and in this case hydraulic failure has been identified as the primary cause of plant death (Anderegg *et al*., 2016). Hydraulic failure results from an interruption of the ascendant water flow by air embolism or xylem occlusion (Zimmermann, 1979; Tyree and Sperry, 1989). Vascular pathogens, which infect the xylem network (Yadeta and Thomma, 2013), are also important drivers of pathogen-mediated plant dieback (Goberville *et al.*, 2016; Pandey *et al*., 2018; Fallon *et al*., 2020).

Vascular pathogens induce wood necrosis, leaf symptoms, and crown defoliation (Beckmann and Roberts, 1995; Pearce, 1996). Their biology and toxic metabolite production has been well studied, in particular using controlled phytotoxicity assays (Andolfi *et al*., 2011; Akpaninyang and Opara, 2017). However, the possible role of hydraulic failure during pathogen-mediated plant dieback has been poorly investigated, and the underlying physiological mechanisms inducing leaf symptoms are not clear yet (Fradin and Thomma, 2006; McDowell *et al*., 2008). Moreover, the long-term impact (over seasons) and relationships between pathogens, leaf symptom presence, and the hydraulic functioning of the plant are still unknown. During vascular pathogenesis, both air (Pérez-Donoso *et al*., 2016) and nongaseous (Sun *et al*., 2013; Czemmel *et al.,* 2015, Pouzoulet *et al*., 2019) embolism have been observed. For example, air embolism is thought to accelerate pathogen progression during Pierce's disease (Pérez-Donoso *et al*., 2016), and nongaseous embolism is associated with occlusion of the xylem conduits by the plant that could slow the disease process while interfering with xylem water transport (Sun *et al*., 2013; Pouzoulet *et al*., 2019).

Xylem occlusion, usually through the production of tyloses and gels, is one of the first plant defense mechanisms against vascular pathogens (Pearce, 1996). Xylem parenchyma cells secrete gels and expand into the vessel lumen, forming tyloses, physically blocking pathogen progression (Zimmermann, 1979). Xylem anatomy plays an important role, both for vascular pathogen development (Martin *et al*., 2009; Martín *et al*., 2013; Venturas *et al*., 2014;

Pouzoulet *et al*., 2017; 2020) and for tylose formation (Bonsen and Kucera, 1990; De Micco *et*

al., 2016; Pouzoulet *et al*., 2019). If effective, this occlusion mechanism allows the plant to

compartmentalize the infected zone and to generate new tissue around it (CODIT model,

Pearce, 1996). Because tyloses can potentially interfere with the hydraulic functioning of the

plant, they could exacerbate disease symptoms (Talboys, 1972). Tyloses are usually observed

in close proximity to pathogens, as shown in artificial inoculation studies (Czemmel *et al*.,

2015; Rioux *et al*., 2018, among others). However, pathogens frequently proliferate in

perennial organs without physically reaching the leaves, thus leaf symptoms are often induced

at a distance (Beckmann and Roberts, 1995). A recent study shows that tyloses can be present

in symptomatic leaves at a distance from the pathogen niches resulting in decreased leaf

hydraulic conductivity (Bortolami *et al*., 2019).

Over the last decades, grapevine (*Vitis vinifera* L.) mortality and yield loss have been reported

in European, American, and South African vineyards due to esca trunk disease (Cloete *et al*.,

2015; Guerin-Dubrana *et al*., 2019). Esca, a vascular disease caused by the infection of multiple

fungal pathogens, affects mostly mature grapevines (more than seven-years-old), and

symptoms include trunk necrosis and leaf symptoms, consisting of "tiger-stripe" necrosis and

leaf wilting (Lecomte *et al*., 2012; Claverie *et al*., 2020), which are not regularly expressed

season-to-season even within individual vines (Guerin-Dubrana *et al*., 2013; Li *et al*., 2017).

While the pathogens responsible for esca-induced trunk necrosis have been identified

(Morales-Cruz *et al*., 2018; Brown *et al*., 2020), the underlying mechanisms of leaf and fruit

symptoms, and plant death are still poorly understood. Bortolami *et al*. (2019) demonstrated

that the two vascular pathogens related to esca (*Phaeomoniella chlamydospora* and

Phaeoacremonium minimum) were never detected in leaves or in stems of the current year, but

always in the trunk (independently from leaf symptom presence). They further showed that

esca symptomatic leaves presented significant losses in hydraulic conductivity due to the

occlusion of the xylem conduits by tyloses. Together, these results reveal that esca impacts leaf

hydraulic functioning, but whether or not there is a corresponding failure in perennial organs,

and the exact timing of this phenomenon, are still unknown. As stems and branches are the

direct connections between the pathogen niche in the trunk and the observed symptoms in the

leaves, the study of stem xylem integrity is crucial in the understanding of esca impact on

grapevine physiology in the current year and across seasons.

In this study, we investigated stem xylem integrity in grapevine during esca leaf symptom
formation asking the following questions: (i) Can esca lead to hydraulic failure in perennial
organs? (ii) Does stem hydraulic failure occur prior to or after leaf symptom expression, and
does it depend on xylem anatomy? (iii) Do long-term symptomatic plants present different
xylem anatomy and levels of hydraulic failure from long-term asymptomatic plants? To answer
these questions, we transplanted 28-years-old grapevines (*Vitis vinifera* L. cv Sauvignon blanc)
from the field into pots to transport, manipulate, and study naturally esca-infected vines. We
coupled *in vivo* visualizations of stem xylem functionality (using synchrotron-based X-ray
microcomputed tomography) with stem specific hydraulic conductivity measurements (k_s),
theoretical hydraulic conductivity estimates (k_{th}), optical observations of vessel occlusions, and
pathogen detection during symptom appearance, while comparing plants with different
symptom history record.

MATERIALS AND METHODS

Plant material

Vitis vinifera cv. Sauvignon blanc grafted onto 101-14 MGt were uprooted in winter 2017,
2018, and 2019 from a vineyard planted in 1992 located at INRAE Bordeaux-Nouvelle
Aquitaine (44°47'24.8"N, 0°34'35.1"W) and transferred into pots. Following plant excavation,
the root system (around 0.125 m3) was immersed under water overnight, and powered with
indole-3-butyric acid. The plants were potted in 20l pots in fine clay medium (Klasmann
Deilmann substrate 4:264) and placed on heating plates at 30 °C for two months. Plants were
then moved to a greenhouse, under natural light conditions, and watered with nutritive solutions
(0.1 mM NH4H2PO4, 0.187 mM NH4NO3, 0.255 mM KNO3, 0.025 mM MgSO4, 0.002 mM
Fe, and oligo-elements [B, Zn, Mn, Cu, and Mo]) until the end of experiment. Since the
plantation these plants have been trained with a double Guyot system. This training system
requires a permanent main trunk and one cane on each side of the trunk which is left every year
to carry the buds that will produce the stems of the year. During the growing season, the stems
of the current year were trimmed at 1.5-2m, the secondary stems and inflorescences were
removed just after bud-break. Each of these plants has been surveyed each year in the field
since 2012 for esca leaf symptom expression following Lecomte *et al*. (2012), and has been
classified yearly as leaf-symptomatic or asymptomatic. Plants were then classified by their

long-term symptomatology record: plants asymptomatic from 2012 to 2018 (pA, previously asymptomatic), and plants that have expressed symptoms at least once between 2012 and 2018 (pS, previously symptomatic).

Esca symptom notation

The evolution of esca leaf symptoms was surveyed twice a week from June to October 2019 on every plant (n=84, Fig. 1). As presented in Fig. 1A, esca symptoms were scored at the stem and whole plant scales. The stems of the current year collected for analyses (both hydraulic measurements or microCT observations) could be noted as: asymptomatic (green leaves and apparently healthy), pre-symptomatic (leaves presenting yellowing or small yellow spots between the veins), tiger-stripe (typical pattern of esca leaf symptoms), or apoplectic (leaves passing from green to wilted in a couple of days). Along the experimentation, entire plants could be noted as asymptomatic (control) or symptomatic (when at least 25% of the canopy was presenting tiger-stripe leaf symptoms). At the end of the experiment (week 40, October 2019) each plant was classified as symptomatic or asymptomatic (control). We were then able to group each stem measured into six different groups (Fig. 1A): one group of stems from control plants (asymptomatic from June to October) and five groups of stems from symptomatic plants: two before symptom appearance (asymptomatic and pre-symptomatic stems); and three after symptom appearance (asymptomatic, tiger-stripe, and apoplectic stems). To clearly differentiate asymptomatic stems collected from symptomatic plants and asymptomatic stems collected from asymptomatic plants, we considered plants (and their stems) that didn't show leaf symptoms during the experiment as control plants (or stems). We investigated whether symptom expression (final symptom notation in October 2019, see Fig. 1) differed between plants with contrasted long-term symptom history (previously asymptomatic vs previously symptomatic, Table 1) using a Chi-square test of independence.

X-ray microCT observation

Synchrotron-based microCT was used to visualize the content of vessels and their functionality in esca tiger-stripe and control stems. Three symptomatic plants (presenting tiger-stripe symptoms for 8, 7, and 3 weeks), and one asymptomatic-control plant were brought to the PSICHE beamline (King *et al*., 2016) at SOLEIL synchrotron facility in September 2019. Stems of the current year (ca. 2 m long) were cut under water and transferred into a solution

containing 75mM of contrasting agent iohexol. The iohexol solution absorbs X-rays very

strongly and appears bright white in X-ray scans above the iodine K-edge at 33.2 keV, and,

once it has been taken up by the transpiration stream, the effective functionality of each vessel

can be confirmed (Pratt and Jacobsen, 2018; Bortolami *et al.*, 2019). These stems were moved

and left outdoor to transpire the solution for at least half a day. The stems were then transferred

to the beamline stage and scanned twice in less than 5 minutes using two different energies of

a high-flux (3×10^{11} photons mm^{-2}) monochromatic X-ray beam: 33.1 keV and 33.3 keV. The

projections were recorded with a sCMOS camera equipped with a 250-mm-thick LuAG

scintillator (Orca Flash, Hamamatsu, Japan). The complete tomographic scan included 1500

projections, and each projection lasted 50 ms. Tomographic reconstructions were performed

using PyHST2 software (Mirone *et al.*, 2014) using the Paganin method (Paganin *et al.*, 2002),

resulting in 32-bit volume reconstructions of 2048 x 2048 x 1024 voxels. The final spatial

resolution was 2.8769 μm^3 voxel^{-1}.

Image analysis of microCT scans

The contrast agent iohexol allowed us to distinguish in intact scans the effective functionality

of each vessel. In the absence of iohexol, X-ray microCT scans are used to distinguish air-filled

vessels (appearing black, corresponding to native PLC) from sap-filled vessels (appearing

grey). The addition of iohexol in the xylem sap allows to distinguish the functional vessels

(they appear bright white when they transport the sap), from the non-functional ones (i.e.

occluded vessels remaining grey, corresponding to occlusion PLC). We could also observe

partially occluded vessels (i.e. vessels with simultaneous presence of air and occlusions, or sap

and occlusions). This specific case was observed by checking the presence of any occlusion in

at least 200 slices in each volume. Partially occluded vessels were considered as occluded,

some examples are presented in Fig. S1. The equivalent-circle diameter of air-filled, occluded,

and functional (iohexol-filled) vessels was measured on the cross sections from the central slice

of the microCT scanned volume using ImageJ software (Schneider *et al.*, 2012). In the high

energy scans recorded at 33.3 keV X-ray beam, iohexol appears bright white but its contrast

can sometimes impede the clear limit of the vessel lumen. Therefore, all vessel diameters were

recorded on the scan recorded at low energy (33.1 keV X-ray beam), then the distinction of

occluded from iohexol-filled vessels was done on the high energy scan (as done by Bortolami

et al. 2019). The theoretical hydraulic conductivity of each vessel (k_{vessel}) [kg m MPa^{-1} s^{-1}] was

calculated using the Hagen-Poiseuille equation:

204
$$k_{vessel} = \frac{(\pi \times \O^4 \times \rho)}{(128 \times \eta)}$$

205 Where: $\O$ is the equivalent circle diameter [m], ρ the density of water [998.2 kg m-3 at 20°C],
206 and η the viscosity of water [1.002 x 10-9 MPa s at 20°C]. The percentage loss of hydraulic
207 conductivity given by native air embolism (native PLC) was calculated by the ratio between
208 the hydraulic conductivity of air-filled vessels and the whole-stem hydraulic conductivity:

209
$$\text{Native PLC (\%)} = 100 \times \frac{(\sum k_{air-filled\ vessels})}{(\sum k_{all\ vessels})}$$

210 The percentage loss of hydraulic conductivity given by occlusions (occlusion PLC) was
211 calculated by the ratio between occluded (plus partially occluded) vessels and the whole-stem
212 hydraulic conductivity:

213
$$\text{Occlusion PLC (\%)} = 100 \times \frac{(\sum k_{occluded\ vessels} + \sum k_{partially\ occluded\ vessels})}{(\sum k_{all\ vessels})}$$

The total percentage loss of hydraulic conductivity (total PLC) was obtained by summing
native PLC with occlusion PLC in each sample. As the first ring of xylem vessels (i.e.
protoxylem) was always non-functional (>90% PLC), both in control and tiger-stripe stems, it
was removed from the analysis.

We investigated whether native PLC, and occlusion PLC differed between control and esca
tiger-stripe plants, using two independent generalized mixed linear models where plants were
treated as a random effect. Proportional data (ranging from 0 to 1, dividing all PLC values by
100) was analyzed to fit a logit link function and binomial distribution as appropriate.

Monitoring stem hydraulic properties over time

Xylem integrity was monitored over time by measuring hydraulic properties in stems produced
on the year of the experiment and collected on control and symptomatic plants along the season
and during esca development. Specific hydraulic conductivity (k_s) was measured on internodes
sampled in the center of the collected stem by the gravity method (Sperry *et al*., 1988), and

compared to its theoretical analog (k_{th}) calculated from xylem anatomical observations on the

same internode or on the one below (see the method described below). When there are observed

differences in k_s among stems, comparisons with theoretical maximums (k_{th}) can show if lower

k_s values result from anatomical differences (i.e. different vessel size distributions) or by

hydraulic failure (in the case of similar vessel size and density). If k_s varies in unity with k_{th},

differences in k_s might result from anatomical differences (e.g. smaller k_s are related to smaller

vessels), otherwise k_s variations are the consequence of hydraulic failure. Each method to

measure k_s, k_{th}, and to observe tyloses is described below.

Sampling started on June 19[th] and finished on September 13[th] 2019 for a total of 10 sampling

dates, 39 stems of the current year from 23 control-asymptomatic plants, and 49 stems of the

current year from 17 symptomatic plants. We randomly sampled control plants and esca

symptomatic plants all along the season through the evolution of esca symptoms, obtaining

measurements from 14 weeks before until 10 weeks after symptom appearance. To explore the

contribution of the experimental design to data analysis, we tested the effect of the year of

uprooting (2018 and 2019), the position of the analyzed internode, and the week of the

measurement (i.e. evolution during the season) on k_s and k_{th} in control plants using separate

generalized linear mixed model with normal distributions and the plant treated as a random

variable (Table S1). A significant impact of the year of uproot was found for k_s and k_{th} values

in control plants (Table S1). This could have resulted from the more favorable conditions (i.e.

climatic stability and nutrient availability) for the greenhouse grown vines (note that plants

uprooted in 2017 were only esca symptomatic and were not included in this analysis). However,

once k_s and k_{th} are plotted together (Fig. S2), all the values lie on the same regression line

without generating outlier values (smaller k_s values correspond to smaller k_{th} values

independently of the uprooting year).

Stem specific hydraulic conductivity (k_s)

k_s measurements were performed on one internode per stem, located in the center of the

collected >1.5m long stem, following Torres-Ruiz *et al*. (2012) gravity method. In the early

morning, each stem was cut at the base under water to avoid air entrance in the stem, maintained

under water and brought to the laboratory. Hydraulic conductivity measurements were always

done before noon, in order to minimize the delay (never more than four hours) from the cut to

the measure. In the laboratory, a representative internode between the 4th to the 10th internode

from the base (i.e. in the center of the stem) was cut underwater with a clean razor blade, the

ends wrapped in tape, and the internode was connected to a pipe system. A flow of 20 mM KCl solution passed through the sample from a reservoir to a precision electronic balance (AS220.R2, RADWAG, Radom, PL) recording the weight every 5 seconds using the WinWedge v3 5.0 software (TAL Technologies, Philadelphia, PA, USA). The solution was passed through the stem at four increasing pressures (ranging from 0.001 to 0.005 MPa), controlled by raising the source height. The average flow for each pressure step was determined after stabilization at a steady-state as the average of 10-15 measures. Hydraulic conductance, k [kg s^{-1} MPa^{-1}] was obtained by the slope generated by the flow and the corresponding pressure. The linear relationship between flow and pressure obtained were always characterized by $R^2 > 0.97$. Stem specific hydraulic conductivity, k_s [kg s^{-1} MPa^{-1} m^{-1}], was calculated as follows:

$$k_S = \frac{(k \times l)}{A}$$

$$A = \left(\left(\frac{d_1}{2}\right)^2 \times \pi\right) - \left(\left(\frac{d_2}{2}\right)^2 \times \pi\right)$$

Where: k is the hydraulic conductance, l is the length of the sample, A is the xylem area, d_1 is the external diameter of the debarked stem, and d_2 is the diameter of the central pith.

Stem theoretical hydraulic conductivity (k_{th}), vessel anatomy, and tylose observation

Just before hydraulic conductivity (k_s) measurements, the lower internode was stored at 4 °C in 80% ethanol for analysis of xylem anatomy. When possible, the same internode of k_s measurements was used for anatomical analysis and k_{th} estimations, otherwise the stored internode was used for the following protocol. 50 μm thick slices were obtained using a GSL-1 microtome (Gärtner *et al.*, 2014). Slices were stained using a 0.5% safranin solution during 5 minutes, and then washed three to four times in ethanol (100%). They were quickly soaked in xylene and mounted on microscope slides with Permount Mounting Medium (Electron Microscopy Science, Hatfield, PA, USA). Images were captured with a stereo microscope SMZ1270 (Nikon, France) mounted with a DS-Fi3 camera (Nikon, France). The theoretical conductivity of each vessel (k_{vessel}) [kg m Mpa^{-1} s^{-1}] was calculated using the Hagen-Poiseuille equation as described above.

Where $\varnothing$ is the equivalent circle diameter [m] (measured with ImageJ software), ρ the density of water [998.2 kg m^{-3} at 20 °C], and η the viscosity of water [1.002 x 10^{-9} MPa s at 20 °C]. k_{th}

289   of the stem [kg s^{-1} m^{-1} Mpa^{-1}] was then calculated by summing every k_{vessel} in the xylem area

290   (*A*) [m²]:

291
$$k_{th} = \frac{\sum k_{vessel}}{A}$$

In the entire cross section of each sample, the physical presence (or absence) of tyloses in vessel

lumina was visually assessed.

Regarding the statistical analysis, stems were grouped in six different categories following their

esca symptomatology (as presented in Fig. 1A). We investigated whether k_s, k_{th}, and total vessel

density differed among these different categories, and how k_s, k_{th}, and total vessel density

differed between stems with and without tyloses (independently from leaf symptom presence),

using independent mixed linear general models. The symptom / tylose category and the year

of uprooting (since it had a significant impact on k_s and k_{th} in control plants, Table S1) were

entered as fixed effects, with the plant treated as a random effect since different stems were

sometimes analyzed from the same plant (88 analyzed stems on 40 different plants). Total

density and densities for each vessel diameter class were log-transformed prior to analysis to

fit normality requirements. For the classes with no vessels (e.g. samples without vessel

diameters above 160 μm), a minimal density of 0.0001 was assigned prior to log

transformation. We investigated whether the frequency of symptomatic stems presenting

tyloses changed with the symptom age (i.e. weeks between first symptom detection and k_s

measurements on the same plant) with a Chi-square test. The relationships between stem k_s and

k_{th} were tested using linear regression models. Finally, we investigated whether k_s and k_{th} in

control stems differed between plants with different symptom history records using

independent mixed linear general models with the plant treated as a random effect.

Fungal detection

313   Detection and quantification of *Phaeomoniella chlamydospora* and *Phaeoacremonium*

314   *minimum* were performed using qPCR in a subsample of stems of the current year (n = 28) and

315   perennial trunks (n = 20 plants) from the same symptomatic and control plants used for

316   hydraulic and anatomical measurements. All along the season, basal internodes, from the same

317   stems sampled for k_s and k_{th} measurements, were directly placed in liquid nitrogen and stored

at -80 °C. At the end of the experiment, a subset of plants was cut at the base for trunk sampling. A 2 cm high section was cut with a sterilized hand saw. The bark was removed and the different tissues of each section (necrotic and apparently healthy wood) were separately collected using ethyl alcohol-sterilized shears in a sterile environment, and immediately placed in liquid nitrogen. All samples were ground in liquid nitrogen using a tissue lyser (Tissuelyser II, Qiagen, Germantown, MD, USA). DNA was extracted from 60mg of ground tissue using the Invisorb Spin Plant Mini Kit (Invitek GmbH, Berlin, Germany)according to the manufacturer's instructions. . Detection and quantification of *P. chlamydospora* and *P. minimum* (previously named *P. aleophilum*) DNA by qPCR (SYBR Green assays) was conducted using the primer sets PchQF (5'-CTCTGGTGTGTAAGTTCAATCGACTC-3')/PchQR (5'-CCATTGTAGCTGTTCCAGATCAG-3') and PalQF(5'-CCGGTGGGGTTTTTACGTCTACAG-3')/ PalQR(5'-CGTCATCCAAGATGCCGAATAAAG-3') (Pouzoulet *et al.* 2013). The qPCR reactions proceeded in a final volume of 25 µl, and the reaction mixtures containing 2 µL of DNA template, 12.5 µl of 2X SYBRGreen Quantitect Master Mix (Qiagen, Venlo, Netherlands), and each primer at a final concentration of 0.4 µM. Experiments were conducted with a Mx3005P Real-Time PCR cycler using MxPro qPCR software (Agilent Technologies). The cycling program, as described in Pouzoulet *et al.* (2017), consisted of an initial denaturation step at 95°C for 15 min, and 40 cycles of 15 s at 95°C (for denaturation) followed by 45 s at 62°C (for both annealing and extension). A melting analysis of 40 min from 60 to 95° was performed to verify reaction's specificity and the absence of byproducts. Preparation and use of standard solutions for the absolute quantification of fungal DNA was realized following Pouzoulet *et al.* (2013) using ten-fold dilutions of fungal DNA extracts obtained from axenic cultures. Reaction efficiencies ranging from 90% and 95% with an $R^2 > 0.99$ (n=15) were obtained for both PchQF/R and PalQF/R primer sets. The average amount of DNA was determined based on three technical replicates (standards and plates) with a detection threshold superior to 95% (i.e. at least three positive amplification out of three replicates) or otherwise discarded (i.e. pathogen DNA was considered absent). Pathogen DNA quantity (average value of three technical replicates, fg/µl) was normalized by the amount of total DNA (ng/µl), measured using a Qubit fluorometer. The results from each trunk sample (i.e. necrotic or apparently healthy wood) were averaged together in order to obtain one quantification per plant. We investigated whether the amount of fungal DNA (both for *P. chlamydospora* and for *P. minimum*) in trunks differed between symptomatic and control plants, and between control plants with different symptom

history records, using generalized linear mixed model with a poisson distribution and a log
likelihood function.

Statistical analysis

All data management and statistical tests were done in SAS software (SAS 9.4; SAS Institute).
We used PROC GLIMMIX for generalized linear mixed models, PROC GLM for generalized
linear models, PROC REG for regression analyses and PROC FREQ for frequency analyses
(Chi-square test of independence). The normality of the response variables was tested using a
Kolmogorov-Smirnov test (PROC UNIVARIATE) prior to analyses. Data were log-
transformed (total density) or appropriate distributions (binomial, poisson) were fitted when
appropriate.

RESULTS

Esca leaf symptom expression within and across seasons

Esca leaf symptoms were recorded in 20 out of the 58 plants followed in this study (35%, Fig.
1, Table 1). The number of symptomatic plants increased gradually with time, from the first
symptom appearance in early June to the last in late September (Fig. 1). There was no effect of
the plant history (previously asymptomatic pA, or previously symptomatic pS) on 2019
symptom expression (n=58, X^2=0.27, P=0.60). On 20 pA plants, six (30%) expressed leaf
symptoms in 2019 (Table 1). On 38 pS plants, fourteen (37%) showed symptoms in 2019
(Table 1). However, pS plants expressed symptoms from June to the end of September, while
pA plants showed leaf symptoms only in September.

In vivo **observations of esca symptomatic stems**

Xylem vessels of control and tiger-stripe stems were observed using three dimensional X-ray
microCT scans in iohexol-fed samples (Fig. 2, 3, Table S2). As shown in Fig. 2, functional and
non-functional vessels can be discriminated through the use of iohexol (functional vessels
appear bright white, non functional vessels appear either black if air-filled or grey if occluded).

We observed almost totally functional stems in all asymptomatic stems (<20% total PLC, Fig. 2A-C), and 40% of tiger-stripe stems (e.g. Fig. 2D-G). Higher levels of PLC (>20% total PLC, Fig H-M) were observed in the remaining tiger-stripe stems, with 40% of tiger-stripe stems exhibiting over 50% total PLC (Fig. 2J-M). When the two components of PLC were disentangled, we observed that the level of native PLC remained low both in control (6.5 ± 2.6%) and in tiger-stripe (12.2 ± 2.9%) stems (Fig. 3A). Occlusion PLC values were virtually zero in control stems (0.7 ± 0.02%) while in tiger-stripe stems the mean occlusion PLC values was 27.5 ± 8.2% (Fig. 3B). Nevertheless, the variability of occlusion PLC across tiger-stripe stems was very high, the values ranging from 0.3% to 72.9% (Fig. 2D-M, and 3B), and occlusion PLC was not correlated to symptom age (n=10, $F_{2,7}$=0.19, P=0.83). Consequently, no statistical differences in native or occlusion PLC were found between control and tiger-stripe stems (Fig. 3). When higher occlusion PLC was measured (Fig. 2H-M), occluded vessels could be organized either on one side of the stem (Fig 2J-L) or randomly distributed across the section (Fig 2H, 2I, 2M). In 90% of symptomatic stems, we observed that the most external vessels were functional. Occlusions were present equally in all vessel diameter classes (Fig. S3).

Tylose development, stem specific (k_s) and theoretical (k_{th}) hydraulic conductivity during esca leaf symptom formation

Tyloses were identified in the xylem vessels of certain tiger-stripe stems and throughout the temporal development of esca leaf symptoms, from the appearance of symptoms to 11 weeks after. All apoplectic stems and 62.5% (15 of 24 analyzed stems) of esca tiger-stripe stems presented tyloses, while all other stems (control, asymptomatic or pre-symptomatic) did not contain these occlusions, even until one week before symptom development. In esca tiger-stripe stems, tyloses were not related to specific plants, or to symptom age (i.e. on the same plant at the same moment, different symptomatic stems could present tyloses, or not, n=24, X^2=7.47, P=0.38).

Overall, no significant impact of esca symptoms was observed on k_s (Fig. 4A), even if tiger-stripe stems were divided between those with and without tyloses. Control stems presented a mean (± SE) k_s of 24.97 ± 1.72 kg s^{-1} MPa^{-1} m^{-1}; all the stems without tyloses measured on symptomatic plants showed the same range of values as control stems (Fig 4A, Table 2): 26.04

± 4.71 for asymptomatic before symptoms appearance, 30.32 ± 4.26 for pre-symptomatic

stems, 19.80 ± 5.18 for asymptomatic stems after symptom appearance on the plant, and 21.29

± 5.40 for tiger-stripe stems without tyloses. Stems with tyloses (tiger-stripe and apoplectic

stems) presented the lowest average k_s values (11.27 ± 2.86 and 2.47 ± 1.45 kg s^{-1} MPa^{-1} m^{-1}

for tiger-stripe and apoplectic, respectively). Regarding k_{th}, no significant impact of esca

symptoms was found (Fig. 4C, Table 2), all the values were in the same range, with average

values ranging from 70.44 (for tiger-stripe stems with tyloses) to 87.88 (for pre-symptomatic

stems) kg s^{-1} MPa^{-1} m^{-1}.

In order to further investigate the impact of esca on stem hydraulics, we explored the

relationship between individual stem k_s and k_{th} in each symptom category (Fig. 4B, S4, Table

2). Significant relationships were found between k_s and k_{th} in all groups except in asymptomatic

stems after symptom appearance and symptomatic stems with the physical presence of tyloses

(Fig. S4, Table 2). The slopes of regression curves between k_s and k_{th} did not vary among

groups in the absence of tyloses (slope values ranged between 0.3 and 0.4, Table 2) while it

was close to 0 in the presence of tyloses (0.17 for tiger-stripe and 0.04 for apoplectic stems).

When k_s and k_{th} are compared in the presence or absence of tyloses, we observed that k_s was

significantly lower when tyloses were present (9.81 ± 2.51 kg s^{-1} MPa^{-1} m^{-1} in the presence of

tyloses vs 25.06 ± 1.46 kg s^{-1} MPa^{-1} m^{-1} in the absence of tyloses, Table 2, n=88, $F_{1,49}$=7.11,

P=0.01) while k_{th} did not significantly differ. Stems without tyloses presented a strong

correlation between k_s and k_{th}, while in the presence of tyloses this relationship was not

significant (Table 2, Fig. 4B).

Total vessel density did not significantly differ between stem symptomatology (comparing all

the seven categories presented in Table 2), even when vessel density was partitioned by vessel

diameter classes (Fig. 4D).

Finally, we tested the impact of disease history (comparing pA and pS plants) on the hydraulic

conductivity and xylem anatomy in control plants. There were no differences between long-

term symptomatic (pS) and long-term asymptomatic (pA) plants in stem k_s, stem k_{th}, or total

vessel density (Table 3).

Fungal detection

The two vascular pathogens associated with esca (*Phaeomoniella chlamydospora* and *Phaeoacremonium minimum*) were never detected in stems of the current year while they were systematically detected in the perennial trunk of both control and symptomatic plants (Table 4). In trunks, a significantly higher quantity of fungal DNA was detected in tiger-stripe symptomatic plants than in controls (Table 4). We found 2.14- and 1.64-fold more of *P. chlamydospora* and *P. minimum* DNA in symptomatic trunks relative to controls. In control plants, different symptom history records impacted the quantity of fungal DNA detected by qPCR, for *Phaeomoniella chlamydospora*, and for *Phaeoacremonium minimum*. We found 1.65- and 2.84-fold more *P. chlamydospora* and *P. minimum* DNA in previously symptomatic trunks relative to previously asymptomatic trunks (Table 3).

DISCUSSION

Our results regarding the impact of esca on stem xylem integrity show that the presence of plant-derived tyloses induced hydraulic failure in 60% of symptomatic stems of the current year. Tyloses were only observed in symptomatic stems, and resulted in more than 50% PLC in 40% of the stems, unrelated to symptom age. We demonstrated that the presence of leaf symptoms during previous seasons had no impact on the likelihood of symptom appearance in the current year, or on stem hydraulic conductivity and xylem anatomy. Vascular fungi were never detected in the same organs as the tyloses (stems of the current year), and although they were present in trunks of both tiger-stripe and control plants, tiger-stripe plants showed higher quantities of fungal DNA. Among control plants that did not express symptoms in the year of the study, we found higher quantities of fungal DNA in trunks of those plants with a long-term history of symptom formation. Albeit xylem occlusions were not observed in the totality of tiger-stripe stems, they could amplify yield loss plant mortality, especially in the context of climate change as they impair water transport in a majority of symptomatic stems.

In vivo xylem integrity observations and hydraulic vulnerability segmentation

Using direct X-ray microCT imaging in esca symptomatic stems, we found that hydraulic conductivity loss was almost entirely associated with the presence of tyloses. Different studies have investigated the link between vascular pathogen development and hydraulic conductivity

in stems (Collins *et al*., 2009; Lachenbruch and Zhao, 2019; Mensah *et al*., 2020). During biotic

stresses, air embolisms have been shown to decrease hydraulic conductivity during bacterial

leaf scorch disease (McElrone *et al*., 2003; 2008), Pierce's disease (Pérez-Donoso *et al*., 2016),

and Pine wilt disease (Yazaki *et al*., 2018). In the case of fungal wilt diseases, the hydraulic

conductivity loss was associated with nongaseous embolism (i.e. tyloses) at the point of

pathogen inoculation (Guerard *et al*., 2000; Sallé *et al*., 2008; Beier *et al*., 2017; Mensah *et al*.,

2020), or with canker presence in naturally infected stems (Lachenbruch and Zhao, 2019).

Using iohexol we were able to visually observe the exact spatial organization of functional

vessels. Interestingly, in some symptomatic samples we found functional vessels surrounding

the non-functional xylem (Fig. 2J-L), suggesting that the plant was able to preserve the more

external vessels from occlusions or to form new functional vessels after the loss of

conductivity. Moreover, the sectoriality of the occlusions observed in Fig. 2J-L was

reminiscent of the sectoriality observed in the distributions of trunk necrosis, especially on the

brown stripe necrosis appearing along the vasculature (Lecomte *et al*., 2012).

Comparing these results with our precedent study using the same technique in leaves, we

showed that esca symptomatic leaves presented higher levels of occlusion PLC (61 ± 7% in

midribs, and 54 ± 9% in petioles, data from Bortolami *et al*., 2019) compared to stems (27 ± 8

%, occlusion PLC), suggesting hydraulic vulnerability segmentation (although PLC in leaves

and stems were measured in different plants and years). The hydraulic segmentation theory

relies on the fact that annual organs (i.e. leaves) are more vulnerable than perennial organs (i.e.

stems) to drought induced air embolism (Tyree and Ewers, 1991). Grapevine is well known for

exhibiting strong hydraulic vulnerability segmentation (Charrier *et al*., 2016; Hochberg *et al*.,

2016; 2017). This is thought to be adaptive, where the higher vulnerability in leaves and

petioles favors embolism formation and leaf shedding prior to embolism formation in stems,

thus protecting the perennial organs. Our observations during esca pathogenesis demonstrate

that, analogous to the hydraulic vulnerability segmentation theory, leaves appear more

vulnerable to the formation of nongaseous embolism as well, which could mitigate the risk of

hydraulic failure in perennial organs. From another perspective, the difference may not be a

direct effect of the specific organ's vulnerability to nongaseous embolism, but a consequence

of a difference in the accumulation of putative toxins and/or elicitors. Indeed, we confirmed

here that esca leaf symptoms occur at a distance from the pathogen niche because vascular

pathogens were never detected in stems of the current year, suggesting that the plant may

transport a signal (i.e. toxins or elicitors) from the infected trunk up to the leaves. If the signal
accumulates in leaves in a higher amount than it does in the stems (water potentials are more
negative in leaves compared to stems), and stimulates occlusion formation, stems would then
be secondarily affected.

Hydraulic conductivity, tyloses, and vessel anatomy

Tyloses could have different impacts, both positive and negative, during wilt disease
pathogenesis: (i) tyloses contribute to pathogen resistance as they aim to seal off vessel lumens
and impede pathogens spread throughout the host (CODIT model, Shigo, 1984). This is the
case regarding the susceptibility of different species or varieties to specific pathogens (Jacobi
and MacDonald, 1980; Ouellette *et al*., 1999; Clérivet *et al.,* 2000; Et-Touil *et al*., 2005;
Venturas *et al*., 2014; Park and Juzwik 2014; Rioux *et al*., 2018), in particular to *Phaeomoniella*
chlamydospora, one of the pathogen associated with esca (Pouzoulet *et al*., 2017; 2020). (ii) In
other studies, it has been shown that tyloses can exacerbate symptoms (Talboys, 1972): they
cause a reduction in stem hydraulic conductivity, sometimes associated with a reduction in
stomatal conductance in leaves and, in the most severe cases, wilting (Parke *et al*., 2007; Beier
et al., 2017; Lachenbruch and Zhao, 2019, Mensah *et al*., 2020 during fungi development; Sun
et al., 2013; Deyett *et al*., 2019 during Pierce's disease). Our results suggest that during esca
tyloses might lead to symptom exacerbation. Esca has also been suggested to lead to a general
reduction in xylem water transport and stomatal conductance (Ouadi *et al*., 2019), and tyloses
could be a major contributor to these phenomena as during winter senescence (Salleo *et al*.,
2002; Sun *et al*., 2008). However, when symptomatic stems have no tyloses (~37% of the stems
with tiger-stripe symptoms), esca leaf symptom formation seems to arise from within the leaf
itself, and may not result from upstream hydraulic failure. Although tyloses were never
detected in asymptomatic stems prior to the onset of leaf symptoms, the time sequence of tylose
and leaf symptom development has still to be determined. Since both the microCT and
anatomical observations visualize relatively narrow regions of the stems, tylose presence could
have been underestimated (i.e. if there was additional tylose development up or downstream of
the stem sections visualized). However, it should be pointed out that if significant
underestimation were present we would expect some loss of conductivity even in internode
sections from which we observed no tyloses in the sampled cross sections. At least when

considering a single internode our direct hydraulic conductivity measurements do not support

the hypothesis that tyloses were underestimated (Fig. 4B).

Xylem is the battleground between vascular pathogens and the plant's defense response

(Yadeta and Thomma, 2013). Even if xylem vessel anatomy is less investigated, it could have

a crucial role in plant resistance and response to vascular pathogens. For example, during Dutch

elm wilt disease (due to *Ophiostoma spp.*) the most sensitive species and varieties present wider

xylem vessels (Elgersma, 1970; Mcnabb *et al.*, 1970; Solla and Gil 2002; Pita *et al.*, 2018).

Smaller vessels could occlude faster, sustaining a more efficient pathogen restriction (Venturas

et al., 2014). Our results on xylem vessel anatomy suggest that stems with tyloses tend to

present higher densities of small vessels, even if we did not observe any differences in total k_{th}

values and microCT scans showed that occlusions appear randomly in every vessel size class

(Fig. S3). It could be possible that tylose formation might be interfering with stem water

relations reducing the carbohydrates available for plant growth, producing smaller vessels in

stems of symptomatic plants. In contrast, artificial inoculations showed that xylem vessel

diameter had a strong impact on esca-related vascular pathogen development (Pouzoulet *et al.*,

2017; 2020), and in the kinetic of vessel occlusion in grapevine stems (Pouzoulet *et al.*, 2019).

The relationships between esca leaf symptoms, xylem anatomy, and tylose presence should be

studied in detail in trunks, where vascular pathogens are present, and among different grapevine

varieties and rootstocks as they are known to show different susceptibility to symptom

expression.

Long-term consequences of esca on leaf symptom expression and stem hydraulic integrity

In field surveys, esca leaf symptoms are often randomly distributed spatially throughout

vineyards and are not consistent from season to season in individual vines (Mugnai *et al.*, 1999;

Surico *et al.*, 2000; Marchi *et al.*, 2006; Guerin-Dubrana *et al.*, 2013; Li *et al.*, 2017). However,

esca-related vine death is strongly related to leaf symptoms as death is usually observed

following a year with symptom expression (Guerin-Dubrana *et al.*, 2013). In agreement with

these field studies, we observed similar percentages of symptomatic plants between those that

had already expressed esca symptoms in the past (from one to seven consecutive years, pS

plants), and those that had never expressed symptoms over the past seven years (pA plants).

However, we also found that pS plants expressed symptoms earlier in the season than pA

plants, suggesting that symptoms might require more time to develop in pA plants. We did not

find any significant differences in k_s and k_{th} values between plants with contrasted long-term

symptom history. This result suggests that esca leaf symptoms may have xylem anatomical

consequences within the year of expression by the production of tyloses, but not across seasons.

Moreover, we showed that DNA pathogen amount (*Phaeoacremonium minimum* and

Phaeomoniella chlamydospora) depends on the symptom expression in the season of sampling,

and on the long-term symptom history. Altogether, these results suggest that a higher amount

of vascular fungi in the trunk represents a higher risk in reproducing leaf symptoms, and

consequently, a higher risk of plant death.

Hydraulic failure and esca leaf symptom pathogenesis

Our results showed that, even if esca-related stem occlusion was extremely variable, 40% of

the microCT analyzed stems presented a total PLC greater than 50%. Under drought conditions

alone, studies suggest that grapevines are not able to recover in the current season from PLC

greater than 50% in stems (Charrier *et al*., 2018). Thus, to what extent these levels of esca-

induced hydraulic failure compromise future vine performance, and/or increase the likelihood

of developing esca leaf symptoms in the future remains an open question.

We showed that, similarly to visual leaf symptoms, tyloses in stems were generated at a

distance from the pathogen niche in the trunk. Comparing our results with Bortolami *et al*.

(2019), we show that the PLC due to the occlusions (hydraulic failure) observed using microCT

in leaves was on average twice higher than the PLC observed in stems in the present work. We

could hypothesize that, following pathogen activities in the trunk, a signal passing through the

xylem network and stimulating tyloses, first accumulates in leaves and then affects the stems.

However, the exact signal and action remain unknown, as we showed that the presence of

tyloses depended upon given symptomatic stems rather than symptomatic plants (i.e. two stems

in the same plant, with same tiger-stripe symptoms, sampled at the same moment, could or

could not present tyloses).

We showed that there were no differences in symptom expression, nor in the stem hydraulic

properties, regarding the long-term symptom history. We can conclude that the processes that

generate tiger-stripe symptoms are largely restricted to the current year of the symptom

expression. However, in plants expressing symptoms for the first time according to our disease

record, these processes could require more time, as they showed symptoms only late in the season. The presence of occlusion, leading to hydraulic failure in stems, could exacerbate leaf symptom expression in the following seasons, possibly contributing to death. We could speculate that a stem expressing extensive hydraulic failure could be more prone to express symptoms in the following year or, in the worst cases, to die. If the level of hydraulic failure could affect the stem mortality in the following year, the choice of stems with a complete absence of failure during the winter pruning could reduce the impact of esca in vineyards. The pruning practices are known to impact the course of infection and leaf symptom development and it has been shown that trunk renewal could be an effective management practice to prevent grapevine trunk diseases in the vineyard (Travadon *et al*. 2016, Kaplan *et al*. 2016, Gramaje *et al*. 2018). In addition, the presence of occlusions could also amplify plant susceptibility to drought-induced hydraulic failure, enhancing the risk of plant mortality in the field as suggested by McDowell *et al*. (2008). It could be speculated that a decrease in soil water potential or a high evaporative demand, concomitant to esca-induced hydraulic failure, could embolize the remaining functional xylem vessels stopping the water flow and desiccating plant tissues (this could be the case in apoplectic plants for example). In perspective, future studies should investigate the link between pathogen activities and occlusion development, especially in trunks, and the subsequent hydraulic failure consequences on whole plant physiology.

SUPPORTING INFORMATION

The following Supporting Information is available for this article:

Fig. S1. Two-dimensional reconstruction of longitudinal cross sections from X-ray microCT volumes of grapevine stems.

Fig. S2. Relationship between k_s and k_{th} in control plants.

Fig. S3. Vessel density and percentage of occluded vessels in tiger-stripe stems for different vessel diameter classes.

Fig. S4. Relationships between k_s and k_{th} in each stem symptom category.

Table S1. Effect of year of uprooting, internode analyzed, and sampling date on k_s and k_{th} in control stems.

Table S2. Calculated theoretical hydraulic conductivity (k_{th} %), and hydraulic conductivity loss (PLC %) from X-ray microCT volumes of intact grapevine stems.

DATA AVAILABILITY STATEMENT

Raw datasets are available in the INRAE dataverse: Bortolami, Giovanni; Farolfi, Elena; Badel, Eric; Burlett, Regis; Cochard, Herve; Ferrer, Nathalie; King, Andrew; Lamarque, Laurent J.; Lecomte, pascal; Marchesseau-Marchal, Marie; Pouzoulet, Jerome; Torres-Ruiz, Jose M.; Trueba, Santiago; Delzon, Sylvain; Gambetta, Gregory A.; Delmas, Chloe E.L., 2021, "Raw data for the paper "Seasonal and long-term consequences of esca on grapevine stem xylem integrity"", https://doi.org/10.15454/U9KJEW, Portail Data INRAE, V1.

ACKNOWLEDGMENTS

We thank the experimental teams of UMR SAVE and UMR EGFV (Bord'O platform, INRAE, Bordeaux, France) and the SOLEIL synchrotron facility (beamline PSICHE) for providing the materials and logistics. Specifically, we thank Jérôme Jolivet and Sebastien Gambier (UMR SAVE) for providing technical knowledge and support for plant transplantation and maintenance. This work was supported by the French Ministry of Agriculture, Agrifood, and Forestry (FranceAgriMer and CNIV) within the PHYSIOPATH project (program Plan National Dépérissement du Vignoble, 22001150-1506) awarded to C.E.L.D., and program Investments for the Future (ANR-10-EQPX-16, XYLOFOREST).

AUTHOR CONTRIBUTIONS

C.E.L.D., G.A.G., G.B., and S.D. designed the experiments;

E.F., G.A.G., S.D., E.B., R.B., H.C., A.K., L.J.L., J.M.T.-R., S.T. participated in synchrotron campaigns;

G.B., C.E.L.D., E.F., and N.F. conducted the esca symptom notations;

G.B., M.M.-M.,and N.F. conducted the histological observations;

E.F. conducted the hydraulic conductivity measurements and participated to data analyses;

N.F., and J.P., conducted the pathogen detection;

G.B. analyzed the microCT, optical images, and analyzed the data;

P.L. provided data on disease history of the plants;

G.B., C.E.L.D., and G.A.G. wrote the article;

all authors edited and agreed on the last version of the article

REFERENCES

Akpaninyang FE, Opara EU. 2017. The Influence of Toxins in Disease Symptom Initiation in Plants: A Review. Journal of Agriculture and Sustainability 10, 29-52.

Aleemullah M., Walsh K. 1996. Australian papaya dieback: Evidence against the calcium deficiency hypothesis and observations on the significance of laticifer autofluorescence. Australian Journal of Agricultural Research 47, 371-385.

Alvindia DG, Gallema FLM. 2017. *Lasiodiplodia theobromae* causes vascular streak dieback (VSD)–like symptoms of cacao in Davao Region, Philippines. Australasian Plant Disease Notes 12, 54.

Anderegg WRL, Kane JM, Anderegg LDL. 2013. Consequences of widespread tree mortality triggered by drought and temperature stress. Nature Climate Change 3, 30-36.

Anderegg WRL, Klein T, Bartlett M, Sack L, Pellegrini AFA, Choat B, Jansen S. 2016. Meta-analysis reveals that hydraulic traits explain cross-species patterns of drought-induced tree mortality across the globe. Proceedings of the National Academy of Sciences 113, 5024-5029.

Andolfi A, Mugnai L, Luque J, Surico G, Cimmino A, Evidente A. 2011. Phytotoxins Produced by Fungi Associated with Grapevine Trunk Diseases. Toxins 3, 1569-1605.

Beckman CH, Roberts EM. 1995. On the Nature and Genetic Basis for Resistance and Tolerance to Fungal Wilt Diseases of Plants. Advances in Botanical Research 21, 35-77.

Beier GL, Held BW, Giblin CP, Cavender-Bares J, Blanchette RA. 2017. American elm cultivars: Variation in compartmentalization of infection by *Ophiostoma novo-ulmi* and its effects on hydraulic conductivity. Forest Pathology 47, 1-11.

Bettenfeld P, Fontaine F, Trouvelot S, Fernandez O, Courty P-E. 2020. Woody Plant Declines. What's Wrong with the Microbiome? Trends in Plant Science 25, 381-394.

Bonsen KJM, Kučera LJ. (1990). Vessel Occlusions in Plants: Morphological, Functional and Evolutionary Aspects. IAWA Journal 11, 393-399.

Bortolami G, Gambetta GA, Delzon S, *et al*. (2019). Exploring the Hydraulic Failure Hypothesis of Esca Leaf Symptom Formation. Plant Physiology 181, 1163-1174.

Brown AA, Lawrence DP, Baumgartner K. 2020. Role of basidiomycete fungi in the grapevine trunk disease esca. Plant Pathology 69, 205-220.

Cailleret M, Jansen S, Robert EMR, *et al*. 2017. A synthesis of radial growth patterns preceding tree mortality. Global Change Biology 23, 1675-1690.

Charrier G, Delzon S, Domec J-C, *et al*. 2018. Drought will not leave your glass empty: Low risk of hydraulic failure revealed by long-term drought observations in world's top wine regions. Science Advances 4, 1-9.

Charrier G, Torres-Ruiz JM, Badel E, *et al*. 2016. Evidence for Hydraulic Vulnerability Segmentation and Lack of Xylem Refilling under Tension. Plant Physiology 172, 1657-1668.

Claverie M, Notaro M, Fontaine F, Wery J. 2020. Current knowledge on Grapevine Trunk Diseases with complex etiology : A systemic approach. Phytopathologia Mediterranea 59, 29-53.

Clérivet A, Déon V, Alami I, Lopez F, Geiger J-P, Nicole M. 2000. Tyloses and gels associated with cellulose accumulation in vessels are responses of plane tree seedlings (*Platanus × acerifolia*) to the vascular fungus *Ceratocystis fimbriata* f. Sp *platani*. Trees 15, 25-31.

Cloete M, Mostert L, Fischer M, Halleen F. 2015. Pathogenicity of South African Hymenochaetales taxa isolated from esca-infected grapevines. Phytopathologia Mediterranea 54, 368-379.

Collins BR, Parke JL, Lachenbruch B, Hansen EM. 2009. The effects of *Phytophthora ramorum* infection on hydraulic conductivity and tylosis formation in tanoak sapwood. Canadian Journal of Forest Research 39, 1766-1776.

Czemmel S, Galarneau ER, Travadon R, McElrone AJ, Cramer GR, Baumgartner K. 2015. Genes expressed in grapevine leaves reveal latent wood infection by the fungal pathogen *Neofusicoccum parvum*. PloS one 10, e0121828.

Desprez-Loustau M-L, Marçais B, Nageleisen L-M, Piou D, Vannini A. 2006. Interactive effects of drought and pathogens in forest trees. Annals of Forest Science 63, 597-612.

Deyett E, Pouzoulet J, Yang J-I, Ashworth VE, Castro C, Roper MC, Rolshausen PE. 2019. Assessment of Pierce's disease susceptibility in *Vitis vinifera* cultivars with different pedigrees. Plant Pathology 68, 1079-1087.

De Micco V, Balzano A, Wheeler EA, Baas P. 2016. Tyloses and gums : A review of structure, function and occurrence of vessel occlusions. IAWA Journal 37, 186-205.

Elgersma DM. 1970. Length and diameter of xylem vessels as factors in resistance of elms to Ceratocystis ulmi. Netherlands Journal of Plant Pathology 76, 179-182.

Eskalen A, Stouthamer R, Lynch SC, Rugman-Jones PF, Twizeyimana M, Gonzalez A, Thibault T. 2013. Host Range of Fusarium Dieback and Its Ambrosia Beetle (Coleoptera : Scolytinae) Vector in Southern California. Plant Disease 97, 938-951.

Et-Touil A, Rioux D, Mathieu FM, Bernier L. 2005. External symptoms and histopathological changes following inoculation of elms putatively resistant to Dutch elm disease with genetically close strains of *Ophiostoma*. Canadian Journal of Botany 83, 656–667.

Fallon B, Yang A, Lapadat C, Armour I, Juzwik J, Montgomery RA, Cavender-Bares J. 2020. Spectral differentiation of oak wilt from foliar fungal disease and drought is correlated with physiological changes. Tree Physiology 40, 377-390.

Fradin EF, Thomma BPHJ. 2006. Physiology and molecular aspects of *Verticillium* wilt diseases caused by *V. dahliae* and *V. albo-atrum*. Molecular Plant Pathology 7, 71-86.

Gärtner H, Lucchinetti S, Schweingruber FH. 2014. New perspectives for wood anatomical analysis in dendrosciences: the GSL1-microtome. Dendrochronologia 32, 47-51.

Goberville E, Hautekèete N-C, Kirby RR, Piquot Y, Luczak C, Beaugrand G. 2016. Climate change and the ash dieback crisis. Scientific Reports 6 doi:10.1038/srep35303.

Gramaje D, Úrbez-Torres JR, Sosnowski MR. 2018. Managing Grapevine Trunk Diseases With Respect to Etiology and Epidemiology : Current Strategies and Future Prospects. Plant Disease 102, 12-39.

Guérard N, Maillard P, Bréchet C, Lieutier F, Dreyer E. 2007. Do trees use reserve or newly assimilated carbon for their defense reactions? A 13C labeling approach with young Scots pines inoculated with a bark-beetle-associated fungus (*Ophiostoma brunneo ciliatum*). Annals of Forest Science 64, 601-608.

Guerin-Dubrana L, Labenne A, Labrousse JC, Bastien S, Rey P, Gegout-Petit A. 2013. Statistical analysis of grapevine mortality associated with esca or Eutypa dieback foliar expression. Phytopathologia Mediterranea 52, 276-288.

Guerin-Dubrana L, Fontaine F, Mugnai L. 2019. Grapevine trunk disease in European and Mediterranean vineyards : Occurrence, distribution and associated disease-affecting cultural factors. Phytopathologia Mediterranea 58, 49-71.

Hochberg U, Albuquerque C, Rachmilevitch S, Cochard H, David-Schwartz R, Brodersen CR, McElrone A, Windt CW. 2016. Grapevine petioles are more sensitive to drought induced embolism than stems : Evidence from *in vivo* MRI and microcomputed tomography observations of hydraulic vulnerability segmentation: Hydraulic vulnerability segmentation in grapevine. Plant, Cell & Environment 39, 1886-1894.

Hochberg U, Windt CW, Ponomarenko A, Zhang Y-J, Gersony J, Rockwell FE, Holbrook NM. 2017. Stomatal Closure, Basal Leaf Embolism, and Shedding Protect the Hydraulic Integrity of Grape Stems. Plant Physiology 174, 764-775.

Jacobi WR, MacDonald WL. 1980. Colonization of resistant and susceptible oaks by *Ceratocystis fagacearum*. Phytopathology 70, 618-623.

Kaplan J, Travadon R, Cooper M, Hillis V, Lubell, Baumgartner K. 2016. Identifying economic hurdles to early adoption of preventative practices_ The case of trunk diseases in California winegrape vineyards. Wine Economics and Policy 15, 127-141.

King A, Guignot N, Zerbino P, *et al.* 2016. Tomography and imaging at the PSICHE beam line of the SOLEIL synchrotron. Review of Scientific Instruments 87, 093704.

Lachenbruch B, Zhao J-P. 2019. Effects of phloem on canopy dieback, tested with manipulations and a canker pathogen in the *Corylus avellana/Anisogramma anomala* host/pathogen system. Tree Physiology 39, 1086-1098.

Lecomte P, Darrieutort G, Liminana J-M, Comont G, Muruamendiaraz A, Legorburu F-J, Choueiri E, Jreijiri F, El Amil R, Fermaud M. 2012. New Insights into Esca of Grapevine: The Development of Foliar Symptoms and Their Association with Xylem Discoloration. Plant Disease 96, 924-934.

Li S, Bonneu F, Chadoeuf J, Picart D, Gégout-Petit A, Guérin-Dubrana L. 2017. Spatial and Temporal Pattern Analyses of Esca Grapevine Disease in Vineyards in France. Phytopathology 107, 59-69.

Martin N, Vesentini D, Rego C, Monteiro S, Oliveira H, Ferreira RB. 2009. *Phaeomoniella chlamydospora* infection induces changes in phenolic compounds content in *Vitis vinifera*. Phytopathologia Mediterranea 48, 101-116.

Martín JA, Solla A, Ruiz-Villar M, Gil L. 2013. Vessel length and conductivity of *Ulmus* branches : Ontogenetic changes and relation to resistance to Dutch elm disease. Trees 27, 1239-1248.

McDowell N, Pockman WT, Allen CD, *et al*. 2008. Mechanisms of plant survival and mortality during drought : Why do some plants survive while others succumb to drought? New Phytologist 178, 719-739.

McElrone AJ, Sherald JL, Forseth IN. 2003. Interactive effects of water stress and xylem-limited bacterial infection on the water relations of a host vine. Journal of Experimental Botany 54, 419-430.

McElrone AJ, Jackson S, Habdas P. 2008. Hydraulic disruption and passive migration by a bacterial pathogen in oak tree xylem. Journal of Experimental Botany 59, 2649-2657.

McNabb HS, Heybroek HM, Macdonald WL. 1970. Anatomical factors in resistance to Dutch elm disease. Netherlands Journal of Plant Pathology 76, 196-205.

Mensah JK, Sayer MAS, Nadel RL, Matusick G, Eckhardt LG. 2020. Physiological response of *Pinus taeda* L. trees to stem inoculation with *Leptographium terebrantis*. Trees 34, 869–880.

Mirone A, Brun E, Gouillart E, Tafforeau P, Kieffer J. 2014. The PyHST2 hybrid distributed code for high speed tomographic reconstruction with iterative reconstruction and a

priori knowledge capabilities. Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms 324, 41-48.

Morales-Cruz A, Allenbeck G, Figueroa-Balderas R, Ashworth VE, Lawrence DP, Travadon R, Smith RJ, Baumgartner K, Rolshausen PE, Cantu D. 2018. Closed-reference metatranscriptomics enables *in planta* profiling of putative virulence activities in the grapevine trunk disease complex: Transcriptomics of pathogen communities. Molecular Plant Pathology 19, 490-503.

Mugnai L, Graniti A, Surico G. 1999. Esca (Black Measles) and Brown Wood-Streaking : Two Old and Elusive Diseases of Grapevines. Plant Disease 83, 404-418.

Ouadi L, Bruez E, Bastien S, Vallance J, Lecomte P, Domec J-C, Rey P. 2019. Ecophysiological impacts of Esca, a devastating grapevine trunk disease, on *Vitis vinifera* L. PloS one 14, 1-20.

Ouellette GB, Baayen RP, Simard M, Rioux D. 1999. Ultrastructural and cytochemical study of colonization of xylem vessel elements of susceptible and resistant *Dianthus caryophyllus* by *Fusarium oxysporum* f.sp. Dianthi. Canadian Journal of Botany 77, 644-663.

Paganin D, Mayo SC, Gureyev TE, Miller PR, Wilkins SW. 2002. Simultaneous phase and amplitude extraction from a single defocused image of a homogeneous object. Journal of Microscopy 206, 33-40.

Pandey S, Rishi RR, Jayaraj R, Giri K, Kumar R, Pandey A, Juwantha R, Madaan S, Bhandari MS. 2019. *Fusarium equiseti* is associated with the wilt and dieback of *Aquilaria malaccensis* in Northeast India. Forest Pathology 49, e12489.

Park J-H, Juzwik J. 2014. *Ceratocystis smalleyi* colonization of bitternut hickory and host responses in the xylem. Forest Pathology 44, 282-292.

Parke JL, Oh E, Voelker S, Hansen EM, Buckles G, Lachenbruch B. 2007. *Phytophthora ramorum* Colonizes Tanoak Xylem and Is Associated with Reduced Stem Water Transport. Phytopathology 97, 1558-1567.

Pearce RB. 1996. Antimicrobial defences in the wood of living trees. New Phytologist 132, 203-233.

Pérez-Donoso AG, Lenhof JJ, Pinney K, Labavitch JM. 2016. Vessel embolism and tyloses in early stages of Pierce's disease. Australian Journal of Grape and Wine Research 22, 81-86.

Pita P, Rodríguez-Calcerrada J, Medel D, Gil L. 2018. Further insights into the components of resistance to *Ophiostoma novo-ulmi* in *Ulmus minor* : Hydraulic conductance, stomatal sensitivity and bark dehydration. Tree Physiology 38, 252-262.

Pouzoulet J, Mailhac N, Couderc C, Besson X, Daydé J, Lummerzheim M, Jacques A. 2013. A method to detect and quantify *Phaeomoniella chlamydospora* and *Phaeoacremonium aleophilum* DNA in grapevine-wood samples. Applied Microbiology and Biotechnology 97, 10163-10175.

Pouzoulet J, Scudiero E, Schiavon M, Rolshausen PE. 2017. Xylem Vessel Diameter Affects the Compartmentalization of the Vascular Pathogen *Phaeomoniella chlamydospora* in Grapevine. Frontiers in Plant Science 8, 1-13.

Pouzoulet J, Scudiero E, Schiavon M, Santiago LS, Rolshausen PE. 2019. Modeling of xylem vessel occlusion in grapevine. Tree Physiology 39, 1438-1445.

Pouzoulet J, Rolshausen PE, Charbois R, Chen J, Guillaumie S, Ollat N, Gambetta GA, Delmas CEL. 2020. Behind the Curtain of the Compartmentalization Process : Exploring How Xylem Vessel Diameter Impacts Vascular Pathogen Resistance. Plant, Cell & Environment doi: 10.1111/pce.13848.

Pratt RB, Jacobsen AL. 2018. Identifying which conduits are moving water in woody plants: A new HRCT-based method. Tree Physiology 38, 1200-1212.

Rioux D, Blais M, Nadeau-Thibodeau N, Lagacé M, DesRochers P, Klimaszewska K, Bernier L. 2018. First Extensive Microscopic Study of Butternut Defense Mechanisms Following Inoculation with the Canker Pathogen *Ophiognomonia clavigignenti-juglandacearum* Reveals Compartmentalization of Tissue Damage. Phytopathology 108, 1237-1252.

Salle A, Ye H, Yart A, Lieutier F. 2008. Seasonal water stress and the resistance of *Pinus yunnanensis* to a bark-beetle-associated fungus. Tree Physiology 28, 679-687.

Salleo S, Nardini A, Lo Gullo MA, Ghirardelli LA. 2002. Changes in stem and leaf hydraulics preceding leaf shedding in *Castanea sativa* L. Biologia plantarum 45, 227-234.

Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ : 25 years of image analysis. Nature Methods 9, 671-675.

Shigo AL. 1984. Compartmentalization: A conceptual framework for understanding how trees grow and defend themselves. Annual Review of Phytopathology, 22, 189–214.

Sperry JS, Donnelly JR, Tyree MT. 1988. A method for measuring hydraulic conductivity and embolism in xylem. Plant, Cell & Environment 11, 35-40.

Solla A, Gil L. 2002. Xylem vessel diameter as a factor in resistance of *Ulmus minor* to *Ophiostoma novo-ulmi*. Forest Pathology 32, 123-134.

Sun Q, Rost TL, Matthews MA. 2008. Wound-induced vascular occlusions in *Vitis vinifera* (Vitaceae) : Tyloses in summer and gels in winter. American Journal of Botany 95, 1498-1505.

Sun Q, Sun Y, Walker MA, Labavitch JM. 2013. Vascular Occlusions in Grapevines with Pierce's Disease Make Disease Symptom Development Worse. Plant Physiology 161, 1529-1541.

Surico G, Marchi G, Ferrandino A, Braccini P, Mugnai L. 2000. Analysis of the spatial spread of esca in some Tuscan vineyards (Italy). Phytopathologia Mediterranea 39, 211-224.

Talboys PW. 1972. Resistance to Vascular Wilt Fungi. Proceedings of the Royal Society of London. Series B, Biological Science 181, 319-332.

Torres-Ruiz JM, Sperry JS, Fernández JE. 2012. Improving xylem hydraulic conductivity measurements by correcting the error caused by passive water uptake. Physiologia Plantarum 146, 129-135.

Travadon R, Lecomte P, Diarra B, Lawrence DP, Renault D, Ojeda H, Rey P, Baumgartner K. 2016. Grapevine pruning systems and cultivars influence the diversity of wood-colonizing fungi. Fungal Ecology 24 82-93.

Tyree MT, Sperry JS. 1989. Vulnerability of Xylem to Cavitation and Embolism. Annual Review of Plant Physiology and Plant Molecular Biology 40, 19-38.

Tyree MT, Ewers FW. 1991. The hydraulic architecture of trees and other woody plants. New Phytologist 119, 345-360.

Úrbez-Torres JR, Peduto F, Vossen PM, Krueger WH, Gubler WD. 2013. Olive Twig and Branch Dieback: Etiology, Incidence, and Distribution in California. Plant Disease 97, 231-244.

Venturas M, López R, Martín JA, Gascó A, Gil L. 2014. Heritability of *Ulmus minor* resistance to Dutch elm disease and its relationship to vessel size, but not to xylem vulnerability to drought. Plant Pathology 63, 500-509.

Yadeta KA, Thomma BPHJ. 2013. The xylem as battleground for plant hosts and vascular wilt pathogens. Frontiers in Plant Science 4 doi:10.3389/fpls.2013.00097.

Yazaki K, Takanashi T, Kanzaki N, Komatsu M, Levia DF, Kabeya D, Tobita H, Kitao M, Ishida A. 2018. Pine wilt disease causes cavitation around the resin canals and irrecoverable xylem conduit dysfunction. Journal of Experimental Botany 69, 589-602.

Zimmermann MH. 1979. The Discovery of Tylose Formation by a Viennese lady in 1845. IAWA Bulletin 2, 51-56.

TABLES

Table 1. Esca leaf symptom observations over the experimental season on *Vitis vinifera* cv Sauvignon blanc.

Symptom notation before 2019	All plants	Previously asymptomatic (pA)	Previously symptomatic (pS)
Symptom notation in 2019			
Esca-symptomatic	35 % (20/58)	30 % (6/20)	37 % (14/38)
Control-asymptomatic	65 % (38/58)	70 % (14/20)	63 % (24/38)

Plants are grouped by their symptom history: previously asymptomatic (pA, plants that have never expressed leaf symptoms between 2012 and 2018) and previously symptomatic (pS, plants that have expressed leaf symptoms at least once since 2012). Ratios present the number of plants in each symptom category (esca-symptomatic or control-asymptomatic) over the total number of plants of the category.

Table 2. Values for specific stem hydraulic conductivity (k_s), theoretical stem hydraulic conductivity (k_{th}) and equations of regression lines between k_s and k_{th} for control and esca symptomatic stems.

Tyloses	Esca	k_s [kg s^{-1} m^{-1} MPa^{-1}]	k_{th} [kg s^{-1} m^{-1} MPa^{-1}]	n (stem - plant)	Regression
Absence	Control	24.97 ± 1.72	78.36 ± 4.51	39 - 23	$k_s = 0.3$ x $k_{th} + 1.6$ R²=0.61 P<0.0001
	Asymptomatic before symptoms	26.04 ± 4.71	74.75 ± 11.80	6 - 6	$k_s = 0.36$ x $k_{th} - 0.94$ R²=0.82 P=0.013
	Pre-symptomatic	30.32 ± 4.26	87.88 ± 8.54	11 - 7	$k_s = 0.37$ x $k_{th} - 1.9$ R²=0.54 P=0.010
	Asymptomatic after symptoms	19.80 ± 5.18	72.58 ± 12.64	5 - 2	$k_s = 0.33$ x $k_{th} - 4$ R²=0.64 P=0.104
	Esca (tiger-stripe)	21.29 ± 5.40	72.85 ± 10.41	9 - 5	$k_s = 0.45$ x $k_{th} - 11.26$ R²=0.74 P=0.003
Presence	Esca (tiger-stripe)	11.27 ± 2.86	70.44 ± 7.81	15 - 5	$k_s = 0.17$ x $k_{th} -0.84$ R²=0.22 P=0.077
	Esca (apoplectic)	2.47 ± 1.45	74.80 ± 33.48	3 - 2	$k_s = 0.04$ x $k_{th} - 0.2$ R²=0.68 P=0.385
Absence	All	25.06 ± 1.46	78.42 ± 3.37	70 - 37	$k_s = 0.34$ x $k_{th} - 1.90$ R²=0.63 P<0.0001
Presence	All	9.81 ± 2.51	71.16 ± 8.00	18 - 7	$k_s = 0.12$ x $k_{th} - 1.28$ R²=0.15 P=0.117

Values represent mean ± SE. n = sample size, (including the number of analyzed stems and number of analyzed plants, respectively). See text and Fig. 4 for statistical analysis. A detailed esca symptom notation is provided in Fig. 1A. Bivariate plots of each regression are presented in Fig. S4.

Table 3. Long-term impact of symptom presence (i.e. comparing plants with different disease history record) in control plants on specific stem hydraulic conductivity (k_s), theoretical stem hydraulic conductivity (k_{th}), stem total vessel density, and amount of *Phaeomoniella chlamydospora* and *Phaeoacremonium minimum* DNA in trunks of plants without foliar symptoms.

	Previously asymptomatic (pA)	Previously asymptomatic (pS)	Type III Tests of Fixed Effects (pA vs pS)
k_s [kg s^{-1} m^{-1} MPa^{-1}]	23.76 ± 2.30	26.54 ± 2.61	n=39, $F_{1,16}$=1.19, P=0.29
k_{th} [kg s^{-1} m^{-1} MPa^{-1}]	72.22 ± 4.85	86.30 ± 7.98	n=39, $F_{1,16}$=3.01, P=0.10
total vessel density [count mm^{-2}]	57.28 ± 4.03	52.61 ± 3.25	n=39, $F_{1,16}$=0.72, P=0.41
P. chlamydospora [pg ng^{-1}]	6.14 ± 1.90	10.15 ± 3.41	**n=13, $F_{1,11}$=5900.06, P<0.0001**
P. minimum [pg ng^{-1}]	9.27 ± 6.97	26.40 ± 13.83	**n=13, $F_{1,11}$=51014, P<0.0001**

Values represent means ± SE. Pathogen quantification was estimated as: pg fungal DNA ng^{-1} total DNA. Statistical tests used are individual generalized linear mixed models to compare pA vs pS plants (fixed effect) with the individual plants entered as a random effect in the models and the year of uprooting as a co-variable (fixed effect). Statistically significant results (P<0.05) are shown in bold.

Table 4. Quantification by qPCR of *Phaeomoniella chlamydospora* and *Phaeoacremonium minimum* DNA in stems and trunks of different esca symptomatology.

Organ	Esca	n	*P. chlamydospora* [pg ng^{-1}]	*P. minimum* [pg ng^{-1}]
Stem	Control	8	0	0
	Pre-symptomatic	3	0	0
	Asymptomatic (after symptoms)	3	0	0
	Tiger-stripe (without tyloses)	4	0	0
	Tiger-stripe (with tyloses)	8	0	0
	Apoplectic	2	0	0
Trunk	Control	13	7.37 ± 1.67 (12/13)[*]	14.54 ± 6.51 (12/13)[*]
	Symptomatic	7	15.80 ± 3.12 (7/7)[*]	23.90 ± 8.82 (7/7)[*]

[*] Number of samples positive for the pathogen over the total number of analyzed samples.

Pathogen quantification was estimated as: pg fungal DNA per ng total DNA. Values represent means ± SE, n = sample size. Trunks of symptomatic plants presented higher amount of both *P. chlamydospora* and *P. minimum*, compared to control (n=20, $F_{1,18}$=29806.11.25, P<0.0001 and n=20, $F_{1,18}$=21925.4, P<0.0001, respectively). See text for statistical methods.

FIGURES

Fig. 1. Representation of esca symptom notation during the experimental season. (**A**) Single stems could be noted as esca asymptomatic, pre-symptomatic, tiger-stripe, or apoplectic. Whole plants have been noted as control (asymptomatic from June to October) or symptomatic (with tiger-stripe symptoms at the end of the season). (**B**) Proportion of plants in each symptom category over the experimental season (n=58). The blue area corresponds to control plants, green area to esca symptomatic plants before symptom appearance, and red area to esca symptomatic plants.

Fig. 2. Two-dimensional reconstruction of cross-sections from X-ray microCT volumes of grapevine stems. Each panel represents a cross section of different stems for control (**A-C**) and esca symptomatic (**D-M**) plants. Iohexol appears white bright in functional vessels; air-filled vessels (i.e. native PLC) appear black; occluded vessels (i.e. occlusion PLC) appear grey. Total PLC (i.e. native PLC plus occlusion PLC) values are given for the presented samples. Scale bars = 1000µm.

Fig. 3. (**A**) Mean values of native PLC in control (blue) and esca tiger-stripe (red) stems of grapevine plants using X-ray microCT imaging. Differences were not significant (n=13, $F_{1,9}$=0.07, P=0.79). (**B**) Mean values of occlusion PLC in control (blue) and esca tiger-stripe (red) stems of grapevine plants using X-ray microCT imaging. Differences were not significant (n=13, $F_{1,9}$=0.33, P=0.58). Boxes and bars show the median, quartiles and extreme values, circles show mean values. N represents the sample size (number of analyzed stems) for each group.

Fig. 4. Relationships between specific stem hydraulic conductivity (k_s), theoretical stem hydraulic conductivity (k_{th}), and vessel density in control and esca symptomatic grapevine plants. (**A**) k_s values for control (blue); asymptomatic (dark green) and pre-symptomatic (yellow) stems in plants before symptom appearance; asymptomatic (light green), tiger-stripe (red), and apoplectic (grey) stems in plants after symptom appearance, differences were not significant (n=88, $F_{5,45}$=1.30 P=0.28). Boxes and bars show the median, quartiles and extreme values, circles within boxes correspond to means, and circles outside boxes to outlier values. (**B**) Relationships between k_s and k_{th}. Symbols represent the absence (circles) or presence (triangles) of tyloses in xylem vessels. Colors represent esca symptomatology (as in panel A). The dashed line represents the regression for stems in which no tyloses were observed in xylem vessels, and the solid line represents the regression for samples with tyloses. R^2 for the regression lines are indicated (see Table 2 and Fig. S4 for detailed analyses). (**C**) k_{th} values for the different stem categories as presented in panel a. Differences were not significant (n=88, $F_{5,45}$=0.58, P=0.71). (**D**) Relationships between mean values of xylem vessel density and their diameters. Differences in total vessel density and in vessel size distributions were not significant (n=88, $F_{6,45}$=0.77, P=0.60; n=792 (88 samples for 9 vessel classes), $F_{48,693}$=1.19, P=0.18). Colors and markers are the same as panel B.